



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Tuning of synaptic responses: an organizing principle for optimization of neural circuits

Citation for published version:

O'Donnell, C & Nolan, MF 2011, 'Tuning of synaptic responses: an organizing principle for optimization of neural circuits', *Trends in Neurosciences*, vol. 34, no. 2, pp. 51-60. <https://doi.org/10.1016/j.tins.2010.10.003>

Digital Object Identifier (DOI):

[10.1016/j.tins.2010.10.003](https://doi.org/10.1016/j.tins.2010.10.003)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Trends in Neurosciences

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Tuning of synaptic responses: an organizing principle for optimization of neural circuit function

Cian O'Donnell ¹ and Matthew F. Nolan ²

¹ Neuroinformatics Doctoral Training Centre, University of Edinburgh, EH8 9AB, United Kingdom.

² Centre for Integrative Physiology, University of Edinburgh, EH8 9XD, United Kingdom.

Corresponding author: Nolan, M.F. (mattnolan@ed.ac.uk)

Abstract

Neuron types are classically defined by anatomical and physiological properties that determine how synaptic inputs are integrated. Here, we provide an overview of the evidence that among neurons of a single type, integration of synaptic responses is further tuned according to the particular function individual neurons carry out. Recent data suggest that tuning of synaptic responses is not restricted to sensory pathways, but extends to cognitive and motor circuits. We propose that tuning of synaptic integration results from general cellular mechanisms for optimization of information processing that are distinct from, but complementary to, homeostasis and memory storage. These cellular tuning mechanisms may be critical for distributed computations underlying sensory, motor and cognitive functions.

Introduction

Organizing principles for the development and function of nervous systems often divide neurons into distinct types on the basis of their morphology and intrinsic electrophysiological properties [1-4]. While this heterogeneity between neuronal types is well established and accepted as functionally important, diversity among neurons of the same type has received less attention. In topographically organized neural circuits (Glossary), encoded environmental features can be 'mapped' onto anatomically defined locations within the brain. For example, nearby neurons in the primary visual cortex often respond preferentially to nearby visual stimuli, creating a cortical map of the visual field [5]. Investigations of topographically organized auditory pathways [6, 7], and more recently cognitive circuits [8], suggest that among neurons of the same type, synaptically driven changes in membrane potential are tuned according to the specific function that each neuron carries out. Thus, just as the strings of a piano are tuned so that each key produces a different note, transformation of synaptic input into patterns of action potential output is tuned according to the particular information processing or behavioral task a neuron

contributes to. By determining each neuron's influence on its downstream targets, these systematic differences between neurons of the same type are likely to be critical to the function of neural circuits. This form of organization may also be of clinical significance. For example, deficits in tuning properties have recently been identified in a mouse model of Fragile X syndrome [9].

In this Opinion article, we argue that tuning of synaptic integration among neurons of the same type is a general principle of nervous system organization that extends beyond the early stages of sensory processing, to motor circuits and to brain areas that mediate complex cognitive tasks, such as navigation and memory. We will suggest how tuning could optimize encoding or computation by neural circuits and will identify predictions for circuit function that follow from this idea. While we focus on cellular properties that determine integration of responses downstream from synaptic activation of neurotransmitter receptors, similar principles may also apply to tuning of synaptic transmission [7, 10, 11].

Configuring synaptic integration, topographic organization of neural representations and experimental evaluation of response tuning

How a neuron responds to, or integrates, its synaptic inputs will dictate its contribution to the function of the circuit in which it is embedded. Different neuronal types configure their responses to synaptic input in diverse ways [12]. This diversification is achieved by differences in morphology and in the complement and localization of non-synaptic ion channels [12]. For example, neurons in auditory pathways employ low-threshold potassium channels to curtail synaptic responses and ensure timing of signals with sub-millisecond accuracy [13]. In contrast, these mechanisms are absent from many neocortical neurons, which often generate highly irregular output [14].

During many sensory, motor and cognitive processes, the brain represents information as patterns of activity distributed within populations of neurons of the same type [15-18]. For example, complex sounds are encoded in parallel by neurons that differ in the sound frequency to which they are most sensitive [19, 20]. Since individual neurons within a single population encode distinct features of a distributed representation, each neuron might be expected to configure the way it responds to synaptic input so as to best perform its specific role in the circuit. Investigation of this idea is more difficult than comparing different neuron types, which are relatively easy to identify, because it is challenging to determine in the same experiment both the salient information that a particular neuron encodes, or the specific computation that it carries out, and its cellular properties.

In a topographically organized neural circuit, this problem can be addressed instead by making intracellular recordings from neurons at defined locations and recording changes in the neuron's membrane potential caused by synaptic stimulation [6-8, 21]. If synaptic responses are tuned according to a neuron's particular function, then they should follow a topographic organization similar to the functional organization of the circuit (Figure 1A). Alternatively, if all neurons of a single type respond in a similar way to activation of their synaptic inputs, then there should not be any systematic anatomical organization of synaptic responses (Figure 1B). In this case, heterogeneity between neurons may instead reflect random variation. Similar tuning mechanisms might also exist in circuits without an easily defined topographic organization, but they will not be detectable by experimental methods that rely on correlating cellular properties with anatomical location (Figure 1C).

Topographically organized distributed representations have been identified in many brain areas, including auditory, somatosensory, visual, and motor control circuits [17, 22-27]. Topographically

organized encoding was also recently discovered in cognitive circuits that represent spatial location [28, 29] (Figure 2). We outline below studies that, by taking advantage of the topographical organization of neural circuits, provide evidence for functional tuning of synaptic responses within populations of a single neuron type.

Evidence for tuning of synaptic responses

Sensory and motor circuits

The importance of cellular mechanisms that determine precisely how neurons in auditory circuits respond to synaptic input has been recognized for some time [13, 30]. While many studies have focused on specialization of particular neuron types, there is now also considerable evidence for tuning of synaptic responses among auditory neurons of the same type (Table 1). For example, neurons in the spiral ganglion encode sound with a characteristic frequency that corresponds to their location along the tonotopic axis of the cochlea. Comparison of spiral ganglion neurons from different parts of the cochlea reveals a tonotopic organization of voltage- and calcium-gated potassium channels [31, 32]. In neurons that encode high frequency sounds, elevated expression of delayed rectifier potassium channels and calcium-activated potassium channels accelerates repolarization of action potentials and suppresses repetitive firing [31-33]. These and further observations [32-34] suggest that synaptic responses of spiral ganglion neurons are tuned to sound frequency. Nevertheless, the consequences for synaptic responses of spiral ganglion neurons are not completely clear, as these experiments focused on properties known to affect synaptic integration, but did not measure synaptically-driven changes in membrane potential. Indeed, with the exception of data obtained from neurons in the chick [6, 7, 11], most functional evidence for synaptic response tuning in auditory circuits relies on extrapolation from measures of ion channel expression and membrane potential responses to injection of current through a recording electrode, rather than direct observation of synaptic responses.

Evidence for tuning of synaptic integration also comes from investigations of sensory and motor circuits in fish. Electric fish detect electric fields signaling the presence of nearby electric fish, or their aquatic prey, using sensory receptors with axons that project to neurons in three segments of their electrosensory lateral line lobe (ELL) [22]. In each division, distinct pyramidal cell types called E-cells can be identified by anatomical and electrophysiological criteria [35]. The waveform, pattern and input frequency selectivity of action potentials fired by E-cells differs between segments of the ELL [35]. These differences may be explained by corresponding segmental differences in ion channel expression (e.g. Refs. [36, 37]). Since each ELL segment receives identical sensory input, but has distinct behavioral roles [38], these data suggest that responses to synaptic activation may be tuned according to a neuron's ultimate behavioral output as well as to its original sensory input. Consistent findings have also been demonstrated in zebrafish spinal cord. Here, motor neurons and premotor interneurons are topographically organized according to the minimal swimming frequency at which they become active [39]. Neurons that are active during lower swimming frequencies have a higher membrane resistance, suggesting they are more sensitive to synaptic input, compared with neurons that are active only during higher swimming frequencies [39].

Cognitive circuits

Does tuning of synaptic responses extend to cognitive circuits? Recent demonstrations of topographical organization of the resolution of spatial firing fields within the hippocampus [29] and medial entorhinal cortex (MEC) [28, 40, 41] (Figure 2) enable mechanisms important for encoding of spatial location to be addressed by comparing properties of neurons at different anatomically defined positions within these structures (e.g. Refs. [8, 42, 43]). Comparison of excitatory post-synaptic potentials (EPSPs) recorded from neurons at different locations along the

dorsal-ventral axis of layer II of the MEC, reveals that neurons in parts of the MEC that represent an animal's location at low spatial resolution generate EPSPs with a slower time course than EPSPs recorded from neurons at positions that represent location at a higher spatial resolution [8] (Figure 2). Thus, encoding of an animal's location may benefit from tuning of synaptic integration. This organization of synaptic responses modifies temporal integration of synaptic input and is explained by differences in the amplitude of currents mediated by hyperpolarization-activated cyclic nucleotide-gated (HCN) channels and leak potassium channels [8].

In addition to synaptic responses, a similar organization of intrinsic theta frequency oscillatory activity and resonance characteristics has been described for neurons in layer II of the MEC [42-45]. Although artificial conductances designed to mimic *in vivo* synaptic activity abolish intrinsic theta oscillatory activity [46], the dorsal-ventral organization of resonance frequency could reflect more robust differences in temporal filtering introduced by HCN channels [47]. A preliminary report suggests a similar organization of currents through HCN channels in hippocampal CA1 pyramidal neurons [48], indicating that tuning of synaptic responses may extend to circuits beyond the MEC.

Together, these data suggest that in auditory, motor and cognitive circuits, neurons tune integration of their synaptic inputs according to their particular function. It is not yet clear if similar organizational principles extend to other brain areas. Since visual pathways follow a topographical organization [5, 15], this may be relatively straightforward to test. In other cortical regions, for example frontal regions important for higher cognitive processes, it is not clear whether distributed representations follow any clear macroscopic topography (e.g. Figure 1C). In these brain areas, evaluation of synaptic tuning will require the direct measurement of both the behaviorally relevant information that a neuron encodes and its responses to synaptic input, for

example using patch-clamp recordings from neurons in awake or anesthetized preparations. Finally, some neuron types may benefit from random rather than tuned synaptic responses properties. For example, in the olfactory bulb, apparently random diversity in synaptic integration by mitral cells may improve population coding by de-correlating action potential firing [49].

Molecular mechanisms for tuning of synaptic integration

What molecular mechanisms coordinate tuning of synaptic integration? Investigations of tonotopic organization of auditory neurons suggest a number of possible mechanisms. First, differences in physiological properties of neurons are correlated with altered mRNA as well as protein levels [6, 21, 50], suggesting that tuning may involve control of transcription or mRNA stabilization. Second, tuning of synaptic responses can involve coordinated differences in more than one class of ion channel and can involve increases in one conductance accompanied by a reduction in another conductance (e.g. Refs. [31, 32]). This suggests that tuning is not a result of non-specific cell-wide up- or down-regulation of mRNA or protein levels. Third, tuning of synaptic responses appears to emerge relatively late in development and can be dependent on neuronal activity [9, 51, 52]. In the cochlea, opposing gradients in expression of brain-derived neurotrophic factor and neurotrophin-3 appear to drive activity-dependent changes in ion channel expression that underlie the tonotopic organization of firing properties of spiral ganglion neurons [31]. Fourth, tonotopic tuning of integrative properties can vary between species. For example, in the rat and gerbil medial nucleus of the trapezoid body (MNTB) [52, 53], there does not appear to be any tuning of sub-threshold properties, while in the mouse MNTB the amplitude of HCN currents appears to be larger in neurons that encode low frequency sounds compared to neurons that encode higher frequency sounds [51]. It is not yet clear if similar mechanisms are used by neurons in the MEC or hippocampus to implement tuning of synaptic integration.

Establishing how molecular signaling components interact to tune synaptic integration could provide important clues to the functions of tuning. In principle, tuning could be implemented in two ways. First, in common with molecular mechanisms used to establish differences between neuron types [4], gene regulatory pathways might also be configured to tune ion channel expression among neurons of the same type. Configuration of gene regulatory pathways could be established during development and then maintained in a cell autonomous fashion in mature animals, or it could be driven by signals used as position cues in the brain [54]. In a second scenario, all neurons of a single type may express the same genetically encoded regulatory machinery, with tuning instead being driven by the synaptic input a neuron receives. In this model, sensors are required to detect differences between the neuron's current state and its target state. Activation of the sensors should then lead to changes in ion channel function that reconfigure the neuron towards its target state. These two scenarios are not necessarily mutually exclusive. For example, core gene regulatory networks for establishing tuning may be subject to modulation by synaptic activity.

Contributions of tuning to neural circuit function

What possible functions does tuning of synaptic responses achieve? Here, we consider three general functions that may lead to the topographically organized tuned synaptic responses observed experimentally: storage of memories [55, 56] (Figure 3A); homeostatic stabilization of neuronal activity [57-60] (Figure 3B); and optimization of representation, transmission or processing of information [61-64] (Figure 4). Although these models are not necessarily mutually exclusive, they nevertheless make distinct functional predictions and so we consider them separately.

Memory storage

Topographically organized differences between neurons in their integrative properties might, in principle, serve as memory traces. While memories are usually thought to be stored by modification of synaptic connections, plasticity of ion channels that influence integration of synaptic inputs can play a similar role [55] (Figure 3A). For example, memory for a conditioned stimulus can be stored by an increase in the responsiveness of a neuron to synaptic input (e.g. Refs. [65, 66]). We consider this an unlikely explanation for tuning of synaptic integration for two reasons. First, the examples of tuning we discuss above have global effects on the excitability of a neuron, making them an inefficient way to store memories. In contrast, the high memory capacity of neural tissue is likely to rely on synapse-specific plasticity [67, 68]. Second, although synaptic response tuning is found in cortical circuits important for memory [8], it also occurs in circuits responsible for early stages of auditory processing. These circuits are generally believed to be hard-wired and therefore unlikely loci for memory storage. Nevertheless, there is evidence for plasticity of brainstem auditory circuits [69] and tonotopic tuning of Kv3.1 expression in the auditory brainstem is dependent on neural activity [9]. If tuning is a readout of stored memory traces, then it should be possible to show that tuning occurs during learning, that experimentally inducing tuning mimics learning, and that preventing tuning blocks learning.

Homeostasis

Homeostatic processes that return neural activity to a previous set-point following a perturbation are important for stabilizing neural circuits [57-59, 70]. For example, increasing or reducing the activity of cultured cortical neurons causes cell-wide scaling of synaptic strength that returns activity towards its previous level [71]. Integrative properties of cortical neurons can also be adjusted homeostatically [60, 72] (Figure 3B). In this case, perturbations that reduce neural activity cause neurons to modify their response properties so that smaller synaptic currents are sufficient to drive spike firing, whereas perturbations that increase activity have the opposite

effect (Figure 3B). Experimental observations of tuning appear to be inconsistent with purely homeostatic models. For example, homeostatic models predict a similar organization of cellular properties across multiple maps encoding the same sensory modality. In contrast, tuning of synaptic response properties can differ between sensory maps that receive similar inputs (c.f. Refs. [51, 53, 73]). Opposing organization of ion channels that control sub- and supra-threshold responses to synaptic input is also difficult to reconcile with purely homeostatic models (e.g. Ref. [32]). Nevertheless, there is currently very little direct evidence to rule out the possibility that examples of topographically organized synaptic responses reflect the outcome of homeostatic mechanisms for stabilization of neural activity.

Optimization

Optimization is the selection of a method or parameter that produces the best possible outcome with respect to some measure. For example, when fitting an equation to data, the optimal solution is the set of parameters that minimizes the difference between the data and the best-fit line. In this sense, homeostasis is a form of optimization in which the difference between actual neural activity and an ideal set point is actively minimized. More generally, optimization principles may apply to other functionally important aspects of neural circuit function. Optimization by tuning of synaptic integration could contribute to at least four distinct functions (Figure 4).

First, the idea that coding or transmission of information is optimized provides an attractive explanation for the role of tuning in sensory pathways. An important example of optimization is Barlow's efficient coding hypothesis, which states that sensory neurons recode signals so that the unused capacity for transmission of information is reduced [74]. This recoding may involve neurons tuning synaptic integration according to statistical properties of their inputs [64, 75]

(Figure 4A). In principle, this can be achieved by tuning the identity and number of ion channels that neurons express [61].

Second, optimization of synaptic responses may be important to ensure neurons can transmit signals effectively. For example, because auditory neurons transmit signals that vary rapidly in time, tonotopic tuning of synaptic integration may reflect differing requirements for accurate timing of synaptic responses required to encode sounds of different frequencies [11, 32] (Figure 4B).

Third, temporal filtering of synaptic responses can improve the signal to noise ratio for information that is of use to a circuit [74, 76] (Figure 4C). Optimal filtering would minimize unwanted components of a signal without affecting information that is useful. For example, tuning of synaptic response properties of neurons in the ELL causes neurons from each segment to extract different frequency components from sensory signals [35]. Optimal activation of sensory neurons by stimulus ensembles that correspond to only a subset of an animal's natural environment is also consistent with this possibility [76]. At a cellular level, such band-pass filtering can be implemented by expression of ion channels that both oppose changes in membrane potential and activate with relatively slow kinetics [77]. For example, slow gating of both HCN and certain voltage-gated potassium channels (e.g. KCNQ channels) enables them to attenuate low frequency components of input signals [47, 78, 79]. In the MEC, filtering by HCN1 channels causes a form of resonance in which neurons are most sensitive to inputs with frequencies between approximately 4 and 10 Hz [47]. This resonant frequency is tuned according to the resolution of spatial firing fields at the position along the dorsal-ventral axis at which the neuron is located [45].

A fourth cellular function that may be a target for optimization by tuning of synaptic integration is the detection of coincident inputs to a neuron. When two inputs to a neuron occur at similar times, their responses are summed so that they more effectively trigger action potentials and / or synaptic plasticity (Figure 4D). By setting the time window for effective summation of synaptic responses, optimization of coincidence detection will set the interval for identifying associations that are useful to the circuit. In the MEC, topographical organization of synaptic integration results in a corresponding tuning of synaptic coincidence detection [8]. In other neuron types, voltage-gated sodium channels play important roles in tuning coincidence detection [80].

A somewhat related possibility is that tuning of synaptic integration controls the induction of synaptic plasticity. For example, by influencing summation of synaptic responses, ion channels such as HCN1 and voltage-gated potassium channels (e.g. Kv4.2) can gate induction of synaptic plasticity (e.g. Refs. [78, 81]), and therefore may determine the conditions for storage of information in a circuit. While there is evidence for topographical organization of synaptic plasticity in the hippocampal circuit [82, 83], it is not yet clear if organization of synaptic integration plays any role.

Finally, tuning of synaptic responses might also reflect processes that cooperate with associative synaptic plasticity to extract high order patterns from distributed synaptic inputs [62]. In this scenario, plasticity of non-synaptic membrane properties acts in parallel with synaptic plasticity. These interactions can be used to learn efficient representations of stimuli. This may be particularly useful for cognitive processes mediated by associational areas of the cerebral cortex, although no direct experimental evidence currently exists for such a function.

What is the relationship between tuning of synaptic integration, cellular computations such as those we outline above, and computations at a systems level? Although systems level principles for optimization of coding in sensory systems are well established (e.g. Refs. [19, 20, 63, 64]), and optimization principles provide a general theoretical framework for understanding auditory coding [84, 85], relatively few studies have directly addressed the cellular mechanisms (e.g. [11, 61, 64, 80]). Indeed, we are not aware of experimental evidence, from auditory or other systems, to directly distinguish the possibility that tuning of synaptic integration is required for efficient coding or transmission of information, from the possibility that it reflects the outcome of homeostatic processes. Since homeostasis relies on responses to sensors that detect neural activity, evidence that tuning of synaptic integration is independent of neural activity would strongly support cell autonomous optimization models. On the other hand, activity-dependent tuning of synaptic integration is consistent with memory, homeostasis and optimization roles. In the latter case, activity-dependent optimization could be implemented through mechanisms analogous to homeostatic plasticity [59], except that the activity-sensor should instead be responsive to information coding or computation rather than to average neuronal activity [61]. At present it's unclear how these sensors could be implemented biologically. Identifying or ruling out the existence of sensors for activity-dependent optimization of information processing could be critical for distinguishing optimization models from homeostatic models.

Concluding remarks

Mechanisms that tune synaptic responses among populations of the same neuronal type are found in several distinct brain areas and diverse species. This suggests a common underlying principle for organizing neuronal circuits. Nevertheless, much remains unclear about both the cellular mechanisms and the functional roles of synaptic response tuning (Box 1). Further evaluation of the relationship between cellular tuning and the forms of optimization outlined

above will benefit from future experiments that uncover the dendritic distribution and temporal patterns of synaptic input that neurons receive *in vivo*, as well as methods to control the high-order statistics of these inputs. Experimental testing of optimization models will also require tools to selectively manipulate tuned membrane properties, while recording neuronal responses to well-defined, behaviorally relevant patterns of synaptic input. It will be particularly important to establish the consequences of manipulations that disrupt tuning without affecting other aspects of neuronal function. For example, to investigate the roles of tuning in the MEC it will be important to flatten or reverse the graded organization of both HCN and leak potassium conductances together. Finally, if tuning of synaptic responses is important for sensory, motor and cognitive processes, then it is likely that deficits in synaptic response tuning will manifest as clinical disorders. For example, genetic deletion of the Fragile X mental retardation protein (FMRP) in mice disrupts the tonotopic organization of the voltage-gated potassium channel Kv3.1b in the MNTB [9]. As future studies reveal more about the mechanisms that tune synaptic responses they may suggest novel approaches to understanding disorders caused by deficits in neural information processing.

Acknowledgements

We thank Mayank Dutia, Mike Ludwig, Timothy O'Leary, Arianna Rinaldi, Mark van Rossum, Sally Till and David Wyllie for comments on the manuscript. This work was supported by the Medical Research Council (M.F.N.), a Marie Curie Excellence grant (M.F.N.) and the Engineering and Physical Sciences Research Council (C O'D).

References

1. Ascoli, G.A., *et al.* (2008) Petilla terminology: nomenclature of features of GABAergic interneurons of the cerebral cortex. *Nat Rev Neurosci* 9, 557-568
2. McCormick, D.A., *et al.* (1985) Comparative electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. *J Neurophysiol* 54, 782-806
3. Sugino, K., *et al.* (2006) Molecular taxonomy of major neuronal classes in the adult mouse forebrain. *Nat Neurosci* 9, 99-107
4. Hobert, O., *et al.* (2010) The molecular and gene regulatory signature of a neuron. *Trends Neurosci*, 33:435-445.
5. Chklovskii, D.B., and Koulakov, A.A. (2004) Maps in the brain: what can we learn from them? *Annu Rev Neurosci* 27, 369-392
6. Yamada, R., *et al.* (2005) Hyperpolarization-activated cyclic nucleotide-gated cation channels regulate auditory coincidence detection in nucleus laminaris of the chick. *J Neurosci* 25, 8867-8877
7. Kuba, H., *et al.* (2005) Tonotopic specialization of auditory coincidence detection in nucleus laminaris of the chick. *J Neurosci* 25, 1924-1934
8. Garden, D.L., *et al.* (2008) Tuning of synaptic integration in the medial entorhinal cortex to the organization of grid cell firing fields. *Neuron* 60, 875-889
9. Strumbos, J.G., *et al.* (2010) Fragile X mental retardation protein is required for rapid experience-dependent regulation of the potassium channel Kv3.1b. *J Neurosci* 30, 10263-10271
10. Meyer, A.C., *et al.* (2009) Tuning of synapse number, structure and function in the cochlea. *Nature neuroscience* 12, 444-453
11. Slee, S.J., *et al.* (2010) Tonotopic tuning in a sound localization circuit. *J Neurophysiol* 103, 2857-2875

12. Nusser, Z. (2009) Variability in the subcellular distribution of ion channels increases neuronal diversity. *Trends Neurosci* 32, 267-274
13. Trussell, L.O. (1999) Synaptic mechanisms for coding timing in auditory neurons. *Annu Rev Physiol* 61, 477-496
14. Shadlen, M.N., and Newsome, W.T. (1994) Noise, neural codes and cortical organization. *Curr Opin Neurobiol* 4, 569-579
15. Douglas, R.J., and Martin, K.A. (2007) Mapping the matrix: the ways of neocortex. *Neuron* 56, 226-238
16. Albright, T.D., *et al.* (2000) Neural science: a century of progress and the mysteries that remain. *Neuron* 25 Suppl, S1-55
17. Schreiner, C.E., and Winer, J.A. (2007) Auditory cortex mapmaking: principles, projections, and plasticity. *Neuron* 56, 356-365
18. Hubel, D.H., and Wiesel, T.N. (1977) Ferrier lecture. Functional architecture of macaque monkey visual cortex. *Proc R Soc Lond B Biol Sci* 198, 1-59
19. Smith, E.C., and Lewicki, M.S. (2006) Efficient auditory coding. *Nature* 439, 978-982
20. Harper, N.S., and McAlpine, D. (2004) Optimal neural population coding of an auditory spatial cue. *Nature* 430, 682-686
21. Fukui, I., and Ohmori, H. (2004) Tonotopic gradients of membrane and synaptic properties for neurons of the chicken nucleus magnocellularis. *J Neurosci* 24, 7514-7523
22. Metzner, W. (1999) Why are there so many sensory brain maps? *Cell Mol Life Sci* 56, 1-4
23. Young, E.D. (1998) Parallel processing in the nervous system: evidence from sensory maps. *Proceedings of the National Academy of Sciences of the United States of America* 95, 933-934
24. Buonomano, D.V., and Merzenich, M.M. (1998) Cortical plasticity: from synapses to maps. *Annual review of neuroscience* 21, 149-186

25. Kandler, K., *et al.* (2009) Tonotopic reorganization of developing auditory brainstem circuits. *Nature neuroscience* 12, 711-717
26. Luo, L., and Flanagan, J.G. (2007) Development of continuous and discrete neural maps. *Neuron* 56, 284-300
27. Sharma, K., and Izpisua Belmonte, J.C. (2001) Development of the limb neuromuscular system. *Curr Opin Cell Biol* 13, 204-210
28. Hafting, T., *et al.* (2005) Microstructure of a spatial map in the entorhinal cortex. *Nature* 436, 801-806
29. Kjelstrup, K.B., *et al.* (2008) Finite scale of spatial representation in the hippocampus. *Science* 321, 140-143
30. Oertel, D. (1999) The role of timing in the brain stem auditory nuclei of vertebrates. *Annu Rev Physiol* 61, 497-519
31. Adamson, C.L., *et al.* (2002) Opposite actions of brain-derived neurotrophic factor and neurotrophin-3 on firing features and ion channel composition of murine spiral ganglion neurons. *J Neurosci* 22, 1385-1396
32. Adamson, C.L., *et al.* (2002) Firing features and potassium channel content of murine spiral ganglion neurons vary with cochlear location. *J Comp Neurol* 447, 331-350
33. Liu, Q., and Davis, R.L. (2007) Regional specification of threshold sensitivity and response time in CBA/CaJ mouse spiral ganglion neurons. *Journal of neurophysiology* 98, 2215-2222
34. Bakondi, G., *et al.* (2009) Hyperpolarization-activated, cyclic nucleotide-gated, cation non-selective channel subunit expression pattern of guinea-pig spiral ganglion cells. *Neuroscience* 158, 1469-1477
35. Mehaffey, W.H., *et al.* (2008) Intrinsic frequency tuning in ELL pyramidal cells varies across electrosensory maps. *Journal of neurophysiology* 99, 2641-2655

36. Ellis, L.D., *et al.* (2007) SK channels provide a novel mechanism for the control of frequency tuning in electrosensory neurons. *J Neurosci* 27, 9491-9502
37. Deng, Q., *et al.* (2005) A C-terminal domain directs Kv3.3 channels to dendrites. *J Neurosci* 25, 11531-11541
38. Metzner, W., and Juranek, J. (1997) A sensory brain map for each behavior? *Proceedings of the National Academy of Sciences of the United States of America* 94, 14798-14803
39. McLean, D.L., *et al.* (2007) A topographic map of recruitment in spinal cord. *Nature* 446, 72-75
40. Brun, V.H., *et al.* (2008) Progressive increase in grid scale from dorsal to ventral medial entorhinal cortex. *Hippocampus* 18, 1200-1212
41. Fyhn, M., *et al.* (2008) Grid cells in mice. *Hippocampus* 18, 1230-1238
42. Giocomo, L.M., *et al.* (2007) Temporal frequency of subthreshold oscillations scales with entorhinal grid cell field spacing. *Science* 315, 1719-1722
43. Boehlen, A., *et al.* (2010) The range of intrinsic frequencies represented by medial entorhinal cortex stellate cells extends with age. *J Neurosci* 30, 4585-4589
44. Giocomo, L.M., and Hasselmo, M.E. (2008) Time constants of h current in layer ii stellate cells differ along the dorsal to ventral axis of medial entorhinal cortex. *J Neurosci* 28, 9414-9425
45. Giocomo, L.M., and Hasselmo, M.E. (2009) Knock-out of HCN1 subunit flattens dorsal-ventral frequency gradient of medial entorhinal neurons in adult mice. *J Neurosci* 29, 7625-7630
46. Fernandez, F.R., and White, J.A. (2008) Artificial synaptic conductances reduce subthreshold oscillations and periodic firing in stellate cells of the entorhinal cortex. *J Neurosci* 28, 3790-3803

47. Nolan, M.F., *et al.* (2007) HCN1 channels control resting and active integrative properties of stellate cells from layer II of the entorhinal cortex. *J Neurosci* 27, 12440-12451
48. Diaz L.M. and Johnston D. (2009) Pharmacological dissociation of h-channels in CA1 neurons between dorsal and ventral hippocampus of adult rats. Program No. 716.14. 2009 Neuroscience Meeting Planner. Chicago, IL: Society for Neuroscience, 2009. Online.
49. Padmanabhan, K., and Urban, N.N. (2010) Intrinsic biophysical diversity decorrelates neuronal firing while increasing information content. *Nat Neurosci*, 13:1276-1282
50. Li, W., *et al.* (2001) Localization of two high-threshold potassium channel subunits in the rat central auditory system. *J Comp Neurol* 437, 196-218
51. Leao, R.N., *et al.* (2006) Topographic organization in the auditory brainstem of juvenile mice is disrupted in congenital deafness. *The Journal of physiology* 571, 563-578
52. Hassfurth, B., *et al.* (2009) Sensory deprivation regulates the development of the hyperpolarization-activated current in auditory brainstem neurons. *The European journal of neuroscience* 30, 1227-1238
53. Brew, H.M., and Forsythe, I.D. (2005) Systematic variation of potassium current amplitudes across the tonotopic axis of the rat medial nucleus of the trapezoid body. *Hear Res* 206, 116-132
54. Chang, H.Y. (2009) Anatomic demarcation of cells: genes to patterns. *Science* 326, 1206-1207
55. Zhang, W., and Linden, D.J. (2003) The other side of the engram: experience-driven changes in neuronal intrinsic excitability. *Nat Rev Neurosci* 4, 885-900
56. Narayanan, R., and Johnston, D. (2008) The ascent of channels with memory. *Neuron* 60, 735-738
57. Turrigiano, G.G. (2008) The self-tuning neuron: synaptic scaling of excitatory synapses. *Cell* 135, 422-435

58. Yu, L.M., and Goda, Y. (2009) Dendritic signalling and homeostatic adaptation. *Current opinion in neurobiology* 19, 327-335
59. Davis, G.W. (2006) Homeostatic control of neural activity: from phenomenology to molecular design. *Annu Rev Neurosci* 29, 307-323
60. Desai, N.S., *et al.* (1999) Plasticity in the intrinsic excitability of cortical pyramidal neurons. *Nat Neurosci* 2, 515-520
61. Stemmler, M., and Koch, C. (1999) How voltage-dependent conductances can adapt to maximize the information encoded by neuronal firing rate. *Nature neuroscience* 2, 521-527
62. Triesch, J. (2007) Synergies between intrinsic and synaptic plasticity mechanisms. *Neural computation* 19, 885-909
63. Barlow, H.B. (1961) Possible principles underlying the transformations of sensory messages. In *Sensory Communication* (Rosenblith, W.A., ed), 217-234, MIT Press
64. Laughlin, S. (1981) A simple coding procedure enhances a neuron's information capacity. *Z Naturforsch [C]* 36, 910-912
65. Aizenman, C.D., and Linden, D.J. (2000) Rapid, synaptically driven increases in the intrinsic excitability of cerebellar deep nuclear neurons. *Nat Neurosci* 3, 109-111
66. Nelson, A.B., *et al.* (2003) Long-lasting increases in intrinsic excitability triggered by inhibition. *Neuron* 40, 609-620
67. Poirazi, P., and Mel, B.W. (2001) Impact of active dendrites and structural plasticity on the memory capacity of neural tissue. *Neuron* 29, 779-796
68. Stepanyants, A., *et al.* (2002) Geometry and structural plasticity of synaptic connectivity. *Neuron* 34, 275-288
69. Tzounopoulos, T., and Kraus, N. (2009) Learning to encode timing: mechanisms of plasticity in the auditory brainstem. *Neuron* 62, 463-469

70. Turrigiano, G.G., and Nelson, S.B. (2004) Homeostatic plasticity in the developing nervous system. *Nat Rev Neurosci* 5, 97-107
71. Turrigiano, G.G., et al. (1998) Activity-dependent scaling of quantal amplitude in neocortical neurons. *Nature* 391, 892-896
72. O'Leary, T., et al. Homeostasis of intrinsic excitability in hippocampal neurones: dynamics and mechanism of the response to chronic depolarization. *J Physiol* 588, 157-170
73. Barnes-Davies, M., et al. (2004) Kv1 currents mediate a gradient of principal neuron excitability across the tonotopic axis in the rat lateral superior olive. *Eur J Neurosci* 19, 325-333
74. Barlow, H.B. (1961) Possible principles underlying the transformations of sensory messages. In *Sensory Communication* (Rosenblith, W.A., ed), 217-234, MIT Press
75. Dayan, P., and Abbott, L.F. (2001) *Theoretical Neuroscience*. MIT Press
76. Machens, C.K., et al. (2005) Testing the efficiency of sensory coding with optimal stimulus ensembles. *Neuron* 47, 447-456
77. Hutcheon, B., and Yarom, Y. (2000) Resonance, oscillation and the intrinsic frequency preferences of neurons. *Trends Neurosci* 23, 216-222
78. Nolan, M.F., et al. (2004) A Behavioral Role for Dendritic Integration: HCN1 Channels Constrain Spatial Memory and Plasticity at Inputs to Distal Dendrites of CA1 Pyramidal Neurons. *Cell* 119, 719-732
79. Hu, H., et al. (2002) Two forms of electrical resonance at theta frequencies, generated by M-current, h-current and persistent Na⁺ current in rat hippocampal pyramidal cells. *J Physiol* 545, 783-805
80. Kuba, H., et al. (2006) Axonal site of spike initiation enhances auditory coincidence detection. *Nature* 444, 1069-1072

81. Chen, X., *et al.* (2006) Deletion of Kv4.2 gene eliminates dendritic A-type K⁺ current and enhances induction of long-term potentiation in hippocampal CA1 pyramidal neurons. *J Neurosci* 26, 12143-12151
82. Maruki, K., *et al.* (2001) Differences in paired-pulse facilitation and long-term potentiation between dorsal and ventral CA1 regions in anesthetized rats. *Hippocampus* 11, 655-661
83. Papatheodoropoulos, C., and Kostopoulos, G. (2000) Decreased ability of rat temporal hippocampal CA1 region to produce long-term potentiation. *Neurosci Lett* 279, 177-180
84. Dean, I., *et al.* (2008) Rapid neural adaptation to sound level statistics. *J Neurosci* 28, 6430-6438
85. Dean, I., *et al.* (2005) Neural population coding of sound level adapts to stimulus statistics. *Nat Neurosci* 8, 1684-1689
86. Yu, F.H., *et al.* (2005) Overview of molecular relationships in the voltage-gated ion channel superfamily. *Pharmacol Rev* 57, 387-395
87. London, M., and Hausser, M. (2005) Dendritic computation. *Annu Rev Neurosci* 28, 503-532
88. Spruston, N. (2008) Pyramidal neurons: dendritic structure and synaptic integration. *Nat Rev Neurosci* 9, 206-221
89. Reyes, A. (2001) Influence of dendritic conductances on the input-output properties of neurons. *Annu Rev Neurosci* 24, 653-675
90. Jack, J.J.B., *et al.* (1975) *Electric current flow in excitable cells*. Clarendon Press
91. Rall, W. (1962) Theory of physiological properties of dendrites. *Ann N Y Acad Sci* 96, 1071-1092
92. Beisel, K.W., *et al.* (2005) Differential expression of KCNQ4 in inner hair cells and sensory neurons is the basis of progressive high-frequency hearing loss. *J Neurosci* 25, 9285-9293
93. Kuba, H., and Ohmori, H. (2009) Roles of axonal sodium channels in precise auditory time coding at nucleus magnocellularis of the chick. *J Physiol* 587, 87-100

94. Ellis, L.D., *et al.* (2008) Differential distribution of SK channel subtypes in the brain of the weakly electric fish *Apteronotus leptorhynchus*. *J Comp Neurol* 507, 1964-1978
95. von Hehn, C.A., *et al.* (2004) Loss of Kv3.1 tonotopicity and alterations in cAMP response element-binding protein signaling in central auditory neurons of hearing impaired mice. *J Neurosci* 24, 1936-1940

Figure 1. Scenarios for topographical organization and tuning of synaptic responses in neural circuits. Each schematic illustrates synaptic inputs from an upstream brain area (colored box) to three neurons of the same type within a downstream network. Information represented by the neurons is color coded. Membrane potential responses of each downstream neuron to activation of a synaptic input are shown to the right. **(a)** and **(b)**. A topographically organized input network sends projections to three neurons of the same type. In the example in **(a)**, ion channel expression is tuned according to the origin of the synaptic input that each neurons receives from the upstream network. This causes the waveform of synaptic responses to differ according to the upstream inputs that each neuron receives. In this schematic, the depicted ion channels correspond to potassium or chloride channels, which typically attenuate synaptic responses. Similar changes in the EPSP waveform could be achieved by removal of voltage-gated sodium channels, which typically amplify synaptic responses. Responses of the middle neuron (grey traces) are overlaid on responses of the adjacent neuron (black traces). In the example in **(b)**, the responses of the downstream neurons are not tuned. Rather each neuron has a similar number and distribution of non-synaptic ion channels and generates synaptic responses with similar response waveforms. **(c)** The tuning of synaptic responses is also possible when the upstream network does not demonstrate gross topographical organization. In this case, ion channel expression and synaptic responses of downstream neurons are tuned according to the identity of upstream neurons that they receive synaptic inputs from.

Figure 2. Topographical organization of spatial encoding and synaptic responses in the medial entorhinal cortex (MEC). **(a)** During exploration, neurons in layer II of the MEC encode spatial information. The schematic arena (left) illustrates the locations at which two example MEC neurons fire action potentials as a rodent explores. The firing fields have a grid like organization. The fields indicated in red are from a neuron in the dorsal part of the animal's MEC. Since the locations where the neuron is active are relatively small and close together, spatial information can be encoded by this neuron with relatively high resolution. Fields indicated in blue are from a neuron in the ventral MEC, which encodes spatial information with relatively low resolution. **(b)** The MEC (colored region in the sagittal brain section) is organized topographically, so that neurons that encode location at relatively high resolution are found more dorsally (red area), whereas neurons that encode location with a lower spatial resolution are found more ventrally (blue area) [28]. **(c)** Key cellular properties of neurons in layer II of the MEC follow a similar topographical organization. More dorsal neurons have a higher density of membrane currents mediated by HCN channels and leak potassium channels [8]. Importantly, these differences in ion channel function modify the way neurons respond to synaptic input. **(d)** Following activation of synaptic inputs (arrows), there is greater summation of EPSPs in ventral compared to dorsal neurons, whereas the underlying synaptic currents do not differ [8]. Images modified, with permission, from Ref. [8].

Figure 3. Models of memory storage and homeostasis mediated by modification of synaptic response properties. **(a)** An example of how information can be stored by modifying synaptic integration. Before learning (left), activation of a synaptic input elicits a single action potential. During learning, the neuron reduces its expression of ion channels that usually attenuate synaptic responses. Subsequent activation of the same synaptic input elicits more action potentials. **(b)** An example of homeostatic control implemented by a change in synaptic integration. When the input to a neuron is reduced, the output from the neuron is also initially reduced (centre). To return its output to a homeostatic set-point, the neuron removes ion channels from its membrane that normally attenuate synaptic responses (right). As a result, the output from the neuron returns to its previous level.

Figure 4. Models for the optimization of neuronal function by tuning synaptic integration. A number of cellular functions may be targets for optimization by tuning synaptic integration. **(a)** Efficient coding minimizes redundancy in a neuron's output. This requires optimal tuning of the relationship between the input to a neuron and the output that it produces. In this example, a neuron receives 1000 inputs. The probability of any particular number of inputs being active at a given time is plotted to the left. How the neuron transforms its synaptic input into spike output is described by its 'input-output' relationship (centre). This shows the frequency of action potentials that it generates in response to a particular number of active inputs. An optimal input-output relationship (black line) will enable the neuron to make equal use of all of the available output frequencies (right) [64]. The shape of the optimal input-output relationship depends on the distribution of active inputs and therefore must be tuned in some way. Input-output relationships that are too shallow (blue line) or too steep (red line) result in some output frequencies being under-utilized (right). A neuron can control the shape of its input relationship by adjusting the number or properties of ion channels that influence its response to synaptic inputs [61]. **(b)** The timing of a neuron's response to synaptic input may also be a target for tuning of synaptic integration. The delay from activation of a synaptic input to initiation of an action potential is controlled by post-synaptic ion channels and can differ between trials because of noise. In this example, the delay before action potential initiation and the variable timing of action potential responses to simulated synaptic inputs are both reduced simply by increasing the density of ion channels that mediate action potential firing (purple traces). **(c)** Temporal filtering is another cellular function that may be a target for optimization by tuning synaptic integration. The input signals that neurons receive fluctuate continuously. The amplitude of a neuron's response depends on the rate at which the input fluctuates and the ion channels that it expresses [77]. This example shows membrane potential responses (upper traces) of a neuron to input currents that fluctuate at a low (lower left trace) or high frequency (lower right trace). Removal from the neuron

of HCN channels, which gate slowly and tend to oppose slow changes in membrane potential, causes the amplitude of responses to low frequency inputs to be increased, but has little effect on the amplitude of responses to higher frequency inputs. **(d)** Neurons can act as coincidence detectors that preferentially respond to inputs that are active close together in time. The exact time window for coincidence detection can be set by adjusting the density of ion channels in the neuron's membrane. In this example, a neuron with a low density of ion channels effectively sums responses to two inputs that are 5 ms apart (green traces). This neuron has a relatively broad time-window for coincidence detection compared to a neuron with a higher density of ion channels, which does not sum responses that are 5 ms apart (black traces).

Table 1. Topographic evidence for tuning of synaptic responses. Ion channels for which there is evidence of a topographical organization include voltage-gated potassium (Kv) channels, calcium-activated potassium (KCa) channels and hyperpolarization-activated cyclic nucleotide-gated (HCN) channels. Abbreviations for channel names are based on the International Union of Pharmacology guidance [86].

Brain Area	Feature	Evidence	Species/ Refs.	
Sensory	Cochlea spiral ganglion	Sound frequency	Action potential duration decreases and adaptation increases with sound frequency	Mouse [31-33]
			KCa1.1 (slo), Kv1.1, Kv1.2, Kv3.1 and Kv7.3 immunohistochemical labeling increases with sound frequency	Mouse [31, 32, 92]
			Membrane potential sag response to injected current decreases with sound frequency	Mouse [32, 33] and guinea pig [34]
			Kv4.2, HCN1 and HCN3 immunohistochemical labeling decreases with sound frequency	Mouse [32] and guinea pig [34]
			EPSP duration and time window for synaptic coincidence detection is shortest for intermediate sound frequencies	Chick [6, 7]
Nucleus laminaris	Sound frequency	Membrane input resistance and time constant is smallest for intermediate sound frequencies	Chick [6, 7]	
		Kv1.2 immunohistochemical labeling is highest for intermediate frequencies	Chick [7]	
		HCN currents and HCN1 (but not HCN2) mRNA and immunohistochemical labeling decreases with sound frequency	Chick [6]	
		Na ⁺ channel localization and axonal site of action potential initiation is further from the soma with higher sound frequencies.	Chick [80]	
		Gain of band pass filtered responses decreases with sound frequency. This leads to improved detection of interaural time differences.	Chick [11]	
Nucleus	Sound	Kv1.1 mRNA and protein increases with sound frequency.	Chick [21]	

magnocellularis	frequency	Na ⁺ channels localization to the axon initial segment decreases and spike threshold increases with sound frequency.	Chick [93]
Medial nucleus of the trapezoid body	Sound frequency	Labeling Kv1.1, Kv3.1 and Kv3.3, voltage-gated potassium current amplitude and action potential frequency adaptation increases with sound frequency	Rat and mouse [9, 50, 51, 53, 95]
		Amplitude of HCN currents and HCN4 immunohistochemical labeling decreases with sound frequency	Mouse [51], but not rat [53] or gerbil [52]
Lateral superior olive	Sound frequency	Kv1.1 immunohistochemistry, spike frequency adaptation and voltage-gated potassium currents decreases with sound frequency	Rat [73]
		Voltage-dependence of HCN currents is shifted to more negative potentials with increasing frequency	Gerbil [52]
Electrosensory lateral line lobe E-cells	Behavioral response	Action potential waveform, pattern and input frequency selectivity is tuned to specific sensory maps	Electric fish [35]
		Kv3.1 mRNA and immunohistochemical labeling, SK1 and SK2 mRNA and immunohistochemical labeling is tuned to specific sensory maps	Electric fish [36, 37, 94]
Motor			
Spinal motor neurons and interneurons	Swimming speed	Membrane input resistance decreases with swimming frequency	Zebrafish [39]
Cognitive			
Medial entorhinal cortex layer II stellate neurons	Resolution of spatial firing fields	EPSP duration, summation and time window for synaptic coincidence detection decreases with swimming frequency	Mouse [8]
		Frequency of intrinsic oscillatory and resonance properties increase with increasing spatial resolution	Rat [42-44] and mouse [45]
		Membrane input resistance and time constant decrease with increasing spatial resolution	Mouse [8] and rat [43] (but see [44]).
		HCN and leak K ⁺ current amplitude increase with increasing spatial resolution	Mouse [8], but not HCN in rat [44]
		HCN current activation time constant increases with increasing spatial resolution	Rat [44], but not mouse [8]

Glossary

Distributed representation

Representation of information that requires a population of neurons. For example, because speech or other natural sounds usually contain many different frequencies of sound waves, and as individual auditory neurons respond preferentially to a single characteristic frequency, the neural representation of natural sounds is distributed across many neurons with different characteristic frequencies. A distributed computation involves simultaneous processing of information encoded within a distributed neural representation.

HCN channels

Cation permeable ion channels that are opened by hyperpolarization of the membrane potential. These channels are unlike most other voltage-gated ion channels that are opened by depolarization.

Leak potassium currents

Currents mediated by potassium channels that are open at rest and typically have conductance that is only weakly dependent on membrane potential.

Resonance

In general this is the preferential response of a system to inputs with a particular frequency - the resonant frequency of the system. In neurons, resonance often manifests as a larger membrane potential change, or a greater number of spikes, in response to inputs that oscillate at a particular range of frequencies, compared with responses to inputs that oscillate at higher or lower frequencies [77].

Sub-threshold responses

Synaptic potentials that are below the threshold for initiation of action potentials. Integration of these responses is typically determined by neuronal morphology and by ion channels that are open at membrane potentials negative to the threshold for action potential firing.

Supra-threshold responses

When synaptic activation causes sufficient depolarization of the membrane potential to initiate action potentials. The number, frequency and time-course of the resulting action potentials are determined by ion channels activated during the action potential and its after-hyperpolarization.

Synaptic integration

In most neurons, activation of a single synapse is insufficient to trigger firing of an action potential. Instead, neurons are constantly integrating excitatory and inhibitory post synaptic potentials. If excitation sufficiently outweighs inhibition then an action potential is triggered. This process of synaptic integration depends not only on the current passing through ligand-gated ion channels, but also on neuronal morphology, the location of active synapses, and the complement and sub-cellular distribution of non-synaptic ion channels that amplify or attenuate synaptic potentials [12, 87-91].

Tonotopic organization

When the frequency of encoded sound can be systematically 'mapped' onto neurons at anatomically defined locations.

Topographic organization

When an encoded variable can be systematically 'mapped' onto neurons at anatomically defined locations so that nearby neurons respond to similar stimuli.


Tuning of synaptic responses

When synaptic responses are configured according to the functional roles of a particular neuron, for example, by setting the density of non-synaptic ion channels that influence the integration of synaptic responses.

Box 1. Outstanding questions

- What are the behavioral functional outcomes that result from the tuning of synaptic integration?
- How can storage, homeostasis and optimization models be distinguished experimentally?
- Is tuning hard-wired or dependent on activity?
- Do the multiple examples of synaptic response tuning reflect a single molecular mechanism, or are they implemented through several distinct mechanisms?
- Does tuning extend to other populations of neurons that encode distributed representations, for example in the visual cortex?
- Does tuning have a common function in different brain networks, or is it tailored to the specific tasks a circuit performs?
- Do deficits in synaptic response tuning manifest as neurological disorders?


Figure 1
[Click here to download high resolution image](#)



Original Figure 1 File

[Click here to download Original Figure File: NeuronTopography.svg](#)


Figure 2
[Click here to download high resolution image](#)



Original Figure 2 File

[Click here to download Original Figure File: GridFigNewMouse.svg](#)


Figure 3
[Click here to download high resolution image](#)



Original Figure 3 File

[Click here to download Original Figure File: Storage.svg](#)

Figure 4
[Click here to download high resolution image](#)



Original Figure 4 File

[Click here to download Original Figure File: TINS10v3.pxp](#)