THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Bridged beta(3)-Peptide Inhibitors of p53-hDM2 Complexation: Correlation between Affinity and Cell Permeability

Citation for published version:

Bautista, AD, Appelbaum, JS, Craig, CJ, Michel, J & Schepartz, A 2010, 'Bridged beta(3)-Peptide Inhibitors of p53-hDM2 Complexation: Correlation between Affinity and Cell Permeability', *Journal of the American Chemical Society*, vol. 132, no. 9, pp. 2904-+. <https://doi.org/10.1021/ja910715u>

Digital Object Identifier (DOI):

[10.1021/ja910715u](https://doi.org/10.1021/ja910715u)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of the American Chemical Society

Publisher Rights Statement:

Copyright © 2010 by the American Chemical Society. All rights reserved.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Published in final edited form as:

J Am Chem Soc. 2010 March 10; 132(9): 2904–2906. doi:10.1021/ja910715u.

Bridged β^3 -Peptide Inhibitors of p53-hDM2 Complexation— Correlation Between Affinity and Cell Permeability

Arjel D. Bautista[#], Jacob S. Appelbaum[§], Cody J. Craig[#], Julien Michel[#], and Alanna Schepartz^{#,¶}

Alanna Schepartz: alanna.schepartz@yale.edu

[#]Department of Chemistry, Yale University, New Haven, Connecticut 06520-8107

[§]Department of Cell Biology, and Molecular, Yale University, New Haven, Connecticut 06520-8107

[¶]Department of Cellular and Developmental Biology, Yale University, New Haven, Connecticut 06520-8107

Abstract

β -peptides possess several features that are desirable in peptidomimetics; they are easily synthesized, fold into stable secondary structures in physiologic buffers, and resist proteolysis. They can also bind to a diverse array of proteins to inhibit their interactions with α -helical ligands. β -peptides are not usually cell permeable, however, and this feature limits their utility as research tools and potential therapeutics. Appending an Arg₈ sequence to a β -peptide improves uptake but adds considerable mass. We reported that embedding a small cationic patch within a PPII, α - or β -peptide helix improves uptake without the addition of significant mass. In another mass-neutral strategy, Verdine, Walensky, and others have reported that insertion of a hydrocarbon bridge between the *i* and *i*+4 positions of an α -helix also increases cell uptake. Here we describe a series of β -peptides containing diether and hydrocarbon bridges and compare them on the basis of cell uptake and localization, affinities for hDM2, and 14-helix structure. Our results highlight the relative merits of cationic patch and hydrophobic bridge strategies for improving β -peptide uptake and identify a surprising correlation between uptake efficiency and hDM2 affinity.

β -peptides^{1–4} possess several features that are desirable in peptidomimetics;^{5,6} they are easily synthesized, fold into helices^{1–3,7} in physiologic buffers,⁸ and resist proteolysis.⁹ They also bind *in vitro* to proteins such as hDM2,^{10–14} hDMX,¹⁰ gp41,^{15,16} and others,^{17–19} and inhibit their interactions with α -helical ligands. β -peptides are not usually cell permeable, however, and this feature limits their utility as research tools and potential therapeutics. Appending an Arg₈ sequence to a β -peptide can improve uptake^{20,21} but adds considerable mass. We reported that embedding a small cationic patch within a PPII,²² α -²³ or β -peptide¹¹ helix improves uptake without the addition of significant mass.^{24,25} Similarly, Verdine, Walensky, and others^{26–33} reported that insertion of a hydrocarbon bridge (a “staple”) between the *i* and *i*+4 positions of an α -helix³⁴ increases uptake.^{26,29,32,34–38} Here we describe a variety of β -peptides containing diether- and hydrocarbon bridges and compare them on the basis of cell uptake and localization, affinity for hDM2, and 14-helix structure. Our results highlight the relative merits of cationic patch and hydrophobic bridge strategies for improving β -peptide uptake and identify an unprecedented correlation between uptake efficiency and hDM2 affinity *in vitro*.

Correspondence to: Alanna Schepartz, alanna.schepartz@yale.edu.

Supporting Information **Available:** β -peptide synthesis, binding and cell uptake assays, and confocal microscopy images. This material is available free of charge on the Internet at <http://pubs.acs.org>.

Our studies began with an analysis of available x-ray^{39,40} and NMR structures^{13,41} of β -peptide 14-helices to identify those position pairs that would best tolerate an ether^{42,43} or hydrocarbon³⁴ bridge. This analysis, supported by recent work of Perlmutter⁴² and Seebach⁴⁴ suggested that a 21-atom bridge could be accommodated between most *i* and *i*+3 positions of a 14-helix. To test this prediction, we synthesized an analog of β -peptide **2**⁷ containing (O-allyl)- β^3 -L-Ser at positions 3 and 6 (**2(3-6)**, Figure 1), and subjected it to on-resin ring-closing metathesis using bis(tricyclohexylphosphine)benzylidene ruthenium (IV) dichloride³⁴ to generate **2(3-6)s**.⁴⁵ The circular dichroism (CD) spectra of **2**, **2(3-6)** and **2(3-6)s** were identical (Figure S1), indicating that this 21-atom diether bridge is accommodated between positions 3 and 6. Introduction of the diether bridge did not significantly increase or decrease the extent of 14-helix structure as judged by CD.

In order to evaluate the relative uptake of bridged β -peptides in the context of a functional molecule of diverse sequence, we synthesized a series of variants of β **53-12**,¹⁰ an inhibitor of p53-hDM2 complexation (Figure 1). These variants contained either (O-allyl)- β^3 -L-Ser (to generate a diether bridge) or (*S*)-3-aminooct-7-enoic acid (to generate a hydrocarbon bridge) at *i* and *i*+3 positions 2 and 5 (**25.O-s** and **25.C-s**, respectively) or 4 and 7 (**47.O-s** and **47.C-s**, respectively). According to the CD spectra (Figure 2), all bridged β -peptides assumed a 14-helical structure and were modestly more helical than unbridged analogs (Figure S2).

As a prelude to evaluating cell uptake and localization, we employed a direct fluorescence polarization assay to compare hydrocarbon and diether bridged β -peptides on the basis of affinity for hDM2₁₋₁₈₈ (Figure 2B). β -peptides containing a diether or hydrocarbon bridge between positions 4 and 7 bound hDM2₁₋₁₈₈ 2-fold better ($K_d = 53.9 \pm 22.7$ and 94.1 ± 18.4 nM, respectively) than the corresponding unbridged analogs ($K_d = 114 \pm 28$ and 253 ± 75 nM, respectively), in line with analogous comparisons in an α -peptide context.³⁵ By contrast, β -peptides containing a diether or hydrocarbon bridge between positions 2 and 5 bound hDM2₁₋₁₈₈ between 4 and 8-fold worse ($K_d = 548 \pm 58$ and 546 ± 96 nM, respectively) than unbridged analogs ($K_d = 139 \pm 13$ and 68.1 ± 7.8 nM, respectively). *In silico* analysis suggests that the lower hDM2₁₋₁₈₈ affinity of β -peptides **25.C-s** and **25.O-s** results from steric hindrance between the hydrocarbon bridge and the hDM2 surface that is absent in the complex with peptides **47.C-s** and **47.O-s** (Figure 3, compare A and B).

We next set out to monitor the mammalian cell uptake and sub-cellular localization of diether- and hydrocarbon bridged β -peptides based on β **53-12**. Uptake was monitored using flow cytometry, whereas sub-cellular localization was assessed using confocal microscopy (Figure 4). β -peptides containing diether or hydrocarbon bridges between positions 4 and 7 were taken up significantly more efficiently (MCF = 8.21 ± 0.45 and 8.63 ± 0.77 , respectively) than unbridged analogs (MCF = 3.23 ± 0.31 and 2.63 ± 0.32 , respectively), irrespective of bridge structure. By contrast, β -peptides containing diether or hydrocarbon bridges between positions 2 and 5 were taken up poorly, irrespective of bridge structure, and behaved much like the unbridged analogs. In all cases, as judged by flow cytometry, the greatest uptake was observed with β -peptide β **53-12SB3**, which contains a cationic patch on one 14-helix face but no bridge of any kind (Figure 4AB).

The localization of bridged β -peptides upon cell uptake was explored in more detail using confocal microscopy. HeLa cells were treated with fluorescently labeled β -peptide (green) as well as Alexa Fluor® 647 labeled transferrin and Hoechst 33342 to visualize recycling endosomes^{46,47} (red) and nuclei (blue). β -peptides containing a diether or hydrocarbon bridge between positions 4 and 7 are distributed widely among Tf+ and Tf- endosomes, as well as nuclear and cytosolic compartments, whereas those containing the analogous bridge between positions 2 and 5 are not (Figure 3). Indeed, β -peptides containing a diether or hydrocarbon bridge between positions 2 and 5 are taken up more poorly than the unbridged analog (Figure

S4). These results highlight an intriguing correlation between hDM2 affinity and cell uptake; it is possible that the structural features that lower hDM2 affinity (Figure S3) also lower uptake efficiency. Indeed, it appears that for these β -peptides, an increase in 14-helix secondary structure does not necessarily confer increased cell uptake.²⁶

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the NIH (GM 74756), the National Foundation for Cancer Research, and a Marie Curie International Outgoing Fellowship within the 7th European Community Framework Programme (J.M.). A.D.B. is grateful to Bristol-Myers Squibb for a graduate research fellowship.

References

1. Cheng RP, Gellman SH, DeGrado WF. *Chem Rev* 2001;101:3219–3232. [PubMed: 11710070]
2. DeGrado WF, Schneider JP, Hamuro Y. *J Pept Res* 1999;54:206–217. [PubMed: 10517158]
3. Gellman SH. *Acc Chem Res* 1998;31:173–180.
4. Seebach D, Overhand M, Kunhle FNM, Martinoni B, Oberer L, Hommel U, Widmer H. *Helv Chim Acta* 1996;79:913–941.
5. Bautista AD, Craig CJ, Harker EA, Schepartz A. *Curr Opin Chem Biol* 2007;11:685–592. [PubMed: 17988934]
6. Kritzer JA, Stephens OM, Guarracino DA, Reznik SK, Schepartz A. *Bioorg Med Chem* 2004;13:11–16. [PubMed: 15582447]
7. Kritzer JA, Tirado-Rives J, Hart SA, Lear JD, Jorgensen WL, Schepartz A. *J Am Chem Soc* 2005;127:167–178. [PubMed: 15631466]
8. Hart SA, Bahadoor ABF, Matthews EE, Qiu XJ, Schepartz A. *J Am Chem Soc* 2003;125:4022–4023. [PubMed: 12670203]
9. Frackenpohl J, Arvidsson PI, Schreiber JV, Seebach D. *ChemBioChem* 2001;2:445–455. [PubMed: 11828476]
10. Harker EA, Daniels DS, Guarracino DA, Schepartz A. *Bioorg Med Chem* 2009;17:2038–2046. [PubMed: 19211253]
11. Harker EA, Schepartz A. *ChemBioChem* 2009;10:990–993. [PubMed: 19266537]
12. Kritzer JA, Lear JD, Hodsdon ME, Schepartz A. *J Am Chem Soc* 2004;126:9468–9469. [PubMed: 15291512]
13. Kritzer JA, Luedtke NW, Harker EA, Schepartz A. *J Am Chem Soc* 2005;127:14584–14585. [PubMed: 16231906]
14. Murray JK, Gellman SH. *Pept Sci* 2007;88:657–686.
15. Bautista AD, Stephens OM, Wang L, Damaoal RA, Anderson KS, Schepartz A. *Bioorg Med Chem Lett* 2009;19:3736–3738. [PubMed: 19497744]
16. Stephens OM, Kim S, Welch BD, Hodsdon ME, Kay MS, Schepartz A. *J Am Chem Soc* 2005;127:13126–13127. [PubMed: 16173723]
17. English EP, Chumanov RS, Gellman SH, Compton T. *J Biol Chem* 2006;281:2661–2667. [PubMed: 16275647]
18. Lee EF, Sadowsky JD, Smith BJ, Czabotar PE, Peterson-Kaufman KJ, Colman PM, Gellman SH, Fairlie WD. *Angew Chem Int Ed* 2009;48:4318–4322.
19. Sadowsky JD, Fairlie WD, Hadley EB, Lee HS, Umezawa N, Nikolovska-Coleska Z, Wang S, Huang DCS, Tomita Y, Gellman SH. *J Am Chem Soc* 2007;129:139–154. [PubMed: 17199293]
20. Jones SW, Christison R, Bundell K, Joyce CJ, Brockbank SMV, Newham P, Lindsay MA. *Br J Pharmacol* 2005;145:1093–1102. [PubMed: 15937518]
21. Tung CH, Weissleder R. *Adv Drug Deliv Rev* 2003;55:281–294. [PubMed: 12564981]

22. Daniels DS, Schepartz A. *J Am Chem Soc* 2007;129:14578–14579. [PubMed: 17983240]
23. Smith BA, Daniels DS, Coplin AE, Jordan GE, McGregor LM, Schepartz AJ. *Am Chem Soc* 2008;130:2948–2949.
24. Lawrence MS, Phillips KJ, Liu DR. *J Am Chem Soc* 2007;129:10110–10112. [PubMed: 17665911]
25. McNaughton BR, Cronican JJ, Thompson DB, Liu DR. *Proc Natl Acad Sci U S A* 2009;106:6111–6116. [PubMed: 19307578]
26. Kim YW, Verdine GL. *Bioorg Med Chem Lett* 2009;19:2533–2536. [PubMed: 19332370]
27. Kutchukian PS, Yang JS, Verdine GL, Shakhnovich EI. *J Am Chem Soc* 2009;131:4622–4627. [PubMed: 19334772]
28. Madden MM, Vera CIR, Song W, Lin Q. *Chem Commun* 2009:5588–5590.
29. Moellering RE, Cornejo M, Davis TN, Bianco CD, Aster JC, Blacklow SC, Kung AL, Gilliland DG, Verdine GL, Bradner JE. *Nature* 2009;462:182–188. [PubMed: 19907488]
30. Whelan J. *Drug Discov Today* 2004;9:907–907. [PubMed: 15501720]
31. Bhattacharya S, Zhang H, Debnath AK, Cowburn D. *J Biol Chem* 2008;283:16274–16278. [PubMed: 18417468]
32. Danial NN, et al. *Nat Med* 2008;14:144–153. [PubMed: 18223655]
33. Henchey LK, Jochim AL, Arora PS. *Curr Opin Chem Biol* 2008;12:692–697. [PubMed: 18793750]
34. Schafmeister CE, Po J, Verdine GL. *J Am Chem Soc* 2000;122:5891–5892.
35. Bernal F, Tyler AF, Korsmeyer SJ, Walensky LD, Verdine GL. *J Am Chem Soc* 2007;129:2456–2457. [PubMed: 17284038]
36. Walensky LD, Kung AL, Escher I, Malia TJ, Barbuto S, Wright RD, Wagner G, Verdine GL, Korsmeyer SJ. *Science* 2004;305:1466–1470. [PubMed: 15353804]
37. Walensky LD, Pitter K, Morash J, Oh KJ, Barbuto S, Fisher J, Smith E, Verdine GL, Korsmeyer SJ. *Mol Cell* 2006;24:199–210. [PubMed: 17052454]
38. Zhang H, Zhao Q, Bhattacharya S, Waheed AA, Tong X, Hong A, Heck S, Curreli F, Goger M, Cowburn D, Freed EO, Debnath AK. *J Mol Biol* 2008;378:565–580. [PubMed: 18374356]
39. Daniels DS, Petersson EJ, Qiu JX, Schepartz A. *J Am Chem Soc* 2007;129:1532–1533. [PubMed: 17283998]
40. Goodman JL, Petersson EJ, Daniels DS, Qiu JX, Schepartz A. *J Am Chem Soc* 2007;129:14746–14751. [PubMed: 17985897]
41. Kritzer JA, Hodsdon ME, Schepartz A. *J Am Chem Soc* 2005;127:4118–4119. [PubMed: 15783163]
42. Bergman YE, Del Borgo MP, Gopalan RD, Jalal S, Unabia SE, Ciampini M, Clayton DJ, Fletcher JM, Mulder RJ, Wilce JA, Aguilar MI, Perlmutter P. *Org Lett* 2009;11:4438–4440. [PubMed: 19719157]
43. Blackwell HE, Grubbs RH. *Angew Chem Int Ed* 1998;37:3281–3284.
44. Ebert MO, Gardiner J, Ballet S, Abell AD, Seebach D. *Helv Chim Acta* 2009;2643–2658.
45. See Supporting Information for details.
46. Ghosh R, Gelman D, Maxfield F. *J Cell Sci* 1994;107:2177–2189. [PubMed: 7983176]
47. Hopkins C, Gibson A, Shipman M, Strickland D, Trowbridge I. *J Cell Biol* 1994;125:1265–1274. [PubMed: 7515888]

Figure 1.

Helical net representation of β -peptides studied herein. β^3 -homoamino acids are identified by the single-letter code used for the corresponding α -amino acid. Orn represents ornithine. Z represents 3-(S)-3-amino-4-(2-trifluoromethylphenyl)-butyric acid.

Figure 2. CD analysis of β -peptides containing hydrocarbon or diether bridges between residues (A) 2 and 5 or (B) 4 and 7. Fluorescence polarization (FP) analysis of hDM2 binding by β -peptides containing (C) hydrocarbon or (D) diether bridges.

Figure 3.
Computational model of hDM2 (grey) in complex with (A) **25.C-s** or (B) **47.C-s**.⁴⁵

Figure 4.

HeLa cell uptake and localization of Flu-labeled β -peptides. (A,B) HeLa cells were incubated with 2 μ M β -peptide for 4 h, treated with 0.25% trypsin for 10 min, washed with cold DMEM and PBS, and analyzed using flow cytometry. (C) Confocal microscopy of HeLa cells treated with 20 μ M of the indicated β -peptide (green), 5 $\text{mg}\cdot\text{mL}^{-1}$ Alexa Fluor 647-transferrin (red) and 150 nM Hoescht 33342 (blue).