



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Somatic evolution and global expansion of an ancient transmissible cancer lineage

Citation for published version:

Baez-Ortega, A, Gori, K, Strakova, A, Allen, JL, Allum, KM, Bansse-Issa, L, Bhutia, TN, Bisson, JL, Briceño, C, Castillo Domracheva, A, Corrigan, AM, Cran, HR, Crawford, JT, Davis, E, de Castro, KF, B de Nardi, A, de Vos, AP, Delgadillo Keenan, L, Donelan, EM, Espinoza Huerta, AR, Faramade, IA, Fazil, M, Fotopoulou, E, Fruean, SN, Gallardo-Arrieta, F, Glebova, O, Gouletsou, PG, Häfelin Manrique, RF, Henriques, JJGP, Horta, RS, Ignatenko, N, Kane, Y, King, C, Koenig, D, Krupa, A, Kruzeniski, SJ, Kwon, Y-M, Lanza-Perea, M, Lazyan, M, Lopez Quintana, AM, Losfelt, T, Marino, G, Martínez Castañeda, S, Martínez-López, MF, Meyer, M, Migneco, EJ, Nakanwagi, B, Neal, KB, Neunzig, W, Ní Leathlobhair, M, Nixon, SJ, Ortega-Pacheco, A, Pedraza-Ordoñez, F, Peleteiro, MC, Polak, K, Pye, RJ, Reece, JF, Rojas Gutierrez, J, Sadia, H, Schmeling, SK, Shamanova, O, Sherlock, AG, Stammnitz, M, Steenland-Smit, AE, Svitich, A, Tapia Martínez, LJ, Thoya Ngoka, I, Torres, CG, Tudor, EM, van der Wel, MG, Vițălaru, BA, Vural, SA, Walkinton, O, Wang, J, Wehrle-Martinez, AS, Widdowson, SAE, Stratton, MR, Alexandrov, LB, Martincorena, I & Murchison, EP 2019, 'Somatic evolution and global expansion of an ancient transmissible cancer lineage', *Science*, vol. 365, no. 6452. <https://doi.org/10.1126/science.aau9923>

Digital Object Identifier (DOI):

[10.1126/science.aau9923](https://doi.org/10.1126/science.aau9923)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Science

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Somatic evolution and global expansion of an ancient transmissible cancer lineage

Adrian Baez-Ortega¹, Kevin Gori^{1*}, Andrea Strakova^{1*}, Janice L. Allen², Karen M. Allum³, Leontine Banske-Issa⁴, Thinlay N. Bhutia⁵, Jocelyn L. Bisson¹, Cristóbal Briceño⁶, Artemio Castillo Domracheva⁷, Anne M. Corrigan⁸, Hugh R. Cran⁹, Jane T. Crawford¹⁰, Eric Davis¹¹, Karina F. de Castro¹², Andriago B. de Nardi¹³, Anna P. de Vos¹⁴, Laura Delgadillo Keenan¹⁵, Edward M. Donelan², Adela R. Espinoza Huerta¹⁶, Ibikunle A. Faramade¹⁷, Mohammed Fazil¹⁸, Eleni Fotopoulou¹⁹, Skye N. Fruean²⁰, Fanny Gallardo-Arrieta²¹, Olga Glebova²², Pagona G. Gouletsou²³, Rodrigo F. Häfelin Manrique²⁴, Joaquim J. G. P. Henriques²⁵, Rodrigo S. Horta²⁶, Natalia Ignatenko²⁷, Yaghouba Kane²⁸, Cathy King³, Debbie Koenig³, Ada Krupa²⁹, Steven J. Kruzeniski¹⁶, Young-Mi Kwon¹, Marta Lanza-Perea⁸, Mihran Lazyan³⁰, Adriana M. Lopez Quintana³¹, Thibault Losfelt³², Gabriele Marino³³, Simón Martínez Castañeda³⁴, Mayra F. Martínez-López³⁵, Michael Meyer³⁶, Edward J. Migneco³⁷, Berna Nakanwagi³⁸, Karter B. Neal³⁹, Winifred Neunzig³, Máire Ní Leathlobhair¹, Sally J. Nixon⁴⁰, Antonio Ortega-Pacheco⁴¹, Francisco Pedraza-Ordoñez⁴², Maria C. Peleteiro⁴³, Katherine Polak⁴⁴, Ruth J. Pye⁴⁵, John F. Reece⁴⁶, Jose Rojas Gutierrez⁴⁷, Haleema Sadia⁴⁸, Sheila K. Schmeling⁴⁹, Olga Shamanova⁵⁰, Alan G. Sherlock⁴⁵, Maximilian Stammnitz¹, Audrey E. Steenland-Smit⁴, Alla Svitch⁵¹, Lester J. Tapia Martínez¹⁶, Ismail Thoya Ngoka⁵², Cristian G. Torres⁵³, Elizabeth M. Tudor⁵⁴, Mirjam G. van der Wel⁵⁵, Bogdan A. Vițălaru⁵⁶, Sevil A. Vural⁵⁷, Oliver Walkinton⁴⁵, Jinhong Wang¹, Alvaro S. Wehrle-Martinez⁵⁸, Sophie A. E. Widdowson⁵⁹, Michael R. Stratton⁶⁰, Ludmil B. Alexandrov⁶¹, Iñigo Martincorena⁶⁰, Elizabeth P. Murchison^{1§}

¹ Transmissible Cancer Group, Department of Veterinary Medicine, University of Cambridge, Cambridge, UK

² Animal Management in Rural and Remote Indigenous Communities (AMRRIC), Darwin, Australia

³ World Vets, Gig Harbor, USA

⁴ Animal Shelter, Stichting Dierenbescherming Suriname, Paramaribo, Suriname

⁵ Sikkim Anti-Rabies and Animal Health Programme, Department of Animal Husbandry, Livestock, Fisheries and Veterinary Services, Government of Sikkim, India

⁶ ConserLab, Animal Preventive Medicine Department, Faculty of Animal and Veterinary Sciences, University of Chile, Santiago, Chile

- 37 ⁷ Corozal Veterinary Hospital, University of Panamá, Panama City, Republic of Panama
38 ⁸ St. George's University, True Blue, Grenada
39 ⁹ The Nakuru District Veterinary Scheme Ltd, Nakuru, Kenya
40 ¹⁰ Animal Medical Centre, Belize City, Belize
41 ¹¹ International Animal Welfare Training Institute, UC Davis School of Veterinary Medicine,
42 Davis, USA
43 ¹² Centro Universitário de Rio Preto (UNIRP), São José do Rio Preto, São Paulo, Brazil
44 ¹³ Department of Clinical and Veterinary Surgery, São Paulo State University (UNESP), São
45 Paulo, Brazil
46 ¹⁴ Ladybrand Animal Clinic, Ladybrand, South Africa
47 ¹⁵ Veterinary Clinic Sr. Dog's, Guadalajara, Mexico
48 ¹⁶ World Vets Latin America Veterinary Training Center, Granada, Nicaragua
49 ¹⁷ National Veterinary Research Institute, Vom, Nigeria
50 ¹⁸ Animal Clinic, Mombasa, Kenya
51 ¹⁹ Intermunicipal Stray Animals Care Centre (DIKEPAZ), Perama, Greece
52 ²⁰ Animal Protection Society of Samoa, Apia, Samoa
53 ²¹ Faculty of Veterinary Science, University of Zulia, Maracaibo, Venezuela
54 ²² Veterinary Clinic BIOCONTROL, Moscow, Russia
55 ²³ Faculty of Veterinary Medicine, School of Health Sciences, University of Thessaly,
56 Karditsa, Greece
57 ²⁴ Veterinary Clinic El Roble, Animal Healthcare Network, Faculty of Animal and Veterinary
58 Sciences, University of Chile, Santiago de Chile, Chile
59 ²⁵ OnevetGroup, Hospital Veterinário Berna, Lisboa, Portugal
60 ²⁶ Universidade Vila Velha, Vila Velha, Brazil
61 ²⁷ Veterinary Clinic Zoovetservis, Kiev, Ukraine
62 ²⁸ École Inter-états des Sciences et Médecine Vétérinaires de Dakar, Dakar, Senegal
63 ²⁹ Department of Small Animal Medicine, Faculty of Veterinary Medicine, Utrecht University,
64 Utrecht, The Netherlands
65 ³⁰ Vetexpert Veterinary Group, Yerevan, Armenia
66 ³¹ Veterinary Clinic Lopez Quintana, Maldonado, Uruguay
67 ³² Clinique Veterinaire de Grand Fond, Saint Gilles les Bains, Reunion, France
68 ³³ Department of Veterinary Sciences, University of Messina, Messina, Italy
69 ³⁴ Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma del Estado de
70 México, Toluca, Mexico
71 ³⁵ School of Veterinary Medicine, Universidad de las Américas, Quito, Ecuador
72 ³⁶ Touray & Meyer Vet Clinic, Serrekunda, The Gambia
73 ³⁷ Hillside Animal Hospital, St. Louis, USA

74 ³⁸ The Kampala Veterinary Surgery, Kampala, Uganda
75 ³⁹ Asavet Veterinary Charities, Tucson, United States
76 ⁴⁰ Vets Beyond Borders, Bylakuppe, India
77 ⁴¹ Faculty of Veterinary Medicine, Autonomous University of Yucatan, Merida, Mexico
78 ⁴² Laboratorio de Patología Veterinaria, Universidad de Caldas, Manizales, Colombia
79 ⁴³ Interdisciplinary Centre of Research in Animal Health (CIISA), Faculty of Veterinary
80 Medicine, University of Lisbon, Lisboa, Portugal
81 ⁴⁴ Four Paws International, Vienna, Austria
82 ⁴⁵ Vets Beyond Borders, The Rocks, Australia
83 ⁴⁶ Help in Suffering, Jaipur, India
84 ⁴⁷ Veterinary Clinic Dr José Rojas, Los Andes, Chile
85 ⁴⁸ Department of Biotechnology, Balochistan University of Information Technology,
86 Engineering and Management Sciences, Quetta, Pakistan
87 ⁴⁹ Corozal Veterinary Clinic, Corozal Town, Belize
88 ⁵⁰ Veterinary Clinic Vetmaster, Ramenskoye, Russia
89 ⁵¹ State Hospital of Veterinary Medicine, Dniprodzerzhynsk, Ukraine
90 ⁵² Jomo Kenyatta University of Agriculture and Technology, Juja, Kenya
91 ⁵³ Laboratory of Biomedicine and Regenerative Medicine, Department of Clinical Sciences,
92 Faculty of Animal and Veterinary Sciences, University of Chile, Santiago, Chile
93 ⁵⁴ Faculty of Veterinary and Agricultural Sciences, University of Melbourne, Melbourne,
94 Australia
95 ⁵⁵ Animal Anti Cruelty League, Port Elizabeth, South Africa
96 ⁵⁶ Clinical Sciences Department, Faculty of Veterinary Medicine Bucharest, Bucharest,
97 Romania
98 ⁵⁷ Department of Pathology, Faculty of Veterinary Medicine, Ankara University, Ankara,
99 Turkey
100 ⁵⁸ Faculty of Veterinary Sciences, National University of Asuncion, San Lorenzo, Paraguay
101 ⁵⁹ Lilongwe Society for Protection and Care of Animals (LSPCA), Lilongwe, Malawi
102 ⁶⁰ Wellcome Sanger Institute, Hinxton, UK
103 ⁶¹ Department of Cellular and Molecular Medicine, University of California, San Diego, La
104 Jolla, USA
105 * These authors contributed equally
106 [§]Corresponding author, epm27@cam.ac.uk

107 **Structured Abstract**

108 INTRODUCTION

109 The canine transmissible venereal tumour (CTVT) is a sexually transmitted cancer that
110 manifests as genital tumours in dogs. This cancer first arose in an individual 'founder dog'
111 several thousand years ago, and has since survived by transfer of living cancer cells to new
112 hosts during coitus. Today, CTVT affects dogs around the world and is the oldest and most
113 prolific known cancer lineage. CTVT thus provides a unique opportunity to explore the
114 evolution of cancer over the long-term, and to track the unusual biological transition from
115 multicellular organism to obligate conspecific asexual parasite. Furthermore, the CTVT
116 genome, acting as a living biomarker, has recorded the changing mutagenic environments
117 experienced by this cancer throughout millennia and across continents.

118 RATIONALE

119 To capture the genetic diversity of the CTVT lineage, we analysed somatic mutations
120 extracted from the protein-coding genomes (exomes) of 546 globally distributed CTVT
121 tumours. We inferred a time-resolved phylogenetic tree for the clone and used this to trace
122 the worldwide spread of the disease and to select subsets of mutations acquired at known
123 geographical locations and time-periods. Computational methods were applied to extract
124 mutational signatures and to measure their exposures across time and space. In addition, we
125 assessed the activity of selection using ratios of non-synonymous and synonymous variants.

126 RESULTS

127 The CTVT phylogeny reveals that the lineage first arose from its founder dog 4,000–8,500
128 years ago, likely in Asia, with the most recent common ancestor of modern globally distributed
129 tumours occurring ~1,900 years ago. CTVT underwent a rapid global expansion within the last
130 500 years, likely aided by intensification of human maritime travel. We identify a highly specific
131 mutational signature dominated by C>T mutations at GTCCA pentanucleotide contexts which
132 operated in CTVT up until ~1,000 years ago. The number of mutations caused by ultraviolet
133 light exposure is correlated with latitude of tumour collection, and we identify CTVTs with
134 heritable hyperactivity of an endogenous mutational process. Several 'driver' mutation
135 candidates are identified in the basal trunk of the CTVT tree, but there is little evidence for
136 ongoing positive selection. Although negative selection is detectable, its effect is largely
137 confined to genes with known essential functions, thus implying that CTVT predominantly
138 evolves via neutral processes.

139 CONCLUSION

140 We have traced the evolution of a transmissible cancer over several thousand years, tracking
141 its spread across continents and contrasting the mutational processes and selective forces
142 that moulded its genome with those described in human cancers. The identification of a highly
143 context-specific mutational process that operated in the past but subsequently vanished, as

144 well as correlation of ultraviolet light-induced DNA damage with latitude, highlight the potential
145 for long-lived, widespread clonal organisms to act as biomarkers for mutagenic exposures.
146 Our results suggest that neutral genetic drift is the dominant evolutionary force operating on
147 cancer over the long-term, in contrast to the ongoing positive selection which is often observed
148 in short-lived human cancers. The weakness of negative selection in this asexual lineage may
149 be expected to lead to the progressive accumulation of deleterious mutations, invoking
150 Muller's Ratchet and raising the possibility that CTVT may be declining in fitness despite its
151 global success.

152

153 **Abstract**

154 The canine transmissible venereal tumour (CTVT) is a cancer lineage that arose several
155 millennia ago and survives by 'metastasising' between hosts via cell transfer. The somatic
156 mutations in this cancer record its phylogeography and evolutionary history. We constructed
157 a time-resolved phylogeny from 546 CTVT exomes and describe the lineage's worldwide
158 expansion. Examining variation in mutational exposure, we identify a highly context-specific
159 mutational process that operated early but subsequently vanished, correlate ultraviolet-light
160 mutagenesis with tumour latitude, and describe tumours with heritable hyperactivity of an
161 endogenous mutational process. CTVT displays little evidence of ongoing positive selection,
162 and negative selection is detectable only in essential genes. We illustrate how long-lived clonal
163 organisms capture changing mutagenic environments, and reveal that neutral drift is the
164 dominant feature of long-term cancer evolution.

165

166 **Introduction**

167 Transmissible cancers are malignant somatic cell clones that spread between individuals via
168 direct transfer of living cancer cells. Analogous to the metastasis of cancer to distant tissues
169 within a single body, transmissible cancers 'metastasise' as allogeneic grafts between
170 individuals within a population (1). Such clones have been observed only eight times in nature,
171 suggesting that they arise rarely; however, once established, transmissible cancers can
172 spread rapidly and widely and persist through time (1, 2). Such cancers provide a unique
173 opportunity to explore the evolution of cancer over the long-term, and to track the unusual
174 biological transition from multicellular organism to obligate conspecific asexual parasite.

175

176 The canine transmissible venereal tumour (CTVT) is the oldest and most prolific known
177 contagious cancer (2, 3). It is a sexually transmitted clone that manifests as genital tumours
178 in dogs. This cancer first arose from the somatic cells of an individual 'founder dog' that lived
179 several thousand years ago (2). The cancer survived beyond the death of this original host by
180 transfer of cancer cells to new hosts. Subsequently, this cancer has spread around the world,

181 and is a common disease in dog populations globally, although it declined and largely
182 disappeared from many Western countries during the twentieth century due to the
183 management and removal of free-roaming dogs (4).

184

185 Similar to cancers that remain in a single individual, CTVT accumulates somatic mutations.
186 These result from the activities of endogenous and exogenous mutational processes, and
187 genetically imprint a cancer's history of mutagenic exposures (5). Thus, the CTVT genome
188 can be considered a living biomarker that records the changing mutagenic environments
189 experienced by this cancer throughout millennia and across continents. Although most
190 somatic mutations in cancer have no functional effect and are considered neutral 'passenger'
191 mutations, a subset of mutations are positively selected 'driver' mutations that confer the
192 proliferation and survival advantages that spur cancer growth (6). Ordinary cancers, which
193 remain in a single host, often acquire additional driver mutations during tumour progression
194 (7); however, it is unknown whether transmissible cancers that survive for hundreds or
195 thousands of years similarly continue to adapt. It seems possible that the evolution of long-
196 lived cancers such as CTVT may instead be dominated by negative selection acting to remove
197 deleterious mutations. Finally, in addition to recording a history of exposures and signatures
198 of selection, somatic mutations provide a tool for tracing CTVT phylogeography, potentially
199 revealing how dogs, together with humans, moved around the world over the last centuries.
200 Here, we use somatic mutations extracted from the protein-coding genomes (exomes) of 546
201 globally distributed CTVT tumours to trace the history, spread, diversity, mutational exposures
202 and evolution of the CTVT clone.

203

204 **CTVT phylogeny**

205 We sequenced the exomes (43.6 megabases, Mb; mean sequencing depth ~132×) of 546
206 CTVT tumours collected between 2003 and 2016 from 43 countries across all inhabited
207 continents (Data sets S1 and S2). Candidate somatic mutations were defined as single
208 nucleotide variants (SNVs) or short insertions and deletions (indels) identified in one or more
209 CTVT tumours, but not found in 495 normal dog exomes from the CTVT tumours' matched
210 hosts. This approach yielded 160,207 variants (148,030 SNVs, 3,392 per Mb; 12,177 indels,
211 279 per Mb; Table S1). The features of this set, including its variant allele fraction distribution,
212 phylogenetic structure, comparison with the distribution of private germline variants in the dog
213 population, mutational signature composition, and non-synonymous to synonymous mutation
214 ratio (details in (8)), suggest that it is very highly enriched for somatic mutations. However,
215 some minimal germline variation may remain, possibly including rare germline variants from
216 the founder dog and residual contaminating alleles from matched hosts.

217

218 We identified the subset of the candidate somatic mutations belonging to a clock-like
219 mutational process (specifically, cytosine-to-thymine (C>T) substitutions at CpG sites (8, 9)),
220 and used these to construct a time-resolved phylogenetic tree for the CTVT lineage (Fig. 1A).
221 The mutation rate was inferred by applying a Bayesian Poisson model to previously
222 ascertained empirical observations (10), and was estimated as 6.87×10^{-7} C>T mutations per
223 CpG site per year (8). The topology of the CTVT phylogenetic tree reveals a long basal trunk
224 (Fig. 1A), representing the chain of CTVT transmissions from its origin ~6,220 years ago (95%
225 highest posterior density interval, HPDI, 4,148–8,508 years ago) to the earliest detected node
226 ~1,938 years ago (95% HPDI 993–3,055 years ago). This node splits a set of five tumours
227 collected in India from the remaining population (groups labelled 57 and 58; Fig. 1A). The
228 second and third most basal nodes (respectively ~1,004 years ago, 95% HPDI 497–1,570
229 years ago, and ~829 years ago, 95% HPDI 424–1,310 years ago) separate sixteen tumours
230 from Eastern Europe and the Black Sea region, and three tumours from Northern India, from
231 the remaining set, respectively (groups labelled 54–56 and 1; Fig. 1A). Together with evidence
232 that the founder dog shared ancestry with ancient dog remains recovered in North-East
233 Siberia and North America (10), the CTVT phylogeny supports a model whereby CTVT
234 originated ~4,000–8,500 years ago in Central or Northern Asia, and remained within the area
235 for the subsequent 2,000–6,000 years. Starting less than ~2,000 years ago, CTVT escaped
236 from its founding population, perhaps due to contact between previously isolated dog groups,
237 and spread to several locations in Asia and Europe (Fig. 1B).

238

239 The more recent history of CTVT is marked by rapid global expansion (11) (Figs. 1C and S1).
240 CTVT was introduced to the Americas with early colonial contact (~500 years ago, 95% HPDI
241 284–888 years ago), probably initially to Central America, and further into North and South
242 America (red sublineage 1; Fig. 1, A and C). About 300 years ago, this sublineage spread out
243 of the Americas in an almost polytomous global sweep which brought CTVT into Africa at least
244 five times and re-introduced the disease to Europe and Asia (black sublineage 1; Fig. 1, A and
245 C). In parallel, a second tumour sublineage spread out of Asia or Europe into Australia and
246 the Pacific (sublineage 2; Fig. 1, A and D). This second sublineage is also detected in North
247 America, and its tumours were introduced to Africa on at least two occasions. By ~100 years
248 ago, CTVT was present in dog populations worldwide, establishing local lineages that have
249 since remained largely *in situ*. The CTVT phylogeny thus suggests that dogs, together with
250 their neoplastic parasites, were extensively transported around the world in the fifteenth to
251 early twentieth centuries, probably via sea travel.

252

253 **Mutational processes in CTVT**

254 The CTVT mutational spectrum, a representation of the six substitution types together with
255 their immediate 5' and 3' base contexts, is dominated by C>T mutations, as previously
256 described (12, 13) (Fig. 2A). Applying Markov chain Monte Carlo sampling on a Bayesian
257 model of mutational signatures (8, 14), we extracted signatures of five mutational processes
258 from the CTVT mutation load. These include three signatures that closely resemble COSMIC
259 (15) signatures 1, 5 and 7 (Fig. 2B). These signatures, which have previously been described
260 in CTVT (12), reflect endogenous mutational processes (signatures 1 and 5) and exposure to
261 ultraviolet light (UV, signature 7) (5). A fourth signature displaying some similarity (cosine
262 similarity 0.81) to COSMIC signature 2, which is associated with activity of APOBEC enzymes
263 (5), was also detected (labelled signature 2*, Fig. 2B).

264

265 The fifth signature extracted from CTVT does not resemble any previously described
266 mutational pattern. This signature, which we designate signature A, is characterised by C>T
267 mutations at NCC contexts and shows striking pentanucleotide sequence preference for
268 GTCCA (TGGAC on the complementary strand; Figs. 2, B and C, and S2). This extended
269 sequence preference is markedly more pronounced than previously reported pentanucleotide
270 context biases, such as those associated with UV light or DNA polymerase epsilon deficiency
271 (Fig. 2C) (16-18), and is not explained by the sequence composition of the canine exome (Fig.
272 S3). It is possible that signature A's causative mutagen is highly context-specific, or,
273 alternatively, that this signature's associated repair processes are ineffective at certain
274 sequence contexts ('repair shielding') (19). In addition, signature A displays strong
275 transcriptional strand bias, with more mutations of guanine on the untranscribed compared to
276 the transcribed strand of genes, indicating that its causative lesion is likely a guanine adduct
277 subject to transcription-coupled repair (TCR). Interestingly, the guanine-directed
278 transcriptional strand bias of signature A at TCC contexts counteracts the cytosine-directed
279 transcriptional strand bias of signature 7 at TCC, such that no overall transcriptional strand
280 bias is observed at this context in the CTVT mutational spectrum (Fig. 2A).

281

282 Using the CTVT phylogenetic tree to isolate subsets of mutations, we explored variation in
283 mutational signature exposure across time and space (Figs. S4 and S5, and Data set S3).
284 Remarkably, this revealed that signature A was highly active prior to ~2,000 years ago
285 (causing ~35% of mutations in the basal trunk of the tree, branch A1), and persisted in parallel
286 at lower levels in the two basal branches after the first node (~12% and ~9% of mutations in
287 branches A2 and A3, respectively), but then abruptly vanished (Figs. 2C and S5). Importantly,
288 signature A is not detectable within the germline of a global population of 495 dogs (Fig. S6).
289 It is possible that signature A reflects the activity of an exogenous mutagen that was uniquely
290 present in the environment that CTVT inhabited prior to its escape from its founding

291 population. Alternatively, it is plausible that signature A may result from an endogenous DNA-
292 damaging agent that occurred in CTVT cells early during the lineage's history, but which
293 ceased to accumulate from ~1,000 years ago, perhaps due to a cellular metabolic change.
294 Although the nature of such a change is unknown, the replacement of possibly defective
295 mitochondrial DNA by horizontal transfer, which likely occurred in parallel in branches A2 and
296 A3 within the last ~1,690 years (11), may have altered the metabolic environment within CTVT
297 cells.

298

299 Although CTVT usually occurs within the internal genital tract, it may sometimes protrude from
300 the genital orifice or spread to perineal skin, resulting in sporadic exposure to solar UV
301 radiation (12, 13). The amount of UV radiation reaching the Earth, however, varies significantly
302 across global environments (20). We investigated whether latitude influenced the degree of
303 UV exposure in CTVT tumours by estimating signature 7 contribution within subsets of
304 mutations acquired at known latitudes. Indeed, qualitative assessment of mutational spectra
305 of location-specific CTVT mutation subsets suggests substantial variation in UV exposure; for
306 example, the mutational spectra of tumours collected in Mauritius show considerably more
307 evidence of signature 7 compared with those of tumours collected in Russia (Fig. S4). Using
308 CC>TT dinucleotide mutations (21) as a proxy for signature 7 (Fig. S7), we identified a non-
309 linear association between latitude and UV exposure (Spearman's correlation -0.40 , 95%
310 HPDI $[-0.65, -0.14]$; Fig. 2D). By fitting CC>TT mutations observed in the basal trunk of the
311 CTVT tree to this curve, we estimated the latitude of the CTVT founder population (Fig. 2, D
312 and E) (8).

313

314 Examining the contribution of signature 5 across the CTVT lineage, we observed three
315 independent phylogenetic groups of tumours that appear to have acquired signature 5-
316 hyperactivity phenotypes (groups labelled 12–16, 20 and 40; Figs. 2, F and G, S4 and S5). In
317 one case, involving tumours collected in several South and Central American countries
318 (groups 12–16), the phenotype has been maintained for ~150 years. This phenotype is likely
319 to result from signature 5, and not from the double-strand DNA repair deficiency-mediated
320 COSMIC signature 3, which presents a similar mutational profile (5, 22), as we failed to
321 observe the enrichment for indels which co-occurs with signature 3 (22, 23). It is, however,
322 possible that these tumours were exposed to another, as yet undescribed, mutational process.
323 Signature 5 is widespread in cancer and normal tissues and has unknown aetiology, although
324 it may be partly associated with endogenously generated adducts subject to nucleotide
325 excision repair (5, 9, 18). We annotated non-synonymous mutations occurring in the three
326 groups' respective clonal ancestors, providing a catalogue of genes which may play a role in
327 generation or suppression of signature 5 (Data set S4).

328

329 **CTVT mutations and gene expression**

330 The prevalence of substitution mutations in CTVT decreases with increasing gene expression,
331 likely reflecting the activity of TCR operating on DNA damage associated with signatures 7
332 and A, as well as a signature 1 preference for genes with lower expression (16, 24, 25) (Fig.
333 S8, A and B). We observed that exons have a higher substitution prevalence than introns,
334 possibly due to sequence context (Figs. S8A and S9). The prevalence of indels is positively
335 correlated with increasing gene expression, as has been observed in human cancers, and
336 may reflect transcription-associated damage (26) (Fig. S8A).

337

338 We assessed the contribution of TCR in two temporally distinct subsets of mutations: those
339 acquired prior to the earliest detectable node in the phylogenetic tree (~8,500–2,000 years
340 ago; branch A1 in Fig. 1A), and those acquired subsequent to this node (~2,000 years ago to
341 present). Interestingly, although C>T mutations acquired at TCC contexts in highly expressed
342 genes in branch A1 have little strand bias, likely due to the opposing transcriptional strand
343 preferences of signatures 7 and A at this context, those genes with very low expression
344 predominantly show the transcriptional strand bias associated with signature A (Fig. S8C).
345 Assuming that the transcriptional strand bias observed in these low-expressed genes reflects
346 earlier expression and subsequent silencing of genes, this suggests that there may have been
347 an early period in CTVT evolution when the lineage was exposed to signature A more intensely
348 than it was to signature 7. This may reflect variation in the climate or environment to which
349 CTVT was exposed early in its history.

350

351 **Selection in CTVT**

352 CTVT has a massive mutation burden, which exceeds that observed in even the most highly
353 mutated human cancer types (Fig. 3A). Each CTVT tumour carries on average 37,800 SNVs
354 across its predominantly diploid (12) exome (~2 million SNVs genome-wide; Table S2).
355 Indeed, the tally of somatic mutations that have accumulated in CTVT since it departed its
356 original host is comparable with the number of germline variants that distinguish some pairs
357 of outbred dogs (Fig. S10). Within the set of 546 tumours, 14,412 (~73%) protein-coding genes
358 carry at least one non-synonymous mutation, and 5,704 (~29%) have mutations predicted to
359 cause protein truncation (Fig. 3B).

360

361 We searched for evidence of positive selection in CTVT. The driver mutations which initially
362 caused CTVT, and which promoted its transmissible phenotype, will have occurred in the
363 basal trunk of the CTVT tree. *SETD2*, *CDKN2A*, *MYC* (previously described (12)), *PTEN* and
364 *RB1*, known cancer genes that frequently harbour driver mutations in human cancers (15),

365 carry biallelic loss-of-function or potential activating mutations in the trunk and may be early
366 drivers of CTVT (Fig. 3C and Table S3). To search for late drivers, which may have been
367 acquired in more recent parallel CTVT lineages, we identified independent mutations that
368 occurred repeatedly across the tree, and measured the normalised ratio of non-synonymous
369 to synonymous mutations (dN/dS) per gene after correcting for mutational biases and context
370 effects (8). This approach only yielded two uncharacterised genes with $dN/dS > 1$ (q -value $<$
371 0.05), predicted to encode a neuroligin precursor and a roundabout homologue (Data set S5).
372 The potential for these genes to act as late drivers in CTVT cannot be assessed, and it is
373 possible that local sequence structures may result in higher than expected recurrent mutation
374 rates at these loci (27). Overall, we find little evidence that CTVT is continuing to adapt to its
375 environment.

376

377 Negative selection, which acts to remove deleterious mutations, is very weak in human
378 cancers (17, 28, 29). Human cancers have short life-spans, and their evolution is dominated
379 by sweeps of strong positive selection, thus reducing the potential for negative selection to act
380 (17). Given its long life-span, high mutation burden and lack of ongoing positive selection, it is
381 possible that negative selection may be a more dominant force in CTVT evolution. Further,
382 unlike in ordinary cancers, in CTVT inter-tumour competition may offer more opportunities for
383 negative selection to manifest, purging lineages less able to infect new hosts and spread
384 through the host population. Indeed, negative selection has been detected operating on CTVT
385 mitochondrial genomes (11). Our analysis of dN/dS in CTVT across all genes, however,
386 yielded $dN/dS \approx 1$ for both missense and nonsense mutations, indicating near-neutral
387 evolution (Fig. 3D and Data set S5). Similarly, dN/dS did not differ from neutrality in genes
388 categorised by expression level (Fig. 3D). Negative selection, acting both on missense and
389 nonsense mutations, could be detected, however, in sets of genes with known essential
390 functions (Fig. 3D), and was particularly pronounced for nonsense mutations in essential
391 genes occurring in haploid regions ($dN/dS = 0.33$, p -value $< 10^{-4}$). A slight signal of negative
392 selection acting on nonsense mutations in haploid regions ($dN/dS = 0.88$, p -value = 0.027) is
393 explained by 269 essential genes, as negative selection was not detected after removal of
394 these genes (Fig. 3D and Data set S5). These results imply that CTVT largely evolves via
395 neutral genetic drift. This may partly reflect functional obsolescence of many mammalian
396 genes in this relatively simple parasitic cancer, as well as the buffering effect of CTVT's largely
397 diploid genome (12). However, it is also likely that transmission bottlenecks between hosts
398 render weak selection inefficient. This may be expected to lead to the progressive
399 accumulation of deleterious mutations in the population (Muller's ratchet) (30), raising the
400 possibility that CTVT may be declining in fitness despite its global success.

401

402 **Discussion**

403 Studies of cancer evolution typically focus on how malignant clones alter during the first years,
404 or perhaps decades, of their existence. We have tracked the evolution of a cancer over several
405 thousand years, and compared the mutational processes and selective forces that moulded
406 its genome with those described in short-lived human cancers.

407

408 Our results suggest that neutral genetic drift may be the dominant evolutionary force operating
409 on cancer over the long-term, in contrast to the ongoing positive selection which is often
410 observed in human cancers (7, 17). Thus, our results suggest that CTVT may have optimised
411 its adaptation to the transmissible cancer niche early. Subsequently acquired advantageous
412 mutations may have offered incremental change of minimal benefit, such that they were
413 insufficient to overcome the neutral effects of drift. Importantly, since the 1980s, CTVT has
414 been routinely treated with vincristine, a cytotoxic microtubule inhibitor (31). Despite the strong
415 selection pressure imposed by vincristine treatment, we find no evidence of convergent
416 evolution of vincristine resistance mechanisms in CTVT at the level of point mutations or
417 indels.

418

419 The mechanisms whereby CTVT is tolerated by the host immune system, despite its status
420 as an allogeneic graft, are poorly understood (32, 33). The weakness of negative selection
421 beyond genes essential for cell viability implies that there are negligible selective pressures
422 imposed via immunoediting of somatic neoepitopes at a genome-wide level. This is perhaps
423 unsurprising, given the massive antigenic burden already presented by allogeneic epitopes.
424 These findings support evidence that CTVT largely circumvents the adaptive immune system,
425 at least during its initial stages of progressive tumour growth, perhaps in part via down-
426 regulation of major histocompatibility complex molecules (13, 33-35).

427

428 Our analyses reveal a mutational signature, signature A, which occurred in the past, but
429 ceased to be active from about 1,000 years ago. Interestingly, a recent study (36) detected
430 evidence for an excess of C>T mutations at TCC contexts, the mutation type most prevalent
431 in signature A, accumulating in the human germline between 15,000 and 2,000 years ago. If
432 this human mutation pulse is due to signature A, it could indicate a shared environmental
433 exposure which was once widespread, but which has now disappeared. However, we find no
434 evidence of an excess of C>T mutations at GTCCA pentanucleotides in the dog germline,
435 suggesting that dogs as a whole were not systemically exposed to signature A in their past.
436 Further research will be required to elucidate the biological origin of signature A and the
437 mechanism of its striking pentanucleotide sequence bias; however, this study highlights the

438 potential for long-lived, widespread clonal organisms to act as biomarkers for the activity of
439 mutational processes.

440

441 Genomic instability and ongoing positive selection are often considered key hallmarks of
442 carcinogenesis (37). CTVT does not have an intrinsically high point mutation rate ('genomic
443 instability'), at least at the level of SNVs, and its vast mutation burden simply reflects the
444 lineage's age. We find no clear evidence for continued positive selection beyond initial truncal
445 events. Thus, CTVT illustrates that, once spawned and sufficiently well-adapted to its niche,
446 neither hallmark is necessary to sustain cancer over the long term.

447

448 CTVT is a remarkable biological entity. It is the oldest, most prolific and most divergent cancer
449 lineage known in nature; it has spread throughout the globe and has seeded its tumours in
450 many thousands of dogs. Here, we have traced this cancer's route through the steppes of
451 Asia and Europe and as an unwelcome stowaway on global voyages. We have observed the
452 patterns in its mutational profiles reflecting the dynamics of its exogenous and endogenous
453 environment. Further, we have shown that CTVT largely evolves via neutral processes, and
454 that the mutations that it continues to acquire may pose a threat, rather than an advantage, to
455 its long-term fitness.

456

457 REFERENCES AND NOTES

- 458 1. M. J. Metzger, S. P. Goff, A Sixth Modality of Infectious Disease: Contagious Cancer
459 from Devils to Clams and Beyond. *PLoS Pathog* **12**, e1005904 (2016).
- 460 2. A. Strakova, E. P. Murchison, The cancer which survived: insights from the genome
461 of an 11000 year-old cancer. *Curr Opin Genet Dev* **30**, 49-55 (2015).
- 462 3. D. P. Blaine, *A domestic treatise on the diseases of horses and dogs*. (T. Boosey,
463 London, 1810).
- 464 4. A. Strakova, E. P. Murchison, The changing global distribution and prevalence of
465 canine transmissible venereal tumour. *BMC Vet Res* **10**, 168 (2014).
- 466 5. L. B. Alexandrov *et al.*, Signatures of mutational processes in human cancer. *Nature*
467 **500**, 415-421 (2013).
- 468 6. M. R. Stratton, P. J. Campbell, P. A. Futreal, The cancer genome. *Nature* **458**, 719-
469 724 (2009).
- 470 7. N. McGranahan, C. Swanton, Clonal Heterogeneity and Tumor Evolution: Past,
471 Present, and the Future. *Cell* **168**, 613-628 (2017).
- 472 8. Materials and methods are available as supplementary materials.
- 473 9. L. B. Alexandrov *et al.*, Clock-like mutational processes in human somatic cells. *Nat*
474 *Genet* **47**, 1402-1407 (2015).

- 475 10. M. Ní Leathlobhair *et al.*, The evolutionary history of dogs in the Americas. *Science*
476 **361**, 81-85 (2018).
- 477 11. A. Strakova *et al.*, Mitochondrial genetic diversity, selection and recombination in a
478 canine transmissible cancer. *Elife* **5**, (2016).
- 479 12. E. P. Murchison *et al.*, Transmissible dog cancer genome reveals the origin and
480 history of an ancient cell lineage. *Science* **343**, 437-440 (2014).
- 481 13. B. Decker *et al.*, Comparison against 186 canid whole-genome sequences reveals
482 survival strategies of an ancient clonally transmissible canine tumor. *Genome Res*
483 **25**, 1646-1655 (2015).
- 484 14. K. C. Gori, A. Baez-Ortega, sigfit: flexible Bayesian inference of mutational
485 signatures. *bioRxiv* **372896**, (2018).
- 486 15. S. A. Forbes *et al.*, COSMIC: exploring the world's knowledge of somatic mutations
487 in human cancer. *Nucleic Acids Res* **43**, D805-811 (2015).
- 488 16. E. D. Pleasance *et al.*, A comprehensive catalogue of somatic mutations from a
489 human cancer genome. *Nature* **463**, 191-196 (2010).
- 490 17. I. Martincorena *et al.*, Universal Patterns of Selection in Cancer and Somatic
491 Tissues. *Cell* **171**, 1029-1041 e1021 (2017).
- 492 18. L. B. Alexandrov *et al.*, The Repertoire of Mutational Signatures in Human Cancer.
493 *bioRxiv*, (2018).
- 494 19. S. S. Lange, K. M. Vasquez, HMGB1: the jack-of-all-trades protein is a master DNA
495 repair mechanic. *Mol Carcinog* **48**, 571-580 (2009).
- 496 20. R. L. McKenzie, P. J. Aucamp, A. F. Bais, L. O. Bjorn, M. Ilyas, Changes in
497 biologically-active ultraviolet radiation reaching the Earth's surface. *Photochem*
498 *Photobiol Sci* **6**, 218-231 (2007).
- 499 21. G. P. Pfeifer, Y. H. You, A. Besaratinia, Mutations induced by ultraviolet light. *Mutat*
500 *Res* **571**, 19-31 (2005).
- 501 22. S. Nik-Zainal *et al.*, Landscape of somatic mutations in 560 breast cancer whole-
502 genome sequences. *Nature* **534**, 47-54 (2016).
- 503 23. S. Morganella *et al.*, The topography of mutational processes in breast cancer
504 genomes. *Nat Commun* **7**, 11383 (2016).
- 505 24. N. J. Haradhvala *et al.*, Mutational Strand Asymmetries in Cancer Genomes Reveal
506 Mechanisms of DNA Damage and Repair. *Cell* **164**, 538-549 (2016).
- 507 25. E. Letouze *et al.*, Mutational signatures reveal the dynamic interplay of risk factors
508 and cellular processes during liver tumorigenesis. *Nat Commun* **8**, 1315 (2017).
- 509 26. S. Jinks-Robertson, A. S. Bhagwat, Transcription-associated mutagenesis. *Annu Rev*
510 *Genet* **48**, 341-359 (2014).

- 511 27. X. Zou *et al.*, Short inverted repeats contribute to localized mutability in human
512 somatic cells. *Nucleic Acids Res* **45**, 11213-11221 (2017).
- 513 28. D. Weghorn, S. Sunyaev, Bayesian inference of negative and positive selection in
514 human cancers. *Nat Genet* **49**, 1785-1788 (2017).
- 515 29. J. Van den Eynden, S. Basu, E. Larsson, Somatic Mutation Patterns in Hemizygous
516 Genomic Regions Unveil Purifying Selection during Tumor Evolution. *PLoS Genet*
517 **12**, e1006506 (2016).
- 518 30. H. J. Muller, The Relation of Recombination to Mutational Advance. *Mutat Res* **106**,
519 2-9 (1964).
- 520 31. E. I. Amber, R. A. Henderson, J. B. Adeyanju, E. O. Gyang, Single-drug
521 chemotherapy of canine transmissible venereal tumor with cyclophosphamide,
522 methotrexate, or vincristine. *J Vet Intern Med* **4**, 144-147 (1990).
- 523 32. D. Frampton *et al.*, Molecular Signatures of Regression of the Canine Transmissible
524 Venereal Tumor. *Cancer Cell* **33**, 620-633 e626 (2018).
- 525 33. H. V. Siddle, J. Kaufman, Immunology of naturally transmissible tumours.
526 *Immunology* **144**, 11-20 (2015).
- 527 34. C. Murgia, J. K. Pritchard, S. Y. Kim, A. Fassati, R. A. Weiss, Clonal origin and
528 evolution of a transmissible cancer. *Cell* **126**, 477-487 (2006).
- 529 35. A. Fassati, N. A. Mitchison, Testing the theory of immune selection in cancers that
530 break the rules of transplantation. *Cancer Immunol Immunother* **59**, 643-651 (2010).
- 531 36. K. Harris, J. K. Pritchard, Rapid evolution of the human mutation spectrum. *Elife* **6**,
532 (2017).
- 533 37. P. C. Nowell, The clonal evolution of tumor cell populations. *Science* **194**, 23-28
534 (1976).
- 535 38. M. P. Hoepfner *et al.*, An improved canine genome and a comprehensive catalogue
536 of coding genes and non-coding transcripts. *PLoS One* **9**, e91172 (2014).
- 537 39. H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler
538 transform. *Bioinformatics* **25**, 1754-1760 (2009).
- 539 40. A. Rimmer *et al.*, Integrating mapping-, assembly- and haplotype-based approaches
540 for calling variants in clinical sequencing applications. *Nat Genet* **46**, 912-918 (2014).
- 541 41. M. E. Skinner, A. V. Uzilov, L. D. Stein, C. J. Mungall, I. H. Holmes, JBrowse: a next-
542 generation genome browser. *Genome Res* **19**, 1630-1638 (2009).
- 543 42. D. Jones *et al.*, cgpCaVEManWrapper: Simple Execution of CaVEMan in Order to
544 Detect Somatic Single Nucleotide Variants in NGS Data. *Current protocols in*
545 *bioinformatics* **56**, 15.10. 11 (2016).
- 546 43. R. Rahbari *et al.*, Timing, rates and spectra of human germline mutation. *Nat Genet*
547 **48**, 126-133 (2016).

- 548 44. W. McLaren *et al.*, The Ensembl Variant Effect Predictor. *Genome Biol* **17**, 122
549 (2016).
- 550 45. A. Stamatakis, RAxML version 8: a tool for phylogenetic analysis and post-analysis
551 of large phylogenies. *Bioinformatics* **30**, 1312-1313 (2014).
- 552 46. P. Skoglund, E. Ersmark, E. Palkopoulou, L. Dalen, Ancient wolf genome reveals an
553 early divergence of domestic dog ancestors and admixture into high-latitude breeds.
554 *Curr Biol* **25**, 1515-1519 (2015).
- 555 47. L. R. Botigue *et al.*, Ancient European dog genomes reveal continuity since the Early
556 Neolithic. *Nat Commun* **8**, 16082 (2017).
- 557 48. B. Carpenter *et al.*, Stan: A probabilistic programming language. *Journal of statistical*
558 *software* **76**, (2017).
- 559 49. A. J. Drummond, M. A. Suchard, D. Xie, A. Rambaut, Bayesian phylogenetics with
560 BEAUti and the BEAST 1.7. *Mol Biol Evol* **29**, 1969-1973 (2012).
- 561 50. A. J. Drummond, A. Rambaut, B. Shapiro, O. G. Pybus, Bayesian coalescent
562 inference of past population dynamics from molecular sequences. *Mol Biol Evol* **22**,
563 1185-1192 (2005).
- 564 51. L. B. Alexandrov, S. Nik-Zainal, D. C. Wedge, P. J. Campbell, M. R. Stratton,
565 Deciphering signatures of mutational processes operative in human cancer. *Cell Rep*
566 **3**, 246-259 (2013).
- 567 52. A. Baez-Ortega, K. Gori, Computational approaches for discovery of mutational
568 signatures in cancer. *Brief Bioinform*, (2017).
- 569 53. K. Suzuki, D. Tonien, K. Kurosawa, K. Toyota, Birthday paradox for multi-collisions.
570 *IEICE Transactions on Fundamentals of Electronics, Communications and Computer*
571 *Sciences* **91**, 39-45 (2008).
- 572 54. R. Patro, G. Duggal, M. I. Love, R. A. Irizarry, C. Kingsford, Salmon provides fast and
573 bias-aware quantification of transcript expression. *Nat Methods* **14**, 417-419 (2017).
- 574 55. G. Tischler, S. Leonard, biobambam: tools for read pair collation based algorithms on
575 BAM files. *Source Code for Biology and Medicine* **9**, 13 (2014).
- 576 56. C. Sonesson, M. I. Love, M. D. Robinson, Differential analyses for RNA-seq:
577 transcript-level estimates improve gene-level inferences. *F1000Res* **4**, 1521 (2015).
- 578 57. J. Rainer, L. Gatto, C. X. Weichenberger, ensemblDb: an R package to create and
579 use Ensembl-based annotation resources. *Bioinformatics*, (2019).
- 580 58. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion
581 for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
- 582 59. S. Durinck, P. T. Spellman, E. Birney, W. Huber, Mapping identifiers for the
583 integration of genomic datasets with the R/Bioconductor package biomaRt. *Nat*
584 *Protoc* **4**, 1184-1191 (2009).

- 585 60. R. Nielsen, Z. Yang, Likelihood models for detecting positively selected amino acid
586 sites and applications to the HIV-1 envelope gene. *Genetics* **148**, 929-936 (1998).
- 587 61. L. D. Hurst, The Ka/Ks ratio: diagnosing the form of sequence evolution. *Trends*
588 *Genet* **18**, 486 (2002).
- 589 62. C. Roadmap Epigenomics *et al.*, Integrative analysis of 111 reference human
590 epigenomes. *Nature* **518**, 317-330 (2015).
- 591 63. T. Hart, K. R. Brown, F. Sircoulomb, R. Rottapel, J. Moffat, Measuring error rates in
592 genomic perturbation screens: gold standards for human functional genomics. *Mol*
593 *Syst Biol* **10**, 733 (2014).
- 594 64. V. A. Blomen *et al.*, Gene essentiality and synthetic lethality in haploid human cells.
595 *Science* **350**, 1092-1096 (2015).
- 596 65. T. Wang *et al.*, Identification and characterization of essential genes in the human
597 genome. *Science* **350**, 1096-1101 (2015).
- 598 66. J. D. Arroyo *et al.*, A Genome-wide CRISPR Death Screen Identifies Genes
599 Essential for Oxidative Phosphorylation. *Cell Metab* **24**, 875-885 (2016).
- 600 67. P. A. Futreal *et al.*, A census of human cancer genes. *Nat Rev Cancer* **4**, 177-183
601 (2004).
- 602 68. D. Hanahan, R. A. Weinberg, Hallmarks of cancer: the next generation. *Cell* **144**,
603 646-674 (2011).

604

605 **ACKNOWLEDGMENTS**

606 We acknowledge the Core Sequencing Facility, IT groups and members of the Cancer
607 Genome Project at the Wellcome Sanger Institute. We thank the following individuals for
608 useful information and for their help obtaining samples for this project: Ilona Airikkala-Otter,
609 Juliana Alzate-Ocampo, Diana Argüello, Jose Ignacio Arias, Clara Lee Arnold, Sue Barrass,
610 Ekaterina Batrakova, Rafaela Bortolotti Viéra, Nikki Brown, Fernando Constantino Casas,
611 John Cooper, Amici Cannis Cotacachi, Stephen M. Cutter, Johan de Vos, Lytvynenko Dmytro,
612 Phillip Farnham, Ariberto Fassati, Andres Fernandez-Riomalo, Ricardo Gaitan, David
613 Hanzlíček, Rafael Ricardo Hupples, John M. Igundu, Matilde Jimenez-Coello, Debra
614 Kamstock, Patrick Kelly, Tatiana Korytina, Anna Kuznetsova, Gleidice Eunice Lavalle
615 (Universidade Federal de Minas Gerais), Olakunle AbdulRasaq Lawal, Thabo Lerotholi, Marco
616 Lima-Maigua, Jimmy Loayza-Feijoo, Mayra López-Bucheli, Mwangi Maina, Margarita
617 Mancero-Albuja, Cynthia Marchiori Bueno, Luis Martínez-López, Alfredo Martínez-Meza,
618 Bedan M. Masuruli, Talita Mariana Morata Raposo, Jude Mulholland, Claudio Murgia, Alvira
619 Murison Swartz, Fran Nargi, Marsden M. Onsare, Edwin Ortiz-Rodríguez, Elisabeth Peach,
620 Lisa Pellegrini, Gerry Polton, Freddy Proaño-Pérez, Juan Carlos Ramirez-Ante, Cameron
621 Raw, Ceseltina Semedo, Ivan Stoikov, Irina Swarisch, Mirela Tinucci Costa, Emily Turitto, M.

622 Rifat Vural, David Walker, Robin Weiss, Kevin Xie, Maurice Zandvliet, staff at Animal Medical
623 Centre Belize City (Belize), veterinary surgeons and staff at Help in Suffering (Jaipur, India),
624 staff at Hopkins Belize Humane Society (Belize), veterinary workers at Pet Centre (UVAS,
625 Lahore, Pakistan), students from St. George's University (True Blue, Grenada, West Indies)
626 who assisted with sample collection, staff at Veterinary Clinic "El Roble" (Chile), staff and
627 volunteers at World Vets (Gig Harbor, USA) and staff at the WVS International Training Centre
628 in Ooty (India). We are grateful to the following organisations for helpful information: American
629 College of Veterinary Internal Medicine (ACVIM), Animal Balance, Animal Care Association
630 (The Gambia), Animal Management in Rural and Remote Indigenous Communities
631 (AMRRIC), Associacao Bons Amigos de Cabo Verde, Humane Society of Cozumel, Humane
632 Society Veterinary Medical Association–Rural Area Veterinary Services (HSVMA–RAVS),
633 Israel Veterinary Medical Association, Italian Veterinary Oncology Society, Rural Vets South
634 Africa, Veterinary Cancer Society, Veterinary Society of Surgical Oncology (VSSO),
635 VetPharma, Vets Beyond Borders, ViDAS and Coco's Animal Welfare, The Spanky Project,
636 VWB/VSF Canada, West Arnhem Land Dog Health Program (WALDHeP), World Small
637 Animal Veterinary Association (WSAVA), МИП ВЕТЕРИНАРИИ (World Veterinary Medicine).
638 **Funding:** This work was supported by Wellcome (102942/Z/13/A) and by a Philip Leverhulme
639 Prize from the Leverhulme Trust. A.S. was supported by a Postgraduate Student Award from
640 the Kennel Club Charitable Trust. **Author contributions:** E.P.M. designed and directed the
641 project. A.B.-O. developed methods and led computational data analysis. K.G. developed
642 methods and assisted with computational analysis. A.St. collected samples, performed
643 laboratory work, designed exome probes, oversaw sequencing and provided conceptual
644 advice. J.L.A., K.M.A., L.B.-I., T.N.B., J.L.B., C.B., A.C.D., A.M.C., H.R.C., J.T.C., E.D.,
645 K.F.d.C., A.B.D.N., A.P.d.V., L.D.K., E.M.D., A.R.E.H., I.A.F., M.F., E.F., S.N.F, F.G.-A., O.G.,
646 P.G.G., R.F.H.M., J.J.G.P.H., R.S.H., N.I., Y.K., C.K., D.K., A.K., S.J.K., M.L.-P., M.L.,
647 A.M.L.Q., T.L., G.M., S.M.C., M.F.M.-L., M.M., E.J.M., B.N., K.N., W.N., S.J.N., A.O.-P., F.P.-
648 O., M.C.P., K.P., R.J.P., J.F.R., J.R.G., H.S., S.K.S., O.S., A.G.S., A.E.S.-S., A.Sv., L.J.T.M.,
649 I.T.N., C.G.T., E.M.T., M.G.v.d.W., B.A.V., S.A.V., O.W., A.S.W.-M. and S.A.E.W. provided
650 clinical samples. Y.-M.K., M.N.L. and M.S. assisted with analysis and contributed to
651 interpretation of results. J.W. contributed to sample management and curation. M.R.S., L.B.A.
652 and I.M. provided technical advice and assisted with interpretation of results. A.B.-O. and
653 E.P.M. wrote the manuscript and designed the figures. All authors commented on the
654 manuscript. **Competing interests:** The authors declare no competing interests. **Data and**
655 **materials availability:** Whole-exome sequence data have been deposited in the European
656 Nucleotide Archive (ENA; <https://www.ebi.ac.uk/ena>) under overarching accession number
657 ERP109580. Variant calling data and other data supporting analyses have been deposited in
658 the University of Cambridge Repository (<https://www.repository.cam.ac.uk>) under DOI

659 10.17863/CAM.24962. Custom algorithms employed for data processing and analysis are
660 available in GitHub (<https://github.com/baezortega/TCG2019>).

661

662 **SUPPLEMENTARY MATERIALS**

663 Materials and Methods

664 Figs. S1 to S16

665 Tables S1 to S4

666 Data sets S1 to S6

667 References (38–68)

668

669 **Main figure legends**

670 **Fig. 1. Phylogeny and geographical expansion of CTVT. (A)** Time-resolved phylogenetic
671 tree inferred from clock-like exonic somatic variation in CTVT. Each tip is a tumour and
672 sampling locations are labelled. Numbers refer to phylogenetic groups displayed on maps in
673 **B–D**. Sublineages 1 and 2, referred to in **C** and **D** respectively, are marked. Three groups of
674 ancestral somatic variation (A1, A2, A3) and their respective numbers of single nucleotide
675 variants (SNVs) are indicated. The estimated age of the CTVT founder tumour and the earliest
676 detected node are indicated in years before present (BP), with grey error bars depicting
677 Bayesian 95% HPDI. **(B to D)** Maps presenting likely routes of early (prior to ~500 years BP)
678 and late (from ~500 years BP) expansion of CTVT. Numbered circles indicate the
679 geographical locations of phylogenetic groups labelled in **A**; arrows represent inferred
680 geographical movements. Circle and arrow colours indicate different sets of geographical
681 movements, as labelled in **A**. Thin arrows indicate expansion routes for which there is limited
682 phylogenetic evidence; dots without numbers denote tumours that are not represented in the
683 tree. C.V., Cape Verde; Gr., Greece; Guat., Guatemala; Hond., Honduras; Ken., Kenya; Rom.,
684 Romania; Tan., Tanzania; Tur., Turkey.

685

686 **Fig. 2. Mutational processes in CTVT. (A)** Trinucleotide-context mutational spectrum of
687 somatic SNVs in a single CTVT tumour. Horizontal axis presents 96 mutation types displayed
688 in pyrimidine context. Relevant trinucleotide mutation contexts are indicated. **(B)** Trinucleotide-
689 context mutational spectra of extracted mutational signatures 1, 5, 2*, A and 7, with relevant
690 trinucleotide mutation contexts indicated. **(C)** Pentanucleotide-context mutational spectra of
691 signature A (top) and signature 7 (bottom). Horizontal axis presents 256 C>T mutation types
692 with relevant mutation contexts indicated. The inset tree shows the phylogenetic branches
693 with exposure to signature A. **(D)** Bayesian logarithmic regression and Spearman's correlation
694 between absolute mean latitude and normalised CC>TT mutations in phylogenetic groups
695 shown in Fig. 1A. Normalised CC>TT mutations represent the ratio between group-unique

696 CC>TT mutations and group-unique C>T changes at CpG dinucleotides. The black line and
697 shadowed area indicate the regression curve and associated 95% HPDI. The orange dot and
698 bars represent predicted absolute mean latitude and associated 90% prediction interval for
699 the basal trunk ancestral variation (group A1). Posterior median and 95% HPDI of the
700 correlation coefficient are shown. (E) Map showing the latitude range corresponding to the
701 90% prediction interval for group A1, presented in D, in the northern hemisphere. (F)
702 Trinucleotide-context mutational spectra of a phylogenetic tumour group showing evidence of
703 signature 5 hyperactivity (top) and a closely related group without signature 5 hyperactivity
704 (bottom). (G) Diagram indicating the phylogenetic situation of the tumour groups displaying
705 signature 5 hyperactivity.

706
707 **Fig. 3. Selection in CTVT.** (A) Somatic SNV prevalence across six human cancer types and CTVT.
708 Dots represent individual tumours; red lines indicate median SNV prevalence. ALL, acute
709 lymphoblastic leukaemia. (B) Bars showing the percentage of protein-coding genes in the CTVT
710 genome harbouring ≥ 1 non-synonymous somatic mutation (SNV or indel; 14,412 genes) and ≥ 1
711 somatic protein-truncating somatic mutation (5,704 genes). (C) Diagram presenting the putative
712 driver events found in the set of basal trunk ancestral variants (group A1, Fig. 1A). A description of
713 each somatic alteration is shown next to the corresponding gene symbol. (D) Exome-wide dN/dS
714 ratios estimated for somatic SNVs in all protein-coding genes (left) and in sets of genes defined
715 according to gene essentiality, copy number state and expression level. Estimates of dN/dS are
716 presented for missense (blue) and nonsense (orange) mutations in each gene group. The dashed
717 line indicates dN/dS = 1 (neutrality); error bars indicate 95% confidence intervals.





