THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Chlorometallate and palladium cluster complexes of wide-span diimine and diamine ligands

Citation for published version:

Hart, JS, Parsons, S & Love, JB 2011, 'Chlorometallate and palladium cluster complexes of wide-span diimine and diamine ligands', *Dalton Transactions*, vol. 40, no. 45, pp. 12025-12027.
<https://doi.org/10.1039/c1dt10861a>

Digital Object Identifier (DOI):

[10.1039/c1dt10861a](https://doi.org/10.1039/c1dt10861a)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Dalton Transactions

Publisher Rights Statement:

Copyright © 2011 by the Royal Society of Chemistry. All rights reserved.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Post-print of a peer-reviewed article published by the Royal Society of Chemistry.

Published article available at: <http://dx.doi.org/10.1039/C1DT10861A>

Cite as:

Hart, J. S., Parsons, S., & Love, J. B. (2011). Chlorometallate and palladium cluster complexes of wide-span diimine and diamine ligands. *Dalton Transactions*, 40(45), 12025-12027.

Manuscript received: 08/05/2011; Accepted: 10/06/2011; Article published: 04/07/2011

Chlorometallate and palladium cluster complexes of wide-span diimine and diamine ligands**

John S. Hart,¹ Simon Parsons¹ and Jason B. Love^{1,*}

^[1]EaStCHEM, School of Chemistry, Joseph Black Building, University of Edinburgh, West Mains Road, Edinburgh, EH9 3JJ, UK.

^[*]Corresponding author; e-mail: jason.love@ed.ac.uk, fax: (+) 44 131 6504743

^[**]We thank the University of Edinburgh and the UK EPSRC for financial support.

Supporting information:

† Electronic supplementary information (ESI) available: Full experimental data and crystal data. CCDC reference numbers 824566–824571. For ESI and crystallographic data in CIF or other electronic format see <http://dx.doi.org/10.1039/C1DT10861A>

Graphical abstract:

Keywords:

organometallic oxides; crystal-structure; reduction; macrocycle; mechanism; design; anions

Abstract

Diimine and diamine ligands that are unable to coordinate to a single metal favour the formation of unusual, high-nuclearity Zn chlorometallate and palladium chloride complexes.

Introduction

Ligands that can promote the assembly of multiple metal cations into precisely-structured arrangements are highly desirable as these multimetallic motifs impact on a diverse range of chemical topics such as catalysis,¹ bio-inorganic chemistry,² medicine,³ and single molecule magnetism.⁴ In particular, ligand designs incorporating 1,4-disubstituted arenes have been intensively investigated as platforms that favour bi- and multinuclear complex formation due to the inability of the two donors to chelate to a single metal cation.⁵ For example, imine and amine acyclic ligands,⁶ macrocycles,⁷ and cryptands⁸ derived from 1,4-disubstituted arenes have been developed and show a propensity to form polynuclear complexes, although in simple systems cyclometallation reactions at the central aryl ring are observed, particularly with late transition metals.⁹

We are interested in developing straightforwardly-synthesised and inexpensive ligands for the assembly of multinuclear complexes of metals from across the Periodic Table,¹⁰ in particular to access new chemistry that can occur within well-defined cavities such as found in Pacman complexes,¹¹ and related supramolecular assemblies.¹² We report here on the use of new wide-span diimine and diamine ligands as outer and inner-sphere coordination platforms for the generation of new, high-nuclearity chlorometallate and palladium chloride complexes. The use of durene in the ligand design ensures that undesired cyclometallation reactions are minimised.

Scheme 1. Synthesis of wide-span diimine (L^{Im}) and diamine (L^{Am}) ligands and their chlorometallate and cluster complexes. Reagents and conditions: (i) H_2NR/Ar , MeCN; (ii) H_2NR/Ar , MeCN; (iii) $NaBH_4$, MeOH, $R = ^tBu$; (iv) 2 HCl, Et_2O , $CoCl_2$; (v) 2 HCl, Et_2O , 4 $ZnCl_2$; (vi) 3 $PdCl_2(MeCN)_2$, CH_2Cl_2 .

The pro-ligands L^{Im} and L^{Am} were synthesised in high yield from the readily available precursors **1** and **2** through straightforward bromide substitution and Schiff-base condensation routes, respectively (Scheme 1). Borohydride reduction of diimine L^{ImR} proved to be the most reproducible route to diamine L^{AmR} . The 2,6-diisopropylaryl-substituted diimine L^{ImAr} and diamine L^{AmAr} compounds were crystallised from saturated solutions of MeCN and their X-ray crystal structures determined (Fig. 1). In the solid state, both L^{ImAr} and L^{AmAr} form extended structures through intermolecular hydrogen-bonding interactions and display no solvent incorporation. In L^{ImAr} (Fig. 1, top), the imine N1-C13 bonds (1.236(2) Å) adopt an anti-conformation (torsion angle = 180.0(2)°) and are offset by 29.8(3)° with respect to the central aryl ring. The sterically-demanding iPr substituents on the terminal aryl groups result in an approximately orthogonal orientation (72.7(2)°) of this aryl ring to the imine N=C group. This orientation results in an intermolecular interaction between the C10 isopropyl methyl hydrogens and the central aryl ring (C10...C16 = 3.745(4) Å), so forming a chain motif in the extended structure. Reduction of the N=C double bond in L^{ImAr} to form L^{AmAr} allows free rotation around the N1-C13 single bond (1.489(2) Å) and results in a different structure in the solid state (Fig. 1, bottom) in which the N1-C13 bonds adopt a syn-conformation (torsion angles = 79.9(2)/84.0(2)°). As with L^{ImAr} , the steric demand of the $iPr_2(C_6H_3)$ substituent results in an approximately orthogonal orientation of this group in relation to the amine N1-C13 bond (76.1(2)/68.2(2)°), but in this case the presence of the amino hydrogen results in a hydrogen bond between the two amines (N1...N1' = 3.264(2) Å); π -H interactions are also seen in the extended structure.

Figure 1. Ball and stick representations of the X-ray crystal structures of diimine L^{ImAr} (left) and diamine L^{AmAr} (right). For clarity, all hydrogen atoms except those involved in hydrogen bonding are omitted.

The coordination chemistry of these ligands was investigated (Scheme 1). Reactions between L^{Im} or L^{Am} and either $CoCl_2$ or $ZnCl_2$ in THF showed little change in the resonances for L in the 1H NMR spectra. However, the use of $CDCl_3$ as the NMR solvent resulted in the crystallisation of the chlorometallate salts $[H_2L^{AmR}][CoCl_4]$ and $[H_2L^{ImAr}][Zn_4Cl_{10}]$, presumably as a result of the presence of adventitious HCl in

CDCl₃. Scaled-up reactions were carried out in which the ligands were protonated by HCl (2 eq) in Et₂O prior to the addition of CoCl₂ (1 eq) or ZnCl₂ (4 eq) in Et₂O. This latter procedure resulted in the formation of analytically pure materials. The X-ray crystal structures of both [H₂L^{AmR}][CoCl₄] and [H₂L^{ImAr}][Zn₄Cl₁₀] were determined (Fig. 2) and show that the protonated ligands interact with the chlorometallate salts through a series of hydrogen-bonding interactions; all hydrogens involved in hydrogen-bonding were located in the difference Fourier map and refined with riding thermal parameters and bond distance restraints.

Figure 2. Ball and stick representations of the X-ray crystal structures of [H₂L^{AmR}][CoCl₄](H₂O)(CHCl₃) (top) and [H₂L^{ImAr}][Zn₄Cl₁₀] (bottom). For clarity, CHCl₃ solvent of crystallisation and all hydrogen atoms except those involved in hydrogen bonding are omitted.

In the solid state structure of [H₂L^{AmR}][CoCl₄] (Fig 2., top) the amine appendages adopt a syn-conformation (torsion angle = 6.5(3)^o) that results in a hydrogen-bonded cavity suited for chlorometallate recognition (over chloride), with N-H⋯Cl and C-H⋯Cl hydrogen-bonding interactions ranging in distance from 3.165(3) to 3.357(3) Å. The chelation of the cobaltate anion by the wide-span ligand causes some strain, as evidenced by the 10^o curvature of the central diaminoarene and the inclusion of a molecule of water that interacts both with the protonated amine (N1⋯O1 2.817(5) Å) and the cobaltate (Cl4⋯O1 3.165(3) Å); as with the free ligands above, other intra- and intermolecular close contacts (to CHCl₃ solvent of crystallisation) are present in the extended structure.

In the solid state structure of [H₂L^{ImAr}][Zn₄Cl₁₀] (Fig. 2, bottom), the protonated imine arms adopt an anti-conformation (torsion angle = 176.0(2)^o) and the terminal aryl groups, and to a lesser extent the

central arene, are oriented-orthogonally to the N=C bond (torsion angles = 88.1(3) and 53.4(4)°, respectively). As with the cobaltate above, a variety of weak interactions exist between the doubly-protonated diimine and the zincate, with N-H...Cl and C-H...Cl hydrogen bonding interactions ranging in distance from 3.152(3) to 3.759(3) Å. However, in this case, the wide-span ligand framework has favoured the isolation of an unusual $\text{Zn}_4\text{Cl}_{10}^{2-}$ anionic cluster. In this cluster, each zinc cation is tetrahedral with one terminal chloride and three chlorides bridging to adjacent zinc cations. This results in a rectangular arrangement of zinc cations in which two opposing sides of the rectangle comprise either a single bridging chloride or two bridging chlorides, with Zn...Zn distances of 3.8345(5) Å and 3.1405(5) Å and Zn-Cl-Zn angles of 114.25(3) and 85.30(2)°. Also, the anti-conformation of the imines allows for an intermolecular N1...Cl3 hydrogen bond (3.152(3) Å) from which a linear chain of alternately sandwiched ligand and metallate clusters grows. While $\text{Zn}_4\text{Cl}_{10}^{2-}$ anionic clusters have been observed before, to our knowledge they adopt adamantane or open-chain structural motifs and not the new rectangular arrangement of Zn cations seen here.¹³ Furthermore, metal chloride clusters that adopt a similar structural motif are rare, and limited primarily to *o*-phenylene-metallated Pt_4 chlorides,¹⁴ $\text{Mo}_4\text{Cl}_{12}^{3-}$ clusters,¹⁵ and the chlorobismuthate and antimonates $\text{M}_4\text{Cl}_{18}^{6-}$ (M = Bi, Sb).¹⁶ Significantly, the recognition of high-nuclearity chlorometallates by L^{Im} and L^{Am} may impact on the high-efficiency, hydrometallurgical extraction of base and precious metals.¹⁷

The reaction between $\text{PdCl}_2(\text{MeCN})_2$ and either L^{ImAr} or L^{ImR} in CH_2Cl_2 led to the sole formation of the trinuclear palladium chloride cluster $[\text{Pd}_3\text{Cl}_6(\text{L})]$; this formulation was supported by elemental analysis. The X-ray crystal structures of $[\text{Pd}_3\text{Cl}_6(\text{L}^{\text{ImAr}})]$ and $[\text{Pd}_3\text{Cl}_6(\text{L}^{\text{ImR}})]$ were determined (Fig. 3) and were found to be isostructural; as such, only the former structure will be discussed (Fig.3, top). In a similar manner to $[\text{H}_2\text{L}^{\text{AmR}}][\text{CoCl}_4]$, the imine appendages adopt a syn-conformation (torsion angle = 6(2)°) that allows the ligand to chelate to the single Pd_3Cl_6 cluster through the imine nitrogens N1 and N2. Also, the terminal and central aryl rings are similarly tilted with respect to the N=C bond by 67(4) and 73(4)°, respectively. Each Pd cation is square planar but the Pd_3Cl_6 cluster is not linear due to the constraints imposed by the chelating diimine ligand. This feature results in a cradle arrangement in which the Pd1-Pd2-Pd3 angle is 127.4(4)° with Pd(μ -Cl)Pd angles of approximately 90°.

The solid state structure of $[\text{Pd}_3\text{Cl}_6(\text{L}^{\text{ImAr}})]$ is retained in solution. In the ^1H NMR spectrum (Fig. S1, Supplementary information), only one imine resonance is seen at 8.38 ppm, whereas two distinct CH_3 environments are seen at 3.75 and 2.09 ppm for the central arene, which reflects the asymmetry of binding to the Pd_3Cl_6 cluster. Furthermore, free rotation about the N-C bond for the terminal 2,6-diisopropyl aryl substituent is hindered, as reflected by the presence of two isopropyl CH resonances at 3.56 and 3.49 ppm and four resonances for the CH_3 environments. The ^1H NMR spectrum of $[\text{Pd}_3\text{Cl}_6(\text{L}^{\text{ImR}})]$ is similar (Fig. S2), indicating that the solid state structure of this complex is also retained in solution, but shows only one resonance for the ^tBu substituent at 1.22 ppm. Furthermore, the ESI mass

spectrum of $[\text{Pd}_3\text{Cl}_6(\text{L}^{\text{ImR}})]$ shows an ion at m/z 682 for the partially-intact cluster $[\text{Pd}_3\text{Cl}_5(\text{L}^{\text{ImR}})]^+$, and indicates structural retention in solution.

Figure 3. Ball and stick representations of the X-ray crystal structures of $[\text{Pd}_3\text{Cl}_6(\text{L}^{\text{ImAr}})]$ and $[\text{Pd}_3\text{Cl}_6(\text{L}^{\text{ImR}})]$. For clarity, disorder components, solvent of crystallisation, and all hydrogen atoms are omitted.

High nuclearity complexes of Pd chloride are rare and the $[\text{Pd}_3\text{Cl}_6(\text{L})]$ complexes can be viewed as short fragments of polymeric $(\text{PdCl}_2)_n$. While postulated through indirect evidence,¹⁸ only a single example of the crystallographic characterisation of a PdCl_2 trimer end-capped by organic ligands has been described;¹⁹ in this example, the unusual Pd_3Cl_6 motif was stabilised through the use of a large cone-angled, bowl-shaped triarylphosphine. Trinuclear PdCl structures are also known for allyl-complexes, although in these cases one chloride has been substituted by the allyl ligand.²⁰ The related dianion $\text{Pd}_3\text{Cl}_8^{2-}$ has also been characterised structurally as its Bu_4N ammonium salt.²¹ In all of these previous structures, the Pd_3Cl_6 motif is linear, in contrast to the cradle structure seen by us in $[\text{Pd}_3\text{Cl}_6(\text{L})]$. Presumably, this new bent geometry for the Pd_3Cl_6 cluster is necessary to allow a good fit to the wide-span diimine ligands. Furthermore, it is evident that methyl substitution of the central arene ligand has ensured that the formation of cyclometallated products is inhibited.

Notes and references

- [1] M. Delferro and T. J. Marks, *Chem. Rev.*, 2011, **111**, 2450; H. Dau, C. Limberg, T. Reier, M. Risch, S. Roggan and P. Strasser, *ChemCatChem*, 2010, **2**, 724; R. M. Haak, S. J. Wezenberg and A. W. Kleij, *Chem. Commun.*, 2010, **46**, 2713; M. J. Wiester, P. A. Ulmann and C. A. Mirkin, *Angew. Chemie. Int. Ed.*, 2010, **50**, 114.
- [2] J. P. Collman, R. Boulatov, C. J. Sunderland and L. Fu, *Chem. Rev.*, 2004, **104**, 561; R. H. Holm, P. Kennepohl and E. I. Solomon, *Chem. Rev.*, 1996, **96**, 2563.
- [3] Z. Guo and P. J. Sadler, *Angew. Chemie. Int. Ed.*, 1999, **38**, 1512; T. Storr, K. H. Thompson and C. Orvig, *Chem. Soc. Rev.*, 2006, **35**, 534.
- [4] T. Glaser, *Chem. Commun.*, 2011, **47**, 116; G. Aromí and E. K. Brechin, *Struct. Bond.*, 2006, **122**, 1.
- [5] A. L. Gavrilova and B. Bosnich, *Chem. Rev.*, 2004, **104**, 349.
- [6] F. Zhang, M. C. Jennings and R. J. Puddephatt, *Organometallics*, 2004, **23**, 1396.
- [7] S. Brooker and H. L. C. Feltham, *Coord. Chem. Rev.*, 2009, **235**, 1458; P. D. Frischmann and M. J. MacLachlan, *Comments Inorg. Chem.*, 2008, **29**, 26; J. Vela, L. Zhu, C. J. Flaschenriem, W. W. Brennessel, R. J. Lachicotte and P. L. Holland, *Organometallics*, 2007, **26**, 3416; P. Comba, T. W. Hambley, P. Hilfenhaus and D. T. Richens, *J. Chem. Soc., Dalton Trans.*, 1996, 533; P. Comba, A. Fath, T. W. Hambley and D. T. Richens, *Angew. Chemie. Int. Ed.*, 1995, **34**, 1883.
- [8] G. Morgan, V. McKee and J. Nelson, *Inorg. Chem.*, 1994, **33**, 4427; Y. Shu-Yan, L. Qin-Hui, S. Meng-Chang and H. Xiao-Yun, *Inorg. Chem.*, 1994, **33**, 1251.
- [9] Y.-F. Han, H. Li, L.-H. Weng and G.-X. Jin, *Chemical Communications*, 2010, **46**, 3556; B. J. O'Keefe and P. J. Steel, *Organometallics*, 1998, **17**, 3621.
- [10] J. B. Love, *Chem. Commun.*, 2009, 3154
- [11] A. M. J. Devoille, P. Richardson, N. L. Bill, J. L. Sessler and J. B. Love, *Inorg. Chem.*, 2011, **50**, 3116; P. L. Arnold, E. Hollis, F. J. White, N. Magnani, R. Caciuffo and J. B. Love, *Angew. Chem., Int. Ed.*, 2011, **50**, 887; J. W. Leeland, A. M. Z. Slawin and J. B. Love, *Organometallics*, 2010, **29**, 714; E. Askarizadeh, S. B. Yaghoob, D. M. Boghaei, A. M. Z. Slawin and J. B. Love, *Chem. Commun.*, 2010, **46**, 710; P. L. Arnold, A.-F. Pécharman, E. Hollis, A. Yahia, L. Maron, S. Parsons and J. B. Love, *Nat. Chem.*, 2010, **2**, 1056; E. Askarizadeh, A. M. J. Devoille, D. M. Boghaei, A. M. Z. Slawin and J. B. Love, *Inorg. Chem.*, 2009, **48**, 7491; P. L. Arnold, D. Patel, C. Wilson and J. B. Love, *Nature*, 2008, **451**, 315.

- [12] J. W. Leeland, F. J. White and J. B. Love, *J. Am. Chem. Soc.*, 2011, **133**, 7320; J. S. Hart, F. J. White and J. B. Love, *Chem. Commun.*, 2011, **47**, 5711; J. W. Leeland, F. J. White and J. B. Love, *Chem. Commun.*, 2011, **47**, 4132; P. L. Arnold, N. A. Potter, C. D. Carmichael, A. M. Z. Slawin, P. Roussel and J. B. Love, *Chem. Commun.*, 2010, **46**, 1833.
- [13] F. Bottomley, P. D. Boyle, S. Karslioglu and R. C. Thompson, *Organometallics*, 1993, **12**, 4090; F. Bottomley and S. Karslioglu, *J. Chem. Soc., Chem. Commun.*, 1991, 222; J. Estager, P. Nockemann, K. R. Seddon, M. Swadźba-Kwaśny and S. Tyrrell, *Inorg. Chem.*, 2011, **50**, 5258.
- [14] P. M. Cook, L. F. Dahl and D. W. Dickerhoof, *J. Am. Chem. Soc.*, 1972, **94**, 5511.
- [15] B. A. Aufdembrink and R. E. McCarley, *J. Am. Chem. Soc.*, 1986, **108**, 2474.
- [16] N. Leblanc, W. Bi, N. Mercier, P. Auban-Senzier and C. Pasquier, *Inorg. Chem.*, 2010, **49**, 5824; A. Piecha, R. Jakubas, V. Kinzhybalo and T. Lis, *J. Mol. Struct.*, 2008, **887**, 194.
- [17] R. J. Ellis, J. Chartres, K. C. Sole, T. G. Simmance, C. C. Tong, F. J. White, M. Schröder and P. A. Tasker, *Chem. Commun.*, 2009, 583.
- [18] P. M. Maitlis, D. Pollock, M. L. Games and W. J. Pryde, *Can. J. Chem.*, 1965, **43**, 470.
- [19] Y. Ohzu, K. Goto and T. Kawashima, *Angew. Chemie. Int. Ed.*, 2003, **42**, 5714; Y. Ohzu, K. Goto, H. Sato and T. Kawashima, *J. Organomet. Chem.*, 2005, **690**, 4175.
- [20] P. M. Bailey, E. A. Kelley and P. M. Maitlis, *J. Organomet. Chem.*, 1978, **144**, C52; B. T. Donovan, R. P. Hughes, P. P. Spara and A. L. Rheingold, *Organometallics*, 1995, **14**, 489; M. Parra-Hake, M. F. Rettig and R. M. Wing, *Organometallics*, 1983, **2**, 1013.
- [21] S. Schwarz, J. Strähle and U. Weisser, *Z. Anorg. Allg. Chem.*, 2002, **628**, 2495.