



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Definition: Optic ataxia

Citation for published version:

Rossetti, Y, Pisella, L & McIntosh, R 2019, 'Definition: Optic ataxia', *Cortex*.
<https://doi.org/10.1016/j.cortex.2019.09.004>

Digital Object Identifier (DOI):

[10.1016/j.cortex.2019.09.004](https://doi.org/10.1016/j.cortex.2019.09.004)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Cortex

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Definition: Optic ataxia

Rossetti Y, Pisella L, McIntosh R

Yves Rossetti (a), Laura Pisella (b), Rob McIntosh (c)

(a) Lyon Medical School, Inserm – CNRS- Université de Lyon, 16 Avenue Lépine, Bron, France, 69676.
Corresponding author.

(b) Lyon Medical School, Inserm – CNRS- Université de Lyon, 16 Avenue Lépine, Bron, France, 69676.

(c) The University of Edinburgh, Psychology Department, 7 George Square, Edinburgh, Scotland, EH8 9JZ

Definition: Optic ataxia

Rossetti Y, Pisella L, McIntosh R

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Optic ataxia is an impairment of visually guided reaching that is attributable neither to primary visual, primary proprioceptive or motor deficits, nor to cerebellar or cognitive impairments. It is one of the triad of symptoms that make up Bálint's syndrome, though it can be observed without the full syndrome following either bilateral or unilateral stroke, progressive degeneration or seizure, usually involving the posterior parietal cortex. Optic ataxia was named because the reaching problem was considered to be a disorder of aiming movements (ataxia) specific to visual targets (optic).

Patients with optic ataxia can usually reach and grasp accurately to targets that they fixate directly, although the ability to make fast trajectory corrections is severely deteriorated. This impairment slows them down but does not prevent them to interact with their environment. Visual eccentricity is a crucial determinant of the impairment, with larger errors for more eccentric targets and for goal-directed actions of the limbs monitored in peripheral vision such as when shaking hands or approaching a staircase, although this does not give rise to spontaneous complaints.

In unilateral cases, misreaching can be restricted to the contralesional visual field (field effect) or hand (hand effect). Errors associated with the field effect typically show a bias toward the point of fixation. Errors associated with the hand effect are most readily observed if visual feedback from the hand is unavailable during the reach. Reaching with the contralesional hand may then be inaccurate even for fixated targets.

In bilateral cases, misreaching may occur with either hand, and for targets throughout the visual field. In extreme cases, 'magnetic misreaching' can occur, such that the person reaches toward the point of gaze, rather than to the object presented in peripheral vision.