THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Total Synthesis of KingianinsA, D, and F

Citation for published version:

Drew, SL, Lawrence, A & Sherburn, MS 2013, 'Total Synthesis of KingianinsA, D, and F', *Angewandte Chemie International Edition*, vol. 52, no. 15, pp. 4221–4224. <https://doi.org/10.1002/anie.201210084>

Digital Object Identifier (DOI):

[10.1002/anie.201210084](https://doi.org/10.1002/anie.201210084)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Angewandte Chemie International Edition

Publisher Rights Statement:

Copyright © 2013 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim. All rights reserved.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

This is the peer-reviewed version of the following article:

Drew, S. L., Lawrence, A., & Sherburn, M. S. (2013). Total Synthesis of Kingianins A, D, and F. *Angewandte Chemie International Edition*, 52(15), 4221–4224.

which has been published in final form at <http://dx.doi.org/10.1002/anie.201210084>

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for self-archiving (<http://olabout.wiley.com/WileyCDA/Section/id-817011.html>).

Manuscript received: 18/12/2012; Article published: 06/03/2013

Total Synthesis of Kingianins A, D, and F**‡

Samuel L. Drew,¹ Andrew L. Lawrence^{1,*} and Michael S. Sherburn^{1,*}

^[1]Australian Research Council Centre of Excellence for Free Radical Chemistry and Biotechnology, Research School of Chemistry, Australian National University, Canberra, ACT 0200, Australia.

^[*]Corresponding author; A.L.L. (current address) a.lawrence@ed.ac.uk, EaStCHEM, School of Chemistry, Joseph Black Building, University of Edinburgh, West Mains Road, Edinburgh, EH9 3JJ, UK; M.S.S.: sherburn@rsc.anu.edu.au

^[**]This work was supported by the Australian Research Council. A.L.L. gratefully acknowledges financial support from the Australian Research Council in the form of a Discovery Early Career Researcher Award (DE120102113). S.L.D. gratefully acknowledges financial support from the Rickards family through the Rodney Rickards Scholarship. We thank Prof. Litaudon (CNRS, France) for kindly providing copies of the original NMR data. We also thank Mr. Tony Herlt (ANU) for assistance with HPLC and Mr. Chris Blake (ANU) for NMR measurements.

^[‡]In Memory of Rodney W. Rickards.

Supporting information:

Supporting information for this article is available online at <http://dx.doi.org/10.1002/anie.201210084>

Graphical abstract:

Keywords:

biomimetic synthesis; domino reactions; natural products; polyynes; radical cations

Abstract

A synthesis fit for a king: The total synthesis of (±)-kingianins A, D, and F has been achieved in ten steps. Key features include the gram-scale synthesis and partial reduction of a conjugated tetrayne to a (Z,Z,Z,Z)-tetraene, the domino 8 π –6 π electrocyclic ring closure of a (Z,Z,Z,Z)-tetraene, and the radical-cation-catalyzed formal Diels–Alder dimerization of functionalized bicyclo[4.2.0]octadiene precursors.

Main text

The kingianin natural products are a unique group of complex racemic bicyclo[4.2.0]octadiene dimers, isolated from the bark of *Endiandra kingiana* (Lauraceae) by Litaudon and co-workers.^[1] The first reported kingianin, (±)-kingianin A (**1**),^[1a] formulates as a dimer of bicyclo[4.2.0]octadiene **2**, and the Litaudon group proposed a biosynthesis involving spontaneous (non-enzyme-mediated) Diels–Alder dimerization (Scheme 1).^[1] Several reports, however, describe the need for temperatures in excess of 150 °C for Diels–Alder dimerization of 1,3-cyclohexadiene.^[2] The notion that a structural feature within compound **2** may lower the barrier to thermal Diels–Alder dimerization was investigated by Moses and co-workers in 2011.^[3] An elegant synthesis of monomer **2** was achieved by the Moses group, but all attempts to induce thermal dimerization failed.^[3] Inspired by the pioneering work of Bauld and co-workers,^[4] we hypothesized that a radical cation Diels–Alder dimerization could explain the formation of the kingianins in nature.

Scheme 1. Diels–Alder biosynthetic pathway to (±)-kingianin A (**1**), as proposed by Litaudon et al.^[1]

The bicyclo[4.2.0]octadiene framework present within Litaudon's proposed biosynthetic monomer **2** is a skeletal feature found in several natural products.^[5–9] The endiandric acids, which were isolated in racemic form in the early 1980s by Black and colleagues, were the first reported examples.^[5] Black proposed that the bicyclo[4.2.0]octadiene structure was formed through a spontaneous 8 π –6 π domino electrocyclization of either an (E,Z,Z,E)-tetraene or a (Z,Z,Z,Z)-tetraene (Scheme 2).^[5b–f] Beautiful biomimetic syntheses of various

bicyclo[4.2.0]octadiene natural products by Nicolaou,^[10] Trauner,^[11] Baldwin,^[12] Parker,^[13] and Moses^[14] have successfully utilized the proposed (*E,Z,Z,E*)-tetraene precursors. Evidently, the difficulty associated with preparing conjugated all-(*Z*)-polyenes has precluded their use in synthesis. In fact, (2*Z*,4*Z*,6*Z*,8*Z*)-decatetraene is both the highest all-(*Z*)-conjugated polyene and the only (*Z,Z,Z,Z*)-tetraene synthesized thus far.^[15]

Given the unprecedented structure and puzzling biosynthetic origin of the kingianin natural products,^[1] we decided to embark upon efforts towards their synthesis. The wealth of synthetic work in the literature utilizing (*E,Z,Z,E*)-tetraene precursors to access bicyclo[4.2.0]octadiene structures^[3, 10–14] convinced us that we should take this opportunity to investigate the alternative biosynthetic precursor, namely the (*Z,Z,Z,Z*)-tetraene (Scheme 2).^[5b–f] Although initially drawn to the sp^2 – sp^2 cross-coupling strategy utilized by Negishi for the synthesis of (*Z,Z,Z*)-trienes,^[16] we elected instead to investigate the feasibility of a four-fold stereoselective partial reduction of a conjugated tetrayne. We anticipated that if this unprecedented^[17] and highly challenging^[18] synthetic transformation were realized then a remarkably short synthesis of the kingianins could be achieved.

Scheme 2. The 8π – 6π biosynthesis of bicyclo[4.2.0]octadiene structures, as proposed by Black et al.^[5b–f]

The application of previously reported methods^[19] for the synthesis of unsymmetrical tetraynes was met with great difficulties. The instability of the requisite intermediates and problems associated with scaling up these approaches led us to develop a new scalable synthesis of unsymmetrical tetraynes. It is well known that steric bulk can stabilize polyene structures.^[20] We took advantage of this fact by targeting TBS-protected (TBS=*tert*-butyldimethylsilyl) alcohol tetrayne **3**,^[21] using Mori–Hiyama conditions for TMS-alkyne (TMS=trimethylsilyl) dimerization,^[22] thereby avoiding unstable halogenated and terminal polyynes. The two requisite diynes **4** and **5** were successfully prepared in three and two steps, respectively, on a multi-gram scale (Scheme 3). Thus, an Alami modified^[23] Cadiot–Chodkiewicz coupling of known bromobutynol **6**^[24] with ethynyltrimethylsilane afforded TMS-diyne **7**, which was converted into TBS-ether **4** under standard conditions.^[25] Meanwhile, known benzyl bromide **8**^[26] was employed in a Negishi reaction^[27] with organozinc reagent **9**,^[28] which was derived from 1,4-bis(trimethylsilyl)buta-1,3-diyne.^[29] Following extensive optimization, tetrayne **3** was isolated in 40 % yield on a gram scale.^[30] This is the first reported crossed Mori–Hiyama coupling reaction^[22] and the first gram-scale synthesis of an unsymmetrical tetrayne.^[19]

With significant quantities of tetrayne **3** now available, investigation into the daunting four-fold reduction could begin.^[17, 18] Following extensive experimentation, it was found that Rieke zinc in ethanol afforded (*Z,Z,Z,Z*)-tetraene **10** in a completely chemoselective and highly diastereoselective manner (Scheme 4).^[17, 31] A solution of tetraene **10** in toluene was immediately heated to 100 °C, which triggered the domino 8 π –6 π electrocyclization sequence.^[32] Following deprotection, the two diastereomeric alcohols **11** and **12** were isolated in a combined yield of 21 % from tetrayne **3** (Scheme 4).

Scheme 3. Gram-scale synthesis of unsymmetrical tetrayne **3**. dppf=1,1'-bis(diphenylphosphino)ferrocene, NBS=*N*-bromosuccinimide, TBS=*tert*-butyldimethylsilyl, TMS=trimethylsilyl.

We were delighted to find that both alcohols **11** and **12** underwent fast radical cation Diels–Alder dimerizations using catalytic quantities of the Ledwith–Weitz aminium salt, (*p*-BrC₆H₄)₃N⁺·SbCl₆[−] (**13**; Scheme 4).^[33] Amide **2**, the proposed biosynthetic precursor to (±)-kingianin A (**1**),^[1a] failed to dimerize under these reaction conditions (Scheme 4).

Scheme 4. Completion of the total synthesis of (±)-kingianins A, D, and F. EDC=1-ethyl-3-(3-dimethylaminopropyl) carbodiimide, HOBt=hydroxybenzotriazole, NMO=*N*-methylmorpholine-*N*-oxide, TBAF=tetrabutylammonium fluoride, TPAP=tetrapropylammonium perruthenate.

The synthesis of (±)-kingianins A (**1**) and D (**14**) was eventually optimized to a sequence involving oxidation of alcohol **11** using the tetrapropylammonium perruthenate/ *N*-methylmorpholine-*N*-oxide (TPAP/NMO) conditions of Stark et al.,^[34] with the product directly subjected to radical cation Diels–Alder dimerization using the Ledwith–Weitz salt (**13**; 5 mol %).^[4, 33] The resultant mixture of diastereomeric diacids was directly converted into the corresponding diamides. Column chromatography afforded a mixture of three dimeric diamides in 17 % yield over the three steps from alcohol **11**. Reverse-phase preparative HPLC allowed the isolation of analytically pure samples of (±)-kingianin A (**1**), (±)-kingianin D (**14**) and a third, as yet undetermined, structure.^[1] This radical cation Diels–Alder dimerization is a remarkably selective reaction, with only three of the potential thirty-two isomeric products isolated. Both (±)-kingianin A (**1**), a homochiral dimer, and (±)-kingianin D (**14**), a heterochiral dimer, are the result of *endo*-selective formal Diels–Alder reactions occurring at the convex faces of both diene and dienophile. Previous studies have shown that the radical cation Diels–Alder dimerization of 1,3-cyclohexadiene is *endo* selective,^[4] however, a full explanation of the site and orientational regioselectivity observed in the present study will require further investigation. The natural product (±)-kingianin F (**15**) was similarly obtained by dimerization of the other bicyclo[4.2.0]octadiene diastereomer **12**, followed by double oxidation and diamide formation.^[35]

In summary, our highly divergent biomimetic strategy has resulted in the total synthesis of ($\pm$)-kingianins A (**1**), D (**14**) and F (**15**), in a longest linear sequence of ten steps. The noteworthy synthetic aspects of our successful approach include the gram-scale preparation of an unsymmetrical tetrayne, the unprecedented reduction of a conjugated tetrayne to a (*Z,Z,Z,Z*)-tetraene, and radical cation Diels–Alder dimerization of functionalized bicyclo[4.2.0]octadienes. From these studies, we conclude that the kingianins are not formed through spontaneous Diels–Alder dimerization. Instead, we propose that nature uses a SET-mediated cycloaddition analogous to the approach described herein.^[36, 37] Our results, in conjunction with previous biomimetic syntheses,^[3, 10–14] demonstrate that (*E,Z,Z,E*)-tetraenes, and not their all-(*Z*) congeners,^[32] are the likely biosynthetic precursors to bicyclo[4.2.0]octadiene natural products.^[38]

Notes and references

- [1] a) A. Leverrier, M. E. Tran Huu Dau, P. Retailleau, K. Awang, F. Guéritte, M. Litaudon, *Org. Lett.* 2010, **12**, 3638; b) A. Leverrier, K. Awang, F. Guéritte, M. Litaudon, *Phytochemistry* 2011, **72**, 1443.
- [2] For experimental and theoretical studies regarding thermal dimerization of 1,3-cyclohexadiene, see: a) F. Hofmann, P. Damm, *Chem. Zentr.* 1926, **1**, 2342; b) K. Alder, G. Stein, *Justus Liebigs Ann. Chem.* 1932, **496**, 197; c) D. Valentine, N. J. Turro, Jr., G. S. Hammond, *J. Am. Chem. Soc.* 1964, **86**, 5202; d) G. R. de Maré, G. Huybrechts, M. Toth, P. Goldfinger, *Trans. Faraday Soc.* 1971, **67**, 1397; e) F.-G. Klärner, B. M. J. Dogan, O. Ermer, W. von E. Doering, M. P. Cohen, *Angew. Chem.* 1986, **98**, 109; *Angew. Chem. Int. Ed. Engl.* 1986, **25**, 108; f) D. H. Ess, A. E. Hayden, F.-G. Klärner, K. N. Houk, *J. Org. Chem.* 2008, **73**, 7586.
- [3] P. Sharma, D. J. Ritson, J. Burnley, J. E. Moses, *Chem. Commun.* 2011, **47**, 10605. No spontaneous dimerization of compounds **2**, **11**, **12** (or the corresponding oxidation products) was observed during the present study (see the Supporting Information for details), nor has any spontaneous dimerization of any bicyclo[4.2.0]octadiene ever been reported.
- [4] D. J. Bellville, D. W. Wirth, N. L. Bauld, *J. Am. Chem. Soc.* 1981, **103**, 718.
- [5] For endiandric acids, see: a) W. M. Bandaranayake, J. E. Banfield, D. St. C. Black, G. D. Fallon, B. M. Gatehouse, *J. Chem. Soc. Chem. Commun.* 1980, 162; b) W. M. Bandaranayake, J. E. Banfield, D. St. C. Black, *J. Chem. Soc. Chem. Commun.* 1980, 902; c) W. M. Bandaranayake, J. E. Banfield, D. St. C. Black, G. D. Fallon, B. M. Gatehouse, *Aust. J. Chem.* 1981, **34**, 1655; d) W. M. Bandaranayake, J. E. Banfield, D. St. C. Black, *Aust. J. Chem.* 1982, **35**, 557; e) W. M. Bandaranayake, J. E. Banfield, D. St. C. Black, G. D. Fallon, B. M. Gatehouse, *Aust. J. Chem.* 1982, **35**, 567; f) J. E. Banfield, D. St. C. Black, S. R. Johns, R. I. Willing, *Aust. J. Chem.* 1982, **35**, 2247.
- [6] For SNF4435 compounds, see: a) K. Kurosawa, K. Takahashi, E. Tsuda, *J. Antibiot.* 2001, **54**, 541; b) K. Takahashi, E. Tsuda, K. Kurosawa, *J. Antibiot.* 2001, **54**, 548; c) K. Kurosawa, K. Takahashi, E. Tsuda, A. Tomida, T. Tsuruo, *Jpn. J. Cancer Res.* 2001, **92**, 1235; d) K. Kurosawa, K. Takahashi, N. Fujise, Y. Yamashita, N. Washida, E. Tsuda, *J. Antibiot.* 2002, **55**, 71.
- [7] For ocellapyrones A and B, see: E. Manzo, M. L. Ciavatta, M. Gavagnin, E. Mollo, S. Wahidulla, G. Cimino, *Tetrahedron Lett.* 2005, **46**, 465.
- [8] For elysiapyrones A and B, see: M. Cueto, L. D'Croz, J. L. Maté, A. San-Martín, J. Darias, *Org. Lett.* 2005, **7**, 415.

- [9] For shimalactones A and B, see: a) H. Wei, T. Itoh, M. Kinoshita, N. Kotoku, S. Aoki, M. Kobayashi, *Tetrahedron* 2005, **61**, 8054; b) H. Wei, T. Itoh, N. Kotoku, M. Kobayashi, *Heterocycles* 2006, **68**, 111.
- [10] a) K. C. Nicolaou, N. A. Petasis, R. E. Zipkin, J. Uenishi, *J. Am. Chem. Soc.* 1982, **104**, 5555; b) K. C. Nicolaou, N. A. Petasis, J. Uenishi, R. E. Zipkin, *J. Am. Chem. Soc.* 1982, **104**, 5557; c) K. C. Nicolaou, R. E. Zipkin, N. A. Petasis, *J. Am. Chem. Soc.* 1982, **104**, 5558; d) K. C. Nicolaou, N. A. Petasis, R. E. Zipkin, *J. Am. Chem. Soc.* 1982, **104**, 5560.
- [11] a) C. M. Beaudry, D. Trauner, *Org. Lett.* 2002, **4**, 2221; b) J. E. Barbarow, A. K. Miller, D. Trauner, *Org. Lett.* 2005, **7**, 2901; c) C. M. Beaudry, D. Trauner, *Org. Lett.* 2005, **7**, 4475; d) A. K. Miller, D. Trauner, *Angew. Chem.* 2005, **117**, 4678; *Angew. Chem. Int. Ed.* 2005, **44**, 4602.
- [12] a) J. E. Moses, J. E. Baldwin, R. Marquez, R. M. Adlington, *Org. Lett.* 2002, **4**, 3731; b) M. F. Jacobsen, J. E. Moses, R. M. Adlington, J. E. Baldwin, *Org. Lett.* 2005, **7**, 2473; c) J. E. Moses, R. M. Adlington, R. Rodriguez, S. J. Eade, J. E. Baldwin, *Chem. Commun.* 2005, 1687.
- [13] K. A. Parker, Y.-H. Lim, *J. Am. Chem. Soc.* 2004, **126**, 15968.
- [14] S. J. Eade, M. W. Walter, C. Byrne, B. Odell, R. Rodriguez, J. E. Baldwin, R. M. Adlington, J. E. Moses, *J. Org. Chem.* 2008, **73**, 4830.
- [15] D. Holme, E. R. H. Jones, M. C. Whiting, *Chem. Ind.* 1956, 928.
- [16] G. Wang, S. Mohan, E.-i. Negishi, *Proc. Natl. Acad. Sci. USA* 2011, **108**, 11344.
- [17] The reduction of conjugated diynes represents the current state of the art, with no reports on the reduction of higher-order conjugated polyynes appearing in the literature. For a recent example of the use of diyne reduction in total synthesis, see: N. J. Matovic, P. Y. Hayes, K. Penman, R. P. Lehmann, J. J. De Voss, *J. Org. Chem.* 2011, **76**, 4467.
- [18] Over-reduction, poor stereoselectivity, and low yields are commonly encountered during the reduction of conjugated alkynes. For examples, see Ref. [16] and: a) E. N. Marvell, J. Tashiro, *J. Org. Chem.* 1965, **30**, 3991; b) L. A. Paquette, J. Chang, Z. Liu, *J. Org. Chem.* 2004, **69**, 6441; c) P. Kumar, S. V. Naidu, P. Gupta, *J. Org. Chem.* 2005, **70**, 2843; d) C. L. Benson, F. G. West, *Org. Lett.* 2007, **9**, 2545.
- [19] For examples of unsymmetrical tetrayne syntheses, see: a) B. W. Gung, G. Kumi, *J. Org. Chem.* 2003, **68**, 5956; b) S. A. Vail, P. J. Krawczuk, D. M. Guldi, A. Palkar, L. Echegoyen, J. P. C. Tomé, M. A. Fazio, D. I. Schuster, *Chem. Eur. J.* 2005, **11**, 3375; c) G. Sabitha, Ch. S. Reddy, J. S. Yadav, *Tetrahedron Lett.* 2006, **47**, 4513; d) E. Métay, Q. Hu, E.-i. Negishi, *Org. Lett.* 2006, **8**, 5773; e) S. Kim, Y. M. Lee, H. R. Kang, J. Cho, T. Lee, D. Kim, *Org. Lett.* 2007, **9**, 2127; f) J. Kendall, R. McDonald, M. J. Ferguson, R. R. Tykwinski, *Org. Lett.* 2008, **10**, 2163; g) M. Abe, A. Kubo, S. Yamamoto, Y. Hatoh, M. Murai, Y.

- Hattori, H. Makabe, T. Nishioka, H. Miyoshi, *Biochemistry* 2008, **47**, 6260; h) B. B. Frank, M. Kivala, B. C. Blanco, B. Breiten, W. B. Schweizer, P. R. Laporta, I. Biaggio, E. Jahnke, R. R. Tykwinski, C. Boudon, J.-P. Gisselbrecht, F. Diederich, *Eur. J. Org. Chem.* 2010, 2487; i) K. Aikawa, Y. Hioki, K. Mikami, *Org. Lett.* 2010, **12**, 5716; j) C. R. Kumar, C.-H. Tsai, Y.-S. Chao, J.-C. Lee, *Chem. Eur. J.* 2011, **17**, 8698.
- [20] a) W. A. Chalifoux, R. R. Tykwinski, *C. R. Chim.* 2009, **12**, 341; b) W. A. Chalifoux, R. R. Tykwinski, *Nat. Chem.* 2010, **2**, 967.
- [21] The free-alcohol congener of **3** was significantly less stable.
- [22] a) Y. Nishihara, K. Ikegashira, K. Hirabayashi, J.-i. Ando, A. Mori, T. Hiyama, *J. Org. Chem.* 2000, **65**, 1780. These Mori–Hiyama conditions have previously been used to make a symmetrical tetrayne; see: b) V. Fiandanese, D. Bottalico, G. Marchese, A. Punzi, *Tetrahedron* 2008, **64**, 53.
- [23] M. Alami, F. Ferri, *Tetrahedron Lett.* 1996, **37**, 2763.
- [24] J. M. Montierth, D. R. DeMario, M. J. Kurth, N. E. Schore, *Tetrahedron* 1998, **54**, 11741.
- [25] E. J. Corey, A. Venkateswarlu, *J. Am. Chem. Soc.* 1972, **94**, 6190.
- [26] D. Imperio, T. Pirali, U. Galli, F. Pagliai, L. Cafici, P. L. Canonico, G. Sorba, A. A. Genazzani, G. C. Tron, *Bioorg. Med. Chem.* 2007, **15**, 6748.
- [27] M. Qian, E.-i. Negishi, *Tetrahedron Lett.* 2005, **46**, 2927.
- [28] T. N. Hoheisel, H. Frauenrath, *Org. Lett.* 2008, **10**, 4525.
- [29] A. B. Holmes, G. E. Jones, *Tetrahedron Lett.* 1980, **21**, 3111.
- [30] The more readily accessible TMS-diyne **5** was used in excess to increase the yield of the cross-coupled product. A 1:1 ratio of TMS-diynes **4** and **5** typically afforded unsymmetrical tetrayne **3** in yields of 25–30 %.
- [31] W.-N. Chou, D. L. Clark, J. B. White, *Tetrahedron Lett.* 1991, **32**, 299.
- [32] We believe that the 100 °C temperature required for the electrocyclization of **10** discounts the all-(*Z*) isomer as a likely biosynthetic precursor. The previously reported 8 π –6 π electrocyclization of (2*Z*,4*Z*,6*Z*,8*Z*)-decatetraene required 14 h at 65 °C to go to completion, see: R. Huisgen, A. Dahmen, H. Huber, *J. Am. Chem. Soc.* 1967, **89**, 7130.
- [33] F. A. Bell, A. Ledwith, D. C. Sherrington, *J. Chem. Soc. C* 1969, 2719.

- [34] A.-K. C. Schmidt, C. B. W. Stark, *Org. Lett.* 2011, **13**, 4164.
- [35] The three-step sequence to produce (±)-kingianin F proved higher yielding when the radical cation Diels–Alder dimerization was carried out first, followed by double oxidation, and diamide formation.
- [36] To the best of our knowledge, this constitutes only the second reported example of a biomimetic radical cation Diels–Alder reaction; see: S. Lin, M. A. Ischay, C. G. Fry, T. P. Yoon, *J. Am. Chem. Soc.* 2011, **133**, 19350.
- [37] While this manuscript was under review, Parker and Lim published an article detailing the total synthesis of (±)-kingianin A, by a tether-mediated intramolecular radical cation Diels–Alder strategy; see: H. N. Lim, K. A. Parker, *Org. Lett.* 2013, **15**, 398.
- [38] Although, it should be noted that enzyme participation in the kingianin biosynthesis cannot be ruled out.