THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Assignment of the TIMP gene to the murine X-chromosome using an inter-species cross

Citation for published version:

Jackson, I, LeCras, TD & Docherty, AJ 1987, 'Assignment of the TIMP gene to the murine X-chromosome using an inter-species cross', *Nucleic Acids Research*, vol. 15, no. 10, pp. 4357.

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Nucleic Acids Research

Publisher Rights Statement:

© 1987 IRL Press Limited,

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Assignment of the TIMP gene to the murine X-chromosome using an inter-species cross

Ian J.Jackson¹, T.D.LeCras² and A.J.P.Docherty²

¹MRC Clinical and Population Cytogenetics Unit, Western General Hospital, Crewe Road, Edinburgh EH4 2XU and ²Celltech Ltd, 244-250 Bath Road, Slough SL1 4DY, UK
Submitted April 3, 1987

The glycoprotein tissue inhibitor of metalloproteinases (TIMP) is a physiological inhibitor of collagenase, stromelysin and gelatinase, which are thought to be involved in extracellular matrix turnover (1). It inhibits to a lesser extent the PMN leucocyte metalloproteinases (1). The gene for human TIMP has been cloned (2), and it was shown to be identical to a cDNA previously identified as erythroid potentiating activity. This human gene has been mapped to the X-chromosome (3,4). We have isolated the mouse TIMP gene from an embryonic cDNA library (provided by Dr. Brigid Hogan) by cross-hybridisation to the human gene. DNA sequencing of the mouse coding sequence shows it to have 72% amino-acid identity with its human counterpart (T.D. Le Cras, unpublished).

Linkage of particular genes to the X-chromosome is conserved throughout the mammals. We therefore anticipated X-linkage of the mouse TIMP gene. To establish this linkage we utilised a cross between two mouse species, *M. musculus* and *M. spretus*. As the two species are 4-5 million years diverged, almost all DNA probes reveal a difference in restriction fragment length after gel electrophoresis and Southern blot hybridisation. The Figure shows that the mouse TIMP probe identifies 2 fragments in EcoR1-digested DNA from *M. musculus* (labelled mu) and two different fragments in *M. spretus* DNA (sp). Hybridisation to EcoR1 digested DNA from male and female F1 offspring of a cross between a male *M. spretus* and a female *M. musculus* is shown. The female offspring inherits the fragments from both parents, whilst the male offspring inherits only the maternal (*musculus*) fragments. This clearly demonstrates the the TIMP gene is on the murine X-chromosome, and has no cross-hybridising homologues on the Y-chromosome or the autosomes.

This method is generally applicable and provides a rapid means of demonstrating X-linkage of any DNA probe.

REFERENCES

- 1) Cawston, T.E. (1986) in *Proteinase Inhibitors*, Barrett, A.J. and Salvesen, G. eds., Elsevier, N.Y. pp 589-606.
- 2) Docherty, A.J.P., Lyons, A., Smith, B.J., Wright, E.M. Stephens, P.E., Harris, T.J.R., Murphy, G. and Reynolds, J.J. (1985) *Nature*, **318** 66-69.
- 3) Spurr, N.K., Goodfellow, P.N. Craig, I.W. and Docherty, A.J.P. (1987) *Ann. Hum. Genetics*, (in press)
- 4) Huebner, K., Isobe, M., Gasson, J.C., Golde, D.W. and Croce, C.M. (1986) *Am. J. Hum. Genet.* **38** 819-826.