

# THE UNIVERSITY of EDINBURGH

# Edinburgh Research Explorer

### Retention ç pesticide Endosulfan by nanofiltration

**Citation for published version:** De Munari, A, Semiao, AJC & Antizar-Ladislao, B 2013, 'Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions', *Water Research*, vol. 47, no. 10, pp. 3484-3496. https://doi.org/10.1016/j.watres.2013.03.055

## Digital Object Identifier (DOI): 10.1016/j.watres.2013.03.055

### Link:

Link to publication record in Edinburgh Research Explorer

Document Version: Peer reviewed version

Published In: Water Research

**General rights** Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy** The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess @ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Retention of pesticide Endosulfan by nanofiltration: Influence of

organic matter-pesticide complexation and solute-membrane

interactions

Submitted to Water Research

### ANNALISA DE MUNARI<sup>1</sup>, ANDREA JOANA CORREIA SEMIAO<sup>2</sup>, BLANCA ANTIZAR-

### LADISLAO<sup>1,\*</sup>

<sup>1</sup> School of Engineering, The University of Edinburgh,

Edinburgh, EH9 3JL, United Kingdom

<sup>2</sup>School of Chemical and Bioprocess Engineering, University College Dublin, Dublin 4, Ireland

\*Corresponding author: Blanca Antizar-Ladislao, Institute for Infrastructure and Environment, School of Engineering, The University of Edinburgh, Edinburgh EH9 3JL, United Kingdom, E-mail: B.Antizar-Ladislao@ed.ac.uk, Tel. +44(0)131 650 5712 De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

1 Abstract

2 Nanofiltration (NF) is a well-established process used in drinking water production to effectively 3 remove Natural Organic Matter (NOM) and organic micropollutants. The presence of NOM has been shown to have contrasting results on micropollutant retention by NF membranes and removal 4 5 mechanisms are to date poorly understood. The permeate water quality can therefore vary during 6 operation and its decrease would be an undesired outcome for potable water treatment. It is hence 7 important to establish the mechanisms involved in the removal of organic micropollutants by NF 8 membranes in the presence of NOM. In this study, the retention mechanisms of pesticide 9 Endosulfan (ES) in the presence of humic acids (HA) by two NF membranes, TFC-SR2 and TFC-10 SR3, a "loose" and a "tight" membrane, respectively, were elucidated. The results showed that two 11 mechanisms were involved: (1) the formation of ES-HA complexes (solute-solute interactions). 12 determined from solid-phase micro-extraction (SPME), increased ES retention, and (2) the 13 interactions between HA and the membrane (solute-membrane interactions) increased membrane molecular weight cut-off (MWCO) and decreased ES retention. HA concentration, pH, and the ratio 14 15 between micropollutant molecular weight (MW) and membrane MWCO were shown to influence 16 ES retention mechanisms. In the absence of HA-membrane interactions at pH 4, an increase of HA 17 concentration increased ES retention from 60% to 80% for the TFC-SR2 and from 80% to 95% for 18 the TFC-SR3 due to ES-HA complex formation. At pH 8, interactions between HA and the loose 19 TFC-SR2 increased the membrane MWCO from 460 to 496 g/mol and ES retention decreased from 20 55% to 30%, as HA-membrane interactions were the dominant mechanism for ES retention. In 21 contrast, for the "tight" TFC-SR3 membrane the increase in the MWCO (from 165 to 179 g/mol), 22 was not sufficient to decrease ES retention which was dominated by ES-HA interactions. 23 Ouantification of the contribution of both solute-solute interactions and solute-membrane interactions is hence fundamental in understanding the removal mechanisms of micropollutant by 24 25 NF membranes in the presence of NOM in order to optimize the treatment process.

- 26
- 27 Keywords: nanofiltration, micropollutant, natural organic matter, solute-solute interactions, solute-
- 28 membrane interactions
- 29
- 30

### 31 1. Introduction

Removal of organic micropollutants from water intended for human consumption is a growing concern for the drinking water industry due to increasing occurrence of these compounds in surface water and groundwater, as well as their adverse effects on human health (Schwarzenbach et al. 2006). Application of nanofiltration (NF) processes for the removal of micropollutants has become widespread due to an effective rejection achieved by full scale plants (Cyna et al. 2002) and the extensive research carried out in explaining the micropollutant removal mechanisms (Bellona et al. 2004, Van der Bruggen et al. 2006).

39

40 Natural organic matter (NOM) is ubiquitously present in water and it influences the fate and the 41 behaviour of micropollutants and their removal by NF (Plakas and Karabelas 2009, Salvaterra et al. 42 2011). NOM has shown to increase, decrease or have negligible effect on micropollutant removal 43 by NF (Agbekodo et al. 1996, Devitt et al. 1999, Jin et al. 2007, 2010, Plakas et al. 2006, Xu et al. 44 2005). Contrasting results have been attributed to the different types of micropollutants, membranes 45 and organic matter used in previous studies and to the complexity of the retention mechanisms 46 (Bellona et al. 2004, Boussahel et al. 2002, Jin et al. 2007, Sanches et al. 2011).

47

48 The increase of micropollutant retention in the presence of NOM has been attributed to the 49 formation of micropollutant-NOM complexes of bigger size and higher charge compared to the 50 micropollutant on its own (Agbekodo et al. 1996, Boussahel et al. 2002, Devitt et al. 1999, Jin et al. De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

- 51 2007, 2010, Plakas et al. 2006, Schäfer et al. 2010). Jin et al. (2007) attempted to estimate estrone
- 52 removal by NF in the presence of humic acids (HA) using the percentage of the fraction of estrone
- 53 bound to HA from literature. Poor estimation was attributed to the difference in estrone 54 concentration between the experiments and the literature. Jin et al. (2007) also inferred that most of 55 the estrone-HA interactions took place on the membrane rather than in solution, explaining further 56 the poor removal estimation. Neale and Schäfer (2012) were able to quantify the contribution of 57 hormone-HA interactions on increased hormone removal by ultrafiltration (UF) in the presence of 58 HA by determining organic matter-water partition coefficients. Increasing HA concentration led to 59 more hormone partition onto HA and in turn to greater hormone removal by UF, demonstrating the role of micropollutant-NOM interactions on micropollutant removal. Increase of micropollutant 60 61 removal by NF in the presence of NOM was also attributed to the modification of the membrane 62 surface caused by adsorption of NOM. NOM deposition can cause pore clogging and change 63 membrane surface charge, resulting in improved retention by steric and electrostatic mechanisms (Jin et al. 2007, Xu et al. 2005). 64
- 65

66 Several mechanisms have been inferred to explain reduced or similar retention of micropollutants in 67 the presence of NOM. Lack of binding between the studied micropollutants and NOM was 68 considered the reason for unchanged retention (Boussahel et al. 2002, Jin et al. 2007, 2010). 69 Reduced micropollutant retention was attributed to increased membrane molecular weight cut-off 70 (MWCO) caused by the presence of organic matter (Xu et al. 2005, Xu et al. 2006). Organic matter 71 has been shown to enhance the negative charge of NF membranes, so it was inferred that the 72 increased electrostatic repulsion within the membrane pores increased membrane MWCO 73 (Braghetta et al. 1997, Childress and Elimelech 1996). Schäfer et al. (2010) inferred that membrane 74 hydrophilization might be responsible for the observed reduction in estrone removal when organic 75 matter was present and for the increase in pure water flux after organic matter filtration. The 76 decrease in micropollutant retention in the presence of organic matter was observed for "loose"

77 membranes (*i.e.* membrane whose MWCO is bigger than micropollutant molecular weight, MW)
78 (Schäfer et al. 2010, Xu et al. 2005, Xu et al. 2006). From the literature, it can be hypothesized that
79 a correlation might exist between micropollutant removal mechanisms in the presence of NOM and
80 micropollutant MW/membrane MWCO ratio. In the cited studies, the mechanisms responsible for
81 reduced micropollutant retention in the presence of NOM were presented as hypotheses without
82 being quantified, indicating the need to perform thorough investigations to demonstrate for which
83 conditions the mechanisms apply.

84

85 For micropollutants that adsorb onto the membranes, retention decreases with time until the membrane is saturated with the compound and real retention is reached (Kimura et al. 2003, 86 87 Nghiem et al. 2004, Semião and Schäfer 2011). In some case studies there is no indication that 88 membrane saturation was achieved. If the membrane is not saturated with the compound, retention 89 can therefore be overestimated and the comparison of retention results with and without NOM can 90 lead to erroneous conclusions. For example, lower micropollutant retention in the presence of NOM 91 has been linked to adsorption competition on the membrane between the micropollutant and NOM. 92 When NOM is present, it adsorbs on the membrane and decreases the membrane adsorption sites 93 available for the micropollutant. It was therefore inferred that the decrease in micropollutant adsorption in the presence of NOM caused a lower retention of the same micropollutant (Yoon et al. 94 95 2006). Vice versa, higher micropollutant adsorption was inferred to cause lower retention because 96 adsorbed micropollutants might experience higher diffusion to the permeate side (Boussahel et al. 97 2002). These two opposing results might be linked to the above-mentioned lack of saturation of the 98 membrane by the micropollutant. It is hence important to ensure membrane saturation during the 99 experiments in order to be able to compare retention results in the presence and absence of NOM. 100 These findings reflect the complexity of the mechanisms involved in the removal of micropollutants 101 by NF membranes in the presence of NOM and invite a systematic investigation of NOM influence 102 on micropollutant removal.

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

103

104 Three main mechanisms can influence micropollutant retention by NF in the presence of NOM

105 solute-solute interactions, solute-membrane interactions and solute-foulant interactions. The impact

106 of fouling by NOM on micropollutant retention has been extensively studied (Hajibabania et al.

107 2001, Ng and Elimelech 2004, Plakas et al. 2006, Xu et al. 2006), so it was not further investigated

- 108 in this study.
- 109

110 In this work, commercial HA have been chosen as a representative of the bigger fraction of NOM 111 because they have been extensively characterized in the literature and they were shown to have 112 similar organic matter retention properties than real Scottish NOM (Chin et al. 1994, De Munari 113 2012, Shin et al. 1999). Pesticide Endosulfan (ES) has been employed as a model micropollutant 114 because it is believed to interact with NOM and it is neutral over a large pH range (Banasiak 2009, 115 Forman et al. 1965). Moreover, ES is extensively used worldwide, persistent in the environment, 116 toxic to aquatic life and has estrogenic properties similar to DDT (German Federal Environment 117 Agency 2007, Soto et al. 1994). Two commercial NF membranes, TFC-SR2 and TFC-SR3 118 provided by Koch, have been selected as their MWCO is respectively higher and lower than the ES 119 molecular weight (MW).

120

121 The aim of this study is to elucidate the mechanisms responsible for the different retention trends 122 obtained for the micropollutants in the presence of NOM. The objectives were to: (1) quantify the 123 contribution of solute-solute interactions (i.e. formation of ES-HA complexes) on ES removal; (2) 124 investigate the contribution of ES-membrane interactions on ES retention; and (3) investigate the 125 contribution of HA-membrane interactions on ES retention for a "tight" and a "loose" membrane. 126 For the first time, the relative contribution of solute-solute interactions and solute-membrane 127 interactions to micropollutant retention by NF membranes in the presence of organic matter will be 128 evaluated to explain micropollutant retention mechanisms.

129

130 In order to quantify the influence of solute-solute interactions on ES retention, the method 131 developed by Neale and Schäfer for UF (2012) was adapted to NF and applied to quantify the 132 contribution of ES-HA complexes on ES removal. HA-water partition coefficient K<sub>HA</sub> for ES was determined and the calculated fraction of ES bound to HA was used to estimate ES removal due to 133 134 complex formation. Neale and Schäfer (2012) did not consider the contribution of solute-membrane interactions since the dominant mechanism for hormone removal by UF was deemed to be solute-135 136 solute interactions only. In this study, the contribution of solute-membrane interactions on 137 micropollutant retention was investigated by evaluating ES adsorption to the membranes (ESmembrane interactions) and the influence of HA on membrane characteristics (HA-membrane 138 139 interactions).

140

### 141 **2. Materials and methods**

### 142 2.1 Stirred cells equipment and filtration protocol

Experiments were performed with a stainless steel stirred cell with a volume of 990 mL and a diameter of 70 mm, resulting in a membrane area of 38.5 cm<sup>2</sup>. The cell was pressurised at 5 bar with filtered lab air to create similar conditions as full scale membrane plants; the pressure was kept constant and automatically measured every minute with a pressure transducer (Omega Engineering, UK). Temperature inside the cell and permeate collected on an electronic balance (Advancer Pro, Ohaus, UK) were measured every minute. A digital magnetic stirrer (Fisher Scientific, UK) was used at a speed of 300 rpm.

150

Membrane coupons were compacted for an hour at 15 bar and pure water flux was measured at 5
bar, before and after the experiments. A 900 mL feed solution was prepared prior to the experiments
and stirred at 150 rpm for 24 hours to ensure formation of ES-HA complexes. The experiments

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

- 154 were carried out in concentration mode in two phases. In the first phase, 480 mL of feed were
- 155 filtered in order to ensure membrane saturation by ES, where a feed, a permeate and a concentrate
- 156 sample were collected to determine ES sorption by mass balance. Filtration of 480 mL of solution
  157 was determined as sufficient to reach ES saturation for both membranes. In the second phase, the
- 158 permeate was returned to the stirred cell, a feed, a concentrate and four permeate samples of 60 mL
- 159 each were collected. Mass of ES sorbed to the membranes was determined with a mass balance on
- 160 the total solution filtered during the experiment and no adsorption of micropollutant occurred
- 161 during the filtration of the 4×60 mL. Retention was determined with a mass balance on the last
- 162 permeate collected; retention was constant for the last four permeates since equilibrium had been
- 163 reached.
- 164

### 165 2.2 Membrane characteristics

166 TFC-SR2 and TFC-SR3 membranes provided by Koch Membrane Systems (San Diego, USA) were extensively characterized. Membrane MWCO was determined by filtering aqueous solutions 167 168 containing dioxane, xylose, dextrose and polyethylene glycol (PEG) of different MW (Teixeira et 169 al. 2005). Pore radius was estimated using the hydrodynamic model (Nghiem et al. 2004). Point of 170 zero charge was measured with an electrokinetic analyser (EKA, Anton Paar KG, Austria) with the 171 same background electrolyte used in the experiments. TFC-SR2 and TFC-SR3 membranes are 172 amphoteric, positively charged at acidic pH and negatively charged with increasing pH. Contact 173 angle measurements (CAM 100, KSV Instrument Ltd, US) were performed using the sessile drop 174 method. At least three measurements per membrane were performed and the average of the 175 measurements is reported (standard deviation  $\pm 2^{\circ}$ ). In the case of membranes with HA deposits, 176 which presented zones of different colour, measurements were repeated in triplicate for different 177 zones and the average was reported. The variation between different zones of the membranes with 178 HA deposit was  $\pm 2^{\circ}$ , comparable with the variation obtained by repeated measurements on virgin 179 membranes. Roughness was estimated by atomic force microscopy (AFM) using a multimode AFM

system (VEECO, presently Bruker Corporation, USA) by contact mode in liquid (scan size 2.0×2.0 180

- μm). The cantilever (Mikromasch CSC38/AIBS type B) was made of Si with a tip radius of 10 nm. 181
- 182
- 183 Since a new membrane coupon was used for each experiment, the reproducibility of retention and 184 adsorption results was mainly affected by the variability of different membrane coupons. For 185 quality control, only coupons with a pure water flux varying  $\pm 15\%$  were employed in the 186 experiments.
- 187

### 188 2.3 Chemicals and solution chemistry

Experiments were conducted in ultra-pure water obtained by PuraLab Ultra (Elga LabWater, UK). 189 190 All chemicals were of analytical grade. Inert organics for MWCO and pore size determination 191 consisted of 25 mgC/L of dioxane, dextrose PEG 400, PEG 600, PEG 100 (Fisher Scientific, UK) 192 and xylose (Acros Organics, UK). HA (Sigma Aldrich, UK) stock solutions containing 250 mgC/L 193 were conserved in the fridge in the dark for one week maximum. HA was used at a concentration of 194 12.5 mgC/L because it represents typical NOM concentration found in upland natural water in Scotland (De Munari 2012). Experiments with HA ranging from 5 to 250 mgC/L were performed to 195 196 study the effects of HA concentration on ES-HA complex formation. Background electrolyte solution consisted of 1 mM NaHCO<sub>3</sub> and 20 mM NaCl (Fisher Scientific, UK). pH was adjusted 197 198 with 1M of HCl or NaOH (Fisher Scientific, UK). Radiolabeled [2,3-14C] ES (>95% purity; 18.5 199 MBq solid form) was purchased from the Institute of Isotopes Co., Ltd. (Hungary). Feed solutions 200 of 10 µg/L of radiolabeled ES were prepared from stock solutions (10,000 µg/L) made in methanol 201 (CH<sub>3</sub>OH) (Fisher Scientific, UK).

202

ES is an organochloride insecticide of the cyclodiene group commercially produced as a 7:3 203 204 isomeric mixture of  $\alpha$  and  $\beta$  forms (Table 1). ES isomers degrade by chemical hydrolysis to ES-diol 205 at pH > 8 with half time of less than 12 hours (Peterson and Batley 1993), comparable with the time 9

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", Water Research 47(10):3484-3496, doi:10.1016/i.watres.2013.03.055

- used in the filtration experiments. ES-sulfate can also be a hydrolysis product of ES isomers in the 206
- 207 presence of microbiological activity or strong chemical oxidants but it was not detected in
- 208 "artificial" samples after eight weeks (Peterson and Batley 1993). Therefore, it was inferred that
- 209 ES-sulfate was not present in the solutions prepared for the experiments.
- 210
- 211 (TABLE 1)
- 212

### 213 2.4 Analytical methods

- 214 Radiolabeled ES was measured with a liquid scintillation counter (Beckman LS 6500, USA); 1mL
- 215 of sample was analysed in 20 mL scintillation vials containing 7 mL of scintillation liquid (Ultima 216 Gold LLT, UK). Inert organics (dioxane, dextrose PEG 400, PEG 600, PEG 100) were measured with a Total Organic Carbon (TOC)  $V_{CPH/CPN}$  Shimadzu analyser with the high sensitivity catalyst 217 218 in a non-purgeable organic carbon mode. Calibration standards were made using potassium 219 hydrogen phthalate (Acros Organics, UK). Samples were analysed for TOC the same day of the 220 experiments. HA concentration was determined by ultraviolet absorbance at 254 nm measured with 221 an UV Visible Spectrophometer Carv 100 (Varian, UK). Samples were analysed within a few hours 222 from collection and stored in the dark at controlled room temperature. Conductivity and pH 223 measurements were conducted for the feed, permeate and concentrate samples using a pH/Cond 224 340i meter (WTW, Germany).
- 225

### 2.5 Calculation of HA-water partitioning coefficient K<sub>HA</sub> 226

227 HA-water partitioning coefficient K<sub>HA</sub> for ES was determined as a function of pH and HA 228 concentration using the negligible-depletion solid phase micro-extraction (nd-SPME) method 229 developed by Neale et al. (2008). The SPME method allows estimating the fraction of micropollutant bound to NOM, irrespective of the specific interactions (e.g. hydrophobic, hydrogen 230 231 bonding, etc.) between micropollutant and NOM. The method measures the freely dissolved

232 concentration of a micropollutant in water at equilibrium and it is based on the valid assumption

233 that only the freely dissolved micropollutant and not the fraction bound to the organic matter is

- 234 measured by the SPME fibre (Poerschmann and Kopinke 2001).
- 235
- 236 The SPME fibres were calibrated to establish the time necessary to reach equilibrium and calculate 237 the fibre-water partition coefficient  $K_{fw}$  for ES as a function of pH. Seven polyacrylate (PA) fibres (Polymcro Technology, US) of 5 cm length each were introduced in 100 mL solutions containing 238 239 10, 25, 50, 100 µg/L of ES and background electrolyte (1 mM NaHCO<sub>3</sub> and 20 mM NaCl) with pH varying from 4 to 12. The fibres were shaken at 200 rpm at 25 °C in a Certomat BS-1 incubator 240 shaker (Sartorius, Germany) and extracted at regular time intervals. The fibres were cut into three 241 242 pieces, added to a scintillation vial with 7 mL of scintillation liquid, manually shaken and left 243 overnight to desorb before being analysed by scintillation counter to determine m<sub>F</sub>, the mass of ES 244 in the fibre.

246 A kinetic approach was used to quantify K<sub>fw</sub>:

247 
$$K_{fw} = \frac{C_F}{C_W} = \frac{m_F}{m_W} \cdot \frac{V_W}{V_F}$$
(1)

where  $C_F$  (µg/L) is the concentration of ES in the fibre,  $C_W$  (µg/L) is the concentration of freely dissolved ES in aqueous solution,  $m_w$  (µg) is the mass of freely dissolved ES in aqueous solution as sampled at the end of the experiments,  $V_w$  (L) is the volume of the aqueous solution,  $V_F$  (L) is the volume of the fibre (0.77 µL). Log K<sub>fw</sub> was calculated from the linear regression of logC<sub>F</sub> as a function of logC<sub>w</sub> obtained with the four different concentrations of ES in solution.

253

In order to determine the HA-water partitioning coefficient  $K_{HA}$  for ES as a function of pH and HA concentration, 100 mL solutions containing 10, 25, 50, 100 µg/L of ES, 5, 12.5, 50, 125, 250 mgC/L of HA and background electrolyte were prepared at pH varying from 4 to 12 and shaken for 24 De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

- 257 hours at 200 rpm to allow complex formation. Fibres were then introduced in the solution and
- shaken for 96 hours, chosen as equilibrium time. Following, the fibres were extracted, cut into three
- 259 pieces, added to a scintillation vial with 7 mL of scintillation liquid, manually shaken and left
- 260 overnight to desorb before being analysed by scintillation counter to determine m<sub>F</sub>.
- 261 In the presence of HA,  $m_w$  was determined from the ES-fibre partitioning coefficient,  $K_{fw}$ , obtained
- 262 during fibre calibration as follows:

263 
$$m_W = \frac{m_F}{K_{fw}} \cdot \frac{V_W}{V_F}$$
(2)

264 K<sub>HA</sub> (L/kg) was then calculated as:

265 
$$K_{HA} = \frac{c_{HA}}{C_W} = \frac{m_{HA}}{m_{DOM}} \cdot \frac{V_W}{m_W}$$
 (3)

- where  $c_{HA}$  (µg/kg) is the concentration of ES sorbed to HA,  $m_{HA}$  (µg) is the mass of ES sorbed to
- 267 HA, m<sub>DOM</sub> (kg) is the total mass of HA dissolved in solution.
- 268
- 269 In order to estimate m<sub>HA</sub> a full mass balance was carried out:
- 270  $m_{HA} = m_{TOT} m_F m_W$  (4)
- 271 where m<sub>TOT</sub> is the initial mass of ES in solution before the fibre is added.
- 272 The fraction of ES bound to HA  $f_{HA}$  (%) was determined as (Neale and Schäfer 2012):

273 
$$f_{HA} = \frac{1}{\frac{V_W}{m_{DOM} \cdot K_{HA}} + 1}$$
(5)

- 274 The error in the determination of  $K_{fw}$ ,  $K_{HA}$  and  $f_{HA}$  was calculated from the instrumental error in the
- 275 determination of m<sub>F</sub>, m<sub>W</sub>, m<sub>TOT</sub> and m<sub>DOM</sub> using the error propagation rule (Miller and Miller 2000).
- 276

277 **3 Results and discussion** 

### 278 3.1 ES-HA complex formation

The formation of ES-HA complexes was quantified as a function of pH and HA concentration in order to evaluate the influence of solute-solute interactions on ES retention in the presence of HA. The HA-water partition coefficient  $K_{HA}$  for ES was determined using the nd-SPME method described in Section 2.5 and the fraction of ES bound to HA,  $f_{HA}$  hence complex formation, was calculated with equation 5. In order to determine  $K_{HA}$ , the SPME fibres were calibrated to obtain the fibre-water partition coefficient  $K_{fw}$  for ES as a function of pH.

285

### 286 3.1.1. Fibre calibration and determination of fibre partitioning coefficient

287 SPME has been largely employed as a technique for extracting organic micropollutants from 288 aqueous solutions and values of  $K_{fw}$  for  $\alpha$ -ES and  $\beta$ -ES are available in the literature (Valor et al. 289 2001). However, there are no data published on the variation of K<sub>fw</sub> with pH for ES. Moreover, the 290 free-standing fibre method used in this study differs from the traditionally employed techniques 291 using fibre holders. Fibre calibration was therefore required to establish equilibrium time and 292 determine K<sub>fw</sub>. PA fibres reached equilibrium after 48 hours and 96 hours were chosen for determination of the partitioning coefficients. No adsorption of ES to the glass was measured in 293 294 control samples.

295

PA fibre uptake was constant with pH up to pH 8 and then decreased rapidly (Figure 1). Since ES does not dissociate, this was attributed to the hydrolysis of α-ES and β-ES to ES-diol, which has lower log K<sub>ow</sub> and therefore less affinity to the fibres (DiFilippo and Eganhouse 2010). However, the decrease at pH 10 and 12 can be considered negligible once the logarithm is calculated, since the difference was less than 0.5 log units, and log K<sub>fw</sub> for ES can be considered constant with pH (Figure 1). For the pH range studied the negligible-depletion condition  $K_{fw} \cdot \frac{V_F}{V_W} <<<1$  was fulfilled

13

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

- 302 (Vaes et al. 1997). Values of log  $K_{fw}$  obtained for ES are in agreement (maximum difference 0.3
- 303 log units) with the values obtained in the literature for PA coated fibres at neutral pH (Valor et al.
- 304 2001), confirming the validity of the nd-SPME technique employed in this study.
- 305
- 306 (FIGURE 1)
- 307
- 308
- 309 3.1.2. Determination of HA partition coefficient and percentage of ES bound to HA

310	The HA partition coefficient for ES, $K_{HA}$ , was estimated as a function of pH and HA concentration.
311	While $K_{oc}$ values for ES have been determined for various soils (Hillenbrand et al. 2006, Ministerio
312	de Agricultura Pesca Y Alimentacion 1999, Organization for Economic Cooperation and
313	Development 1995), there are no published data on partitioning coefficient of ES and aquatic
314	organic matter. Values obtained for soil organic carbon cannot be used for aquatic HA as partition
315	coefficients in water could be 2-3 orders of magnitude greater (Devitt and Wiesner 1998). It is
316	fundamental to quantify the partition coefficients at the same conditions used during the
317	experiments with membranes as environmental conditions have shown to influence considerably the
318	K <sub>HA</sub> values (DiFilippo and Eganhouse 2010).

319

As shown in Figure 2, the partitioning coefficient for ES between HA and water and the fraction of ES bound to HA can be considered constant with pH. Albeit HA deprotonates at pH above 4, the complexation of ES and HA did not decrease at high pH as ES is neutral in the whole pH range, in agreement with findings for other organochloride pesticides (Prosen et al. 2002). Log  $K_{HA}$ decreased with increasing HA concentration indicating that complexation is limited by the mass of ES available. The percentage of ES bound to HA, calculated with equation 5, increased with HA concentration.

328

### (FIGURE 2)

329

### 330 3.2 Influence of pH and HA concentration on ES retention

The influence of HA on the removal of ES for different micropollutant MW/membrane MWCO 331 332 ratio was investigated. Firstly, ES retention was determined with and without HA for TFC-SR2 and 333 TFC-SR3 as a function of pH and secondly, ES retention was determined for the TFC-SR2 and TFC-SR3 as a function of HA concentration. The study of ES retention as a function of pH is 334 335 important because, while pH was shown not to influence ES-HA complex formation and the 336 percentage of ES bound to HA (Figure 2a), it can influence membrane charge within the pores (Braghetta et al. 1997, Dalwani et al. 2011, 2010). ES retention in the presence of HA was expected 337 338 to be affected by HA concentration since HA concentration influenced the formation of ES-HA 339 complexes and the percentage of ES bound to HA (Figure 2b).

340

### 341 <u>3.2.1. Influence of pH on ES retention</u>

The permeate flux  $J_v/J_o$  ratio, an indicator of membrane fouling, the ES mass adsorption on the membrane and the ES retention were determined for a pH range between 4 and 12, as shown in Figure 3. The use of new membrane coupons with pure water flux varying ±15%, resulted in ±8% variability in retention results and ±0.02 µg/cm<sup>2</sup> variability in adsorption results as determined by repeating some experiments in duplicate and triplicate with different membrane coupons. The determined variability was used for all the experiments in this study.

348 349

(FIGURE 3)

350

Figure 3a shows that, for both membranes,  $J_v/J_o$  was constant with pH when ES was filtered alone, showing no effect of pH on the membrane permeate flux. When ES and HA were both filtered, fouling (i.e.  $J_v/J_o$  decline) was not observed for both membranes when compared to  $J_v/J_o$  with ES De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

- 354 only. For the "tighter" TFC-SR3 (Table 2) the presence of HA did not influence the flux ratio, while
- 55 for the "looser" TFC-SR2 (Table 2) HA increased the flux ratio at pH 6, 8 and 10 up to 40%.
- 356 Increased permeate flux in the presence of NOM at pH > 4 has been previously observed for
- 357 "loose" NF membranes (Braghetta et al. 1997, Schäfer et al. 2010). Since NOM enhances the
- 358 negative charge of the NF membranes, which increases the electrostatic repulsion within the
- 359 membrane pores, it was inferred that NOM might increase membrane MWCO (Braghetta et al.
- 360 1997, Childress and Elimelech 1996).
- 361 362
  - (TABLE 2)
- 363

ES adsorption onto the TFC-SR3 was low (average over pH: 0.01  $\mu$ g/cm<sup>2</sup>) and independent of pH and the presence of HA (Figure 3b). ES adsorption was higher (average over pH: 0.03  $\mu$ g/cm<sup>2</sup>) for the more hydrophobic TFC-SR2 (Table 2) compared to the TFC-SR3. At pH>8, ES adsorption decreased with and without HA for the TFC-SR2. In this pH range ES hydrolyses to ES-diol, which is less hydrophobic than  $\alpha$ -ES and  $\beta$ -ES (Table 1) and might adsorb less on the membrane (Banasiak 2009).

370

For both membranes in the absence of HA ES retention varied with pH (Figure 3c). It is inferred that the variation of membrane surface characteristics with pH (i.e., negative surface charge increase with increase of pH) combined with hydrolization of ES isomers to ES-diol, which has a lower molecular weight and lower hydrophobicity, decreased ES retention by the TFC-SR3 membrane with increasing pH. The same occurred for the TFC-SR2 membrane with the exception of pH 12. At this pH it is possible that some of the ES diol is dissociated (pKa value in Table 1) which might have an enhanced retention by the negatively charged membrane.

378

For TFC-SR3 (micropollutant MW/membrane MWCO ratio > 1) HA increased ES retention at every pH. This increase in retention is not caused by pore clogging from HA, since  $J_v/J_o$  did not decrease when HA was present compared to filtration without HA (Figure 3a). A possible explanation for the increase in ES retention is the formation of ES-HA complexes: as HA is highly retained by the NF membrane, ES molecules that complex with HA will also be retained. The influence of complexation on ES retention will be studied in detail in the next section.

In contrast, ES retention in the presence of HA decreased at pH 6, 8 and 10 and was similar at pH 4 386 and 12 for the TFC-SR2 (micropollutant MW/membrane MWCO ratio < 1). Decrease in ES 387 retention in the presence of HA was observed for the "loose" membrane, confirming the inferred 388 389 correlation between micropollutant MW/membrane MWCO ratio and micropollutant removal in the 390 presence of NOM. Decreased ES retention in the presence of HA at pH 6, 8 and 10 for TFC-SR2 391 corresponded to an increase in flux ratio (Figure 3a), as observed in previous studies (Boussahel et 392 al. 2002, Schäfer et al. 2010), and to an increase in ES adsorption (Figure 3b). At pH 12, the flux 393 was unchanged (Figure 3a) and the ES adsorption on the membrane was lower, translating into an 394 increased retention of ES.

395

Two mechanisms of ES removal in the presence of HA can be identified so far: (1) for TFC-SR3 the presence of HA increased ES retention, potentially caused by the formation of ES-HA complexes (2) for TFC-SR2 the presence of HA decreased ES retention and increased  $J_v/J_o$  for certain pH conditions (pH 6 to 10), indicating a change of membrane performance when HA is present, whilst for others (pH 4 and 12) it increased ES retention, potentially caused by the formation of ES-HA complexes.

402

403 3.2.2. Influence of HA concentration on ES retention

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

404 In order to carry out a more in depth investigation of the role of ES-HA complexes on ES removal,

- 405 retention of ES as a function of HA concentration was studied. For both membranes ES retention as
- 406 a function of HA concentration was studied at pH 4 because membranes and HA are neutral and
- 407 charge effects are minimal. ES retention was also investigated at pH 8 for the TFC-SR2, where the
- 408 presence of HA was found to decrease ES retention (Figure 3c). ES retention as a function of HA
- 409 concentration at pH 8 was not studied for the TFC-SR3 as ES retention in the presence of HA is
- 410 independent of pH (Figure 3c).
- 411

412 At pH 4, flux ratio decline occurred for both membranes for HA > 50 mgC/L (Figure 4a). For these experimental conditions the high concentrations of HA caused the membrane to foul by pore 413 414 blocking (Xu et al. 2005). In this case, ES retention can be affected by pore blocking, competition 415 of adsorption sites between HA deposits and ES (Yoon et al. 2006), and interactions between ES 416 and the fouling layer. The complexity of the mechanisms when the HA fouling layer is present does 417 not allow the study of the impact of ES-HA complexes on ES retention. Therefore, the mechanisms 418 of ES retention in the presence of HA when the HA fouling layer is present will not be discussed in 419 detail further since an investigation of the impact of fouling mechanisms in micropollutant retention 420 is outside the scope of this study. For HA  $\leq$  50 mgC/L there was no effect on the flux ration J<sub>y</sub>/J<sub>0</sub> for 421 both TFC-SR2 and TFC-SR3. 422

424

423

For the TFC-SR2 at pH 8, the flux ratio increased from 0.97 to 1.21 for HA > 5 mgC/L confirming what was observed previously in Figure 3a. Some studies in the literature have reported an increase in permeate flux after NOM filtration and inferred that it could be linked to increase in MWCO (Braghetta et al. 1997, Childress and Elimelech 1996, Xu et al. 2005, Xu et al. 2006). The influence of HA filtration on MWCO will be studied in detail in the next section.

(FIGURE 4)

430

431	ES adsorption was independent of HA concentration for TFC-SR3, while for TFC-SR2 it increased
432	with HA concentration up to HA 50 mgC/L and then decreased for a concentration higher than 50
433	mgC/L (Figure 4b). The decrease in ES adsorption for HA > 50 mgC/L might be caused either by a
434	higher percentage of ES bound to HA (Figure 2 b) leaving less freely dissolved ES to adsorb on the
435	membrane or due to fouling of the membrane by HA (Figure 4 a) which competes for adsorption
436	sites on the membrane (Yoon et al. 2006). An increase in ES adsorption for $HA > 5 mgC/L$ can be
437	caused by the inferred increase in the MWCO of the membrane due to HA filtration.
438	Micropollutants have been shown to adsorb inside the NF membrane pores (Kimura et al. 2003,
439	Semião and Schäfer 2012). Hence whilst HA fouling blocks the pores and does not allow for the ES
440	to adsorb inside the TFC-SR2, an increase of MWCO increases the internal area for more ES
441	adsorption to occur.

442

At pH 4 ES retention increased with HA concentration for the looser TFC-SR2 from 60.0% to 84.4% while it was constant for the tighter TFC-SR3 with a retention of 96.0%. (Figure 4c). For TFC-SR2 at pH 8 ES retention decreased with HA concentration up to HA 50 mgC/L from 55% to 30%, confirming what was observed for HA 12.5 mgC/L at pH 6, 8 and 10 (Figure 3c).

447

454

These results confirmed that the presence of HA increased the retention of pesticide ES for TFC-SR3, whose MWCO is smaller than ES MW, and decreased ES retention for TFC-SR2 at pH 8, whose MWCO is bigger than ES MW. However, ES retention by TFC-SR2 was found to increase at pH 4 with increase of HA concentration. HA concentration and pH played a role in the different ES retention in the presence of HA. HA concentration influenced ES-HA interactions and pH affected HA-membrane interactions. De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

- 455 ES adsorption on the membranes (ES-membrane interactions) was not believed to influence ES
- 456 retention in the presence of HA. ES adsorption values on TFC-SR2 were similar at pH 4 and at pH
- 457 8 (Figure 4b) for HA concentrations <12.5 mgC/L while ES retention in the presence of HA
- 458 increased at pH 4 and decreased at pH 8 (Figure 4c). To elucidate the mechanisms responsible for
- 459 increase/decrease of ES retention in the presence of HA, the contributions of solute-solute
- 460 interactions (ES-HA complexes) and solute-membrane (HA-membrane) interactions on ES removal
- 461 have been investigated and described in Section 3.3.
- 462

### 463 3.3 Mechanisms of ES retention in presence of HA

In order to quantify the contribution of ES-HA interactions on ES removal, ES retention in the 464 465 presence of HA was estimated from the calculated partition coefficient,  $K_{HA}$ , and the fraction of ES 466 bound to HA,  $f_{HA}$ . The approach developed by Neale and Schäfer (2012) for UF was modified for 467 NF. Neale and Schäfer (2012) determined hormone retention in the presence of HA as proportional to the retention of HA and to the fraction of micropollutant bound to HA only, due to the low 468 469 retention of hormones by UF membranes. In the case of NF, micropollutant retention cannot be 470 ignored and the method was modified and improved by carrying out a mass balance in order to take this term into account. 471 472 473 Some assumptions were made to determine the fraction of ES bound to HA and therefore estimate

- 474 retention due to complex formation:
- 475 (1) The total mass of ES in solution ( $m_{ES}$ ) is either freely dissolved ( $m_{fES}$ ) or complexed to HA
- 476  $(m_{ES-HA})$  after the membrane saturation has been reached:
- $m_{\rm ES} = m_{\rm ES-HA} + m_{\rm fES} \quad (6)$
- 478 (2) ES complexed with HA is retained together with the HA to which it is partitioned
- 479 (3) ES freely dissolved in HA solution has the same retention as ES during experiments without
- 480 HA carried out at the same conditions of pH, pressure and background electrolyte.

481

While the first two assumptions can be considered generally valid, the last assumption will not be true if the presence of HA influences the retention of freely dissolved ES. HA-membrane interactions can influence the retention of freely dissolved ES in the HA solution and result in different overall ES retention than the estimated retention. HA can adsorb to the membrane surface resulting in pore blocking (Xu et al. 2005) and increasing membrane charge (Hong and Elimelech 1997, Xu et al. 2005), membrane MWCO (Xu et al. 2006) and membrane hydrophilicity (Hong and Elimelech 1997, Schäfer et al. 2010).

489

490 The total ES retention due to the retention of ES complexed with HA and freely dissolved ES has 491 been estimated by mass balance from the total mass of ES in the feed ( $m_{ESfeed}$ ) and in the permeate 492 ( $m_{ESperm}$ ):

493  $m_{\text{ESfeed}} = m_{\text{ES-HAfeed}} + m_{\text{fESfeed}} = f_{HA} \cdot m_{\text{ESfeed}} + (1 - f_{HA}) \cdot m_{\text{ESfeed}}$ (7)

494  $m_{\text{ESperm}} = m_{\text{ES-HAperm}} + m_{\text{fESperm}} = (1 - R_{\text{HA}}) \cdot m_{\text{ES-HAfeed}} + (1 - R_{\text{ES}}) \cdot m_{\text{fESfeed}}$ (8)

495

where  $R_{HA}$  is the retention of HA and  $R_{ES}$  is the retention of ES during experiments without HA. For TFC-SR3,  $R_{HA}$  was 99% and could be considered constant with pH and HA concentration (± 2% variability); for TFC-SR2, the average HA retention over the pH was 88%, increasing from 75.8% at pH 4 to 98.5% at pH 12 and from 71.5% for HA concentration 5 mgC/L to 99.1% for HA 250 mgC/L.

501

502 The retention estimated with this methodology is indicated with a dotted line in Figure 3c and 503 Figure 4c, and was compared with the retention obtained during ES-HA experiments. Similar 504 experimental and estimated retention would indicate that solute-membrane interactions are 505 negligible (i.e. the third hypothesis is verified) and solute-solute interactions are dominant. When De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

506 solute-membrane interactions are important the retention predicted from K<sub>HA</sub> is expected to be

- 507 different from the retention obtained experimentally. This methodology depends on the fraction of
- 508 micropollutant bound to NOM, on micropollutant retention and on NOM retention and it can be
- 509 applied to any micropollutant that forms complexes with NOM to estimate the contribution of
- 510 micropollutant-NOM interactions on micropollutant retention by NF, irrespective of the type of
- 511 interactions and the compound properties.
- 512

513 As shown in Figure 3c, ES retention in the presence of HA estimated with K<sub>HA</sub> was similar 514 (maximum 10% difference) to the experimental retention for the TFC-SR3, showing that for this 515 membrane solute-solute interactions (i.e. the formation of ES-HA complexes) are the dominant 516 mechanism affecting ES retention. For TFC-SR2 the estimated retention was higher than the 517 experimental retention for all the pH range studied, with the exception of pH 4 and 12, indicating 518 that solute-membrane interactions (i.e. HA-membrane interactions) played a role. For both 519 membranes at pH 4 the estimated ES retention was similar to the experimental retention (Figure 4c), 520 indicating that at this pH solute-solute interactions are the dominant mechanism.

521

522 For TFC-SR2 at pH 8 ES retention obtained experimentally was lower than the estimated ES 523 retention for any HA concentration (Figure 4c). At this pH the role of solute-membrane interactions 524 on ES removal was more predominant than the formation of ES-HA complexes. The U-shape of the 525 retention curve indicated that solute-membrane interactions were dominant for lower HA 526 concentrations while at high HA concentrations both ES-HA interactions and pore blocking by HA 527 fouling (evidenced from a decrease in permeate flux in Figure 4a) became more important. However, even at high HA concentrations ES retention was lower than the estimated retention 528 529 (dotted line) since solute-membrane interactions could not be considered negligible.

531	In order to establish how solute-membrane interactions affected ES retention in the presence of HA,
532	the characteristics of both membranes were evaluated for membranes pre-filtered with 900 mL of
533	ultra-pure water and background electrolyte and membranes pre-filtered with 900 mL of solution
534	containing 12.5 mgC/L of HA and background electrolyte at pH 8. Membrane MWCO and contact
535	angle were determined in order to evaluate change in membrane pore size and hydrophilicity caused
536	by the presence of HA (Table 3).
537	
538	(TABLE 3)
539	
540	HA deposited preferentially on the TFC-SR3, forming a visible yellowish layer on the membrane,
541	while visible HA deposits were not observed on the TFC-SR2. This different behaviour does not

542 seem to be linked with the different roughness of the two membranes (Table 2), as the membrane 543 with the highest roughness had no visible deposits on its surface. MWCO increased in the presence 544 of HA for both membranes, confirming what was inferred in previous studies (Xu et al. 2005, Xu et 545 al. 2006). The increase in MWCO for TFC-SR3 did not influence ES retention as the higher MWCO (179 g/mol) was still lower than ES MW (407 g/mol). In the case of TFC-SR2, for which 546 547 micropollutant MW/membrane MWCO ratio < 1 and ES is partially retained, the increase in MWCO from 460 to 496 g/mol decreased the ratio further, in turn decreasing ES retention. 548

549

550 After HA were filtered the observed contact angle decreased for TFC-SR2 and increased for TFC-SR3 (Table 3). For membranes fouled by NOM, contact angle measurements have been shown to 551 552 be representative of the fouling layer: in the case of fouling by HA, contact angles indicated the 553 adhesion of a layer of intermediate hydrophobicity (Jucker and Clark 1994, Lee et al. 2004, Palacio 554 et al. 1999, Xu et al. 2006). For TFC-SR3 the contact angle is thought to be determined by the HA laver visibly deposited on the membrane after filtration; the roughness of the HA deposits were also 555 556 likely to increase the observed contact angle (de Gennes 1985). For TFC-SR2, for which HA

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", Water Research 47(10):3484-3496, doi:10.1016/i.watres.2013.03.055

557 deposits were not visible after filtration, the contact angle is thought to reflect membrane

- 558 hydrophilization due to filtration of charged HA (Hong and Elimelech 1997). Increase in
- 559 hydrophilization can also explain the observed increase in permeate flux (Figure 4a).
- 560

579

580

561 A schematic of the proposed mechanisms is presented in Figure 5. At pH 4, when membranes and 562 HA are neutrally charged, size exclusion and solute-solute interactions (i.e. ES-HA interactions) 563 dominated ES retention. At neutral pH, both HA and the membranes are negatively charged and solute-membrane interactions (i.e. HA-membrane interactions) influence membrane MWCO and 564 565 hydrophilicity. The filtration of HA increased the MWCO for both membranes (Table 3), but different trends were obtained. For TFC-SR3, whose micropollutant MW/membrane MWCO ratio 566 567 > 1, the small increase in MWCO from 165 to 179 g/mol was not sufficient to allow for increase in 568 ES passage. This low HA impact in MWCO increase was further evidenced by a lack of increase in 569 permeate flux (Figure 3 a and Figure 4 a). For TFC-SR3, the formation of ES-HA complexes was 570 the dominant mechanism responsible for the increase of ES retention and it overcame the increase 571 in MWCO (Figure 3 c). For TFC-SR2, interactions between charged HA and the membrane also 572 resulted in an increase of the MWCO (Table 3), which is further evidenced from an increase in 573 permeate flux (Figure 3 a and Figure 4 a). As TFC-SR2 has a micropollutant MW/membrane MWCO < 1, a further increase in the MWCO had a high impact in ES retention by decreasing it 574 575 considerably. For TFC-SR2, HA-membrane interaction was the dominant mechanism responsible 576 for the decrease of ES retention when HA are present since the formation of HA-ES complexes did not overcome the increase of membrane MWCO. 577 578

(FIGURE 5)

581 **4 Conclusions** 

582 The mechanisms involved in the removal of micropollutants by NF membranes in the presence of

583 NOM were identified and evaluated in this study. For the first time, the contribution of both solute-

solute interactions and solute-membrane interactions were evaluated to explain micropollutant

585 retention mechanisms.

586

587 Results showed that in the absence of HA-membrane interactions, ES-HA complexation was the 588 governing mechanism in ES removal, enhancing ES removal. ES-membrane interactions did not 589 contribute to the variation of ES retention in the presence of HA. In contrast, HA-membrane 590 interactions affected membrane performance by increasing the MWCO and decreasing ES retention. 591

592 This work improves the understanding of the fundamentals involved in the removal mechanisms by 593 NF and demonstrates that the evaluation of the contributions of both solute-solute interactions and 594 solute-membrane interactions is fundamental for understanding micropollutant removal 595 mechanisms by NF

### 596 Acknowledgments

597 The authors would like to thank Koch Membranes Systems for kindly providing the membranes 598 used during the experiments, Dr. Alexander Bismarck and Dr. Kingsley Ho, Imperial College, UK, for assistance with zeta potential measurements, Dott. Flaminia Gallo and Dr. Alberto Figoli ITM-599 600 CNR, Italy, for contact angle measurements, Ime Akanaveti and Dr. Nhan T Pham, The University of Edinburgh, UK, for AFM measurements, Dr. Peta Neale, The University of Queensland, for 601 602 useful discussions and proof reading and Dr. Helfrid Schulte-Herbrüggen, The University of 603 Edinburgh, for paper proof reading. Special thanks are given to the Drinking Water Quality 604 Regulator for Scotland (DWQR) and the School of Engineering of The University of Edinburgh for 605 funding the PhD studentship for De Munari.

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

607 References

Agbekodo, K.M., Legube, B. and Dard, S. (1996) Atrazine and simazine removal mechanisms by
nanofiltration: Influence of natural organic matter concentration. Water Research 30(11), 25352542.

611 Banasiak, L.J. (2009) Removal of inorganic and trace organic contaminants by electrodialysis. PhD

612 Thesis, The University of Edinburgh.

613 Bellona, C., Drewes, J.E., Xu, P. and Amy, G. (2004) Factors affecting the rejection of organic

614 solutes during NF/RO treatment - a literature review. Water Research 38(12), 2795-2809.

Boussahel, R., Montiel, A. and Baudu, M. (2002) Effects of organic and inorganic matter on
pesticide rejection by nanofiltration. Desalination 145(1-3), 109-114.

Braghetta, A., Di Giano, F.A. and Ball, W.B. (1997) Nanofiltration of natural organic matter: pH
and ionic strength effects. Journal of Environmental Engineering 123(7), 628-641.

619 Childress, A.E. and Elimelech, M. (1996) Effect of solution chemistry on the surface charge of 620 polymeric reverse osmosis and nanofiltration membranes. Journal of Membrane Science 119(2),

621 253-268.

622 Chin, Y., Aiken, G. and O'Loughlin, E. (1994) Molecular weight, polydispersity, and spectroscopic

623 properties of aquatic humic substances. Environmental Science and Technology 28(11), 1853-1858.

624 Cyna, B., Chagneau, G., Bablon, G. and Tanghe, N. (2002) Two years of nanofiltration at the Méry625 sur-Oise plant, France. Desalination 147(1–3), 69-75.

626 Dalwani, M., Benes, N.E., Bargeman, G., Stamatialis, D. and Wessling, M. (2011) Effect of pH on

627 the performance of polyamide/polyacrylonitrile based thin film composite membranes. Journal of

628 Membrane Science 372(1–2), 228-238.

Dalwani, M., Benes, N.E., Bargeman, G., Stamatialis, D. and Wessling, M. (2010) A method for
characterizing membranes during nanofiltration at extreme pH. Journal of Membrane Science
363(1–2), 188-194.

de Gennes, P.G. (1985) Wetting: statics and dynamics. Reviews of Modern Physics 57(3), 827-863.

De Munari, A. (2012) Removal mechanisms of organic and inorganic solutes in raw, upland
drinking water by nanofiltration: Influence of solute-solute and solute-membrane interactions. PhD
Thesis, The University of Edinburgh.

636 Devitt, E.C., Ducellier, F., Côté, P. and Wiesner, M. (1999) Effects of natural organic matter and
637 the raw water matrix on the rejection of atrazine by pressure-driven membranes. Water Research
638 32(9), 2563-2568.

639 Devitt, E.C. and Wiesner, M.R. (1998) Dialysis investigations of atrazine-organic matter
640 interactions and the role of a divalent metal. Environmental Science and Technology 32(2), 232641 237.

DiFilippo, E.L. and Eganhouse, R.P. (2010) Assessment of PDMS-water partition coefficients:
Implications for passive environmental sampling of hydrophobic organic compounds.
Environmental Science and Technology 44(18), 6917-6925.

Forman, S., Durbetaki, A., Choen, M. and Olofson, R.A. (1965) Conformational equilibria in cyclic
sulfites and sulfates. The configurations and conformations of the two isomeric thiodans. Journal of
Organic Chemistry 30(1), 169-175.

German Federal Environment Agency (2007) Draft dossier prepared in support of a proposal of
endosulfan to be considered as a candidate for inclusion in the annexes of the Stockholm
Convention.

Hajibabania, S., Verliefde, A., McDonald, J.A., Khan, S.J. and Le-Clech, P. (2001) Fate of trace
organic compounds during treatment by nanofiltration. Journal of Membrane Science 373(1-2),
130-139.

Hansch, C., Leo, A. and Hoekman, D. (1995) Exploring QSAR: hydrophobic, electronic and steric
constants, American Chemical Society, Washington DC, USA.

Hillenbrand, T., Marscheider-Weidemann, F., Strauch, M. and Heitmann, K. (2006) PrioritäreStoffe der Wasserrahmenrichtlinie Datenblatt Nr. 14: Endosulfan (in German).

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

- Hong, S. and Elimelech, M. (1997) Chemical and physical aspects of natural organic matter (NOM)
- fouling of nanofiltration membranes. Journal of Membrane Science 132(2), 159-181.
- 660 Jin, X., Hu, J. and Ong, S.L. (2007) Influence of dissolved organic matter on estrone removal by NF
- membranes and the role of their structures. Water Research 41(14), 3077-3088.
- 662 Jin, X., Hu, J. and Ong, S.L. (2010) Removal of natural hormone estrone from secondary effluents
- using nanofiltration and reverse osmosis. Water Research 44(2), 638-648.
- 664 Jucker, C. and Clark, M.M. (1994) Adsorption of aquatic humic substances on hydrophobic
- 665 ultrafiltration membranes. Journal of Membrane Science 97, 37-52.

Kimura, K., Amy, G., Drewes, J. and Watanabe, Y. (2003) Adsorption of hydrophobic compounds
onto NF/RO membranes: an artifact leading to overestimation of rejection. Journal of Membrane
Science 221(1-2), 89-101.

- 669 Lee, N., Amy, G., Croué, J.-P. and Buisson, H. (2004) Identification and understanding of fouling
- in low-pressure membrane (MF/UF) filtration by natural organic matter (NOM). Water Research38(20), 4511-4523.

Miller, J.N. and Miller, J.C. (2000) Statistics and chemometrics for analytical chemistry, PearsonEducation Limited, Harlow.

674 Ministerio de Agricultura Pesca y Alimentacion (1999) Endosulfan - Vol 1 - Monograph prepared

in the context of the inclusion of the following active substance in Annex I of the Council Directive91/414/EEC.

- 677 Neale, P.A., Escher, B.I. and Schäfer, A.I. (2008) Quantification of solute-solute interactions using
- 678 negligible-depletion solid-phase microextraction: measuring the affinity of estradiol to bulk organic
- 679 matter. Environmental Science and Technology 42(8), 2886-2892.
- Neale, P.A. and Schäfer, A.I. (2012) Quantification of solute–solute interactions in steroidal
  hormone removal by ultrafiltration membranes. Separation and Purification Technology 90, 31-38.

682 Ng, H.Y. and Elimelech, M. (2004) Influence of colloidal fouling on rejection of trace organic

- 683 contaminants by reverse osmosis. Journal of Membrane Science 244(1-2), 215-226.
- 684 Nghiem, L.D., Schäfer, A.I. and Elimelech, M. (2004) Removal of natural hormones by
- nanofiltration membranes: Measurement, modeling, and mechanisms. Environmental Science andTechnology 38(6), 1888-1896.

687 Organization for Economic Cooperation and Development (1995) Environment Monograph No.

688 108: Final report on the OECD project to compare pesticides data reviews.

Palacio, L., Calvo, J.I., Prádanos, P., Hernández, A., Väisänen, P. and Nyström, M. (1999) Contact
angles and external protein adsorption onto UF membranes. Journal of Membrane Science 152(2),
189-201.

Peterson, S.M. and Batley, G.E. (1993) The fate of endosulfan in aquatic ecosystems.Environmental Pollution 82(2), 143-152.

Plakas, K.V. and Karabelas, A.J. (2009) Triazine retention by nanofiltration in the presence of
organic matter: The role of humic substance characteristics. Journal of Membrane Science 336(1–
2), 86-100.

Plakas, K.V., Karabelas, A.J., Wintgens, T. and Melin, T. (2006) A study of selected herbicides
retention by nanofiltration membranes - the role of organic fouling. Journal of Membrane Science
284(1-2), 291-300.

Poerschmann, J. and Kopinke, F.-D. (2001) Sorption of very hydrophobic organic compounds
(VHOCs) on dissolved humic organic matter (DOM). 2. Measurement of sorption and application
of a Flory-Huggins concept to interpret the data. Environmental Science and Technology 35(6),
1142-1148.

Prosen, H., Troha, A. and Zupančič - Kralj, L. (2002) Studies of interaction between some
organochlorine insecticides and humic acid using solid-phase microextraction and gas
chromatography. Acta Chimica Slovenica 49(3), 561-573.

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055

- 707 Salvaterra, A.F., Sarmento, G., Minhalma, M. and de Pinho, M.N. (2011) Nanofiltration of surface
- 708 water for the removal of endocrine disruptors. Desalination and Water Treatment 35(1-3), 54-61.
- 709 Sanches, S., Penetra, A., Granado, C., Cardoso, V.V., Ferreira, E., Benoliel, M.J., Barreto Crespo,
- 710 M.T., Pereira, V.J. and Crespo, J.G. (2011) Removal of pesticides and polycyclic aromatic
- 711 hydrocarbons from different drinking water sources by nanofiltration. Desalination and Water
- 712 Treatment 27(1-3), 141-149.
- 713 Schäfer, A.I., Nghiem, L.D., Meier, A. and Neale, P.A. (2010) Impact of organic matrix compounds
- 714 on the retention of steroid hormone estrone by a "loose" nanofiltration membrane. Separation and
- 715 Purification Technology 73(2), 179-187.

Schwarzenbach, R.P., Escher, B.I., Fenner, K., Hofstetter, T.B., Johnson, C.A., von Gunten, U. and
Wehrli, B. (2006) The challenge of micropollutants in aquatic systems. Science 313(5790), 10721077.

Semião, A.J.C. and Schäfer, A.I. (2011) Estrogenic micropollutant adsorption dynamics onto
 nanofiltration membranes. Journal of Membrane Science 381(1-2), 132-141.

Semião, A.J.C. and Schäfer, A.I. (2012) Removal of adsorbing estrogenic micropollutants by
 nanofiltration membranes. Part A - experimental evidence. Journal of Membrane Science 431: 244 256.

Shin, H.-S., Monsallier, J.M. and Choppin, G.R. (1999) Spectroscopic and chemical
characterizations of molecular size fractionated humic acid. Talanta 50(3), 641-647.

Soto, A., Chung, K. and Sonnenschein, C. (1994) The pesticide endosulfan, toxaphene and dieldrin
have estrogenic effects on human estrogen-sensitive cells. Environmental Health Perspectives
102(4), 380-383.

- Teixeira, M.R., Rosa, M.J. and Nyström, M. (2005) The role of membrane charge on nanofiltration
   performance. Journal of Membrane Science 265(1–2), 160-166.
- 731 Vaes, W.H.J., Ramos, E.U., Hamwijk, C., van Holsteijn, I., Blaauboer, B.J., Seinen, W., Verhaar,
- 732 H.J.M. and Hermens, J.L.M. (1997) Solid phase microextraction as a tool to determine

733 membrane/water partition coefficients and bioavailable concentrations in vitro systems. Chemical

734 Research in Toxicology 10(10), 1067-1072.

Valor, I., Perez, M., Cortada, C., Apraiz, D., Molto, J.C. and Font, G. (2001) SPME of 52 pesticides
and polychlorinated biphenyls: Extraction efficiencies of the SPME coatings
poly(dimethylsiloxane), polyacrylate, poly(dimethylsiloxane)-divinylbenzene,
carboxenpoly(dimethylsiloxane) and carbowaxdivinylbenzene. Journal of Separation Science 24(1),
39-48.

Van der Bruggen, B., Verliefde, A., Braeken, L., Cornelissen, E., Moons, K., Verberk, J., Van Dijk,
H. and Amy, G.L. (2006) Assessment of a semi-quantitative method for estimation of the rejection
of organic compounds in aqueous solution in nanofiltration. Journal of Chemical Technology and
Biotechnology 81(7), 1166-1176.

Xu, P., Drewes, J.E., Bellona, C., Amy, G.L., Kim, T.U., Adam, M. and Heberer, T. (2005)
Rejection of emerging organic micropollutants in nanofiltration–reverse osmosis membrane
applications. Water Environment Research 77(1), 40-48.

Xu, P., Drewes, J.E., Kim, T.-U., Bellona, C. and Amy, G. (2006) Effect of membrane fouling on
transport of organic contaminants in NF/RO membrane applications. Journal of Membrane Science
279(1-2), 165-175.

Yoon, Y., Westerhoff, P., Snyder, S.A. and Wert, E.C. (2006) Nanofiltration and ultrafiltration of
endocrine disrupting compounds, pharmaceuticals and personal care products. Journal of Membrane
Science 270(1-2), 88-100.

753

754

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055 List of Figures

Figure 1 Mass of ES in the nd-SPME fibre for different ES concentrations and Log  $K_{fw}$  values for ES as function of pH. ES 10, 25, 50, 100  $\mu$ g/L, background electrolyte 1 mM NaHCO<sub>3</sub> and 20 mM NaCl.

Figure 2 HA-water partition coefficient and fraction of ES bound to HA (a) as a function of pH, HA 12. 5 mgC/L and (b) as a function of HA concentration. ES 10, 25, 50, 100  $\mu$ g/L, background electrolyte 1 mM NaHCO3 and 20 mM NaCl.

**Figure 3** (a) Ratio of permeate flux  $J_v$  and initial pure water flux  $J_0$  (b) ES adsorption (c) and ES retention as a function of pH for TFC-SR2 and TFC-SR3. Dotted lines indicate ES retention in presence of HA estimated using the calculated partition coefficient K<sub>HA</sub>. ES 10 µg/L, HA 12.5 mgC/L, pressure 5 bar, background electrolyte 1 mM NaHCO3 and 20 mM NaCl.

**Figure 4 (a)** Ratio of permeate flux  $J_v$  and initial pure water flux  $J_0$  (**b**) ES adsorption and (**c**) ES retention as a function of HA concentration for TFC-SR2 and TFC-SR3. Dotted lines indicate ES retention in presence of HA estimated using the calculated partition coefficient K<sub>HA</sub>. ES 10 µg/L, pressure 5 bar, HA 5, 7.5, 12.5, 25 50, 125 250 mgC/L, background electrolyte 1 mM NaHCO<sub>3</sub> and 20 mM NaCl.

**Figure 5** Conceptual sketch (to scale) of main retention mechanisms for freely dissolved ES (MW 407 g/mol) and ES-HA complexes by TFC-SR2 and TFC-SR3. The presence of HA increases the MWCO for both membranes at pH 8 (changes indicated by the lighter area on the membranes). For TFC-SR3, MWCO increases from 165 to 179 g/mol but the passage of ES decreases due to the formation of ES-HA complexes: solute-solute interactions are the main mechanism of ES retention

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055 in the presence of HA. For TFC-SR2, MWCO increases from 460 to 496 g/mol causing an increase

in the passage of ES despite the formation of ES-HA complexes: solute-membrane interactions are

the dominant mechanism.

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055 List of Tables

 Table 1 – Characteristics of pesticide Endosulfan

Table 2 - Characteristics of TFC-SR2 and TFC-SR3

Table 3 – Membrane characteristics for a membrane pre-filtered with 900 mL of ultra-pure water and background electrolyte and a membrane through which 12.5 mgC/L of HA and background electrolyte have been filtered at pH 8

	a-ES	β-ES	Endosulfan diol
	$\begin{array}{c} Cl & H & H \\ H & & \\ Cl & & \\ H & \\ \end{array} = 0$	$\begin{array}{c} Cl \\ H \\ Cl \\ H \\ Cl \\ H \\ $	CI CI CI CI CI CI CI CI CI CI CI CI CI C
Formula	C <sub>9</sub> H <sub>6</sub> Cl <sub>6</sub> O <sub>3</sub> S	$C_9H_6Cl_6O_3S$	$C_9H_8Cl_6O_2$
Molecular Weight (g/mol)	406.93	406.93	360.88
Log K <sub>ow</sub>	3.83 (Hansch et al. 1995)	3.83 (Hansch et al. 1995)	3.68 (Banasiak 2009)
pKa	-	-	14.62–15.22 (Banasiak 2009)
Dipole Moment	1.02 (Forman et al. 1965)	3.18 (Forman et al. 1965)	Not available

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055 **Table 2** 

	Material	MWCO <sup>1</sup> (g/mol)	Estimated pore radius (nm)	Point of zero charge <sup>2</sup> (pH)	Contact Angle (°)	Roughness <sup>3</sup> (nm)
TFC-SR2	Polyamide on	460	0.52	4.25	57	17.9
TFC-SR3	polysulfone	165	0.38	3.84	44	5.2

<sup>1</sup>pressure 5 bar, neutral pH

<sup>2</sup>Electrolyte solution 1 mM NaHCO<sub>3</sub> and 20 mM NaCl

<sup>3</sup> Average roughness, R<sub>a</sub>

	TFC-SR2		TFC-SR3		
Pre-filtration Method	Ultra-pure water and	HA and electrolyte <sup>1</sup>	Ultra-pure water and	HA and electrolyte <sup>1</sup>	
	electrolyte <sup>1</sup>		electrolyte <sup>1</sup>		
MWCO (g/mol)	460	496	165	179	
Contact angle	57	49	44	59	

<sup>1</sup>Electrolyte solution 1 mM NaHCO<sub>3</sub> and 20 mM NaCl

De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055 Figure 1



### Figure 2



De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", Water Research 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055 Figure 3



Figure 4



De Munari, A., Semiao, A.J.C., Antizar-Ladislao, B. (2013) "Retention of pesticide Endosulfan by nanofiltration: Influence of organic matter-pesticide complexation and solute-membrane interactions", *Water Research* 47(10):3484-3496. doi:10.1016/j.watres.2013.03.055 Figure 5

