THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Biochar amendment improves alpine meadows growth and soil health in Tibetan plateau over a three year period

Citation for published version:

Rafiq, MK, Yanfu, B, Aziz, R, Rafiq, MT, Masek, O, Bachmann, RT, Joseph, S, Shahbaz, M, Qayyum, A, Zhanhuan, S, Danaee, M & Ruijun, L 2020, 'Biochar amendment improves alpine meadows growth and soil health in Tibetan plateau over a three year period', *Science of the Total Environment*, vol. 717.
<https://doi.org/10.1016/j.scitotenv.2019.135296>

Digital Object Identifier (DOI):

[10.1016/j.scitotenv.2019.135296](https://doi.org/10.1016/j.scitotenv.2019.135296)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Science of the Total Environment

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Biochar amendment improves alpine meadows growth and soil health in Tibetan plateau over a three year period

Muhammad Khalid Rafiq^{a,b,c,1,*} Bai Yanfu^{b,1} Raksanda Aziz^d, Muhammad Tariq Rafiq^e, Ondřej Mašek^a, Robert Thomas Bachmann^f, Stephen Joseph^g, Maqbool Shahbaz^c, Abdul Qayyum^h, Shang Zhanhuan^{b,*}, Mahmoud Danaee^g, Long Ruijun^b

^a UK Biochar Research Centre, School of GeoSciences, University of Edinburgh, Crew Building, King's Buildings, Edinburgh EH9 3FF, United Kingdom

^b State Key Laboratory of Grassland Agro-ecosystems, International Centre for Tibetan Plateau Ecosystem Management, School of Life Sciences, Lanzhou University, Lanzhou 730000, China

^c Rangeland Research Institute, National Agricultural Research Center, Islamabad 44000, Pakistan

^d Department of Environmental Science, International Islamic University, Islamabad, 44000 Pakistan

^e Center for Interdisciplinary Research in Basic Sciences, International Islamic University, 44000 Islamabad, Pakistan

^f Malaysian Institute for Chemical and Bioengineering Technology (MICET), University Kuala Lumpur (UniKL), Lot 1988, Taboh Naning, 78000 Alor Gajah, Melaka, Malaysia

^g University of Newcastle, School of Environmental and Life Sciences, Office C325, Chemistry, Callaghan, New South Wales 2308, Australia

^h Department of Agronomy, University of Haripur, 22620, Pakistan.

*** Corresponding author's e.mail :** muhammad.k.rafiq@ed.ac.uk

shangzhh@lzu.edu.cn

1. [These authors contributed equally to this work.](#)

ABSTRACT

Previous biochar research has primarily focused on agricultural annual cropping systems with very little attention given to highly fragile, complex and diverse natural alpine grassland ecosystems. The present study investigated the effect of biochar on the growth of alpine meadows and soil health. This study was conducted in the Qinghai Tibetan Plateau over a three year period to investigate the effect of three rice husk biochar application rates alone and combination with high and low NPK fertilizer dosages on alpine meadow productivity, soil microbial diversity as well as pH, carbon and nitrogen content at 0-10 cm and 10-20 cm depth. At the end of the 3rd year soil samples were analysed and assessed by combined analysis of variance. The results showed that biochar application in combination with nitrogen (N), phosphorus (P) and potassium (K) fertilizer had a significant increase in fresh and dry biomass during the second and third year of the study as compared to control and alone biochar application ($p \leq 0.05$). Biochar alone and in combination with NPK fertilizer resulted in a significant increase in the soil pH and carbon contents of the soil. XPS results, the SEM imaging and EDS analysis of aged biochar demonstrated that the biochar has undergone complex changes over the 3 years as compared to fresh biochar. This research suggests that biochar has positive effect on alpine meadow growth and soil health and may be an effective tool for alpine meadow restoration.

Keywords Biochar · Alpine meadows · Soil health

Introduction

Grasslands are the largest extended biome on earth and play a significant role as carbon sink (He et al. 2009). The grasslands store about 34 % of the global terrestrial carbon and are highly fragile in terms of carbon stability (Cheng et al. 2011). The carbon stocks in grasslands have been notably driven by land-use changes and management measures (Cheng et al. 2011; Sun et al. 2011; Sousa et al. 2012). Since soil carbon and nitrogen cycles closely interact, it is important to examine how anthropogenic factors such as overgrazing affect both C and N stocks and their interactions in the soil (Houghton et al. 1999; He et al. 2008).

The Tibetan Plateau is a main watershed region for China, India, and Pakistan representing a distinct cryospheric environment (Wang et al. 2007; Shi et al. 2010). The plateau is a source of usable water for nearly 40 % of the world's population, including China and India (United Nations Environment Programme (UNEP). 2007). The plateau has the largest biome plateau area on the Eurasian continent and represents a major ecological region with the lowest-latitude permafrost in the globe (Wang et al. 2002). Diverse types of grasslands extending from the Tibetan Plateau to Inner Mongolia and the mountains of the Xinjiang province, thus constitute the third biggest grassland ecological unit on earth (Yang et al. 2012). About 85 % of the plateau consists of alpine grasslands serving as a major source for livestock grazing (Dong et al. 2010; Harris. 2010), predominantly yak and Tibetan sheep. Alpine grasslands provide additional vital ecosystem services such as carbon capture, biodiversity, soil and water conservations (Yang et al. 2004; Chen et al. 2008; Wang et al. 2009). The C stored in soils of the plateau (33.5×10^9 t C) makes up 2.4 % of total world soil C (Wang et al. 2002) but due to poor land management this carbon is being lost at an increasing rate.

Similar to other ecosystems, Tibetan plateau grasslands have been experiencing considerable deposition of atmospheric N in the form of nitric acid over the past three decades (Yang et al. 2012). Persistent acidification of the soil decreased pH and increased

base cation loss, resulting in enhanced aluminium toxicity and loss of soil productivity (Bowman et al. 2008). Sustained longer acidification of soil could also modify formation and function of grasslands ecologies, such as plant biodiversity loss, loss of biomass productivity, and fractional inhibition of C and N cycling (Liu et al. 2011; Yang et al. 2012).

Furthermore, raising more yaks and removal of yak dung results huge carbon and nitrogen losses in Tibetan grasslands. In 2006, 40 million tons of yak dung was produced and 60% of that was collected for household energy needs. The removal of yak dung from grasslands results a loss of 16 million tons of carbon, 0.8 million tons of N and 0.2 tons of P on annual basis, not only altering the C and N cycles on plateau but also causes grassland degradation (Cai et al., 2013; Ni, 2002; Tian et al., 2006; Lu et al., 2015).

Previous management measures including fencing of pastures, reduction in numbers of livestock and fertilizer applications have been practiced to restore these degraded grasslands (Akiyama and Kawamura. 2007). However, these management practices have not been demonstrated to restore the extremely degraded grasslands of the plateau (Wu et al. 2010b).

Fertilizer application improves grassland productivity and restores degraded grasslands. Research investigations have shown that N-P-K fertilizer can enhance grassland production and its forage quality. However, due to grassland degradation phenomenon, there is less nutrient maintenance in grassland vegetation and nutrients are more prone to leaching.

Biochar, produced by thermal decomposition of organic material (Lehmann and Joseph. 2009), has been shown to improve low fertility soils as well as sequester carbon to mitigate global warming (Lehmann et al. 2006; Sohi et al. 2010; Woolf et al. 2010). Biochar applications to low fertility soils have improved yields in different cropping patterns worldwide (Glaser et al. 2002; Jeffery et al. 2011; Kammann et al. 2011; Vaccari et al. 2011; Spokas et al. 2012; Wang et al. 2012). Additionally, biochar application, reduce soil acidity (Knowles et al., 2011), increase cation exchange capacity (CEC) of soil (Mikan and Abrams,

1996) and reduce concentrations of pollutants. Significant reduction of leaching of fertilizer N from soil has been reported as a result of amendment with biochar produced from forest residues (Manolikaki & Diamadopoulos, 2017). Reduction of nitrate leaching from soil amended by biochar produced from pecan shells has been demonstrated over 25 and 67 days (Chaplot & Cooper, 2015).

Yak dung clay blended biochar and yak manure biochar has been proved to enhance production of blue grass in an artificial pasture and highland barley crop in short term in Tibetan plateau (Rafiq et al., 2017 and Zhang et al., 2018), however yak manure has been used for household cooking purposes and have competitive uses for its conversion to biochar.

Rice is one of the most widely cultivated agricultural crops in China. In China, approximately 54 million tons of rice husk is produced every year. The high volumes of rice husks that are considered as waste after milling are not appropriately treated. Rice husk is one of the main feedstock used to produce bio-oil by fluidized-bed reactors or other fast pyrolysis systems in China (Wang and Liu, 2018). Abundant biochar produced during the process of fast pyrolysis as by-product in China could be a potential application for grassland restoration.

Keeping in view, this study therefore aims to investigate the dosage effect of surface-applied rice husk biochar and NPK fertilizer on fresh and dry yield of grassland biomass under field conditions over a period of three years. Changes in pH, C and N content at 0-10 cm and 10-20 cm depth as well as microbial functional diversity are also elucidated.

Materials and methods

Experimental field site

The field study was carried out at Dawu village, Maqin County, of the Golou Tibetan Autonomous Prefecture of Qinghai Province, China (34° 28'11" N, 100° 12'39"E). The alpine meadow is located at 4200 m above sea level. The soil type of the study field is silt-clay, an alpine meadow soil as declared by Chinese System for Soil Classification. The average

annual temperature of the area is -0.6°C , ranging from -10°C during the month of January to 11.7°C in the month of July. The annual mean precipitation is 513 mm occurring during the months of May to September. There is no entirely frost-free period. The primary vegetation type in the area is alpine meadows dominated by *Kobresia spp*, *Polygonum spp*. and *Poa spp*.

Characterization of experimental biochar

Rice husk was obtained from Hangzhou, Zhejiang Province (China) and converted to biochar at a pyrolysis temperature of 500°C using a vertical furnace with continuous feeding (Jiaxing JIAHUA Animal Husbandry Co., Ltd., Zhejiang, China). The physico-chemical characteristics of the biochar such as pH, ash content, total nitrogen, total carbon, total hydrogen, total phosphorous, total potassium, calcium, magnesium, sodium were analysed. The pH of biochar was measured in deionized water at the ratio of 1:5 wt/wt with a calibrated Orion 720 pH meter (Enders et al., 2012). Ash content was analyzed by heating biochar samples at 500°C for 8 h in a muffle furnace (Dai et al., 2013). The elemental composition was determined according to Enders et al. (2012) using an elemental analyzer from Elementar Analysensysteme GmbH (varioELcube). Nutrient elements Ca, K, Mg, Na, and P were measured using an inductively coupled plasma-atomic emission spectrometer (IRIS ER/S). Before analysis, the biochar sample (about 0.05 g) was first digested by the concentrated $\text{HNO}_3/\text{H}_2\text{O}_2$ solutions (Dai et al., 2013).

BET (N_2) surface area, FTIR and thermo gravimetric analysis (TGA) were determined prior to field application according to techniques reported by Rafiq et al. (2016). X-ray photoelectron spectroscopy (XPS) spectra were collected from biochar powders with a thermo ESCALAB 250 spectrometer using an Al Ka monochromatized source and a multidetection analyzer under a 10^{-8} Pa residual pressure. Surface charging effects were corrected with C 1s peak at 284.6 eV as a reference. Examination of the biochars before and

after the field trials was carried out using a Zeiss Sigma SEM with a Bruker X-ray dispersive spectrometer (EDS) detector.

Experimental design

The size of each experimental plot was 2×4 m. There was a distance of 50 cm between the experimental plots to serve as a buffer zone (Qi et al., 2015). There were twelve treatments in this experiment carried out in triplicate under randomized complete block design (RCBD). Biochar was applied at 3 application rates: low (2 t/ha, BC_L), medium (4 t/ha, BC_M) and high (6 t/ha, BC_H) to the grassland. Biochar application rates were selected based on the recommendations (Clare et al., 2014) that due to higher biochar production costs, it needs to apply around 1-5 t/ha to realise plant response. Furthermore, Rafiq et al., applied yak blended biochar @ 3 tons/ha on pasture areas in Tibetan plateau. Two levels of NPK fertilizer were applied (30N, 15 P and 10 K kg/ha) and (60 N, 30 P and 20 K kg/ha) and designated as NPK_L and NPK_H, respectively. Higher level of NPK fertilizer corresponds to the recommendations of (Yu li et., al 2015). The NPK fertilizer was applied in the form of urea for N, single superphosphate for P and potassium chloride for K. The detailed plan of the treatments applied include: T₁ = CK (Control, no amendment), T₂ = BC_L, T₃ = BC_M, T₄ = BC_H, T₅ = NPK_L, T₆ = NPK_H, T₇ = BC_L+NPK_L, T₈ = BC_L+NPK_H, T₉ = BC_M+NPK_L, T₁₀ = BC_M+NPK_H, T₁₁ = BC_H+NPK_L and T₁₂ = BC_H+NPK_H. The biochar and NPK were applied through surface applications. The experiment commenced at the third week of June, 2014.

Vegetation and soil sampling

At the end of August 2014, 2015 and 2016, biomass samples were collected approximately 1 cm from the ground using 50×50 cm quadrat (Qi et al.,2015) while soil samples were collected at depths of 0 - 10 and 10 - 20 cm with the help of auger and placed into plastic bags and brought to the laboratory for further analysis of pH, carbon and nitrogen. At the end of August 2016, soil samples at a depth of 0 - 10 cm were collected for selected treatments as

control T₁, T₃, T₆ and T₁₀ to test effect of biochar and fertilizer on microbial functional diversity. In addition, biochar samples were subjected to microscopic and XPS analysis to investigate changes on the surface.

Biomass and soil measurements

Fresh biomass of the collected grass samples was weighed and recorded in the field soon after harvesting. The fresh samples were then put into paper bags and brought to laboratory for dry weight measurements. Biomass samples were dried at 65°C for 48 hrs in oven (Pérez-Suárez et al., 2014) and their dry biomass recorded. After cleaning and sieving with a 2 mm sieve, the air-dried soil samples (dried till constant weight) were tested for pH, C and N. The pH value of the experimental soils was tested using 1 : 2.5 soil : water suspension (Thiele-Bruhn et al. 2015) with an Orion 720 pH meter with a combination electrode. Total carbon and nitrogen of the soil was determined using elemental analyzer (Elementer Analyse systeme GmbH, varioEL-cube).

Separation of aged biochars from soils

Biochar particles present in soil were collected from the experimental fields after three years during August 2016 and brought to laboratory. Biochar samples were shaken to remove soil particles in DI water solution at a ratio of 1:10 w/v. The biochar was then washed four times with distilled water and dried at 60 °C (Koide et al., 2011) for further XPS and SEM analysis.

Incubation experiment for microbial functional diversity analysis

The microbial functional diversity of soil microbial population was determined using the Biolog EcoPlate™ (BIOLOG Inc., CA, USA). The soil samples were mixed with 90 mL of sterilized 0.85% (w/v) NaCl solution and shaken for 20 min followed by pre-incubation for 24 hours to initiate microbial utilization of soluble organic compound present in the soil. Samples were brought to 10⁻³ final dilutions before inoculation. Biolog EcoPlate™ has 96-

wells with three repeats, each one consisting of 31 sole carbon sources and a control with water. The consumption rate of carbon sources was tested by the reduction in tetrazolium dye which turns from color less to purple. The optical density (OD) of incubated plates was measured at 590 nm and 25°C with a plate reader and monitored every 24 hr for 7 days. The Procedure adopted by Rafiq et al., 2017 was followed to to investigate the microbial diversity and activity in this study. Average well color development (AWCD) was calculated using the equation,

$$AWCD = \Sigma(C-R)/31$$

where C is optical density (OD) of every well of carbon and R is the OD value of control with water only.

Negative (C-R) values were excluded from further analysis.

Microbial functional diversity was measured with the Shannon index (H') as follows,

$$H' = -\Sigma P_i \ln(P_i),$$

where P_i was determined by subtracting control OD from OD of every other well. After that it is divided by the total OD for all 31 substrates.

Data analysis

Analysis of variance was conducted and Duncan's Multiple Range Test (DMRT) at 5% level of probability was employed to compare means. Computer based statistical package MSTATC following Steel et al. (1997) was applied for this statistical analysis. To evaluate the cumulative effect of twelve treatments over the three year period on fresh biomass (FB), dry biomass (DB) and soil properties PHA, PHB are pH values at 0-10 cm and 10-20 cm soil depths, NA, CA are nitrogen and carbon content at 0-10 cm and NB and CB indicates nitrogen and carbon content at 10- 20 cm soil depth., were analyzed and prior to data analysis all variables were subjected to normality test and found that data for all of the variables were distributed normally. Mean comparison was done using Duncan test for each

dependent variable separately at 0.05 level. The data were subjected to principal component analysis (PCA) in a Multivariate analysis.

Results

Physico-chemical characterisation of rice husk biochar

The TG curves and FTIR spectra are provided in Figure S1. Most of the carbon (94 %) in the rice husk biochar remained even when heated to 700°C, indicating a highly stable carbon in the material. Fourier Transform Infrared (FTIR) spectroscopy revealed a broad peak at 3432 cm⁻¹ and 577 as well as sharp peaks at 2922, 2880, 1644, 1421, 887 and x cm⁻¹. A weakly defined peak was also detected at 1122 cm⁻¹.

Table 1 summarises the physico-chemical characteristics of rice husk biochar used in the experiment. The pH value of biochar was 10.4 with a carbon content of 40.8 wt.%. The biochar had an ash content of 39.7 wt.% and trace amounts of N, P, K and other elements necessary for plant growth. The biochar has a BET surface area of 3.19 g/m² and an average pore width of 10.6 nm.

Biomass responses to biochar and fertilizer application on alpine meadow

Table 2 shows the biomass productivity response of the alpine meadow in Tibetan plateau as a result of biochar application from 2014-16. It was found that an increasing biochar application rate resulted in an increase in fresh and dry biomass yield during the first year of biochar application.. However, this increase in biomass was not statistically significant at the $p = 0.05$ level probably.. When biochar was applied together with NPK the best yield was observed for BC_M throughout the study period. The fresh biomass yields of the treatments like BC_H+NPK_H, BC_M+NPK_H, BC_L+NPK_H were significantly greater than biochar treatments ($p \leq 0.05$). The largest dry biomass yield was measured for the NPK_H treatment in 2014. During the second year, there was no significant difference in fresh and dry biomass

yield between the control and those from the biochar applications alone. In contrast, NPK fertilizer application showed a significant increase for both fresh and dry grass yield as compared to the control and pure biochar applications. However, the greatest significant increase in fresh biomass yield was measured in the BC_M+NPK_H treatment. The fresh and dry biomass productivity of the meadow in the third year was significantly greater for all treatments compared to control. However maximum fresh and dry biomass yield was observed for the (BC_M+NPK_H) treatment throughout the study period. The increase in biomass as biochar and fertilizer application together indicates that responses of alpine meadows to addition of biochar and fertilizer were additive and positive.

Amelioration effects of biochar and fertilizer application on soil pH, carbon and nitrogen content of alpine meadow

The soil pH data at two soil depth levels for a period of three years is presented in Table 3. The addition of biochar led to increase the soil pH value significantly over a three years. The data indicate that soil had a lower pH at surface level (0-10cm) as compared to 10-20 cm of soil depth. Biochar application alone or combined with NPK fertilizer showed significantly higher pH values at 0-10 cm soil depth as compared to control and alone NPK fertilizer treatments ($p \leq 0.05$) during the first year of study. However, biochar addition results higher pH levels in the both soil depth levels during the second and third years of the study. This indicates that effect remained to persistent over time. The nitrogen content of alpine meadow soil at two depth (0-10 and 10-20) during 2014-16 is provided in Table 4. Application of biochar and fertilizer led to effective addition of nitrogen in soil. Nitrogen concentrations increased in the meadow soil with biochar application. A greater nitrogen content was observed at 0-10 cm soil depth level as compared to 10-20 cm. The greatest total nitrogen content (0.55 wt.%) in year one was observed for NPK_H treatment and there were significant increases between the control and the other treatments (except BC_L) at 0-10 cm depth during

the first year of study ($p \leq 0.05$). In year 2, the most significant increase in N soil content (0.66 wt.%) was measured in the NPK_H treatment for the 0-10cm soil profile and in the BC_L⁺ NPK_H for the 10-20cm profile.

Changes in soil carbon content over the three years for all treatments is given in Table 5. There was little change in the C content of the control at both depths over the 3 years. The addition of BC_H⁺ NPK_H and BC_H⁺ NPK_L resulted in the largest increase in C in the top soil profile in year one; BC_M⁺ NPK_H in year 2, and BC_L⁺ NPK_H in year 3. For the samples taken at depths between 10-20 cm, the greatest C content was measured in BC_M, BC_H⁺ NPK_H and BC_M⁺ NPK_H for year 1, BC_M⁺ NPK_H in year 2 and BC_L in year 3.

Cumulative impact of different treatment on biomass and soil properties over three years. Based on a combined ANOVA over three years, results revealed that there was a significant effect of time on all dependent variables (Table 6). The results also show that all dependent variables were statistically significant in all treatments. Results of mean comparison among treatments using Duncan Multiple range test indicated that the greatest mean for fresh biomass (FB) was observed in BCM+NPKH ($179 \pm 18 \text{ g/m}^2$) which was significantly higher than other treatments and the lowest level of FB belonged to BCL, BCM, BCH which were not statistically different from the control group. For DB, results of mean comparison showed that the highest mean of BD was observed for BC_M + NPK_H ($114 \pm 15 \text{ g/m}^2$) which was higher than other treatments.

The highest level of soil pH (A) (Table 7) was measured in treatment BCH (6.99 ± 0.05), which was significantly different from other treatments and the lowest level of pH (A) was observed for control group (6.63 ± 0.04). For pH (B) results of mean comparison showed that three treatments had the highest level including BCH (7.02 ± 0.03), BCL+NPKL (7.02 ± 0.04) and BCL+NPKH (7.01 ± 0.04), which were not statistically different. The lowest pH (B) belonged to control group (6.76 ± 0.03) and BCM+NPKH (6.84 ± 0.06). These results

300 indicated that the level of N (B), NPKL (0.41 ± 0.02) was significantly higher than other
301 treatments except BCM, BCL+NPKH, BCM+NPKH and BCH+NPKL. Results of mean
302 comparison for C(A) and C(B) revealed that BCM+NPKH had the highest means score for
303 both C(A) (7.37 ± 0.39) and C(B) (4.3 ± 0.3) two variable.
304

The data were subjected to principal component analysis (PCA) in a multivariate analysis. Biomass productivity and soil characters have been used to define patterns on the impacts of treatments applied. Results showed that three components with Eigen values more than one were extracted and these three components explained 74.1 % of total variability (Figure 1). This shows great variation among biomass productivity and soil characteristics under investigation. The first principal component (PC1) comprising of PHB, PHA and NB explained 25.0 % of total variability (Table 8). The characters with greatest positive weight on PC2 were CB and CA and these components explained 24.7 % of total variance among all data. DB, NA and FB were associated with the third principal component (PC3) which explained 24.4 % of total variance.

Effect of biochar and fertilizer on functional and microbial diversity

Figure 2 shows that $BC_M^+ NPK_H$ had the highest AWCD values at 144 hours as compared to CK, BC_M and NPK. The results showed that biochar and NPK fertilizer applied in combination had positive impacts on the microbial activity as compared to control or other selected treatments.

The values of microbial diversity (H') at incubation of 144 h against different treatments showed that biochar application (BC_M) alone and in combination with NPK fertilizer ($BC_M^+ NPK_H$) had higher Shannon Index values indicating that biochar addition can improve soil microbial diversity (Table 9).

X-Ray photoelectron and electron dispersive spectroscopy of original and aged biochar

XPS and SEM-EDS analysis of original and aged biochar shows that the biochar has undergone complex changes over the 3 years (Table 10, Figure 3 A,B). There has been a decrease in the aromatic carbon and an increase in organic compounds yielding a higher content of C/O and C/N functional groups, K, Si, Ca, Mg, N, S and Fe atomic % than the control soil and stored biochar (Table 10). The $-C=C-$ functional group constituted 63.4 mol

- % in original biochar, while biochar extracted from the soil had 51.1 mol-%. The functional groups – C-OH, C-O-C=, C-O-R and – C-N, C=O increased upon aging in soil.

SEM-EDS results show that the surface of original biochar has a relatively large content of Si, no detectable Fe and only relatively small concentrations of K, Ca, S, Al, P and Cl. The aged biochar, on the other hand, contained higher concentrations of K, Fe and Mn and Al. These images and elemental and functional group measurements are indicative of the formation of organo-mineral clusters on the surface of the biochar.

Discussion

It has been observed in several studies that biochar addition to soils due to its various properties has improved soil fertility and thus increased crop yields on agricultural lands (Marris, 2006; Chan et al. 2007). The characterization for pH, C, N and ash content were within the range reported for rice husk biochars used by Manickam et al. (2015). BET (N₂) surface area of rice husk biochar used in this experiment was lower than rice husk biochar produced in gasifiers (Manickam et al. 2012) as well as the peanut biochar used by Du et al. (2018). The observed variability is attributed to differences in process conditions primarily temperature (Rafiq et al. 2016) as well as feedstock type.

Observed FTIR peaks are in close agreement with biochars produced Sharma et al. (2004) from lignin at pyrolysis temperatures $\geq 450^{\circ}\text{C}$. FTIR peaks at wavelengths 3432 and 1122 cm^{-1} are attributed to -OH and C-O stretching vibration of phenolic compounds (Sharma et al. 2004; Ma et al. 2017). The appearance of peaks at 887 and 790 cm^{-1} are not only indicative of aromatic C-H but also evidence of formation of fused ring systems (Sharma et al. 2004). Sharma et al. (2004) observed a slow decrease in aliphatic CH stretch (2800-3000 cm^{-1}) with increase in pyrolysis temperature. . The presence of aliphatic CH was also observed in rice husk biochar used in this study suggesting that it originated from lignin.

The H/C molar ratio of 0.26 was well below 0.7 as required by IBI standards and EU guidelines (2012). The O/C molar ratio was 0.33 which meets the standards of the EU guidelines (2012). Similar H/C and O/C ratios have been reported for rice husk biochar in literature (Manickam et al. 2012). The molar H/C_{org} ratio can be used to predict the relative amount of organic biochar carbon that remains after 100 years incubation in soil (Budai et al. 2013). The organic carbon content in our rice husk biochar (Table 1) was assumed to be the same as total carbon since the carbonate content in wood and grass based biochars was found to be negligible (Enders et al. 2012). Hence, 91 wt.% of the rice husk biochar carbon can be expected to remain in alpine meadow soil after 100 years barring other factors such as loss due to erosion.

The application of rice husk biochar showed positive effects on alpine meadows biomass productivity over three years with and without NPK fertilizers. Crop productivity is often reported to increase with biochar application to soils but not always consistently (Jeffery et al. 2011; Subedi et al. 2016). The results from soil trials demonstrated that biochar/NPK fertilizer can assist in alpine meadow restoration. The biochar and NPK application did not show a significant impact on biomass yield during the first year of application however, in the subsequent years as in second and third years biomass yield was observed having a significant increase with the application of biochar with and without fertilizer (Table 2). Delayed impacts of biochar application on biomass improvements, till one or two years, have been reported in the literature (Haefele et al. 2011; Carvalho et al. 2016). These findings are consistent with the findings of this experiment. We observed biomass improvements during second and third years of the biochar application. Furthermore, results showed that significantly improved the biomass productivity of meadows during the second and third years. Persistent increase in crop productivity following biochar inputs are a good indicator of economic viability for scaling up the applications (Liu et al., 2013). Similar results were also

reported by Adam et al (2013) and Slavich et al (2013) who observed that biochar has the ability to improve prairie growth and prairie restoration. Possible reasons for the nonsignificant effects of biochar on forage biomass during the first year may be related to lower biochemical processes in alpine areas having lower temperatures, in the presence or absence of biochar, plus slower biochar degradation and its interaction with soil and consequently delaying its beneficial effects on soil properties and plant productivity (Verheijen et al. 2010; Fang et al. 2015). Several mechanisms for increase in biomass yield after biochar applications have been discussed in the literature. These include liming effects of biochar, improved water holding capacity of soils, nutrient use efficiency and reduced leaching, improved soils structure and porosity and increased surface area for nutrient adsorption. Many studies shown that over time aging of biochar in soils have more produced effects of biochar on soil moisture content (Paetsch et al., 2018). This increased moisture content and improvement of soil structure amended with biochar leads to effective root system development for water and nutrient supply. Perhaps, these factors contributed to the improved biomass productivity of alpine meadow after biochar application in this experiment.

The results showed that biochar application improved the soil pH values in the alpine meadows. The plant feedstock materials that are used to produce biochar contain base cations and these cations are transferred to biochars during pyrolysis of organic materials. The rice husk biochar contains high concentrations of soluble oxides, hydroxides and carbonates of Ca, Mg and K (Table 1), which may have contributed to the increase in soil pH, as observed in our study (Table 3). Increase in soil pH values has also been reported by Laird *et al.* (2010), where biochar with high ash content (14-56%), similar to present study (37% ash), were used. The alkalinity character is enhanced with pyrolysis temperature allowing rice husk biochar to act as a liming agent (Lehman et al. 2007; Wang et al. 2014).

405 Similar findings were reported in previous studies (Demirbas et al. 2004; Chan et al. 2007;
406 Revell et al. 2012; Wang et al. 2014). The application of biochar due to its ability to act as a
407 liming agent improved soil pH levels. Similar findings were reported in previous studies
408 (Demirbas et al. 2004; Chan et al. 2007; Revell et al. 2012; Wang et al. 2014). Similarly,
409 Novak et al. (2009) found that biochar enhanced soil pH in the southern United States. Wang
410 et al (2014) also showed that biochar application could increase the carbon content in soil.
411 Similarly it have be investigated that use of biochar application in prairie rehabilitation
412 initiatives and proved biochar addition not only enhances improve the growth of prairie
413 species, but also sequestered carbon (Lehman et al. 2007) and accelerated the recovery of
414 carbon pool in these soils improve the growth of prairie species, but also sequestered carbon
415 (Lehman et al. 2007) and accelerated the recovery of carbon pool in these soils. The AWCD
416 value in the well of an EcoPlate™ is a key indicator of microbial functional diversity,
417 because it indicates the capability of soil microorganisms to utilize various carbon substrates.
418 Previous findings have shown that the application of organic matter to soil can enhance
419 microbial populations their diversity and activities (Gomez et al., 2006).The results of this
420 experiment showed that biochar and NPK fertilizer applied in combination had positive
421 impacts on the microbial activity and diversity. The biochar upon aging has shown (table:10,
422 fig:4), that there are increased c/o functional groups in biochar. These characteristics have
423 been proved to increase the abundance of beneficial microorganisms in soil (Ye et al., 2017).
424 The findings are consistent with published studies that found that microbial activity enhanced
425 with biochar application (Kolb et al. 2009; Liang et al. 2010). Liao et al (2016) also found
426 that biochar application has positive effects on soil microbial diversity. XPS results and the
427 SEM imaging and EDS analysis shows that the biochar has undergone complex changes over
428 the 3 years and these changes are similar to those describe by Joseph et al (2010), Archanjo et
429 al (2017) and Hagemann et al (2017). These images and elemental and functional group

measurements are indicative of the formation of organo-mineral clusters on the surface of the biochar. Previous research (Joseph et al. 2010; Archanjo et al. 2017; Hagemann et al. 2017) has shown that these clusters with high content or redox active Fe and Mn minerals that are bonded by organic compounds that have a high concentration of C/O functional groups can increase the ability of plants to take up nutrients.

Conclusion

This study has demonstrated that biochar can have significant effects on biomass production, soil acidification and carbon sequestration. In addition, biochar showed positive effects on microbial diversity and activity. Application of biochar to natural, wasteland and degraded systems could be a potential strategy to sequester carbon (Woolf et al. 2010). Further research is required to evaluate the long-term effects of biochar species diversity plant and detailed soil dynamics like nutrient mineralization, availability and transfer to plant. Additionally, biochar application methods and biochar erosion aspects need to be investigated for its appropriate testing mechanism. More research work is also required to develop and test biochar from the local feed stocks and to enrich it with heterogeneous nutrient material like attapulgite clay for its cost effectivity and wider acceptability.

Acknowledgements

This study was supported by the national key research and development project (2016YFC0501906), the fundamental research funds for the central Universities(lzujbky-2017-k16), natural science foundation of China (41671508; 31870433), the second stage's research and technique extending project of Sanjiangyuan ecological protection and building

of 2017 in Qinghai (2017-S-1-06), Key R&D and transformation program of Qinghai (2017-NK-149-2), and Qinghai innovation platform construction project (2017-ZJ-Y20).

References

Adams KM, Benjamin TJ, Emery NC et al (2013) The Effect of Biochar on Native and Invasive Prairie Plant Species. *Invasive Plant Science and Management* 6:197–207.

Akiyama T, Kawamura K (2007) Grassland degradation in China, methods of monitoring, management and restoration. *Grassl. Sci* 53(1):1–17.

Archanjo BS, Mendoza M E, Albu M et al (2017) Nanoscale analyses of the surface structure and composition of biochars extracted after composting and from field trials using advanced analytical electron microscopy. *Geoderma* 294:70–79.

Bowman WD, Cleveland CC, Halada L et al (2008) Negative impact of nitrogen deposition on soil buffering capacity. *Nature Geoscience* 1:767–770.

Cai, YJ, Wang, XD, Ding WX et al (2013). Potential short-term effects of yak and Tibetan sheep dung on greenhouse gas emissions in two alpine grassland soils under laboratory conditions. *Biol. Fertil. Soils* 49, 1215–1226. [http:// dx.doi.org/10.1007/s00374-013-0821-7](http://dx.doi.org/10.1007/s00374-013-0821-7).

Carvalho MTM, Madari BE, Bastiaans, L et al (2016) Properties of a clay soil from 1.5 to 3.5 years after biochar application and the impact on rice yield. *Geoderma* 276:7–18.

Chan KY, Van Zwieten L, Meszaros I et al (2007) Agronomic values of greenwaste biochar as a soil amendment. *Soil Res* 45: 629–634.

Chaplot V, Cooper M Soil aggregate stability to predict organic carbon outputs from soils. *Geoderma*, 243 (2015), pp. 205-213

Chen J, Yamamura Y, Hori Y et al (2008) Small-scale species richness and its spatial variation in an alpine meadow on the Qinghai–Tibet Plateau. *Ecol. Res* 23: 657–663.

480 Cheng J, Wu GL, Zhao LP et al (2011) Cumulative effects of 20-year exclusion of livestock
 481 grazing on above- and belowground biomass of typical steppe communities in arid areas
 482 of the Loess Plateau, China. *Plant Soil Environ* 57: 40–44.

483 Clare A, Shackley SS, Joseph J (2014) Competing uses for China's straw: the economic and
 484 carbon abatement potential of biochar. *GCB Bioenergy* (2014), 10.1111/gcbb.12220.

485 Dai Z, Meng J, Muhammad N, Liu X et al. (2013). The potential feasibility for soil
 486 improvement, based on the properties of biochars pyrolyzed from different feedstocks. *J.*
 487 *Soils Sediments*. 13, 989–1000.

488 Demirbas B (2004) Effects of temperature and particle size on bio-char yield from pyrolysis
 489 of agricultural residues. *J Anal Appl Pyrolysis* 72(2): 243–248.

490 Dong SK, Wen L, Zhu L et al (2010) Implication of coupled natural and human systems in
 491 sustainable rangeland ecosystem management in HKH region. *Frontiers of Earth*
 492 *Science in China* 4: 42–50.

493 Du Z, Xiao Y, Qi X et al (2018) Peanut-Shell Biochar and Biogas Slurry Improve Soil
 494 Properties in the North China Plain: A Four-Year Field Study. *Scientific Reports* 8:1–9.

495 Enders A, Hanley K, Whitman T, Joseph S, Lehmann J (2012) Characterization of biochars
 496 to evaluate recalcitrance and agronomic performance. *Bioresource Technol* 114:644–
 497 653

498 Glaser B, Lehmann J, Zech W (2002) Ameliorating physical and chemical properties of
 499 highly weathered soils in the tropics with charcoal—a review. *Biology and Fertility of*
 500 *Soils* 35:219–230.

501 Gomez E, Ferreras L, Toresani S (2006) Soil bacterial functional diversity as influenced by
 502 organic amendment application. *Bioresour. Technol.*, 97 (2006), pp. 1484-1489

503 Haefele SM, Konboon Y, Wongboon W et al (2011). Effects and fate of biochar from rice
 504 residues in rice-based systems. *Field Crop Research* 121: 430-440.

505 Hagemann N, Harter J, Behrens S (2016) Elucidating the impacts of biochar applications on
 506 nitrogen cycling microbial communities. In: Ralebitso-Senior, T.K., Orr, C.H. (Eds.),
 507 Biochar Application. Elsevier, Amsterdam, pp. 163–198

508 Harris RB (2010) Rangeland degradation on the Qinghai–Tibetan plateau, a review of the
 509 evidence of its magnitude and causes. *J Arid Environ* 74:1–12.

510 He NP, Wu L, Wang YS et al (2009) Changes in carbon and nitrogen in soil particle-size
 511 fractions along a grassland restoration chronosequence in northern China. *Geoderma*
 512 302–308.

513 He NP, Yu Q, Wu L et al (2008) Carbon and nitrogen store and storage potential as affected
 514 by land-use in a *Leymus chinensis* grassland of northern China. *Soil Biol Biochem*
 515 40:2952–2959.

516 Hossain MK, Strezov V, Chan KY et al (2010). Agronomic properties of waste water sludge
 517 biochar and bioavailability of metals in production of cherry tomato
 518 (*Lycopersicon esculentum*). *Chemosphere* 78: 1167–1171.

519 Houghton RA, Hackler JL, Lawrence KT (1999) The U. S. carbon budget, contributions in
 520 sustainable rangeland ecosystem management in HKH region. *Front Earth Sci China* 4:
 521 42–50.

522 Jeffery S, Verheijen FGA, Velde MVD et al (2011) A quantitative review of the effects of
 523 biochar application to soils on crop productivity using meta-analysis. *Agriculture,*
 524 *Ecosystems and Environment* 144:175–187.

525 Jones DL, Rousk J, Edwards-Jones G et al (2012) Biochar mediated changes in soil quality
 526 and plant growth in a three year field trial. *Soil Biology and Biochemistry* 45:113–124.

527 Joseph SD, Camps-Arbestain M, Lin Y et al (2010) An investigation into the reactions of
 528 biochar in soil. *Aust J Soil Res* 501–515.

529 Kammann CI, Linsel S, Gößling JW et al (2011) Influence of biochar on drought tolerance of
 530 *Chenopodium quinoa* Willd and on soil–plant relations. *Plant and Soil* 345(1-2):195–
 531 210.

532 Kolb SE, Fermanich KJ, Dornbush ME (2009) Effect of charcoal quantity on microbial
 533 biomass and activity in temperate soils. *Soil Sci Soc Am J* 73(4):1173.

534 Koide RT, Petprakob K, Peoples M (2011). Quantitative analysis of biochar in field soil. *Soil*
 535 *Biol. Biochem.* 43:1563–1568.

536 Knowles OA, Robinson BH, Contangelo A et al (2011) Biochar for the mitigation of nitrate
 537 leaching from soil amended with biosolids. *Sci. Total Environ.*, 409 pp. 3206-3210

538 Lehmann J, Gaunt J, Rondon M (2006) Bio-char sequestration in terrestrial ecosystems-a
 539 review. *Mitigation and Adaptation Strategies for Global Change* 11:395–419.

540 Li Yu, Xiao-Long S, Jian-ning Z et al (2015), Responses of plant diversity and primary
 541 productivity to nutrient addition in a *Stipa baicalensis* grassland, China. *Journal of*
 542 *Integrative Agriculture.* 14(10): 2099–2108.

543 Liang B, Lehmann J, Sohi SP et al (2010) Black carbon affects the cycling of non-black
 544 carbon in soil. *Org Geochem* 41: 206e213.

545 Liao N, Li Q, Zhang W et al (2016) Effects of biochar on soil microbial community
 546 composition and activity in drip-irrigated desert soil. *European Journal of Soil Biology*
 547 72:27-34.

548 Liu XJ, Duan L, Mo JM et al (2011) Nitrogen deposition and its ecological impact in China:
 549 an overview. *Environ Pollut* 159: 2251–2264.

550 Liu Y, Yang M, Wu Y et al (2011) Reducing CH₄ and CO₂ emissions from waterlogged
 551 paddy soil with biochar. *Journal of Soils and Sediments* 11:930–939.

552 Lu X, Yan Y, Sun J (2015). Short-term grazing exclusion has no impact on soil properties

and nutrients of degraded alpine grassland in Tibet, China. *Solid Earth* 6, 1195–1205.

<http://dx.doi.org/10.5194/se-61195-2015>

Ma Z, Yang V, Ma Q et al (2017) Evolution of the chemical composition, functional group, pore structure and crystallographic structure of bio-char from palm kernel shell pyrolysis under different temperatures. *Journal of Analytical and Applied Pyrolysis* 127:350–359.

Manickam T, Bachmann RT, Illani ZI et al (2012) Characterization of Local Mill Rice Husk Charcoal and Its Effect on Compost Properties. *Malaysian Journal of Soil Science* 16:89–102.

Manickam T, Cornelissen G, Bachmann RT et al (2015) Biochar application in Malaysian sandy and acid sulfate soils: Soil amelioration effects and improved crop production over two cropping seasons. *Sustainability (Switzerland)* 7:16756–16770.

Manolikaki I, E. Diamadopoulos E (2017) Ryegrass yield and nutrient status after biochar application in two Mediterranean soils *Archives of Agronomy and Soil Science*, 63 (8), pp. 1093-1107

Mikan CJ, Abrams AD (1996) Mechanisms inhibiting the forest development of historic charcoal hearths in southeastern Pennsylvania *Can. J. For. Res.*, 26 (1996), pp. 1893-1898

Ni J, 2002. Carbon storage in grasslands of China. *J. Arid Environ.* 50, 205–218. <http://dx.doi.org/10.1006/jare.2001.0902>.

Novak JM, Busscher WJ, Laird D et al (2009) Impact of biochar amendment on fertility of a Southeastern coastal plain soil. *Soil Sci* 174:105–112.

Paetsch L, Mueller CW, Kögel-Knabner I (2018) Effect of in-situ aged and fresh biochar on soil hydraulic conditions and microbial C use under drought conditions. *Sci. Rep.*, 8 (1) (2018), p. 6852

578 Pérez-Suárez M, Castellano MJ, Kolka R et al (2014) Nitrogen and carbon dynamics in
 579 prairie vegetation strips across topographical gradients in mixed Central Iowa
 580 agroecosystems. *Agric Ecosyst Environ* 188:1–11.

581 Qi J, Nie Z, Jiao T et al (2015) Phosphorus and Defoliation Interact and Improve the Growth
 582 and Composition of the Plant Community and Soil Properties in an Alpine Pasture of
 583 Qinghai-Tibet Plateau. *PLoS ONE* 10(10): e0141701.

584 Rafiq MK, Bachmann RT, Rafiq MT et al (2016) Influence of Pyrolysis Temperature on
 585 Physico-Chemical Properties of Corn Stover (*Zea mays* L.) Biochar and Feasibility for
 586 Carbon Capture and Energy Balance. *PLOS ONE* 11:e0156894.

587 Rafiq MK, Joseph SD, Li F et al (2017). Pyrolysis of attapulgite clay blended with yak dung
 588 enhances pasture growth and soil health: characterization and initial field trials. *Sci.*
 589 *Total Environ.* 607–608 (2017), pp. 184-194.

590 Revell KT, Maguire RO, Agblevor FA (2012) Field trials with poultry litter biochar and its
 591 effect on forages, green peppers, and soil properties. *Soil Sci* 177:573–579.

592 Sharma RK, Wooten JB, Baliga VL et al (2004) Characterization of chars from pyrolysis of
 593 lignin. *Fuel* 83:1469–1482.

594 Shi XM, Li XG, Long RJ et al (2010) Dynamics of soil organic carbon and nitrogen
 595 associated with physically separated fractions in a grassland-cultivation sequence in the
 596 Qinghai–Tibetan Plateau. *Biol Fertil Soils* 46:103–111.

597 Singh BP, Hatton BJ, Singh B et al (2010) Influence of biochars on nitrous oxide emission
 598 and nitrogen leaching from two contrasting soils. *Journal of Environment Quality*
 599 39:1224.

600 Slavich PG, Sinclair K, Morris SG et al (2003) Contrasting effects of manure and green waste
 601 biochars on the properties of an acidic ferralsol and productivity of a subtropical pasture.
 602 *Plant and Soil* 366(1-2):213-227.

603 Sohi S, E Krull, E Lopez-Capel et al (2010) A review of biochar and its use and function in
604 soil. *Advances in Agronomy* 105:47–82.

605 Sousa FP, Ferreira TO, Mendonca ES et al (2012) Carbon and nitrogen in degraded Brazilian
606 semi-arid soils undergoing desertification. *Agric Ecosyst Environ* 148:11–21.

607 Spokas KA, Cantrell KB, Novak JM et al (2012) Biochar: A synthesis of its agronomic
608 impact beyond carbon sequestration. *J Environ Qual* 41:973–989 (this issue).

609 Steel RGD, Torrie JH, Dickey DA (1997) *Principles and procedures of statistics: A*
610 *biometrical approach*. 3rd ed. McGraw Hill Book Co. Inc. New York: 400-428.

611 Subedi R, Taupe N, Pelissetti S et al (2016) Greenhouse gas emissions and soil properties
612 following amendment with manure derived biochars: influence of pyrolysis temperature
613 and feedstock type. *Journal of Environment Management* 166:73–83.

614 Sun DS, Wesche K, Chen DD (2011) Grazing depresses soil carbon storage through changing
615 plant biomass and composition in a Tibetan alpine meadow. *Plant Soil Environ* 57:271–
616 278.

617 Thiele-Bruhn S, Wessel-Bothe S, Aust MO (2015) Time-resolved in-situ pH measurement in
618 differently treated, saturated and unsaturated soils. *J Pl Nutr Soil Sci* 178(3):425-432.

619 Tian HQ, Wang SQ, Liu JY et al (2006). Patterns of soil nitrogen storage in China. *Global*
620 *Biogeochem. Cycles* 20, GB1001. [http://dx. doi.org/10.1029/2005GB002464](http://dx.doi.org/10.1029/2005GB002464).

621 Vaccari FP, Baronti S, Lugato E et al (2011) Biochar as a strategy to sequester carbon and
622 increase yield in durum wheat. *European Journal of Agronomy* 34:231–238.

623 Van Zwieten L, Kimber S, Morris S et al (2010) Effects of biochar from slow pyrolysis of
624 papermill waste on agronomic performance and soil fertility. *Plant and Soil* 327:235–
625 246.

626 Wang CT, Long RJ, Wang QL et al (2009) Changes in plant diversity, biomass and soil C, in
627 alpine meadows at different degradation stages in the Headwater Region of Three
628 Rivers, China. *Land Degrad Dev* 20:187–198.

629 Wang GX, Qian J, Cheng G D et al (2002) Soil organic carbon pool of grassland soils on the
630 Qinghai-Tibetan plateau and its global implication. *The Science of the Total*
631 *Environment* 291:207–217.

632 Wang GX, Qian J, Cheng GD et al (2002) Soil organic carbon pool of grassland soils on the
633 Qinghai–Tibetan Plateau and its global implication. *Sci Total Environ* 291:207–217.

634 Wang GX, Wang YB, Li YS et al (2007) Influences of alpine ecosystem responses to
635 climatic change on soil properties on the Qinghai–Tibetan Plateau, China. *Catena*
636 70:506–514.

637 Wang J, Pan X, Liu Y et al (2012) Effects of biochar amendment in two soils on greenhouse
638 gas emissions and crop production. *Plant and Soil* 360:287–298.

639 Wang Y, Liu R (2018). Improvement of Acidic Soil Properties by Biochar from Fast
640 Pyrolysis. *Environmental Progress & Sustainable Energy*. Vol.37:5.1743-1749.
641 <https://aiche.onlinelibrary.wiley.com/doi/epdf/10.1002/ep.12825>

642 Wang Y, Yin R, Liu R (2014) Characterization of biochar from fast pyrolysis and its effect
643 on chemical properties of the tea garden soil. *J Anal Appl Pyrol* 110:375–381.

644 Woolf, D, Amonette JE, Street-Perrott FA et al (2010) Sustainable biochar to mitigate global
645 climate change. *Nature Communications* 1:56.

646 Wu GL, Li ZH, Zhang L et al (2010b) Effects of artificial grassland establishment on soil
647 nutrients and carbon properties in a black-soil-type degraded grassland. *Plant Soil*
648 333:469–479.

649 Yang YH, Ji CJ, Ma WH et al (2012) Significant soil acidification across northern China's
650 grasslands during 1980s–2000s. *Glob. Change Biol* 18:2292–2300.

651 Yang YH, Rao S, Hu HF et al (2004) Plant species richness of alpine grasslands in relation to
652 environmental factors and biomass on the Tibetan Plateau. *Biodivers Sci* 12:200–205.

653 Ye J, Joseph S, Ji M et al (2017) Chemolithotrophic processes in the bacterial communities
654 on the surface of mineral-enriched biochars *ISME J.* (2017), [10.1038/ismej.2016.187](https://doi.org/10.1038/ismej.2016.187)

655 Zhang Y, Li B, Zheng D (2002) A discussion on the boundary and area of the Tibetan
656 plateau in China. *Geographical Resources* 21:1–8.

657 Zhang J, Huang B, Chen L et al (2018) Characteristics of biochar produced from yak
658 manure at different pyrolysis temperatures and its effects on the yield and growth of
659 highland barley. *Chemical Speciation & Bioavailability* 30 (1): 57–67.

660 Zhang LH, Xu CB, Champagne P (2010) Overview of recent advances in thermo-chemical
661 conversion of biomass, *Energy Conversion and Management* 51 :969–982

662

Figure.1 Scatter plot of the first three principal components of the PCA

Figure.2 AWCD of metabolized substrates in Biolog EcoPlates using four different soil samples (n=3)

Figure 3. A. Secondary electron images and elemental analysis of the surface of fresh rice husk biochar

Figure 3. B Secondary electron images and elemental analysis of the surface of f 3 year aged rice husk biochar

696

697 **Table 1.** Major properties of rice husk biochar

Parameter	Value	Parameter	Value
Ash (wt.%)	39.7 ± 0.5	Fe (wt.%)	0.73 ± 0.02
pH	10.38 ± 0.02	P (mg/l)	10.3 ± 0.13
C (wt.%)	40.8 ± 1.3	K (mg/l)	47.9 ± 0.4
N (wt.%)	0.32 ± 0.03	Ca (mg/l)	11.0 ± 0.2
H (wt.%)	0.89 ± 0.21	Mg (mg/l)	6.20 ± 0.1
O (wt.%)	17.9 ± 0.7	Na (mg/l)	2.06 ± 0.06
S (wt.%)	0.41 ± 0.08	BET (N ₂) surface area (m ² /g)	3.19
Si (wt.%)	11.92 ± 0.11	Average pore width (nm)	10.6

698

699

700 **Table 2.** Effect of biochar and fertilizer application on alpine meadow productivity over three
701 years (2014-2016)

Treatments	Fresh Biomass (g/m ²)			Dry Biomass (g/m ²)		
	2014	2015	2016	2014	2015	2016
CK	101.47 abc	108.67 d	119.33 h	42.36 abc	49.73 e	58.93 g
BC _L	90.48 d	106.50 d	128.29 g	34.56 c	48.60 e	69.21 f
BC _M	97.13 bcd	119.93 d	129.92 g	37.06 bc	61.00 e	68.69 f
BC _H	96.27 cd	119.60 d	132.29 g	39.86 bc	64.87 e	70.70 f
NPK _L	101.77 abc	165.67 c	155.33 f	49.76 abc	84.30 d	75.82 f
NPK _H	104.43 abc	169.53 c	164.67 e	60.73 a	90.27 cd	84.68 e
BC _L ⁺ NPK _L	107.07 ab	177.67 bc	174.29 d	50.06 abc	97.47 bcd	104.31 d
BC _L ⁺ NPK _H	110.50 a	164.03 c	174.20 d	51.63 abc	95.20 bcd	100.62 d
BC _M ⁺ NPK _L	102.50 abc	170.33 c	191.21 c	45.33 abc	104.00 bc	128.66 b
BC _M ⁺ NPK _H	109.33 a	197.00 a	229.24 a	54.90 ab	136.27 a	152.31 a
BC _H ⁺ NPK _L	105.87 abc	175.80 bc	188.47 c	51.56 abc	103.80 bc	104.91 d
BC _H ⁺ NPK _H	108.63 a	190.37 ab	204.07 b	56.10 ab	111.97 b	118.33 c

702 Column means presented with different letters indicate significance differences ($p \leq 0.05$)

703

704

705

706

Table 3. Effect of biochar and fertilizer application on soil pH over three years (2014-2016)

Treatments	pH					
	2014		2015		2016	
	0-10 cm	10-20-cm	0-10 cm	10-20 cm	0-10 cm	10-20 cm
CK	6.54 ef	6.64 d	6.62 f	6.79 e	6.72 g	6.83f
BC _L	6.77 abc	6.71 cd	6.84 de	6.87 de	6.92 ef	7.00 e
BC _M	6.87 a	6.71 cd	7.02 ab	7.02 ab	7.03 abc	7.11 ab
BC _H	6.80 ab	6.95 a	7.08 a	6.97 abc	7.09 a	7.13 a
NPK _L	6.49 f	6.72 cd	6.79 e	6.87 de	7.00 cde	7.01 e
NPK _H	6.53 de	6.75 c	6.90 bcde	6.91 cd	6.91 f	7.03 de
BC _L ⁺ NPK _L	6.77 abc	6.88 ab	6.99 abc	7.02 ab	7.08 ab	7.13 a
BC _L ⁺ NPK _H	6.75 bc	6.88 ab	6.98 abc	7.05 a	7.03 abc	7.09 abc
BC _M ⁺ NPK _L	6.69 cd	6.74 cd	6.91 bcd	6.92 cd	6.97 cdef	7.07 bcd
BC _M ⁺ NPK _H	6.78 abc	6.66 d	6.88 cde	6.81 e	6.94 def	7.05 cde
BC _H ⁺ NPK _L	6.79 ab	6.79 bc	6.87 cde	6.94 bcd	7.00 bcde	7.07 bcd
BC _H ⁺ NPK _H	6.85 a	6.93 a	6.94 bcd	6.96 abc	7.00 bcd	7.04 cde

Column means presented with different letters indicate significance differences ($p \leq 0.05$)

Table 4. Effect of biochar and fertilizer application on soil nitrogen content over three years (2014-2016) period

Treatments	Nitrogen (wt.%)					
	2014		2015		2016	
	0-10 cm	10-20 cm	0-10 cm	10-20 cm	0-10 cm	10-20 cm
CK	0.37 ef	0.29 b	0.35 f	0.27 e	0.39 c	0.27 f
BC _L	0.40 def	0.31 b	0.35 ef	0.31 bcde	0.44 c	0.33 cdef
BC _M	0.48 ab	0.34 b	0.50 cd	0.33 abcde	0.47 bc	0.36 bcde
BC _H	0.48 ab	0.43b	0.46 de	0.37 abc	0.44 c	0.32 def
NPK _L	0.50bcd	0.30 b	0.32 f	0.29 de	0.59 a	0.41 ab
NPK _H	0.55 a	0.42 b	0.66 a	0.39 ab	0.60 a	0.40 abc
BC _L ⁺ NPK _L	0.44 cde	0.34 b	0.37 ef	0.31 cde	0.46 bc	0.30 ef
BC _L ⁺ NPK _H	0.50 bcd	0.33 b	0.58 bc	0.39 a	0.55 ab	0.36 bcde
BC _M ⁺ NPK _L	0.53 bc	0.27 b	0.34 f	0.27 e	0.47 bc	0.38 bcde
BC _M ⁺ NPK _H	0.51 bc	0.34 b	0.62 ab	0.34 abcde	0.62 a	0.46 a
BC _H ⁺ NPK _L	0.52 bc	0.35 b	0.48 cd	0.35 abcd	0.55 ab	0.39 abcd
BC _H ⁺ NPK _H	0.54 a	0.35 b	0.51 cd	0.30 cde	0.45 bc	0.33 cdef

Column means presented with different letters indicate significance differences at ($p \leq 0.05$)

Table 5. Effect of biochar and fertilizer application on soil carbon content over time

Treatments	Carbon (%)					
	2014		2015		2016	
	0-10 cm	10-20 cm	0-10 cm	10-20 cm	0-10 cm	10-20 cm
CK	4.05 e	3.47 ab	3.89 e	3.17 efg	4.45 de	3.51 abcd
BC _L	4.78 de	3.82 ab	4.07 e	3.36 efg	5.24 c	4.12 a
BC _M	5.90 d	4.10 a	6.87 bc	3.89 cd	6.23 ab	3.51 abcd
BC _H	6.42 c	3.79 ab	5.72 d	3.19 efg	6.19 ab	2.99 cd
NPK _L	4.06 e	2.51 c	5.43 d	3.07 fg	4.93 cd	2.88 d
NPK _H	4.07 e	3.07 bc	3.67 e	2.87 g	4.17 e	3.26 bcd
BC _L ⁺ NPK _L	5.47 d	3.14 bc	6.14 cd	3.51 def	6.31 ab	3.32 bcd
BC _L ⁺ NPK _H	6.82 c	3.84 ab	7.54 b	4.49 b	6.92 a	3.94 ab
BC _M ⁺ NPK _L	7.08 bc	3.72 ab	7.08 bc	3.57 cde	6.55 ab	3.93 ab
BC _M ⁺ NPK _H	7.18 bc	4.23 a	8.65 a	5.47 a	6.29 ab	3.19 cd
BC _H ⁺ NPK _L	8.52 a	4.15 a	7.25 b	3.31 efg	6.70 ab	3.32 bcd
BC _H ⁺ NPK _H	8.63 a	3.56 ab	6.97 bc	4.00 bc	6.07 b	3.68 abc

Mean values presented in columns with different letters indicate significant differences at $p \leq 0.05$

Table 6. Summary of ANOVA (MS) for effect of time in years and treatments on all dependent variables

Source	FB	DB	PHA	PHB	NA	NB	CA	CB
Year	40444**	24051**	0.59**	0.63**	0.009*	8.69	1.10*	0.34**
Treatment	4917**	3614**	0.09**	0.057**	0.012**	8.70	14.23**	1.59**
Y* T	872**	549**	0.01**	0.007**	0.005*	8.70	1.27**	0.61**

* $p < 0.05$; ** $p < 0.01$

767 **Table 7.** Cumulative impact of different treatment on biomass and soil properties over three

Treatment	FB (g/m ²)	DB(g/m ²)	pHA	pHB	NA (%)	NB (%)	CA (%)	CB(%)
Control	109.82±2.91h	50.34±2.51 f	6.63±0.04 i	6.76±0.03 f	0.37±0.01 f	0.28±0.01 e	4.13±0.19 e	3.38±0.12 def
BCL	108.42±5.61 h	50.79±5.17f	6.85±0.03 fg	6.86±0.05 de	0.4±0.02 ef	0.32±0.01 cde	4.7±0.21 d	3.77±0.14 bcd
BCM	115.66±4.97 g	55.59±4.95ef	6.98±0.03 ab	6.95±0.06 bc	0.52±0.03 c	0.39±0.02 ab	7.01±0.32 ab	3.84±0.21 bc
BCH	116.05±5.64 g	58.48±4.89 e	6.99±0.05 a	7.02±0.03 a	0.5±0.04 cd	0.34±0.23bcd	6.12±0.2 c	3.33±0.17 ef
NPKL	140.92±10 f	69.86±5.32 d	6.76±0.07 h	6.87±0.04 de	0.66±0.03 a	0.41±0.02 a	5.08±0.13 d	2.82±0.11 g
NPKH	146.21±10.52 ef	74.11±6.92 d	6.8±0.06 gh	6.9±0.04 cd	0.44±0.04 de	0.34±0.02 bcd	3.97±0.13 e	3.07±0.08 fg
BCL+NPKL	148.56±10.65 e	83.95±8.69 c	6.95±0.05 abc	7.02±0.04 a	0.43±0.02 ef	0.32±0.01 cde	5.98±0.17 c	3.33±0.13 ef
BCL+NPKH	149.58±10.02 de	82.48±8.25 c	6.92±0.05 bcd	7.01±0.04 a	0.55±0.02 bc	0.37±0.01abc	7.1±0.17 ab	4.09±0.15 ab
BCM+NPKL	154.68±13.5 cd	92.66±12.45b	6.86±0.05 de	6.91±0.05 cd	0.38±0.03 ef	0.31±0.02 de	6.81±0.14 b	3.74±0.18 bcde
BCM+NPKH	178.52±18.03 a	114.49±15.2a	6.87±0.03 ef	6.84±0.06 f	0.59±0.03 b	0.38±0.02 ab	7.37±0.39 a	4.3±0.34 a
BCH+NPKL	156.71±13 c	86.76±8.92 c	6.89±0.03 ef	6.94±0.04 bc	0.52±0.02 c	0.36±0.03abc	6.99±0.12 ab	3.6±0.18 cde
BCH+NPKH	167.69±15.03 b	95.46±10.1b	6.93±0.03abc	6.98±0.02 ab	0.53±0.03 bc	0.33±0.02 cde	7.23±0.44 ab	3.75±0.12 bcde

768

769 Values are mean ± SE of three replication, Means with letters are not significantly different at $p = 0.01$ according to Duncan's multiple range test

770

771

Table 8. Principal components (PCs) for 8 traits biomass productivity and soil characteristics
(Varimax rotation)

Traits	Component		
	1	2	3
pHB	0.881	0.204	0.133
pHA	0.82	0.443	0.044
NB	0.604	-0.257	-0.103
CB	-0.024	0.89	-0.047
CA	0.391	0.797	0.305
DB	-0.102	-0.087	0.88
NA	0.162	0.098	0.798
FB	-0.017	0.479	0.642
Eigenvalue	2.00	1.98	1.95
Proportion $\sigma^2\%$	25.03	24.74	24.36
Cumulative $\sigma^2\%$	25.03	49.77	74.13

Table 9. Impact of application of biochar on soil microbial diversity (the Shannon index)

Treatment	Shannon Index of Diversity
CK	3.24±0.004
BC _M	3.25±0.003
NPK _H	3.23±0.002
BC _M ⁺ NPK _H	3.28±0.007

793
794

Table 10. X-ray photoelectron spectroscopy analysis of original and aged rice husk biochar

Name	Structure	Biochar stored at room temperatures for three years		Biochar extracted from the soil after three years	
		Peak BE	At%	Peak BE	At%
C1s A	– C=C- non-functionalised sp ² C	284.84	63.41	284.82	51.06
C1s B	– C-OH, C-O-C=, C-O-R	286.46	8.57	286.26	10.85
C1s C	– C-N, C=O	288.33	3.21	288.5	3.83
C1s d	– C=N, -N=C-O-	ND	ND	289.17	1.54
N1sA	Pyridne N	398.8	0.45	398.8	0.40
N1sB	N-H	400.7	0.55	400.7	0.60
Al2p		72.44	0.73	75.31	0.58
Ca2p		352.88	0.39	348.41	0.82
Fe2p		724.34	0.38	712.74	0.93
	FeOOH	711.2	0.30	711.2	0.65
	Fe(SO ₄) ₃	715.9	0.20	715.9	0.35
O1s		533.61	15.29	533.61	30.50
Mg1s		1305.35	0.37	1303.35	0.74
N1s		401.3	1.56	400.66	2.27
K2p		293.66	0.31	294.39	1.06
S2p		169.52	0.2	170.07	0.22
Si2p		104.69	5.58	103.61	11.19

795
796
797