THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Neuroticism, depressive symptoms and white-matter integrity in the Lothian Birth Cohort 1936

Citation for published version:

McIntosh, AM, Bastin, ME, Luciano, M, Munoz-Maniega, S, Valdes Hernandez, M, Royle, NA, Hall, J, Murray, C, Lawrie, SM, Starr, JM, Wardlaw, JM & Deary, IJ 2013, 'Neuroticism, depressive symptoms and white-matter integrity in the Lothian Birth Cohort 1936', *Psychological Medicine*, vol. 43, no. 6, pp. 1197-1206. <https://doi.org/10.1017/S003329171200150X>

Digital Object Identifier (DOI):

[10.1017/S003329171200150X](https://doi.org/10.1017/S003329171200150X)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Psychological Medicine

Publisher Rights Statement:

© McIntosh, A. M., Bastin, M. E., Luciano, M., Maniega, S. M., Del C Valdés Hernández, M., Royle, N. A., Hall, J., Murray, C., Lawrie, S. M., Starr, J. M., Wardlaw, J. M., & Deary, I. J. (2013). Neuroticism, depressive symptoms and white-matter integrity in the Lothian Birth Cohort 1936. *Psychological Medicine*, 43(6), 1197-1206doi: 10.1017/S003329171200150X

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Neuroticism, depressive symptoms and white-matter integrity in the Lothian Birth Cohort 1936

A. M. McIntosh^{1,2,3*}, M. E. Bastin^{2,3,4,5}, M. Luciano^{2,6}, S. Muñoz Maniega^{2,3,5},
M. del C. Valdés Hernández^{2,3,5}, N. A. Royle^{2,3,5}, J. Hall^{1,3}, C. Murray⁶, S. M. Lawrie^{1,3}, J. M. Starr^{2,7},
J. M. Wardlaw^{2,3,5} and I. J. Deary^{2,6}

¹ Division of Psychiatry, University of Edinburgh, UK

² Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, UK

³ SINAPSE (Scottish Imaging Network, A Platform for Scientific Excellence) collaboration

⁴ Division of Health Sciences (Medical Physics), University of Edinburgh, UK

⁵ Brain Research Imaging Centre, Division of Clinical Neurosciences, University of Edinburgh, UK

⁶ Department of Psychology, University of Edinburgh, UK

⁷ Geriatric Medicine Unit, University of Edinburgh, UK

Background. Clinical depression is associated with reductions in white-matter integrity in several long tracts of the brain. The extent to which these findings are localized or related to depressive symptoms or personality traits linked to disease risk remains unclear.

Method. Members of the Lothian Birth Cohort 1936 (LBC936) were assessed in two waves at mean ages of 70 and 73 years. At wave 1, they underwent assessments of depressive symptoms and the personality traits of neuroticism and extraversion. Brain diffusion magnetic resonance imaging (MRI) data were obtained at the second wave and mood assessments were repeated. We tested whether depressive symptoms were related to reduced white-matter tract fractional anisotropy (FA), a measure of integrity, and then examined whether high neuroticism or low extraversion mediated this relationship.

Results. Six hundred and sixty-eight participants provided useable data. Bilateral uncinate fasciculus FA was significantly negatively associated with depressive symptoms at both waves (standardized $\beta=0.12$ – 0.16). Higher neuroticism and lower extraversion were also significantly associated with lower uncinate FA bilaterally (standardized $\beta=0.09$ – 0.15) and significantly mediated the relationship between FA and depressive symptoms.

Conclusions. Trait liability to depression and depressive symptoms are associated with reduced structural connectivity in tracts connecting the prefrontal cortex with the amygdala and anterior temporal cortex. These effects suggest that frontotemporal disconnection is linked to the etiology of depression, in part through personality trait differences.

Received 9 March 2012; Revised 22 May 2012; Accepted 11 June 2012; First published online 11 July 2012

Key words: Depression, DTI, extraversion, MRI, neuroticism.

Introduction

Clinical depression is a common disorder that affects approximately 15% of the population at some time in their lives and shows a tendency to become recurrent or persistent (Thornicroft & Sartorius, 1993). The disorder is predicted to become the leading cause of worldwide disability (Murray & Lopez, 1997) and has significant costs for affected individuals, their families and society at large. Despite the considerable burden, the neurobiology of depression is poorly understood

although genetic factors and stressful life events both play a part in its etiology (Levinson, 2006).

Brain imaging has provided important insights into the biological basis of depression, with evidence of modest gray-matter reductions in frontotemporal regions using both region of interest and voxel-based methods (Drevets *et al.* 1997; Videbech, 1997; Kumar *et al.* 1998; Videbech & Ravnkilde, 2004). Depression has not, however, typically been associated with a single cortical or subcortical region (Arnone *et al.* 2012) and this could suggest that the disorder is caused by abnormal interactions between different brain regions due to disrupted connectivity. In support of this idea, some studies have found evidence of reduced white-matter integrity in the prefrontal and temporal lobes, and in several other regions (Taylor *et al.* 2007;

* Address for correspondence: Professor A. M. McIntosh, Division of Psychiatry, University of Edinburgh, Kennedy Tower, Royal Edinburgh Hospital, Edinburgh EH10 5HF, UK.
(Email: andrew.mcintosh@ed.ac.uk)

Cullen *et al.* 2010; Dalby *et al.* 2010). Many of these white-matter tracts, for example the uncinate fasciculus and cingulum, connect prefrontal areas with the medial temporal lobe. Disrupted connectivity of these emotion-related areas may therefore lead to emotional dysregulation and possibly an increased risk of depression. The extent to which these findings are secondary to the effects of clinical psychiatric illness or its treatment is unclear.

To overcome the issue of confounding by treatment-related factors, an alternative approach to studying clinical depression directly is to examine the relationship of white matter tract integrity to depressive symptoms and known risk factors in a non-clinical population cohort. Trait liability to depression has long been associated with neuroticism (Kendler *et al.* 1993a, 1994, 2002; Matthews *et al.* 2009), a heritable and relatively stable personality trait. Family and twin studies indicate that there is a high genetic correlation between neuroticism, clinical depression (Kendler *et al.* 2006) and depressive symptoms in non-clinical population samples (Ivkovic *et al.* 2007), with an overlap of more than 50% (Kendler *et al.* 1993b; Levinson, 2006) in their underlying architecture. Lower extraversion has also been proposed as a risk factor for depression, although the literature (Barnett *et al.* 2011) implicating this dimension of personality in the etiology of depression is sparse.

In the current study we first examined the relationship of white-matter tract fractional anisotropy (FA), derived from diffusion tensor imaging (DTI) as a proxy for tract integrity, with depressive symptoms in individuals from a population-based sample, the Lothian Birth Cohort 1936 (LBC1936). Second, we examined whether trait liability to depression was also associated with reductions in white-matter tract FA in the same tracts using the personality traits of (higher) neuroticism and (lower) extraversion. Third, because we hypothesized that reduced frontotemporal white-matter tract FA confers greater risk of depressive symptoms through effects on trait liability, we conducted a mediation analysis specifically to test this directional hypothesis.

Method

Subjects

The LBC1936 comprises 1091 community-dwelling relatively healthy individuals without dementia, residing around the city of Edinburgh, Scotland, most of whom had participated in the Scottish Mental Survey 1947 (SMS1947). The LBC1936 undertook personality, medical and cognitive testing at a mean age of about 70 years (for full details on participant recruitment

and cognitive testing, see Deary *et al.* 2007). Subjects were then reassessed approximately 3.2 years later at which time they received additional investigations, including detailed structural brain magnetic resonance imaging (MRI) (Wardlaw *et al.* 2011), and retook the medical and cognitive assessments, including the Mini Mental State Examination (MMSE) of cognitive function (Folstein *et al.* 1975). Participants lived independently in the community around the city of Edinburgh, Scotland, and were able to travel on both occasions to the Wellcome Trust Clinical Research Facility in Edinburgh for testing. The study was approved by the Lothian (REC 07/MRE00/58) and Scottish Multicentre (MREC/01/0/56) Research Ethics Committees and all subjects gave written informed consent.

Depressive symptoms and personality traits

Hospital Anxiety and Depression Scale (HADS; Zigmond & Snaith, 1983) scores were available from the assessments performed at 70 and 73 years of age. These assessments were available separately as depression (HADS-D) and anxiety (HADS-A) measures with seven self-report items for each subscale. Neuroticism and extraversion were measured at the first assessment using the Neuroticism, Extraversion and Openness Five Factor Inventory (NEO-FFI; Costa & McCrae, 1989), a self-report scale consisting of 12 items for each NEO-FFI personality trait. The other three traits (Openness, Agreeableness and Conscientiousness) were not included in the analyses in the current study.

Diffusion MRI

Diffusion MRI was performed on a GE Signa HDXt 1.5-T clinical scanner (General Electric, USA) using a self-shielding gradient set with maximum gradient strength of 33 mT m^{-1} , and a manufacturer-supplied eight-channel phased-array head coil. Single-shot, spin-echo echo-planar diffusion-weighted volumes ($b = 1000 \text{ s mm}^{-2}$) were acquired in 64 non-collinear directions, along with seven T_2 -weighted volumes ($b = 0 \text{ s mm}^{-2}$). Seventy-two contiguous axial slices of thickness 2 mm were acquired with a field of view of $256 \times 256 \text{ mm}$ and matrix size of 128×128 , giving a resolution of $2 \times 2 \times 2 \text{ mm}$. The repetition time was 16.5 s and echo time 95.5 ms, producing a total scan time of approximately 20 min (see Wardlaw *et al.* 2011 for full details of the imaging protocol).

White-matter tractography

Datasets were converted into NIfTI-1 (<http://nifti.nimh.nih.gov/nifti-1>) format and preprocessed using

Fig. 1. Representative white-matter tracts obtained from a single individual.

FSL tools (FMRIB, Oxford, UK; www.fmrib.ox.ac.uk/) to extract the brain, remove bulk motion and eddy current-induced artifacts and estimate water diffusion tensor parameters. Brain connectivity data were created using the BEDPOSTX/ProbTrackX tractography algorithm with a two-fiber model and 5000 streamlines to reconstruct tracts of interest. An automatic tract selection method with good reproducibility (Clayden *et al.* 2009a), based on a model of tract topology (Clayden *et al.* 2007), was used to generate equivalent tracts of interest in each subject. This technique, termed probabilistic neighborhood tractography (PNT) and implemented in the TractoR package (<http://github.com/jonclayden/tractor/wiki/>), optimizes the choice of seed point for tractography by estimating the best matching tract from a series of candidate tracts generated from a neighborhood of voxels placed around a voxel transferred from standard space against a reference tract that was derived from a digital human white-matter atlas (Mori & van Zijl, 2007). The topological tract model was also used to reject any false positive connections (Clayden *et al.* 2009b), thereby significantly improving tract segmentation (Maniega *et al.* 2008). As displayed in Fig. 1, 12 white-matter tracts were segmented, namely the genu and splenium of corpus callosum, cingulum cingulate gyri (CCG) (Clayden *et al.* 2007), anterior thalamic radiations (ATR), uncinate, arcuate and inferior longitudinal fasciculi (ILF). For each subject, the seed point that produced the best match tract to the reference for each of the 12 pathways was determined, with the resulting tractography mask applied to each subject's FA volume. Tract-averaged FA values were calculated from these masks and used in all subsequent analyses. To ensure that the segmented tracts

were anatomically plausible representations of the fasciculi of interest, a researcher (S.M.M.) visually inspected all masks blind to the other study variables and excluded tracts with aberrant or truncated pathways.

Statistical analyses

Statistical analyses were conducted in the R statistical software package (www.r-project.org/) using linear regression. To meet underlying statistical assumptions, we applied a square root transformation to the HADS symptom measures. Initially, we entered the transformed HADS scores as the dependent variable and age at symptom assessment, sex and white-matter tract FA as the independent variables (analysis 1). To correct for multiple testing, we applied a false discovery rate (FDR; Benjamini & Hochberg, 1995) correction to these initial associations. Subsequently, we examined whether higher neuroticism or lower extraversion were also related to white-matter tract FA using the same model, substituting these personality measures as the dependent variables (analysis 2). We then examined whether previous antidepressant treatment or a history of stroke confounded any significant relationships between FA and depression or personality measures by conducting a series of planned additional analyses. We then adjusted for the effects of antidepressant treatment, current or past cigarette use or stroke by including these as covariates in the statistical models. Finally, to assess whether neuroticism/extraversion could account for any associations between depression and white-matter tract FA, we examined the influence of these variables on analysis 1. Specifically, we noted the

Table 1. Demographic and historical information

Male sex, <i>n</i> (%)	356 (53)	
Age at assessment 1 (years), mean (s.d.)	69.5 (0.8)	
Age at assessment 2 (imaging) (years), mean (s.d.)	72.7 (0.7)	
Antidepressants at assessment 1, <i>n</i> (%)	23 (3.4)	
Antidepressants at assessment 2, <i>n</i> (%)	30 (4.5)	
Diabetes, <i>n</i> (%)	72 (10.8)	
Hypertension, <i>n</i> (%)	323 (48.4)	
MMSE score < 22, <i>n</i> (%)	1 (0.1)	
MMSE score < 27, <i>n</i> (%)	43 (6.4)	
MMSE score, median (IQR)	29 (28–30)	
HADS Depression – assessment 1, mean (s.d.)	2.6 (2.2)	
HADS Depression – assessment 2, mean (s.d.)	2.6 (2.2)	
HADS Anxiety – assessment 1, mean (s.d.)	4.8 (3.1)	
HADS Anxiety – assessment 2, mean (s.d.)	4.4 (3.2)	
NEO-FFI Neuroticism – assessment 1, mean (s.d.)	16.6 (7.6)	
NEO-FFI Extraversion – assessment 1, mean (s.d.)	27.3 (5.9)	
<i>Tractography data (n = 668)</i>		
Tract	Complete <i>n</i> (%)	Mean FA (s.d.)
Genu of corpus callosum	646 (96.7)	0.41 (0.046)
Splenium of corpus callosum	663 (99.3)	0.49 (0.069)
Left arcuate fasciculus	639 (95.7)	0.45 (0.042)
Right arcuate fasciculus	580 (86.8)	0.43 (0.043)
Left anterior thalamic radiation	556 (83.2)	0.32 (0.033)
Right anterior thalamic radiation	643 (96.3)	0.33 (0.047)
Left uncinate fasciculus	567 (84.9)	0.33 (0.030)
Right uncinate fasciculus	628 (94.0)	0.33 (0.032)
Left inferior longitudinal fasciculus	663 (99.3)	0.40 (0.048)
Right inferior longitudinal fasciculus	664 (99.4)	0.38 (0.048)
Left cingulum cingulate gyrus	641 (96.0)	0.44 (0.047)
Right cingulum cingulate gyrus	650 (97.3)	0.39 (0.044)

MMSE, Mini Mental State Examination; HADS, Hospital Anxiety and Depression Scale; NEO-FFI, Neuroticism, Extraversion and Openness Five Factor Inventory; FA, fractional anisotropy; s.d., standard deviation; IQR, interquartile range.

attenuation in the effect of white-matter tract FA after neuroticism/extraversion was included in the model, and conducted a formal mediation analysis using the Sobel test (Baron & Kenny, 1986).

Results

A total of 668 individuals from the LBC1936 provided useable tractography data from the 740 who presented for MRI. As indicated in Table 1, the average age of the subjects was 72.7 (s.d.=0.7) years at the time of the imaging examination and 356 (53%) were male. The rates of antidepressant prescribing were lower than in the general Scottish population (see www.isdscotlandarchive.scot.nhs.uk/isd/6517.html), approximately 10% of the sample reported a diagnosis of diabetes and 48.4% reported a diagnosis of hypertension. MMSE scores (Folstein *et al.* 1975) showed that only one subject met a specific threshold for organic brain disorder whereas 6.4% of the sample met a sensitive threshold. At both assessments, the

HADS-D score correlated moderately and positively with neuroticism (Pearson's $r=0.35$ at age 69.5, and $r=0.35$ at age 72.7) and negatively with extraversion (Pearson's $r=-0.38$ at age 69.5, and $r=-0.34$ at age 72.7). Table 1 also shows that the number of individual tracts that did not meet quality criteria (truncation or failed to follow expected path) ranged from 0.6% for the right inferior longitudinal fasciculus to 16.8% for the left anterior thalamic radiation.

We first examined the relationship of the HADS-D subscale scores with FA in each tract (Table 2). Depressive symptoms measured at the time of the imaging assessment were significantly and negatively associated with FA in the left and right uncinate fasciculus (standardized $\beta=-0.12$, $p=0.0038$ in the left hemisphere; standardized $\beta=-0.13$, $p=0.0011$ in the right hemisphere). Depressive symptoms measured at the first assessment 3 years earlier were also significantly and negatively associated with bilateral uncinate FA, with similar effect sizes (standardized $\beta=-0.16$, $p=0.0002$ in the left

Table 2. Association between white-matter tract FA and symptoms of depression

	Assessment wave 1					Assessment wave 2				
	β	S.E.(β)	Standard β	t value	Pr(> t)	β	S.E.(β)	Standard β	t value	Pr(> t)
Genu	-0.09	0.59	-0.01	-0.153	0.878	0.53	0.60	0.03	0.877	0.381
Splenium	-0.85	0.40	-0.08	-2.159	0.031	-0.46	0.40	-0.04	-1.149	0.251
LArc	-0.57	0.65	-0.03	-0.876	0.381	-0.88	0.66	-0.05	-1.325	0.186
RArc	-0.74	0.69	-0.04	-1.071	0.285	-0.45	0.69	-0.03	-0.658	0.511
LATR	-0.57	0.89	-0.03	-0.641	0.522	-0.38	0.92	-0.02	-0.411	0.681
RATR	-1.24	0.84	-0.06	-1.479	0.140	-1.06	0.85	-0.05	-1.251	0.211
LCCG	-1.13	0.58	-0.08	-1.933	0.054	-1.26	0.60	-0.08	-2.122	0.034
RCCG	-1.03	0.62	-0.07	-1.663	0.097	-1.51	0.63	-0.09	-2.383	0.017
LUnc	-3.57	0.97	-0.16	-3.700	<0.001*	-2.89	0.99	-0.12	-2.904	0.004*
RUnc	-3.22	0.86	-0.15	-3.752	<0.001*	-2.86	0.87	-0.13	-3.293	0.001*
LILF	-1.36	0.57	-0.09	-2.364	0.018	-0.71	0.59	-0.05	-1.211	0.226
RILF	-0.56	0.57	-0.04	-0.983	0.326	-0.61	0.58	-0.04	-1.040	0.299

FA, Fractional anisotropy; L, left; R, right; Arc, arcuate fasciculus; ATR, anterior thalamic radiation; CCG, cingulum cingulate gyrus; Unc, uncinate fasciculus; ILF, inferior longitudinal fasciculus; S.E., standard error.

Bold type indicates effects that were statistically significant at both study assessments.

* False discovery rate (FDR) <0.05.

hemisphere; standardized $\beta = -0.15$, $p = 0.0002$ in the right hemisphere). The association between uncinate fasciculus FA and depressive symptoms survived FDR correction bilaterally at both assessments. There were no significant associations between HADS-A scores and FA in any tract (Supplementary Table S1).

We next examined the relationship between neuroticism and extraversion, trait measures of liability to depression that were measured at the first assessment, and mean white-matter FA in each tract determined at the second assessment 3 years later (Table 3). There was a significant and negative association between neuroticism scores and FA in the uncinate fasciculus bilaterally (standardized $\beta = -0.12$, $p = 0.0083$ in the left hemisphere; standardized $\beta = -0.09$, $p = 0.032$ in the right hemisphere). Extraversion was significantly and positively correlated with uncinate fasciculus FA in the right hemisphere only (standardized $\beta = 0.13$, $p = 0.0029$), although the association with FA in the left uncinate fasciculus narrowly failed to reach conventional statistical significance (standardized $\beta = 0.09$, $p = 0.053$). Two of these associations, namely a negative association between left uncinate FA and neuroticism and a positive association between right uncinate FA and extraversion, remained significant after FDR adjustment across all 12 tracts assessed.

We then reanalyzed the data after the graphical detection and subsequent removal of a single outlying observation with an FA value of <0.2, with no effect on any of our significant findings, all of which remained significant. We then adjusted for the effects

of antidepressants prescribed at time 1 or time 2, the presence of a self-reported history of stroke, current alcohol use at the time of the scan, and whether the individual reported a history of past or current tobacco use. In each case the relationship between uncinate white-matter integrity remained in the same direction as reported in the whole sample (see online Supplementary material). However, the relationship between right uncinate white-matter integrity and neuroticism became non-significant.

We then assessed the effect of adding neuroticism or extraversion to a statistical model where tract FA and age and sex were already fitted as independent variables and depression was the dependent variable. The purpose of this analysis was to examine whether the addition of neuroticism/extraversion attenuated the effect of uncinate tract FA on depressive symptoms, indicating that neuroticism was more closely associated with depression and potentially mediating the association of tract FA with depressive symptoms. The addition of neuroticism led to a significantly improved model fit in both the left (see online Supplementary Table S2) and right hemispheres. The regression coefficient estimates for the effect of uncinate tract FA on depressive symptoms were somewhat attenuated in both the left hemispheres after the inclusion of neuroticism in the model (see Supplementary Table S2). Similarly, the addition of extraversion to a model where depressive symptoms was the dependent variable and right uncinate FA was the independent variable also improved the model fit whereas the effect of tract FA on depressive symptoms

Table 3. Association between white-matter tract FA, neuroticism and extraversion

	Neuroticism					Extraversion				
	β	S.E.(β)	Standard β	t value	Pr(> t)	β	S.E.(β)	Standard β	t value	Pr(> t)
Genu	-4.98	6.84	-0.03	-0.729	0.467	4.21	5.45	0.03	0.773	0.440
Splenium	-6.64	4.54	-0.06	-1.463	0.144	2.65	3.60	0.08	0.737	0.462
LArc	-6.26	7.30	-0.04	-0.857	0.392	5.12	5.82	0.04	0.879	0.380
RArc	-1.27	7.70	-0.01	-0.165	0.869	4.08	6.17	0.03	0.661	0.509
LATR	-19.93	9.88	-0.09	-2.017	0.044	17.60	7.86	0.10	2.240	0.025
RATR	-18.59	9.41	-0.08	-1.974	0.049	2.13	7.46	0.01	0.286	0.779
LCCG	-21.78	6.53	-0.14	-3.334	<0.001*	9.22	5.29	0.07	1.742	0.082
RCCG	-17.41	7.21	-0.10	-2.417	0.016	11.98	5.78	0.09	2.074	0.039
LUnc	-28.61	10.80	-0.12	-2.65	0.008*	16.70	8.61	0.09	1.939	0.053
RUnc	-21.04	9.77	-0.09	-2.154	0.032	23.31	7.78	0.13	2.996	0.003*
LILF	-7.09	6.58	-0.04	-1.078	0.282	14.26	5.22	0.11	2.730	0.007*
RILF	-7.75	6.49	-0.05	-1.194	0.233	10.41	5.09	0.08	2.047	0.041

FA, Fractional anisotropy; L, left; R, right; Arc, arcuate fasciculus; ATR, anterior thalamic radiation; CCG, cingulum cingulate gyrus; Unc, uncinate fasciculus; ILF, inferior longitudinal fasciculus.

Bold text used to highlight effects on Unc.

* False discovery rate (FDR) <0.05.

was also somewhat attenuated (see supplementary Table S2).

Mediation analysis supported our hypothesis that neuroticism and extraversion partly mediate the relationship between tract FA and depression. Specifically, Sobel tests showed a significant mediating effect of neuroticism between lower uncinate fasciculus FA and depressive symptoms at the first wave of assessment in both the left (test for mediation: $Z = -2.38$, $p = 0.017$) and right (test for mediation: $Z = -2.03$, $p = 0.042$) hemispheres. Extraversion also significantly partly mediated the relationship between right uncinate fasciculus FA and depressive symptoms in the right hemisphere (test for mediation: $Z = -2.90$, $p = 0.004$) but not the left (test for mediation: $Z = -1.91$, $p = 0.056$). In each case, the mediating effect of neuroticism was partial, as the (albeit attenuated) effect of white-matter integrity on depressive symptoms remained significant in each case.

Discussion

Using data from the LBC1936 we found that there were significant negative associations between depressive symptoms and white-matter integrity in the bilateral uncinate fasciculus. This relationship was highly significant when the depressive symptoms were rated either at the time of the imaging assessment or 3 years earlier. Subsequently we found that the personality traits of neuroticism and extraversion also have significant associations with white-matter integrity in the uncinate, consistent with their

risk-conferring effects on depression. The association between depressive symptoms and white-matter integrity was slightly and significantly attenuated after the inclusion of neuroticism or extraversion in the model. Sobel analyses confirmed that neuroticism and extraversion partially mediated the relationship between white-matter integrity and depression. These associations were significant after adjustment for age and sex, after correction for multiple testing, and were robust to planned sensitivity analyses that took into account previous antidepressant treatment or stroke. These findings suggest that the effect of reduced uncinate white-matter integrity on depressive symptoms is conferred partially through effects on extraversion and neuroticism that have been shown to increase liability to depression in other studies.

No significant associations were found between depressive symptoms and white-matter integrity in any other tract, nor were there any associations between anxiety rating scale scores and white-matter integrity. By contrast, there was a significant negative association (surviving FDR correction) between left cingulum white-matter integrity and neuroticism. Similarly, there was a significant positive association between left ILF integrity and extraversion. To the best of our knowledge, neither of these associations has been reported previously in the literature. The absence of a relationship with anxiety is superficially surprising, as anxiety frequently accompanies depressive symptoms in the clinical disorder. The absence of a relationship between white-matter integrity and anxiety implies a more specific relationship with

depressive symptoms, and suggests that anxiety symptoms are associated with a smaller detectable signal or that their neural correlates are more detectable as changes in gray-matter structure or brain function.

The uncinate fasciculus connects the prefrontal cortex to limbic and emotional regions of the temporal lobe, such as the amygdala and anterior temporal cortex. These areas are implicated in the encoding of rewarding or punishing stimuli and in normal emotional processing and regulation in both the animal (Cardinal *et al.* 2002) and human (Phillips *et al.* 2003) literature. Abnormalities in these tracts have been found in studies of unipolar depression (Taylor *et al.* 2007; Cullen *et al.* 2010; Dalby *et al.* 2010) and people at high familial risk of major mental illness (Huang *et al.* 2011). There is also a growing body of literature showing abnormal frontotemporal activation and connectivity in depression (Anand *et al.* 2005a,b) and in a variety of other mental disorders (Lawrie *et al.* 2002; Chepenik *et al.* 2010) for which neuroticism is an identified risk factor (Maier *et al.* 1994; van Os & Jones, 2001; Angst *et al.* 2003; Barnett *et al.* 2011).

Few studies have directly examined the relationship of white-matter tract FA to neuroticism, although recently a negative association between harm avoidance and measures of tract integrity was reported in several long white-matter tracts (Westlye *et al.* 2011). Structural and functional abnormalities of frontotemporal brain regions have, however, been linked to specific genetic risk factors for depression and other mental illnesses. These studies include several examining the role of the 5-HTTLPR polymorphism (Ball *et al.* 1997), in which the s allele confers reduced frontotemporal functional connectivity (Pezawas *et al.* 2005), increased neuroticism (Barnett *et al.* 2011) and an increased risk of depression (Caspi *et al.* 2003). The current findings extend that literature to imply that the personality traits of neuroticism and extraversion may mediate the effects of reduced frontotemporal connectivity on the risk of depressive symptoms.

Although no relationship between left ILF FA and depressive symptoms was found in the current study, left ILF FA was significantly and positively related to extraversion. The ILF connects occipital brain areas with the temporal lobe whereas the cingulum connects the medial prefrontal cortex to the hippocampus. There is evidence that the integrity of these tracts is also associated with liability to depression (Sheline *et al.* 2008), bipolar disorder (Chaddock *et al.* 2009; Zanetti *et al.* 2009; Sprooten *et al.* 2011) and schizophrenia. It is also possible that reductions in tract integrity in these regions may have important consequences for cognitive function in non-demented

adults (Kantarci *et al.* 2011), a hypothesis that could be addressed in the current cohort at a later imaging assessment.

The current study is the largest investigation to date of depressive symptoms and white-matter tract FA as a proxy for integrity in any non-clinical sample. It should be noted that <10 people in the current study met the specific HADS-D threshold for clinical depression. These thresholds have been largely determined in healthcare-seeking populations, so the positive predictive value is likely to be higher in the current sample where the prevalence of depression is likely to be lower. The association of depressive symptoms with white-matter FA in a non-clinical cohort is likely to have broader applicability to clinical depression for several reasons. First, the symptoms of clinical depression are identical to those captured by the items of most depression-rating scales, albeit without the usual 2-week duration criterion needed to make a clinical diagnosis. However, the choice of a 2-week duration criterion for depression is arbitrary and has limited validity, with shorter duration of symptoms predicting depression in both affected subjects and co-twins (Kendler & Gardner, 1998). Depressive symptoms measured at a single episode also show similar heritability estimates to clinically defined syndromal depression (McGue & Christensen, 1997) and higher estimates of heritability when measured repeatedly over time (McGue & Christensen, 2003). Furthermore, there is evidence that major and subthreshold 'minor' depression share a common neural basis (Kumar *et al.* 1998). Overall, these findings suggest that scale-measured depressive symptoms are associated with trait liability to depression and are not merely measures of current mental state.

A further limitation of this study is the relatively small effect sizes that were found for the relationships between white-matter integrity, depression and the personality traits of neuroticism and depression. All scales for personality and depressive symptoms have strengths and limitations and each only approximates the measurement of the underlying latent variables. The estimates given here may therefore underestimate the true relationships, which cannot be measured in an error-free way. Finally, in the present sample, neuroticism and extraversion were correlated with one another (Pearson's $r = -0.37$) and it is possible that both measures may be measuring a common underlying trait. Future studies should explore the facets of each personality measure that are independently associated with white-matter integrity, and also perhaps mediate the association with symptoms of depression.

The current study is the first large-scale investigation showing an association between frontotemporal white-matter tract FA, depression and trait

liability to clinical mood disorder. These findings are not due to differences in age, sex or the secondary effects of antidepressant treatment, and suggest that variation in frontotemporal interactions may underlie depressive symptoms at a population level, and may also apply to clinical populations.

Supplementary material

For supplementary material accompanying this paper visit <http://dx.doi.org/10.1017/S003329171200150X>.

Acknowledgments

Funding from the following bodies is acknowledged: The Disconnected Mind Project (www.disconnect.edmind.ed.ac.uk/) funded by Age UK and the UK Medical Research Council (MRC), the Row Fogo Trust, the Carnegie Trust for the Universities of Scotland, the Scottish Funding Council (SFC), which funds the SINAPSE Collaboration (www.sinapse.ac.uk), and The Centre for Cognitive Ageing and Cognitive Epidemiology (G0700704/84698; www.ccace.ed.ac.uk/), part of the cross-council Lifelong Health and Wellbeing Initiative (LLWH). Funding from the Biotechnology and Biological Sciences Research Council (BBSRC), the Engineering and Physical Sciences Research Council (EPSRC), the Economic and Social Research Council (ESRC) and the MRC is also gratefully acknowledged. Personality, medical and cognitive testing was performed at the Wellcome Trust Clinical Research Facility, Edinburgh (www.wtcrf.ed.ac.uk/), and MRI was performed at the Brain Research Imaging Centre, University of Edinburgh (www.sbirc.ed.ac.uk/). M.L. is a Royal Society of Edinburgh/Lloyds TSB Foundation for Scotland Personal Research Fellow. J.H. is supported by a Scottish Senior Clinical Fellowship. A.M.M. is supported by the Brain and Behavior Research Fund (formerly NARSAD) through an Independent Investigator Award and by the Health Foundation through a Clinician Scientist Fellowship.

Declaration of Interest

A.M.M. and S.M.L. report having received research support from Pfizer Pharmaceuticals and consultancy fees from Roche Pharmaceuticals. J.H. reports having received research funding from Pfizer Pharmaceuticals.

References

Anand A, Li Y, Wang Y, Wu J, Gao S, Bukhari L, Mathews VP, Kalnin A, Lowe MJ (2005a). Activity and

connectivity of brain mood regulating circuit in depression: a functional magnetic resonance study. *Biological Psychiatry* **57**, 1079–1088.

Anand A, Li Y, Wang Y, Wu J, Gao S, Bukhari L, Mathews VP, Kalnin A, Lowe MJ (2005b). Antidepressant effect on connectivity of the mood-regulating circuit: an fMRI study. *Neuropsychopharmacology* **30**, 1334–1344.

Angst J, Gamma A, Endrass J (2003). Risk factors for the bipolar and depression spectra. *Acta Psychiatrica Scandinavica* (Suppl.) **418**, 15–19.

Arnold D, McIntosh AM, Ebmeier KP, Munafò MR, Anderson IM (2012). Magnetic resonance imaging studies in unipolar depression: systematic review and meta-regression analyses. *European Neuropsychopharmacology* **22**, 1–16.

Ball D, Hill L, Freeman B, Eley TC, Strelau J, Riemann R, Spinath FM, Angleitner A, Plomin R (1997). The serotonin transporter gene and peer-rated neuroticism. *NeuroReport* **8**, 1301–1304.

Barnett JH, Huang J, Perlis RH, Young MM, Rosenbaum JF, Nierenberg AA, Sachs G, Nimgaonkar VL, Miklowitz DJ, Smoller JW (2011). Personality and bipolar disorder: dissecting state and trait associations between mood and personality. *Psychological Medicine* **41**, 1593–1604.

Baron RM, Kenny DA (1986). The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology* **51**, 1173–1182.

Benjamini Y, Hochberg Y (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**, 289–300.

Cardinal RN, Parkinson JA, Hall J, Everitt BJ (2002). Emotion and motivation: the role of the amygdala, ventral striatum, and prefrontal cortex. *Neuroscience and Biobehavioral Reviews* **26**, 321–352.

Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H, McClay J, Mill J, Martin J, Braithwaite A, Poulton R (2003). Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science* **301**, 386–389.

Chaddock CA, Barker GJ, Marshall N, Schulze K, Hall MH, Fern A, Walshe M, Bramon E, Chitnis XA, Murray R, McDonald C (2009). White matter microstructural impairments and genetic liability to familial bipolar I disorder. *British Journal of Psychiatry* **194**, 527–534.

Chepenik LG, Raffo M, Hampson M, Lacadie C, Wang F, Jones MM, Pittman B, Skudlarski P, Blumberg HP (2010). Functional connectivity between ventral prefrontal cortex and amygdala at low frequency in the resting state in bipolar disorder. *Psychiatry Research* **182**, 207–210.

Clayden JD, King MD, Clark CA (2009a). Shape modelling for tract selection. *Medical Image Computing and Computer-Assisted Intervention* **12**, 150–157.

Clayden JD, Storkey AJ, Bastin ME (2007). A probabilistic model-based approach to consistent white matter tract segmentation. *IEEE Transactions on Medical Imaging* **26**, 1555–1561.

- Clayden JD, Storkey AJ, Muñoz Maniega S, Bastin ME (2009b). Reproducibility of tract segmentation between sessions using an unsupervised modelling-based approach. *NeuroImage* **45**, 377–385.
- Costa PT, McCrae RR (1989). *The NEO-PI/NEO-FFI Manual Supplement*. Psychological Assessment Resources: Odessa, FL.
- Cullen KR, Klimes-Dougan B, Muetzel R, Mueller BA, Camchong J, Hourii A, Kurma S, Lim KO (2010). Altered white matter microstructure in adolescents with major depression: a preliminary study. *Journal of the American Academy of Child and Adolescent Psychiatry* **49**, 173–83e1.
- Dalby RB, Frandsen J, Chakravarty MM, Ahdidan J, Sorensen L, Rosenberg R, Videbech P, Ostergaard L (2010). Depression severity is correlated to the integrity of white matter fiber tracts in late-onset major depression. *Psychiatry Research* **184**, 38–48.
- Deary IJ, Gow AJ, Taylor MD, Corley J, Brett C, Wilson V, Campbell H, Whalley LJ, Visscher PM, Porteous DJ, Starr JM (2007). The Lothian Birth Cohort 1936: a study to examine influences on cognitive ageing from age 11 to age 70 and beyond. *BMC Geriatrics* **7**, 28.
- Drevets WC, Price JL, Simpson Jr. JR, Todd RD, Reich T, Vannier M, Raichle ME (1997). Subgenual prefrontal cortex abnormalities in mood disorders. *Nature* **386**, 824–827.
- Folstein MF, Folstein SE, McHugh PR (1975). 'Mini-mental state'. A practical method for grading the cognitive state of patents for the clinician. *Journal of Psychiatric Research* **12**, 189–198.
- Huang H, Fan X, Williamson DE, Rao U (2011). White matter changes in healthy adolescents at familial risk for unipolar depression: a diffusion tensor imaging study. *Neuropsychopharmacology* **36**, 684–691.
- Ivkovic V, Vitart V, Rudan I, Janicijevic B, Smolej-Narancic N, Skaric-Juric T, Barbalic M, Polasek O, Kolcic I, Biloglav Z, Visscher PM, Hayward C, Hastie ND, Anderson N, Campbell H, Wright AF, Rudan P, Deary IJ (2007). The Eysenck personality factors: psychometric structure, reliability, heritability and phenotypic and genetic correlations with psychological distress in an isolated Croatian population. *Personality and Individual Differences* **42**, 123–133.
- Kantarci K, Senjem ML, Avula R, Zhang B, Samikoglu AR, Weigand SD, Przybelski SA, Edmonson HA, Vemuri P, Knopman DS, Boeve BF, Ivnik RJ, Smith GE, Petersen RC, Jack Jr. CR (2011). Diffusion tensor imaging and cognitive function in older adults with no dementia. *Neurology* **77**, 26–34.
- Kendler KS, Gardner CO (1998). Boundaries of major depression: an evaluation of DSM-IV criteria. *American Journal of Psychiatry* **155**, 172–177.
- Kendler KS, Gardner CO, Prescott CA (2002). Toward a comprehensive developmental model for major depression in women. *American Journal of Psychiatry* **159**, 1133–1145.
- Kendler KS, Gatz M, Gardner CO, Pedersen NL (2006). Personality and major depression: a Swedish longitudinal, population-based twin study. *Archives of General Psychiatry* **63**, 1113–1120.
- Kendler KS, Neale MC, Kessler RC, Heath AC, Eaves LJ (1993a). The lifetime history of major depression in women. Reliability of diagnosis and heritability. *Archives of General Psychiatry* **50**, 863–870.
- Kendler KS, Neale MC, Kessler RC, Heath AC, Eaves LJ (1993b). A longitudinal twin study of personality and major depression in women. *Archives of General Psychiatry* **50**, 853–862.
- Kendler KS, Neale MC, Kessler RC, Heath AC, Eaves LJ (1994). The clinical characteristics of major depression as indices of the familial risk to illness. *British Journal of Psychiatry* **165**, 66–72.
- Kumar A, Jin Z, Bilker W, Udupa J, Gottlieb G (1998). Late-onset minor and major depression: early evidence for common neuroanatomical substrates detected by using MRI. *Proceedings of the National Academy of Sciences USA* **95**, 7654–7658.
- Lawrie SM, Buechel C, Whalley HC, Frith CD, Friston KJ, Johnstone EC (2002). Reduced frontotemporal functional connectivity in schizophrenia associated with auditory hallucinations. *Biological Psychiatry* **51**, 1008–1011.
- Levinson DF (2006). The genetics of depression: a review. *Biological Psychiatry* **60**, 84–92.
- Maier W, Minges J, Lichtermann D, Heun R, Franke P (1994). Personality variations in healthy relatives of schizophrenics. *Schizophrenia Research* **12**, 81–88.
- Maniega SM, Bastin ME, McIntosh AM, Lawrie SM, Clayden JD (2008). Atlas-based reference tracts improve automatic white matter segmentation with neighbourhood tractography. In *Proceedings of the 16th Scientific Meeting of the ISMRM*, p. 3318. International Society for Magnetic Resonance in Medicine, Berkeley, CA, USA.
- Matthews G, Deary IJ, Whiteman MC (2009). *Personality Traits*. Cambridge University Press: Cambridge.
- McGue M, Christensen K (1997). Genetic and environmental contributions to depression symptomatology: evidence from Danish twins 75 years of age and older. *Journal of Abnormal Psychology* **106**, 439–448.
- McGue M, Christensen K (2003). The heritability of depression symptoms in elderly Danish twins: occasion-specific versus general effects. *Behavior Genetics* **33**, 83–93.
- Mori S, van Zijl P (2007). Human white matter atlas. *American Journal of Psychiatry* **164**, 1005.
- Murray CJL, Lopez AD (1997). Alternative projections of mortality and disability by cause 1990–2020: Global Burden of Disease Study. *Lancet* **349**, 1498–1504.
- Pezawas L, Meyer-Lindenberg A, Drabant EM, Verchinski BA, Munoz KE, Kolachana BS, Egan MF, Mattay VS, Hariri AR, Weinberger DR (2005). 5-HTTLPR polymorphism impacts human cingulate-amygdala interactions: a genetic susceptibility mechanism for depression. *Nature Neuroscience* **8**, 828–834.
- Phillips ML, Drevets WC, Rauch SL, Lane R (2003). Neurobiology of emotion perception I: The neural basis of normal emotion perception. *Biological Psychiatry* **54**, 504–514.
- Sheline YI, Price JL, Vaishnavi SN, Mintun MA, Barch DM, Epstein AA, Wilkins CH, Snyder AZ, Couture L, Schechtman K, McKinstry RC (2008). Regional white matter hyperintensity burden in automated segmentation

- distinguishes late-life depressed subjects from comparison subjects matched for vascular risk factors. *American Journal of Psychiatry* **165**, 524–532.
- Sproueten E, Sussmann JE, Clugston A, Peel A, McKirdy J, William T, Moorhead J, Anderson S, Shand AJ, Giles S, Bastin ME, Hall J, Johnstone EC, Lawrie SM, McIntosh AM** (2011). White matter integrity in individuals at high genetic risk of bipolar disorder. *Biological Psychiatry* **70**, 350–356.
- Taylor WD, MacFall JR, Gerig G, Krishnan RR** (2007). Structural integrity of the uncinate fasciculus in geriatric depression: relationship with age of onset. *Neuropsychiatric Disease and Treatment* **3**, 669–674.
- Thornicroft G, Sartorius N** (1993). The course and outcome of depression in different cultures: 10-year follow-up of the WHO collaborative study on the assessment of depressive disorders. *Psychological Medicine* **23**, 1023–1032.
- van Os J, Jones PB** (2001). Neuroticism as a risk factor for schizophrenia. *Psychological Medicine* **31**, 1129–1134.
- Videbech P** (1997). MRI findings in patients with affective disorder: a meta-analysis. *Acta Psychiatrica Scandinavica* **96**, 157–168.
- Videbech P, Ravnkilde B** (2004). Hippocampal volume and depression: a meta-analysis of MRI studies. *American Journal of Psychiatry* **161**, 1957–1966.
- Wardlaw JM, Bastin ME, Valdes Hernandez MC, Muñoz Maniega S, Royle NA, Morris Z, Clayden JD, Sandeman EM, Eadie E, Murray C, Starr JM, Deary IJ** (2011). Brain ageing, cognition in youth and old age, and vascular disease in the Lothian Birth Cohort 1936: rationale, design and methodology of the imaging protocol. *International Journal of Stroke* **6**, 547–559.
- Westlye LT, Bjørnebekk A, Grydeland H, Fjell AM, Walhovd KB** (2011). Linking an anxiety-related personality trait to brain white matter microstructure: diffusion tensor imaging and harm avoidance. *Archives of General Psychiatry* **68**, 369–377.
- Zanetti MV, Jackowski MP, Versace A, Almeida JR, Hassel S, Duran FL, Busatto GF, Kupfer DJ, Phillips ML** (2009). State-dependent microstructural white matter changes in bipolar I depression. *European Archives of Psychiatry and Clinical Neuroscience* **259**, 316–328.
- Zigmond AS, Snaith RP** (1983). The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica* **67**, 361–370.