



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Specialty experience in geriatric medicine is associated with a small increase in knowledge of delirium

Citation for published version:

Jenkin, RPL, Musonda, P, Maclulich, AMJ, Myint, PK & Davis, DHJ 2014, 'Specialty experience in geriatric medicine is associated with a small increase in knowledge of delirium', *Age and Ageing*, vol. 43, no. 1, pp. 141-4. <https://doi.org/10.1093/ageing/aft159>

Digital Object Identifier (DOI):

[10.1093/ageing/aft159](https://doi.org/10.1093/ageing/aft159)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Age and Ageing

Publisher Rights Statement:

© The Author 2013. Published by Oxford University Press on behalf of the British Geriatrics Society.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



12. Kramer BJ. Differential predictors of strain and gain among husbands caring for wives with dementia. *Gerontologist* 1997; 37: 239–49.
13. Quinn C, Clare L, Woods RT. What predicts whether caregivers of people with dementia find meaning in their role? *Int J Geriatric Psychiatry* 2012; 27: 1195–202.
14. Cohen CA, Colantonio A, Vernich L. Positive aspects of caregiving: rounding out the caregiver experience. *Int J Geriatric Psychiatry* 2002; 17: 184–8.
15. National Center for Chronic Disease Prevention and Health Promotion. Healthy aging at a glance 2011: helping people to live long and productive lives and enjoy a good quality of life. In. Atlanta, GA; 2011.
16. Administration on Aging. A profile of older Americans: 2010. Department of Health and Human Services. Available at http://www.aoa.gov/aoaroot/aging_statistics/Profile/2010/docs/2010_profile.pdf.
17. Centers for Disease Control and Prevention/National Centers of Health Statistics. Depression among adults 20 years of age and over, by age and percent of poverty level: United States, 2005–2010. Health, United States, 2011. Data From the National Health and Nutrition Examination Survey. Available at <http://www.cdc.gov/nchs/data/has/2011/fig33.pdf>.
18. Federal Interagency Forum on Aging-Related Statistics. Older Americans 2010: Key Indicators of Well-Being. Washington, DC: U.S. Government Printing Office. Available at http://www.agingstats.gov/Main_Site/Data/2010_Documents/docs/OA_2010.pdf.

Received 29 April 2013; accepted in revised form 28 July 2013

Age and Ageing 2014; 43: 141–144

doi: 10.1093/ageing/aft159

Published electronically 16 October 2013

© The Author 2013. Published by Oxford University Press on behalf of the British Geriatrics Society.

This is an Open Access article distributed under the terms of the Creative Commons

Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits

unrestricted reuse, distribution, and reproduction in any medium, provided

the original work is properly cited.

Specialty experience in geriatric medicine is associated with a small increase in knowledge of delirium

RODRIC PETER LLEWELYN JENKIN¹, PATRICK MUSONDA², ALASDAIR M. J. MACLULLICH³, PHYO KYAW MYINT⁴, DANIEL H. J. DAVIS⁵

¹Department of Medicine for the Elderly, St Mary's Hospital, Praed Street, London W2 1NY, UK

²Norwich Medical School, University of East Anglia, Centre for Infectious Disease Research, Norwich UK

³Edinburgh Delirium Research Group, University of Edinburgh, Edinburgh, UK

⁴Academic Department of Medicine for the Elderly, University of East Anglia, Norwich, Norfolk, UK

⁵Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

Address correspondence to: R. P. L. Jenkin. Tel: 02077008428; Email: rodricjenkin@nhs.net; rodricjenkin@hotmail.com

Abstract

Background: delirium is underdiagnosed and undertreated. Understanding of delirium among doctors in medical and ICU settings has previously been shown to be low. We hypothesised that junior doctors who had gained experience in geriatrics, neurology or psychiatry may have an increased knowledge of delirium.

Methods: we used data from a large multi-centre study of junior doctors conducted between December 2006 and January 2007 which is, to date, the largest survey of understanding of delirium among junior doctors. The original survey used a questionnaire within which certain key items led to a correct or incorrect answer. Total correct answers were recorded giving a maximum total knowledge score of 17 for each participant. The relationship between total knowledge score achieved on the questionnaire and time since qualification; specialty experience in geriatric medicine, psychiatry and/or neurology and self-reported experience with the Confusion Assessment Method (independent variables) were modelled using linear regression.

Results: around half (53.2%; 399 of 750) of those surveyed stated that they had experience in geriatric medicine. In contrast only 4.1 and 8.0% of respondents had experience in psychiatry and neurology, respectively. Experience in geriatric medicine was significantly associated with a modest increase in correct answers (4.7 versus 4.3 points, $P = 0.020$). No other variables were significantly associated with better scores.

Conclusion: experience in geriatric medicine leads to a small improvement in understanding of delirium among junior doctors.

Keywords: *delirium, survey, geriatric, doctor, specialty, older people*

Introduction

Delirium is an acute neuropsychiatric syndrome, occurring in 11–42% of elderly in-patients [1]. The presence of delirium is associated with prolonged hospital stay, higher rates of institutionalisation and increased mortality [2]. Furthermore, delirium is associated with accelerated cognitive decline in those with pre-existing dementia and a significantly increased risk for the development of dementia in those with normal cognition [3, 4]. Despite its high prevalence and major clinical significance, delirium is consistently under-recognised [1, 5].

Previous studies have examined doctors' understanding of delirium in the medical inpatient and ICU settings [6–8]. These studies have shown that the understanding of delirium appears to be poor in comparison with other common medical conditions such as acute coronary syndrome or pneumonia. In addition, there is a consistent discrepancy between participants' perception of the importance of delirium and their understanding of delirium; while the respondents acknowledged delirium is highly prevalent and important, they lacked understanding and were unfamiliar with delirium assessment tools.

Whether the knowledge of delirium among junior doctors is directly related to duration of their medical training or specialties in which they have worked has not previously been examined. We hypothesised that attachments in geriatrics, neurology or psychiatry and total time working as a doctor may be associated with a greater knowledge and understanding of delirium.

Subjects and methods

We used data from a multicentre survey conducted in the UK between 1 December 2006 and 31 January 2007 which is, to date, the largest survey of doctors' understanding of delirium. The methods of the survey have been previously reported [7]. In brief, a questionnaire was designed to test knowledge of delirium prevalence, diagnostic criteria using DSM-IV, use of specific screening tools, association with adverse outcomes and pharmacological management. A convenience sample of junior doctors working in acute medical specialties and emergency medicine was taken at each participating trust and responses from 784 doctors were received in total. The 34 Acute trusts involved were recruited through

contacts in the British Geriatrics Society. Participants filled in the questionnaire at the point of approach and no site investigator reported any refusals.

In addition to answering the questionnaire, participants were asked to report whether they had postgraduate specialty experience in geriatric medicine, psychiatry and/or neurology and the number of months since they originally qualified as a doctor.

Statistical analyses

Within the questionnaire, certain key items led to a correct or incorrect answer, for each either a score of 1 (for a correct answer) or a score of 0 (for an incorrect answer) was given from which a total score out of 17 was derived. Typical knowledge questions included: 'Using DSM-IV, which of the following features are essential to a diagnosis of delirium?'; 'Patients with delirium most commonly display reduced motor activity and lethargy (true/false?)'. 'Assuming no-contraindications, what would you consider to be an appropriate starting dose of haloperidol in a 70-year-old with severe agitation in whom behavioural management has been unsuccessful?' Answers were given equal weighting in the score; missing answers were regarded as incorrect. The unadjusted relationships between specialty exposure and median knowledge scores were assessed using the Mann–Whitney U test. The relationship between total knowledge score (dependent variable) and time since qualification; specialty experience in geriatric medicine, psychiatry and/or neurology and self-reported experience with the Confusion Assessment Method (CAM) (independent variables) were modelled using linear regression, with random-effects allowed for recruitment centre to account for the clustered nature of the data. All analyses were conducted in Stata 12.1 (StataCorp, USA).

Results

Around half (53.2%; 399 of 750) of those surveyed stated that they had experience in geriatric medicine. In contrast only 4.1 and 8.0% of respondents had experience in psychiatry and neurology, respectively. Use of the CAM was reported by 8.6%.

Scores on the knowledge of delirium items followed a normal distribution. The median score was 4 out of a

maximum possible score of 17. Figure 1 shows the unadjusted relationships between specialty experience and knowledge score. Experience in geriatric medicine was significantly associated with a modest increase in correct answers (4.7 versus 4.3 points, $P = 0.020$). No other variables were significantly associated with better scores.

The linear regression models showed that self-reported experience in geriatric medicine was significantly associated with an extra 0.29 correct answers after adjusting for centre when compared with those without specialty experience ($P = 0.029$) (Table 1). In addition, months since qualification and experience in psychiatry were associated with slightly better scores before adjusting for centre. However, after accounting for clustering in the data, these variables were no longer significant. Post-estimation of testing of model residuals showed no violation of the model assumptions, with non-constant residual variance ($P = 0.21$).

Discussion

We confirm that the overall knowledge of delirium appeared low in junior doctors at the time the survey was conducted. Experience in geriatric medicine was associated with a higher level of knowledge; however, the difference was small. Experience in neurology, psychiatry, use of the CAM and total months since qualification were not significantly associated with an improved knowledge of delirium. The small proportion with experience in neurology and psychiatry may have resulted in lack of power to detect any differences in knowledge.

This is the first study to examine an association between experience in geriatric medicine and knowledge of delirium. The study benefits from its large sample size and that the participants were drawn from trusts across the UK.

There are a number of limitations. It was a convenience sample so there is a possibility of selection bias, although this




Figure 1. Average delirium scores by specialty experience. Bar charts showing differences in knowledge score, by specialty exposure and previous use of the Confusion Assessment Method.

Table 1. Summary values of key characteristics and estimates from linear mixed models, with and without adjustment for survey centre

Explanatory variable	Answering 'Yes', <i>n</i> (%)	Unadjusted, slope (95% CI)	<i>P</i> -value	Adjusted, slope (95% CI)	<i>P</i> -value
No. of months experience	N/A	0.006 (0.002, 0.010)	0.001	0.004 (−0.000, 0.008)	0.063
Experience in geriatrics	399/750 (53.2%)	0.306 (0.073, 0.539)	0.010	0.291 (0.030, 0.553)	0.029
Experience in psychiatry	29/711 (4.1%)	0.600 (−0.000, 1.194)	0.050	0.479 (−0.187, 1.145)	0.159
Experience in neurology	57/712 (8.0%)	0.066 (−0.369, 0.501)	0.766	−0.157 (−0.672, 0.358)	0.549
Previous use of CAM	64/748 (8.6%)	0.176 (−0.241, 0.593)	0.408	0.015 (−0.456, 0.487)	0.949

CAM, Confusion Assessment Method. The adjusted slopes account for clustering by centre. The slopes show the coefficients from the model output and can be interpreted as the additional number of correct answers given self-reported clinical experience: per extra month of experience since qualification; for previous experience in (geriatric medicine/psychiatry/neurology) compared with no experience in (geriatric medicine/psychiatry/neurology) and for person with previous experience of the Confusion Assessment Method (CAM) compared with those without previous experience of the CAM.

may be mitigated to an extent by the large sample size. Although 'experience in geriatrics' referred to experience since qualification rather than exposure as a student, we did not ask participants to clarify what their specific experience had entailed. It is possible that a subset of doctors who have worked in posts with an emphasis on acute geriatric care or worked in specific acute geriatric units may have had a greater increase in level of knowledge of delirium and our analysis would not have detected this. We also did not ask how long ago they worked in geriatric medicine or whether they are currently working in geriatric medicine. As such, it is possible that a placement in geriatric medicine provides a more significant increase in knowledge of delirium, but that this is not sustained.

It is possible that insufficient emphasis was placed on the diagnosis of delirium in medical practice in general at the time of our survey. In the UK, the 2009 curriculum for Core Medical Trainees (including revisions up to August 2012) required knowledge coming under the heading of 'Acute confusion/Delirium'. Although it specifically states that trainees should know the predisposing and precipitating factors for delirium, and the initial and subsequent investigations required, no mention is made of making the diagnosis of delirium or knowing the diagnostic criteria. This is in contrast to the preceding section in the same document where for chest pain, an appropriate emphasis on the diagnostic process is included ([http://www.jrcptb.org.uk/trainingandcert/Documents/2009%20CMT%20framework%20\(revised%20Aug%202012\).pdf](http://www.jrcptb.org.uk/trainingandcert/Documents/2009%20CMT%20framework%20(revised%20Aug%202012).pdf)).

The curriculum for more senior trainees specialising in geriatric medicine has 'diagnostic criteria for delirium' as part of the expected knowledge to be acquired (http://www.gmc-uk.org/geriatric_curriculum_2010.pdf_32486221.pdf). Although it is questionable how much the specialty curricula directly influence junior doctors' learning, it would seem appropriate to include the diagnostic criteria for delirium as a knowledge requirement for CMT doctors in the next iteration of the curriculum. This would parallel the expectation that a core medical trainee should know how to diagnose a heart attack, stroke or pneumonia.

The publication of the 2010 NICE guidelines for delirium may have provided a stimulus to greater understanding of delirium among trainees, giving a clear national reference for the identification, investigation and management of patients with delirium (<http://guidance.nice.org.uk/CG103/NICEGuidance/pdf/English>). A further significant policy shift initiative which may lead to an improvement in the recognition of delirium is the implementation of the dementia CQUIN (Commissioning for Quality and Innovation) target. This requires trusts to perform a cognitive assessment on all patients admitted as an emergency who are over the age of 75 with an aim to increase identification of patients with delirium and dementia. As the summary guidance for the dementia CQUIN points out, '[identifying] all patients with cognitive impairment will improve the detection of delirium' (<https://www.wp.dh.gov.uk/cno/files/2012/05/CQUINN-safety-thermometer-guidance.pdf>).

In summary, specialty experience in geriatric medicine provides a small improvement in knowledge of delirium, and

increased emphasis on training in this area may lead to this effect being strengthened. Repeating the original survey in light of the NICE guidelines and increasing profile of delirium would be of great interest.

Key points

- Delirium is underdiagnosed and undertreated.
 - Understanding of delirium among junior doctors is low.
 - Experience of Geriatric Medicine leads to a modest increase in understanding of delirium.
-

Acknowledgements

The authors would like to thank the members of the National Delirium Survey group, listed in the online appendix.

Conflicts of interest

None declared.

Funding

D.D. is funded by the Wellcome Trust as a Research Training Fellow. R.J., P.M., A.M. and P.K.M. have no funding acknowledgments to declare.

References

1. Siddiqi N, House AO, Holmes JD. Occurrence and outcome of delirium in medical in-patients: a systematic literature review. *Age Ageing* 2006; 35: 350–64.
2. Witlox J, Eurelings LS, de Jonghe JF, Kalisvaart KJ, Eikelenboom P, van Gool WA. Delirium in elderly patients and the risk of postdischarge mortality, institutionalization, and dementia: a meta-analysis. *JAMA* 2010; 304: 443–51.
3. Fong TG, Jones RN, Shi P *et al.* Delirium accelerates cognitive decline in Alzheimer disease. *Neurology* 2009; 72: 1570–5.
4. Davis DH, Muniz Terrera G, Keage H *et al.* Delirium is a strong risk factor for dementia in the oldest-old: a population-based cohort study. *Brain* 2012; 135(Pt 9): 2809–16.
5. Collins N, Blanchard MR, Tookman A, Sampson EL. Detection of delirium in the acute hospital. *Age Ageing* 2010; 39: 131–5.
6. Young RS, Hinami K, Arseven A, Jain B, Williams MV. Hospitalists' lack of knowledge of the Confusion Assessment Method: a barrier to systematic validated delirium surveillance. *Hosp Pract (Minneapolis)* 2012; 40: 56–63.
7. Davis D, MacLulich A. Understanding barriers to delirium care: a multicentre survey of knowledge and attitudes amongst UK junior doctors. *Age Ageing* 2009; 38: 559–63.
8. Ely EW, Stephens RK, Jackson JC *et al.* Current opinions regarding the importance, diagnosis, and management of delirium in the intensive care unit: a survey of 912 healthcare professionals. *Crit Care Med* 2004; 32: 106–12.

Received 29 April 2013; accepted in revised form 28 July 2013