



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Seven new loci associated with age-related macular degeneration

Citation for published version:

D Gene Consortium, Fritsche, LG, Chen, W, Schu, M, Yaspan, BL, Yu, Y, Thorleifsson, G, Zack, DJ, Arakawa, S, Cipriani, V, Ripke, S, Igo, RP, Buitendijk, GHS, Sim, X, Weeks, DE, Guymer, RH, Merriam, JE, Francis, PJ, Hannum, G, Agarwal, A, Armbrecht, AM, Audo, I, Aung, T, Barile, GR, Benchaboune, M, Bird, AC, Bishop, PN, Branham, KE, Brooks, M, Brucker, AJ, Cade, WH, Cain, MS, Campochiaro, PA, Chan, C-C, Cheng, C-Y, Chew, EY, Chin, KA, Chowers, I, Clayton, DG, Cojocaru, R, Conley, YP, Cornes, BK, Daly, MJ, Dhillon, B, Edwards, AO, Evangelou, E, Fagerness, J, Ferreyra, HA, Friedman, JS, Geirsdottir, A & Wright, AF 2013, 'Seven new loci associated with age-related macular degeneration', *Nature Genetics*, vol. 45, no. 4, pp. 433-439. <https://doi.org/10.1038/ng.2578>

Digital Object Identifier (DOI):

[10.1038/ng.2578](https://doi.org/10.1038/ng.2578)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Nature Genetics

Publisher Rights Statement:

Published in final edited form as:

Nat Genet. 2013 April; 45(4): 433–439e2.

Published online 2013 March 3. doi: 10.1038/ng.2578

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Published in final edited form as:

Nat Genet. 2013 April ; 45(4): 433–439e2. doi:10.1038/ng.2578.

Seven New Loci Associated with Age-Related Macular Degeneration

The AMD Gene Consortium

Abstract

Age-related macular degeneration (AMD) is a common cause of blindness in older individuals. To accelerate understanding of AMD biology and help design new therapies, we executed a collaborative genomewide association study, examining >17,100 advanced AMD cases and >60,000 controls of European and Asian ancestry. We identified 19 genomic loci associated with AMD with $p < 5 \times 10^{-8}$ and enriched for genes involved in regulation of complement activity, lipid metabolism, extracellular matrix remodeling and angiogenesis. Our results include 7 loci reaching $p < 5 \times 10^{-8}$ for the first time, near the genes *COL8A1/FILIP1L*, *IER3/DDR1*, *SLC16A8*, *TGFBR1*, *RAD51B*, *ADAMTS9/MIR548A2*, and *B3GALT1*. A genetic risk score combining SNPs from all loci displayed similar good ability to distinguish cases and controls in all samples examined. Our findings provide new directions for biological, genetic and therapeutic studies of AMD.

AMD is a highly heritable progressive neurodegenerative disease that leads to loss of central vision through death of photoreceptors^{1,2}. In developed countries, AMD is the leading cause of blindness in those >65 years³. Genes in the complement pathway^{4–11} and a region of chromosome 10^{12,13} have now been implicated as the major genetic contributors to disease. Association has also been demonstrated with several additional loci^{14–20}, each providing an entry-point into AMD biology and potential therapeutic targets.

To accelerate the pace of discovery in macular degeneration genetics, 18 research groups from across the world formed the AMD Gene Consortium in early 2010, with support from the National Eye Institute (Table 1, Supplementary Table 1, Supplementary Note). To extend the catalog of disease associated common variants, we first organized a meta-analysis of genomewide association scans (GWAS) – combining data for >7,600 cases with advanced disease (geographic atrophy, neovascularization, or both) and >50,000 controls. Each study was first subject to GWAS quality control filters (customized taking into account study specific features²¹ as detailed in Supplementary Table 2) and standardized to the HapMap reference panel and statistical genotype imputation^{22–25}. Results were combined through meta-analysis²⁶ and thirty-two variants representing loci with promising evidence of association were genotyped in an additional >9,500 cases and >8,200 controls (Supplementary Tables 1–3; summary meta-analysis results available online). Our overall analysis of the most promising variants thus included >17,100 cases and >60,000 controls.

Our meta-analysis evaluated evidence for association at 2,442,884 SNPs (Figure 1). Inspection of Q-Q plots (Supplementary Figure 1) and the genomic control value ($\lambda_{GC}=1.06$) suggest that unmodeled population stratification does not significantly impact our findings (Supplementary Table 4 for details). Joint analysis of discovery and follow-up

studies²⁷ resulted in 19 loci reaching $p < 5 \times 10^{-8}$ (Figure 1, Table 2, Supplementary Table 5). These 19 loci include all susceptibility loci previously reaching $p < 5 \times 10^{-8}$, except the 4q12 gene cluster for which association was reported in a Japanese population. In addition, the set includes seven loci reaching $p < 5 \times 10^{-8}$ for the first time.

We evaluated heterogeneity between studies using the I^2 statistic, which compares variability in effect size estimates between studies to chance expectations²⁸. We observed significant ($p < .05/19$) heterogeneity only for loci near *ARMS2* ($I^2 = 75.7\%$, $p < 1 \times 10^{-6}$) and near *CFH* ($I^2 = 85.4\%$, $p < 1 \times 10^{-6}$). Although these two loci were significantly associated in every sample examined, the magnitude of association varied more than expected. To explore sources of heterogeneity, we carried out a series of sub-analyses: we repeated the genomewide meta-analysis adding an age-adjustment, separating neovascular (NV) and geographic atrophy (GA) cases, in men and women, and in European- and Asian-ancestry samples separately (Figure 3, Supplementary Figure 2). These sub-analyses of the full GWAS dataset did not uncover additional loci reaching $p < 5 \times 10^{-8}$; furthermore heterogeneity near *CFH* and *ARMS2* remained significant in all sub-analyses ($I^2 > 65\%$, $p < .001$). Consistent with previous reports^{17,29,30}, separate analysis of NV and GA cases showed *ARMS2* risk alleles preferentially associated with risk of NV ($OR_{NV} = 2.97$, $OR_{GA} = 2.50$, $p_{\text{difference}} = .0008$) whereas *CFH* risk alleles preferentially associated with risk of GA ($OR_{NV} = 2.34$, $OR_{GA} = 2.80$, $p_{\text{difference}} = .0006$). We also observed large differences in effect sizes when stratifying by ethnicity, with variants near *CFH* exhibiting stronger evidence for association among Europeans ($p = .0000001$) and those near *TNFRSF10A* among East Asians ($p = .002$). Potential explanations include differences in linkage disequilibrium between populations or differences in environmental or diagnostic factors that modify genetic effects.

Identifying the full spectrum of allelic variation that contributes to disease in each locus will require sequencing of AMD cases and controls. To conduct an initial evaluation of the evidence for multiple AMD risk alleles in the nineteen loci described here, we repeated genomewide association analyses conditioning on the risk alleles listed in Table 2. We then examined each of the 19 implicated loci for variants with independent association (at $p < .0002$, corresponding for an estimate of ~250 independent variants per locus). This analysis resulted in the identification of the previously well documented independently associated variants near *CFH* and *C2/CFB*^{8,10,31,32} and of additional independent signals near *C3*, *CETP*, *LIPC*, *FRK/COL10A1*, *IER3/DDR1*, *RAD51B* (Supplementary Table 6). In four of these loci, the independently associated variants mapped very close (within <60kb) to the original signal. This shows each locus can harbor multiple susceptibility alleles, encouraging searches for rare variants that elucidate gene function in these regions^{33,34}.

To prioritize our search for likely causal variants, we examined each locus in detail (see LocusZoom³⁵ plots in Supplementary Figure 3) and investigated whether AMD risk alleles were associated with changes in protein sequence, copy number variation or insertion deletion polymorphisms. One quarter of associated variants altered protein sequence, either directly ($N = 2$) or through linkage disequilibrium ($r^2 > .6$; $N = 3$) with a nearby non-synonymous variant (Supplementary Table 7). Some coding variants point to well-studied genes (*ARMS2*, *C3* and *APOE*) while others help prioritize nearby genes for further study. In chromosome 4q25, index SNP rs4698775 is in strong linkage disequilibrium ($r^2 = .88$) with a potentially protein damaging variant in *CCDC109B*³⁶, a coiled coil domain containing protein that may be involved in the regulation of gene expression. In chromosome 6q22, index SNP rs3812111 is a perfect proxy for a coding variant in *COL10A1*, a collagen protein that could be important in maintaining the structure and function of the extra-cellular matrix. Interestingly, rs1061170 (NP_000177.2[CFH]:p.His402Tyr) was not in disequilibrium with rs10737680, the most strongly associated SNP in the *CFH* region but, instead, was tagged by a secondary and

weaker association signal (Supplementary Tables 6&7). This is consistent with prior haplotype analyses of the locus^{10,31,32,34,37}.

We used publicly available data^{38,39} to check whether any of our index SNPs might be proxies for copy number variants or insertion-deletion polymorphisms (indels), which are hard to directly interrogate with genotyping arrays. This analysis identified a single strong association ($r^2=.99$), between rs10490924, a coding variant in the *ARMS2* gene which is the peak of association in 10q26, and a 3' UTR indel polymorphism associated with *ARMS2* mRNA instability⁴⁰. Since index SNP rs10490924 is also in strong disequilibrium ($r^2=.90$) with a nearby SNP, rs11200638, that regulates *HTRA1*⁴¹, our data does not directly answer whether *HTRA1* or *ARMS2* is the causal gene in this locus. Although a common deletion of the *CFHR1* and *CFHR3* genes has been proposed^{42,43}, there was only modest signal in this study which is likely due to linkage disequilibrium with our most significantly associated variants in the locus ($r^2=.31$ between rs10737680 and 1000 Genomes Project MERGED_DEL_2_6731) as previously suggested³⁴.

Using RNA-sequencing⁴⁴, we examined mRNA levels of 85 genes within 100 kb of our index SNPs in post-mortem human retina (Supplementary Table 8). Of 19 independent risk loci, three had no genes with expressed transcripts in either old or young retina. Two genes showed differential expression between post-mortem retina of young (ages 17–35) and elderly (ages 75 and 77) individuals: *CFH* ($p=.009$) and *VEGFA* ($p=.003$), both with increased expression in older individuals. Using previously published data⁴⁵, we also examined the expression of associated genes in fetal and adult retinal pigment epithelium (RPE). This revealed increased *C3* expression in adult RPE compared to fetal RPE ($p=.0008$). *CFH*, *VEGFA* and *C3* are thus up-regulated with aging, and their role in AMD may indicate an accelerated aging process. In addition to *C3* and *CFH*, all the complement genes with detectable expression in the retina or RPE experiments showed higher expression levels in older tissue.

To identify biological relationships among our genetic association signals, we catalogued the genes within 100kb of the variants in each association peak ($r^2>0.8$ with the index SNP listed in Table 1). Ingenuity Pathway Analysis (Ingenuity Systems, Redwood, CA) highlighted several biological pathways, particularly the complement system and atherosclerotic signaling, to be enriched in the resulting set of 90 genes (Table 3, Supplementary Table 9). To account for features of genomewide association studies (such as the different number of SNPs in each gene), we carried out two additional analyses. First, we repeated our analysis for 50 sets of 19 control loci drawn from the NHGRI GWAS catalog⁴⁶. In these 50 control sets, Ingenuity enrichment p-values for the complement system and for atherosclerosis signaling genes were exceeded 16% and 32% of the time respectively (although these two specific pathways were never implicated in a control dataset). We also repeated our enrichment analyses using the INRICH tool⁴⁷, which is specifically designed for the analysis of genomewide association studies – but accesses a more limited set of annotations. The INRICH analyses showed enrichment for genes encoding collagen and extra-cellular region proteins (both with $p=10^{-5}$ and after adjustment for multiple testing $p_{\text{adjust}}=.0006$), complement and coagulation cascades ($p=.0002$, $p_{\text{adjust}}=.03$), lipoprotein metabolism ($p=.0003$, $p_{\text{adjust}}=.04$), and regulation of apoptosis ($p=.0009$, $p_{\text{adjust}}=.09$) (Supplementary Table 10).

To explore the connections between our genetic association signals, we tested for interaction between pairs of risk alleles – looking for situations where joint risk was different than expected based on marginal effects. This analysis resulted in 171 tests of interaction, of which 9 were nominally significant ($p<.05$, see Supplementary Table 11), consistent with chance expectations. The strongest observed interaction involved risk alleles at rs10737680

(near *CFH*) and rs429608 (near *C2/CFB*), the only association that remained significant after adjusting for multiple testing ($p=.000052$, $<0.05/171=.00029$). Individuals carrying risk alleles at both these loci were at slightly higher risk of disease than expected.

The proportion of the variability in the risk of AMD that is due to genes, or heritability, has been estimated at 45–70%². Estimating the proportion of disease risk explained by the susceptibility loci identified⁴⁸ depends greatly on the disease prevalence - which is difficult to estimate in our sample, as it includes cases and controls of different ages and collected through a variety of ascertainment schemes. Using a model that assumes an underlying normally distributed but unobserved disease risk score or liability⁴⁹, the nineteen loci described here account for between 10% (if AMD prevalence is close to 1%) and 30% (if AMD prevalence is closer to 10%) of the variability in disease risk (corresponding to 15–65% of the total genetic contribution to AMD). The variants representing the peak of association at loci previously reaching genomewide significance account for the bulk of this variability: the new loci identified here account for 0.5–1.0% of the total heritability of AMD whereas secondary signals at novel and known loci account for 1.5–3.0% of the total heritability.

We report here the most comprehensive genetic association study of macular degeneration yet conducted, involving 18 international research groups, and a large set of cases and controls. Our data reveal 19 susceptibility loci, including 7 loci reaching $p<5\times 10^{-8}$ for the first time, nearly doubling the number of known AMD loci outside the complement pathway. Our results show some susceptibility alleles exhibit different association across ethnic groups and may be preferentially associated with specific subtypes of disease. As with other GWAS meta-analysis, differences in genotyping methods, quality control steps and imputation strategies between samples might have a minor effect in our results – future studies may document that more uniform approaches across larger sample sizes might uncover more signals. A conundrum of macular degeneration genetics remains that the loci identified to date contribute to both GA and NV, two different phenotypes of advanced disease. In our sample, subtype specific GWAS analyses considering GA or NV cases only did not identify additional loci. Consistent with observations for other complex diseases³⁹, the majority of common disease susceptibility alleles do not alter protein sequences and are not associated with insertions or deletions of coding sequence or with copy number variation. We expect that the loci identified here will provide an ideal starting point for studies of rare variation^{33,34}.

In contrast to most other complex diseases, a risk score combining information across our 19 loci, can distinguish cases and controls relatively well (Figure 4, area under the ROC curve [AUC]=.52 including only new loci or AUC=.74 including new and previously reported loci; Supplementary Figure 4). It may be possible to use similar scores to identify and prioritize at risk individuals so they receive preventative treatment prior to the onset of disease⁵⁰. Monotherapies are increasingly utilized to manage neovascular disease, but offer only a limited repertoire of treatment options to patients. Identification of novel genes and pathways enables us to pursue a larger range of disease-specific targets for development of new therapeutic interventions. We expect that future therapies directed at earlier stages of the disease process will allow patients to retain visual function for longer periods, improving the quality of life for individuals with AMD.

ONLINE METHODS

GENOME-WIDE SCAN FOR LATE AMD ASSOCIATION INCLUDING FOLLOW-UP

Study-specific association analysis for discovery—Genotyping was performed on a variety of different platforms summarized in Supplementary Table 2. Each group

submitted results from association tests using genotyped and imputed data where the allelic dosages were computed with either MACH²⁵, IMPUTE²³, BEAGLE²⁴, or snpStats⁵² using the HapMap2 reference panels. The CEU panel was used as a reference for imputation-based analyses for most samples (predominantly of European ancestry), with two exceptions: for the JAREDS samples (predominantly of East Asian ancestry), the CHB+JPT panel was used as a reference; for the VRF samples (predominantly of South Asian ancestry) the combined CEU and CHB+JPT panels were used^{22,53}. For most data sets association tests were run under a logistic regression model using either Plink⁵⁴, Mach2dat²⁵, ProbABEL⁵⁵, or snpStats⁵², though for one dataset containing related individuals the generalized estimating equations algorithm⁵⁶ as implemented in R^{57,58}. In addition to the primary analysis which tested for SNP associations with advanced AMD unadjusted for age, an age-adjusted sensitivity analysis was conducted by each group with available age. Each group also provided stratified results by sex or AMD subtype (GA or NV) as long as the sample size per stratum exceeded 50 subjects. For all analyses, study-specific control for population stratification was conducted (Supplementary Table 4).

Study-specific association analysis for follow-up—Genotyping of the selected SNPs was performed on different platforms; the same models, sensitivity and stratified analyses were computed by each follow-up partner, while SNPs with insufficient call rate were excluded based on study-specific thresholds. If the index SNP could not be genotyped, a highly correlated proxy was used whenever possible (Supplementary Tables 2&3).

Quality control before meta-analysis—Before meta-analysis, all study-specific files underwent quality control procedures to check for completeness and plausible descriptive statistics on all variables as well as for compliance of allele frequencies with HapMap⁵⁹. In addition, we excluded SNP results of a study into meta-analysis (i) for discovery: if imputation quality measures were too low (MACH & PLINK <0.3; SNPTEST <0.4) or if effect sizes ($|\beta|$) or standard errors were too extreme (> 5) indicating instability of the estimates, (ii) for follow-up: if Hardy-Weinberg equilibrium was violated ($p < 0.05/32$).

Meta-analyses—For both discovery and follow-up, we performed meta-analyses using the inverse variance weighted fixed effect model, which pools the effect size and standard error of each participated GWAS. Using an alternative weighted z-score method, which is based on a weighted sum of z-score statistics, we obtained a very similar set of test statistics (correlation of $-\log_{10}(p\text{-value}) > 0.98$). All analyses were performed using METAL²⁶ and R. For the discovery, we applied two rounds of genomic control corrections to each individual GWAS and the combined meta results, respectively⁵¹. All results were analyzed and validated among four independent teams.

EXTENDED ANALYSES FOR THE IDENTIFIED AMD LOCI

Extended analyses were conducted on the identified loci and particularly on the top SNP of each locus.

Second signal analysis—To detect potential independent signals within the identified AMD loci, each study partner with genotypes for all identified SNPs available re-analyzed their data for all SNPs in the respective loci (index SNP $\pm 1\text{Mb}$) using a logistic regression model containing all identified index SNPs. Quality control procedures were performed as before. The beta estimates for each SNP were meta-analyzed applying the effective sample size weighted z-score method and two rounds of genomic control correction. The significance threshold ($p < 0.05$) for an independent association signal within any of the identified loci was Bonferroni-adjusted using the average effective number of SNPs

involved across the identified loci determined by SNPSpD⁶⁰. To this analysis, 13 studies contributed including 7,489 cases and 51,562 controls.

Interaction analysis—Utilizing a pre-specified R-scripts (see URLs), GWAS partners performed 171 logistic regression analyses modeling the pair-wise interaction of the 19 index SNPs assuming an additive model for main and interaction effects. Study-specific covariates were included to the model if required. Per study, quality control included a check for consistency of SNP main effects between discovery and interaction analysis. SNPs with low imputation quality measures and pairs with $|\beta| > 5$ or standard errors > 5 were excluded before meta-analyzing the interaction effects with the inverse variance weighted fixed effect model in METAL. To this analysis, 12 studies contributed including 6,645 cases and 49,410 controls.

GENETIC RISK SCORE

The meta-analyzed effect sizes, β_j , for each of the 19 SNPs were calculated in the meta-

analysis described above and normalized: $\hat{\beta}_j = \beta_j / \sum_{k=1}^{19} \beta_k$, $j=1, \dots, 19$. Using these as weights, each study partner with all 19 SNPs available computed the individuals' genetic risk score as a normalized weighted sum of the AMD risk increasing alleles among the identified SNPs as

$$S_i = \sum_j \hat{\beta}_j x_{ij},$$

where x_{ij} is the genotype of the i th individual at the j th SNP, so S_i ranges from 0 to 2. This data was available from 12 studies including 7,195 cases and 49,149 controls.

For each study, we used a leave-one-out cross-validation to assess the prediction of the risk score. For the k th subject, we fitted a logistic regression model from all subjects in the study

excluding the k th subjects: $\log\left(\frac{y_i}{1-y_i}\right) = \alpha + \gamma S_i$, $i \neq k$, α is the intercept and γ is the effect of the genetic risk score. The fitted probability of the k th subject was then estimated as $\hat{y}_k = 1 / (1 + e^{-(\alpha + \gamma S_k)})$. We sorted the fitted probabilities and calculated sensitivity and specificity by varying the risk threshold (the value compared with the fitted probability to dichotomize the subject into case or control) from 0 to 1. These were utilized to compute the area-under-the curve (AUC) of the receiver-operating-curve (ROC).

IDENTIFICATION OF CORRELATED CODING VARIANTS AND TAGGED NON-SNP VARIATION

LD estimates were calculated using genotype data of the identified risk loci (index SNPs ± 500 kb) of individuals with European ancestry from the 1000 Genomes Project (March 2012 release)⁶¹ or from HapMap (release #28)⁵⁹. Variants correlated ($r^2 > 0.6$) with one of the GWAS index SNPs were identified using PLINK⁵⁴. To filter coding variants, all correlated variants were mapped against RefSeq transcripts using ANNOVAR⁶².

GENE EXPRESSION

We evaluated expression of genes within 100kb of one of the 19 index SNPs, as well as of several retina-specific, RPE-specific and housekeeping genes unrelated to AMD for comparison in retina (RNA-sequencing data from three young [17–35 yrs age] and two old

individuals [75 and 77 yrs age]) as well as in fetal and adult retinal pigment epithelium (RPE; published data in the Gene Expression Omnibus database⁴⁵). Expression was analyzed using previously described protocols⁴⁴ (Supplementary Table 8).

PATHWAY ANALYSES

Functional enrichment analysis was performed using the Ingenuity Pathway Analysis software (IPA, Ingenuity® Systems). Any gene located within 100kb of a SNP in high LD ($r^2 > 0.8$) with one of the index SNPs was considered a potential AMD risk associated gene and considered for subsequent pathway enrichment analysis. LD estimates were calculated as described above. Applying the above inclusion filters, 90 genes appear to be implicated by our 19 replicated AMD SNPs (Supplementary Table 8). Because genes with related function sometimes cluster in the same locus, we trimmed gene lists during analysis so that only one gene per locus was used to evaluate enrichment for each pathway. The P-value of the association between our implicated gene list and any of the canonical pathways and/or functional gene sets as annotated by IPA's Knowledge Base was computed using a one-sided Fisher's exact test. The Benjamini-Hochberg method was used to estimate False Discovery Rates. To evaluate significance of observed enrichment, we repeated our Ingenuity analysis starting with 50 lists of 19 SNPs randomly drawn from the NHGRI GWAS catalog⁴⁶ and, again, using the INRICH tool⁶³. When using INRICH, we used gene sets defined in the Broad's Molecular Signatures database⁴⁷ (ver3.0) representing manually curated canonical pathway, Gene Ontology biological process, cellular component and molecular function gene sets (C2:CP, C5:BP, C5:CC and C5:MF). We provided INRICH with our full GWAS SNP list and allowed it to carry out 100,000 permutations, matching selected loci in terms of gene count, SNP density and total number of SNPs.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are indebted to all the participants who volunteered their time, DNA and information to make this research study possible. We are also in great debt to the clinicians, nurses and research staff who participated in patient recruitment and phenotyping. We thank Dr. Hemin Chin for constant support and encouragement, which helped us bring this project to completion. We thank Drs. Sheldon Miller and Jennifer Barb for access to RPE expression data and the MIGEN study group for use of their genotype data. We thank C. Pappas, N. Miller, J. Hageman, W. Hubbard, L. Lucci, A. Vitale, P. Bernstein and N. Amin for technical and clinical assistance. We thank E. Roitchchina, A.C. Viswanathan, J. Xie, M. Inouye, E.G. Holliday, J. Attia, R.J. Scott for contributions to the Blue Mountains Eye Study GWAS. We thank members of the Genetic Factors in AMD Study Group, the Scottish Macula Society Study Group, the Wellcome Trust Clinical Research facility at Southampton General Hospital. We thank Tunde Peto and colleagues at the Reading Centre, Moorfields Eye Hospital, London, and C. Brussee and A. Hooghart for help in patient recruitment and phenotyping. Full details of funding sources can be found in the Supplementary Information.

References

1. Swaroop A, Chew EY, Rickman CB, Abecasis GR. Unravelling a Late-Onset Multifactorial Disease: From Genetic Susceptibility to Disease Mechanisms for Age-related Macular Degeneration. *Annu Rev Genomics Hum Genet.* 2009; 10:19–43. [PubMed: 19405847]
2. Seddon JM, Cote J, Page WF, Aggen SH, Neale MC. The US twin study of age-related macular degeneration: relative roles of genetic and environmental influences. *Archives of Ophthalmology.* 2005; 123:321–7. [PubMed: 15767473]
3. Friedman DS, et al. Prevalence of age-related macular degeneration in the United States. *Archives of Ophthalmology.* 2004; 122:564–72. [PubMed: 15078675]

4. Edwards AO, et al. Complement factor H polymorphism and age-related macular degeneration. *Science*. 2005; 308:421–4. [PubMed: 15761121]
5. Haines JL, et al. Complement factor H variant increases the risk of age-related macular degeneration. *Science*. 2005; 308:419–21. [PubMed: 15761120]
6. Klein RJ, et al. Complement factor H polymorphism in age-related macular degeneration. *Science*. 2005; 308:385–9. [PubMed: 15761122]
7. Yates JR, et al. Complement C3 variant and the risk of age-related macular degeneration. *N Engl J Med*. 2007; 357:553–61. [PubMed: 17634448]
8. Gold B, et al. Variation in factor B (BF) and complement component 2 (C2) genes is associated with age-related macular degeneration. *Nat Genet*. 2006; 38:458–62. [PubMed: 16518403]
9. Fagerness JA, et al. Variation near complement factor I is associated with risk of advanced AMD. *Eur J Hum Genet*. 2009; 17:100–4. [PubMed: 18685559]
10. Hageman GS, et al. A common haplotype in the complement regulatory gene factor H (HF1/CFH) predisposes individuals to age-related macular degeneration. *Proc Natl Acad Sci U S A*. 2005; 102:7227–32. [PubMed: 15870199]
11. Maller JB, et al. Variation in complement factor 3 is associated with risk of age-related macular degeneration. *Nat Genet*. 2007; 39:1200–1. [PubMed: 17767156]
12. Rivera A, et al. Hypothetical LOC387715 is a second major susceptibility gene for age-related macular degeneration, contributing independently of complement factor H to disease risk. *Hum Mol Genet*. 2005; 14:3227–36. [PubMed: 16174643]
13. Jakobsdottir J, et al. Susceptibility genes for age-related maculopathy on chromosome 10q26. *Am J Hum Genet*. 2005; 77:389–407. [PubMed: 16080115]
14. Klaver CC, et al. Genetic association of apolipoprotein E with age-related macular degeneration. *American Journal of Human Genetics*. 1998; 63:200–6. [PubMed: 9634502]
15. Souied EH, et al. The epsilon4 allele of the apolipoprotein E gene as a potential protective factor for exudative age-related macular degeneration. *American Journal of Ophthalmology*. 1998; 125:353–9. [PubMed: 9512153]
16. McKay GJ, et al. Evidence of association of APOE with age-related macular degeneration: a pooled analysis of 15 studies. *Human Mutation*. 2011; 32:1407–16. [PubMed: 21882290]
17. Chen W, et al. Genetic variants near TIMP3 and high-density lipoprotein-associated loci influence susceptibility to age-related macular degeneration. *Proceedings of the National Academy of Sciences of the United States of America*. 2010; 107:7401–6. [PubMed: 20385819]
18. Neale BM, et al. Genome-wide association study of advanced age-related macular degeneration identifies a role of the hepatic lipase gene (LIPC). *Proceedings of the National Academy of Sciences of the United States of America*. 2010; 107:7395–400. [PubMed: 20385826]
19. Yu Y, et al. Common variants near FRK/COL10A1 and VEGFA are associated with advanced age-related macular degeneration. *Human Molecular Genetics*. 2011; 20:3699–709. [PubMed: 21665990]
20. Arakawa S, et al. Genome-wide association study identifies two susceptibility loci for exudative age-related macular degeneration in the Japanese population. *Nature Genetics*. 2011; 43:1001–4. [PubMed: 21909106]
21. McCarthy MI, et al. Genome-wide association studies for complex traits: consensus, uncertainty and challenges. *Nat Rev Genet*. 2008; 9:356–69. [PubMed: 18398418]
22. Li Y, Willer CJ, Sanna S, Abecasis GR. Genotype Imputation. *Annu Rev Genomics Hum Genet*. 2009; 10:387–406. [PubMed: 19715440]
23. Marchini J, Howie B, Myers S, McVean G, Donnelly P. A new multipoint method for genome-wide association studies by imputation of genotypes. *Nature Genetics*. 2007; 39:906–13. [PubMed: 17572673]
24. Browning BL, Browning SR. A unified approach to genotype imputation and haplotype-phase inference for large data sets of trios and unrelated individuals. *American Journal of Human Genetics*. 2009; 84:210–23. [PubMed: 19200528]
25. Li Y, Willer CJ, Ding J, Scheet P, Abecasis GR. MaCH: using sequence and genotype data to estimate haplotypes and unobserved genotypes. *Genetic Epidemiology*. 2010; 34:816–34. [PubMed: 21058334]

26. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. 2010; 26:2190–1. [PubMed: 20616382]
27. Skol AD, Scott LJ, Abecasis GR, Boehnke M. Joint analysis is more efficient than replication-based analysis for two-stage genome-wide association studies. *Nat Genet*. 2006; 38:209–13. [PubMed: 16415888]
28. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003; 327:557–60. [PubMed: 12958120]
29. Sobrin L, et al. ARMS2/HTRA1 locus can confer differential susceptibility to the advanced subtypes of age-related macular degeneration. *American Journal of Ophthalmology*. 2011; 151:345–52. e3. [PubMed: 21122828]
30. Seddon JM, et al. Association of CFH Y402H and LOC387715 A69S with progression of age-related macular degeneration. *JAMA*. 2007; 297:1793–800. [PubMed: 17456821]
31. Li M, et al. CFH haplotypes without the Y402H coding variant show strong association with susceptibility to age-related macular degeneration. *Nature Genetics*. 2006; 38:1049–54. [PubMed: 16936733]
32. Maller J, et al. Common variation in three genes, including a noncoding variant in CFH, strongly influences risk of age-related macular degeneration. *Nature Genetics*. 2006; 38:1055–9. [PubMed: 16936732]
33. Nejentsev S, Walker N, Riches D, Egholm M, Todd JA. Rare variants of IFIH1, a gene implicated in antiviral responses, protect against type 1 diabetes. *Science*. 2009; 324:387–9. [PubMed: 19264985]
34. Raychaudhuri S, et al. A rare penetrant mutation in CFH confers high risk of age-related macular degeneration. *Nature Genetics*. 2011; 43:1232–6. [PubMed: 22019782]
35. Pruim RJ, et al. LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics*. 2010; 26:2336–7. [PubMed: 20634204]
36. Adzhubei IA, et al. A method and server for predicting damaging missense mutations. *Nat Methods*. 2010; 7:248–9. [PubMed: 20354512]
37. Sivakumaran TA, et al. A 32 kb Critical Region Excluding Y402H in CFH Mediates Risk for Age-Related Macular Degeneration. *PLoS ONE*. 2011; 6:e25598. [PubMed: 22022419]
38. The Wellcome Trust Case Control Consortium. Genome-wide association study of CNVs in 16, 000 cases of eight common diseases and 3, 000 shared controls. *Nature*. 2010; 464:713–20. [PubMed: 20360734]
39. The 1000 Genomes Project. A map of human genome variation from population scale sequencing. *Nature*. 2010; 467
40. Fritsche LG, et al. Age-related macular degeneration is associated with an unstable ARMS2 (LOC387715) mRNA. *Nat Genet*. 2008; 40:892–6. [PubMed: 18511946]
41. Dewan A, et al. HTRA1 promoter polymorphism in wet age-related macular degeneration. *Science*. 2006; 314:989–92. [PubMed: 17053108]
42. Hughes AE, et al. A common CFH haplotype, with deletion of CFHR1 and CFHR3, is associated with lower risk of age-related macular degeneration. *Nat Genet*. 2006; 38:1173–7. [PubMed: 16998489]
43. Fritsche LG, et al. An imbalance of human complement regulatory proteins CFHR1, CFHR3 and factor H influences risk for age-related macular degeneration (AMD). *Human Molecular Genetics*. 2010; 19:4694–704. [PubMed: 20843825]
44. Brooks MJ, Rajasimha HK, Roger JE, Swaroop A. Next-generation sequencing facilitates quantitative analysis of wild-type and Nrl(–/–) retinal transcriptomes. *Molecular Vision*. 2011; 17:3034–54. [PubMed: 22162623]
45. Strunnikova NV, et al. Transcriptome analysis and molecular signature of human retinal pigment epithelium. *Human Molecular Genetics*. 2010; 19:2468–86. [PubMed: 20360305]
46. Hindorf LA, et al. Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. *Proceedings of the National Academy of Sciences of the United States of America*. 2009; 106:9362–7. [PubMed: 19474294]

47. Subramanian A, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences of the United States of America*. 2005; 102:15545–50. [PubMed: 16199517]
48. Manolio TA, et al. Finding the missing heritability of complex diseases. *Nature*. 2009; 461:747–53. [PubMed: 19812666]
49. So HC, Gui AH, Cherny SS, Sham PC. Evaluating the heritability explained by known susceptibility variants: a survey of ten complex diseases. *Genetic Epidemiology*. 2011; 35:310–7. [PubMed: 21374718]
50. Seddon JM, Reynolds R, Yu Y, Daly MJ, Rosner B. Risk models for progression to advanced age-related macular degeneration using demographic, environmental, genetic, and ocular factors. *Ophthalmology*. 2011; 118:2203–11. [PubMed: 21959373]
51. Devlin B, Roeder K. Genomic control for association studies. *Biometrics*. 1999; 55:997–1004. [PubMed: 11315092]
52. Wallace C, et al. The imprinted DLK1-MEG3 gene region on chromosome 14q32.2 alters susceptibility to type 1 diabetes. *Nature Genetics*. 2010; 42:68–71. [PubMed: 19966805]
53. Huang L, et al. Genotype-imputation accuracy across worldwide human populations. *American Journal of Human Genetics*. 2009; 84:235–50. [PubMed: 19215730]
54. Purcell S, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*. 2007; 81:559–75. [PubMed: 17701901]
55. Aulchenko YS, Struchalin MV, van Duijn CM. ProbABEL package for genome-wide association analysis of imputed data. *BMC Bioinformatics*. 2010; 11:134. [PubMed: 20233392]
56. Zeger SL, Liang KY. Longitudinal data analysis for discrete and continuous outcomes. *Biometrics*. 1986; 42:121–30. [PubMed: 3719049]
57. Carey, VJ. R package version 4.13–18. 2012. Ported to R by Thomas Lumley and Brian Ripley. Note that maintainers are not available to give advice on using a package they did not author. gee: Generalized Estimation Equation solver.
58. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing; Vienna, Austria: 2012.
59. The International HapMap Consortium et al. Integrating common and rare genetic variation in diverse human populations. *Nature*. 2010; 467:52–8. [PubMed: 20811451]
60. Nyholt DR. A simple correction for multiple testing for single-nucleotide polymorphisms in linkage disequilibrium with each other. *Am J Hum Genet*. 2004; 74:765–9. [PubMed: 14997420]
61. The 1000 Genomes Project Consortium et al. An integrated map of genetic variation from 1,092 human genomes. *Nature*. 2012; 491:56–65. [PubMed: 23128226]
62. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res*. 2010; 38:e164. [PubMed: 20601685]
63. Lee PH, O'Dushlaine C, Thomas B, Purcell SM. INRICH: interval-based enrichment analysis for genome-wide association studies. *Bioinformatics*. 2012; 28:1797–9. [PubMed: 22513993]

Appendix

AUTHOR LIST

Lars G Fritsche [^{1,2,*}], Wei Chen [^{2,3,*}], Matthew Schu [^{4,*}], Brian L Yaspan [^{5,6,*}], Yi Yu [^{7,*}], Gudmar Thorleifsson [⁸], Donald J Zack [^{9,10,11,12}], Satoshi Arakawa [¹³],

¹Institute of Human Genetics, University of Regensburg, Regensburg, Germany.

²Department of Biostatistics, Center for Statistical Genetics, University of Michigan, Ann Arbor, Michigan, USA.

*These authors contributed equally to this work.

³Division of Pediatric Pulmonary Medicine, Allergy and Immunology, Department of Pediatrics, Children's Hospital of Pittsburgh of UPMC, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA.

⁴Department of Medicine (Section of Biomedical Genetics), Boston University Schools of Medicine and Public Health, Boston, Massachusetts, USA.

⁵Center for Human Genetics Research, Vanderbilt University Medical Center, Nashville, Tennessee, USA.

⁶Department of Molecular Physiology and Biophysics, Vanderbilt University School of Medicine, Nashville, Tennessee, USA.

Valentina Cipriani [¹⁴], Stephan Ripke [^{15,16}], Robert P Igo, Jr. [^{17,18}], Gabriëlle H S Buitendijk [^{19,20}], Xueling Sim [^{2,21}], Daniel E Weeks [^{22,23}], Robyn H Guymmer [²⁴], Joanna E Merriam [²⁵], Peter J Francis [²⁶], Gregory Hannum [²⁷], Anita Agarwal [^{28,29}], Ana Maria Armbrrecht [³⁰], Isabelle Audo [^{10,31,32,33}], Tin Aung [^{34,35}], Gaetano R Barile [²⁵], Mustapha Benchaboune [³⁶], Alan C Bird [¹⁴], Paul N Bishop [^{37,38}], Kari E Branham [³⁹], Matthew Brooks [⁴⁰], Alexander J Brucker [⁴¹], William H Cade [^{42,43}], Melinda S Cain [²⁴], Peter A Campochiaro [^{11,44}], Chi-Chao Chan [⁴⁵], Ching-Yu Cheng [^{34,35,46,47}], Emily Y Chew [⁴⁸], Kimberly A Chin [⁷], Itay Chowers [⁴⁹], David G Clayton [⁵⁰], Radu Cojocaru [⁴⁰], Yvette P Conley [⁵¹], Belinda K Cornes [³⁴], Mark J Daly [¹⁵], Baljean Dhillon [³⁰], Albert O Edwards [⁵²], Evangelos Evangelou [⁵³], Jesen

⁷Ophthalmic Epidemiology and Genetics Service, Tufts Medical Center, Boston, Massachusetts, USA.

⁸deCODE genetics, Reykjavik, Iceland.

⁹Department of Molecular Biology and Genetics, Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA.

¹⁰Department of Genetics, Institut de la Vision, UPMC Univ Paris 06, UMR_S 968, Paris, France.

¹¹Department of Neuroscience, Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA.

¹²Institute of Genetic Medicine, Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA.

¹³Laboratory for Genotyping Development, Research Group for Genotyping, Center for Genomic Medicine (CGM), RIKEN, Yokohama, Japan.

¹⁴Moorfields Eye Hospital and Institute of Ophthalmology, University College London, London, UK.

¹⁵Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, Massachusetts, USA.

¹⁶Stanley Center for Psychiatric Research, Broad Institute of Harvard and Massachusetts Institute of Technology, Cambridge, Massachusetts, USA.

¹⁷Department of Epidemiology, Case Western Reserve University, Cleveland, Ohio, USA.

¹⁸Department of Biostatistics, Case Western Reserve University, Cleveland, Ohio, USA.

¹⁹Department of Epidemiology, Erasmus Medical Center, Rotterdam, the Netherlands.

²⁰Department of Ophthalmology, Erasmus Medical Center, Rotterdam, the Netherlands.

²¹Centre for Molecular Epidemiology, National University of Singapore, Singapore, Singapore.

²²Department of Human Genetics, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.

²³Department of Biostatistics, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.

²⁴Centre for Eye Research Australia, University of Melbourne, Royal Victorian Eye and Ear Hospital, East Melbourne, Australia.

²⁵Department of Ophthalmology, Columbia University, New York, New York, USA.

²⁶Macular Degeneration Center, Casey Eye Institute, Oregon Health & Science University, Portland, Oregon, USA.

²⁷Department of Bioengineering, University of California San Diego, La Jolla, California, USA.

²⁸Vanderbilt Eye Institute, Vanderbilt University Medical Center, Nashville, Tennessee, USA.

²⁹Department of Ophthalmology & Visual Sciences, Vanderbilt University School of Medicine, Nashville, Tennessee, USA.

³⁰Department of Ophthalmology, University of Edinburgh and Princess Alexandra Eye Pavilion, Edinburgh, UK.

³¹INSERM, U968, Paris, France.

³²CNRS, UMR_7210, Paris, France.

³³Institute of Ophthalmology, University College London, London, UK.

³⁴Singapore Eye Research Institute, Singapore National Eye Centre, Singapore, Singapore.

³⁵Department of Ophthalmology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore.

³⁶Centre Hospitalier National d'Ophthalmologie des Quinze-Vingts, INSERM-DHOS CIC 503, Paris, France.

³⁷School of Biomedicine, Faculty of Medical and Human Sciences, University of Manchester, Manchester, UK.

³⁸Central Manchester University Hospitals NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, UK.

³⁹Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, Michigan, USA.

⁴⁰Neurobiology Neurodegeneration & Repair Laboratory (N-NRL), National Eye Institute, National Institutes of Health, Bethesda, Maryland, USA.

⁴¹Penn Presbyterian Medical Center, Scheie Eye Institute, Philadelphia, Pennsylvania, USA.

⁴²John P. Hussman Institute for Human Genomics, University of Miami Miller School of Medicine, Miami, Florida, USA.

⁴³Dr. John T. Macdonald Foundation Department of Human Genetics, University of Miami Miller School of Medicine, Miami, Florida, USA.

⁴⁴Department of Ophthalmology, Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA.

⁴⁵Immunopathology Section, Laboratory of Immunology, National Eye Institute, National Institutes of Health, Bethesda, Maryland, USA.

⁴⁶Saw Swee Hock School of Public Health, National University of Singapore, Singapore, Singapore.

⁴⁷Centre for Quantitative Medicine, Office of Clinical Sciences, Duke-NUS Graduate Medical School, Singapore, Singapore.

⁴⁸Division of Epidemiology and Clinical Applications, the Clinical Trials Branch, National Eye Institute, National Institutes of Health, Bethesda, Maryland, USA.

⁴⁹Department of Ophthalmology, Hadassah-Hebrew University Medical Center, Jerusalem, Israel.

⁵⁰Department of Medical Genetics, Cambridge Institute for Medical Research, University of Cambridge, Cambridge, UK.

⁵¹Department of Health Promotion and Development, School of Nursing, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.

⁵²Institute for Molecular Biology, University of Oregon, Eugene, Oregon, USA.

⁵³Department of Hygiene and Epidemiology, University of Ioannina Medical School, Ioannina, Greece.

Fagerness [^{54,55}], Henry A Ferreyra [⁵⁶], James S Friedman [⁴⁰], Asbjorg Geirsdottir [⁵⁷], Ronnie J George [⁵⁸], Christian Gieger [⁵⁹], Neel Gupta [⁴⁰], Stephanie A Hagstrom [⁶⁰], Simon P Harding [⁶¹], Christos Haritoglou [⁶²], John R Heckenlively [³⁹], Frank G Holz [⁶³], Guy Hughes [^{56,64}], John P A Ioannidis [^{65,66,67}], Tatsuro Ishibashi [⁶⁸], Peronne Joseph [^{17,18}], Gyungah Jun [^{4,69,70}], Yoichiro Kamatani [⁷¹], Nicholas Katsanis [^{72,73,74}], Claudia N Keilhauer [⁷⁵], Jane C Khan [^{50,76,77}], Ivana K Kim [^{78,79}], Yutaka Kiyohara [⁸⁰], Barbara E K Klein [⁸¹], Ronald Klein [⁸¹], Jaclyn L Kovach [⁸²], Igor Kozak [⁵⁶], Clara J Lee [^{56,64}], Kristine E Lee [⁸¹], Peter Lichtner [⁸³], Andrew J Lotery [⁸⁴], Thomas Meitinger [^{83,85}], Paul Mitchell [⁸⁶], Saddek Mohand-Saïd [^{31,32,36,87}], Anthony T Moore [¹⁴], Denise J Morgan [⁸⁸], Margaux A Morrison [⁸⁸], Chelsea E Myers [⁸¹], Adam C Naj [^{42,43}], Yusuke Nakamura [⁸⁹], Yukinori Okada [⁹⁰], Anton Orlin [⁹¹], M Carolina Ortube [^{92,93}], Mohammad I Othman [³⁹], Chris Pappas [⁸⁸], Kyu Hyung Park [⁹⁴], Gayle J T Pauer [⁶⁰], Neal S Peachey [^{60,95}], Olivier Poch [⁹⁶], Rinki Ratna Priya [⁴⁰], Robyn Reynolds [⁷], Andrea J Richardson [²⁴], Raymond Ripp [⁹⁶], Guenther Rudolph [⁶²],

⁵⁴Center for Human Genetic Research, Massachusetts General Hospital, Boston, Massachusetts, USA.

⁵⁵Program in Medical and Population Genetics, Broad Institute of Harvard and Massachusetts Institute of Technology, Cambridge, Massachusetts, USA.

⁵⁶Department of Ophthalmology and Shiley Eye Center, University of California San Diego, La Jolla, California, USA.

⁵⁷Department of Ophthalmology, National University Hospital, Reykjavik, Iceland.

⁵⁸Department of Glaucoma, Vision Research Foundation, Sankara Nethralaya, Chennai, India.

⁵⁹Institute of Genetic Epidemiology, Helmholtz Zentrum München, Deutsches Forschungszentrum für Gesundheit und Umwelt, Neuherberg, Germany.

⁶⁰Cole Eye Institute, Cleveland Clinic Foundation, Cleveland, Ohio, USA.

⁶¹Department of Eye and Vision Science, Institute of Ageing and Chronic Disease, University of Liverpool, Liverpool, UK.

⁶²Augenklinik, Ludwig-Maximilians-Universität München, München, Germany.

⁶³Department of Ophthalmology, University of Bonn, Bonn, Germany.

⁶⁴Institute for Genomic Medicine, University of California San Diego, La Jolla, California, USA.

⁶⁵Stanford Prevention Research Center, Department of Medicine, Stanford University School of Medicine, Stanford, California, USA.

⁶⁶Department of Health Research and Policy, Stanford University School of Medicine, Stanford, California, USA.

⁶⁷Department of Statistics, Stanford University School of Humanities and Sciences, Stanford, California, USA.

⁶⁸Department of Ophthalmology, Graduate School of Medical Science, Kyushu University, Fukuoka, Japan.

⁶⁹Department of Ophthalmology, Boston University Schools of Medicine and Public Health, Boston, Massachusetts, USA.

⁷⁰Department of Biostatistics, Boston University Schools of Medicine and Public Health, Boston, Massachusetts, USA.

⁷¹Fondation Jean Dausset, Centre d'Etude du Polymorphisme Humain (CEPH), Paris, France.

⁷²Center for Human Disease Modeling, Duke University, Durham, North Carolina, USA.

⁷³Department of Cell Biology, Duke University, Durham, North Carolina, USA.

⁷⁴Department of Pediatrics, Duke University, Durham, North Carolina, USA.

⁷⁵Department of Ophthalmology, Julius-Maximilians-Universität, Würzburg, Germany.

⁷⁶Department of Ophthalmology, Royal Perth Hospital, Perth, Australia.

⁷⁷Centre for Ophthalmology and Visual Science, University of Western Australia, Perth, Australia.

⁷⁸Department of Ophthalmology, Harvard Medical School, Boston, Massachusetts, USA.

⁷⁹Massachusetts Eye and Ear Infirmary, Boston, Massachusetts, USA.

⁸⁰Department of Environmental Medicine, Graduate School of Medical Science, Kyushu University, Fukuoka, Japan.

⁸¹Department of Ophthalmology and Visual Sciences, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, USA.

⁸²Bascom Palmer Eye Institute, University of Miami, Miller School of Medicine, Miami, Florida, USA.

⁸³Institute of Human Genetics, Helmholtz Zentrum München, Deutsches Forschungszentrum für Gesundheit und Umwelt, Neuherberg, Germany.

⁸⁴Faculty of Medicine, Clinical and Experimental Sciences, University of Southampton, Southampton, UK.

⁸⁵Institute of Human Genetics, Technische Universität München, München, Germany.

⁸⁶Centre for Vision Research, Department of Ophthalmology and the Westmead Millennium Institute, University of Sydney, Sydney, Australia.

⁸⁷Department of Therapeutics, Institut de la Vision, UPMC Univ Paris 06, UMR_S 968, Paris, France.

⁸⁸Department of Ophthalmology and Visual Sciences, University of Utah, John A. Moran Eye Center, Salt Lake City, Utah, USA.

⁸⁹Laboratory of Molecular Medicine, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo, Japan.

⁹⁰Laboratory for Statistical Analysis, Center for Genomic Medicine (CGM), RIKEN, Yokohama, Japan.

⁹¹Department of Ophthalmology Weill Cornell Medical College, Weill Cornell Medical College, New York, New York, USA.

⁹²Department of Ophthalmology, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California, USA.

⁹³Jules Stein Eye Institute, Los Angeles, California, USA.

⁹⁴Department of Ophthalmology, Seoul National University Bundang Hospital, Gyeonggi, Rep. of Korea.

⁹⁵Research Service, Louis Stokes Veteran Affairs Medical Center, Cleveland, Ohio, USA.

⁹⁶Laboratory of Integrative Bioinformatics and Genomics, IGBMC, Illkirch, France.

Euijung Ryu [⁹⁷], José-Alain Sahel [^{10,31,32,33,36,98,99}], Debra A Schaumberg [^{78,100}], Hendrik P N Scholl [^{44,63}], Stephen G Schwartz [⁸²], William K Scott [^{42,43}], Humma Shahid [^{50,101}], Haraldur Sigurdsson [^{57,102}], Giuliana Silvestri [¹⁰³], Theru A Sivakumaran [¹⁰⁴], R Theodore Smith [^{25,105}], Lucia Sobrin [^{78,79}], Eric H Souied [¹⁰⁶], Dwight E Stambolian [¹⁰⁷], Hreinn Stefansson [⁸], Gwen M Sturgill-Short [⁹⁵], Atsushi Takahashi [⁹⁰], Nirubol Tosakulwong [⁹⁷], Barbara J Truitt [^{17,18}], Evangelia E Tsirosi [¹⁰⁸], André G Uitterlinden [^{19,109}], Cornelia M van Duijn [¹⁹], Lingam Vijaya [⁵⁸], Johannes R Vingerling [^{19,20}], Eranga N Vithana [^{34,35}], Andrew R Webster [¹⁴], H.-Erich Wichmann [^{110,111,112,113}], Thomas W Winkler [¹¹⁴], Tien Y Wong [^{24,34,35}], Alan F Wright [¹¹⁵], Diana Zelenika [¹¹⁶], Li Zhang [^{56,64,117}], Ling Zhao [^{56,64}], Kang Zhang [^{56,64,117}], Michael L Klein [²⁶], Gregory S Hageman [^{88,118}], G Mark Lathrop [^{71,116}], Kari Stefansson [^{8,102}], Rando Allikmets [^{25,119,+}], Paul N Baird [^{24,+}], Michael B Gorin [^{92,93,120,+}], Jie Jin Wang [^{24,86,+}], Caroline C W Klaver [^{19,20,+}], Johanna M Seddon [^{7,121,+}], Margaret A Pericak-Vance [^{42,43,+}], Sudha K Iyengar [^{17,18,122,123,124,+}], John R W Yates [^{14,50,+}], Anand Swaroop [^{39,40,+}], Bernhard H F Weber [^{1,+}], Michiaki Kubo [^{13,+}], Margaret M DeAngelis [^{88,+}], Thierry Léveillard [^{10,31,32,+}], Unnur Thorsteinsdottir [^{8,102,+}], Jonathan L Haines [^{5,6,+}], Lindsay A Farrer [^{4,69,70,125,126,+}], Iris M Heid [^{59,114,+}], Gonçalo R Abecasis [^{2,+}]

AFFILIATIONS

- ¹Institute of Human Genetics, University of Regensburg, Regensburg, Germany.
²Department of Biostatistics, Center for Statistical Genetics, University of Michigan, Ann Arbor, Michigan, USA.

⁹⁷Division of Biomedical Statistics and Informatics, Mayo Clinic, Rochester, Minnesota, USA.

⁹⁸Fondation Ophtalmologique Adolphe de Rothschild, Paris, France.

⁹⁹Académie des Sciences-Institut de France, Paris, France.

¹⁰⁰Division of Preventive Medicine, Brigham and Women's Hospital, Boston, Massachusetts, USA.

¹⁰¹Department of Ophthalmology, Addenbrooke's Hospital, Cambridge, UK.

¹⁰²Faculty of Medicine, University of Iceland, Reykjavik, Iceland.

¹⁰³Centre for Vision and Vascular Science, Queen's University, Belfast, UK.

¹⁰⁴Division of Human Genetics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA.

¹⁰⁵Department of Biomedical Engineering, Columbia University, New York, New York, USA.

¹⁰⁶Hôpital Intercommunal de Créteil, Hôpital Henri Mondor, Université Paris Est, Créteil, France.

¹⁰⁷Department of Ophthalmology and Genetics, University of Pennsylvania, Philadelphia, Pennsylvania, USA.

¹⁰⁸Department of Ophthalmology, University of Thessaly School of Medicine, Larissa, Greece.

¹⁰⁹Department of Internal Medicine, Erasmus Medical Center, Rotterdam, the Netherlands.

¹¹⁰Institute of Epidemiology I, Helmholtz Zentrum München, Deutsches Forschungszentrum für Gesundheit und Umwelt, Neuherberg, Germany.

¹¹¹Institute of Medical Informatics, Ludwig-Maximilians-Universität, and Klinikum Grosshadern, München, Germany.

¹¹²Institute of Biometry, Ludwig-Maximilians-Universität, and Klinikum Grosshadern, München, Germany.

¹¹³Institute of Epidemiology, Ludwig-Maximilians-Universität, and Klinikum Grosshadern, München, Germany.

¹¹⁴Department of Epidemiology and Preventive Medicine, Regensburg University Medical Center, Regensburg, Germany.

¹¹⁵Medical Research Council Human Genetics Unit, Institute of Genetics and Molecular Medicine, Edinburgh, UK.

¹¹⁶Centre National de Génotypage, CEA - IG, Evry, France.

¹¹⁷Molecular Medicine Research Center and Department of Ophthalmology, West China Hospital, Sichuan University, Chengdu, China.

¹¹⁸Center for Translational Medicine, University of Utah, John A. Moran Eye Center, Salt Lake City, Utah, USA.

¹¹⁹Department of Pathology & Cell Biology, Columbia University, New York, New York, USA.

⁺These authors share starred senior authorship. They directed the project or one of its major component studies.

¹²⁰Department of Human Genetics, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California, USA.

¹²¹Tufts University School of Medicine, Boston, Massachusetts, USA.

¹²²Department of Genetics, Case Western Reserve University, Cleveland, Ohio, USA.

¹²³Department of Ophthalmology, Case Western Reserve University, Cleveland, Ohio, USA.

¹²⁴Department of Clinical Investigation, Case Western Reserve University, Cleveland, Ohio, USA.

¹²⁵Department of Neurology, Boston University Schools of Medicine and Public Health, Boston, Massachusetts, USA.

¹²⁶Department of Epidemiology, Boston University Schools of Medicine and Public Health, Boston, Massachusetts, USA.

- ³Division of Pediatric Pulmonary Medicine, Allergy and Immunology, Department of Pediatrics, Children's Hospital of Pittsburgh of UPMC, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA.
- ⁴Department of Medicine (Section of Biomedical Genetics), Boston University Schools of Medicine and Public Health, Boston, Massachusetts, USA.
- ⁵Center for Human Genetics Research, Vanderbilt University Medical Center, Nashville, Tennessee, USA.
- ⁶Department of Molecular Physiology and Biophysics, Vanderbilt University School of Medicine, Nashville, Tennessee, USA.
- ⁷Ophthalmic Epidemiology and Genetics Service, Tufts Medical Center, Boston, Massachusetts, USA.
- ⁸deCODE genetics, Reykjavik, Iceland.
- ⁹Department of Molecular Biology and Genetics, Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA.
- ¹⁰Department of Genetics, Institut de la Vision, UPMC Univ Paris 06, UMR_S 968, Paris, France.
- ¹¹Department of Neuroscience, Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA.
- ¹²Institute of Genetic Medicine, Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA.
- ¹³Laboratory for Genotyping Development, Research Group for Genotyping, Center for Genomic Medicine (CGM), RIKEN, Yokohama, Japan.
- ¹⁴Moorfields Eye Hospital and Institute of Ophthalmology, University College London, London, UK.
- ¹⁵Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, Massachusetts, USA.
- ¹⁶Stanley Center for Psychiatric Research, Broad Institute of Harvard and Massachusetts Institute of Technology, Cambridge, Massachusetts, USA.
- ¹⁷Department of Epidemiology, Case Western Reserve University, Cleveland, Ohio, USA.
- ¹⁸Department of Biostatistics, Case Western Reserve University, Cleveland, Ohio, USA.
- ¹⁹Department of Epidemiology, Erasmus Medical Center, Rotterdam, the Netherlands.
- ²⁰Department of Ophthalmology, Erasmus Medical Center, Rotterdam, the Netherlands.
- ²¹Centre for Molecular Epidemiology, National University of Singapore, Singapore, Singapore.
- ²²Department of Human Genetics, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.
- ²³Department of Biostatistics, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.
- ²⁴Centre for Eye Research Australia, University of Melbourne, Royal Victorian Eye and Ear Hospital, East Melbourne, Australia.
- ²⁵Department of Ophthalmology, Columbia University, New York, New York, USA.
- ²⁶Macular Degeneration Center, Casey Eye Institute, Oregon Health & Science University, Portland, Oregon, USA.
- ²⁷Department of Bioengineering, University of California San Diego, La Jolla, California, USA.
- ²⁸Vanderbilt Eye Institute, Vanderbilt University Medical Center, Nashville, Tennessee, USA.
- ²⁹Department of Ophthalmology & Visual Sciences, Vanderbilt University School of Medicine, Nashville, Tennessee, USA.
- ³⁰Department of Ophthalmology, University of Edinburgh and Princess Alexandra Eye Pavilion, Edinburgh, UK.
- ³¹INSERM, U968, Paris, France.
- ³²CNRS, UMR_7210, Paris, France.

- ³³Institute of Ophthalmology, University College London, London, UK.
- ³⁴Singapore Eye Research Institute, Singapore National Eye Centre, Singapore, Singapore.
- ³⁵Department of Ophthalmology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore.
- ³⁶Centre Hospitalier National d'Ophtalmologie des Quinze-Vingts, INSERM-DHOS CIC 503, Paris, France.
- ³⁷School of Biomedicine, Faculty of Medical and Human Sciences, University of Manchester, Manchester, UK.
- ³⁸Central Manchester University Hospitals NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, UK.
- ³⁹Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, Michigan, USA.
- ⁴⁰Neurobiology Neurodegeneration & Repair Laboratory (N-NRL), National Eye Institute, National Institutes of Health, Bethesda, Maryland, USA.
- ⁴¹Penn Presbyterian Medical Center, Scheie Eye Institute, Philadelphia, Pennsylvania, USA.
- ⁴²John P. Hussman Institute for Human Genomics, University of Miami Miller School of Medicine, Miami, Florida, USA.
- ⁴³Dr. John T. Macdonald Foundation Department of Human Genetics, University of Miami Miller School of Medicine, Miami, Florida, USA.
- ⁴⁴Department of Ophthalmology, Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA.
- ⁴⁵Immunopathology Section, Laboratory of Immunology, National Eye Institute, National Institutes of Health, Bethesda, Maryland, USA.
- ⁴⁶Saw Swee Hock School of Public Health, National University of Singapore, Singapore, Singapore.
- ⁴⁷Centre for Quantitative Medicine, Office of Clinical Sciences, Duke-NUS Graduate Medical School, Singapore, Singapore.
- ⁴⁸Division of Epidemiology and Clinical Applications, the Clinical Trials Branch, National Eye Institute, National Institutes of Health, Bethesda, Maryland, USA.
- ⁴⁹Department of Ophthalmology, Hadassah-Hebrew University Medical Center, Jerusalem, Israel.
- ⁵⁰Department of Medical Genetics, Cambridge Institute for Medical Research, University of Cambridge, Cambridge, UK.
- ⁵¹Department of Health Promotion and Development, School of Nursing, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.
- ⁵²Institute for Molecular Biology, University of Oregon, Eugene, Oregon, USA.
- ⁵³Department of Hygiene and Epidemiology, University of Ioannina Medical School, Ioannina, Greece.
- ⁵⁴Center for Human Genetic Research, Massachusetts General Hospital, Boston, Massachusetts, USA.
- ⁵⁵Program in Medical and Population Genetics, Broad Institute of Harvard and Massachusetts Institute of Technology, Cambridge, Massachusetts, USA.
- ⁵⁶Department of Ophthalmology and Shiley Eye Center, University of California San Diego, La Jolla, California, USA.
- ⁵⁷Department of Ophthalmology, National University Hospital, Reykjavik, Iceland.
- ⁵⁸Department of Glaucoma, Vision Research Foundation, Sankara Nethralaya, Chennai, India.
- ⁵⁹Institute of Genetic Epidemiology, Helmholtz Zentrum München, Deutsches Forschungszentrum für Gesundheit und Umwelt, Neuherberg, Germany.
- ⁶⁰Cole Eye Institute, Cleveland Clinic Foundation, Cleveland, Ohio, USA.
- ⁶¹Department of Eye and Vision Science, Institute of Ageing and Chronic Disease, University of Liverpool, Liverpool, UK.

- ⁶²Augenklinik, Ludwig-Maximilians-Universität München, München, Germany.
- ⁶³Department of Ophthalmology, University of Bonn, Bonn, Germany.
- ⁶⁴Institute for Genomic Medicine, University of California San Diego, La Jolla, California, USA.
- ⁶⁵Stanford Prevention Research Center, Department of Medicine, Stanford University School of Medicine, Stanford, California, USA.
- ⁶⁶Department of Health Research and Policy, Stanford University School of Medicine, Stanford, California, USA.
- ⁶⁷Department of Statistics, Stanford University School of Humanities and Sciences, Stanford, California, USA.
- ⁶⁸Department of Ophthalmology, Graduate School of Medical Science, Kyushu University, Fukuoka, Japan.
- ⁶⁹Department of Ophthalmology, Boston University Schools of Medicine and Public Health, Boston, Massachusetts, USA.
- ⁷⁰Department of Biostatistics, Boston University Schools of Medicine and Public Health, Boston, Massachusetts, USA.
- ⁷¹Fondation Jean Dausset, Centre d'Etude du Polymorphisme Humain (CEPH), Paris, France.
- ⁷²Center for Human Disease Modeling, Duke University, Durham, North Carolina, USA.
- ⁷³Department of Cell Biology, Duke University, Durham, North Carolina, USA.
- ⁷⁴Department of Pediatrics, Duke University, Durham, North Carolina, USA.
- ⁷⁵Department of Ophthalmology, Julius-Maximilians-Universität, Würzburg, Germany.
- ⁷⁶Department of Ophthalmology, Royal Perth Hospital, Perth, Australia.
- ⁷⁷Centre for Ophthalmology and Visual Science, University of Western Australia, Perth, Australia.
- ⁷⁸Department of Ophthalmology, Harvard Medical School, Boston, Massachusetts, USA.
- ⁷⁹Massachusetts Eye and Ear Infirmary, Boston, Massachusetts, USA.
- ⁸⁰Department of Environmental Medicine, Graduate School of Medical Science, Kyushu University, Fukuoka, Japan.
- ⁸¹Department of Ophthalmology and Visual Sciences, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, USA.
- ⁸²Bascom Palmer Eye Institute, University of Miami, Miller School of Medicine, Miami, Florida, USA.
- ⁸³Institute of Human Genetics, Helmholtz Zentrum München, Deutsches Forschungszentrum für Gesundheit und Umwelt, Neuherberg, Germany.
- ⁸⁴Faculty of Medicine, Clinical and Experimental Sciences, University of Southampton, Southampton, UK.
- ⁸⁵Institute of Human Genetics, Technische Universität München, München, Germany.
- ⁸⁶Centre for Vision Research, Department of Ophthalmology and the Westmead Millennium Institute, University of Sydney, Sydney, Australia.
- ⁸⁷Department of Therapeutics, Institut de la Vision, UPMC Univ Paris 06, UMR_S 968, Paris, France.
- ⁸⁸Department of Ophthalmology and Visual Sciences, University of Utah, John A. Moran Eye Center, Salt Lake City, Utah, USA.
- ⁸⁹Laboratory of Molecular Medicine, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo, Japan.
- ⁹⁰Laboratory for Statistical Analysis, Center for Genomic Medicine (CGM), RIKEN, Yokohama, Japan.
- ⁹¹Department of Ophthalmology Weill Cornell Medical College, Weill Cornell Medical College, New York, New York, USA.
- ⁹²Department of Ophthalmology, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California, USA.
- ⁹³Jules Stein Eye Institute, Los Angeles, California, USA.

- 94Department of Ophthalmology, Seoul National University Bundang Hospital, Kyeonggi, Rep. of Korea.
- 95Research Service, Louis Stokes Veteran Affairs Medical Center, Cleveland, Ohio, USA.
- 96Laboratory of Integrative Bioinformatics and Genomics, IGBMC, Illkirch, France.
- 97Division of Biomedical Statistics and Informatics, Mayo Clinic, Rochester, Minnesota, USA.
- 98Fondation Ophtalmologique Adolphe de Rothschild, Paris, France.
- 99Académie des Sciences–Institut de France, Paris, France.
- 100Division of Preventive Medicine, Brigham and Women’s Hospital, Boston, Massachusetts, USA.
- 101Department of Ophthalmology, Addenbrooke’s Hospital, Cambridge, UK.
- 102Faculty of Medicine, University of Iceland, Reykjavik, Iceland.
- 103Centre for Vision and Vascular Science, Queen’s University, Belfast, UK.
- 104Division of Human Genetics, Cincinnati Children’s Hospital Medical Center, Cincinnati, Ohio, USA.
- 105Department of Biomedical Engineering, Columbia University, New York, New York, USA.
- 106Hôpital Intercommunal de Créteil, Hôpital Henri Mondor, Université Paris Est, Créteil, France.
- 107Department of Ophthalmology and Genetics, University of Pennsylvania, Philadelphia, Pennsylvania, USA.
- 108Department of Ophthalmology, University of Thessaly School of Medicine, Larissa, Greece.
- 109Department of Internal Medicine, Erasmus Medical Center, Rotterdam, the Netherlands.
- 110Institute of Epidemiology I, Helmholtz Zentrum München, Deutsches Forschungszentrum für Gesundheit und Umwelt, Neuherberg, Germany.
- 111Institute of Medical Informatics, Ludwig-Maximilians-Universität, and Klinikum Grosshadern, München, Germany.
- 112Institute of Biometry, Ludwig-Maximilians-Universität, and Klinikum Grosshadern, München, Germany.
- 113Institute of Epidemiology, Ludwig-Maximilians-Universität, and Klinikum Grosshadern, München, Germany.
- 114Department of Epidemiology and Preventive Medicine, Regensburg University Medical Center, Regensburg, Germany.
- 115Medical Research Council Human Genetics Unit, Institute of Genetics and Molecular Medicine, Edinburgh, UK.
- 116Centre National de Génotypage, CEA - IG, Evry, France.
- 117Molecular Medicine Research Center and Department of Ophthalmology, West China Hospital, Sichuan University, Chengdu, China.
- 118Center for Translational Medicine, University of Utah, John A. Moran Eye Center, Salt Lake City, Utah, USA.
- 119Department of Pathology & Cell Biology, Columbia University, New York, New York, USA.
- 120Department of Human Genetics, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California, USA.
- 121Tufts University School of Medicine, Boston, Massachusetts, USA.
- 122Department of Genetics, Case Western Reserve University, Cleveland, Ohio, USA.
- 123Department of Ophthalmology, Case Western Reserve University, Cleveland, Ohio, USA.
- 124Department of Clinical Investigation, Case Western Reserve University, Cleveland, Ohio, USA.
- 125Department of Neurology, Boston University Schools of Medicine and Public Health, Boston, Massachusetts, USA.

¹²⁶Department of Epidemiology, Boston University Schools of Medicine and Public Health, Boston, Massachusetts, USA.

*These authors contributed equally to this work.

[†]These authors share starred senior authorship. They directed the project or one of its major component studies.

AUTHOR CONTRIBUTIONS

AMD Gene Analysis Committee: L.G.F., W.C., M.S., B.L.Y., Y.Y., L.A.F., I.M.H. (co-lead), G.R.A. (co-lead);

AMD Gene Steering Committee: B.H.F.W. (chair, senior executive committee), G.R.A. (senior executive committee), M.M.D. (senior executive committee), J.L.H. (senior executive committee), S.K.I. (senior executive committee), M.A.P. (senior executive committee), R.A., P.N.Ba., C.C.W.K., B.E.K.K., M.L.K., M.K., T.L., J.M.S., U.T., D.E.W., J.R.W.Y., K.Z.;

AMD-EU-JHU Study: D.J.Z., I.A., M.Be., A.C.B., P.A.C., I.C., F.G.H., Y.Ka., N.K., A.J.L., S.M., O.P., R.Ri., J-A.S., H.P.N.S., E.H.S., A.R.W., D.Z., G.M.L., T.L. contributed phenotype, genotypes and analysis for the AMD-EU-JHU study;

BDES Study: R.P.I., B.E.K.K., R.K., K.E.L., C.E.M., T.A.S., B.J.T., S.K.I. contributed phenotype, genotypes and analysis for the BDES study;

Blue Mountains Eye Study: X.S., P.M., T.Y.W., J.J.W. contributed phenotype, genotypes and analysis for the BMES study;

BU/Utah Study: M.S., G.S.H., G.J., I.K.K., D.J.M., M.A.M., C.P., K.H.P., D.A.S., G.S., E.E.T., M.M.D., L.A.F. contributed phenotype, genotypes and analysis for the BU/UTAH study;

CCF/VAMC Study: S.A.H., P.J., G.J.T.P., N.S.P., G.M.S., R.P.I., S.K.I. contributed phenotype, genotypes and analysis for the CCF/VAMC study;

CEI Study: P.J.F., M.L.K. contributed phenotype, genotypes and analysis for the CEI study;

Columbia Study: J.E.M., G.R.B., R.T.S., R.A. contributed phenotype, genotypes and analysis for the Columbia study;

deCode: G.T., H.Si., H.St., K.S., U.T. contributed phenotype, genotypes and analysis for the deCode study;

Japan Age Related Eye Diseases Study: S.A., T.I., Y.Ki., Y.N., Y.O., A.T., M.K. contributed phenotype, genotypes and analysis for the JAREDS study;

Melbourne Study: R.H.G., M.R.N.C., A.J.R., P.N.Ba. contributed phenotype, genotypes and analysis for the Melbourne study;

Miami/Vanderbilt Study: B.L.Y., A.A., W.H.C., J.L.K., A.C.N., S.G.S., W.K.S., M.A.P., J.L.H. contributed phenotype, genotypes and analysis for the Miami/Vanderbilt study;

MMAP/NEI Study: W.C., K.E.B., M.Br., A.J.B., C-C.C., E.Y.C., R.C., A.O.E., J.S.F., N.G., J.R.H., A.O., M.I.O., R.R.P., E.R., D.E.S., N.T., A.S., G.R.A. contributed phenotype, genotypes and analysis for the MMAP/NEI study;

Rotterdam Study: G.H.S.B., A.G.U., C.M.v.D., J.R.V., C.C.W.K. contributed phenotype, genotypes and analysis for the Rotterdam study;

SAGE Study: T.A., C-Y.C., B.K.C., E.N.V. contributed phenotype, genotypes and analysis for the SAGe study;

Southern German AMD Study: L.G.F., C.G., C.H., C.N.K., P.L., T.M., G.R., H.-E.W., T.W.W., B.H.F.W., I.M.H. contributed phenotype, genotypes and analysis for the Southern German AMD study;

Tufts/Massachusetts General Hospital Study: Y.Y., S.R., K.A.C., M.J.D., E.E., J.F., J.P.A.I., R.Re., L.S., J.M.S. contributed phenotype, genotypes and analysis for the Tufts/ MGH study;

U.K. Cambridge/Edinburgh Study: V.C., A.M.A., P.N.Bi., D.G.C., B.D., S.P.H., J.C.K., A.T.M., H.Sh., A.F.W., J.R.W.Y. contributed phenotype, genotypes and analysis for the UK Cambridge/Edinburgh study;

University of Pittsburgh/UCLA Study: D.E.W., Y.P.C., M.C.O., M.B.G. contributed phenotype, genotypes and analysis for the Univ. of Pittsburgh/UCLA study;

UCSD Study: G.Ha., H.F., G.Hu., I.K., C.J.L., L.Zhang, L.Zhao, K.Z. contributed phenotype, genotypes and analysis for the USCD study;

VRF Study: R.J.G., L.V., R.P.I., S.K.I. contributed phenotype, genotypes and analysis for the VRF study;

Gene Expression and RNA-Sequencing Data: These data were contributed and analyzed by M.Br., J.S.F., N.G., R.R.P and A.S.

URLs

METAL, <http://www.sph.umich.edu/csg/abecasis/Metal/>; R, <http://www.R-project.org/>; gee, <http://CRAN.R-project.org/package=gee>; Single Nucleotide Polymorphism Spectral Decomposition, <http://gump.qimr.edu.au/general/daleN/SNPSPDLite/>; Pre-specified R-scripts, <http://www.epi-regensburg.de/wp/genepi-downloads>; The 1000 Genomes Project, <http://www.1000genomes.org/>; The HapMap Project, <http://www.hapmap.org/genotypes/>; PolyPhen-2, <http://genetics.bwh.harvard.edu/pph2/>; Ingenuity® Systems, <http://www.ingenuity.com>; NHGRI GWAS catalog, <http://www.genome.gov/gwastudies/>; INRICH, <http://atgu.mgh.harvard.edu/inrich>; Full Result Set, <http://www.sph.umich.edu/csg/abecasis/public/amdgene2012/>

CONFLICT OF INTEREST STATEMENT

A.A., G.R.A., K.E.B., V.C., Y.P.C., M.J.D., A.O.E., L.G.F., M.B.G., J.L.H., A.T.M., D.A.S., W.K.S., J.M.S., A.S., B.H.F.W., D.E.W., and J.R.W.Y. are co-inventors or beneficiaries of patents related to genetic discoveries in AMD. J.L.H. is a shareholder in ArcticDX. S.G.S. is a consultant for Alimera, Bausch + Lomb, and Eyeteck; receives royalties from IC Labs. U.T., K.S., G.T, and H.St. are affiliated and/or employed by deCODE Genetics and own stock and/or stock options in the company. P.M. is on advisory boards for Allergan, Bayer, Novartis, Pfizer and Solvay and has received travel, honorarium and research support from these companies; he has no stocks, equity, contract of employment or named position on company board.




FIGURE 1. Summary of genomewide association scan results

Summary of genomewide association scan results in the discovery GWAS sample.

Previously described loci reaching $p < 5 \times 10^{-8}$ are labeled in blue; new loci reaching $p < 5 \times 10^{-8}$ for the first time after follow-up are labeled in green.




FIGURE 2. Sensitivity analysis

The top left panel compares estimated effect sizes for the original analysis and for an age-adjusted analysis (where age was included as a covariate and samples of unknown age were excluded). The top right panel compares analyses stratified by sex. The bottom left panel evaluates stratification by disease subtype. The bottom right panel evaluates stratification by ethnicity. The size of each marker reflects confidence intervals (with height reflecting confidence interval along the Y axis and width reflecting confidence interval along the X axis). Comparisons reaching $p < 0.05$ are labeled and colored in red.




FIGURE 3. Risk score analysis

We calculated a risk score for each individual, defined as the product of the number of risk alleles at each locus and the associated effect size for each allele (measured on the log-odds scale). The plot summarizes the ability of these overall genetic risk scores to distinguish cases and controls.

TABLE 1
Summary of the Samples Used in Genomewide Discovery and Targeted Follow-Up Analyses

For additional details, including a breakdown of the number of cases and controls in individual samples, see Supplementary Table 1. N_{CASES} includes only cases with geographic atrophy, choroidal neovascularization, or both.

Analysis	Contributing Study Groups	N _{CASES}	%Female	%Neovascular Disease	N _{CONTROLS}	% Female
Genomewide Discovery	15	7,650	53.9	59.2	51,844	45.2
Targeted Follow-up	18	9,531	56.3	57.8	8,230	53.8
Overall	33	17,181	55.2	58.4	60,074	46.3

TABLE 2
Summary of Loci Reaching Genome-Wide Significance

All results reported here include a genomic control correction for individual studies and also for the final meta-analysis⁵¹.

SNP/Risk Allele	Chromosome, Position	Nearby Genes	EAF	Discovery		Follow-up		Joint			
				P	OR	P	OR	P	OR		
Loci Previously Reported With $P < 5 \times 10^{-8}$											
rs10490924/T	10	124.2 Mb	ARMS2/HTRA1	0.30	4×10^{-353}	2.71	2.8×10^{-190}	2.88	4×10^{-540}	2.76	[2.72–2.80]
rs10737680/A	1	196.7 Mb	CFH	0.64	1×10^{-283}	2.40	2.7×10^{-152}	2.50	1×10^{-434}	2.43	[2.39–2.47]
rs429608/G	6	31.9 Mb	C2/CFB	0.86	2×10^{-54}	1.67	2.4×10^{-37}	1.89	4×10^{-89}	1.74	[1.68–1.79]
rs2230199/C	19	6.7 Mb	C3	0.20	2×10^{-26}	1.46	3.4×10^{-17}	1.37	1×10^{-41}	1.42	[1.37–1.47]
rs5749482/G	22	33.1 Mb	TIMP3	0.74	6×10^{-13}	1.25	9.7×10^{-17}	1.45	2×10^{-26}	1.31	[1.26–1.36]
rs4420638/A	19	45.4 Mb	APOE	0.83	3×10^{-15}	1.34	4.2×10^{-7}	1.25	2×10^{-20}	1.30	[1.24–1.36]
rs1864163/G	16	57 Mb	CETP	0.76	8×10^{-13}	1.25	8.7×10^{-5}	1.17	7×10^{-16}	1.22	[1.17–1.27]
rs943080/T	6	43.8 Mb	VEGFA	0.51	4×10^{-12}	1.18	1.6×10^{-5}	1.12	9×10^{-16}	1.15	[1.12–1.18]
rs13278062/T	8	23.1 Mb	TNFRSF10A	0.48	7×10^{-10}	1.17	6.4×10^{-7}	1.14	3×10^{-15}	1.15	[1.12–1.19]
rs920915/C	15	58.7 Mb	LIPC	0.48	2×10^{-9}	1.14	0.004	1.10	3×10^{-11}	1.13	[1.09–1.17]
rs4698775/G	4	110.6 Mb	CFI	0.31	2×10^{-10}	1.16	0.025	1.08	7×10^{-11}	1.14	[1.10–1.17]
rs3812111/T	6	116.4 Mb	COL10A1	0.64	7×10^{-8}	1.13	0.022	1.06	2×10^{-8}	1.10	[1.07–1.14]
Loci Reaching $P < 5 \times 10^{-8}$ for the First Time											
rs13081855/T	3	99.5 Mb	COL8A1/FILIPIL	0.10	4×10^{-11}	1.28	6.0×10^{-4}	1.17	4×10^{-13}	1.23	[1.17–1.29]
rs3130783/A	6	30.8 Mb	IER3/DDR1	0.79	1×10^{-6}	1.15	3.5×10^{-6}	1.16	2×10^{-11}	1.16	[1.11–1.20]
rs8135665/T	22	38.5 Mb	SLC16A8	0.21	8×10^{-8}	1.16	5.6×10^{-5}	1.13	2×10^{-11}	1.15	[1.11–1.19]
rs334353/T	9	101.9 Mb	TGFBR1	0.73	9×10^{-7}	1.13	6.7×10^{-6}	1.13	3×10^{-11}	1.13	[1.10–1.17]
rs8017304/A	14	68.8 Mb	RAD51B	0.61	9×10^{-7}	1.11	2.1×10^{-5}	1.11	9×10^{-11}	1.11	[1.08–1.14]
rs6795735/T	3	64.7 Mb	ADAMTS9/MIR548A2	0.46	9×10^{-8}	1.13	0.0066	1.07	5×10^{-9}	1.10	[1.07–1.14]
rs9542236/C	13	31.8 Mb	B3GALT1	0.44	2×10^{-6}	1.12	0.0018	1.08	2×10^{-8}	1.10	[1.07–1.14]

See Supplementary Table 5 for a summary of all gene name abbreviations used in this Table and elsewhere in the paper. EAF is the allele frequency of the risk increasing allele.

TABLE 3

Pathway Analysis

Ingenuity Canonical Pathways	Enrichment Analysis		
	Nominal p-value	FDR q-value	Pathway Size(N _{genes})
Complement System	0.000012	0.0015	35
Atherosclerosis Signaling	0.00014	0.009	129
VEGF Family Ligand-Receptor Interactions	0.0042	0.150	84
Dendritic Cell Maturation	0.0046	0.150	185
Phospholipid Degradation	0.0058	0.151	102
MIF-mediated Glucocorticoid Regulation	0.0088	0.153	42
Inhibition of Angiogenesis by TSP1	0.0093	0.153	39
Fc Epsilon RI Signaling	0.0098	0.153	111
p38 MAPK Signaling	0.011	0.153	106

* *CFB*, *C2*, *C4A*, and *C4B* all flank rs429608 and thus counted as single hit when determining significance of enrichment.

** *APOC1*, *APOE*, *APOC2*, and *APOC4* all flank rs4420638 and thus counted as single hit when determining significance of enrichment.