

THE UNIVERSITY of EDINBURGH

Edinburgh Research Explorer

Genetics of animal temperament: aggressive behaviour at mixing is genetically associated with the response to handling in pigs

Citation for published version:

D'Eath, RB, Roehe, R, Turner, SP, Ison, SH, Farish, M, Jack, MC & Lawrence, AB 2009, 'Genetics of animal temperament: aggressive behaviour at mixing is genetically associated with the response to handling in pigs', *Animal*, vol. 3, no. 11, pp. 1544-1554. https://doi.org/10.1017/S1751731109990528

Digital Object Identifier (DOI):

10.1017/S1751731109990528

Link:

Link to publication record in Edinburgh Research Explorer

Document Version: Publisher's PDF, also known as Version of record

Published In: Animal

Publisher Rights Statement: Copyright 2009 The Animal Consortium

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Édinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Genetics of animal temperament: aggressive behaviour at mixing is genetically associated with the response to handling in pigs

R. B. D'Eath[†], R. Roehe, S. P. Turner, S. H. Ison, M. Farish, M. C. Jack and A. B. Lawrence

Sustainable Livestock Systems, Scottish Agricultural College, West Mains Road, Edinburgh, EH9 3JG, UK

(Received 30 January 2009; Accepted 22 June 2009; First published online 13 July 2009)

Aggression when pigs are mixed into new social groups has negative impacts on welfare and production. Aggressive behaviour is moderately heritable and could be reduced by genetic selection. The possible wider impacts of selection for reduced aggressiveness on handling traits and activity in the home pen were investigated using 1663 male and female pedigree pigs (898 purebred Yorkshire and 765 Yorkshire \times Landrace). Aggressive behaviour was observed over 24 h after pigs were mixed at 10 weeks of age into groups balanced for unfamiliarity and weight. Aggression was highly heritable (duration of involvement in reciprocal fighting $h^2 = 0.47 \pm 0.03$, and duration of delivering one-sided aggression $h^2 = 0.34 \pm 0.03$). Three weeks after mixing, home pen inactivity (indicated by the frequency of lying) was observed over 24 h. Inactivity was weakly heritable ($h^2 = 0.05 \pm 0.01$) but showed no significant genetic association with aggression. Pigs' behaviour during handling by humans was assessed on entry to, whilst inside and on exit from a weigh crate at both mixing and end of test at 22 weeks. Pigs were generally easy to handle, moving easily into and out of the crate. Scores indicating 'very difficult to move' were rare. Handling scores at weighing were weakly heritable ($h^2 = 0.03$ to 0.17), and moderately correlated across the two weighings ($r_a = 0.28$ to 0.76). Aggressive behaviour at mixing was genetically associated with handling at the end of test weighing: pigs that fought and delivered one-sided aggression had handling scores indicating more active behaviour at weighing (e.g. moving quickly into the crate v. fighting $r_q = 0.41 \pm 0.05$ and v. bullying $r_q = 0.60 \pm 0.04$). Also, there was a genetic association between receiving one-side aggression at mixing and producing high-pitched vocalisations in the weigh crate ($r_q = 0.78 \pm 0.08$). Correlated behavioural responses occurring across different challenging situations (e.g. social mixing and human handling) have been described by the concept of animal temperament (also known as coping styles, personality or behavioural syndromes), but this has rarely been demonstrated at the genetic level in farm animals. These findings may have practical implications for the development of breeding programmes aimed at altering animal temperament. Breeding to reduce aggression could result in some reduction in activity at weighing. This would have consequences for animal production, because pigs which are inactive at weighing take longer to move into and out of the weigh crate, and perhaps also for animal welfare.

Keywords: social behaviour, breeding, coping styles, genetic parameters, behavioural syndromes

Implications

Aggression when pigs are mixed with unfamiliar individuals causes stress. Genetic selection for pigs showing low aggression could reduce stress and improve animal welfare. In a behavioural genetics study on 1663 pigs in one farm, we looked at responses to two challenges: (1) mixing with unfamiliar pigs, which results in aggression and (2) being moved through a weigh crate. These two responses were

moderately associated at the genetic level, meaning that genetic selection to reduce aggression could result in pigs that are somewhat less active (and perhaps calmer?) in the weigh crate at weighing.

Introduction

The behavioural responses of farmed livestock to their social and physical environment can affect animal welfare, production and health. Within a population, individuals show differences in behaviour, which are stable across time

⁺ E-mail: rick.death@sac.ac.uk

and situations. This is referred to as animal temperament, personality or behavioural syndromes (Sih *et al.*, 2004; Réale *et al.*, 2007). Examples of temperament traits in livestock include aggressiveness (D'Eath, 2002), maternal traits (Jarvis *et al.*, 2005), responses to isolation (Boissy *et al.*, 2005), and responses to humans or handling (Phocas *et al.*, 2006).

Although experiences during development are also important (e.g. D'Eath and Lawrence, 2004; D'Eath, 2005), such temperament traits have been shown to be partly under genetic control (Van Oers et al., 2005; Réale et al., 2007). This raises the possibility that genetic selection could be used to alter animal temperament, resulting in animals that are better adapted to the farmed environment. For example in pigs, specific behavioural traits which could be altered by genetic selection include social aggression, tail-biting, savaging of newborn piglets by sows and other aspects of piglet and sow behaviour that contribute to piglet survival (Kanis et al., 2004; Roehe et al., 2009; Turner et al., 2009). Aggressive behaviours following the mixing of pigs are moderately heritable ($h^2 = 0.17$ to 0.24, Løvendahl *et al.*, 2005; $h^2 = 0.37$ to 0.46, Turner *et al.*, 2008; $h^2 = 0.31$ to 0.43, Turner *et al.*, 2009). An associated indicator trait (skin lesion counts 24 h after mixing) is also heritable ($h^2 = 0.22$, Turner *et al.*, 2006; $h^2 = 0.12$ to 0.20, Turner *et al.*, 2008; $h^2 = 0.21$ to 0.26, Turner *et al.*, 2009) and is genetically correlated with aggressive behaviour $(r_{\rm q} = 0.56-0.77, \text{ Turner } et al., 2008; r_{\rm g} = 0.67-0.79, \text{ Turner}$ et al., 2009). This suggests that skin lesions could be used as a proxy measurement, which is the result of aggression, and would be easier to apply in practical animal breeding than observations of behaviour. Aggressive behaviour in pigs has a number of negative impacts on animal welfare including skin injuries, social stress and impaired immune function (D'Eath, 2002; Turner et al., 2006). A number of these traits also have production impacts affecting growth, carcass gradings and meat quality (Warriss et al., 1998; D'Eath, 2002). Selection to reduce aggressive behaviour in pigs thus appears to be both feasible and desirable.

However, selection for one trait may affect other traits, mainly because single genes may have more than one effect (pleiotropy) and because genes do not segregate randomly in the population (linkage disequilibrium; Falconer and Mackay, 1996). The correlation of the breeding values of two traits is known as the genetic correlation (Falconer and Mackay, 1996). However, selection against aggressive behaviour may affect other genetically correlated traits, thereby affecting the feasibility and desirability of adding behavioural traits to a breeding programme.

In the present study, we investigated the genetic relationships between aggressive behaviour at mixing and two further behaviour traits of potential practical and ethical importance, in a pedigree herd of 1663 pigs. For example, lower aggressiveness may be genetically correlated with reduced activity of animals in any situation. To investigate this, we observed activity in the home pen (3 weeks after mixing) over 24 h. Conversely, high aggressiveness in pigs may be associated with increased reactivity in other challenging situations (Thodberg *et al.*, 1999; Ruis *et al.*, 2000; but see D'Eath and Burn, 2002) as found in a number of other species ('coping styles' reviewed by Koolhaas *et al.* (2007)). If there is a genetic relationship between these traits, then selection for reduced aggression would lead to reduced responses to other challenges. To investigate this, we scored responses to a practically relevant challenge: handling during weighing, on two occasions 85 days apart. This situation contains a number of different aspects that could affect how pigs respond: presence of human handlers (Hemsworth *et al.*, 1990), novelty, restraint and partial isolation from penmates.

Material and methods

The work described here was carried out at a farm in Ransta, Sweden as part of a larger study on the genetics of behaviour in pigs, aspects of which have been described in detail in Turner *et al.* (2009). The work was subjected to an ethical appraisal by the Animal Experiments Committee at the Scottish Agricultural College (SAC), and permission for the study was granted by Sweden's animal protection authority (Djur Skydds Myndigheten).

Animals and housing

The genetic analyses were carried out on a pedigree population of pigs (n = 1663) from a dam-line nucleus, consisting of 898 Swedish Yorkshire and 765 Swedish Yorkshire × Swedish Landrace, of which 582 were entire males, 222 castrates and 859 were female. Pigs were the progeny of 86 sires and 251 dams, and their full pedigree back to the great-grandparent generation was known.

Pigs were born in open farrowing pens (5.76 m² with 0.24 m² creep area) and, after weaning, were housed in these pens until 70.5 \pm 4.3 (mean \pm s.d.) days of age, when the mean live-weight was 27.7 \pm 5.5 kg. At this stage, pigs were transported to a different building and mixed into new groups of 15 (details below) in part-slatted pens (29% slats, 71% shallow sawdust bedding over a solid floor) measuring 4.0 m \times 3.2 m (0.85 m² per pig). Dry pelleted food was provided *ad libitum* from a single-space feeder. Pigs remained in these pens until after 156.7 \pm 5.9 days of age (live-weight 105.0 \pm 11.9 kg) when the last measurements were taken. The mean (\pm s.d.) temperature was 19.4 (\pm 2.9)°C.

The farm employed a batch-farrowing system, and the experiment used 105 to 120 pigs from each of 14 batches, spread over a 14-month period.

Observations of aggressive behaviour at mixing

This part of the study has been described in depth (Turner *et al.*, 2009), but briefly, 109 groups of 15 pigs and two groups of 14 pigs were mixed into new groups (n = 1663). These new groups consisted of entirely purebred (Swedish Yorkshire) or crossbred (Swedish Yorkshire × Swedish Landrace)

D'Eath, Roehe, Turner, Ison, Farish, Jack and Lawrence

pigs. Group composition in terms of the number and proportion of pigs from different groups entering a new group can affect the overall level and targets of aggression. Consequently, group composition was kept uniform and balanced: single-sex groups of 15 were composed of three pigs from each of five littermate groups. Since pig size is a predictor of participation and success in aggressive interactions (D'Eath, 2002), weight variation within a group was minimised.

On the day of mixing, each animal was given an individually distinct spray mark and an injection against mycoplasma, which was routine on this farm. Three pig littermate subgroups were selected from their pen and were kept separately in a trailer for transport. They were then transported for approximately 10 min over a distance of 4 km and mixed into their new groups of 15 on arrival. Then, 24-h time-lapse video recordings of these newly mixed groups were made, and each individual's aggressive behaviour was subsequently observed for analysis using Noldus Observer software. The total time each pig spent in reciprocal aggression (in the following referred to as 'fighting') and in both delivery and receipt of non-reciprocal aggression (in the following referred to as 'bullying' and 'being bullied') were recorded. Reciprocal aggression (fighting) was defined as aggressive interactions lasting \geq 1 s in which both pigs were seen to be pushing, head knocking and/or biting each other, and non-reciprocal aggression was the same, but delivered by one pig with no retaliation by the opponent. At 24 h after mixing, the pigs were weighed as detailed below.

Scoring of the response to handling at weighing

Pigs were weighed on two occasions: (i) 24 h after mixing on day 71.5 \pm 4.3 (in the following referred to as 'weighing at mixing'), and (ii) at the end of the test period at 156.7 \pm 5.9 days of age (in the following referred to as 'weighing at end of test'). Handling scores were collected at these times from 1620 pigs (29 pigs had died by the second weighing and 14 pigs were excluded due to ill health, primarily due to lameness). On each occasion, one group of pigs at a time were removed from their pen into one half of a corridor area (28.0 m \times 1.5 m) with a weight crate, suitable for the weight of pigs studied, placed half way along the corridor and positioned so as to block pig access. In an ad-hoc order, pigs were then, (a) moved into the weigh crate, (b) weighed and (c) allowed to leave the weigh crate to access the other half of the corridor. At these three scoring points, each pig's behaviour was scored on a 3- or 5-point scale according to the system given in Table 1, where low values indicate difficult-to-handle pigs and high values indicate easy-to-handle pigs. Also, whether or not the pig produced high-pitched 'squeal' vocalisations whilst it was in the weight crate was recorded (Marchant et al., 2001). The order in which pigs were weighed was noted. After weighing, all pigs were returned to their home pen, and another group weighed and scored in a similar way. Scoring of pigs was always performed by the same individual.

Tabla	1	Scoring	cuctom	ucod	for	handling	during	waighing
lable		Scoring	System	useu	101	nanunny	uunny	weigining

Score	Description
Moving into the c	rate
1	Pig is very difficult to move and is trying to escape
2	Pig is <i>difficult</i> to move into the crate
3	Pig moves into the crate with <i>some assistance</i> from stockperson
4	Pig walks into the crate with <i>little or no</i> encouragement
5	Pig runs forward into the crate
In the crate	
1	Pig <i>moves around a lot</i> during weighing, jumping and crashing around
2	Pig moves around during weighing
3	Pig stands still during weighing
Leaving the crate	
1	Pig resists and is very difficult to push out of the weigh crate
2	Pig moves out of the weigh crate after <i>some</i> pushing
3	Pig leaves <i>of its own accord</i> once the door is opened

Observations of inactivity in the home pen

Aggressive behaviour associated with the formation of a new group is most intense during the first few hours after mixing (D'Eath, 2002) and typically declines to baseline over the next several days. These home pen observations were thus intended to reflect behaviour in a relatively stable social group context rather than during the intense initial aggression associated with group formation. Due to practical constraints, pigs from 81 of the 111 mixed pens (n = 1212) were observed at 18.2 \pm 2.9 (mean \pm s.d.) days after mixing, when pigs were 88.8 ± 5.2 days of age. Each pig was spray-marked enabling individual identification and then time-lapse video-recorded for 24 h. Once in an hour, instantaneous scan-samples of every pig's behaviour were made, enabling an estimate of activity levels. Pigs were defined as either being active (standing or sitting) or inactive (lying): active pigs were either standing upright on all four legs (which included standing still, walking or running), or sitting in a dog-like posture on its hind haunches with its front legs straight. Inactive pigs were lying with both front and rear portions of the ventral or lateral skin surface in contact with the ground.

Genetic analysis

The aggressive behavioural traits (fighting, bullying and being bullied) showed skewed distributions with substantial kurtosis as described in more detail in Turner *et al.* (2008) and Turner *et al.* (2009). To reduce the skewness as well as kurtosis and to approach normality, a log transformation $Y = \log_e (1 + \text{observation})$ was used for these behavioural traits. All other traits (handling scores and home pen

activity) were based on their original scale. To obtain reliable estimates of the genetic parameters, in particular of the genetic correlations, the two lines had to be pooled for the analysis. However, the lines are partly genetically connected because Yorkshire animals were used in the purebred line as well as sire in the crossbred line.

The estimation of genetic and environmental parameters was carried out in three analyses. In the first analysis, handling scores traits at mixing and activity in the home pen, were estimated together with behavioural traits indicating aggressiveness. In the second analysis, genetic and environmental parameters of handling scores at mixing and at end of test were treated as different traits and estimated in a multiple trait model. In the third analysis, handling scores at end of test were analysed with behavioural traits reflecting aggression. The model used for all analysed traits was the same and is described in more detail in the following:

$$\mathbf{y} = \mathbf{X}\mathbf{b} + \mathbf{Z}\mathbf{a} + \mathbf{W}\mathbf{c} + \mathbf{e},\tag{1}$$

where \mathbf{v} includes the vector of observations of the analysed traits. Vector **b** includes the fixed effects of line (purebred Yorkshire, crossbred Yorkshire x Landrace), sex (males, castrates and females), experimental batch (14 separate mixing days reflecting the batch farrowing policy on the farm) and body weight at mixing as a covariable fitted by using linear regression. For handling scores at the end of test, the latter covariable was based on body weight at end of test. Preliminary analysis showed that the sex effect was only significant for the behavioural trait moving into the crate recorded at end of test (indicating that females are moving faster into the crate). However, the gender effect has been kept in the model for all analysed traits to be sure that this factor did not influence the estimates of genetic parameters. The vectors **a**, **c** and **e** represent the additive genetic effects, common environmental pen effects (into which the animals were mixed) and the environmental residual effects, respectively. X, Z and W are incidence matrices linking the effects with y.

The variance-covariance structure was as follows:

$$\mathbf{V}\begin{bmatrix}\mathbf{a}\\\mathbf{c}\\\mathbf{e}\end{bmatrix} = \begin{bmatrix}\mathbf{A}\otimes\mathbf{G} & 0 & 0\\ 0 & \mathbf{I}\otimes\mathbf{M} & 0\\ 0 & 0 & \mathbf{I}\otimes\mathbf{R}\end{bmatrix},$$

where **G** represents the (co)variance matrix due to additive genetic effects, **M** is the (co)variance matrix due to the common environmental pen effects (111 pens) in which the mixing of animals took place and **R** is the (co)variance matrix due to residual effects. **A** denotes the additive genetic relationship matrix (2419 animals) and **I** is the identity matrix.

Genetic and environmental variance components of model (1) were estimated using restricted maximum likelihood (REML) approach as implemented in the programme VCE-5 (Kovac *et al.*, 2003). In all tables of results, standard errors of each estimated parameter are reported in parentheses. Estimated parameters were tested for significance from zero using P < 0.05. Some traits are estimated more than once in different analyses and in some cases their estimates were slightly different. This is because different combinations of traits result in slightly different sampling variance due to their different covariances. Where two different estimates of a trait are available, we took the most accurate estimate and quote the value with the smallest standard error if the estimate is discussed in the text.

Results

Aggressive behaviour at mixing

The heritabilities of, and correlations between aggressive traits have been discussed in detail in Turner *et al.* (2009). In summary (Tables 2 and 6), both fighting and delivery of bullying were moderate to highly heritable traits, and these two behaviours were strongly associated as indicated by a high positive genetic correlation and a moderate residual correlation. This suggests that the same animals are engaged in fighting and bullying. In contrast, receiving bullying showed a much lower heritability. The pen of mixing had a small but significant effect on aggressive behaviour, and there were strong positive correlations between these pen effects, suggesting that in certain pens, all forms of aggression were increased (Tables 3 and 7).

Scoring of the response to handling at weighing

Pigs were generally quite efficient to handle, with the majority receiving intermediate or high scores, although the scoring was not very diverse. In each case, the most commonly used (modal) score was used for 50% of more of the pigs scored (Figure 1). Vocalisations in the crate were quite rare, occurring on 6.0% of occasions during the first weighing at mixing, and 7.0% during the second weighing at end of test.

Genetic analysis comparing the corresponding handling score traits from the two weighing events (e.g. moving into the crate in the weighing at mixing and at end of test), generally revealed weak to moderate genetic correlations (Table 4). The exception to this was that vocalisations on the two occasions were highly correlated. Because of the generally weak genetic correlations, the handling scores recorded on the two occasions were treated as separate traits, rather than repeated observations of the same trait. Heritabilities for the handling traits were quite low, although all estimates were significantly different from zero. Residual correlations between different handling scores were very low. At both weighing events, there was a moderately high positive genetic correlation between the scores for moving into the crate and leaving the crate (weighing at mixing: $r_{\rm g} = 0.64 \pm 0.08$; weighing at end of test: $r_{\rm g} = 0.60 \pm 0.10$; Table 4), suggesting that the behavioural response to these two aspects of the situation

		Handling scores	for weighing at mixing		Home pen		Mixing aggression	
Trait	Moving into crate	In crate	Vocalising $(n = 0, y = 1)$	Leaving crate	Inactivity	Fighting	Being bullied	Bullying
Moving into crate	0.15 (0.02)*	-0.62 (0.08)*	0.02 (0.16)	0.64 (0.15)*	0.26 (0.15)	-0.14 (0.07)*	-0.06 (0.10)	-0.01 (0.07)
In crate	-0.07 (0.02)*	0.17 (0.03)*	-0.31 (0.20)	-0.35 (0.15)*	0.32 (0.16)*	0.18 (0.11)	-0.21 (0.13)	0.02 (0.12)
Vocalising $(n = 0, y = 1)$	-0.06 (0.02)	-0.17 (0.02)*	0.03 (0.01)*	-0.27 (0.23)	0.39 (0.20)	0.12 (0.16)	0.21 (0.28)	-0.20 (0.14)
Leaving crate	0.10 (0.02)*	-0.11 (0.02)*	-0.01 (0.02)	0.06 (0.02)*	0.20 (0.17)	0.23 (0.14)	-0.11 (0.36)	0.33 (0.17)
Inactivity	-0.03 (0.02)	0.01 (0.02)	-0.06 (0.02)	-0.04 (0.02)*	0.05 (0.02)*	0.20 (0.17)	-0.28 (0.20)	-0.09 (0.17)
Fighting	-0.10 (0.03)*	0.14 (0.04)*	0.02 (0.03)	-0.03 (0.03)	-0.01 (0.03)	0.48 (0.05)*	-0.02 (0.13)	0.87 (0.03)*
Being bullied	-0.11 (0.02)*	0.02 (0.02)	-0.04 (0.02)*	0.01 (0.04)	0.01 (0.02)	0.32 (0.03)*	0.08 (0.03)*	-0.29 (0.14)*
Bullying	-0.06 (0.02)*	0.12 (0.03)*	0.05 (0.03)	-0.07 (0.03)*	0.03 (0.03)	0.36 (0.04)*	0.21 (0.03)*	0.34 (0.05)*

5 . ת ٤ <u>ה</u>

П Table 3 Phenotypic fractions of the variance due to pen of mixing (diagonal) and their correlations (above diagonal) among different handling score and aggression traits using the handling scores for weighing at mixing

		Handling scores	for weighing at mixing		Home pen		Mixing aggression	
Trait	Moving into crate	In crate	Vocalising $(n = 0, y = 1)$	Leaving crate	Inactivity	Fighting	Being bullied	Bullying
Moving into crate In crate Vocalising $(n = 0, y = 1)$ Leaving crate	0.06 (0.01)*	-0.01 (0.13) 0.04 (0.01)*	0.51 (0.19)* -0.81 (0.13)* 0.01 (0.01)	0.14 (0.15) 0.37 (0.13)* -0.40 (0.17)* 0.05 (0.01) *	-0.42 (0.20) * -0.50 (0.22) * 0.12 (0.29) 0.26 (0.25) 0.26 (0.25) 0.26 0.25	-0.41 (0.14)* 0.22 (0.17) -0.39 (0.18)* -0.45 (0.18)*	$\begin{array}{c} 0.03 & (0.12) \\ -0.11 & (0.12) \\ -0.14 & (0.20) \\ -0.10 & (0.16) \\ 0.16 & 0.10 \\ 0.16 \\ 0.16 \\ 0.16 \\ 0.16 \end{array}$	$\begin{array}{c} 0.07 & (0.12) \\ -0.01 & (0.13) \\ -0.16 & (0.19) \\ -0.17 & (0.18) \\ 0.14 & (0.51) \end{array}$
macuvuy Fighting Being bullied Bullying						(cz.0) 10.0 0.04 (0.01)*	-0.11 (0.24) 0.46 (0.13)* 0.14 (0.02) *	(c2.0) 41.0 0.63 (0.10) * 0.96 (0.03) * 0.08 (0.02) *
	-							

Standard errors are given in parentheses. The symbol * indicates that the estimate was significantly different from zero at P < 0.05.

D'Eath, Roehe, Turner, Ison, Farish, Jack and Lawrence



Figure 1 Frequency histograms of the pig handling scores recorded during weighing at two time points (first weighing at mixing: light bars; second weighing at end of test: dark bars), for (a) moving into crate (modal score 4 used for 60.2% of pigs at mixing and 51.3% at end of test), (b) in crate (modal score 2 used for 52.9% at mixing and 63.4% at end of test), (c) leaving crate (modal score 3 was used for 73% of pigs at mixing and 91.4% at end of test).

shared a degree of common genetic influence. Therefore, at the genetic level, pigs that entered the crate with little encouragement also exited with little encouragement. The handling score for moving into the crate showed a negative genetic correlation with the score for behaviour in the crate, indicating that pigs that were difficult to move into the crate were subsequently inactive during weighing. However, this genetic association was only significant for the weighing at mixing ($r_{\rm g} = -0.57 \pm 0.04$; Table 4) but not for the end of test weighing ($r_{\rm g} = -0.17 \pm 0.10$; Table 4). The handling score in the crate showed a negative genetic correlation with the score for behaviour on leaving the crate. This correlation was similar in magnitude for both weighing events ($r_{\rm g} = -0.30 \pm 0.10$; $r_{\rm g} = -0.29 \pm 0.11$; Table 4), indicating that pigs that were inactive during

Table 4 Heritabilities	<i>(on diagonal) as well as gene</i>	tic (above diagonal)	and residual corre	lations (below dia	gonal) among diffe	erent handling scor	es assessed at mix	king and at the end	of test weighing
		Ŧ	andling scores for	weighing at mixin	ß	Han	dling scores for w	eighing at end of	est
	Trait	Moving into crate	In crate	Vocalising $(n = 0, y = 1)$	Leaving crate	Moving into crate	In crate	Vocalising $(n = 0, y = 1)$	Leaving crate
Scores at mixing	Moving into crate In crate Vocalising $(n = 0, y = 1)$	0.16 (0.02)* -0.07 (0.02)* -0.06 (0.01)*	-0.57 (0.04)* 0.20 (0.02)* -0.19 (0.01)*	-0.04 (0.10) -0.10 (0.09) 0.04 (0.01) *	0.64 (0.08)* -0.30 (0.10)* -0.21 (0.09)*	0.36 (0.05)* 0.11 (0.09) -0.04 (0.06) 0.73 (0.06)*	-0.18 (0.08)* 0.28 (0.07)* 0.69 (0.10)* -0.36 (0.13)*	-0.29 (0.09)* 0.09 (0.07) 0.76 (0.05)* -0.04 (0.13)	0.14 (0.09) -0.31 (0.11)* 0.38 (0.07)* 0.48 (0.11)*
Scores at end of test	Moving trate Moving into crate In crate Vocalising $(n = 0, y = 1)$ Leaving crate	-0.02 (0.02) -0.01 (0.02) -0.03 (0.02) 0.03 (0.02)	0.10 (0.02)* 0.10 (0.02)* 0.04 (0.02)* -0.03 (0.01)* 0.07 (0.02)	-0.06 (0.02)* -0.08 (0.02)* 0.13 (0.02)* 0.01 (0.01)	0.02 (0.02) 0.05 (0.02)* 0.04 (0.02)* -0.01 (0.01)	0.00 (0.00) 0.14 (0.01)* -0.15 (0.02)* -0.11 (0.02)* 0.08 (0.02)	(01.0) (01.0) (01.0) (01.0) *(0.0) (0.01) -0.23 (0.0) *(100)	0.10 (0.09) 0.10 (0.09) 0.24 (0.15) 0.07 (0.01) -0.05 (0.01)	0.20 (0.10)* 0.60 (0.10)* -0.29 (0.11)* 0.66 (0.09)* 0.07 (0.01)*
Standard errors are given	i in parentheses. The symbol * ii	ndicates that the estin	nate was significantly	y different from zero	e at <i>P</i> < 0.05.				

Genetics of pig temperament

D'Eath, Roehe, Turner, Ison, Farish, Jack and Lawrence

weighing were more likely to need to be pushed in order to leave when the exit gate was opened. Vocalisation in the crate showed the highest genetic correlation between the two weighing events, but generally low genetic correlations with the other handling scores. The exception to this was that pigs which had to be pushed from the crate at the mixing weighing showed a propensity to vocalise ($r_{\rm q} = -0.21 \pm$ 0.09), while conversely on the end of test weighing, it was pigs which left the crate of their own accord that showed a propensity to vocalise ($r_{\rm g} = 0.66 \pm 0.09$). All other genetic and residual correlations were low (Table 4).

Pen effects were responsible for only a small fraction of the variance for each of the handling score traits, although there were some correlations (Table 5).

Inactivity in the home pen

Inactivity in the home pen (measured by the amount of lying) was distributed symmetrically (Figure 2) and showed a low heritability ($h^2 = 0.05 \pm 0.01$; Table 6).

Genetic correlations between behavioural traits

Mixing aggression and handling scores. Estimates of genetic correlations between aggression traits and handling scores from the first weighing were not significantly different from zero (Table 2) with one exception: there was a very weak negative genetic correlation between fighting and moving into the crate, suggesting that pigs that fought needed more encouragement to enter the crate. There were a few significant correlations between the pen effects: pigs from pens in which fighting was high were slower to enter and leave the crate, and less likely to vocalise in the crate (Table 3).

There were a number of significant genetic correlations between handling scores from the end of test weighing and aggression traits (Table 6): in contrast to the finding from the weighing at mixing, genetically aggressive pigs (showing high durations of fighting and bullying) showed a genetic propensity to move more quickly into and out of the weigh crate. The receipt of bullying at mixing showed a positive genetic correlation with vocalising whilst in the crate and a negative correlation with movement into the crate during the end of test weighing, indicating that these bullied pigs required greater encouragement to enter the crate. Residual correlations between handling at the end of test weighing and aggression were low and generally nonsignificant (Table 6, below the diagonal). Pen effects on aggression and handling at the end of test weighing were generally not significantly correlated.

Handling scores and home pen inactivity. The standard error associated with estimates of genetic correlations between inactivity in the home pen and handling scores were large for both the weighing events at mixing and at end of test. Consequently, almost none of these correlations were statistically significant (Tables 2 and 6). The one exception was that there was a significant genetic correlation between inactivity and behaviour in the crate at the

		Ť	andling scores for	weighing at mixin	ğ	Han	dling scores for w	eighing at end of 1	est
	Trait	Moving into crate	In crate	Vocalising $(n = 0, y = 1)$	Leaving crate	Moving into crate	In crate	Vocalising $(n = 0, y = 1)$	Leaving crate
Scores at mixing	Moving into crate In crate	0.06 (0.01)*	-0.02 (0.09) 0.04 (0.01)*	0.42 (0.14)* -0.74 (0.14)*	0.14 (0.08) 0.46 (0.10)*	0.44 (0.07)* -0.47 (0.08)*	-0.56 (0.08)* 0.24 (0.09)*	0.08 (0.09) -0.49 (0.09)*	0.11 (0.19) -0.23 (0.19)
	Vocalising $(n = 0, y = 1)$ Leaving crate			0.02 (0.01)*	-0.32 (0.14)* 0.05 (0.01)*	0.58 (0.09)* -0.18 (0.10)	-0.11 (0.18) 0.04 (0.13)	0.04 (0.19) -0.34 (0.16)*	0.15 (0.18) 0.71 (0.15)*
Scores at end of test	Moving into crate In crate					0.06 (0.01)*	-0.28 (0.09)* 0.06 (0.01)*	0.27 (0.12)* -0.25 (0.12)*	0.02 (0.19) -0.30 (0.18)
	Vocalising $(n = 0, y = 1)$ Leaving crate							0.05 (0.01)*	-0.15 (0.28) 0.01 (0.01)





Figure 2 Frequency histogram of inactivity (frequency of lying) in the home pen 18.2 \pm 2.9 (mean \pm s.d.) days after mixing.

mixing weighing: pigs that were inactive in the home pen had higher scores in the crate indicating that they stood still during weighing.

When the pen effects were considered, there were some significant correlations between inactivity and aspects of behaviour during weighing (Tables 3 and 7).

Mixing aggression and home pen inactivity. None of the genetic correlations between inactivity and aggressive behaviour were statistically significant, as all estimates had large standard errors (Tables 2 and 6). There were also no significant correlations between pen effects on inactivity and aggressive behaviour (Tables 3 and 7).

Discussion

Genetics of aggressive behaviour at mixing

As expected based on earlier work in pigs (Løvendahl *et al.*, 2005; Turner *et al.*, 2006; Turner *et al.*, 2008), aggression had a moderate-to-high heritability, comparable to, or higher than that in reports from other species. Being bullied showed a lower heritability, suggesting that being the target of aggression is genetically determined to some degree. There was a high positive genetic correlation between the two offensive aggressive behaviours: fighting and bullying. The pen effect explained a considerable part of the variation in the environmental component of aggression. This may result from the fact that certain combinations of types of animals result in greater aggression in a pen as a whole (D'Eath, 2002). Aggressive behaviour results from this study are presented and discussed in greater detail in Turner *et al.* (2009).

Genetics of behaviour during handling at weighing

Weighing is a complex challenge for the pig, including aspects of novelty, social separation/reinstatement, confinement and the presence of actively interacting humans. Despite this, the different aspects of behaviour during weighing showed low but significant heritabilities ($h^2 = 0.03$ to 0.17), even though scores were not particularly variable, suggesting that response to handling could be altered by genetic selection. In contrast to the aggressive social behaviours, pen effects on handling were very small.

		Handling scores for	r weighing at end of test		Home pen		Mixing aggression	
Trait	Moving into crate	In crate	Vocalising $(n = 0, y = 1)$	Leaving crate	Inactivity	Fighting	Being bullied	Bullying
Moving into crate	0.16 (0.02)*	-0.24 (0.11)*	0.00 (0.12)	0.56 (0.11)*	-0.03 (0.15)	0.41 (0.05)*	-0.55 (0.09)*	0.60 (0.04)*
n crate	-0.15 (0.03)*	0.10 (0.02)*	-0.07 (0.10)	-0.25 (0.13)	-0.22 (0.16)	-0.58 (0.06)*	-0.19 (0.11)	-0.59 (0.06)*
Vocalising $(n = 0, y = 1)$	-0.11 (0.03)*	-0.22 (0.03)*	0.10 (0.02)*	0.64 (0.11)*	-0.08 (0.15)	0.12 (0.08)	0.78 (0.08)*	0.01 (0.10)
Leaving crate	0.08 (0.03)*	-0.06 (0.03)*	-0.05 (0.03)	0.07 (0.02)*	-0.13 (0.15)	0.19 (0.09)*	0.15 (0.15)	0.37 (0.09)*
nactivity	-0.01 (0.03)	0.03 (0.03)	-0.05 (0.03)	-0.06 (0.03)*	0.05 (0.01)*	0.19 (0.11)	-0.06 (0.17)	-0.17 (0.13)
Fighting	-0.05 (0.04)	0.12 (0.05)*	-0.04 (0.04)	0.01 (0.04)	-0.01 (0.04)	0.47 (0.03)*	-0.01 (0.10)	0.87 (0.04)*
Being bullied	0.05 (0.03)	-0.01 (0.03)	-0.05 (0.03)	-0.03 (0.03)	0.01 (0.03)	0.32 (0.03)*	0.12 (0.02)*	-0.20 (0.11)
Bullying	-0.10 (0.04)*	0.07 (0.04)	0.06 (0.04)	-0.07 (0.04)	0.04 (0.04)	0.36 (0.03)*	0.21 (0.04)*	0.34 (0.03)*

		Handling scores for	weighing at end of test		Home pen		Aggression traits	
Trait	Moving into crate	In crate	Vocalising $(n = 0, y = 1)$	Leaving crate	Inactivity	Fighting	Being bullied	Bullying
Moving into crate	0.07 (0.02)*	-0.22 (0.13)	0.29 (0.14)*	0.04 (0.48)	-0.51 (0.24)*	0.22 (0.14)	0.17 (0.09)	0.35 (0.11)*
In crate		0.05 (0.01)*	-0.16 (0.15)	-0.19 (0.47)	0.56 (0.22)*	-0.05 (0.15)	0.15 (0.10)	0.08 (0.13)
Vocalising $(n = 0, y = 1)$			0.04 (0.01)*	-0.42 (0.44)	0.11 (0.28)	-0.09 (0.15)	-0.17 (0.09)	-0.01 (0.12)
Leaving crate				0.01 (0.01)	-0.39 (0.40)	0.24 (0.17)	-0.25 (0.23)	-0.21 (0.24)
Inactivity					0.02 (0.01)*	0.20 (0.22)	-0.06 (0.16)	-0.04 (0.18)
Fighting						0.05 (0.01)*	0.44 (0.13)*	0.64 (0.12)*
Being bullied							0.13 (0.02)*	0.95 (0.04)*
Bullying								0.08 (0.02)*

Although there are no comparable studies in pigs, similar levels of heritability for handling scores have been found in beef cattle under similar confined circumstances (behaviour in and on exit from a crush; Burrow, 1997; Kadel *et al.*, 2006). Higher heritabilities have been reported for the reaction to humans in a standardised test in a larger space (a handling pen) in both pigs ($h^2 = 0.38 \pm 0.19$, Hemsworth *et al.*, 1990) and beef cattle ($h^2 = 0.06$ to 0.26, Phocas *et al.*, 2006).

A degree of consistency over time in a response is one aspect that defines animal temperament (Sih *et al.*, 2004; Réale *et al.*, 2007). In the present study there were moderate genetic correlations between corresponding handling scores at the two weighing events 85 days apart, suggesting that there were some common genetic influences on behaviour at these two times. However, the correlations were significantly less than one, suggesting that it was not appropriate to treat the two behavioural responses to weighing as a repeated measurement of a single trait in genetic models. Genetic correlations for repeated measures of behavioural traits have rarely been reported in livestock species (Wolf *et al.*, 2008), although repeatability is often found at the phenotypic level (Ruis *et al.*, 2000; D'Eath, 2004).

Within each weighing event, the three handling scores and vocalisations reflect different aspects of the pig's response to the same event and as such correlations between them were expected. At each weighing, moving into and out of the crate was positively genetically correlated, but the genetic correlation between these two and behaviour in the crate was negative. The direction of this relationship was perhaps due to the scoring system used: We took the human handler's perspective in assigning high scores to 'good' handling characteristics: moving quickly in and out of the crate and standing still in the crate. However, the direction of the genetic correlations we found makes better sense from a pig perspective: pigs were either active or inactive, where active pigs moved quickly into and out of the crate and moved around a lot during weighing. This interpretation requires a little caution: score 1 for moving into the crate allows for 'actively' stubborn pigs who try to escape, although in practice this occurred rarely (Figure 1a). The fact that the scores for moving in and out of the crate were positively correlated could suggest that the motivation to move away from the human handler was the common factor influencing behaviour. If fear of the crate had been the most important, the expectation might be that pigs that were slow to enter the crate would be quick to leave it.

Although rapid movement in and out of the crate is desirable from a human handler's perspective, high levels of activity in the crate and on exit may indicate that pigs were more fearful of the weighing situation. In support of this, was the finding that there was a positive genetic correlation between high-pitched 'squeal' vocalisations in the crate and a rapid exit from the crate (after the second weighing). In beef cattle, vigorous behaviour in a weigh crate and a subsequent rapid exit are often positively correlated and taken to indicate a flighty, fearful temperament, which is

D'Eath, Roehe, Turner, Ison, Farish, Jack and Lawrence

undesirable because of its implications for handler safety (Turner and Lawrence, 2007) and its association with poor meat quality (Kadel *et al.*, 2006).

Genetics of home pen inactivity

The estimated heritability for inactivity during a 24-h period in the home pen was very low (0.05 ± 0.01), but significantly different from zero. In contrast to the aggressive social behaviours, pen effects on activity were very small.

As far as we are aware, genetic parameters for activity/ inactivity over a 24-h period in the home pen have not previously been estimated in pigs. Heritability for activity during a 2-min test in an empty pen (open field) has been estimated at 0.16 (Beilharz and Cox, 1967) and at 0.28 ± 0.17 (Hemsworth *et al.*, 1990). However, activity in a 2-min openfield test is unlikely to bear much relation to activity over 24 h in the home pen. As an example, mice selected for high levels of wheel running showed no differences in a 3-min open-field test (Bronikowski *et al.*, 2001).

Estimated heritabilities for 24-h activity in other species are generally higher than that reported here. Examples include wheel running duration in mice ($h^2 = 0.14$, Swallow *et al.*, 1998), activity levels in hyperactive (attention-deficit hyperactivity disorder, ADHD) children ($h^2 = 0.35$ to 0.77, depending on the method of assessment, Wood *et al.*, 2008) and sleep durations in adult humans ($h^2 = 0.17$, Gottlieb *et al.*, 2007). This may (in part) be due to the relatively crude measure of activity (hourly scan samples) used in the present study.

Genetic correlations between traits

Inactivity in the home pen, which had a low heritability, showed no significant genetic correlation with aggression. This suggests that selection to reduce aggression would not alter the general level of activity. Genetic correlations between handling scores and inactivity were also not significant, with the exception of behaviour whilst restrained in the crate at the first weighing, suggesting that selection on either of these traits would have little effect on the other.

Handling (at the second weighing) was genetically correlated with aggression: (i) aggressive pigs (showing higher durations of fighting and bullying) were more active throughout the weighing event (see discussion above). (ii) Pigs that were the targets of bullying were reluctant to enter the crate and vocalized more in the crate. These results show genetic correlations between the behavioural response in different challenging situations: a social challenge (mixing) and a non-social challenge (handling during weighing). The extent of correlated responses across different situations is a source of much debate and interest in the literature on animal temperament (Sih et al., 2004; Koolhaas et al., 2007; Réale et al., 2007). In pigs, this question has been particularly controversial, with some authors finding an association between social and nonsocial challenges (Thodberg et al., 1999; Ruis et al., 2000; Bolhuis et al., 2005) and others not (D'Eath and Burn, 2002; Janczak et al., 2003). However, in contrast to the rodent

literature, which makes extensive use of high and low aggression selection lines (Miczek *et al.*, 2001; Gammie *et al.*, 2006), this research has typically been conducted at the phenotypic level. The present study is the first to identify genetic associations between the behavioural response to social and non-social challenges in pigs.

The practical implications of this finding is that selection for reduced aggression in pigs might result in some degree of correlated reduction in activity during physical restraint (at least in older animals), but an increase in the labour input required to move animals when isolated from their pen mates. There may also be animal welfare implications: reduced activity during weighing itself could reflect a reduction in the level of fear. In beef cattle, rapid exit from a weigh crate or crush is assumed to reflect greater fear of the situation, and a more nervous, flighty temperament (Turner and Lawrence, 2007). If selection for reduced aggression in pigs did result in less fearful animals because of a correlated selection response, this could have a positive effect on animal welfare and also perhaps on meat guality (Kadel et al., 2006). We should perhaps be a little cautious though, because pigs that are reluctant to move during weighing might also be experiencing fear. A further study with more detailed behavioural (and perhaps physiological) assessment of pigs during weighing would be needed to investigate this before deciding whether selection for reduced activity during weighing could improve pig welfare.

Conclusions

This study has shown that general activity in the home pen and ease of handling at weighing in pigs are heritable traits. Estimates of heritability were low but significantly different from zero. There were moderate genetic correlations between aggression when unfamiliar pigs were mixed and handling scores at weighing: aggressive pigs required less encouragement during movement to and from a weigh crate, but were more active when restrained in the weigh crate. Genetic associations between behavioural responses to different challenging situations are consistent with the concept of temperament (behavioural syndromes, coping styles or personality). Selection for reduced aggression in pigs is feasible and desirable, but this study shows that other behaviours could show a correlated response to some degree, with possible implications for animal production and welfare.

Acknowledgements

This work was supported by the Scottish Government Rural and Environmental Research and Analysis Directorate. It is part of the Welfare Quality[®] research project, co-financed by the European Commission, within the 6th Framework Programme, contract No. FOOD-CT-2004-506508. The assistance of Nils Lundeheim and Lotta Rydhmer at SLU Uppsala, Maria Kihlberg at Quality Genetics, and Jan Vallgårda and the staff of Vallrums Lantbruk is gratefully appreciated. Ulla Schmidt, Bjorne Olsson and Elisabeth Merin performed on-farm data collection.

D'Eath, Roehe, Turner, Ison, Farish, Jack and Lawrence

References

Beilharz RG and Cox DF 1967. Genetic analysis of open field behavior in swine. Journal of Animal Science 26, 988–990.

Boissy A, Fisher AD, Bouix J, Hinch GN and Le Neindre P 2005. Genetics of fear in ruminant livestock. Livestock Production Science 93, 23–32.

Bolhuis JE, Schouten WGP, Schrama JW and Wiegant VA 2005. Individual coping characteristics, aggressiveness and fighting strategies in pigs. Animal Behaviour 69, 1085–1091.

Bronikowski AM, Carter PA, Swallow JG, Girard IA, Rhodes JS and Garland T 2001. Open-field behavior of house mice selectively bred for high voluntary wheel-running. Behavior Genetics 31, 309–316.

Burrow HM 1997. Measurements of temperament and their relationships with performance traits in beef cattle. Animal Breeding Abstracts 65, 477–495.

D'Eath RB 2002. Individual aggressiveness measured in a resident-intruder test predicts the persistence of aggressive behaviour and weight gain of young pigs after mixing. Applied Animal Behaviour Science 77, 267–283.

D'Eath RB 2004. Consistency of aggressive temperament in domestic pigs: the effects of social experience and social disruption. Aggressive Behavior 30, 435–448.

D'Eath RB 2005. Socialising piglets before weaning improves social hierarchy formation when pigs are mixed post-weaning. Applied Animal Behaviour Science 93, 199–211.

D'Eath RB and Burn CC 2002. Individual differences in behaviour: a test of 'coping style' does not predict resident-intruder aggressiveness in pigs. Behaviour 139, 1175–1194.

D'Eath RB and Lawrence AB 2004. Early life predictors of the development of aggressive behaviour in the domestic pig. Animal Behaviour 67, 501–509.

Falconer DS and Mackay TFC 1996. Introduction to quantitative genetics, 4th edition. Longman, New York.

Gammie SC, Garland T and Stevenson SA 2006. Artificial selection for increased maternal defense behavior in mice. Behavior Genetics 36, 713–722.

Gottlieb DJ, O'Connor GT and Wilk JB 2007. Genome-wide association of sleep and circadian phenotypes. BMC Medical Genetics 8 (suppl. 1), S9.

Hemsworth PH, Barnett JL, Treacy D and Madgwick P 1990. The heritability of the trait fear of humans and the association between this trait and subsequent reproductive- performance of gilts. Applied Animal Behaviour Science 25, 85–95.

Janczak AM, Pedersen LJ and Bakken M 2003. Aggression, fearfulness and coping styles in female pigs. Applied Animal Behaviour Science 81, 13–28.

Jarvis S, D'Eath RB and Fujita K 2005. Consistency of piglet crushing by sows. Animal Welfare 14, 43–51.

Kadel MJ, Johnston DJ, Burrow HM, Graser HU and Ferguson DM 2006. Genetics of flight time and other measures of temperament and their value as selection criteria for improving meat quality traits in tropically adapted breeds of beef cattle. Australian Journal of Agricultural Research 57, 1029–1035.

Kanis E, van den Belt H, Groen AF, Schakel J and De Greef KH 2004. Breeding for improved welfare in pigs: a conceptual framework and its use in practice. Animal Science 78, 315–329.

Koolhaas JM, de Boer SF, Buwalda B and van Reenen K 2007. Individual variation in coping with stress: a multidimensional approach of ultimate and proximate mechanisms. Brain Behavior and Evolution 70, 218–226.

Kovac M, Groeneveld E and Garcia-Cortez A 2003. VCE-5 User's Guide and Reference Manual Version 5.1.2. Institute of Animal Science and Animal Husbandry, Federal Agricultural Research Centre (Bundesforschungsanstalt für Landwirtscharft, FAL), Mariensee/Neustadt, Germany.

Løvendahl P, Damgaard LH, Nielsen BL, Thodberg K, Su GS and Rydhmer L 2005. Aggressive behaviour of sows at mixing and maternal behaviour are

heritable and genetically correlated traits. Livestock Production Science 93, 73–85.

Marchant JN, Whittaker X and Broom DM 2001. Vocalisations of the adult female domestic pig during a standard human approach test and their relationships with behavioural and heart rate measures. Applied Animal Behaviour Science 72, 23–39.

Miczek KA, Maxson SC, Fish EW and Faccidomo S 2001. Aggressive behavioral phenotypes in mice. Behavioural Brain Research 125, 167–181.

Phocas F, Boivin X, Sapa J, Trillat G, Boissy A and Le Neindre P 2006. Genetic correlations between temperament and breeding traits in Limousin heifers. Animal Science 82, 805–811.

Réale D, Reader SM, Sol D, McDougall PT and Dingemanse NJ 2007. Integrating animal temperament within ecology and evolution. Biological Reviews 82, 291–318.

Roehe R, Shrestha NP, Mekkawy W, Baxter EM, Knap PW, Smurthwaite KM, Jarvis S, Lawrence AB and Edwards SA 2009. Genetic analyses of piglet survival and individual birth weight on first generation data of a selection experiment for piglet survival under outdoor conditions. Livestock Science 121, 173–181.

Ruis MAW, te Brake JHA, Van de Burgwal JA, de Jong IC, Blokhuis HJ and Koolhaas JM 2000. Personalities in female domesticated pigs: behavioural and physiological indications. Applied Animal Behaviour Science 66, 31–47.

Sih A, Bell A and Johnson JC 2004. Behavioral syndromes: an ecological and evolutionary overview. Trends in Ecology and Evolution 19, 372–378.

Swallow JG, Carter PA and Garland T 1998. Artificial selection for increased wheel-running behavior in house mice. Behavior Genetics 28, 227–237.

Thodberg K, Jensen KH and Herskin MS 1999. A general reaction pattern across situations in prepubertal gilts. Applied Animal Behaviour Science 63, 103–119.

Turner SP and Lawrence AB 2007. Relationship between maternal defensive aggression, fear of handling and other maternal care traits in beef cows. Livestock Science 106, 182–188.

Turner SP, Roehe R, Mekkawy W, Farnworth MJ, Knap PW and Lawrence AB 2008. Bayesian analysis of genetic associations of skin lesions and behavioural traits to identify genetic components of individual aggressiveness in pigs. Behavior Genetics 38, 67–75.

Turner SP, Roehe R, D'Eath RB, Ison SH, Farish M, Jack MC, Lundeheim N, Rydhmer L and Lawrence AB 2009. Genetic validation of skin injuries in pigs as an indicator of post-mixing aggressiveness and the relationship with aggression under stable social conditions. Journal of Animal Science (in press).

Turner SP, White IMS, Brotherstone S, Farnworth MJ, Knap PW, Penny P, Mendl M and Lawrence AB 2006. Heritability of post-mixing aggressiveness in grower-stage pigs and the relationship between aggressiveness and production traits. Animal Science 82, 615–620.

Van Oers K, de Jong G, Van Noordwijk AJ, Kempenaers B and Drent PJ 2005. Contribution of genetics to the study of animal personalities: a review of case studies. Behaviour 142, 1185–1206.

Warriss PD, Brown SN, Gade PB, Santos C, Costa LN, Lambooij E and Geers R 1998. An analysis of data relating to pig carcass quality and indices of stress collected in the European Union. Meat Science 49, 137–144.

Wolf BT, McBride SD, Lewis RM, Davies MH and Haresign W 2008. Estimates of the genetic parameters and repeatability of behavioural traits of sheep in an arena test. Applied Animal Behaviour Science 112, 68–80.

Wood AC, Rijsdijk F, Saudino KJ, Asherson P and Kuntsi J 2008. High heritability for a composite index of children's activity level measures. Behavior Genetics 38, 266–276.