



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Measuring anti-Müllerian hormone for the assessment of ovarian reserve

When and for whom is it indicated?

Citation for published version:

Anderson, RA, Nelson, SM & Wallace, WHB 2012, 'Measuring anti-Müllerian hormone for the assessment of ovarian reserve: When and for whom is it indicated?', *Maturitas: An international journal of midlife health and beyond*, vol. 71, no. 1, pp. 28-33. <https://doi.org/10.1016/j.maturitas.2011.11.008>

Digital Object Identifier (DOI):

[10.1016/j.maturitas.2011.11.008](https://doi.org/10.1016/j.maturitas.2011.11.008)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Maturitas: An international journal of midlife health and beyond

Publisher Rights Statement:

© 2011 Elsevier Ireland Ltd

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Review

Measuring anti-Müllerian hormone for the assessment of ovarian reserve: When and for whom is it indicated?

R.A. Anderson^{a,*}, S.M. Nelson^{b,1}, W.H.B. Wallace^{c,2}

^a Division of Reproductive and Developmental Sciences, University of Edinburgh, Edinburgh, Scotland, UK

^b Centre for Population and Health Sciences, University of Glasgow, Glasgow, Scotland, UK

^c Division of Child Life and Health, Department of Reproductive and Developmental Sciences, University of Edinburgh, Scotland, UK

ARTICLE INFO

Article history:

Received 4 November 2011

Accepted 7 November 2011

Keywords:

Ovary
Ovarian reserve
Anti-Müllerian hormone
Menopause
IVF

ABSTRACT

Our understanding of female reproductive function has been hampered by our inability to directly assess the number of non-growing primordial follicles present in the ovary, the ovarian reserve. Female reproductive hormones (FSH and LH, the inhibins and steroids) reflect the activity of the larger growing follicles and thus are largely informative of peri-ovulatory ovarian activity. In contrast anti-Müllerian hormone (AMH) is a product of the granulosa cells of small growing follicles, whose number (and therefore circulating AMH concentrations) is reflective of the ovarian reserve. AMH declines with age in adult women, and emerging data suggest a relationship with remaining reproductive lifespan and age at the menopause. Early studies demonstrated that AMH concentrations are stable across the menstrual cycle, adding to its clinical utility. The most established role for AMH measurement is in women about to start IVF treatment, where it is predictive of the ovarian response and is of clear value in identifying women at risk of ovarian hyperstimulation syndrome or whose response will be poor and thus their expectations can be tailored. AMH is detectable in childhood, and although relationships to puberty are not yet available, it appears that AMH rises to a peak in the early 20s. Developing indications include in assessment and individualisation of the risk to fertility from chemotherapy, in the diagnosis of PCOS and as a tumour marker in granulosa cell tumours. The increasingly routine use of AMH by IVF clinics heralds much wider adoption in a range of clinical situations across the reproductive lifespan.

© 2011 Elsevier Ireland Ltd. All rights reserved.

Contents

1. Introduction	29
1.1. Assessment of ovarian reserve before assisted conception	29
1.2. Age-specific interpretation of AMH as a surrogate measure of ovarian reserve	29
1.3. Assessing and predicting gonadal damage	30
1.4. Diagnosis in oligo/amenorrhoea	31
1.5. Prediction of reproductive lifespan	31
2. Conclusion	31
Contributors	31
Competing interests	32
Provenance and peer review	32
Acknowledgement	32
References	32

* Corresponding author at: MRC Centre for Reproductive Health, The Queen's Medical Research Institute, University of Edinburgh, 49 Little France Crescent, Edinburgh EH16 4TJ, UK. Tel.: +44 131 2426386.

E-mail addresses: richard.anderson@ed.ac.uk (R.A. Anderson), Scott.Nelson@glasgow.ac.uk (S.M. Nelson), hamish.wallace@nhs.net (W.H.B. Wallace).

¹ Address: Muirhead Chair in Obstetrics & Gynaecology, Centre for Population and Health Sciences, University of Glasgow, Level 2 McGregor Building, Western Infirmary, Glasgow G11 6NT, UK. Tel.: +44 141 211 2329.

² Address: Department of Haematology/Oncology, Royal Hospital for Sick Children, 17 Millerfield Place, Edinburgh EH9 1LW, UK. Tel.: +44 131 536 0420.

1. Introduction

Our current understanding of female reproductive function presumes that the ovary contains a finite number of oocytes within primordial follicles, and that their depletion heralds the menopause [1]. This pool of primordial follicles is formed during fetal life from approximately 18 weeks gestation, following oogonial proliferation, entry and arrest in meiosis, and interaction with somatic cells. A small number of this resting pool of primordial follicles is activated into growth every day throughout the reproductive lifespan of a woman (including in childhood) with the vast majority destined to undergo atresia [2]. Indeed before puberty all growing follicles will become atretic, and after puberty only one a month may escape this fate and progress to ovulation. The pool of resting follicles is the true ovarian reserve, but the term ‘ovarian reserve’ is also widely used in the assisted conception literature to mean the number of growing follicles that can be recruited by exogenous FSH to grow to a pre-ovulatory stage, resulting in the potential collection of an oocyte for assisted reproduction. This growing pool is perhaps better termed the functional ovarian reserve. Although the true and functional ovarian reserves reflect different stages of follicular development, they are inherently linked and both decline in parallel with increasing age. Accurate measurement of the ovarian reserve has long been a quest in reproductive medicine and recent years have seen a dramatic increase in research in this field, a large part of which has been fuelled by the recognition that measurement of anti-Müllerian hormone (AMH) in serum is a much more accurate measure of the ovarian reserve than the other hormones that have previously been available to us. Thus while inhibin B shows good prediction of oocyte yield after superovulation [3], it requires measurement in the early follicular phase of the menstrual cycle and only decreases late in the reproductive years [4]. It has therefore not supplanted FSH as the most widely used marker of the ovarian reserve despite the latter hormone’s well-recognised limitations.

AMH was first identified in the 1940s on the basis of its production by the fetal Sertoli cell, resulting in loss of the Müllerian duct system in the male. Production of AMH by granulosa cells in the adult ovary was first reported 30 years ago, in the chicken [5], but research into its function and clinical utility only gathered pace at the turn of the century with the development of commercially available assays. Early studies demonstrated that serum AMH showed a close correlation with the number of oocytes that were obtained following superovulation for IVF [6], and that it declined with age in women [7,8]. These two findings are central to our current understanding of the utility of AMH as a measure of the ovarian reserve.

In contrast to the large amount of clinical research on the utility of measuring AMH, there are remarkably few data regarding its biological functions. Early studies using an AMH knock-out mouse demonstrated that perhaps its most important role is the regulation of initiation of early follicle growth [9,10], and subsequent studies have indicated that it may regulate the responsiveness of growing follicles to FSH [11]. AMH production by cumulus cells is stimulated by the oocyte [12] consistent with additional functions in mediating oocyte/cumulus maturation. In keeping with these roles, AMH expression by granulosa cells of the follicle is initiated at the onset of follicle growth, but equally importantly expression then declines rapidly with little expression in antral follicles beyond 6–8 mm diameter in the human [13]. The lack of production of AMH by the mural granulosa of larger follicles in the lead up to ovulation (or by the corpus luteum) is critical for its clinical utility as it means that its serum concentration varies little over the menstrual cycle [14,15]. Thus for practical purposes it can be assayed at any stage of the cycle.

Recognition of these unique characteristics has led to the rapid adoption of AMH for a range of indications. In this review we discuss the potential uses and recent developments for AMH, and summarise the evidence supporting its adoption into clinical practice.

1.1. Assessment of ovarian reserve before assisted conception

This is the most established indication for measuring AMH, with many clinics now routinely assaying AMH before IVF treatment. As AMH correlates strongly with the number of oocytes obtained following superovulation, a baseline measurement allows individualisation of both the treatment strategy and the patient’s expectations [16,17]. Specifically women with a high AMH are likely to respond excessively to exogenous gonadotrophins and their treatment strategy can be modified to minimise the risk of ovarian hyperstimulation syndrome (OHSS). Conversely women with a low AMH are likely to respond poorly to stimulation with consequently a low chance of pregnancy, and their expectations can be managed appropriately with alternatives like oocyte donation discussed. Although AMH is primarily a measure of oocyte quantity rather than quality, it is positively associated with IVF live birth rates independently of age [16,18,19]. This is primarily due to its relationship with oocyte yield, as at any given age range women with a higher AMH are likely to have more oocytes retrieved and thereby potentially more embryos for selection.

The cost-effectiveness of the use of an AMH-based treatment strategy in IVF has recently been assessed, and proposed to lead to substantial savings [20]. Some of this is through the more accurate identification of women at risk of OHSS who can then be treated with a less intense stimulation regimen, with significant cost savings from fewer hospital admissions for OHSS. A less patient-centred aspect of this analysis was however the exclusion of women with a very low AMH from state funded IVF, which will inevitably result in exclusion of some women with a low-but not zero-chance of success as there is no AMH concentration, even when it is undetectable, that is associated with absolutely no chance of pregnancy.

1.2. Age-specific interpretation of AMH as a surrogate measure of ovarian reserve

The production of AMH by the growing follicle pool means that its concentration varies across childhood, adolescence and adult life [7,21]. Initial adult cohorts clearly demonstrated an age related decline reaching undetectable levels in advance of the menopause [22]. Subsequently AMH nomograms have been reported and validated from very large cohorts of women [23–25]. While these data derive from women attending infertility clinics, subset analysis of women where there was a severe male contribution to the infertility (and where female pathology is therefore less likely) showed no difference from the whole cohort. Longitudinal data are lacking, but the size of the cross-sectional cohorts now reported make it likely that they will indeed reflect longitudinal changes at a population level.

More recently AMH concentrations in populations of children and younger adults have been described, and a compilation of these data now allows a description of AMH concentration from birth to the menopause (Fig. 1) [26]. This confirms a transient neonatal rise in AMH similar to the well-described mini-puberty of the male neonate, with a subsequent rise through childhood and adolescence to a peak in the early 20s. Concentrations then decline from the late 20s. Collectively these data indicate that AMH shows no dramatic change at puberty but shows an increase from late childhood through to early adulthood. These data are derived from cross-sectional analyses and prospective multiple sampling through puberty is required to test and refine this model. The




Fig. 1. Serum AMH from conception to the menopause (note log scale). The red line is the peak model that best fits the 3260 datapoints shown as triangles. The coefficient of determination, r^2 , is 0.34, indicating that 34% of variation in serum AMH levels is due to age alone. Peak serum AMH is at 24.5 years.

Reproduced from doi:10.1371/journal.pone.0022024.

continuing rise through later adolescence suggests that there are ongoing changes in patterns of follicle maturation over perhaps a decade following menarche.

These large studies also confirm the wide variability in AMH between women. This is in fact to be expected, as there is a comparably wide range in primordial and non-growing follicle number [1,27], and in age at menopause. The age-related changes in AMH have also been modelled relative to the non-growing follicle (NGF) pool [28]. During childhood the NGF shows an inverse relationship with AMH (i.e. the former is declining while the latter is rising), but both AMH and the rate of loss of the NGF pool (assumed to reflect largely initiation of follicle growth) rise in parallel. After a peak at age 24, both AMH and NGF activation rate both start to fall, mirroring the fall in non-growing follicle numbers.

1.3. Assessing and predicting gonadal damage

Our ability to interpret AMH in an age-specific manner has allowed the development of new applications. These include accurate assessment of the impact of gonadotoxic therapy, e.g. chemotherapy and radiotherapy, or of ovarian surgery, on the ovarian reserve [29,30]. AMH falls very rapidly with the onset of chemotherapy [31,32]. In lymphoma analysis allowed the clear demonstration of the toxicity of alkylating agent based therapy, as those women showed little or no recovery of AMH following chemotherapy, whereas women receiving non-alkylating agent therapy showed a good recovery to approximately pre-treatment concentrations [32]. AMH is also useful in prepubertal girls, showing a fall during chemotherapy with recovery dependant on the toxicity of the regimen used [33], while other reproductive hormones are not of value in this context. Post-treatment follow-up studies have also shown that AMH concentrations are reduced in women following treatment for cancer, either in childhood or adulthood, with low AMH concentrations most consistently seen

following total body radiation [31,34–37]. Normal concentrations have also been reported in populations of women following certain chemotherapy regimens indicating that those particular therapies do not cause loss of the ovarian reserve that is detectable using this marker [37,38]. This essentially reassuring result in some women requires long term follow up with both fertility and age of menopause as endpoints, and at present data with that degree of completeness are lacking. The ability of the reproductive system to compensate for rapid loss of follicle number is indicated by the only modest change in reproductive lifespan following unilateral oophorectomy [39]. If translatable to the post-chemotherapy situation, this might indicate that only very substantial loss of the ovarian reserve will have a clinically relevant impact. As AMH reflects the number of growing follicles, small changes in the rate of activation may well not be detectable.

Comparable results have been reported following surgery for ovarian endometriosis [30], with a fall in AMH immediately following surgery, and recovery thereafter. Assuming that primordial follicles cannot be replaced and that surgery (and chemotherapy) removes both growing and non-growing follicles, this can only indicate that a smaller non-growing pool re-establishes a similar growing pool to that present pretreatment, thus normalising AMH. It would appear, however, reasonable to reassure a woman regarding future fertility prospects if her AMH is normal following recovery from cancer or other treatments although there is a lack of prospective data to substantiate this fully at present.

A further consideration is whether a higher pre-treatment AMH before chemotherapy makes it more likely that a woman will retain ovarian function following treatment, whereas a woman of the same age but with a lower AMH is more at risk of experiencing a premature menopause. In support of this initial studies demonstrated that pre-treatment AMH was higher in women treated for breast cancer who retained menses 1 year after chemotherapy

[40], and with higher post-treatment AMH concentrations [41], although others have found no difference [42]. A recent analysis of menstrual and ovarian function 5 years after chemotherapy for breast cancer found that pretreatment AMH was indeed a predictor of ongoing menses (with associated biochemical evidence of increased ovarian activity) at that time [43]. Age was also a predictor, as expected, but strikingly in a multivariate analysis only AMH remained predictive, with the effect of age no longer significant. This is consistent with age being a surrogate marker of the ovarian reserve whose value becomes redundant when a direct marker of adequate accuracy is included. This finding has since been replicated in a preliminary report of a further small cohort [44], but should also be replicated in younger cohorts of women with other conditions, e.g. lymphoma, before this test can be recommended for clinical practice. Furthermore it should be recognised that these studies used ovarian function rather than fertility as an outcome and while this is relevant for some long-term outcomes such as osteoporosis and other risks of early ovarian insufficiency, for many women the key question will be whether they will retain their fertility after chemotherapy. Confirmation that AMH provides individualisation of risk can rapidly translate into clinical practice, with more invasive or time-consuming methods of fertility preservation being appropriate for women with a low AMH whereas others with a high AMH may elect to start treatment without delay.

1.4. *Diagnosis in oligo/amenorrhea*

Although we have classically measured gonadotrophins and oestradiol to subclassify oligoamenorrhoea, a single AMH measurement may be a better first line investigation. In oligoamenorrhoeic women with Polycystic Ovarian Syndrome (PCOS), AMH is often dramatically high due to AMH being produced by the large number of antral follicles [45,46]. Conversely women with amenorrhoea due to premature ovarian insufficiency have a low AMH, while women with amenorrhea related to hyperprolactinemia or in hypogonadotrophic hypogonadism often have normal circulating concentrations of AMH [47]. AMH is also very high in women with granulosa cell tumours, where it is of value as a tumour marker [48].

Women with hyperprolactinemia or hypogonadotrophic hypogonadism will have low FSH concentrations and this might therefore be expected to reduce AMH production from the growing follicle pool. However the degree of gonadotropin suppression seems to be largely insufficient to influence AMH production in these two groups of women. Likewise AMH production is reported to be unaffected in women taking the combined contraceptive pill who also have significant gonadotropin suppression [49], although a rise in AMH following discontinuation of the contraceptive pill has also been reported [50]. It appears that more prolonged and complete gonadotropin suppression can cause more readily detectable suppression of AMH, as seen in women on long-term goserelin treatment for breast cancer [31], and this may also underpin the decline in AMH concentrations during pregnancy [51]. Clinical experience indicates that some women with hypogonadotrophic hypogonadism do have a low AMH, and further well-defined cohorts of women with this condition (including follow up to recovery of normal ovarian function) are needed to delineate in detail possible changes in AMH. AMH is thus far not part of the diagnostic work up for PCOS but the criteria for making this diagnosis remains subject to revision and it may well be that AMH becomes part of this in due course [52]. Certainly the markedly elevated AMH concentrations contrast to the frequently normal values of other reproductive hormones in PCOS, thus raising the possibility of a clear biochemical aspect to the diagnosis, which can also only aid further our understanding of this condition.

1.5. *Prediction of reproductive lifespan*

The decline in AMH during the later reproductive years is one of the most established findings, in both prospective and cross-sectional studies, thus raising the possibility of prediction of the menopause [22] and by implication, a woman's remaining reproductive lifespan. In analysis of serial blood samples taken from 50 women followed prospectively from the age of 42 it was demonstrated that AMH declined to undetectable levels some 5 years before the final menstrual period (FMP) [53]. Inhibin B declined similarly becoming undetectable some 4 years before the FMP. AMH was significantly related both to time to and age at FMP, whereas inhibin B was less predictive of both. A recent larger Dutch study has provided further evidence that AMH could provide long-term prediction of age at menopause [54]. The study included 257 ovulatory women aged 21–46, who were reassessed after 11 years. Analysis showed that a low age-specific AMH at initial testing was associated with a shift in the distribution of the menopause to a younger age, whereas a higher age-specific AMH shifted the distribution toward a higher age. Importantly in this study the use of AMH was in the context of age, rather than independent of it. Even in this fairly large study however only some 11% of the women studied had become menopausal during the follow up period and thus larger and longer duration studies will be required to provide accurate estimates of the predictive power of AMH and the variability associated with this. It must be recognised that age at menopause is very strongly genetically determined and although a large number of environmental influences have been identified, their combined effect is modest [55]. It will be interesting in future work to assess whether AMH is a better predictor than knowing the mothers' age at menopause.

2. Conclusion

The last decade has provided a wealth of data confirming the value of AMH as a measure of the ovarian reserve. The clearest data for its clinical utility is in the context of IVF and measurement has been routine practice in some clinics now for several years. The validity of the evidence base for other indications is weaker but rapidly growing. Much of the value of AMH lies in its relation to the declining ovarian pool with age, and thus its potential ability to predict future reproductive outcomes. We must again emphasise how little longitudinal or prospective data there are in this field and therefore its clinical use in advising individual women outside the context of IVF is not well supported. The use of AMH in women with a new cancer diagnosis is promising as it may allow greater individualisation of risk than is currently available, and its value in the differential diagnosis of oligoamenorrhoea seems clear. It is inevitable that AMH will be increasingly measured by women wishing to know their future reproductive lifespan, as societal changes will continue to result in women increasingly delaying childbearing until their later reproductive years. This is likely to go hand in hand with the more widespread use of fertility preservation technologies such as oocyte vitrification for non-medical reasons despite the lack of a clear evidence base. While bearing these caveats in mind it is clear that the increasing use of AMH will be of substantial benefit widely across reproductive medicine.

Contributors

Professor Richard A. Anderson contributed in writing of ms, approval of final version. Professor W. Hamish B. Wallace contributed in writing of ms, approval of final version. Professor Scott M. Nelson contributed in writing of ms, approval of final version.

Competing interests

R.A. Anderson and S.M. Nelson have acted as consultants for Beckman Coulter and Roche Diagnostics.

Provenance and peer review

Commissioned and externally peer reviewed.

Acknowledgement

The author's studies were funded by the Medical Research Council (WBS: U.1276.00.002.00001 to RAA).

References

- Wallace WH, Kelsey TW. Human ovarian reserve from conception to the menopause. *PLoS ONE* 2010;5:e8772.
- McGee EA, Hsueh AJ. Initial and cyclic recruitment of ovarian follicles. *Endocr Rev* 2000;21:200–14.
- Yong PTK, Baird DT, Thong KJ, et al. Prospective analysis of the relationships between the ovarian follicle cohort and basal FSH concentration, the inhibin response to exogenous FSH and ovarian follicle number at different stages of the normal menstrual cycle and after pituitary down-regulation. *Hum Reprod* 2003;18:35–44.
- Welt CK, McNicholl DJ, Taylor AE, et al. Female reproductive aging is marked by decreased secretion of dimeric inhibin. *J Clin Endocrinol Metab* 1999;84:105–12.
- Hutson J, Ikawa H, Donahoe PK. The ontogeny of Müllerian inhibiting substance in the gonads of the chicken. *J Pediatr Surg* 1981;16:822–7.
- Seifer DB, MacLaughlin DT, Christian BP, et al. Early follicular serum Müllerian-inhibiting substance levels are associated with ovarian response during assisted reproductive technology cycles. *Fertil Steril* 2002;77:468–71.
- de Vet A, Laven JSE, de Jong FH, et al. Antimüllerian hormone serum levels: a putative marker for ovarian aging. *Fertil Steril* 2002;77:357–62.
- van Rooij IAJ, Broekmans FJM, Scheffer GJ, et al. Serum antimüllerian hormone levels best reflect the reproductive decline with age in normal women with proven fertility: a longitudinal study. *Fertil Steril* 2005;83:979–87.
- Durlinger AL, Kramer P, Karels B, et al. Control of primordial follicle recruitment by anti-Müllerian hormone in the mouse ovary. *Endocrinology* 1999;140:5789–96.
- Gigli I, Cushman RA, Wahl CM, et al. Evidence for a role for anti-Müllerian hormone in the suppression of follicle activation in mouse ovaries and bovine ovarian cortex grafted beneath the chick chorioallantoic membrane. *Mol Reprod Dev* 2005;71:480–8.
- Durlinger AL, Gruijters MJ, Kramer P, et al. Anti-Müllerian hormone attenuates the effects of FSH on follicle development in the mouse ovary. *Endocrinology* 2001;142:4891–9.
- Salmon NA, Handyside AH, Joyce IM. Oocyte regulation of anti-Müllerian hormone expression in granulosa cells during ovarian follicle development in mice. *Dev Biol* 2004;266:201–8.
- Weenen C, Laven JS, Von Bergh AR, et al. Anti-Müllerian hormone expression pattern in the human ovary: potential implications for initial and cyclic follicle recruitment. *Mol Hum Reprod* 2004;10:77–83.
- La Marca A, Malmusi S, Giulini S, et al. Anti-Müllerian hormone plasma levels in spontaneous menstrual cycle and during treatment with FSH to induce ovulation. *Hum Reprod* 2004;19:2738–41.
- van Disseldorp J, Lambalk CB, Kwee J, et al. Comparison of inter- and intra-cycle variability of anti-Müllerian hormone and antral follicle counts. *Hum Reprod* 2010;25:221–7.
- Nelson SM, Yates RW, Fleming R. Serum anti-Müllerian hormone and FSH: prediction of live birth and extremes of response in stimulated cycles—implications for individualization of therapy. *Hum Reprod* 2007;22:2414–21.
- Nelson SM, Yates RW, Lyall H, et al. Anti-Müllerian hormone-based approach to controlled ovarian stimulation for assisted conception. *Hum Reprod* 2009;24:867–75.
- Majumder K, Gelbaya TA, Laing I, et al. The use of anti-Müllerian hormone and antral follicle count to predict the potential of oocytes and embryos. *Eur J Obstet Gynecol Reprod Biol* 2010;150:166–70.
- La Marca A, Nelson SM, Sighinolfi G, et al. Anti-Müllerian hormone-based prediction model for a live birth in assisted reproduction. *Reprod Biomed Online* 2011;22:341–9.
- Yates AP, Rustamov O, Roberts SA, et al. Anti-Müllerian hormone-tailored stimulation protocols improve outcomes whilst reducing adverse effects and costs of IVF. *Hum Reprod* 2011;26:2353–62.
- Hagen CP, Aksglaede L, Sorensen K, et al. Serum levels of anti-Müllerian hormone as a marker of ovarian function in 926 healthy females from birth to adulthood and in 172 Turner syndrome patients. *J Clin Endocrinol Metab* 2010;93:1356–8.
- van Rooij IA, Tonkelaar I, Broekmans FJ, et al. Anti-Müllerian hormone is a promising predictor for the occurrence of the menopausal transition. *Menopause* 2004;11:601–6.
- Nelson SM, Messow MC, Wallace AM, et al. Nomogram for the decline in serum antimüllerian hormone: a population study of 9,601 infertility patients. *Fertil Steril* 2011;95:736–41.
- Nelson SM, Messow MC, McConnachie A, et al. External validation of nomogram for the decline in serum anti-Müllerian hormone in women: a population study of 15,834 infertility patients. *Reprod Biomed Online* 2011;23:204–6.
- Seifer DB, Baker VL, Leader B. Age-specific serum anti-Müllerian hormone values for 17,120 women presenting to fertility centers within the United States. *Fertil Steril* 2011;95:747–50.
- Kelsey TW, Wright P, Nelson SM, et al. A validated model of serum anti-Müllerian hormone from conception to menopause. *PLoS ONE* 2011;6:e22024.
- Faddy MJ, Gosden RG. A model conforming the decline in follicle numbers to the age of menopause in women. *Hum Reprod* 1996;11:1484–6.
- Kelsey TW, Anderson RA, Wright P, et al. Data-driven assessment of the human ovarian reserve. *Mol Hum Reprod* 2011.
- Meirow D, Biederman H, Anderson RA, et al. Toxicity of chemotherapy and radiation on female reproduction. *Clin Obstet Gynecol* 2010;53:727–39.
- Chang HJ, Han SH, Lee JR, et al. Impact of laparoscopic cystectomy on ovarian reserve: serial changes of serum anti-Müllerian hormone levels. *Fertil Steril* 2010;94:343–9.
- Anderson RA, Themmen APN, Al Qahtani A, et al. The effects of chemotherapy and long-term gonadotrophin suppression on the ovarian reserve in premenopausal women with breast cancer. *Hum Reprod* 2006;21:2583–92.
- Decanter C, Morschhauser F, Pigny P, et al. Anti-Müllerian hormone follow-up in young women treated by chemotherapy for lymphoma: preliminary results. *Reprod Biomed Online* 2010;20:280–5.
- Brougham MFH, Crofton P, Johnson EJ, et al. Anti-Müllerian hormone is a marker of gonadotoxicity in pre- and post-pubertal girls treated for cancer: a prospective study. *Eur J Cancer*, submitted for publication.
- Bath LE, Wallace WH, Shaw MP, et al. Depletion of ovarian reserve in young women after treatment for cancer in childhood: detection by anti-Müllerian hormone, inhibin B and ovarian ultrasound. *Hum Reprod* 2003;18:2368–74.
- van Beek RD, van den Heuvel-Eibrink MM, Laven JS, et al. Anti-Müllerian hormone is a sensitive serum marker for gonadal function in women treated for Hodgkin's lymphoma during childhood. *J Clin Endocrinol Metab* 2007;92:3869–74.
- Lutchman Singh K, Muttukrishna S, Stein RC, et al. Predictors of ovarian reserve in young women with breast cancer. *Br J Cancer* 2007;96:1808–16.
- Lie Fong S, Laven JS, Hakvoort-Cammel FG, et al. Assessment of ovarian reserve in adult childhood cancer survivors using anti-Müllerian hormone. *Hum Reprod* 2009;24:982–90.
- Rosendahl M, Andersen CY, Ernst E, et al. Ovarian function after removal of an entire ovary for cryopreservation of pieces of cortex prior to gonadotoxic treatment: a follow-up study. *Hum Reprod* 2008;23:2475–83.
- Gosden RG, Telfer E, Faddy MJ, et al. Ovarian cyclicity and follicular recruitment in unilaterally ovariectomized mice. *J Reprod Fertil* 1989;87:257–64.
- Anders C, Marcom PK, Peterson B, et al. A pilot study of predictive markers of chemotherapy-related amenorrhea among premenopausal women with early stage breast cancer. *Cancer Invest* 2008;26:286–95.
- Rosendahl M, Andersen CY, la Cour Freiesleben N, et al. Dynamics and mechanisms of chemotherapy-induced ovarian follicular depletion in women of fertile age. *Fertil Steril* 2010;94:156–66.
- Yu B, Douglas N, Ferin MJ, et al. Changes in markers of ovarian reserve and endocrine function in young women with breast cancer undergoing adjuvant chemotherapy. *Cancer* 2010;116:2099–105.
- Anderson RA, Cameron DA. Pretreatment serum anti-Müllerian hormone predicts long-term ovarian function and bone mass after chemotherapy for early breast cancer. *J Clin Endocrinol Metab* 2011;96:1336–43.
- Anderson RA, Cameron DA. Pre-chemotherapy anti-Müllerian hormone is predictive of long-term ovarian function in women with breast cancer. *Endocr Rev* 2011;32, abstract OR17-6.
- Pigny P, Merlen E, Robert Y, et al. Elevated serum level of anti-Müllerian hormone in patients with polycystic ovary syndrome: relationship to the ovarian follicle excess and to the follicular arrest. *J Clin Endocrinol Metab* 2003;88:5957–62.
- Laven JS, Mulders AG, Visser JA, et al. Anti-Müllerian hormone serum concentrations in normoovulatory and anovulatory women of reproductive age. *J Clin Endocrinol Metab* 2004;89:318–23.
- Li HW, Anderson RA, Yeung WS, et al. Evaluation of serum antimüllerian hormone and inhibin B concentrations in the differential diagnosis of secondary oligoamenorrhea. *Fertil Steril* 2011;96:774–9.
- La Marca A, Volpe A. The anti-Müllerian hormone and ovarian cancer. *Hum Reprod Update* 2007;13:265–73.
- Li HW, Wong CY, Yeung WS, et al. Serum anti-Müllerian hormone level is not altered in women using hormonal contraceptives. *Contraception* 2011;83:582–5.
- van den Berg MH, van Dulmen-den Broeder E, Overbeek A, et al. Comparison of ovarian function markers in users of hormonal contraceptives during the hormone-free interval and subsequent natural early follicular phases. *Hum Reprod* 2010;25:1520–7.
- Nelson SM, Stewart F, Fleming R, et al. Longitudinal assessment of antimüllerian hormone during pregnancy-relationship with maternal adiposity, insulin, and adiponectin. *Fertil Steril* 2010;93:1356–8.

- [52] Dewailly D, Gronier H, Poncelet E, et al. Diagnosis of polycystic ovary syndrome (PCOS): revisiting the threshold values of follicle count on ultrasound and of the serum AMH level for the definition of polycystic ovaries. *Hum Reprod* 2011;26:3123–9.
- [53] Sowers MR, Eyvazzadeh AD, McConnell D, et al. Anti-Müllerian hormone and inhibin B in the definition of ovarian aging and the menopause transition. *J Clin Endocrinol Metab* 2008;93:3478–83.
- [54] Broer SL, Eijkemans MJ, Scheffer GJ, et al. Anti-Müllerian hormone predicts menopause: a long-term follow-up study in normoovulatory women. *J Clin Endocrinol Metab* 2011;96:2532–9.
- [55] Kok HS, van Asselt KM, van der Schouw YT, et al. Genetic studies to identify genes underlying menopausal age. *Hum Reprod Update* 2005;11:483–93.