THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Theropod (Dinosauria) diversity from the Potiguar Basin (Early-Late Cretaceous Albian-Cenomanian), northeast Brazil

Citation for published version:

Pereira, PVGC, Ribeiro, TB, Brusatte, S, Candeiro, CRA, Marinho, TS & Bergqvist, LP 2020, 'Theropod (Dinosauria) diversity from the Potiguar Basin (Early-Late Cretaceous Albian-Cenomanian), northeast Brazil', *Cretaceous Research*, vol. 114, 104517. <https://doi.org/10.1016/j.cretres.2020.104517>

Digital Object Identifier (DOI):

[10.1016/j.cretres.2020.104517](https://doi.org/10.1016/j.cretres.2020.104517)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Cretaceous Research

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Cretaceous Research

THEROPOD (DINOSAURIA) DIVERSITY FROM THE POTIGUAR BASIN (EARLY - LATE CRETACEOUS), NORTHEAST BRAZIL

--Manuscript Draft--

Manuscript Number:	YCRES_2019_258R2
Article Type:	Full Length Article
Keywords:	Dinosauria; Potiguar Basin; Theropoda; faunistic richness, Taxonomic description
Corresponding Author:	Paulo Victor Luiz Gomes da Costa Pereira UFRJ Rio de Janeiro, RJ Brazil
First Author:	Paulo Victor Luiz Gomes da Costa Pereira
Order of Authors:	Paulo Victor Luiz Gomes da Costa Pereira Theo Baptista Ribeiro Stephen Brusatte Carlos Roberto dos Anjos Candeiro Thiago da Silva Marinho Lilian Paglarelli Bergqvist
Abstract:	<p>The theropod record from the Cretaceous of northeastern Brazil are rare and consist mostly of isolated and incomplete remains, with only four species described. Here we describe, identify and evaluate the diversity of theropod materials from the Albian-Cenomanian Açu Formation, Potiguar Basin. The material consists of nine isolated vertebrae and a tooth. The vertebrae have been identified as belonging to four theropod groups: Abelisauria, Carcharodontosauria, Spinosauridae, Megaraptora, and Maniraptora. The isolated tooth was classified as belonging to a spinosaurid. One of the significant results so far is the occurrence of Megaraptora in the Potiguar Basin, based on the general morphology, the bones are very similar to Aerosteon and Megaraptor. Another unexpected result is the identification and presence of a maniraptoran caudal vertebrae, very rare in Brazil, with few fossils described. Besides this, other groups already found on isochronous basins of the Northeast region of Brazil and Africa as Carcharodontosauria and Spinosauridae. The presence of these theropod groups in the Açu Formation reveals a dinosaur richness in the Potiguar Basin similar to isochronous basins in Northern Africa and increases the knowledge about the diversity of South American dinosaurs.</p>
Suggested Reviewers:	<p>Manuel Alfredo Medeiros manuel.alfredo@ufma.br Extensive knowledge and experience with brazilian dinosaurs fauna</p> <p>Juan Canale juanignaciocanale@hotmail.com Interest and great knowledge of theropoda fauna in south america</p> <p>Rafael Matos Lindoso rafael.lindoso@ifma.edu.br Has experience with Theropoda materials from Brazil</p>
Opposed Reviewers:	

The suggestions were taken into consideration during the reviewing process and were a valuable contribution for the improvement of the paper.

Almost all the specific revisions and suggestions from reviewer 02 and 03 were made with no exception and are highlighted in red color in the text (in the “with highlights” version). Bellow we answer some of the edito/reviewers question and suggestions.

Editor: Place your Figure captions at the end of the manuscript file, after References.

A: Done

Editor: As the reviewers have commented below, the English of the manuscript needs to be improved (grammar, syntax, structure of sentences, odd phrasing, etc.). It is the authors' responsibility to proof their manuscript for English problems. The revised version should be carefully proofed before you resubmit it. You should seek help from your co-author, Prof. Steve Busatte, to carefully go through the manuscript before submission. This would help to remove the linguistic problems and in dealing with all the key issues mentioned by the reviewers.

A: Done. The English was improved. Prof. Steve Busatte read carefully the manuscript.

Editor: Order of Figures: All figures must to be presented in the same sequence that have their first citations in the manuscript text. Please check and correct carefully the order of figures in the manuscript and their corresponding first citations. Delete out-of order citations and/or rearrange them if necessary. Check also for missing citations

A: Done.

Reviewer 01: All revision in the abstract.

A: All the revisions were accepted.

Reviewer 01: I recommend to change the title for a more realistic one, for example: "Theropod dinosaur remains from..."

A: We understood the statement of Revisor 01, but we preferred to keep the original title, as it summarizes the results seen in the manuscript.

Reviewer 03: Five theropod species only in the northeast, or in all Brazil?

A: The four specimens are only from northeastern Brazil. This detail is better explained in the second paragraph of the Introduction.

Reviewer 03: Spinosauridae was recognized by a tooth, not a vertebra

A: done

Reviewer 03: I suggest to end this sentence with something more "... about the diversity of dinosaursxxx..."

A: Done

Reviewer 01: Please, restrict to Cretaceous record, to address the relevance of the materials here reported.

A: We choose to keep this part of the text to preserve the coesion within of the Introduction, featuring the first works and discoveries on theropod paleontology of Brazil. The next paragraph has a small summary of the Cretaceous northeastern theropoddinosaurs of Brazil.

Reviewer 01: Please, restrict to Cretaceous record

A: Done

Reviewer 01: Confuse

A: We rewrite this paragraph.

Reviewer 03: To this list should be added the recently described *Vespersaurus paranaensis* Langer et al., 2019 and *Gnathovorax cabreirai* Pacheco et al., 2019.

A: We choose to limit this list to the northeastern Brazil's theropod record only to make the text more easy to read/understand

Reviewer 03: The authors do not describe any osteoderm in the text.

A: Done

Reviewer 01: Do you mean the sedimentary infilling is divided into three groups?

A: Yes, we are following the terminology and geological description of the Basin.

Reviewer 03: I suggest to add its collection number

A: Done

Reviewer 03: They were described only five morphotypes in the text.

A: Done

Reviewer 01: This is not necessary to say. Anatomy is based on morphology. Avoid the use of "morphotypes", as if they were discrete biological units. Based on morphology, you identify vertebrae as corresponding to such section of the column, and to which theropod clade it may belong. Identification of "morphotypes" is useless.

A: We agree. We decided to remove the word “morphology” to avoid confusion.

Reviewer 01: Considering that this is not a ms devoted to analyze theropod teeth as a whole, and taking into account that just only one tooth is described, I suggest to remove all these considerations.

A: Done.

Reviewer 03: I suggest to be consistent with the abbreviations, using their English versions as is usual besides FABL: CBW: crown basal width (rather than EST), CH: crown height (rather than ALT), etc. See Hendrickx et al 2015. The dentition of megalosaurid theropods . Acta Paleontologica Polonica 60 (3): 627-642.

A: We followed the reviewer 01 and removed this part.

Reviewer 03: The values of most of these parameters were not specified in the analysis of the tooth recovered. I suggest to add them.

A: We followed the reviewer 01 and removed this part.

Reviewer 03: This measurement has a big fault, given there covered tooth lacks most of its tip.

A: Done. We didn't use this measurement any more.

Reviewer 03: Other abbreviation was used in Material and methods

A: We followed the reviewer 01 and removed this part.

Reviewer 01: Remove this and replace for "Systematic Paleontology"

A: Done

Reviewer 03: I suggest that, before comparing this material with other spinosaurids, the authors should add a summary of characters that allow to assign it to Spinosauridae.

A: Done. We summarize the characteristics of spinosaurids in the Discussion.

Reviewer 03: Given there are two morphotypes assigned to Megaraptora, both showing similar characteristics, the authors should explain why there are not included in a single morphotype.

A: Done. We reorganized the material in the groups and removed the division “Morphotypes”

Reviewer 03: The word “expressive” seems wrong in this context.

A: Done.

Reviewer 01: Which is the systematic relevance of all these ratios?

A: We strongly suggest that the reviewer read the articles about the use of quantitative analysis in the identification of theropod teeth:

-HENDRICKX, C. & MATEUS, O. 2014. Abelisauridae (Dinosauria: Theropoda) from the Late Jurassic of Portugal and dentition-based phylogeny as a contribution for the identification of isolated theropod teeth. *Zootaxa*, 3751, 1–74.

-HENDRICKX, C., MATEUS, O., ARAÚJO, R. 2015. A proposed terminology of theropod teeth (Dinosauria, Saurischia). *Journal of Vertebrate Paleontology* 35 (5), e982797. <http://dx.doi.org/10.1080/02724634.2015.982797>. HENDRICKX, C. & MATEUS, O. 2014. Abelisauridae (Dinosauria: Theropoda) from the Late Jurassic of Portugal and dentition-based phylogeny as a contribution for the identification of isolated theropod teeth. *Zootaxa*, 3751, 1–74.

-HENDRICKX, C., MATEUS, O., ARAÚJO, R., AND CHOINIERE, J. 2019. The distribution of dental features in non-avian theropod dinosaurs: Taxonomic potential, degree of homoplasy, and major evolutionary trends. *Palaeontologia Electronica* 22.3.74 1– 110. <https://doi.org/10.26879/820palaeo-electronica.org/content/2019/2806-dental-features-in-theropods>.

-SMITH, J.B., VANN, D.R., DODSON, P. 2005. Dental morphology and variation in theropod dinosaurs: implications for the taxonomic identification of isolated teeth. *The Anatomical Record Part A* 285: 699-736.

-SANKEY, J.T., BRINKMAN, D.B., GUENTHER M., CURRIE, P.J. 2002. Small theropod and bird teeth from the Late Cretaceous (Late Campanian) Judith River Group, Alberta. *Journal of Paleontology*, 76, 751–763.

Reviewer 01: “Which are the anatomical bases to refer these elements as to Neovenatoridae/Megaraptora? The present manuscript suffers of the lack of anatomical descriptions and comparissons allowing the reader to understand why present authors conclusions.”;

“Please, provide morphological bases to support this referral.”;

“Please, give reasons for referring these elements as to Abelisauroidae”

“Please, explain why this element is referred as to Maniraptora”

“Please, explain why this element is referred as to Carcharodontosauria”

“Please, explain why this element is referred as to Megaraptora”;

A: These parts have been rewrite to increase cohesion and make it clearer to understand. The first paragraph of the "comparisons" part has the characteristics used to allocate the fossil in the specific theropod group and the corresponding bibliography.

Reviewer 01: Please, obviate this detail. It is not morphological!

A: We removed the measurements.

Reviewer 01: Please, clarify

A: Done

Reviewer 01: Before description of any dinosaur bone, it must be glued.

A: Done. Material have been found associated, but there is no clear point of junction between both pieces, with most of the middle portion being lost.

Reviewer 03: Sem-spherical or semicircular?

A: Done. Semicircular

Reviewer 03: The authors should describe (and figure) the materials as a single vertebra, not as two different fragments. It is confusing and does not help with interpretation of the materials.

A: Ok. Done

Reviewer 03: At the beginning of this morphotype description the authors suggest that they are possibly caudal vertebrae.

A: Ok. Done

Reviewer 03: Why anterior? It could not be posterior?

A: We removed this fragment because it is not significant or relevant for this manuscript.

Reviewer 03: Why the authors assign this material to morphotype 1? They only suggest that it belongs to a theropod.

A: We removed this fragment because it is not significant or relevant for this manuscript.

Reviewer 01: I am surprised with this statement: Ceratosauria as members of Tetanurae. This is not a serious manuscript.

A: It was just a confusion when we were arranging the morphotypes. We know that Ceratosauria is not inside Tetanurae. Thanks for the revision...

Reviewer 03: This character used for differentiate this morphotype is clearly related to the position of the vertebra inside the vertebral series; This character is used for differentiate this morphotype is also present in morphotype 1; I suggest to look for other more specific characters to differentiate this morphotype.

A: We revised the attribution of this morphotype in Abelisauria and not found any solid characteristic to sustain this classification. We decided to follow the reviewers and be conservative and put these material at Theropoda indet.

Reviewer 03: I think both are synonyms

A: Done.

Reviewer 03: This sentence is confusing.

A: Done.

Reviewer 03: Think the authors should specify which transition point are referring here (I suppose that proposed by Russel (1972), but this is my guess)

A: Done

Reviewer 03: The shape of the articular face of caudal vertebrae is so variable, it show differences inside the same taxonomic group. For example inside Abelisauridae, Carnotaurus shows semicircular articular surface, and Majungasaurus show oval articular surface, as the authors clearly show in the figure 8. This makes this character not useful for separating morphotypes.

A: We don't use anymore this ratio type of character on the description.

Reviewer 03: The articular surfaces shown in the figure 4 has ovoidal articular faces.

A: Done.

Reviewer 03: hourglass-shaped

A: Done.

Reviewer 03: The lateral surfaces are slightly concave in these taxa, not very.

A: Done

Reviewer 03: Please see Aranciaga-Rolando et al (2018) A supposed Gondwanan oviraptorosaur from the Albian of Brazil represents the oldest South American megaraptoran. Cretaceous Research 84: 107-119.

A: Done

All the reviewers: All the revisions in the discussion.

A: We followed the suggestion of the reviewer 01 and reorganized the discussion. We replaced the most part to results in the comparative morphology and rewrite the discussion based on the importance of the Potiguar's fossils.

FIGURES

We done all the revisions requested by the reviewers and improved the figures.

Highlights – article: THEROPOD (DINOSAURIA) DIVERSITY FROM THE POTIGUAR BASIN (EARLY - LATE CRETACEOUS), NORTHEAST BRAZIL

- These are the first described theropod materials from the Potiguar Basin, Brazil.
- Four morphotypes were described based on morphological and/or diagnostic characters.
- Carcharodontosauria and Spinosauridae were groups identified.
- Rare Megaraptora, and Maniraptora materials were also identified.

**THEROPOD (DINOSAURIA) DIVERSITY FROM THE POTIGUAR BASIN
(EARLY-LATE CRETACEOUS ALBIAN – CENOMANIAN), NORTHEAST
BRAZIL**

**Paulo Victor Gomes da Costa Pereira^{1*}; Theo Baptista Ribeiro¹; Stephen Louis
Brusatte²; Carlos Roberto Dos Anjos Candeiro³; Thiago da Silva Marinho⁴; Lilian
Paglarelli Bergqvist¹**

*corresponding author – paulovictor29@yahoo.com.br

1- CCMN - Universidade Federal do Rio de Janeiro, Ilha do Fundão, Rio de Janeiro (RJ), Brazil.

E-mail: paulovictor29@yahoo.com; theobribeiro1@gmail.com; bergqvist@geologia.ufRJ.br

2 - University of Edinburgh, Grant Institute, School of GeoSciences. Edinburgh, Scotland, UK. E-mail:
stephen.brusatte@ed.ac.uk

3 - Universidade Federal de Goiás (UFG), Laboratório de Paleontologia e Evolução (LABPALEOEVO),
Curso de Geologia. Aparecida de Goiânia, GO, Brazil. E-mail: candeiro@ufg.br

4- Centro de Pesquisas Paleontológicas L. I. Price, Complexo Cultural e Científico Peirópolis, Pró-Reitoria
de Extensão Universitária, Universidade Federal do Triângulo Mineiro, Uberaba, Minas Gerais, Brazil. E-
mail: tsmarinho@icene.ufTM.edu.br

Abstract

The theropod record from the Cretaceous of northeastern Brazil are rare and consist mostly of isolated and incomplete remains, with only four species described. Here we describe, identify and evaluate the diversity of theropod materials from the Albian-Cenomanian Açu Formation, Potiguar Basin. The material consists of seven isolated theropod vertebrae and a tooth. We identify the material as belonging to four theropod

groups: Spinosauroidae, Carcharodontosauria, Megaraptora, and Maniraptora. One of the significant results is the occurrence of Megaraptora in the Potiguar Basin; based on the general morphology, some of the bones we describe are very similar to those of *Aerosteon* and *Megaraptor*. Another unexpected result is the identification and presence of a maniraptoran caudal vertebrae; these dinosaurs are very rare in Brazil, with few fossils previously described. Furthermore, we identify other groups that have already been found in isochronous basins of the Northeast region of Brazil and Africa, including Carcharodontosauria and Spinosauroidae. The presence of these theropod groups in the Açu Formation reveals a dinosaur richness in the Potiguar Basin similar to isochronous basins in Northern Africa and increases knowledge about the diversity of South American dinosaurs.

Keywords: Dinosauria, Potiguar Basin, Theropoda, faunistic richness, Carcharodontosauria, Megaraptora, Spinosauridae, Maniraptora.

Introduction

The first studies in Brazil that attributed vertebrate fossil remains to dinosaurs were published in the nineteenth and mid-twentieth centuries (Marsh, 1869; Derby, 1890; Mawson and Woodward, 1907; Price, 1960, 1961). Since that time, dinosaur fossils have been recorded from three principal localities and ages in Brazil: the Triassic of the Santa Maria and Caturrita formations (Langer et al., 2007a), the mid-Cretaceous of the Araripe, Triunfo and São Luís-Grajaú basins (Frey and Martill 1995; Kellner 1996a, b, 1999; Medeiros et al., 2007; Carvalho et al., 2017), and the Late Cretaceous of the Bauru and Paracatu groups (Franco-Rosas et al., 2004; Kellner et al., 2004; Brusatte et al. 2017).

There are eight theropod dinosaur species formally described from Brazil so far, four from the northeastern region: *Santanaraptor placidus* Kellner, 1999, *Irritator challengeri* Martill, Cruikshank, Frey, Small and Clarke, 2002 and *Mirischia asymmetrica* Naish, Martill and Frey 2004 from the Araripe Basin; and *Oxalaia quilombensis* Kellner, Azevedo, Machado, Carvalho and Henriques, 2011 from the São Luís-Grajaú Basin.

A promising area for new dinosaur discoveries is the rocks of the Açu Formation, in the Potiguar Basin. Until now, the macrofossils of the Açu Formation consisted of bivalve molluscs, small crustaceans, fish scales, and plant remains (Duarte and Santos, 1961). However this changed in the 2000s, when researchers from the Group of Analogs to Oil Reservoirs of the Department of Geology of the Federal University of Rio Grande do Norte, in geological mapping of the Açu 4 operational unit, found large vertebrate fossils.

In the decade after the discovery of these first continental vertebrate fossils in the formation (Santos *et al.*, 2005), no other fieldwork was conducted. However, in 2015 and 2016, this area was again prospected by Laboratório de Macrofósseis of the Universidade Federal do Rio Janeiro and dozens of fossils were found. The aim of the present work is to describe and identify the collected materials attributed to theropod dinosaurs, showing that the Potiguar Basin preserves a large diversity of species and has great potential for future discoveries and studies about the mid-Cretaceous paleoenvironments of the Atlantic margin of Brazil.

Geological Setting and Lithostratigraphy

The Potiguar Basin is located at the eastern continental margin of northeastern Brazil, cropping out in the states of Rio Grande do Norte and Ceará (Fig. 01), with a total

estimated area of 60,000 km², of which 22,000 km² is interpreted as continental (Cassab, 2003). The Potiguar Basin is bounded to the east by Alto de Touros, which separates it from the Pernambuco-Paraíba Basin, to the northwest by the Alto de Fortaleza, which separates it from the Ceará Basin, and to the south and west by crystalline basement rocks (Pessoa-Neto *et al.*, 2007).

The sedimentary units of the Potiguar Basin are divided into three groups: Areia Branca (Pendência and Alagamar formations), Apodi (Açu, Quebradas and Jandaíra formations) and Agulha (Ubarana, Guamaré and Tibau formations) (Araripe and Feijó, 1994). The Açu formation is divided into four subunits according to electric logs, identified from bottom to top as Açu 1, Açu 2, Açu 3 and, Açu 4 (Vasconcelos *et al.*, 1990). The material described here comes from the Açu 4 subunit, which corresponds to a transgressive, coastal-estuarine system.

The Açu-4 Unit consists of sixteen facies, fourteen being siliciclastic and two being hybrid. The siliciclastic facies are grouped into nine associations, namely: (1) lag residual deposits, (2) channel fill deposits, (3) crevasse-splay deposits, (4) floodplain deposits, (5) abandoned channel deposits, (6) upper-flow regime sandflat deposits, (7) lower-flow regime sandflats, (8) sandflat/mudflat deposits of restricted environment, and (9) mudflat deposits. The first five facies associations represent a meandering fluvial system with tidal influence, and the remaining integrate the intermediate and distal sectors of an estuarine complex dominated by tides. The hybrid facies were deposited in a shallow platform adjacent to an estuary (Costa *et al.*, 2014).

The Açu Formation has been suggested to be Albian-Cenomanian in age (Early–Late Cretaceous), based on palynological data (Araripe and Feijó, 1994).

Material and Methods

The fossils were collected from outcrops of the Açú Formation, Potiguar Basin (Ceará state, northeastern Brazil) and are deposited at the Fossil Reptile Collection of the Departamento de Geologia (DG), Universidade Federal do Rio de Janeiro (UFRJ). The material consists of seven isolated theropod vertebrae (UFRJ-DG 521-R, 523-R, 524-R, 528-R, 558-R, 575-R, and 634-R) and a tooth (619-Rd)

The following tooth characteristics were assessed, following the nomenclature proposed by Hendrickx *et al.* (2015): general morphological traits of the dental crown (its overall shape, curvature, ornamentations in the enamel), denticles (presence, size and shape) cross section (compression and shape), orientation of the tooth (lingual, labial, mesial and distal) and measurements.

Systematic paleontology

SAURISCHIA Seeley, 1888

THEROPODA Marsh, 1881

Referred material: UFRJ-DG 532-R and 575-R.

Description:

UFRJ-DG 528-R

Specimen 528-R is a theropod vertebral centrum (Fig. 02, C-E). It is amphicoelous, and slightly higher than long. Its lateral surface is smooth and slightly concave, without marks or other remarkable characteristics, giving the vertebra a straight and somewhat featureless appearance. The ventral surface smooth with no groove or keel and it is slightly concave in lateral view.

The dorsal surface possesses a distinct longitudinal groove extending from one articular facet to the other that can be identified as the neural canal. The articular faces

have nearly straight margins. The anterior facet is somewhat concave, and the posterior is slightly convex and slightly oval in shape; both articular facets have the same general proportions (height longer than length). The anterior articular face presents a deeper concavity, and is slightly larger in size, than the posterior face, which is very flat and without deep depressions.

UFRJ-DG 575-R

Specimen 575-R (Fig. 02, A-B) is a theropod vertebral centrum broken in two: a smaller anterior piece and a larger posterior section. Although the material was found associated there is no clear point of junction between both pieces, as most of the middle portion has been lost. The anterior fragment exhibits a very concave articular face of semi-circular shape and slightly forward-protruding margins.

On the lateral surface of the anterior fragment there is a deep perforation close to the dorsal region that reaches the other lateral surface, which can be described as a pleurocoel. The ventral surface of the anterior fragment is smooth and concave in anterior view. The dorsal surface of the anterior fragment is broken, missing most of the surface above the pleurocoel.

The posterior fragment has a slightly smaller articular surface, which is broken on the anterior portion; it is also concave and of semi-circular shape, with slightly backwards-protruding margins. Its dorsal surface and the dorsal half of the left lateral surface are broken, while the right lateral surface is broken in a slightly more dorsal region in comparison to the left one. The ventral surface of the fragment is smooth and concave in lateral view. Due to the highly fragmentary state of UFRJ-DG 575-R, it is possible to see multiple small pervasive pneumatic chambers, the camellae, in the internal bone.

Comparisons:

The highly pneumatized camellate bone seen in UFRJ-DG 575-R is a characteristic seen in many groups of theropods, from the basal *Ceratosaurus* to tetanuran groups such as carcharodontosaurids and coelurosaurs mainly in its presacral vertebrae (Carrano and Sampson, 2008). This feature, together with the poor preservation of this specimen, which prevents the identification of other more diagnostic characteristics, hinders the classification of this specimen beyond Theropoda.

THEROPODA Marsh, 1881

TETANURAE Gauthier, 1986

? SPINOSAUROIDEA Stromer, 1915

Referred material: UFRJ-DG 619-Rd.

Description:

UFRJ-DG 619-Rd (Fig. 03) is a fragment of a large isolated tooth crown, probably belonging to the middle to almost apical portion of the tooth. The specimen lacks any form of enamel, as it has dentine exposed, what prevents description of external ornamentation such as transversal undulations, flutes and denticulation. The crown is almost completely straight with only a subtle curvature in its lingual surface, while the labial surface remains slightly convex.

The crown fragment has an overall cone-like shape with an almost ovoid cross section. In basal view, it is possible to see concentrically deposited rings of dentine surrounding a small depression, which probably represents the apical-most portion of the dental pulp cavity.

Comparisons:

UFRJ-DG 619-Rd have some characteristics that it shares with the highly specialized teeth of spinosauroid theropods. The most salient of these is the almost straight conical shaped crown, with an ovoid cross section, a feature often seen in piscivorous animals (Mateus, 2011; Hendrickx and Mateus, 2014).

THEROPODA Marsh, 1881

TETANURAE Gauthier, 1986

MANIRAPTORA Gauthier, 1986

Referred material: UFRJ-DG 521-R

Description:

UFRJ-DG 521-R

Specimen UFRJ-DG 521-R (Fig. 04) is an almost complete distal caudal vertebrae of a maniraptoran theropod. It is amphicoelous with a length to height ratio of almost 2.5, making it a least twice longer than tall. The dorsal surface of the centrum is almost complete with half of a dorsal midline ridge reminiscent of reduced neural spine, a well preserved and more dorsally positioned prezygapophysis, and a lost postzygapophysis. The prezygapophysis articular surface is ellipsoid and is reclined 45° laterally. The neural canal is almost completely preserved, having lost only its posterior half.

The lateral surfaces of the centrum are mostly smooth, marked only with a midline ridge reminiscent of a reduced transverse processes. The ventral surface of the centrum has a shallow groove that extends from one articular facet to the other. In the lateral view the ventral surface is slightly concave.

The articular facets of the centrum are both concave, with the anterior facet being more excavated than the posterior facet, and have a semi-circular shape. The articular

margins are almost straight, with the anterior margin being larger than the posterior margin.

Comparisons:

UFRJ-DG 521-R has characteristics of a maniraptoran centrum positioned after the transition point in the tail (Russell, 1972, Gauthier, 1986; Tykoski, 2005), as it is longer than high and possesses a large reduction in both its neural spine and transverse processes, with those structures becoming midline ridges (Senter *et al.*, 2011; Motta *et al.*, 2018). Thus, it is possible to deduce that it is positioned after vertebra 11 of the caudal series as seen in *Buitreraptor*, *Rahonavis*, Dromaeosauridae and Troodontidae (Ostrom, 1969; Forster *et al.*, 1998; Senter *et al.*, 2012; Xu *et al.*, 2017).

The presence of a reduced transverse process forming a midline ridge after the transition point is seen in the distal caudal vertebrae of *Rahonavis* and *Buitreraptor* (Forster *et al.*, 1998; Novas *et al.*, 2017), a characteristic also seen in UFRJ-DG 521-R, which differentiates it from most other paravians as dromaeosaurids, *Archaeopteryx*, *Jeholornis* and *Anchiornis*.

In addition, the 521-R specimen also has dorsally positioned pre-zygapophyses in the same way as in *Buitreraptor*, *Rahonavis* and *Anchiornis* (fig. 05) (Motta *et al.*, 2018). The vertebral centrum has a length-to-height ratio between close to 2.5, a ratio usually seen in dromaeosaurids with exception to *Buitreraptor* but not seen in other maniraptorans as troodontids and microraptorians whose ratio can reach up to 5.0 to 6.0.

THEROPODA Marsh, 1881

TETANURAE Gauthier, 1986

ALLOSAUROIDEA Marsh, 1878

CARCHARODONTOSAURIA Benson, Brusatte and Carrano, 2010

Referred material: UFRJ-DG 523-R and 524-R.

Description:

UFRJ-DG523-R

Specimen 523-R (Fig. 06, D-F) is a theropod vertebral centrum, with the following characteristics: it is amphicoelous, and slightly longer than high. Its lateral surface is very concave and smooth on both sides, with the shape of an hourglass in dorsal view. The ventral surface is mostly smooth on the anterior part, with marks that possibly indicate the articulation with the hemal arch on the posterior part.

The dorsal surface is marked by a long and deep longitudinal canal from one articular face to the other, which widens on the extremities and tapers in the middle. This canal was possibly the space of the neural canal of the vertebra, given the marks of fusion with the neural arch that meet on its borders.

The articular faces are ovoid in shape and have slightly forward-protruding margins, the anterior facet being higher in comparison to the posterior facet. The anterior articular face has a concavity deeper than the posterior one, being also slightly larger in its proportions.

UFRJ-DG524-R

Specimen 524-R (Fig. 06, A-C) is a centrum of a theropod caudal vertebra. It is amphicoelous and is slightly longer than high, which indicates a more proximal position in the caudal series. The lateral surface is smooth and marked by two deep concavities on both lateral faces, giving it an hourglass-like shape.. Additionally, on the most dorsal region of the lateral surface there is a small and shallow longitudinal depression on each side.

The ventral surface is a double keel marked by a very superficial groove extending from the anterior part up to the posterior part. The dorsal surface is marked by the neural canal of the vertebra. Above the anterior part of this canal the entire upper portion of the neural tube is preserved, forming a small arch filled by sediment positioned slightly above the anterior articular face.

The articular faces are semi-circular and somewhat oval, with the anterior one being slightly larger than the posterior, and their margins slightly protrude forward. The anterior articular face has a concavity slightly deeper than the posterior.

Comparisons:

Both UFRJ-DG 523 and 524 present characteristics commonly found in carcharodontosaurids (Fig. 07). For instance, depressions in the most dorsal part of the lateral surface are found in *Giganotosaurus*, *Mapusaurus* and *Tyrannotitan* and in the mid-caudal vertebra Vb-870 found in the Wadi Milk Formation (Coria and Salgado, 1995; Coria and Currie, 2006; Novas *et al.*, 2005a; Canale *et al.*, 2015; Rauhut, 1999). This condition is different from that in *Carcharodontosaurus*, which has pleurocoels in its anterior caudal vertebrae (Stromer, 1931). Furthermore, the strongly waisted centrum morphology, a double keel cut by a longitudinal groove and offset articular facets (although it is a plesiomorphic feature found in *Allosaurus* Gilmore, 1920; Madsen, 1976) are also found in specimens such as the carcharodontosaurid material from Sudan (Rauhut, 1999) and in *Tyrannotitan*, *Mapusaurus* and *Acrocanthosaurus* (Canale *et al.*, 2015; Harris, 1998; Coria and Currie, 2006; Currie and Carpenter, 2000).

THEROPODA Marsh, 1881

NEOVENATORIDAE Benson, Carrano and Brusatte, 2010

MEGARAPTORA Benson, Carrano and Brusatte, 2010

Referred material: UFRJ-DG 558-R e 634-R

Description:

UFRJ-DG 558-R

Specimen 558-R is a centrum of a theropod caudal vertebra, damaged by various cracks (Fig. 08, D-F). It is amphicoelous, and slightly longer than high, indicating a somewhat proximal position within the caudal series. Its ventral surface is very smooth and convex in lateral view, but is very damaged in the region where the base of the posterior articular face would be.

The dorsal surface is marked by a great depression extending longitudinally from one articular face to the other, wider in the extremities, denoting the neural canal. The lateral surfaces are marked by a longitudinal elliptic depression on their medial parts, where there is a pleurocoel on each side. The left lateral pleurocoel is deeper and better defined than the right lateral one.

Its articular faces are semi-circular and have very straight margins. The anterior articular face possesses a more distinctive depression of a slightly greater size than the posterior face and is also in a better state of preservation. The posterior articular face possesses a very slight concavity, making it almost straight, and is in a much more damaged state, presenting cracks and breaches on the ventral base of the face.

UFRJ-DG 634-R

This material is in a worse state of preservation than UFRJ-DG 558-R (Fig. 08, A-C). The ventral centrum portion and anterior articular face are fragmented. On its lateral surface, there is what appears to be the border of the pleurocoel in the same position seen in specimen 558-R.

Different from the other vertebra of this group, part of the neural arch and the transverse process are preserved on the right side of the specimen. The transverse process is positioned upwards at an angle of approximately 45° degrees.

Comparisons

The presence of pleurocoels in the caudal vertebrae is characteristic of megaraptoran neovenatids (Benson *et al.*, 2010). Pneumaticity in the caudal vertebrae is rare in Theropoda, present only in some groups: Megaraptora, Oviraptorosauria, Therizinosauria, and Carcharodontosauridae (Benson *et al.*, 2012). As far as is known, no fossils of therizinosaurs have been found in South America and South American fossils attributed to oviraptorosaurs have been reassigned to other taxa, including to Maniraptora (e.g. Agnolín and Martinelli, 2007, Aranciaga-Rolando *et al.*, 2018). In addition, the caudal vertebrae of Oviraptorosauria have, on the ventral surface, a medial groove delimited by two longitudinal elevations (e.g., Sues, 1997; Xu *et al.*, 2007). Specimen UFRJ-DG 558-R does not have this feature (Fig. 09).

South American carcharodontosaurids (e.g., *Giganotosaurus*, *Mapusaurus*, *Tyrannotitan*) show slightly concave lateral sides in the caudal vertebrae, but do not bear actual pneumatic foramina. Stromer (1931) described an anterior caudal vertebra from northern Africa, which he identified as *Carcharodontosaurus*, which had pneumatic characteristics, including a pleurocoel. However, that vertebra has a different general morphology and proportions when compared with the megaraptorid vertebrae from the Potiguar Basin (length-height ratio is 1 in *Carcharodontosaurus* and approximately 1.48 in UFRJ DG 558-R) and other members of Megaraptora.

Among the Megaraptora group, only *Aerosteon*, *Aoniraptor*, *Orkoraptor* and *Megaraptor* have preserved caudal vertebrae (Fig. 10) (Sereno *et al.*, 2008; Benson *et al.*, 2010; Motta *et al.*, 2016). The height/length ratio of UFRJ DG 558-R is 1.4, consistent

with a median tail position, compared to the ratios of 1.2 and 1.3, respectively, of the medial caudal vertebrae of *Aerosteon* and *Orkoraptor* (Novas *et al.*, 2008). The Potiguar Basin specimens resemble those of *Aoniraptor* (Fig. 07, F) due to the absence of a keel in the ventral region, but are distinguished by the presence of a pair of pneumatic troughs in the lateral region, separated by a septum. Only the first caudal vertebra of *Aoniraptor* presents such fossae, a characteristic present in the other megaraptorans (e.g., Novas *et al.*, 2008; Sereno *et al.*, 2008).

Comparing the morphology of pneumatic foramina, UFRJ DG 558-R (Figure 10, A) is very similar to *Aerosteon* (Figure 10, C), *Megaraptor* (Figure 10, H) and *Orkoraptor* (Figure 10, G) in the presence of a large elliptic foramen and a second smaller circular shaped foramen. In addition, UFRJ-DG 558-R and 634-R has its cavities located on the lateral surface of the vertebral centrum near the base of the neural arch, which does not occur in the other species observed.

UFRJ-DG 558-R and 634-R also presents extensive pneumatization in the vertebral centrum, composed of a camerate internal microstructure (Britt, 1993), with several small chambers, similar to other megaraptorans (e.g., *Aerosteon*, *Megaraptor*; Martinelli *et al.*, 2013).

Discussion

The Açu Formation material and its importance

The fossil potential of Açu Formation was poorly known, with only a few fossils recovered (Duarte and Santos, 1962; Silva-Santos, 1963; Mussa *et al.*, 1984), until the discovery of vertebrae and teeth identified as belonging to Theropoda indet. and Titanosauria (Santos *et al.*, 2005).

No further work was conducted until 2018, when the materials described here were studied in more detail. Thus far, the dinosaur fauna of the Potiguar Basin includes two groups of Sauropoda (Diplodocoidea: Rebbachisauridae, Pereira et al., in press; Titanosauriformes, Barbosa et al., 2018; Titanosauria, Pereira et al., 2018) and four groups of Theropoda (Spinosauroidae, Carcharodontosauridae, Megaraptora and Maniraptora, present work).

The occurrence of these groups (except Megaraptora) in the Potiguar Basin is yet another similarity between the faunas of northeastern Brazil and multiple North Africa Cretaceous units (e.g. Medeiros and Schultz, 2001a, 2002; Sereno and Brusatte, 2008; Contessi, 2009; Candeiro et al., 2011; Candeiro, 2015). Except for the Elrhaz (Niger); Douiret and Ain El Guettar (both in Tunisia) and Chicla (Libya) formations, which were dated to the Early Cretaceous, all other Cretaceous formations from Northern Africa are Albian-Cenomanian in age, roughly equivalent to the Açu Formation (Werner, 1994; Rossetti, 1997; Rossetti and Truckenbrodt, 1997; Smith et al., 2001; Anderson et al., 2007; Sereno and Brusatte, 2008; Cavin et al., 2010). Among the formations, the Alcântara Formation (Brazil), Bahariya Formation (Egypt), Echkar Formation (Niger) and the Waldi Milk Formation (Sudan) have similarities with the Açu Formation's dinosaur fauna.

According to paleobiogeographic models, South America and Africa started separating from each other in the Valanginian (Early Cretaceous), leading to the formation of the South Atlantic Ocean (Viramonte et al., 1999; Jokat et al. 2003; Macdonald et al., 2003). Although the ocean turned into one of the most important continental barriers of the southern hemisphere, faunal interchange among the terrestrial landmasses of western of Gondwana definitely occurred up to the Albian, and possibly

until the Cenomanian (e.g. Petri, 1987; Reyment and Dingle, 1987; Pletsch et al., 2001, Tello Saenz et al., 2003, Guedes et al., 2005, Bodin et al., 2010).

Based on the proposed age and geographic position, the fossil vertebrates of the Açu Formation may have lived during some of the last intervals of continental connection between South America and Western Africa, before the complete formation of the South Atlantic Ocean (Arai, 2009; Castro et al., 2012). This makes them exceedingly important for understanding biogeography and faunal evolution.

More extensive comparisons are still limited by the lack of completeness of the Açu material and the absence of formally described taxa. The continuation of studies on previously collected material (like that described in this paper) and prospecting for new fossils is important in this basin which, while still the subject of only recent research, already exhibits among the greatest diversity of dinosaur groups in Brazil.

Conclusion

In the present work we assigned the material from Açu Formation, Potiguar Basin, to four groups: Spinosauroida, Carcharodontosauria, Maniraptora and Megaraptora (Fig. 11), the two last groups being relatively rare in Brazil. All these groups have already been found in isochronous formations in both Northeastern Brazil and Northern Africa, leading further support for faunal similarities in the “mid”-Cretaceous western Gondwana. These fossils provide the first theropod record from Potiguar Basin and an important opportunity to increase the knowledge on the diversity of this still poorly known basin.

Acknowledgements

The authors thank Prof. Dr. Valdeci dos Santos Júnior Santos for discovering the site where the material were found, and for the support given to the fieldwork. We thank

the students Luciano Vidal for assistance with the figures. PVLGCP was funded by a grant from the Jurassic Foundation and pos doctoral grant by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, number 88882.463232/2019-01). LPB and CRAC were financially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)/Bolsista de Produtividade em Pesquisa. IMMGB and LPB were also supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico [grant 459086/2014-6]. LPB also acknowledge Fundação de Amparo a Pesquisa do Estado do Rio de Janeiro [grants #E-26/202.829/2018]. Our collaborative project was funded also by a grant from the Fundação de Amparo a Pesquisa e Goiás and the Newton Fund, which supported SLB's visit to Brazil to work with PVGCP and CRAC in June–July 2016.

References

- Agnolín, F. L., Martinelli, A.G. 2007. Did oviraptorosaurs (Dinosauria; Theropoda) inhabit Argentina. *Cretaceous Research* 28, 785–790.
- Anderson, P.E., Benton, M.J., Trueman, C.N., Paterson, B.A., Cuny, G., 2007. Palaeoenvironment of vertebrates on the southern shore of Tethys: the nonmarine Early Cretaceous of Tunisia. *Palaeogeography, Palaeoclimatology, Palaeoecology* 243 (1–2), 118–131.
- Arai, M. 2011. Paleogeografia do Atlântico Sul no Aptiano: um novo modelo a partir de dados micropaleontológicos recentes. *Boletim de Geociências da Petrobras*, 17(2): 331-351.
- Aranciaga-Rolando, A. M., Egli, F. B., Sales, M. A., Martinelli, A. G., Canale, J. I., Ezcurra, M. D. 2018. A supposed Gondwanan oviraptorosaur from the Albian of

- 419 Brazil represents the oldest South American megaraptoran. *Cretaceous Research* 84,
420 107-119.
- 421 Araripe, P. T., Feijó, F. J. 1994. Bacia Potiguar. *Boletim de Geociências da Petrobras* 8:
422 27-141.
- 423 Barbosa, F.H.S., Ribeiro, I.C., Pereira, P.V.L.G.C., Bergqvist, L.P. 2018. Vertebral
424 lesions in a titanosaurian dinosaur from the Lower-Upper Cretaceous of Brazil.
425 *Geobios* 51: 385–389.
- 426 Benson, R. B., Carrano, M. T., Brusatte, S. L. 2010. A new clade of archaic large-bodied
427 predatory dinosaurs (Theropoda: Allosauroidae) that survived to the latest Mesozoic.
428 *Naturwissenschaften*, 97(1), 71-78.
- 429 Benson, R. B., Butler, R. J., Carrano, M. T., and O'Connor, P. M. 2012. Air- filled
430 postcranial bones in theropod dinosaurs: physiological implications and the
431 'reptile'–bird transition. *Biological Reviews*, 87(1), 168-193.
- 432 Bodin, S., Petitpierre, L., Wood, J., Elkanouni, I., Redfern, J. 2010. Timing of Early to
433 mid-Cretaceous tectonic phases along North Africa: new insights from the Jeffara
434 escarpment (Libya–Tunisia). *Journal of African Earth Sciences* 58 (3), 489–506.
- 435 Britt, B. B. 1993. Pneumatic postcranial bones in dinosaurs and other archosaurs.
436 *Geology and Geophysics*, University of Calgary.
- 437 Brusatte, S. L., Candeiro C. R. A., Simbras, F. M. 2017. The last dinosaurs of Brazil: The
438 Bauru Group and its implications for the end-Cretaceous mass extinction. *Annals of*
439 *the Brazilian Academy of Sciences* 89: 1465-1485.
- 440 Canale, J. I., Novas, F. E., Pol, D. 2015. Osteology and phylogenetic relationships of
441 *Tyrannotitan chubutensis* Novas, de Valais, Vickers-Rich and Rich, 2005
442 (Theropoda: Carcharodontosauridae) from the Lower Cretaceous of Patagonia,

- 443 Argentina. Historical Biology 27, 1e32. [http://dx.doi.org/10.1080/](http://dx.doi.org/10.1080/08912963.2013.861830)
 444 08912963.2013.861830.
- 445 Candeiro, C. R. A.; Fanti, F.; Therrien, F.; Lamanna, M. C. 2011. Continental fossil
 446 vertebrates from the mid-Cretaceous (Albian-Cenomanian) Alcântara Formation,
 447 Brazil, and their relationship with contemporaneous faunas from North Africa.
 448 Journal of African Earth Sciences 60, 79-92.
- 449 Candeiro, C. R. A. 2015. Middle Cretaceous dinosaur assemblages from northern Brazil
 450 and northern Africa and their implications for northern Gondwanan composition.
 451 Journal of South American Earth Sciences 61, 147-153.
- 452 Carrano, M. T. and Sampson, S. D. 2008. The phylogeny of ceratosauria (Dinosauria:
 453 Theropoda). Journal of Systematic Palaeontology, 6(2), 183-236.
- 454 Carvalho, I. S., Salgado, L., Lindoso, R. M., Araújo-Júnior, H. I., Nogueira, F. C. C.,
 455 Soares, J. A. 2017. A new basal titanosaur (Dinosauria, Sauropoda) from the Lower
 456 Cretaceous of Brazil. Journal of South American Earth Sciences 75, 74-84.
- 457 Cassab, R. C. T. 2003. Paleontologia da Formação Jandaíra, Cretáceo Superior da Bacia
 458 Potiguar, com ênfase na paleobiologia dos gastrópodos. Thesis Doctoral, Instituto de
 459 Geociências, Universidade Federal do Rio de Janeiro, Rio de Janeiro, 186 p.
 460 Unpublished.
- 461 Cavin, L., Tong, H., Boudad, L., Meister, C., Piuz, A., Tabouelle, J., Aarab, M., Amiot,
 462 R., Buffetaut, E., Dyke, G., Hua, S., Leloeuff, J. 2010. Vertebrate assemblages from
 463 the early Late Cretaceous of southeastern Morocco: an overview. Journal of African
 464 Earth Sciences 57, 391–412.
- 465 Contessi, M., 2009. Vertebrate remains from the Barremian–Cenomanian (Early
 466 Cretaceous) coastal deposits of South Tunisia. In: Fanti, F., Spalletta, C. (Eds.),
 467 International Conference on Vertebrate Palaeobiogeography and Continental

- 468 Bridges across Tethys, Mesogea, and Mediterranean Sea – Abstracts Book. Museo
 469 Geologico Giovanni Capellini, Dipartimento di Scienze della Terra e Geologico-
 470 Ambientali, Bologna, Italy, pp. 31–32.
- 471 Coria, R. A., Currie, P. J. 2006. A new carcharodontosaurid (Dinosauria, Theropoda)
 472 from the Upper Cretaceous of Argentina. *Geodiversitas* 28, 71–118.
- 473 Coria, R. A., Salgado, L. 1995. A new giant carnivorous dinosaur from the Cretaceous of
 474 Patagonia. *Nature* 377, 224–226.
- 475 Costa, A. B. C., Córdoba, V. C., Netto, R. G., Lima Filho, F. P. 2014. Registro faciológico
 476 e paleoambiental da transgressão que marca a passagem do Cenomaniano para o
 477 Turoniano na Bacia Potiguar, NE do Brasil. *Comunicações Geológicas*, 101(I), 415-
 478 420.
- 479 Currie, P.J., Carpenter K. 2000. A new specimen of *Acrocanthosaurus atokensis*
 480 (Theropoda, Dinosauria) from the Lower Cretaceous Antlers Formation (Lower
 481 Cretaceous, Aptian) of Oklahoma, USA. *Geodiversitas*. 22: 207 – 246.
- 482 Derby, A. O. 1890. Nota sobre a geologia e paleontologia de Mato Grosso. *Arquivos do*
 483 *Museu Nacional*. 9, 59-88.
- 484 Duarte, L. and Santos, R. 1962. Fósseis do Arenito Açú. *Anais da Academia Brasileira*
 485 *de Ciências* 34(1), 57-68.
- 486 Duarte, L., Santos, R.S. 1961. Novas ocorrências fossilíferas nos estados do Rio Grande
 487 do Norte e Ceará. *Coleção Mossoroense*, 56: 11 p.
- 488 Franco-Rosas A. C., Salgado L, Rosas C. F., Carvalho I. S. 2004. Nuevos materiales de
 489 titanosaurios (Sauropoda) em el Cretácico Superior de Mato Grosso, Brasil. *Rev*
 490 *Bras Paleont* 7, 329–336.
- 491 Frey E., Martill D. M. 1995. A possible oviraptorosaurid theropod from the Santa
 492 Formation (Lower Cretaceous, Albian) of Brazil. *N Jahrb GeolPal.ont M* 7: 397–412.

- 493 Forster, C. A., Sampson, S. D., Chiappe, L.M., Krause, D.W. 1998. The Theropod:
494 Ancestry of Birds: New Evidence from the Late Cretaceous of Madagascar. *Science*
495 16 (5358), 1915-1919.
- 496 Gauthier, J. A. 1986. Saurischian monophyly and the origin of birds; pp. 1–55 in K.
497 Padian (ed.), *The Origin of Birds and the Evolution of Flight*. *Memoirs of the*
498 *California Academy of Sciences*, 8.
- 499 Gilmore, C. W. 1920. Osteology of the carnivorous Dinosauria in the United States
500 National Museum, with special reference to the genera *Antrodemus* (*Allosaurus*) and
501 *Ceratosaurus*. *Bulletin U. S. National Museum*. 110,1-154.
- 502 Guedes, E., Heilbron, M., Vasconcelos, P. M., De Morisson Valeriano, C., Horta De
503 Almeida, J. C., Teixeira, W., Filho, A. T. 2005. K–Ar and ⁴⁰Ar/³⁹Ar ages of dikes
504 emplaced in the onshore basement of the Santos Basin, Resende area, SE Brazil:
505 implications for the south Atlantic opening and Tertiary reactivation. *Journal of*
506 *South American Earth Sciences* 18 (3–4), 371–382.
- 507 Harris, J. D. 1998. Are analysis of *Acrocanthosaurus atokensis*, its phylogenetic status
508 and paleobiogeographic implications, based on a new specimen from Texas. *Bulletin*
509 *of New Mexico Museum of Natural History and Science* 13, 1-75.
- 510 1. Hendrickx, C., Mateus, O. 2014. Abelisauridae (Dinosauria: Theropoda) from the
511 Late Jurassic of Portugal and dentition-based phylogeny as a contribution for the
512 identification of isolated theropod teeth. *Zootaxa*, 3759.
- 513 Hendrickx, C., Mateus, O., Araújo, R. 2015. A proposed terminology of theropod teeth
514 (Dinosauria, Saurischia). *Journal of Vertebrate Paleontology*, 35(5), e982797.
- 515 Kellner, A. W. A. 1996a. Fossilized theropod soft tissue. *Nature* 379, 32–32.
- 516 Kellner, A. W. A. 1996b. Remarks on Brazilian dinosaurs. *Memoirs of Queensland*
517 *Museum* 39, 611–626.

- 518 Kellner, A. W. A. 1999. Short note on a new dinosaur (Theropoda, Coelurosauria) from
519 the Santana Formation (Romualdo Member, Albian), North-eastern Brazil. Boletim
520 do Museu Nacional 49, 01-08.
- 521 Kellner, A. W. A., Azevedo, S. A. K., Carvalho, L. B., Henriques, D. D. R., Costa, T.,
522 Campos, D. A. 2004. Bones out of the jungle: on a dinosaur locality from Mato
523 Grosso, Brazil. Journal of Vertebrate Paleontology 124 (Suppl3): 78a.
- 524 Kellner, A. W. A., Azevedo, S. A., Machado, E. B., Carvalho, L. B. D. Henriques, D. D.
525 2011. A new dinosaur (Theropoda, Spinosauridae) from the Cretaceous
526 (Cenomanian) Alcântara Formation, Cajual Island, Brazil. Anais da Academia
527 Brasileira de Ciências, 83(1), 99-108.
- 528 Jokat, W., Boebel, T., König, M., Meyer, U. 2003. Timing and geometry of early
529 Gondwana breakup. Journal of Geophysical Research 108 (B9, 2428), 1–15.
- 530 Langer, M. C., Ribeiro, A. M., Schultz, C.L. Ferigolo, J. 2007a. The continental tetrapod-
531 bearing Triassic of south Brazil. Bull NMMNHS 41, 201–218.
- 532 Macdonald, D., Gomez-Perez, I., Franzese, J., Spalletti, L., Lawver, L., Gahagan, L.,
533 Dalziel, I., Thomas, C., Trewin, N., Hole, M., Paton, D. 2003. Mesozoic break-up of
534 SW Gondwana: implications for regional hydrocarbon potential of the southern
535 South Atlantic. Marine and Petroleum Geology 20 (3–4), 287–308.
- 536 Madsen, J. H. 1976. *Allosaurus fragilis*, a revised osteology. Utah geological and
537 mineralogical Survey Bulletin 109, 1-163.
- 538 Marsh, O. C. 1869. Notice of some new reptilian remains from the Cretaceous of Brazil.
539 American Journal of Science 47, 390–392.
- 540 Marsh, O. C. 1878. Principal characters of American Jurassic dinosaurs. American
541 Journal of Science 95, 411-416.

- 542 Marsh, O. C. 1881. Principal characters of American Jurassic dinosaurs, part V. American
543 Journal of Science 125, 417-423.
- 544 Martill, D. M., Cruickshank, A. R. I., Frey, E., Small, P. G., Clarke, M. 1996. A new
545 crested maniraptoran dinosaur from the Santana Formation (Lower Cretaceous) of
546 Brazil. Journal of the Geological Society 153(1), 5-8.
- 547 Martinelli, A.G., Ribeiro, L.C.B., Neto, F.M., Méndez, A.H., Cavellani, C.L., Felix, E.,
548 Ferraz, M.L.F., Teixeira, V.P.A., 2013. Insight on the theropod fauna from the
549 Uberaba Formation (Bauru Group), Minas Gerais State: new megaraptoran specimen
550 from the Late Cretaceous of Brazil. Rivista Italiana di Paleontologia e Stratigrafia
551 119, 205–214.
- 552 Mateus, O., Araújo, R., Natário, C., Castanhinha, R. 2011. A new specimen of the
553 theropod dinosaur *Baryonyx* from the early Cretaceous of Portugal and taxonomic
554 validity of *Suchosaurus*. Zootaxa 2827, 54–68.
- 555 Mawson, J., Woodward, A. S. 1907. On the Cretaceous Formation of Bahia (Brazil), and
556 on the vertebrate fossils contained therein. Journal of the Geological Society 63,
557 128–139.
- 558 Medeiros, M. A., Schultz, C. L., 2002. A fauna dinossauriana da Laje do Coringa,
559 Cretáceo médio do Nordeste do Brasil. Arquivos do Museu Nacional 60 (3), 155–
560 162.
- 561 Medeiros, M. A., Freire, P. C., Pereira, A. A., Santos, R. A. B., Lindoso, R. M., Coelho,
562 A. F. A., Passos, E. B., Sousa, E. 2007. Another African dinosaur recorded in the
563 Eocenomanian of Brazil and a revision on the paleofauna of the Laje do Coringa site.
564 In: CARVALHO I.S., CASSAB R. C. T., SCHWANKE C., CARVALHO M.A.,
565 Fernandes A. C. S., Rodrigues M. A. C., Carvalho M. S. S., Arai M. And Oliveira

- 566 M. E. Q. (Eds), *Paleontologia: Cenários de Vida*. Rio de Janeiro, Interciência 1, 413–
567 423.
- 568 Motta, M. J., Aranciaga Rolando, A. M., Rozadilla, S., Agnolín, F. E., Chimento, N. R.,
569 Egli, F. B., Novas, F. E. 2016. New theropod fauna from the upper cretaceous
570 (Huincul Formation) of Northwestern Patagonia, Argentina. *New Mexico Museum*
571 *of Natural History and Science Bulletin*, 71, 231-253.
- 572 Motta, M. J., Egli, F. B., Novas, F. E. 2018. Tail anatomy of *Buitreraptor gonzalezorum*
573 (Theropoda, Unenlagiidae) and comparisons with other basal paravians. *Cretaceous*
574 *Research*, 83, 168-181.
- 575 Mussa, D.; Oliveira, L.D.D., Barcia-Andrade, A. 1984. Fragmentos estélicos de Palmae
576 procedentes da Formação Açú (?), Bacia Potiguar, Brasil. *Boletim do IGC-USP,*
577 *Instituto de Geociências* 15, 129-141.
- 578 Naish, D.; Martill, D. M., Frey, E. 2004. Ecology, systematics and biogeographical
579 relationships of dinosaurs, including a new theropod from the Santana Formation
580 (Albian, Early Cretaceous) of Brazil. *Historical Biology* 18, 01-14.
- 581 Novas, F.E., Ribeiro, L.C.B., Carvalho, I.S., 2005a. Maniraptoran theropod ungual from
582 the Marília Formation (Upper Cretaceous), Brazil. *Revista Museo Argentino*
583 *Ciencias Naturales* 7, 31-36.
- 584 Novas, F. E., Ezcurra, M. D., Lecuona, A. 2008. *Orkoraptor burkei* nov.gen. et sp., a
585 large theropod from the Maastrichtian Pari Aike Formation, Southern Patagonia,
586 Argentina. *Cretaceous Research*, 29(3), 468-480.
- 587 Novas, F. E., Brissón Egli, F., Agnolín, F. L., Gianechini, F. A., and Cerda, I., 2017.
588 Postcranial osteology of a new specimen of *Buitreraptor gonzalezorum* (Theropoda,
589 Coelurosauria). *Cretaceous Research*.

- Ostrom, J. H. 1969. Osteology of *Deinonychus antirrhopus*, an Unusual Theropod from the Lower Cretaceous of Montana (Vol. 30) Peabody Museum of Natural History, Yale University.
- Pereira, P. V. L. G. C., Marinho, T. S., Candeiro, C. R. A., Bergqvist, L. P. 2018. A new titanosaurian (Sauropoda, Dinosauria) osteoderm from the Cretaceous of Brazil and its significance. *Ameghiniana* 55(6), 644-650. (71).
- Pereira, P. V. L. G. C., Veiga, I. M. M. G., Ribeiro, T. B., Cardozo, R. H. B., Candeiro, C. R. A., Bergqvist, L.P. 2020. The Path of Giants: A new occurrence of Rebbachisauridae (Dinosauria, Diplodocoidea) in the Açu Formation, NE Brazil, and its paleobiogeographic implications., *Journal of South American Earth Sciences*, <https://doi.org/10.1016/j.jsames.2020.102515>.
- Pessoa Neto, O. D. C., Soares, U. M., Silva, J. D., Roesner, E. H., Florencio, C. P. and Souza, C. D. 2007. Bacia Potiguar. *Boletim de Geociências da Petrobras* 15:357-369.
- Petri, S. 1987. Cretaceous paleogeographic maps of Brazil. *Palaeogeography Palaeoclimatology and Palaeoecology* 59,117–168.
- Pletsch, T., Erbacher, J., Holbourn, A., Kuhnt, W., Moullade, M., Oboh-Ikuenobede, F., Soding, E., Wagner, T. 2001. Cretaceous separation of Africa and South America: the view from the West African margin (ODP Leg 159). *Journal of South American Earth Sciences* 14, 147–174.
- Price, L. I. 1960. Dentes de Theropoda num testemunho de sonda no estado do Amazonas. *Na Academia Brasileira de Ciências* 32, 79–84.
- Price, L. I. 1961. Sobre os dinossáurios do Brasil. *Na Academia Brasileira de Ciência* 33(3–4), 28-29.
- Rauhut, O. W. M. 1999. A dinosaur fauna from the Late Cretaceous (Cenomanian) of northern Sudan. *Palaeontologia Africana* 35, 61–84.

- 615 Reyment, R., Dingle, R. 1987. Palaeogeography of Africa during the Cretaceous period.
616 Palaeogeography Palaeoclimatology Palaeoecology 59, 93-116.
- 617 Rossetti, D. F., 1997. Internal architecture of mixed tide- and storm-influenced deposits:
618 an example from the Alcântara Formation, northern Brazil. Sedimentary Geology
619 114 (1-4), 163-169. 115
- 620 Rossetti, D. F., Truckenbrodt, W. 1997. Revisão estratigráfica para os depósitos do
621 Albiano-Terciário Inferior (?) na Bacia de São Luís (MA), norte do Brasil. Boletim
622 do Museu Paraense Emílio Goeldi, Série Ciências da Terra 9, 29-41.
- 623 Russell D. A., 1972 – Ostrich dinosaurs from the Late Cretaceous of western Canada.
624 Canadian Journal of Earth Sciences 9, 375-402.
- 625 Santos, M. F. C. F.; Florêncio, C. P.; Reyes-Pérez, Y. A.; Bergqvist, L. P.; Porpino, K.
626 O.; Uchoa, A. F. and Lima-Filho, F. P. 2005. Dinossauros na Bacia Potiguar: o
627 registro da primeira ocorrência. Boletim de Resumos Expandidos do XXI Simpósio
628 de Geologia do Nordeste, Recife, 19, 325-328.
- 629 Senter, P., 2011. Using creation science to demonstrate evolution: morphological
630 continuity within Dinosauria. Journal of evolutionary biology 24(10): 2197-2216. 15.
- 631 Senter, P., Kirkland, J. I., DeBlieux, D. D., Madsen, S., Toth, N., 2012. New
632 dromaeosaurids (Dinosauria: Theropoda) from the Lower Cretaceous of Utah, and
633 the evolution of the dromaeosaurid tail. PLoSOne, 7(5), e36790.
- 634 Sereno, P. C., Martínez, R. N., Wilson, J. A., Varricchio, D. J., Alcober, O. A. 2008.
635 Evidence for Avian Intrathoracic Air Sacs in a New Predatory Dinosaur from
636 Argentina. PLoS One 3, e3303.
- 637 Sereno, P. C., Brusatte, S. L. 2008. Basal abelisaurid and carcharodontosaurid theropods
638 from the lower Cretaceous Elrhaz Formation of Niger. Acta Palaeontologica
639 Polonica 53 (1), 15-6.

- 640 Silva-Santos, R. 1963. Peixes do Cretácico do Rio Grande do Norte. Anais da Academia
641 Brasileira de Ciências 35, 67-74.
- 642 Smith, J., Lamanna, M., Lacoara, K., Dodson, P., Smith, J., Poole, J., Giegengack, R.,
643 Attia, Y. 2001. A giant sauropod from an Upper Cretaceous mangrove deposit in
644 Egypt. Science 292, 1704–1706.
- 645 Stromer, E. 1915. Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wüsten
646 Ägyptens. II. Wirbeltier-Reste der Baharije-Stufe (unterstes Cenoman). 3. Das
647 Original des Theropoden *Spinosaurus aegyptiacus* nov. gen., nov. spec.
648 Abhandlungen der Königlich Bayerischen Akademie der Wissenschaften,
649 Mathematisch-physikalische Classe, 28(3) Abhandlung, 1–32.
- 650 Stromer, E., 1931. Wirbeltierreste der Baharije-Stufe (unterstes Cenoman). 10. Ein
651 Skelett-Rest von *Carcharodontosaurus* nov. gen: Abhandlungen Bayerische
652 Akademie Wissenhafte Atheilung-naturwissenschaften Abteilung Neue Folge, 9, 1–
653 23.
- 654 Sues, H. D. 1997. On Chirostenotes, a Late Cretaceous oviraptorosaur (Dinosauria:
655 Theropoda) from western North America. Journal of Vertebrate Paleontology 17,
656 698–716.
- 657 Tello Saenz, C., Hackspacher, P., Hadler Neto, J., Iunes, P., Guedes, S., Ribeiro, L., Paulo,
658 S. 2003. Recognition of Cretaceous, Paleocene, and Neogene tectonic reactivation
659 through apatite fission-track analysis in Precambrian areas of southeast Brazil:
660 association with the opening of south Atlantic Ocean. J South Am Earth Sci 15,765–
661 774.
- 662 Tykoski R. S., 2005 – Anatomy, Ontogeny, and Phylogeny of Coelophysoid Theropods.
663 Unpublished PhD Thesis, The University of Texas at Austin, 1-553.

- 664 Vasconcelos, E. P., Lima Neto, F. F., Roos, S. 1990. Unidades de correlação da Formação
665 Açu. 36º Congresso Brasileiro de Geologia (Natal), Actas, 227-240.
- 666 Viramonte, J., Kay, S., Becchio, R., Escayola, M., Novitski, I. 1999. Cretaceous rift
667 related magmatism in central-western South America. *Journal of South American*
668 *Earth Sciences* 12, 109–121.
- 669 Werner, C., 1994. Die kontinentale Wirbeltierfauna aus der unteren Oberkreide des Sudan
670 (Wadi Milk Formation). *Berliner Geowissenschaftliche Abhandlungen Reihe E* 13,
671 221–249.
- 672 Xu X., Tan Q., Wang J., Zhao X.-J., Tan L. 2007. Gigantic bird-like dinosaur from the
673 Late Cretaceous of China. *Nature*, 447, 844-847.
- 674 Xu, X., Qin, Z. C. 2017. A new tiny dromaeosaurid dinosaur from the Lower Cretaceous
675 Jehol Group of western Liaoning and niche differentiation among the Jehol
676 dromaeosaurids. *Vertebrata Palasiatica*, 55(2), 129-144.

677

678 Figure captions

679

- 680 Figure 01: Geological map of the continental part of the Potiguar Basin with the region
681 near the Limoeiro do Norte municipality (Ceará state) where the material were discovered
682 (dark star). CE, Ceará state; RN, Rio Grande do Norte state and its capital, Natal.
683 Modified from Cassab (2003).

684

- 685 Figure 02: The theropod vertebrae UFRJ-DG 528-R (A-C) and UFRJ-DG 575-R (D-E).
686 UFRJ-DG 575-R: A, lateral view; B, the anterior articular facet. UFRJ-DG 528-R: C, the
687 lateral view; D, the ventral view; E, anterior articular facet. Note the large pneumatic

foramen on the side of the anterior fragment of UFRJ-DG 575-R. pfr = pneumatic foramen. Scale bar: 2 cm.

Figure 03: Spinosauroid tooth (UFRJ-DG 619-R): A, the labial view; B, the lingual view; and C, the cross section. Scale: 1 cm

Figure 04: Maniraptoran caudal vertebrae (UFRJ-DG 521-R): A, Lateral view; B, ventral view; C, anterior articular facet. Prz, prezygophysis; Nc, neural canal. Scale: 1cm.

Figure 05: Comparison of UFRJ-DG 521-R and other maniraptorans. A, Potiguar's material; B, *Rahonavis*; C, *Buitreraptor*; D, *Anchiornis*. Pr, prezygapophysis; lg, Longitudinal groove. Modified from Motta et al., (2018).

Figure 06: Carcharodontosaurid caudal vertebrae UFRJ-DG 523(A-C) and UFRJ-DG 524-R (D-F). UFRJ-DG 524-R: A, ventral view; B, lateral view; C, anterior articular facet. UFRJ-DG 523-R: D, ventral view; E, lateral view; F, anterior articular facet. Nc, neural canal. Scale: 1cm.

Figure 07: Comparison of UFRJ-DG 523-R and 524-R and other carcharodontosaurids. A and B, UFRJ DG 523-R; C and D, UFRJ DG 524-R; E and F, Kem Kem beds material (from Rauhut, 1999); G, *Tyrannotitan chubutensis* MPEF-PV 1156 (Modified from Canale et al., 2015); H, *Mapusaurus roseae* MCF-PVPH-108.81 (Modified from Coria and Currie, 2006) ; I, *Acrocanthosaurus atokensis* NCSM 14345 (Modified from Currie and Carpenter, 2000). Scale bar = 5 cm.

Figure 08: Caudal vertebrae UFRJ-DG. UFRJ-DG 558-R: A, posterior articular facet; B, lateral view; C, ventral view. UFRJ-DG 634-R: D, anterior articular facet; E, lateral view; F, ventral view. Pfr, Pneumatic foramen. Scale bar: 1cm.

Figure 09: Brazilian megaraptoran vertebrae findings. A and B, UFRJ DG 558-R; C and D, MPMA 08-003-94 (Méndez et al., 2012); E and F, CPPLIP 1324 (Martinelli et al., 2013). A, C e E, lateral view; B, D e F, ventral view. Pfr, Pneumatic foramen. Scale bar = 1cm.

Figure 10: Megaraptoran caudals vertebrae. A and B, UFRJ DG 558-R; C and D, *Aerosteon*; E and F, *Aoniraptor*; G, *Orkoraptor*. H, *Megaraptor*. A, C, E, G e H, lateral view; B, D e F, ventral view. Pfr, pneumatic foramen. Scale bar = 5cm.

Figure 11: Reconstruction of the theropods groups from Açú Formation, Potiguar Basin. In the center, a group of megaraptorans slaughtering a titanosaur; on the right a carcharodontosaurid awakens from its sleep; in the top center, a maniraptoran just watches. Art by Luciano da Silva Vidal.

A

B

pfr

C

D

E

2 cm

A

B

C

1 cm

Click here to access/download
RDM Data Profile XML
DataProfile_3987663.xml

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Author Statement

Paulo Victor Luiz Gomes da Costa Pereira: Conceptualization, Investigation, Writing - Original Draft, Writing - Review & Editing, Supervision, Funding acquisition.

Theo Baptista Ribeiro: Investigation. Writing - Original Draft, Writing - Review & Editing, Visualization.

Stephen Louis Brusatte: Validation, Writing - Review & Editing, Supervision

Carlos Roberto dos Anjos Candeiro: Writing - Original Draft, Writing - Review & Editing, Supervision.

Thiago da Silva Marinho: Investigation, Writing - Review & Editing

Lilian Paglarelli Bergqvist: Resources, Writing - Review & Editing, Supervision, Project administration, Funding acquisition.

Dear Editor

I, Dr. Paulo Victor Luiz Gomes da Costa Pereira, the first author of this manuscript, send this letter in order to request the change of the reviewer 01. I know that such a change can cause a delay in publication, but I believe that is important. We consider all the changes requested by you and the other reviewers and I am absolutely sure that the manuscript improved a lot with the changes proposed by you all. But I have to point out that the criticisms made by the reviewer 01 were often disrespectful and arrogant, a role that does not compete with the duties of a reviewer in a major international magazine.

His comments, did not contribute much to our work, since the reviewer limited himself to not agreeing with our classifications, always without presenting actual arguments or presenting bibliography that disagreed with our results, a totally different behavior when compared to the other two reviewers.

Our work is based on morphological description and comparison between materials, based on several similar articles with African and European fossils that have already been published, including by Cretaceous Research.

The role of our manuscript (and science in general) is to contribute as much as possible with the fossils collected. Nothing prevents better preserved fossils from disagreeing with our considerations in the future.

I'm really upset that I had to wait more than 8 months for a review that did not seek to improve the manuscript at all.

I thank you very much for the attention given by you and the other reviewers to our manuscript. Be sure that our work has improved a lot with your contributions!

Yours sincerely,

Dr. Paulo Victor Luiz Gomes da Costa Pereira

**THEROPOD (DINOSAURIA) DIVERSITY FROM THE POTIGUAR BASIN
(EARLY-LATE CRETACEOUS ~~ALBIAN~~– CENOMANIAN), NORTHEAST
BRAZIL**

**Paulo Victor Gomes da Costa Pereira^{1*}; Theo Baptista Ribeiro¹; Stephen Louis
Brusatte²; Carlos Roberto Dos Anjos Candeiro³; Thiago da Silva Marinho⁴; Lilian
Paglarelli Bergqvist¹**

*corresponding author – paulovictor29@yahoo.com.br

1- CCMN - Universidade Federal do Rio de Janeiro, Ilha do Fundão, Rio de Janeiro (RJ), Brazil.

E-mail: paulovictor29@yahoo.com; theobribeiro1@gmail.com; bergqvist@geologia.ufRJ.br

2 - University of Edinburgh, Grant Institute, School of GeoSciences. Edinburgh, Scotland, UK. E-mail:
stephen.brusatte@ed.ac.uk

3 - Universidade Federal de Goiás (UFG), Laboratório de Paleontologia e Evolução (LABPALEOEVO),
Curso de Geologia. Aparecida de Goiânia, GO, Brazil. E-mail: candeiro@ufg.br

4- Centro de Pesquisas Paleontológicas L. I. Price, Complexo Cultural e Científico Peirópolis, Pró-Reitoria
de Extensão Universitária, Universidade Federal do Triângulo Mineiro, Uberaba, Minas Gerais, Brazil. E-
mail: jsmarinho@icene.ufm.edu.br

Abstract

The theropod records from the Cretaceous of northeastern Northeast of Brazil are rare
and consist mostly of isolated and incomplete remains, with only ~~four~~ ^{five} species
described. Here we describe, identify and evaluate the diversity of theropod materials
from the Albian-Cenomanian Açú Formation, Potiguar Basin. The material consists of
~~seven~~ ^{nineteen} isolated ~~theropod~~ ^{theropod} vertebrae (~~UFRI-DG 521 R, 523 R, 524 R, 528 R, 532~~

Commented [PP1]:
Revisor 01: I recommend to change the title for a more realistic one, for example: "Theropod dinosaur remains from..."
A: We understood the statement of Revisor 01, but we preferred to keep the original title, as it summarizes the results seen in the manuscript.

Commented [BS2R1]: I actually agree with the reviewer. Because we are describing a series of new fossils, 'Theropod dinosaur remains' sounds better. But this is your choice.

Formatted: English (United States)

Formatted: English (United States)

Field Code Changed

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Field Code Changed

Commented [3]: Five theropod species only in the northeast, or in all Brazil?

Commented [LM4]: A: The four specimens are only from northeastern Brazil. This detail is better explained in the second paragraph of the Introduction.

Formatted: No underline, Font color: Auto, English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: No underline, Font color: Auto, English (United States)

R, 547 R, 558 R, 575 R, 587 R and 589 R) and a tooth (UFRI-DG-653 R). ~~FWe identify~~
~~the material, have been identified~~ as belonging to ~~four~~ theropod groups: ~~Abelisauria,~~
~~Carcharodontosauria,~~ Spinosauroidae, ~~Carcharodontosauriidae,~~ Megaraptora, and
~~Maniraptora.~~ The vertebrae were classified into five ~~four~~ morphotypes based on
~~morphological form and/or diagnostic characters and comprised represented at this~~
~~moment by five four groups:~~ ~~Abelisauria,~~ ~~Carcharodontosauria,~~
~~Spinosauridae,~~ Megaraptora, and ~~Maniraptora.~~ ~~We classify the~~ The isolated tooth was
~~classified as belonging to a spinosaurid.~~ One of the ~~significant~~ ~~great~~ results ~~so far~~ is the
occurrence of Megaraptora in the Potiguar Basin; ~~i~~ based on the general morphology,
~~some of the bone~~ ~~elements described we describe~~ are very similar to ~~those of~~ ~~Aerosteon~~
and ~~Megaraptor.~~ Another ~~unexpected~~ ~~remarkable~~ result is the ~~description~~ ~~identification~~
and presence of ~~a maniraptoran~~ ~~a~~ caudal vertebrae ~~of a maniraptoran~~; ~~these dinosaurs are~~
very rare in Brazil, with few ~~fossils~~ ~~remnants~~ ~~previously~~ described. ~~Besides~~
~~this~~ ~~Furthermore, we identify~~ other groups ~~that have~~ already ~~been~~ found ~~on-in~~ isochronous
basins of the Northeast region of Brazil and Africa, ~~including as~~ Carcharodontosauria and
Spinosauridae. The presence of these theropod groups ~~in the~~ ~~at~~ Açú Formation reveals
an ~~unexpected~~ dinosaur richness ~~fauna~~ ~~in the~~ ~~in the~~ Potiguar Basin ~~similar to~~
isochronous basins in Northern Africa and ~~opens up an important opportunity to increase~~
~~the~~ knowledge about ~~the~~ ~~the~~ diversity of South American dinosaurs.

Commented [B55]: Five groups are listed

Formatted: No underline, Font color: Auto, English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Commented [6]: Spinosauridae was recognized by a tooth, not a vertebra

Formatted: No underline, Font color: Auto, Strikethrough

Commented [7]: I suggest to end this sentence with something more "... about the diversity of dinosaurs xxx..."

Commented [PP8]: A: Done

Formatted: No underline, Font color: Auto

Formatted: English (United States)

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Keywords: Dinosauria, Potiguar Basin, Theropoda, faunistic richness, ~~Abelisauria,~~
Carcharodontosauria, Megaraptora, Spinosauridae, ~~and~~ Maniraptora.

Introduction

The first ~~works~~ ~~studies~~ in Brazil that attributed vertebrate fossil remains to

dinosaurs were ~~conducted-published into~~ the nineteenth and mid-twentieth centuries (Marsh, 1869; Derby, 1890; Mawson ~~&and~~ Woodward, 1907; Price, 1960, 1961). Since that time, dinosaur fossils have been recorded ~~in-from~~ three principal localities/ and ages in Brazil: the Triassic of the Santa Maria and Caturrita formations (Langer et al., 2007a), the mid-Cretaceous of the Araripe, Triunfo and São Luís-Grajaú basins (Frey ~~&and~~ Martill 1995; Kellner 1996a, b, 1999; Medeiros *et al.*, 2007; Carvalho *et al.*, 2017), and the Late Cretaceous of the Bauru and Parecis groups (Franco-Rosas *et al.*, 2004; Kellner *et al.*, 2004; Brusatte *et al.* 2017).

~~As of now, there have been five theropod dinosaur species formally described from Brazil: *Staurikosaurus pricei* Colbert, 1970 and *Guaibasaurus candelariensis* Bonaparte, Ferigolo & Ribeiro, 1999 from the Caturrita Formation; *Angaturama limai* Kellner & Campos, 1996, *Santanaraptor placidus* Kellner, 1999, *Irritator challengeri* Martill, Cruikshank, Frey, Small and Clarke, 2002 and *Mirischia asymmetrica* Naish, Martill & Frey 2004 from the Araripe Basin; *Oxalaia quilombensis* Kellner, Azevedo, Machado, Carvalho & Henriques, 2011 from the São Luís Grajaú Basin, and *Pycnonemosaurus nevesi* Kellner & Campos, 2002 from the Bauru Group. There are eight theropod dinosaur species formally described from -for- Brazil so far, four of those described for its from the northeastern region: *Santanaraptor placidus* Kellner, 1999, *Irritator challengeri* Martill, Cruikshank, Frey, Small and Clarke, 2002 and *Mirischia asymmetrica* Naish, Martill and Frey 2004 from the Araripe Basin; and *Oxalaia quilombensis* Kellner, Azevedo, Machado, Carvalho and Henriques, 2011 from the São Luís-Grajaú Basin.~~

A promising area for new dinosaur discoveries is ~~the rocks of the~~ *Açu Formation*, ~~in the~~ *Portuguar Basin*. ~~Recently~~ *Until now*, the macrofossils of the *Açu Formation* ~~(in this~~

Commented [PP9]: Revisor 01: Please, restrict to Cretaceous record, to address the relevance of the materials here reported.

A: We choose to keep this part of the text to preserve the cohesion within of the Introduction, featuring the first works and discoveries on theropod paleontology of Brazil. The next paragraph has a small summary of the Cretaceous northeastern theropod dinosaurs of Brazil.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Commented [10]: To this list should be added the recently described *Vespersaurus paranaensis* Langer et al., 2019 and *Gnathovorax cabreirai* Pacheco et al., 2019

Commented [LM11]: We choose to limit this list to the northeastern Brazil's theropod record only to make the text more easy to read/understand

area were restricted to a few occurrences from outcrops on the western border of the basin (Russas and Tabuleiro do Norte municipalities, Ceará state), consisting of bivalve molluscs, small crustaceans, fish scales, and plant remains (Duarte and Santos, 1961). This situation lasted until However this changed in the 2000s, when researchers from the Group of Analogs to Oil Reservoirs of the Department of Geology of the Federal University of Rio Grande do Norte, in geological mapping of the Açu 4 operational unit, found large vertebrate fossils. These fossils were attributed to Titanosauria and Theropodindet., but were not described in detail (Santos et al., 2005).

In the decade after the discovery of these first continental vertebrate fossils in the formation (Santos *et al.*, 2005), no other fieldwork was conducted. However, in the years 2015 and 2016, this area was again prospected by Laboratório de Macrofósseis of the Universidade Federal do Rio Janeiro and dozens of fossils were collected. The aim of the present work is to describe and identify the collected materials attributed to theropod dinosaurs, showing that the Potiguar Basin preserves a great-large diversity of species and has great potential for future discoveries and studies about the mid-Cretaceous paleoenvironments of the Atlantic margin of Brazil.

Geological Setting and Lithostratigraphy

The Potiguar Basin is located at the eastern continental margin of northeastern Brazil, cropping out in the states of Rio Grande do Norte and Ceará (Fig. 01), with a total estimated area of 60,000 km², of which 22,000 km² is interpreted as continental (Cassab, 2003). The Potiguar Basin is bounded to the east by Alto de Touros, which which separates it from the Pernambuco-Paraíba Basin, to the northwest by the Alto de

Formatted: Font: Italic

Formatted: Line spacing: Double

Commented [BS12]: You need the which

Fortaleza, which ~~bordering-separates~~ it from the Ceará Basin, and to the south and west by crystalline basement rocks (Pessoa-Neto *et al.*, 2007).

~~Figure 01: Geological map of the continental part of the Potiguar Basin with the region near the Limoeiro do Norte municipality (Ceará state) where the osteoderm were was discovered (dark star). CE, Ceará state; RN, Rio Grande do Norte state (Capital is Natal). Modified from Cassab (2003).~~

The ~~sedimentary units of the~~ Potiguar Basin ~~is are~~ divided into three groups: Areia Branca (Pendência and Alagamar formations), Apodi (Açu, Quebradas and Jandaíra formations) and Agulha (Ubarana, Guamaré and Tibau formations) (Araripe ~~& and~~ Feijó, 1994). The Açu formation is divided into four subunits according to electric logs, identified from bottom to top as Açu 1, Açu 2, Açu 3 and, Açu 4 (Vasconcelos *et al.*, 1990). The material described here comes from the Açu 4 subunit, which corresponds to a transgressive, coastal-estuarine system.

The Açu-4 Unit consists of sixteen facies, fourteen being siliciclastic and two being hybrid. The siliciclastic facies ~~were are~~ grouped into nine associations, namely: (1) lag residual deposits, (2) channel fill deposits, (3) crevasse-splay deposits, (4) floodplain deposits, (5) abandoned channel deposits, (6) upper-flow regime sandflat deposits, (7) lower-flow regime sandflats, (8) sandflat/mudflat deposits of restricted environment, and (9) mudflat deposits. The first five facies associations represent a meandering fluvial system with tidal influence, and the remaining integrate the intermediate and distal sectors of an estuarine complex dominated by tides. The hybrid facies were deposited in a shallow platform adjacent to an estuary (Costa *et al.*, 2014).

Commented [13]: The authors do not describe any osteoderm in the text.

Commented [LM14]: Done.

Formatted: English (United States)

Formatted: Indent: First line: 0.49"

Commented [PP15]: Revisor 01: Do you mean the sedimentary infilling is divided into three groups?
A: Yes, we are following the terminology and geological description of the Basin.

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Italic

The Açú Formation has been suggested to be Albian-Cenomanian in age (Early–Late Cretaceous), based on palynological data (Araripe ~~&and~~ Feijó, 1994).

Material and Methods

The fossils were collected from outcrops of ~~the~~ Açú Formation, Potiguar Basin (Ceará state, northeastern Brazil) and are deposited at the Fossil Reptile Collection of the Departamento de Geologia (DG), Universidade Federal do Rio de Janeiro (UFRJ). The material consists of ~~40-seven~~ isolated ~~theropod~~ vertebrae (UFRJ-DG 521-R, 523-R, 524-R, 528-R, ~~532-R~~, ~~547-R~~, 558-R, 575-R, ~~587-R~~ and ~~589634-R~~) and a ~~tooth (619-Rd)~~ ~~tooth~~. The vertebrae were classified into ~~five~~ ~~six~~ morphotypes based on morphological form and/or diagnostic characters.

The following tooth characteristics were ~~analyzed~~ ~~assessed~~, ~~according~~ ~~following~~ ~~to~~ the nomenclature proposed by Hendrickx *et al.* (2015): ~~—describing:~~ general morphological ~~traits~~ ~~ogy~~ of the dental crown (~~geometric shape, relative curvature and surface ornamentation~~ its overall shape, curvature, ornamentations in the enamel), denticles (presence, size and shape) ~~and~~, cross section (compression and shape), orientation of the tooth (lingual, labial, mesial and distal) and measurements ~~and blood~~ grooves (presence and visibility) (Currie *et al.*, 1990; Sankey *et al.*, 2002; Smith *et al.*, 2005; Candeiro, 2007);.

~~AL: Maximum apicobasal extent, of the tooth crown mesial base, measured from the mesial portion at the level of the cervix to the apical most point of the crown (Smith *et al.*, 2005).~~

~~CBL: Maximum mesiodistal extent of the tooth crown at the level of the cervix (Smith *et al.*, 2005). Equivalent to FABL used by some authors (Currie *et al.*, 1990; Farlow *et al.*, 1991; Sankey *et al.*, 2002).~~

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Commented [16]: I suggest to add its collection number

Commented [LM17]: Done.

Commented [18]: They were described only five morphotypes in the text.

Commented [LM19]: Done.

Formatted: English (United States)

Formatted: Font: Italic

Commented [PP20]: Revisor 01: Considering that this is not a ms devoted to analyze theropod teeth as a whole, and taking into account that just only one tooth is described, I suggest to remove all these considerations. A: Done.

Formatted: No underline, Font color: Auto, English (United States)

Formatted: Indent: First line: 0"

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Formatted: English (United States)

CBW: Maximum labiolingual extent of the tooth crown base, perpendicular to the CBL and at the level of the cervix. (Smith *et al.*, 2005).

Formatted: No underline, Font color: Auto, English (United States)

CBR: Ratio expressing the narrowness, the lateral compression of the base of the crown, corresponding to the quotient of CBW by CBL ($CBR = CBW/CBL$, Smith *et al.*, 2005). A strongly labiolingually narrow crown has a quotient of less than 0.4; a moderately narrow tooth is around 0.5-0.6; a weakly narrow crown, with an ovoid cross section, has a ratio fluctuating between 0.6-0.7; and a tooth with a subcircular transversal section has a ratio between 0.9 and 1.1 (Smith *et al.*, 2005).

Formatted: No underline, Font color: Auto, English (United States)

CH: Maximum apicobasal extent of the distal margin of the crown (Smith *et al.*, 2005). Equivalent to the TCH proposed by Farlow *et al.* (1991).

Formatted: No underline, Font color: Auto, English (United States)

CHR: Ratio expressing elongation, the relative size of the tooth, equivalent to the quotient of CH by CBL ($CHR = CH/CBL$, Smith *et al.*, 2005). A short crown tooth has a quotient less than 1.5; a medium crown tooth has a quotient varying from 1.5-2.5 and a strongly elongated crown has a ratio above 2.5.

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

DC: Number of denticles on the distal carina at mid crown per 5mm (Smith *et al.*, 2005). Equivalent to five times the posterior medial carina denticles per millimeter (Buckley *et al.*, 2010). In this study, teeth with less than 20mm had their denticles measured over 1mm, with this value then multiplied by five.

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

MC: Number of denticles on the medial part of the mesial carina per 5mm (Smith *et al.*, 2005). Corresponds to five times the number of denticles of the medial part of the mesial carina per millimeter (Buckley *et al.*, 2010). In this study, teeth smaller than 20mm had their denticles measured over 1mm, and this value was then multiplied by five.

Formatted: English (United States)

Formatted: No underline, Font color: Auto, English (United States)

Anteroposterior basal length—commonly referred to as FABL (fore aft basal length)—a measure taken between the most extreme points of the tooth at its base. It is conceptually represented by a straight line.

Formatted: No underline, Font color: Auto, Strikethrough

Formatted: Strikethrough

175 ~~▲ Total crown height — ALT — the distance between the apex and the base of the~~
176 ~~crown.~~
177 ~~▲ Relative crown height — IAR — division of the total crown height by~~
178 ~~baselineanteroposterior length (ALT/FABL).~~
179 ~~▲ Transverse section thickness — EST — measure of the labial lingual thickness of~~
180 ~~the basal cross section of the tooth.~~
181 ~~▲ Density of denticles — DDA (anterior carina, also known as mesial carina) — DDP~~
182 ~~(posterior carina, also known as distal carina). Measurement of the number of denticles~~
183 ~~per 1 mm on the medial part of the crown, in both carinae.~~

185 Terminology

186 In the present work, the terminologies proposed by Smith et al., (2005) were used
187 to refer to both the dental structures and the positioning of the analyzed teeth. According
188 to these authors, the dental crown is divided into three parts: apical (the most distal part
189 of the tooth), medial (region between the apical and basal), and basal (lower part of the
190 tooth, closer to the root). The authors defined as the labial surface the surface that was in
191 contact with the animal's lips, and the lingual surface to that which faced the tongue of
192 the animal.

194 Results

195 Dental element:

196 ~~Sistematic~~Systematic paleontology

197 Tooth

198 THEROPODA Marsh, 1881

Formatted: No underline, Font color: Auto, Strikethrough

Formatted: Strikethrough

Formatted: No underline, Font color: Auto, Strikethrough

Formatted: Strikethrough

Formatted: No underline, Font color: Auto, Strikethrough

Formatted: Strikethrough

Formatted: No underline, Font color: Auto, Strikethrough

Formatted: Strikethrough

Commented [21]: I suggest to be consistent with the abbreviations, using their English versions as is usual besides FABL: CBW: crown basal width (rather than EST), CH: crown height (rather than ALT), etc. See Hendrickx et al 2015. The dentition of megalosaurid theropods. Acta Paleontologica Polonica 60 (3): 627-642.

Commented [LM22]: Done. We updated the method and nomenclatures according to Hendrickx et al., 2015 A proposed terminology of theropod teeth (Dinosauria, Saurischia). We also cut some redundant parts of the methods, such as the terminology.

Commented [23]: The values of most of these parameters were not specified in the analysis of the tooth recovered. I suggest to add them.

Formatted: Strikethrough

Formatted: Strikethrough

Commented [LM24]: The values of the measurements are now stated in the description of UFRJ 654-Rd.

Formatted: Strikethrough

Formatted: English (United States)

Formatted: Justified, Right: 0"

Formatted: Font: 14 pt, Bold, No underline, English (United States)

Formatted: Font: 14 pt, Bold, No underline, English (United States)

Formatted: Font: 14 pt, Bold, Not Italic, No underline, English (United States)

Formatted: Font: Bold, Not Italic, English (United States)

Formatted: Right: 0", Space After: 10 pt, Line spacing: Multiple 1.15 li

Formatted: Font: Bold

~~TETANURAE~~ Gauthier, 1986

~~SPINOSAURIDAE~~ Stromer, 1915

~~Referred material:~~ UFRJ DG 653 R, a tooth.

~~Description and comparisons:~~

The material consists of an incomplete Spinosauridae crown (Fig. 02, A and B) whose apex has been lost. The crown has a high relative height (HR = 2.4) and is lingually curved. Its cross section is rounded with a Crown Base Ratio (CBR) of 7.8 (Smith et al., 2005).

This crown does not have denticulation in any of its carinae is shared with *Irritator challengeri* and *Spinosaurus aegyptiacus* but differs from the denticulated carinae of *Baryonyx walkeri* (Mateus et al., 2011). On the labial surface, there are no striations, while on the lingual surface there are ca. 9 well defined striations that bifurcate near the base of the crown.

UFRJ DG 653 R d shares many characteristics seen in other spinosaurines as a conical teeth crown with a ovoid shaped crown base, feature usually seen in piscivorous animals; the non serrated carinae, which differentiates it from the other spinosaurid family, the Baryonychinae, whose teeth features a large number of small sized serrations on both of its carinae; flutes (Must see the material again).²

~~Axial elements:~~

Commented [25]: This measurement has a big fault, given the recovered tooth lacks most of its tip.

A: Done. We added to the text that we estimated the total height of the crown based on the works of (CITE)

Commented [26]: Other abbreviation was used in Material and methods

A: Done

Commented [27]: This ratio was not mentioned in Materials and methods

A: Done

Commented [28]: I suggest that, before comparing this material with other spinosaurids, the authors should add a summary of characters that allow to assign it to Spinosauridae.

A: Done. We summarize the characteristics of spinosaurids in the Discussion.

Commented [29]: I suggest to change to "...carinae, a shared condition..."

A: Done.

Formatted: Highlight

Commented [PP30]: A: Review the tooth description and explain why

224 SAURISCHIA Cite Seeley, 1888 year Morphotype 1:

225

226 THEROPODA Marsh, 1881

227 AVEROSTRA Paul, 2002

228

229 TETANURAE Gauthier, 1986

230 NEOVENATORIDAE Benson, Carrano & Brusatte, 2010

231 ? MEGARAPTORA Benson, Carrano & Brusatte, 2010

232

233

234 Referred material: UFRJ-DG 532-R and 575-R and 587-R

235 *Description and comparisons:*

236 Morphotype 1 consists of two specimens (Fig. 02, C-F) based on partial centra

237 and is characterized by semicircular articular surfaces and high pneumaticity. They are

238 possibly caudal vertebrae, but a more conclusive description is difficult because of the

239 poor preservation of the material.

240

241

242

243 Figure 02: Tooth and vertebrae attributed to Morphotype 01: Spinosauridae tooth,

244 A; tooth cross section, B; Articular view, C(UFRJ DG 587 R); D (UFRJ DG 575 R).

245 Lateral view, E and F(UFRJ DG 575 R). Note the large pneumatic foramen on the side

246 of the anterior fragment. pfr = pneumatic foramen. Scale bar: 1cm.

247 UFRJ-DG 528-R

Formatted: Font: Bold, No underline

Formatted: No underline

Formatted: Centered

Formatted: Font: Bold

Formatted: Left

Formatted: English (United States)

Commented [31]: Given there are two morphotypes assigned to Megaraptora, both showing similar characteristics, the authors should explain why there are not included in a single morphotype.
A: Done.

Formatted: English (United States)

Formatted: Centered

Formatted: English (United States)

Formatted: Font: Italic

Formatted: Indent: First line: 0.49"

Formatted: Indent: First line: 0.49"

Commented [32]: Which view?
A: FAZER

Commented [33]: In the figure there is a black circle, I strongly suggest to add the real basal cross-section picture.
A: FAZER

Commented [34]: Those are lateral views of the same fragment, or one of each fragment?
A: FAZERPAULO

Formatted: English (United States)

Formatted: English (United States)

Specimen 528-R is a theropod vertebral centrum (Fig. 023, ~~CA-EB~~~~C~~). It is amphicoelous, and slightly higher than long. Its lateral surface is smooth and slightly concave, without marks or other ~~remareable~~~~remarkable~~ —characteristics, giving the vertebra a straight and ~~cleansomewhat~~ ~~featureless~~ appearance. The ventral surface smooth with no groove or keel and ~~is not very concave~~ it is slightly concave in lateral view, ~~being smooth on the medial region and presenting a groove on the posterior region close to the articular face margin.~~

The dorsal surface possesses a distinct longitudinal groove extending from one articular facet ~~end~~ to the other that, ~~and this groove~~ can be identified as the neural canal. ~~The anterior part of this groove is covered, but this covering is lost from the medial part further posteriorly, exposing the neural canal of this region in dorsal view.~~

The articular faces have ~~almost~~~~nearly~~ straight margins. The anterior facet is somewhat concave, and the posterior is slightly convex and slightly oval in shape; both articular facets have the same general proportions (height longer than length). The anterior articular face presents a ~~deeper concavity~~, and is slightly larger in size, ~~than~~ the posterior face, which is very flat and without deep depressions.

UFRJ-DG 575-R

Specimen 575-R (Fig. 02, ~~AC-BED~~~~F~~) is a theropod vertebral centrum broken in two: a ~~4,5cm long~~ smaller anterior piece and a ~~5,2cm long~~ larger posterior section. ~~Although the material have been~~~~was~~ found associated there is no clear point of junction between both pieces, ~~with~~~~as~~ most of the middle portion ~~being~~~~has been~~ lost. The anterior fragment ~~presents~~~~exhibits~~ a very concave articular face of semi-circular shape and slightly forward-protruding margins.

Formatted: English (United States)

Commented [35]: The word "expressive" seems wrong in this context.
A: Done.

Formatted: Indent: First line: 0.49"

Commented [PP36]: Revisor 01: Before description of any dinosaur bone, it must be glued.
A: Done.

On the lateral surface of the anterior fragment there is a deep perforation close to the dorsal region that reaches the other lateral surface, which can be described as a pleurocoel ~~(=pneumatic foramen) (=pneumatic foramen), (as the left lateral surface close to the dorsal surface is missing a piece, exposing the internal part of the bone).~~ The ventral surface of the anterior fragment is smooth and concave in anterior view. The dorsal surface of the anterior fragment is broken, missing most of the surface above the pleurocoel.

The posterior fragment has a slightly smaller articular surface, ~~which is~~ broken on the anterior portion; it is also concave and of semi-spherical circular shape, with slightly backwards-protruding margins. Its dorsal surface and the dorsal half of the left lateral surface are broken, while the right lateral surface is broken in a slightly more dorsal region in comparison to the left one. The ventral surface of the fragment is smooth and concave in lateral view. Due to the highly fragmentary state of UFRJ-DG 575-R, it is possible to see multiple small pervasive pneumatic chambers, the camellae, in the internal bone.

When both fragments are joined, it is clear that the length of the centrum is even larger than the height on its most complete point, in this case the anterior articular face. This is common in anterior dorsal vertebrae of Allosauroidae (measured next to the pleurocoels, which confirms the more anterior location). Furthermore, the margins of the articular facets have forward protrusions like those found in this group (Gilmore, 1920; Madsen, 1976).

Comparisons:

Commented [37]: Sem-spherical or semicircular? Done.

Commented [38]: The authors should describe (and figure) the materials as a single vertebra, not as two different fragments. It is confusing and does not help with interpretation of the materials.

Commented [39]: At the beginning of this morphotype description the authors suggest that they are possibly caudal vertebrae.

Formatted: Font: Italic

Formatted: Indent: First line: 0"

Formatted: Font: Italic

Formatted: Font: Italic

The highly pneumatized camellate bone seen in UFRJ-DG 575-R is a characteristic seen in many groups of theropods, from the basal *Ceratosaurus* to tetanuran groups such as carcharodontosaurids and coelurosaurs mainly in its presacral vertebrae (Carrano & Sampson, 2008). This feature, together with the poor preservation of this specimen, which prevents the identification of other more diagnostic characteristics, hinders the classification of this specimen beyond *Averostra* Theropoda.

Formatted: Font: Italic

UFRJ-DG587-R

Formatted: Indent: First line: 0.49"

Specimen 587-R is the anterior articular face of a theropod vertebra (Fig. 02, C). It has an semicircular shape, is slightly taller (9,5 cm) than wide (8cm), and has a shallow concavity on its articular surface, possibly indicating it is the anterior face of an amphicoelous vertebra, characteristic of Theropoda. The dorsal surface is indented with a concavity representing the beginning of the neural canal.

Commented [40]: Why anterior? It could not be posterior?

Commented [41]: Why the authors assign this material to morphotype 1? They only suggest that it belongs to a theropod.

Morphotype 2:

THEROPODA Marsh, 1881

TETANURAE Gauthier, 1986

CERATOSAURIA Marsh 1884

Formatted: English (United States)

NEOCERATOSAURIA Novas 1989

ABELISAURIA Novas 1992

Referred material: UFRJ DG 528 R and 532 R.

Description and comparisons:

Formatted: Font: Italic

Morphotype 2 (Fig. 03) includes two vertebrae with a low length/height ratio and semicircular articular faces with little lateral compression.

UFRJ-DG528-R

Specimen 528-R is a theropod vertebral centrum (Fig. 03, A-C). It is amphicoelous, and slightly higher than long (8,5 vs. 6cm). Its lateral surface is smooth and slightly concave, without marks or characteristics of note, giving the vertebra a straight appearance. The ventral surface is not very concave in lateral view, being smooth on the medial region and presenting a breach groove on the posterior region close to the articular face margin.

The dorsal surface possesses a distinct longitudinal groove extending from one articular end to the other, and this groove can be identified as the neural canal. The anterior part of this groove is covered, but this covering is lost from the medial part further posteriorly, exposing the neural canal of this region in dorsal view.

The articular faces have almost straight margins. The anterior face is somewhat concave, and the posterior is slightly convex and slightly oval in shape; both articular faces have the same general proportions (height longer than length). The anterior articular face presents a more expressive concavity, and is slightly larger in size than, the posterior face, which is very flat and without deep depressions.

UFRJ-DG532-R

It is a fragment of a theropod vertebral² articular facet (Figure reference?). Its oval shape in the dorso-ventral direction, due to its height being longer than its length (8cm vs. 6cm), indicates a position in the most proximal region of the caudal vertebrae. The articular surface is slightly concave, presenting only a small depression, which indicates it is the

Commented [42]: This character used for differentiate this morphotype is clearly related to the position of the vertebra inside the vertebral series.

Formatted: Indent: First line: 0"

Commented [43]: This character is used for differentiate this morphotype is also present in morphotype 1

Commented [44]: I suggest to look for other more specific characters to differentiate this morphotype

Formatted: Right: 0.02"

Formatted: Indent: First line: 0"

Commented [45]: This sentence seems wrong, please rephrase
A: Done.

Commented [46]: The word "expressive" seems wrong in this context.
A: Done.

Formatted: Indent: First line: 0"

posterior articular face of the centrum (Bonaparte, 1985; Sampson et al. 1998; Coria et al. 2002, Méndez, 2014 year?). The articular surface is slightly damaged on its lower right part, missing a fragment that goes almost up to the medial part of the articular face.

THEROPODA Marsh, 1881

TETANURAE Gauthier, 1986

? SPINOSAUROIDEA Stromer, 1915

Referred material: UFRJ-DG 619-Rd.

Description:

UFRJ-DG 619-Rd (Fig. 03) is a fragment of a large isolated tooth crown, probably belonging to the middle to almost apical portion of the tooth. The specimen lacks any form of enamel, ~~having as it has all its~~ dentine exposed, what prevents ~~the possibility to~~ description of ~~describe any kind of external~~ ornamentation ~~as such as~~ transversal undulations, ~~and~~ flutes and denticulation. The crown is almost completely straight with only a subtle curvature in its lingual surface, while the labial surface remains slightly convex.

The crown fragment has an overall cone-like shape with an almost ovoid cross section. In ~~the~~ basal view, ~~it's~~ it is possible to see concentrically deposited rings of dentine surrounding a small depression, which probably represents the apical-most portion of the dental pulp cavity.

Comparisons:

UFRJ-DG 619-Rd have some characteristics that it shares with the highly specialized teeth ~~seen in of~~ spinosauroid theropods. The most salient of these is the almost straight

Formatted: English (United States)

Formatted: Font: Not Bold, Italic

Formatted: Indent: First line: 0.49"

Formatted: Font: Italic

Formatted: Font: Italic

370 conical shaped crown, with an ovoid cross section ~~shape~~, a feature ~~usually~~ often seen in
371 piscivorous animals (~~CITE~~ Mateus, 2011; Hendrickx & and Mateus, 2014) ~~being the most~~
372 ~~remarkable of those.~~

Commented [BS47]: Which shape? Cross sectional?

374 ~~The lack of denticulation in on any of its carina is a characteristic seen in spinosaurines~~
375 ~~such as Irritator challengeri and Spinosaurus aegyptiacus (cite~~ Stromer, 1915; Martill et
376 ~~al., 1996) while the highly denticulated carina is characteristic of baryonychines as~~
377 ~~Baryonyx walkeri (Charig & Milner, 1986; Mateus et al., 2011). However, due to the lack~~
378 ~~of enamel in this material it is not possible to distinguish between Baryonychinae and~~
379 ~~Spinosaurinae nor its possible.~~

Commented [BS48]: This paragraph is not necessary because you cannot assess the denticulation. So, just delete it.
A: Done

Formatted: Indent: First line: 0", Right: 0.02"

382 Morphotype 3:

384 **THEROPODA** Marsh, 1881
385 **TETANURAE** Gauthier, 1986
386 **MANIRAPTORA** Gauthier, 1986

Formatted: Portuguese (Brazil)

391 Referred material: UFRJ-DG 521-R

Formatted: Centered

Formatted: Portuguese (Brazil)

Formatted: English (United States)

Formatted: Left

Formatted: Font: Italic

Formatted: Left, Indent: First line: 0.49"

Formatted: Left, Indent: First line: 0"

392 Description: and comparisons:

393 Morphotype 3 (Fig. 03, ~~E?~~ F-G) consists of one vertebra whose ratio between height
394 and its length somewhat greater than six and dorsally positioned prezygapophyses.

length is more than twice its height, and which lacks any processes, indicating to be a distal caudal vertebra.

Commented [49]: Again, using characters related to the vertebral position in the caudal series is

Formatted: Left, Indent: First line: 0.49", Right: 0.02"

UFRJ-DG 521-R

Specimen UFRJ-DG 521-R (Fig. 04) is an almost complete distal caudal vertebrae of a paravianmaniraptoran theropod. It is amphicoelous with a length to height ratio of almost 2.5, making it a least twice longer than tall. The dorsal surface of the centrum is almost complete with half of a dorsal midline ridge reminiscent of reduced neural spine, a well preserved and more dorsally positioned pre-zygapophysis, and a lost post-zygapophysis. The pre-zygapophysis articular surface is ellipsoid and is reclined 45° laterally. The neural canal is almost completely preserved, having lost only its posterior half.

Formatted: English (United States)

Formatted: English (United States)

Commented [BS50]: A paravian? Can you be this specific? If so, this should go in the Systematic Palaeontology section. If not, then just call it a maniraptoran here.

Formatted: English (United States)

Formatted: English (United States)

The lateral surfaces of the centrum are mostly smooth, marked only with a midline ridge reminiscent from of the reduced transverse processes of the vertebrae. The ventral surface of the centrum has a shallow groove that goes extends from one articular facet to the other. In the lateral view the ventral surface is slightly concave.

The articular facets of the centrum are both concave, with the anterior facet being more excavated than the posterior facet, and have a semi-circular shape. The articular margins are almost straight, with the anterior margin being larger than the posterior margin.

Specimen 521-R is a distal caudal vertebra of a theropod. It is almost complete, damaged only in the postzygapophysis region. It is a biconcave, amphicoelous vertebrae with a centrum slightly shorter anteroposteriorly than twice the height near the neural spine remnant and diapophysis, which confirms its more distal position within the caudal series (but not so distant from a medial position).

Formatted: Indent: First line: 0"

Formatted: Subscript

Commented [51]: I think both are synonyms
A: Done.

On the dorsal surface, the neural canal and the prezygapophyses are well preserved on the anterior part of the vertebra, being damaged only on in medial posterior portion. The prezygapophyses are intact. The lateral surfaces of the centrum are smooth and present only a pair of small depressions on the part close to the dorsal surface. The ventral surface is intact and is very smooth and concave in lateral view.

The articular faces are biconcave; i.e. both present very expressive concavities on their surfaces, the anterior being much deeper than the posterior. Their margins are very straight and have a semi-circular shape, the anterior one being slightly larger than the posterior.

-Comparisons:

UFRJ-DG 521-R has characteristics of a paravianmaniraptoran centrum positioned after the transition point in the tail (Russell, 1972, Gauthier, 1986; Tykoski, 2005), as it is being longer than high and possessing possesses a large reduction in both its neural spine and transverse processes, with those structures becoming midline ridges (Senter *et al.*, 2011; Motta *et al.*, 20187). This way it is Thus, it is possible to deduce that it is positioned at least after the vertebra 11 of the caudal series as seen in *Buitreraptor*, *Rahonavis*, *Dromaeosauridae* and *Troodontidae* (Ostrom, 1969; Forster *et al.*, 1998; Senter *et al.*, 2012; Xu *et al.*, 2017).

The presence of a reduced transverse process forming a midline ridges after the transition point is seen seen in some paravians, such as *Microraptor*, *Rahonavis*, *Buitreraptor*, and *Archaeopteryx*, but it is seen in specimens of the distal caudal vertebrae of *Microraptor*, *Rahonavis* and *Buitreraptor*. *Anchiornis* and *Archaeopteryx* (Hwang *et al.*, 2002; Hu *et al.*, 2009; Han *et al.*, 2014; Forster *et al.*, 1998; Novas *et al.*, 2017), a characteristic also seen in UFRJ-DG 521-R, which differentiates it from most other paravians as dromaeosaurids, *Archaeopteryx*, *Jeholornis* and *Anchiornis*.

Commented [52]: This sentence is confusing.
A: Done.

Formatted: Highlight

Formatted: Font: Italic

Commented [53]: I think the authors should specify which transition point are referring here (I suppose that proposed by Russel (1972), but this is my guess)
A: Done

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Commented [B554]: I don't agree with this taxonomy. *Microraptor* and *Buitreraptor* are dromaeosaurids, at least in most phylogenetic analyses. *Rahonavis* probably too. *Anchiornis* may

be a troodontid. The Agnolin and Novas phylogeny has many strange results and is not widely accepted.

So instead, just say that:

'The presence of a transverse process after the transition point is seen in some paravians, such as *Microraptor*, *Rahonavis*, *Buitreraptor*, and *Archaeopteryx*.'

Don't use the term *Averaptor*.

Formatted: Font: Italic, English (United States)

Formatted: English (United States)

Formatted: Font: Italic, English (United States)

Formatted: English (United States)

Formatted: Font: Italic, English (United States)

Formatted: English (United States)

The presence of the transverse processes after the transition point of the tail is not observed in more basal paravian groups of like dromaeosaurids and troodontids that have smooth lateral surfaces in their centra (Ostrom, 1969; Norell & Makovicky, 1999; Currie & Dong, 2001; Xu et al., 2012), but it is seen in specimens of *Microraptor*, *Rahonavis*, *Buitreraptor*, *Anchiornis* and *Archaeopteryx* (Hwang et al., 2002; Hu et al., 2009; Han et al., 2014; Novas et al., 2017), all belonging to the group Averaptora.

In addition, the 521-R specimen also has the dorsally positioned prezygapophyses more dorsally positioned in the same way as in *Buitreraptor*, *Rahonavis* and *Anchiornis* (fig. 059) (Motta et al., 20188). The vertebral centrum has a length-to-height ratio between its height and its length is close to 2.5, a ratio usually seen in dromaeosaurids with exception to *Buitreraptor* but not seen in other maniraptorans as troodontids and microraptorians whose ratio can reach up to 5.0 to 6.0. a ratio seen in more basal paravians such as dromaeosaurids, in contrast to troodontids and more derived averaptorans whose ratio can be as much as 6.0.

Figure 03: Caudal vertebrae attributed to Morphotype 02 (A-D) and Morphotype 03 (E-G). UFRJ-DG-528-R: A, ventral view; B, Lateral view; C, anterior articular facet. UFRJ-DG-532-R: D, anterior articular facet. UFRJ-DG-521-R: E, Lateral view; F, ventral view; G, anterior articular facet. Prz, prezygophysis; Ne, neural canal. Scale: 1cm.

Morphotype 4:

THEROPODA Marsh, 1881

Formatted: Indent: First line: 0.49"

Commented [BS55]: I don't agree with this taxonomy. *Microraptor* and *Buitreraptor* are dromaeosaurids, at least in most phylogenetic analyses. *Rahonavis* probably too. *Anchiornis* may be a troodontid. The Agnolin and Novas phylogeny has many strange results and is not widely accepted.

So instead, just say that:

'The presence of a transverse process after the transition point is seen in some paravians, such as *Microraptor*, *Rahonavis*, *Buitreraptor*, and *Archaeopteryx*.'

Don't use the term Averaptor.

Formatted: Font: Italic

Formatted: Font: Italic

Commented [BS56]: Again, don't use averaptorans. Instead, just say that long vertebrae like these are seen in some, but not all, paravians.

Formatted: Highlight

Formatted: Indent: First line: 0.49", Right: 0"

Commented [57]: I think the aspect of the figure is not proper: it shows different fonts, the letters are unaligned, and shows two scalebars. The same applies for Figure 4

Formatted: Justified, Indent: First line: 0.49"

Formatted: Indent: First line: 0.49", Right: 0"

470 TETANURAE Gauthier, 1986
471 ALLOSAUROIDEA Marsh, 1878
472 CARCHARODONTOSAURIA Benson, Brusatte and Carrano, 2010
473

474 Referred material: UFRJ-DG 523-R and 524-R.

475 *Description and comparisons:*
476 Morphotype 4 (Fig. 04) is formed by two vertebrae with low length/height ratio,
477 close to morphotype 1, and oval articular facet with strongly waisted centra with lateral
478 depressions in the dorsal half of the lateral surface, a double keel cut by a longitudinal
479 groove and offset articular facets.

480
481 Figure 04: Caudal vertebrae attributed to Morphotype 4. UFRJ DG 524 R: A, ventral
482 view; B, lateral view; C, anterior articular facet. UFRJ DG 523 R: D, ventral view; E,
483 lateral view; F, anterior articular facet. Nc, neural canal. Scale: 1 cm.

484
485 UFRJ-DG523-R

486 Specimen 523-R (Fig. 064, D-F) is a theropod vertebral centrum, with the
487 following characteristics: it is amphicoelous, and slightly longer than high (8.9 cm vs.
488 6.5 cm). Its lateral surface is very concave and smooth on both sides, with the shape of an
489 hourglass in dorsal view. The ventral surface is mostly smooth on the anterior part, with
490 breaches and marks that possibly indicate the articulation fusion of with the hemal arch
491 on the posterior part.

492 The dorsal surface is marked by a long and expressive deep longitudinal canal
493 from one articular face to the other, which widens on the extremities and tapers in the

Formatted: Font: Italic

Formatted: Indent: First line: 0.49"

Commented [58]: Again, this character varies with the position inside the caudal series.

Commented [59]: The shape of the articular face of caudal vertebrae is so variable, it show differences inside the same taxonomic group. For example inside Abelisauridae, Carnotaurus shows semicircular articular surface, and Majungasaurus show oval articular surface, as the authors clearly show in the figure 8. This makes this character not useful for separating morphotypes.

Formatted: Font: Not Bold, Pattern: Clear (White)

Formatted: Indent: First line: 0.49", Right: 0.02"

Formatted: Highlight

middle. This canal was possibly the space of the neural canal of the vertebra, given the marks of fusion with the neural arch that meet on its borders.

The articular faces are ~~ovoidal~~ ~~semi-circular~~ in shape and ~~with~~ have slightly forward-protruding margins ~~of the articular faces~~, the anterior facet being higher in comparison to the posterior facet. The anterior articular face has a concavity deeper than the posterior one, being also slightly larger in its proportions.

Commented [60]: The articular surfaces shown in the figure 4 has ovoidal articular faces.
A: Done.

UFRJ-DG524-R

Specimen 524-R (Fig. 064, A-C) is a centrum of a theropod caudal vertebra. It is amphicoelous and is slightly longer than high (~~7,5 cm vs. 8,5 cm~~), which indicates a more proximal position ~~between them in the~~ caudal ~~vertebrae series~~. The lateral surface is smooth and marked by two ~~deep~~ ~~expressive~~ concavities on both lateral faces, ~~giving to it an~~ ~~hourglass-like shape~~ ~~with a shape like an hourglass~~. Additionally, on the most dorsal region of the lateral surface ~~it is possible to notice~~ ~~there is~~ a small and shallow longitudinal depression on each side.

Formatted: English (United States)

Formatted: Highlight

Commented [61]: hourglass-shaped
A: Done.

The ventral surface is a double keel marked by a very superficial groove extending from the anterior part up to the posterior part. The dorsal surface is marked by the neural canal of the vertebrae. Above the anterior part of this canal the entire ~~cover~~ ~~upper~~ ~~portion~~ ~~art~~ of the ~~neural~~ tube is preserved, forming a small arch filled by sediment positioned slightly above ~~of~~ the anterior articular face.

The articular faces are semi-circular and somewhat oval, with the anterior one being slightly larger than the posterior, and their margins slightly protrude forward. The anterior articular face has a concavity slightly deeper than the posterior, ~~which is more~~ ~~superficial~~.

Comparisons:

Formatted: Font: Italic

Both vertebrae of morphotype 4 UFRJ-DG 523 and AND-524 present characteristics commonly found in carcharodontosaurids (Fig. 0740). For instance, depressions in the most dorsal part of the lateral surface are found in *Giganotosaurus*, *Mapusaurus* and *Tyrannotitan* and in the mid-caudal vertebrae Vb-870 found in the Wadi Milk Formation (Coria and Salgado, 1995; Coria and Currie, 2006; Novas *et al.*, 2005a; Canale *et al.*, 2015; Rauhut, 1999), which This condition is different from that in *Carcharodontosaurus*, that which has pleurocoels in their anterior caudal vertebrae (Stromer, 1931). Furthermore, the strongly waisted centrum morphology, a double keel cut by a longitudinal groove and offset articular facets (although it is a plesiomorphic feature found in *Allosaurus* Gilmore, 1920; Madsen, 1976) are also found in specimens such as the carcharodontosaurid material from Sudan (Rauhut, 1999) and in *Tyrannotitan*, *Mapusaurus* and *Acrocanthosaurus* (Canale *et al.*, 2015; Harris, 1998; Coria and Currie, 2006; Currie and Carpenter, 2000).

532

533

534

Morphotype 5:

536

537

THEROPODA Marsh, 1881

538

NEOVENATORIDAE Benson, Carrano and Brusatte, 2010

539

MEGARAPTORA Benson, Carrano and Brusatte, 2010

540

541

Referred material: UFRJ-DG 558-R e 634-R

542

Description and comparisons:

Commented [BS62]: I thought you were not using the morphotype terminology any more?

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Italic

Formatted: Indent: First line: 0.49"

Morphotype 5 (Fig. 05) includes two amphicoelous caudal vertebrae with pleurocoels and high pneumaticity.

Formatted: Indent: First line: 0.49"

Figure 05: Caudal vertebrae attributed to Morphotype 05. UFRJ-DG-634 R: A, posterior articular facet; B, lateral view; C, ventral view. UFRJ-DG-558 R: D, anterior articular facet; E, lateral view; F, ventral view. Pfr, Pneumatic foramen. Scale bar: 1cm.

Formatted: Indent: First line: 0.49", Right: 0.02"

UFRJ-DG-558-R

Formatted: Indent: First line: 0.49"

Specimen 558-R is a centrum of a theropod caudal vertebra, damaged by various cracks (Fig. 085, D-F). It is amphicoelous, and slightly longer than high (6cm vs. 7cm), indicating a somewhat proximal position within the caudal series. Its ventral surface is very smooth and convex in lateral view, but is very damaged in the region where the base of the posterior articular face would be.

Formatted: English (United States)

The dorsal surface is marked by a great depression extending longitudinally from one articular face to the other, wider in the extremities, denoting the neural canal. The lateral surfaces are marked by a longitudinal elliptic depression on their medial parts, where there is a pleurocoel on each side. The left lateral pleurocoel is deeper and better defined than the right lateral one. ~~The presence of pleurocoels in the caudal vertebrae is characteristic of megaraptoran neovenatorids the Megaraptora group of the Neovenatoridae family (Benson et al., 2011).~~

Commented [B563]: This shouldn't be here, but in the comparisons section, below.

Its articular faces are semi-circular and have very straight margins. The anterior articular face possesses a more distinctive depression of a slightly greater size than the posterior face and is also in a better state of preservation. The posterior articular face

possesses a very slight concavity, making it almost straight, and is in a much more damaged state, presenting cracks and breaches on the ventral base of the face.

UFRJ-DG 634-R

—This material is in a worse state of preservation than UFRJ-DG 558-R (Fig. 085, A-C). The ventral centrum portion and anterior articular face are fragmented. On its lateral surface, there is what appears to be the border of the pleurocoel in the same position seen in specimen 558-R, leading to the attribution of this vertebra to this morphotype.

Differently from the other vertebrae of this morphotype group, part of the neural arch and the transverse process ~~are~~ preserved on the right side of the specimen. The transverse process is positioned upwards at an angle of approximately 45° degrees.

Comparisons

The presence of pleurocoels in the caudal vertebrae is characteristic of megaraptoran neovenatids (Benson *et al.*, 2010). Pneumaticity in the caudal vertebrae is rare in Theropoda, present only in some groups: Megaraptora, Oviraptorosauria, Therizinosauria, and Carcharodontosauridae (Benson *et al.*, 2012+). As far as is known, no fossils of therizinosaurs have been found in South America and South American fossils attributed to oviraptorosaurs have been reassigned to other taxa, including to Maniraptora (e.g. see Agnolín & Martinelli, 2007, Aranciaga-Rolando *et al.*, 2018). In addition, the caudal vertebrae of Oviraptorosauria have, on the ventral surface, a medial groove delimited by two longitudinal elevations (e.g., Sues, 1997; Xu *et al.*, 2007). Specimen UFRJ-DG 558-R does not have this feature (Fig. 09).

South American carcharodontosaurids (e.g., *Giganotosaurus*, *Mapusaurus*, *Tyrannotitan*) show a slightly concave lateral sides in the caudal vertebrae, whilebut do

Formatted: Indent: First line: 0.49", Right: 0"

Formatted: Indent: First line: 0"

Formatted: English (United States)

Formatted: Indent: First line: 0"

Commented [BS64]: This shouldn't be here, but in the comparisons section, below.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Not Highlight

not ~~being~~^{bear} actual pneumatic foramina. Stromer (1931) described an anterior caudal vertebra from northern Africa, which he identified as *Carcharodontosaurus*, which had pneumatic characteristics, including a pleurocoel. However, that vertebra has a different general morphology and proportions when compared with ~~morphotype~~^{the megaraptorid} ~~vertebrae~~^{from the Potiguar Basin (length-height ratio is 1 in *Carcharodontosaurus* and approximately 1.48 in UFRJ DG 558-R) and other members of Megaraptora.}

Commented [BS65]: Again, I thought you were not using the morphotype descriptions any more.

Among the Megaraptora group, only *Aerosteon*, *Aoniraptor*, *Orkoraptor* and *Megaraptor* have preserved caudal vertebrae (Figure 1007) (Serenó *et al.*, 2008; Benson *et al.*, 2010; Motta *et al.*, 2016). The height/length ratio of UFRJ DG 558-R is 1.4, consistent with a median tail position, compared to the ratios of 1.2 and 1.3, respectively, of the medial caudal vertebrae of *Aerosteon* and *Orkoraptor* (Novas *et al.*, ~~al.~~²⁰⁰⁸). The Potiguar Basin specimens resemble those of *Aoniraptor* (Fig. 07, F) due to the absence of a keel in the ventral region, but ~~is~~^{are} distinguished by the presence of a pair of pneumatic ~~septal~~^{troughs} in the lateral region, separated by a septum. Only the first caudal vertebra of *Aoniraptor* presents such fossae, a characteristic present in the other megaraptorans (e.g., Novas *et al.*, 2008; Sereno *et al.*, 2008).

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Comparing the morphology of pneumatic foramina, UFRJ DG 558-R (Figure 1007, A) is very similar to *Aerosteon* (Figure 1007, C), *Megaraptor* (Figure 0710, H) and *Orkoraptor* (Figure 1007, G) ~~by~~ⁱⁿ the presence of a large elliptic foramen and a second smaller circular shaped foramen. In addition, ~~morphotype 6~~^{UFRJ-DG 558-R and 634-R} has its cavities located on the lateral surface of the vertebral centrum near the base of the neural arch, which does not occur in the other species ~~analyzed~~^{observed}.

Formatted: Font: Italic

Formatted: Font: Italic

Commented [BS66]: Same comment as above.

UFRJ-DG 558-R and 634-R The ~~morphotype 06~~^{vertebrae} also presents extensive pneumatization in the vertebral centrum, composed of a camerate internal microstructure

Commented [BS67]: Same comment

(Britt, 1993), with several small chambers, similar to other megaraptorans (e.g., *Aerosteon, Megaraptor*; Martinelli *et al.*, 2013).

Formatted: Font: Italic

Discussion

Formatted: Indent: First line: 0"

The Açú Formation material and ~~your~~its importance

The fossil potential of Açú Formation was poorly known, with only a few fossils recovered (Duarte and Santos, 1962; Silva-Santos, 1963; Mussa *et al.*, 1984), until the discovery of vertebrae and teeth identified as belonging to Theropoda indet. and Titanosauria (Santos *et al.*, 2005).

No further work was conducted until 2018, when the materials described here were studied in more detail. Thus far, the dinosaur fauna of the Potiguar Basin includes two groups of Sauropoda (Diplodocoidea: Rebbachisauridae, Pereira *et al.*, in press; Titanosauriformes, Barbosa *et al.*, 2018; Titanosauria, Pereira *et al.*, 2018) and four groups of Theropoda (Spinosauroidea, Carcharodontosauridae, Megaraptora and Maniraptora, present work). ~~The fossil potential of Açú Formation was poorly known, with only a few fossils recovered (Duarte & Santos, 1962; Silva-Santos, 1963; Mussa *et al.*, 1984), until the discovery of vertebrae and teeth identified as belonging to Theropoda indet. and Titanosauria (Santos *et al.*, 2005).~~

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

The occurrence of these groups (except Megaraptora) in the Potiguar Basin is yet another similarity between the faunas of northeastern Brazil and multiple North Africa Cretaceous units (e.g. Medeiros and Schultz, 2001a, 2002; Sereno and Brusatte, 2008; Contessi, 2009; Candeiro *et al.*, 2011; Candeiro, 2015). Except for the Elrhaz (Niger); Douiret and Ain El Guettar (both in Tunisia) and Chicla (Libya) formations, which were

dated to the Early Cretaceous, all other Cretaceous formations from Northern Africa are Albian-Cenomanian in age, roughly equivalent to the Açu Formation (Werner, 1994; Rossetti, 1997; Rossetti and Truckenbrodt, 1997; Smith et al., 2001; Anderson et al., 2007; Sereno and Brusatte, 2008; Cavin et al., 2010). Among the formations, the Alcântara Formation (Brazil), Bahariya Formation (Egypt), Echkar Formation (Niger) and the Waldi Milk Formation (Sudan) have similarities with the Açu Formation's dinosaur fauna.

According to paleobiogeographic models, South America and Africa started separating from each other in the Valanginian (Early Cretaceous), leading to the formation of the South Atlantic Ocean (Viramonte et al., 1999; Jokat et al. 2003; Macdonald et al., 2003). Although the ocean turned into one of the most important continental barriers of the southern hemisphere, faunal interchange among the terrestrial landmasses of western of Gondwana definitely occurred up to the Albian, and possibly until the Cenomanian (e.g. Petri, 1987; Reymont and Dingle, 1987; Pletsch et al., 2001, Tello Saenz et al., 2003, Guedes et al., 2005, Bodin et al., 2010).

Based on the proposed age and geographic position, the fossil vertebrates of the Açu Formation may have lived during some of the last intervals of continental connection between South America and Western Africa, before the complete formation of the South Atlantic Ocean (Arai, 2009; Castro et al., 2012). This makes them exceedingly important for understanding biogeography and faunal evolution.

More extensive comparisons are still limited by the lack of completeness of the Açu material and the absence of formally described taxa. The continuation of studies on previously collected material (like that described in this paper) and prospecting for new fossils is important in this basin which, while still the subject of only recent research, already exhibits among the greatest diversity of dinosaur groups in Brazil.

No further work was conducted until 2018, where when the materials described here began to be described were studied in more detail. For now, thus far, the dinosaur fauna of the Potiguar Basin has so far includes two groups of Sauropoda (Diplodocoidea: Rebbachisauridae, Pereira *et al.* in press; Titanosauriformes, Barbosa *et al.*, 2018; Titanosauria, Pereira *et al.*, 2018) and four groups of Theropoda (Spinosaurioidea, Carcharodontosauridae, Megaraptora and Paraves, present work).

The occurrence of these groups (except Megaraptora) in the Potiguar Basin is one yet another of the numerous similarities between the faunas of northeastern Brazil and multiple North Africa Cretaceous units (e.g., Medeiros & Schultz, 2001a, 2002; Sereno & Brusatte, 2008; Contessi, 2009; Candeiro *et al.*, 2011; Candeiro, 2015). Except for the Elrhaz (Niger) formations; Douiret and Ain El Guettar (both in Tunisia) and Chiela (Libya) formations, which were dated to belong to the Early Cretaceous, all other Cretaceous formations from Northern Africa are Albian-Cenomanian in age, roughly equivalent to the Açu Formation (Werner, 1994; Rossetti, 1997; Rossetti & Truckenbrodt, 1997; Smith *et al.*, 2001; Anderson *et al.*, 2007; Sereno & Brusatte, 2008; Cavin *et al.*, 2010). Among the formations, the Alcântara Formation (Brazil), Bahariya Formation (Egypt), Echkar Formation (Niger) and the Waldi Milk Formation (Sudan) may have special attention due to have similarities with the Açu Formation's dinosaur fauna.

According to paleobiogeographic models, South America and Africa started separating from each other in the Valanginian (Early Cretaceous), leading to the formation of the South Atlantic Ocean (Viramonte *et al.*, 1999; Jokat *et al.*, 2003; Macdonald *et al.*, 2003). Although the ocean turned into one of the most important continental barriers of the southern hemisphere, faunal interchange among the terrestrial landmasses of western of Gondwana definitely occurred up to the Albian, and possibly

Formatted: Font: Italic

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Commented [BS68]: I am not convinced this is a paravian bone. It might be, but what are the clear diagnostic characters that say it must be a dromaeosaurid/troodontid/bird and not a member of another maniraptoran group?

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Commented [BS69]: The Albian is Early Cretaceous

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

until the Cenomanian (e.g. Petri, 1987; Reymont & Dingle, 1987; Pletsch *et al.*, 2001; Tello Saenz *et al.*, 2003; Guedes *et al.*, 2005; Bodin *et al.*, 2010).

Based on the proposed age and geographic position, the fossil vertebrates of the Açu Formation may have lived during some of the last moments/intervals of continental connection between South America and Western Africa, before the complete formation of the South Atlantic Ocean (Arai, 2009; Castro *et al.*, 2012). This makes them exceedingly important for understanding biogeography and faunal evolution.

More extensive comparisons are still limited by the lack of completeness of the Açu material and the absence of formally described taxa. The continuation of the studies on the already previously collected material (like that described in this paper) and the prospecting for new fossils is important in this basin which, while the research still the subject of only recent research, it is already a formation that shows one of exhibits among the greatest diversity of dinosaur groups in Brazil.

The first described megaraptoran was *Megaraptor namunhuaiquii* from the Turonian of Patagonia/Argentina (Novas, 1998). Recently, new findings have increased our knowledge about the anatomy and taxonomic diversity of these animals (Calvo *et al.*, 2004; Novas *et al.*, 2008; Hocknull *et al.*, 2009; Novas, 2009).

In 2010, Benson *et al.* created the name *Megaraptora* for a newly recognized clade of theropods including taxa found in Argentina (*Acrosteon*, *Megaraptor*, *Aoniraptor* and *Orkoraptor*), Australia (*Australovenator*) and Japan (*Fukuiraptor*). One of the most striking features of the group is the presence of pneumatic anterior-caudal vertebrae (Calvo *et al.*, 2004).

Pneumaticity in the caudal vertebrae is rare in Theropoda, present only in some groups: *Megaraptora*, *Oviraptorosauria*, *Therizinosauria* and *Carcharodontosauridae* (Benson *et*

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Not Highlight

al., 2011). As far as is known, no fossils of therizinosaurs have been found in South America and South American fossils attributed to oviraptorosaurs have been reassigned to other taxa (see Agnolín & Martinelli, 2007). In addition, the caudal vertebrae of Oviraptorosauria have, on the ventral surface, a medial groove delimited by two longitudinal elevations (e.g., Sues, 1997; Xu et al., 2007). Specimen UFRJ-DG-558-R does not have this feature.

South American carcharodontosaurids (e.g., *Giganotosaurus*, *Mapusaurus*, *Tyrannotitan*) show ~~very deeply concave lateral sides in the caudal vertebrae, but not apneumatic foramen~~. Stromer (1931) described an anterior caudal vertebra from northern Africa, which he identified as *Carcharodontosaurus*, which had pneumatic characteristics, including a pleurocoel. However, that vertebra has a different general morphology and proportions when compared with morphotype 6 from the Portuguese Basin (length-height ratio is 1 in *Carcharodontosaurus* and approximately 1.48 in UFRJ-DG-558-R) and other members of Megaraptora.

~~In Brazil, there are only two previous records (Fig. 06) attributed to Megaraptora, both in the Bauru Group. Méndez et al. (2012) described an isolated caudal vertebral centrum (MPMA 08-003-94), found in the municipality of Ibirá, São Paulo (Maastrichtian, Late Cretaceous). The authors compared their specimen with the megaraptorids *Aerosteon* and *Megaraptor*, and found important differences, such as the absence of a median longitudinal keel on the ventral surface and its more elongated proportions. Martinelli et al. (2013) described another isolated caudal vertebra found in Uberaba (Campanian, Late Cretaceous) as belonging to Megaraptora.~~

~~In a recent work, Motta et al. (2016) considered that both specimens are in fact sacral vertebrae, due to their more elongated proportions, rough articular face and anteroposteriorly expanded transverse processes. Due to this fact, a comparison with the~~

Commented [70]: The lateral surfaces are slightly concave in these taxa, not very.
A: Done

Commented [71]: Please see Aranciaga-Rolando et al (2018) A supposed Gondwanan oviraptorosaur from the Albian of Brazil represents the oldest South American megaraptoran. Cretaceous Research 84: 107-119.

material analyzed here becomes difficult, and the Potiguar vertebrae are thus the first caudal vertebrae of megaraptorans from Brazil.

Figure 06: Brazilian Megaraptoran Vertebrae findings. A and B, UFRJ DG 558 R; C and D, MPMA 08-003-94 (Méndez et al., 2012); E and F, CPPLIP 1324 (Martinelli et al., 2013). A, C e E, lateral view; B, D e F, ventral view. Pfr, Pneumatic foramen. Scale bar = 1cm.

Formatted: English (United States)

Among the Megaraptora group, only *Aerosteon*, *Aoniraptor*, *Orkoraptor* and *Megaraptor* have preserved caudal vertebrae (Figure 07) (Sereno et al., 2008; Benson et al., 2010; Motta et al., 2016).

The height/length ratio of UFRJ DG 558 R is 1.4, consistent with a median tail position, compared to the ratios of 1.2 and 1.3, respectively, of the medial caudal vertebrae of *Aerosteon* and *Orkoraptor* (Novas et al. al., 2008). The Potiguar Basin specimens resemble those of *Aoniraptor* (Fig. 07, F) due to the absence of a keel in the ventral region, but is distinguished by the presence of a pair of pneumatic septal troughs in the lateral region, separated by a septum. Only the first caudal vertebra of *Aoniraptor* presents such fossae, a characteristic present in the other megaraptorans (e.g., Novas et al., 2008; Sereno et al., 2008).

Comparing the morphology of pneumatic foramina, UFRJ DG 558 R (Figure 07, A) is very similar to *Aerosteon* (Figure 07, C), *Megaraptor* (Figure 07, H) and *Orkoraptor* (Figure 07, G) by the presence of a large elliptic foramen and a second smaller circular shaped foramen. In addition, morphotype 6 has its cavities located on the lateral surface

of the vertebral centrum near the base of the neural arch, which does not occur in the other species analyzed.

Figure 07: Megaraptoran caudals vertebrae. A and B, UFRJ DG 558 R; C and D, *Aerosteon*; E and F, *Aoniraptor*; G, *Orkoraptor*; H, *Megaraptor*. A, C, E, G e H, lateral view; B, D e F, ventral view. Pfr, pneumatic foramen. Scale bar = 5cm.

Formatted: English (United States)

The morphotype 06 vertebrae also presents extensive pneumatization in the vertebral centrum, composed of a camerate internal microstructure (Britt, 1993), with several small chambers, similar to other megaraptorans (e.g., *Aerosteon*, *Megaraptor*; Martinelli et al., 2013). Based on the general morphology, the elements described herein possibly belong to a form closer to *Aerosteon* and *Megaraptor* than to *Aoniraptor*.

Commented [72]: This is a too risky statement given the scarcity of the recovered materials.

The Potiguar material is also one of the oldest records of the group in South America, together with *Aoniraptor* from the Early Cenomanian mid-Turonian of Argentina (Motta et al., 2016).

Commented [73]: Please see Aranciaga-Rolando et al (2018) A supposed Gondwanan oviraptorosaur from the Albian of Brazil represents the oldest South American megaraptoran. Cretaceous Research 84: 107-119.

The abelisauroid fossil record from Brazil is known from fragmentary specimens. Recently, Silva (2013) and Santucci et al. (2018) described incomplete abelisaurid cranial and postcranial Abelisauria remains from Barremian-Aptian age of Quiricó Formation (São Francisco Basin) of northern Minas Gerais state.

Brazilian noosaurids were known only by teeth from Albian-Cenomanian of Alcântara Formation, where *Masiakasaurus* like teeth were recorded. However this changed when *Vespersaurus paranaensis*, a desert dwelling monodactyl noosaurid, was described in the

Formatted: Font: Italic

Formatted: Font: Italic

Late Cretaceous Rio Paraná Formation (Langer et al., 2019). From Albian-Cenomanian of Alcântara formation bed have produced one tooth reported as *Masiakasaurus* like (Noosauridae) by Lindoso et al. (2012).

Commented [74]: You can add *Vespersaurus*. A: After revising the discussion, the comparison to *Masiakasaurus* and other Noosaurids seemed to have no place anymore. In the end we restricted to the abelisaurid record only, not to the abelisauroid.

~~The first abelisaurid species discovered in Brazil was *Peyronemosaurus nevesi* was the first and unique specie of abelisaurid from Brazil (Kellner & Campos, 2002). It was described based on postcranial remains from the Upper Cretaceous of the Ribeirão Boiadeiro Group and represents the most complete collection known to Brazil. Additionally, Bittencourt and Kellner (2002) described nine Abelisauria teeth from the same locality of *Peyronemosaurus*. The second abelisaurid discovered in Brazil was *Thanos simonattoi* (Delecourt & Iori, 2018), whose description was based on an almost complete axis with an axial intercentrum.~~

~~The first record of Abelisauridae known to Brazil was reported by Bertini (1996) and was discovered in the Adamantina Formation, western São Paulo State. Later, other authors reported isolated teeth from the Adamantina Formation from western São Paulo state (e.g., Candeiro et al., 2004; Azevedo et al., 2007) and from Minas Gerais state (e.g., Candeiro et al., 2006; Oliveira et al., 2012). Also, there are some known postcranial records from the São José do Rio Preto and Marília formations (Méndez et al., 2014) were recently described by Méndez et al. (2014) as well as other abelisaurid materials from the Adamantina Formation, São Paulo State (a partial femur, Brum et al., 2016).~~

~~However, the most abundant materials of abelisaurid are from the Marília Formation that outcrops in the region of Peirópolis, municipality of Uberaba. Innumerable teeth (Candeiro et al. 2012) and postcranial materials (Novas et al. 2008, Machado et al. 2013) from this locality were already described.~~

~~Even though most of the abelisaurid axial characteristics are mainly in their uniquely shaped transverse processes in the caudal vertebrae (e.g. Méndez, 2014), there are some characteristics that can be seen in their caudal centra.~~

~~The mid-caudal vertebrae of Abelisauria have, as basic characteristics, an amphicoelous condition with subcircular articular facets, a centrum twice as long as tall, well marked~~

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Italic

Formatted: Font: Italic

Commented [75]: As was stated before, this character varies a lot inside Abelisauridae.

facets, and a ventral concavity in its lateral view as seen in *Majungasaurus*, *Aucasaurus*, and *Ilokelesia* (Coria & Salgado, 1998), however much these features are also seen in Abelisauroides and even in basal ceratosaurs (Méndez, 2014). All these characteristics can be observed in the specimens of morphotype 2, with specimen 547 R standing out from the others for the presence of two small depressions on its lateral surface, a characteristic not very common within the group.

The anterior caudal vertebrae of Abelisauria present a set of striking features that can be observed in specimens of morphotype 3 (Fig. 08). First is its oval, taller than wide, facet that both the 528 R specimen and the 532 R have. In addition, there are features found in the centrum of specimen 528 R, such as the slight centrum? central? or constriction (Méndez, 2014) and a concave anterior margin while the posterior convex margin is seen in the first four caudal vertebrae of *Aucasaurus*, *Carnotaurus*, *Majungasaurus*, *Ekrixinatosaurus*, *Rajasaurus* and *Rahiolisaurus* (Bonaparte, 1985; Sampson et al. 1998; Coria et al. 2002). Unlike *Aucasaurus*, *Carnotaurus* and *Ekrixinatosaurus*, specimen 528 R also does not exhibit any forms of depression or pneumaticity on its lateral surface. These are the oldest record of Abelisauria at Brazil.

Figure 08: Comparison of morphotype 02 and other abelisaurids. A and B, *Majungasaurus*; C and D, *Carnotaurus*; E and F, *Ekrixinatosaurus*; G and H, *Aucasaurus*; I and J, morphotype 02. Scale bar: 5 cm.

The record of maniraptorans is rare in Brazil (see Delcourt & Grillo, 2014). It is based mostly on isolated teeth from several localities and postcranial elements, namely, a manual ungual and scapula from the Serra da Galga Member of Peirópolis, Uberaba,

Commented [76]: Which facets?

Commented [77]: None of the characters mentioned are exclusive of Ceratosauria, but they are present in other theropods. Moreover, the specimens described and figured for this morphotype are anterior caudals, not middle.

Commented [78]: Which is the specimen 547-R? It was not described or figured!

Commented [79]: Morphotype 3 was assigned to Maniraptora in the text, not Abelisauria.

Commented [80]: Again, this character is so variable. Please compare anterior articular surface of *Carnotaurus* and *Majungasaurus* in the figure 8.
A: Done.

Commented [81]: It is not clear to which margins the authors refers.
A: Done.

Commented [82]: I feel that the evidence given by the authors in the text is not enough to assign this material to Abelisauria, so this statement is too risky.

Minas Gerais state (Marília Formation, Maastrichtian), and an unenlagioid dorsal vertebra and fragmentary remains of an undetermined maniraptoran from the Adamantina Formation (Late Cretaceous) of São Paulo state (Novas et al., 2005; Machado et al., 2008; Candeiro et al., 2012; Delcourt & Grillo, 2014). The material here described is the first post-cranial remain of a maniraptoran outside Bauru Basin and from the mid-Cretaceous, shedding new light on the biogeography of this group in South America and western Gondwana.

Specimen 521-R has characteristics of a vertebral centrum positioned after the transition point, being longer than high possessing a large reduction in both its neural spine and transverse processes (Senter et al., 2011). This way it is possible to deduce that it is at least after the vertebra 11 of the caudal series as seen in *Buitreraptor*, *Rahonavis*, *Dromaeosauridae* and *Troodontidae* (Ostrom, 1969; Forster et al., 1998; Senter et al., 2012; Xu et al., 2017).

The maintenance of the transverse processes is not observed in more basal groups of Paraves like dromaeosaurids and troodontids that have smooth lateral surfaces in their centra (Ostrom, 1969; Norell & Malkovicky, 1999; Currie & Dong, 2001; Xu et al., 2012), but is seen in specimens of *Microraptora*, *Rahonavis*, *Buitreraptor*, *Anchiornis* and *Archaeopteryx* (Hwang et al., 2002; Hu et al., 2009; Han et al., 2014; Novas et al., 2017), all belonging to the group Aves. In addition, the 521-R specimen also presents the prezygapophyses more dorsally positioned in the same way as in *Buitreraptor*, *Rahonavis* and *Anchiornis* (fig. 09) (Motta et al., 2018). The vertebral centrum has a ratio between its height and its length somewhat greater than six which is seen in almost all the groups of Paraves except for *Dromaeosauridae* that displays a smaller ratio being the only 3 times longer than high.

Commented [83]: I think the authors should specify which transition point are referring here (I suppose that proposed by Russel (1972), but this is my guess)
A: Done

Figure 09: Comparison of Morphotype 3 and other paravians. A, Potiguar's material; B, *Rahonavis*; C, *Buitreraptor*; D, *Anchiornis*. Pr, prezygapophysis; lg, Longitudinal groove. Modified from Motta et al., (2018).

Commented [84]: In the references the date of the publication is 2017

Carcharodontosaurid were amongst the largest and some of the most widespread theropods during the Kimmeridgian-Turonian (Candeiro, 2015; Delecourt & Grillo, 2017). In Brazil, carcharodontosaurids were recorded based on isolated teeth and putative remains from São Luís-Grajaú Basin (Cenomanian) and Bauru Group (Late Cretaceous) (Vilas Bôas et al., 1999; Medeiros, 2001; Azevedo et al., 2013). The Potiguar Basin materials lie within the age-range of occurrence of carcharodontosaurids and, might be related to the São Luís-Grajaú and western Africa fauna where these are common findings.

Both vertebrae of morphotype 4 present characteristics commonly found in carcharodontosaurids (Fig. 10). For instance, depressions in the most dorsal part of their lateral surface are found in *Giganotosaurus*, *Mapusaurus* and *Tyrannotitan* and in the mid-caudal vertebrae Vb 870 found in the Wadi Milk Formation (Coria & Salgado, 1995; Coria & Currie, 2006; Novas et al., 2005; Canale et al., 2015; Rauhut, 1999), which is different from *Carcharodontosaurus*, which has pleurocoels in their anterior-caudal vertebrae (Stromer, 1931). Furthermore, the strongly waisted centrum morphology, a double-keel cut by a longitudinal groove and offset articular facets (although it is a plesiomorphic feature found in *Allosaurus* Gilmore, 1920; Madsen, 1976) are also found in specimens such as the carcharodontosaurid material from Sudan (Vb 870) and in *Tyrannotitan*, *Mapusaurus* and *Acrocanthosaurus* (Canale et al., 2015; Harris, 1998; Coria & Currie, 2006; Currie & Carpenter, 2000).

Commented [85]: Cite?
A: Done.

Formatted: Indent: First line: 0"

Figure 10: Comparison of Morphotype 4 and *Carcharodontosauria* indet. A and B, UFRJ DG 523 R; C and D, UFRJ DG 524 R; E and F, Kem Kem beds material (from Rauhut, 1999). G? H? I?. Scale bar = 5 cm.

Commented [86]: The item G of the figure has a very poor resolution. Also G, H, I were taken from the literature, and the figure caption lack the proper cite.

Formatted: Indent: First line: 0"

The record of maniraptorans is rare in Brazil (see Delcourt & Grillo, 2014). It is based mostly on isolated teeth from several localities and postcranial elements, namely, a manual ungual and scapula from the Serra da Galga Member of Peirópolis, Uberaba, Minas Gerais state (Marília Formation, Maastrichtian), and an unenlagioid dorsal vertebra and fragmentary remains of an undetermined maniraptoran from the Adamantina Formation (Late Cretaceous) of São Paulo state (Novas et al., 2005; Machado et al., 2008; Candeiro et al., 2012; Delcourt & Grillo, 2014). The material here described is the first postcranial remain of a maniraptoran outside Bauru Basin and from the mid-Cretaceous, shedding new light on the biogeography of this group in South America and western Gondwana.

Specimen 521 R has characteristics of a vertebral centrum positioned after the transition point, being longer than high possessing a large reduction in both its neural spine and transverse processes (Senter et al., 2011). This way it is possible to deduce that it is at least after the vertebra 11 of the caudal series as seen in *Buitreraptor*, *Rahonavis*, *Dromaeosauridae* and *Troodontidae* (Ostrom, 1969; Forster et al., 1998; Senter et al., 2012; Xu et al., 2017).

The maintenance of the transverse processes is not observed in more basal groups of Paraves like dromaeosaurids and troodontids that have smooth lateral surfaces in their centra (Ostrom, 1969; Norell & Malkovicky, 1999; Currie & Dong, 2001; Xu et al., 2012), but is seen in specimens of *Microraptor*, *Rahonavis*, *Buitreraptor*, *Anchiornis* and *Archaeopteryx* (Hwang et al., 2002; Hu et al., 2009; Han et al., 2014; Novas et al., 2017),

all belonging to the group Averoaptora. In addition, the 521-R specimen also presents the prezygapophyses more dorsally positioned in the same way as in *Buitreraptor*, *Rahonavis* and *Anchiornis* (fig. 10) (Motta et al., 2018). The vertebral centrum has a ratio between its height and its length somewhat greater than six which is seen in almost all the groups of Paraves except for Dromaeosauridae that displays a smaller ratio being the only 3 times longer than high.

The spinosaurid record of Brazil is known by two species: *Oxalaia quilombensis* (CITE) and *Irritator challengeri* (CITE), both spinosaurines from the "mid"-Cretaceous strata from the Cenomanian Aleantra Formation and from the Aptian-Albian Santana Formation respectively. Other than those two species, there are many unidentified isolated spinosaurid post cranial elements and teeth that range from Berriasian-Valanginian Feliz Deserto Formation (CITE) to the "mid"-Cretaceous Santana and Aleantra Formations (CITE).

There is no discussion about the recovered tooth, and the traits that allow assigning it to Spinosauridae.

Conclusion **REVISARRRRR**

In the present work we assigned the material from Açú Formation, Potiguar Basin, to four groups: Spinosauroida, Carcharodontosauria, Maniraptora and Megaraptora (Fig. 11), the two last groups being relatively rare in Brazil. All this groups have already been found in isochronous formations in both Northeastern Brazil and Northern Africa, leading further support for faunal similarities in the "mid"-Cretaceous western Gondwana. These fossils provide the first theropod record from Potiguar Basin and an important opportunity to increase the knowledge on the diversity of this still poorly known basin. We describe

Commented [87]: This text is repeated to that of the lines 539-569!
A: Done.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: English (United States)

Formatted: Indent: First line: 0"

Formatted: Highlight

~~several newly discovered dinosaur fossils, which constitute A a new dinosaur fauna was discovered and described from the Açu Formation (Potiguar Bbasin), Northeast region of Brazil. The vertebrae were are classified into five morphotypes based on morphological form and/or diagnostic characters and comprised at this moment by can be assigned to five four groups: Abelisauria, Carcharodontosauria, Spinosauridae, Megaraptora, and Maniraptora. While the teeth was recovered as a Spinosaurid Furthermore, a single tooth is attributed to Spinosauridae Besides These groups were have already been found on in isochronous basins of the Northeast region of Brazil and Africa, lending further support for faunal similarities between these regions.~~

Commented [88]: This group was not recognized by any vertebra, but a tooth.
A: Done.

~~the presence of these theropod groups at Açu Formation The new fossils reveals an unexpected dinosaur richness fauna at from the Potiguar Basin (fig. 11) and opens up an provide an important opportunity to increase the knowledge about on the diversity and palaeobiogeography of this important vertebrate group these animals during a time of Gondwanan fragmentation.~~

~~Figure 11: Reconstruction of the theropods groups described in this present study to the Açu Formation, Potiguar Basin. In the water, a Spinosauridae. On the ground, on the left an Abelisauridae; in the center, a group of megaraptorans Megaraptora and a slaughtered atitanosaur Titanosauriasauropod while to the right, a Carcharodontosauridae awakens from its sleep; in the top center, a Paraves just watches. Drawing of Luciano da Silva Vidal.~~

Formatted: English (United States)

Acknowledgements

Formatted: No underline

The authors thank Prof. Dr. Valdeci dos Santos Júnior Santos for discovering the site where the material were found, and for the support given to the fieldwork. We thank the students Luciano Vidal for assistance with the figures. PVLGCP was funded by a grant from the Jurassic Foundation and pos doctoral grant by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, number 88882.463232/2019-01). LPB and CRAC were financially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)/Bolsista de Produtividade em Pesquisa. IMMGB and LPB were also supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico [grant 459086/2014-6]. LPB also acknowledge Fundação de Amparo a Pesquisa do Estado do Rio de Janeiro [grants #E-26/202.829/2018]. This research was funded partially by Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). Our collaborative project was funded also by a grant from the Fundação de Amparo a Pesquisa e Goiás and the Newton Fund, which supported SLB's visit to Brazil to work with PVLGCP and CRAC in June–July 2016.

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

References

- Agnolín, F. L., Martinelli, A.G. 2007. Did oviraptorosaurs (Dinosauria: Theropoda) inhabit Argentina. *Cretaceous Research* 28, 785–790.
- Anderson, P.E., Benton, M.J., Trueman, C.N., Paterson, B.A., Cuny, G., 2007. Palaeoenvironment of vertebrates on the southern shore of Tethys: the nonmarine Early Cretaceous of Tunisia. *Palaeogeography, Palaeoclimatology, Palaeoecology* 243 (1–2), 118–131.

Formatted: Font: 14 pt, Bold, No underline, English (United States)

- Arai, M. 2011. Paleogeografia do Atlântico Sul no Aptiano: um novo modelo a partir de dados micropaleontológicos recentes. *Boletim de Geociências da Petrobras*, 17(2): 331-351.
- Aranciaga-Rolando, A. M., Egli, F. B., Sales, M. A., Martinelli, A. G., Canale, J. I., Ezcurra, M. D. 2018. A supposed Gondwanan oviraptorosaur from the Albian of Brazil represents the oldest South American megaraptoran. *Cretaceous Research* 84, 107-119.
- Araripe, P. T., Feijó, F. J. 1994. Bacia Potiguar. *Boletim de Geociências da Petrobras* 8: 27-141.
- Barbosa, F.H.S., Ribeiro, I.C., Pereira, P.V.L.G.C., Bergqvist, L.P. 2018. Vertebral lesions in a titanosaurian dinosaur from the Lower-Upper Cretaceous of Brazil. *Geobios* 51: 385–389.
- Benson, R. B., Carrano, M. T., Brusatte, S. L. 2010. A new clade of archaic large-bodied predatory dinosaurs (Theropoda: Allosauroidea) that survived to the latest Mesozoic. *Naturwissenschaften*, 97(1), 71-78.
- Benson, R. B., Butler, R. J., Carrano, M. T., and O'Connor, P. M. 2012. Air-filled postcranial bones in theropod dinosaurs: physiological implications and the 'reptile'–bird transition. *Biological Reviews*, 87(1), 168-193.
- Bodin, S., Petitpierre, L., Wood, J., Elkanouni, I., Redfern, J. 2010. Timing of Early to mid-Cretaceous tectonic phases along North Africa: new insights from the Jeffara escarpment (Libya–Tunisia). *Journal of African Earth Sciences* 58 (3), 489–506.
- Britt, B. B. 1993. Pneumatic postcranial bones in dinosaurs and other archosaurs. *Geology and Geophysics*, University of Calgary.

Formatted: Portuguese (Brazil)

- Brusatte, S. L., Candeiro C. R. A., Simbras, F. M. 2017. The last dinosaurs of Brazil: The Bauru Group and its implications for the end-Cretaceous mass extinction. *Annals of the Brazilian Academy of Sciences* 89: 1465-1485.
- Canale, J. I., Novas, F. E., Pol, D. 2015. Osteology and phylogenetic relationships of *Tyrannotitan chubutensis* Novas, de Valais, Vickers-Rich and Rich, 2005 (Theropoda: Carcharodontosauridae) from the Lower Cretaceous of Patagonia, Argentina. *Historical Biology* 27, 1e32. <http://dx.doi.org/10.1080/08912963.2013.861830>.
- Candeiro, C. R. A.; Fanti, F.; Therrien, F.; Lamanna, M. C. 2011. Continental fossil vertebrates from the mid-Cretaceous (Albian-Cenomanian) Alcântara Formation, Brazil, and their relationship with contemporaneous faunas from North Africa. *Journal of African Earth Sciences* 60, 79-92.
- Candeiro, C. R. A. 2015. Middle Cretaceous dinosaur assemblages from northern Brazil and northern Africa and their implications for northern Gondwanan composition. *Journal of South American Earth Sciences* 61, 147-153.
- Carrano, M. T. and Sampson, S. D. 2008. The phylogeny of ceratosauria (Dinosauria: Theropoda). *Journal of Systematic Palaeontology*, 6(2), 183-236.
- Carvalho, I. S., Salgado, L., Lindoso, R. M., Araújo-Júnior, H. I., Nogueira, F. C. C., Soares, J. A. 2017. A new basal titanosaur (Dinosauria, Sauropoda) from the Lower Cretaceous of Brazil. *Journal of South American Earth Sciences* 75, 74-84.
- Cassab, R. C. T. 2003. Paleontologia da Formação Jandaíra, Cretáceo Superior da Bacia Potiguar, com ênfase na paleobiologia dos gastrópodos. Thesis Doctoral, Instituto de Geociências, Universidade Federal do Rio de Janeiro, Rio de Janeiro, 186 p. Unpublished.

Formatted: English (United States)

- Cavin, L., Tong, H., Boudad, L., Meister, C., Piuze, A., Tabouelle, J., Aarab, M., Amiot, R., Buffetaut, E., Dyke, G., Hua, S., Leloeuff, J. 2010. Vertebrate assemblages from the early Late Cretaceous of southeastern Morocco: an overview. *Journal of African Earth Sciences* 57, 391–412.
- Contessi, M., 2009. Vertebrate remains from the Barremian–Cenomanian (Early Cretaceous) coastal deposits of South Tunisia. In: Fanti, F., Spalletta, C. (Eds.), *International Conference on Vertebrate Palaeobiogeography and Continental Bridges across Tethys, Mesogea, and Mediterranean Sea – Abstracts Book*. Museo Geologico Giovanni Capellini, Dipartimento di Scienze della Terra e Geologico-Ambientali, Bologna, Italy, pp. 31–32.
- Coria, R. A., Currie, P. J. 2006. A new carcharodontosaurid (Dinosauria, Theropoda) from the Upper Cretaceous of Argentina. *Geodiversitas* 28, 71–118.
- Coria, R. A., Salgado, L. 1995. A new giant carnivorous dinosaur from the Cretaceous of Patagonia. *Nature* 377, 224–226.
- Costa, A. B. C., Córdoba, V. C., Netto, R. G., Lima Filho, F. P. 2014. Registro faciológico e paleoambiental da transgressão que marca a passagem do Cenomaniano para o Turoniano na Bacia Potiguar, NE do Brasil. *Comunicações Geológicas*, 101(I), 415–420.
- Currie, P.J., Carpenter K. 2000. A new specimen of *Acrocanthosaurus atokensis* (Theropoda, Dinosauria) from the Lower Cretaceous Antlers Formation (Lower Cretaceous, Aptian) of Oklahoma, USA. *Geodiversitas*. 22: 207 – 246.
- Derby, A. O. 1890. Nota sobre a geologia e paleontologia de Mato Grosso. *Arquivos do Museu Nacional*. 9, 59-88.
- Duarte, L. and Santos, R. 1962. Fósseis do Arenito Açú. *Anais da Academia Brasileira de Ciências* 34(1), 57-68.

Formatted: English (United States)

- Duarte, L., Santos, R.S. 1961. Novas ocorrências fossilíferas nos estados do Rio Grande do Norte e Ceará. *Coleção Mossoroense*, 56: 11 p.
- Franco-Rosas A. C., Salgado L, Rosas C. F., Carvalho I. S. 2004. Nuevos materiales de titanosaurios (Sauropoda) em el Cretácico Superior de Mato Grosso, Brasil. *Rev Bras Paleont* 7, 329–336.
- Frey E., Martill D. M. 1995. A possible oviraptorosaurid theropod from the Santa Formation (Lower Cretaceous, Albian) of Brazil. *N Jahrb GeolPal.ont M* 7: 397–412.
- Forster, C. A., Sampson, S. D., Chiappe, L.M., Krause, D.W. 1998. The Theropod: Ancestry of Birds: New Evidence from the Late Cretaceous of Madagascar. *Science* 16 (5358), 1915-1919.
- Gauthier, J. A. 1986. Saurischian monophyly and the origin of birds; pp. 1–55 in K. Padian (ed.), *The Origin of Birds and the Evolution of Flight*. *Memoirs of the California Academy of Sciences*, 8.
- Gilmore, C. W. 1920. Osteology of the carnivorous Dinosauria in the United States National Museum, with special reference to the genera *Antrodemus* (*Allosaurus*) and *Ceratosaurus*. *Bulletin U. S. National Museum*. 110,1-154.
- Guedes, E., Heilbron, M., Vasconcelos, P. M., De Morisson Valeriano, C., Horta De Almeida, J. C., Teixeira, W., Filho, A. T. 2005. K–Ar and ⁴⁰Ar/³⁹Ar ages of dikes emplaced in the onshore basement of the Santos Basin, Resende area, SE Brazil: implications for the south Atlantic opening and Tertiary reactivation. *Journal of South American Earth Sciences* 18 (3–4), 371–382.
- Harris, J. D. 1998. Are analysis of *Acrocantnosaurus atokensis*, its phylogenetic status and paleobiogeographic implications, based on a new specimen from Texas. *Bulletin of New Mexico Museum of Natural History and Science* 13, 1-75.

1. [Hendrickx, C., Mateus, O. 2014. Abelisauridae \(Dinosauria: Theropoda\) from the Late Jurassic of Portugal and dentition-based phylogeny as a contribution for the identification of isolated theropod teeth. *Abelisauridae \(Dinosauria: Theropoda\) from the Late Jurassic of Portugal and dentition-based phylogeny as a contribution for the identification of isolated theropod teeth*. *Zootaxa*, 3759.](#)

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: List Paragraph, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"

[Hendrickx, C., Mateus, O., Araújo, R. 2015. A proposed terminology of theropod teeth \(Dinosauria, Saurischia\). *Journal of Vertebrate Paleontology*, 35\(5\), e982797.](#)

Formatted: Portuguese (Brazil)

[Kellner, A. W. A. 1996a. Fossilized theropod soft tissue. *Nature* 379, 32–32.](#)

[Kellner, A. W. A. 1996b. Remarks on Brazilian dinosaurs. *Memoirs of Queensland Museum* 39, 611–626.](#)

[Kellner, A. W. A. 1999. Short note on a new dinosaur \(Theropoda, Coelurosauria\) from the Santana Formation \(Romualdo Member, Albian\), North-eastern Brazil. *Boletim do Museu Nacional* 49, 01-08.](#)

[Kellner, A. W. A., Azevedo, S. A. K., Carvalho, L. B., Henriques, D. D. R., Costa, T., Campos, D. A. 2004. Bones out of the jungle: on a dinosaur locality from Mato Grosso, Brazil. *Journal of Vertebrate Paleontology* 124 \(Suppl3\): 78a.](#)

[Kellner, A. W. A., Azevedo, S. A., Machado, E. B., Carvalho, L. B. D. Henriques, D. D. 2011. A new dinosaur \(Theropoda, Spinosauridae\) from the Cretaceous \(Cenomanian\) Alcântara Formation, Cajual Island, Brazil. *Anais da Academia Brasileira de Ciências*, 83\(1\), 99-108.](#)

[Jokat, W., Boebel, T., König, M., Meyer, U. 2003. Timing and geometry of early Gondwana breakup. *Journal of Geophysical Research* 108 \(B9, 2428\), 1–15.](#)

Formatted: Portuguese (Brazil)

[Langer, M. C., Ribeiro, A. M., Schultz, C.L. Ferigolo, J. 2007a. The continental tetrapod-bearing Triassic of south Brazil. *Bull NMMNHS* 41, 201–218.](#)

Formatted: English (United States)

- Macdonald, D., Gomez-Perez, I., Franzese, J., Spalletti, L., Lawver, L., Gahagan, L., Dalziel, I., Thomas, C., Trewin, N., Hole, M., Paton, D. 2003. Mesozoic break-up of SW Gondwana: implications for regional hydrocarbon potential of the southern South Atlantic. *Marine and Petroleum Geology* 20 (3–4), 287–308.
- Madsen, J. H. 1976. *Allosaurus fragilis*, a revised osteology. Utah geological and mineralogical Survey Bulletin 109, 1-163.
- Marsh, O. C. 1869. Notice of some new reptilian remains from the Cretaceous of Brazil. *American Journal of Science* 47, 390–392.
- Marsh, O. C. 1878. Principal characters of American Jurassic dinosaurs. *American Journal of Science* 95, 411-416.
- Marsh, O. C. 1881. Principal characters of American Jurassic dinosaurs, part V. *American Journal of Science* 125, 417-423.
- Martill, D. M., Cruickshank, A. R. I., Frey, E., Small, P. G., Clarke, M. 1996. A new crested maniraptoran dinosaur from the Santana Formation (Lower Cretaceous) of Brazil. *Journal of the Geological Society* 153(1), 5-8.
- Martinelli, A.G., Ribeiro, L.C.B., Neto, F.M., Méndez, A.H., Cavellani, C.L., Felix, E., Ferraz, M.L.F., Teixeira, V.P.A., 2013. Insight on the theropod fauna from the Uberaba Formation (Bauru Group), Minas Gerais State: new megaraptoran specimen from the Late Cretaceous of Brazil. *Rivista Italiana di Paleontologia e Stratigrafia* 119, 205–214.
- Mateus, O., Araújo, R., Natário, C., Castanhinha, R. 2011. A new specimen of the theropod dinosaur *Baryonyx* from the early Cretaceous of Portugal and taxonomic validity of *Suchosaurus*. *Zootaxa* 2827, 54–68.

Formatted: Portuguese (Brazil)

Mawson, J., Woodward, A. S. 1907. On the Cretaceous Formation of Bahia (Brazil), and on the vertebrate fossils contained therein. *Journal of the Geological Society* 63, 128–139.

Formatted: English (United States)

Medeiros, M. A., Schultz, C. L., 2002. A fauna dinossauriana da Laje do Coringa, Cretáceo médio do Nordeste do Brasil. *Arquivos do Museu Nacional* 60 (3), 155–162.

Medeiros, M. A., Freire, P. C., Pereira, A. A., Santos, R. A. B., Lindoso, R. M., Coelho, A. F. A., Passos, E. B., Sousa, E. 2007. Another African dinosaur recorded in the Eocenomanian of Brazil and a revision on the paleofauna of the Laje do Coringa site. In: CARVALHO I.S., CASSAB R. C. T., SCHWANKE C., CARVALHO M.A., Fernandes A. C. S., Rodrigues M. A. C., Carvalho M. S. S., Arai M. And Oliveira M. E. Q. (Eds), *Paleontologia: Cenários de Vida*. Rio de Janeiro, Interciência 1, 413–423.

Motta, M. J., Aranciaga Rolando, A. M., Rozadilla, S., Agnolín, F. E., Chimento, N. R., Egli, F. B., Novas, F. E. 2016. New theropod fauna from the upper cretaceous (Huincul Formation) of Northwestern Patagonia, Argentina. *New Mexico Museum of Natural History and Science Bulletin*, 71, 231-253.

Motta, M. J., Egli, F. B., Novas, F. E. 2018. Tail anatomy of *Buitreraptor gonzalezorum* (Theropoda, Unenlagiidae) and comparisons with other basal paravians. *Cretaceous Research*, 83, 168-181.

Formatted: English (United States)

Mussa, D.; Oliveira, L.D.D., Barcia-Andrade, A. 1984. Fragmentos estélicos de *Palmae* procedentes da Formação Açu (?), Bacia Potiguar, Brasil. *Boletim do IGC-USP, Instituto de Geociências* 15, 129-141.

- Naish, D.; Martill, D. M., Frey, E. 2004. Ecology, systematics and biogeographical relationships of dinosaurs, including a new theropod from the Santana Formation (Albian, Early Cretaceous) of Brazil. *Historical Biology* 18, 01-14.
- Novas, F.E., Ribeiro, L.C.B., Carvalho, I.S., 2005a. Maniraptoran theropod ungual from the Marília Formation (Upper Cretaceous), Brazil. *Revista Museo Argentino Ciencias Naturales* 7, 31-36.
- Novas, F. E., Ezcurra, M. D., Lecuona, A. 2008. *Orkoraptor burkei* nov.gen. et sp., a large theropod from the Maastrichtian Pari Aike Formation, Southern Patagonia, Argentina. *Cretaceous Research*, 29(3), 468-480.
- Novas, F. E., Brissón Egli, F., Agnolin, F. L., Gianechini, F. A., and Cerda, I., 2017. Postcranial osteology of a new specimen of *Buitreraptor gonzalezorum* (Theropoda, Coelurosauria). *Cretaceous Research*.
- Ostrom, J. H. 1969. Osteology of *Deinonychus antirrhopus*, an Unusual Theropod from the Lower Cretaceous of Montana (Vol. 30) Peabody Museum of Natural History, Yale University.
- Pereira, P. V. L. G. C., Marinho, T. S., Candeiro, C. R. A., Bergqvist, L. P. 2018. A new titanosaurian (Sauropoda, Dinosauria) osteoderm from the Cretaceous of Brazil and its significance. *Ameghiniana* 55(6), 644-650. (71).
- Pereira, P. V. L. G. C., Veiga, I. M. M. G., Ribeiro, T. B., Cardozo, R. H. B., Candeiro, C. R. A., Bergqvist, L.P. 2020. The Path of Giants: A new occurrence of Rebbachisauridae (Dinosauria, Diplodocoidea) in the Açu Formation, NE Brazil, and its paleobiogeographic implications., *Journal of South American Earth Sciences*, <https://doi.org/10.1016/j.jsames.2020.102515>.
- Pessoa Neto, O. D. C., Soares, U. M., Silva, J. D., Roesner, E. H., Florencio, C. P. and Souza, C. D. 2007. Bacia Potiguar. *Boletim de Geociências da Petrobras* 15:357-369.

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

- Petri, S. 1987. Cretaceous paleogeographic maps of Brazil. *Palaeogeography Palaeoclimatology and Palaeoecology* 59,117–168.
- Pletsch, T., Erbacher, J., Holbourn, A., Kuhnt, W., Moullade, M., Oboh-Ikuenobede, F., Soding, E., Wagner, T. 2001. Cretaceous separation of Africa and South America: the view from the West African margin (ODP Leg 159). *Journal of South American Earth Sciences* 14, 147–174.
- Price, L. I. 1960. Dentes de Theropoda num testemunho de sonda no estado do Amazonas. *Na Academia Brasileira de Ciências* 32, 79–84.
- Price, L. I. 1961. Sobre os dinossáurios do Brasil. *Na Academia Brasileira de Ciências* 33(3–4), 28–29.
- Rauhut, O. W. M. 1999. A dinosaur fauna from the Late Cretaceous (Cenomanian) of northern Sudan. *Palaeontologia Africana* 35, 61–84.
- Reyment, R., Dingle, R. 1987. Palaeogeography of Africa during the Cretaceous period. *Palaeogeography Palaeoclimatology Palaeoecology* 59, 93–116.
- Rossetti, D. F., 1997. Internal architecture of mixed tide- and storm-influenced deposits: an example from the Alcântara Formation, northern Brazil. *Sedimentary Geology* 114 (1–4), 163–169. 115
- Rossetti, D. F., Truckenbrodt, W. 1997. Revisão estratigráfica para os depósitos do Albiano–Terciário Inferior (?) na Bacia de São Luís (MA), norte do Brasil. *Boletim do Museu Paraense Emílio Goeldi, Série Ciências da Terra* 9, 29–41.
- Russell D. A., 1972 – Ostrich dinosaurs from the Late Cretaceous of western Canada. *Canadian Journal of Earth Sciences* 9, 375–402.
- Santos, M. F. C. F.; Florêncio, C. P.; Reyes-Pérez, Y. A.; Bergqvist, L. P.; Porpino, K. O.; Uchoa, A. F. and Lima-Filho, F. P. 2005. Dinossauros na Bacia Potiguar: o

Formatted: English (United States)

[registro da primeira ocorrência. Boletim de Resumos Expandidos do XXI Simpósio de Geologia do Nordeste, Recife, 19, 325-328.](#)

[Senter, P., 2011. Using creation science to demonstrate evolution: morphological continuity within Dinosauria. Journal of evolutionary biology 24\(10\): 2197-2216. 15.](#)

[Senter, P., Kirkland, J. I., DeBlieux, D. D., Madsen, S., Toth, N., 2012. New dromaeosaurids \(Dinosauria: Theropoda\) from the Lower Cretaceous of Utah, and the evolution of the dromaeosaurid tail. PLoSOne, 7\(5\), e36790.](#)

[Serenó, P. C., Martínez, R. N., Wilson, J. A., Varricchio, D. J., Alcober, O. A. 2008. Evidence for Avian Intrathoracic Air Sacs in a New Predatory Dinosaur from Argentina. PLoS One 3, e3303.](#)

[Serenó, P. C., Brusatte, S. L. 2008. Basal abelisaurid and carcharodontosaurid theropods from the lower Cretaceous Elrhaz Formation of Niger. Acta Palaeontologica Polonica 53 \(1\), 15-6.](#)

[Silva-Santos, R. 1963. Peixes do Cretácico do Rio Grande do Norte. Anais da Academia Brasileira de Ciências 35, 67-74.](#)

[Smith, J., Lamanna, M., Lacoara, K., Dodson, P., Smith, J., Poole, J., Giegengack, R., Attia, Y. 2001. A giant sauropod from an Upper Cretaceous mangrove deposit in Egypt. Science 292, 1704-1706.](#)

[Stromer, E. 1915. Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wüsten Ägyptens. II. Wirbeltier-Reste der Baharije-Stufe \(unterstes Cenoman\). 3. Das Original des Theropoden *Spinosaurus aegyptiacus* nov. gen., nov. spec. Abhandlungen der Königlich Bayerischen Akademie der Wissenschaften, Mathematisch-physikalische Classe, 28\(3\) Abhandlung, 1-32.](#)

[Stromer, E., 1931. Wirbeltierreste der Baharije-Stufe \(unterstes Cenoman\). 10. Ein Skelett-Rest von *Carcharodontosaurus* nov. gen: Abhandlungen Bayerische](#)

Formatted: Portuguese (Brazil)

[Akademie Wissenschafte Atheilung-naturwissenschaften Abteilung Neue Folge, 9, 1–23.](#)

[Sues, H. D. 1997. On Chirosstenotes, a Late Cretaceous oviraptorosaur \(Dinosauria: Theropoda\) from western North America. Journal of Vertebrate Paleontology 17, 698–716.](#)

[Tello Saenz, C., Hackspacher, P., Hadler Neto, J., Iunes, P., Guedes, S., Ribeiro, L., Paulo, S. 2003. Recognition of Cretaceous, Paleocene, and Neogene tectonic reactivation through apatite fission-track analysis in Precambrian areas of southeast Brazil: association with the opening of south Atlantic Ocean. J South Am Earth Sci 15,765–774.](#)

[Tykoski R. S., 2005 – Anatomy, Ontogeny, and Phylogeny of Coelophysoid Theropods. Unpublished PhD Thesis, The University of Texas at Austin, 1-553.](#)

[Vasconcelos, E. P., Lima Neto, F. F., Roos, S. 1990. Unidades de correlação da Formação Açu. 36º Congresso Brasileiro de Geologia \(Natal\), Actas, 227-240.](#)

[Viramonte, J., Kay, S., Becchio, R., Escayola, M., Novitski, I. 1999. Cretaceous rift related magmatism in central-western South America. Journal of South American Earth Sciences 12, 109–121.](#)

[Werner, C., 1994. Die kontinentale Wirbeltierfauna aus der unteren Oberkreide des Sudan \(Wadi Milk Formation\). Berliner Geowissenschaftliche Abhandlungen Reihe E 13, 221–249.](#)

[Xu X., Tan Q., Wang J., Zhao X.-J., Tan L. 2007. Gigantic bird-like dinosaur from the Late Cretaceous of China. Nature, 447, 844-847.](#)

[Xu, X., Qin, Z. C. 2017. A new tiny dromaeosaurid dinosaur from the Lower Cretaceous Jehol Group of western Liaoning and niche differentiation among the Jehol](#)

Formatted: English (United States)

Formatted: Justified, Indent: Left: 0", Hanging: 0.32", Space After: 0 pt, Line spacing: Double, Font Alignment: Baseline

dromaeosaurids. *Vertebrata Palasiatica*, 55(2), 129-144.

Agnolín, F. L. & Martinelli, A.G. 2007. Didoviraptorosaurs (Dinosauria; Theropoda) inhabit Argentina? *Cretaceous Res.* 28, 785-790.

Araripe, P. T. & Feijó, F. J. 1994. Bacia Potiguar. *Boletim de Geociências da Petrobras* 8: 27-141.

Azevedo, R. P., Vasconcellos, P. L., Candeiro, C. R. A. & Bergqvist, L.P. 2007. Restos microscópicos de vertebrados fósseis do Grupo Bauru (Neocretáceo), no oeste do Estado de São Paulo, Brasil. In: Carvalho, I.S.; Cassab, R.C.; Schwanke, C.; Cavalho, M.A.; Fernandes, A.C.; Rodrigues, M.A.C.; Carvalho, M.S.; Arai, M.; Oliveira, M.E.Q.. (Org.). *Paleontologia: Cenários da Vida*. 1ed. Rio de Janeiro: Interciências, v. 2, p. 541-549.

Azevedo, R. P. F., Simbras, F. M., Furtado, M. R., Candeiro, C. R. A. & Bergqvist, L. P. 2013. First Brazilian carcharodontosaurid and other new theropod dinosaur fossils from the Campanian-Maastrichtian Presidente Prudente Formation, São Paulo State, southeastern Brazil. *Cretaceous Research*, 40, 131-142.

Benson, R. B., Carrano, M. T. & Brusatte, S. L. 2010. A new clade of archaic large-bodied predatory dinosaurs (Theropoda: Allosauroidae) that survived to the latest Mesozoic. *Naturwissenschaften*, 97(1), 71-78.

Benson, R. B. J., Butler, R. J., Carrano, M. T. & O'Connor, P. M. 2011. Air-filled postcranial bones in theropod dinosaurs: physiological implications and the "reptile" bird transition. *Biol. Rev.* number?

Bertini, R. J.. Evidências de Abelisauridae (Carnosauria: Saurischia) do Neocretáceo da bacia do Paraná. In: IV Simpósio sobre o Cretáceo do Brasil, 1996, Águas de São Pedro. *Boletim IV Simpósio sobre o Cretáceo do Brasil. Águas de São Pedro, 1996*, p. 267-271.

- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Portuguese (Brazil)
- Formatted: Font: (Default) Times New Roman, 12 pt, (Asian) Japanese
- Formatted: Indent: Left: 0", First line: 0"
- Formatted: Portuguese (Brazil)

- Formatted: German (Germany)
- Formatted: Portuguese (Brazil)

- 1276 ~~Bonaparte, J. F. 1985. A horned Cretaceous carnosaur from Patagonia. National~~
 1277 ~~Geographic Research 1: 140-151.~~
- 1278 ~~Bonaparte, J. F. & Novas, F. E., 1985, *Abelisaurus comahuensis*, n. gen., n. sp.~~
 1279 ~~Carnosauriadel Cretáceo tardio de Patagonia: Ameghiniana, v. 21, p. 259-265. Owen, R.~~
 1280 ~~1842. Report on British fossil reptiles, Pt. II. Reptiles. Report of the British Association~~
 1281 ~~for the Advancement of Science, 1841:60-204.~~
- 1282 ~~Bonaparte, J. F.; Ferigolo, J. & Ribeiro, A. M. 1999. A new Early Late Triassic~~
 1283 ~~saurischian dinosaur from Rio Grande do Sul State, Brazil. Proceedings of the Gondwana,~~
 1284 ~~v.15, p.01-10.~~
- 1285 ~~Britt, B. B. 1993. Pneumatic posterianal bones in dinosaurs and other archosaurs. Geology~~
 1286 ~~and Geophysics, University of Calgary.~~
- 1287 ~~Brum, A. S., Machado, E. B., Campos, D. A. & Kellner, A. W. A. 2016. Morphology and~~
 1288 ~~internal structure of two new abelisaurid remains (Theropoda, Dinosauria) from the~~
 1289 ~~Adamantina Formation (Turonian-Maastrichtian), Bauru Group, Paraná Basin, Brazil.~~
 1290 ~~Cretaceous Research, v. 60, p. 287-296.~~
- 1291 ~~Brum, A. S., Machado, E. B., Campos, D. A. & Kellner, A. W. A. 2018. Description of~~
 1292 ~~uncommon pneumatic structures of a noasaurid (Theropoda, Dinosauria) cervical~~
 1293 ~~vertebra to the Bauru Group (Upper Cretaceous), Brazil. Cretaceous Research, v. 85, p.~~
 1294 ~~193-206.~~
- 1295 ~~Brusatte, S. L., C.R.A. Candeiro, and F.M. Simbras. 2017. The last dinosaurs of Brazil:~~
 1296 ~~The Bauru Group and its implications for the end-Cretaceous mass extinction. Annals of~~
 1297 ~~the Brazilian Academy of Sciences 89: 1465-1485.~~
- 1298 ~~Calvo, J. O., Porfiri, J. D., Veralli, C., Novas, F. E. & Poblete, F. 2004. Phylogenetic~~
 1299 ~~status of *Megaraptornamunhuaiquii* based on a new specimen from Neuquén, Patagonia,~~
 1300 ~~Argentina. Ameghiniana 41, 565e575.~~

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

- Canale, J. I., Novas, F. E. & Pol, D. 2015. *Osteology and phylogenetic relationships of *Tyrannotitanchubutensis* Novas, de Valais, Vickers Rich and Rich, 2005 (Theropoda: Carcharodontosauridae) from the Lower Cretaceous of Patagonia, Argentina. *Historical Biology* 27, 1e32. <http://dx.doi.org/10.1080/08912963.2013.861830>.*
- Candeiro, C. R. A., Santos, A.R., Rich, T., Marinho, T.S. & Oliveira, E.C. 2006. *Vertebrate fossils from the Adamantina Formation (Late Cretaceous), Prata paleontological district, Minas Gerais State, Brazil. GEOBIOS, v. 39, p. 319-327.*
- Candeiro, C. R. 2007. *Padrões morfológicos dos dentes de Abelisauroides e Carcharodontosauridae (Theropoda, Dinosauria) do Cretáceo da América do Sul. Tese de Doutorado, Departamento de Geologia, Universidade Federal do Rio de Janeiro, 180 pp.*
- Candeiro, C.R.A., Cau, A., Fanti, F., Nava, W., Novas, F.E., 2012a. *First evidence of an unenlagiid (Dinosauria, Theropoda, Maniraptora) from the Bauru Group, Brazil. Cretaceous Research 37, 223e226.*
- Carvalho, I. S., Salgado, L., Lindoso, R. M., Araújo Júnior, H. I., Nogueira, F. C. C., & Soares, J. A. 2017. *A new basal titanosaur (Dinosauria, Sauropoda) from the Lower Cretaceous of Brazil. Journal of South American Earth Sciences, 75, 74-84.*
- Carvalho, J. C. & Santucci, R. M. 2018. *New dinosaur remains from the Quiricó Formation, Sanfranciscana Basin (Lower Cretaceous), Southwestern Brazil. CRETACEOUS RESEARCH, v. 85, p. 20-27.*
- Cassab, R. C. T. 2003. *[Paleontologia da Formação Jandaíra, Cretáceo Superior da Bacia Potiguar, com ênfase na paleobiologia dos gastrópodos. Tesis Doctoral, Instituto de Geociências, Universidade Federal do Rio de Janeiro, Rio de Janeiro, 186 p. Unpublished.*
- Colbert, E. H. 1970. *A saurischian dinosaur from the Triassic of Brazil. American Museum Novitates, v.2.405, p.01-39.*

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

- Coria, R. A., Chiappe, L. M., and Dingus L. 2002. A new close relative of *Carnotaurussastrei* Bonaparte 1985 (Theropoda: Abelisauridae) from the Late Cretaceous of Patagonia. *Journal of Vertebrate Paleontology* 22: 460–465.
- Coria, R. A. & Currie, P. J. 2006. A new carcharodontosaurid (Dinosauria, Theropoda) from the Upper Cretaceous of Argentina. *Geodiversitas* 28: 71–118.
- Coria, R. A. & Salgado, L. 1995. A new giant carnivorous dinosaur from the Cretaceous of Patagonia. *Nature* 377: 224–226.
- Coria, R. A. & Salgado, L. 1998. A basal Abelisauria Novas, 1992 (Theropoda–Ceratosauria) from the Cretaceous of Patagonia, Argentina. *Gaia* 15: 89–102.
- Costa, A. B. C., Córdoba, V. C., Netto, R. G., & Lima Filho, F. P. 2014. Registro faciológico e paleoambiental da transgressão que marca a passagem do Cenomaniano para o Turoniano na Bacia Potiguar, NE do Brasil. *Comunicações Geológicas*, 101(1): 415–420.
- Currie, P. J. & Carpenter K. 2000. A new specimen of *Acrocanthosaurus atokensis* (Theropoda, Dinosauria) from the Lower Cretaceous Antlers Formation (Lower Cretaceous, Aptian) of Oklahoma, USA. *Geodiversitas* 22: 207–246.
- Currie, P. J., Dong, Z., 2001. New information on Cretaceous troodontids (Dinosauria, Theropoda) from the People's Republic of China. *Canadian Journal of Earth Sciences*, 38(12), 1753–1766.
- Currie, P. J.; Rigby, J. K. Jr. & Sloan, R. E. 1990. Theropod teeth from the Judith River Formation of southern Alberta, Canada. // *Dinosaur systematics: approaches and perspectives*. London: Cambridge University Press: 108–125.
- Delcourt, R. & Grillo, O. N. 2014. Onmaniraptoran material (Dinosauria: Theropoda) from Vale do Rio do Peixe formation, baurugroup, Brazil. *Revista Brasileira de Paleontologia*. 17(3):307–316.

Formatted: Portuguese (Brazil)

- Derby, A. O. 1890. Nota sobre a geologia e paleontologia de Mato Grosso. Arquivos do Museu Nacional. 9, 59-88.
- Duarte, L. & Santos, R.S. 1961. Novas ocorrências fossilíferas nos estados do Rio Grande do Norte e Ceará. Coleção Mossoroense, 56, 11 p.
- Franco-Rosas A. C., Salgado L, Rosas C. f. & Carvalho I. S. 2004. Nuevos materiales de titanosaurios (Sauropoda) em el Cretáceo Superior de Mato Grosso, Brasil. Rev Bras Paleont 7: 329-336.
- Frey E. & Martill D. M. 1995. A possible oviraptorosaurid theropod from the Santa Formation (Lower Cretaceous, Albian) of Brazil. N Jahrb Geol Pal. ont M 7: 397-412.
- Forster, C. A., Sampson, S. D., Chiappe, L.M., & Krause, D.W. 1998. The Theropod: Ancestry of Birds: New Evidence from the Late Cretaceous of Madagascar. Science 16 (5358): pp. 1915-1919.
- Gauthier, J. A. 1986. Saurischian monophyly and the origin of birds; pp. 1-55 in K. Padian (ed.), The Origin of Birds and the Evolution of Flight. Memoirs of the California Academy of Sciences, 8.
- Gilmore, C. W. 1920. Osteology of the carnivorous Dinosauria in the United States National Museum, with special reference to the genera *Antrodemus* (*Allosaurus*) and *Ceratosaurus*. Bull. U. S. Natl Mus. 110, 1-154.
- Han, G., Chiappe, L. M., Ji, S. A., Habib, M., Turner, A. H., Chinsamy, A., & Han, L. 2014. A new raptorial dinosaur with exceptionally long feathering provides insights into dromaeosaurid flight performance. Nature Communications, 5, 4382.
- Harris, J. D. 1998. A reanalysis of *Acrocanthosaurus atokensis*, its phylogenetic status and paleobiogeographic implications, based on a new specimen from Texas. Bull New Mexico Mus Nat Hist Sci. 13: 1-75.

Formatted: Portuguese (Brazil)

Formatted: German (Germany)

Formatted: Portuguese (Brazil)

- Hoeknull, S. A., White, M. A., Tischler, T. R., Cook, A. G. & Calleja, N. D. 2009. New
Mid-Cretaceous (Latest Albian) Dinosaurs from Winton, Queensland, Australia. PLoS
ONE 4(7).
- Hu, D., Hou, L., Zhang, L., & Xu, X., 2009. A pre-Archaeopteryx troodontid theropod
from China with long feathers on the metatarsus. Nature 461: 640–643.
- Hwang, S. H., Norell, M. A., Qiang, J. L., & Kellner, A. W. A., 2002. New specimens of
Microraptor zhaoianus (Theropoda: Dromaeosauridae) from northeastern China.
American Museum Novitates, 1–44.
- Kellner, A. W. A. 1996a. Fossilized theropod soft tissue. Nature 379:32–32.
- Kellner, A. W. A. 1996b. Remarks on Brazilian dinosaurs. Mem. Mus. Nat. Hist. Rio de Janeiro 39:611–626.
- Kellner, A. W. A. 1999. Short note on a new dinosaur (Theropoda, Coelurosauria) from
the Santana Formation (Romualdo Member, Albian), North-eastern Brazil. Boletim do
Museu Nacional, v.49, p.01–08.
- Kellner, A. W. A. & Azevedo, S. A. K. 1999. A new sauropod dinosaur (Titanosauria)
from the Late Cretaceous of Brazil. National Science Museum Monographs, v.15, p.111–
142.
- Kellner, A. W. A. & Campos, D. A. 1996. First Early Cretaceous theropod dinosaur from
Brazil with comments on Spinosauridae. Neues Jahrbuch für Geologie und
Paläontologie Abhandlungen, 199, 151–166.
- Kellner, A. W. A. & Campos, D. A. 2002. On a theropod dinosaur (Abelisauria) from the
continental Cretaceous of Brazil. Arquivo do Museu Nacional, v.60 (3), p.163–170.
- Kellner, A. W. A., Azevedo, S. A. K., Carvalho, L. B., Henriques, D. D. R., Costa, T., &
Campos, D. A. 2004. Bones out of the jungle: on a dinosaur locality from Mato Grosso,
Brazil. J. Vertebr. Paleontol. 24 (Suppl.3): 78a.

Formatted: German (Germany)

Formatted: Portuguese (Brazil)

Formatted: German (Germany)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

- 1398 ~~Kellner, A. W. A., Azevedo, S. A., Machado, E. B., Carvalho, L. B. D. & Henriques, D.~~
 1399 ~~D. 2011. A new dinosaur (Theropoda, Spinosauridae) from the Cretaceous (Cenomanian)~~
 1400 ~~Alcântara Formation, Cajual Island, Brazil. Anais da Academia Brasileira de Ciências,~~
 1401 ~~83(1), 99–108.~~
- 1402 ~~Langer, M. C., Ribeiro, A. M., Schultz, C.L. & Ferigolo, J. 2007a. The continental~~
 1403 ~~tetrapod-bearing Triassic of south Brazil. Bull NMMNHS 41: 201–218.~~
- 1404 ~~Machado, E. B.; Campos, D. A. & Kellner, A.W.A. 2008. On a Theropoda scapula (Upper~~
 1405 ~~Cretaceous) from the Marília Formation, Bauru Group,~~
 1406 ~~Brazil. Paläontologische Zeitschrift, 82:308–313. doi:10.1007/BF02988897.~~
- 1407 ~~Madsen, J. H. 1976. *Allosaurus fragilis*, a revised osteology. Utah geol. min. Surv. Bull.~~
 1408 ~~109, 1–163.~~
- 1409 ~~Marsh, O. C. 1869. Notice of some new reptilian remains from the Cretaceous of Brazil.~~
 1410 ~~American Journal of Science, 47: 390–392.~~
- 1411 ~~Marsh, O. C. 1881. Principal characters of American Jurassic dinosaurs, part V. American~~
 1412 ~~Journal of Science, (125), 417–423.~~
- 1413 ~~Marsh, D. C. 1884. Principal characters of American Jurassic dinosaurs. Am. J. Sci.,~~
 1414 ~~27(3): 329–340.~~
- 1415 ~~Martinelli, A. G., Ribeiro, L. C. B., Neto, F. M., Méndez, A. H., Cavellani, C. L., Felix,~~
 1416 ~~E., Ferraz, M. L. F. & Teixeira, V. P. A. 2013, Insight on the theropod fauna from the~~
 1417 ~~Uberaba Formation (Bauru Group), Minas Gerais State: New megaraptoran specimen~~
 1418 ~~from the Late Cretaceous of Brazil. Revista Italiana di Paleontologia e Stratigrafia, v. 119,~~
 1419 ~~p. 205–214.~~
- 1420 ~~Sues, H. D., Frey, E., Martill, D. M., & Scott, D. M. 2002. *Irritator challengeri*, a~~
 1421 ~~spinosaurid (Dinosauria: Theropoda) from the Lower Cretaceous of Brazil.~~
 1422 ~~Journal of Vertebrate Paleontology, 22(3), 535–547.~~

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Commented [89]: This reference is wrongly placed
 A: Done

Formatted: Portuguese (Brazil)

- 1423 [Mateus O., Araújo R., Natário C. & Castanhinha R. 2011. A new specimen of the theropod](#)
 1424 [dinosaur *Baryonyx* from the early Cretaceous of Portugal and taxonomic validity of](#)
 1425 [Suchosaurus. Zootaxa 2827: 54–68.](#)
- 1426 [Mawson, J. & Woodward, A.S. 1907. On the Cretaceous Formation of Bahia \(Brazil\),](#)
 1427 [and on the vertebrate fossils contained therein. Q J Geol Soc 63:128–139.](#)
- 1428 [Medeiros, M.A., 2001. A Laje do Coringa \(Ilha do Cajual, Bacia de São Luís, Baía de](#)
 1429 [São Marcos, MA\): conteúdo fossilífero, bioestratigrafia, diagênese e implicações na](#)
 1430 [paleobiogeografia do Mesocretáceo do Nordeste Brasileiro. Instituto de Geociências da](#)
 1431 [Universidade Federal do Rio Grande do Sul, Porto Alegre, Doctoral dissertation, 107 pp.](#)
- 1432 [Medeiros, M.A., Freire, P.C., Pereira, A. A., Santos, R.A.B., Lindoso, R.M., Coelho, A.](#)
 1433 [F. A., Passos, E. B. & Sousa, E. 2007. Another African dinosaur recorded in the](#)
 1434 [Eocenomanian of Brazil and a revision on the paleofauna of the Laje do Coringa site. In:](#)
 1435 [CARVALHO I.S., CASSAB R. C. T., SCHWANKE C., CARVALHO M.A.,](#)
 1436 [FERNANDES A. C. S., RODRIGUES M. A. C., CARVALHO M. S. S., ARAI M. &](#)
 1437 [OLIVEIRA M. E. Q. \(Eds\), Paleontologia: Cenários de Vida. Rio de Janeiro, Interciência](#)
 1438 [1:413–423.](#)
- 1439 [Méndez, A.H., Novas, F.E. & Iori, F.V., 2012. First record of Megaraptora \(Theropoda,](#)
 1440 [Neovenatoridae\) from Brazil. Comptes Rendus Palevol 11, 251e256.](#)
- 1441 [Méndez, A. H., Novas, F. E. & Iori, F. V. 2014. New record of abelisauroid theropods](#)
 1442 [from the Bauru Group \(Upper Cretaceous\), São Paulo State, Brazil. Revista Brasileira de](#)
 1443 [Paleontologia, v. 17, p. 23–32,](#)
- 1444 [Motta, M., Rolando, A., Rozadilla, S., Agnolin, F., Chimento, N., Brissón, E., F., &](#)
 1445 [Novas, F. 2016. New theropod fauna from the upper cretaceous \(Huincul formation\) of](#)
 1446 [northwestern patagonia, Argentina. New Mexico Museum of Natural History and Science](#)
 1447 [Bulletin, number?](#)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

- Motta, M. J., Egli, F. B., & Novas, F. E. 2017. Tail anatomy of *Buitreraptor gonzalezorum* (Theropoda, Unenlagiidae) and comparisons with other basal paravians. *Cretaceous Research*, Number?
- Naish, D.; Martill, D. M. & Frey, E. 2004. Ecology, systematics and biogeographical relationships of dinosaurs, including a new theropod from the Santana Formation (Albian, Early Cretaceous) of Brazil. *Historical Biology*, v.18, p.01-14.
- Norell, M. A., & Makovicky, P. J., 1999. Important features of the dromaeosaurid skeleton II: information from newly collected specimens of *Velociraptor mongoliensis*. *American Museum Novitates*, no. 3282.
- Novas, F. E. 1989. Los Dinosaurios Carnívoros de la Argentina. Ph.D. Thesis. Univ. Nac. La Plata, Fac. Ciencias Naturales, 510 pp. (unpublished).
- Novas, F. E. 1992. La Evolución de los Dinosaurios Carnívoros, in SANZÓ J.L. & BUSCALIONI, A.D. (Eds.), Los Dinosaurios y su entorno biótico, Actas II Curso de Paleontología de Cuenca, Instituto "Juan de Valdes", Ayuntamiento de Cuenca, Spain, pp.125-163.
- Novas, F.E., Ribeiro, L.C.B. & Carvalho, I.S., 2005a. Maniraptoran theropod ungual from the Marília Formation (Upper Cretaceous), Brazil. *Revista Museo Argentino Ciencias Naturales* 7, 31-36.
- Novas, F. E., de Valais S., Vickers Rich P. & Rich T. 2005b. A large Cretaceous theropod from Patagonia, Argentina, and the evolution of carcharodontosaurids. *Naturwissenschaften*, 92 (5): 226-230.
- Novas, F. E., Ezeurra, M. D., & Lecuona, A. 2008. *Orkoraptor burkei* nov. gen. et sp., a large theropod from the Maastrichtian Pari Aike Formation, Southern Patagonia, Argentina. *Cretaceous Research*, 29(3), 468-480.

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

- Novas, F. E. 2009. *The Age of Dinosaurs in South America*. Indiana University Press, Bloomington, p. 452.
- Novas, F. E., BrissónEgli, F., Agnolin, F. L., Gianechini, F. A., & Cerda, I., 2017. *Postcranial osteology of a new specimen of Buitreraptor gonzalezorum (Theropoda, Coelurosauria)*. *Cretaceous Research*.
- Oliveira, E. C., Candeiro, C. R. A., & Faria, A. M. 2012. Técnica para desagregação de rochas sedimentares para recuperar restos de vertebrados das formações Adamantina e Marília (Grupo Bauru, Cretáceo Superior). *Revista do Instituto Geológico*, v. 33, p. 49-55.
- Ostrom, J. H. 1969. *Osteology of Deinonychus antirrhopus, an Unusual Theropod from the Lower Cretaceous of Montana (Vol. 30)*. Peabody Museum of Natural History, Yale University.
- Pessoa Neto, O. D. C., Soares, U. M., Silva, J. D., Roesner, E. H., Florencio, C. P. & Souza, C. D. 2007. Bacia Potiguar. *Boletim de Geociências da Petrobras* 15:357-369.
- Peyer, B. 1988. *Comparative odontology*. Chicago, University of Chicago Press, Ltd, 347 p.
- Price, L. I. 1960. Dentes de Theropoda num testemunho de sonda no estado do Amazonas. *Na Acad Bras Cienc* 32: 79-84.
- Price, L. I. 1961. Sobre os dinossáurios do Brasil. *Na Acad Bras Cienc* 33(3-4): xxviii-xxix.
- Rauhut, O. W. M. 1999. A dinosaur fauna from the Late Cretaceous (Cenomanian) of northern Sudan. *Palaeontologia Africana*, 35, 61-84.
- Russell D. A., 1972. Ostrich dinosaurs from the Late Cretaceous of western Canada. *Canadian Journal of Earth Sciences*, 9: 375-402.

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

- 1496 ~~Sampson, S. D., Witmer, L. M., Forster, C. A., Krause, D. W., O'Connor, P. M., Dodson,~~
 1497 ~~P., & Ravoavy, F. 1998. Predatory dinosaur remains from Madagascar: implications for~~
 1498 ~~the Cretaceous biogeography of Gondwana. *Science* 280: 1048–1051.~~
- 1499 ~~Sankey, J. T., Brinkman, D. B., Guenther & P.J. Currie. 2002. Small theropod and bird~~
 1500 ~~teeth from the Judith River Group (late Campanian), Alberta. *Journal of Paleontology*~~
 1501 ~~76(4):751–763.~~
- 1502 ~~Santos, M. F. C. F.; Florêncio, C. P.; Reyes Pérez, Y. A.; Bergqvist, L. P.; Porpino, K.~~
 1503 ~~O.; Uchoa, A. F. & Lima Filho, F. P. 2005. Dinossauros na Bacia Potiguar: o registro da~~
 1504 ~~primeira ocorrência. *Boletim de Resumos Expandidos do XXI Simpósio de Geologia do*~~
 1505 ~~*Nordeste, Recife*, 19: 325–328.~~
- 1506 ~~Seeley, H. G. 1888. On the classification of the fossil animals commonly named~~
 1507 ~~Dinosauria. *Proceedings of the Royal Society of London*, 43:165–171.~~
- 1508 ~~Senter, P., 2011. Using creation science to demonstrate evolution: morphological~~
 1509 ~~continuity within Dinosauria. *Journal of evolutionary biology*, 24(10), 2197–2216.~~ 15.
- 1510 ~~Senter, P., Kirkland, J. I., DeBlieux, D. D., Madsen, S., & Toth, N., 2012. New~~
 1511 ~~dromaeosaurids (Dinosauria: Theropoda) from the Lower Cretaceous of Utah, and the~~
 1512 ~~evolution of the dromaeosaurid tail. *PLoS One*, 7(5), e36790.~~
- 1513 ~~Sereno, P. C., Martínez, R. N., Wilson, J. A., Varricchio, D. J. & Alcober, O. A. 2008.~~
 1514 ~~Evidence for Avian Intrathoracic Air Sacs in a New Predatory Dinosaur from Argentina.~~
 1515 ~~*PLoS One* 3: e3303.~~
- 1516 ~~Smith, J. & Dodson, P. 2003. A proposal for standard terminology of anatomical notation~~
 1517 ~~and orientation in fossil vertebrate dentitions. *Journal of Vertebrate Paleontology*, v.23,~~
 1518 ~~n.1 p. 1–12.~~

Formatted: Portuguese (Brazil)

Formatted: German (Germany)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Smith, J. B., Vann, D. R. & Dodson, P. 2005. Dental morphology and variation in theropod dinosaurs: implications for the taxonomic identification of isolated teeth. *The Anatomical Record*, 285(2), 699-736.

Formatted: German (Germany)

Stromer, E. 1915. Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wüsten Ägyptens. II. Wirbeltier-Reste der Baharije Stufe (unterstes Cenoman). 3. Das Original des Theropoden *Spinosaurus aegyptiacus* nov. gen., nov. spec. *Abhandlungen der Königlich Bayerischen Akademie der Wissenschaften, Mathematisch-physikalische Classe*, 28(3) Abhandlung: 1-32.

Stromer, E., 1931. Wirbeltierreste der Baharije Stufe (unterstes Cenoman). 10. Ein Skelett-Rest von *Carcharodontosaurus* nov. gen. *Abhandlungen Bayerische Akademie Wissenschaften Abteilung Naturwissenschaften Abteilung Neue Folge*, v. 9, p. 1-23.

Sues, H. D. 1997. On *Chirostenotes*, a Late Cretaceous oviraptorosaur (Dinosauria: Theropoda) from western North America. *Journal of Vertebrate Paleontology* 17: 698-716.

Formatted: Portuguese (Brazil)

Sues, H. D., Frey, E., Martill, D. M., & Scott, D. M. 2002. *Irritator challengeri*, a spinosaurid (Dinosauria: Theropoda) from the Lower Cretaceous of Brazil. *Journal of Vertebrate Paleontology*, 22(3), 535-547.

Commented [90]: This reference is wrongly placed

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Tykoski R. S., 2005. Anatomy, Ontogeny, and Phylogeny of Coelophysoid Theropods. Unpublished PhD Thesis, The University of Texas at Austin: 1-553.

Vasconcelos, E. P., Lima Neto, F. F., & Roos, S. 1990. Unidades de correlação da Formação Açu. 36º Congresso Brasileiro de Geologia (Natal), Actas: 227-240.

Vilas Bôas, I., Carvalho, I.S., Medeiros, M.A. & Pontes, H., 1999. Dentes de *Carcharodontosaurus* (Dinosauria, Tyrannosauridae) do Cenomaniano, Bacia de São Luís (Norte do Brasil). *Anais da Academia Brasileira de Ciências* 71 (4 D), 846-847.

~~Xu, X., Norell, M. A., Wang, X. L., Makovicky, P. J., & Wu, X. C., 2002. A basal troodontid from the Early Cretaceous of China. *Nature*, 415(6873), 780-784.~~

Formatted: Portuguese (Brazil)

~~Xu X., Tan Q., Wang J., Zhao X. J. & Tan L. 2007. Gigantic bird-like dinosaur from the Late Cretaceous of China. *Nature*, 447: 844-847.~~

Formatted: German (Germany)

Formatted: Portuguese (Brazil)

~~Xu, X., & Qin, Z. C., 2017. A new tiny dromaeosaurid dinosaur from the Lower Cretaceous Jehol Group of western Liaoning and niche differentiation among the Jehol dromaeosaurids. *Vertebrata Palasiatica*, 55(2), 129-144.~~

Figure captions

Figure 01: Geological map of the continental part of the Potiguar Basin with the region near the Limoeiro do Norte municipality (Ceará state) where the material were discovered (dark star). CE, Ceará state; RN, Rio Grande do Norte state and its capital, Natal. Modified from Cassab (2003).

Formatted: Portuguese (Brazil)

Figure 02: The ~~avirostran~~theropod vertebrae UFRJ-DG 52875-R (A-~~CB~~) and UFRJ-DG 57528-R (~~DC~~-E). UFRJ-DG 575-R: A, lateral view; B, the anterior articular facet. UFRJ-DG 528-R: C, the lateral view; D, the ventral view; E, anterior articular facet. Note the large pneumatic foramen on the side of the anterior fragment of UFRJ-DG 575-R. pfr = pneumatic foramen. Scale bar: 2 cm.

Figure 03: Spinosauroid tooth (UFRJ-DG 619-R): A, the labial view; B, the lingual view; and C, the cross section. Scale: 1 cm

Figure 04: Maniraptoran caudal vertebrae (UFRJ-DG 521-R): A, Lateral view; B, ventral view; C, anterior articular facet. Prz., prezygophysis; Nc, neural canal. Scale: 1cm.

Commented [91]: I think the aspect of the figure is not proper: it shows different fonts, the letters are unaligned, and shows two scale bars. The same applies for Figure 4
A: Done.

Figure 05: Comparison of UFRJ-DG 521-R and other maniraptorans. A, Potiguar's material; B, *Rahonavis*; C, *Buitreraptor*; D, *Anchiornis*. Pr, prezygapophysis; lg, Longitudinal groove. Modified from Motta et al., (2018).

Commented [92]: In the references the date of the publication is 2017
A: Done.

Figure 06: Carcharodontosaurid caudal vertebrae UFRJ-DG 5234(A-C) and UFRJ-DG 5243-R (D-F). UFRJ-DG 524-R: A, ventral view; B, lateral view; C, anterior articular facet. UFRJ-DG 523-R: D, ventral view; E, lateral view; F, anterior articular facet. Nc, neural canal. Scale: 1cm.

Figure 07: Comparison of UFRJ-DG 523-R and 524-R and other carcharodontosaurids. A and B, UFRJ DG 523-R; C and D, UFRJ DG 524-R; E and F, Kem Kem beds material (from Rauhut, 1999); G, *Tyrannotitan chubutensis* MPEF-PV 1156 (Modified from Canale et al., 2015); H, *Mapusaurus roseae* MCF-PVPH-108.81 (Modified from Coria & Currie, 2006) ; I, *Acrocanthosaurus atokensis* NCSM 14345 (Modified from Currie & Carpenter, 2000). Scale bar = 5 cm.

Commented [93]: The item G of the figure has a very poor resolution. Also G, H, I were taken from the literature, and the figure caption lack the proper cite.

Figure 08: Caudal vertebrae UFRJ-DG. UFRJ-DG 558-R-634-R: A, posterior articular facet; B, lateral view; C, ventral view. UFRJ-DG 634-R-558-R: D, anterior articular facet; E, lateral view; F, ventral view. Pfr, Pneumatic foramen. Scale bar: 1cm.

Figure 09: Brazilian megaraptoran vertebrae findings. A and B, UFRJ DG 558-R; C and D, MPMA 08-003-94 (Méndez et al., 2012); E and F, CPPLIP 1324 (Martinelli et al., 2013). A, C e E, lateral view; B, D e F, ventral view. Pfr, Pneumatic foramen. Scale bar = 1cm.

Formatted: Highlight

Figure 10: Megaraptoran caudals vertebrae. A and B, UFRJ DG 558-R; C and D, *Aerosteon*; E and F, *Aoniraptor*; G, *Orkoraptor*. H, *Megaraptor*. A, C, E, G e H, lateral view; B, D e F, ventral view. Pfr, pneumatic foramen. Scale bar = 5cm.

Formatted: Portuguese (Brazil)

Figure 11: Reconstruction of the theropods groups from Açú Formation, Potiguar Basin. In the center, a group of megaraptorans slaughtering a titanosaur; on the right a carcharodontosaurid awakens from its sleep; in the top center, a maniraptoran just watches. DrawingArt by Luciano da Silva Vidal.

Figure captions

Figure 01: Geological map of the continental part of the Potiguar Basin with the region near the Limoeiro do Norte municipality (Ceará state) where the material were discovered (dark star). CE, Ceará state; RN, Rio Grande do Norte state. Modified from Cassab (2003).

Formatted: English (United States)

Figure 02: The avirostran vertebrae UFRJ DG 575 R (A-B) and UFRJ DG 528 R (C-E). UFRJ DG 575 R: A, lateral view; B, the anterior articular facet. UFRJ DG 528 R: C, the lateral view; D, the ventral view; E, anterior articular facet. Note the large pneumatic foramen on the side of the anterior fragment of UFRJ DG 575 R. pfr = pneumatic foramen. Scale bar: 2 cm.

Figure 03: Spinosauroid tooth (UFRJ DG 619 R): A, the labial view; B, the lingual view; and C, the cross section. Scale: 1 cm

Figure 04: Maniraptoran caudal vertebrae (UFRJ DG 521 R): A, Lateral view; B, ventral view; C, anterior articular facet. Prz., prezygophysis; Ne, neural canal. Scale: 1cm.

Commented [94]: I think the aspect of the figure is not proper: it shows different fonts, the letters are unaligned, and shows two scale bars. The same applies for Figure 4 A: Done.

Formatted: English (United States)

Figure 05: Carcharodontosaurid caudal vertebrae UFRJ DG 524 (A-C) and UFRJ DG 523 R (D-F). UFRJ DG 524 R: A, ventral view; B, lateral view; C, anterior articular facet. UFRJ DG 523 R: D, ventral view; E, lateral view; F, anterior articular facet. Ne, neural canal. Scale: 1cm.

Figure 06: Caudal vertebrae UFRJ DG. UFRJ DG 634 R: A, posterior articular facet; B, lateral view; C, ventral view. UFRJ DG 558 R: D, anterior articular facet; E, lateral view; F, ventral view. Pfr, Pneumatic foramen. Scale bar: 1cm.

Figure 07: Brazilian megaraptoran vertebrae findings. A and B, UFRJ DG 558 R; C and D, MPMA-08-003-94 (Méndez et al., 2012); E and F, CPPLIP 1324 (Martinelli et al., 2013). A, C e E, lateral view; B, D e F, ventral view. Pfr, Pneumaticforamen. Scale bar = 1cm.

Formatted: English (United States)

Figure 08: Megaraptoran caudals vertebrae. A and B, UFRJ DG 558 R; C and D, Aerosteon; E and F, Aoniraptor; G, Orkoraptor. H, Megaraptor. A, C, E, G e H, lateral view; B, D e F, ventral view. Pfr, pneumatic foramen. Scale bar = 5cm.

Figure 09: Comparison of Morphotype 3 and other paravians. A, Potiguar's material; B, *Rahonavis*; C, *Buitreraptor*; D, *Anchiornis*. Pr, prezygapophysis; lg, Longitudinal groove. Modified from Motta et al., (2018).

Commented [95]: In the references the date of the publication is 2017
A: Done.

Figure 10: Comparison of UFRJ DG 523 R and 524 R and other carcharodontosaurids. A and B, UFRJ DG 523 R; C and D, UFRJ DG 524 R; E and F, Kem Kem beds material (from Rauhut, 1999); G, *Tyrannotitan chubutensis* MPEF PV 1156 (Modified from Canale et al., 2015); H, *Mapusaurus roseae* MCF PVPH 108.81 (Modified from Coria & Currie, 2006) ; I, *Aerocanthosaurus atokensis* NCSM 14345 (Modified from Currie & Carpenter, 2000). Scale bar = 5 cm.

Commented [96]: The item G of the figure has a very poor resolution. Also G, H, I were taken from the literature, and the figure caption lack the proper cite.

- Formatted: Font: Italic
- Formatted: Font: Italic
- Formatted: Font: Not Bold, English (United States)
- Formatted: Font: Italic
- Formatted: Font: Italic
- Formatted: English (United States)
- Formatted: Font: Italic
- Formatted: Font: Italic
- Formatted: English (United States)
- Formatted: English (United States)

Figure 11: Reconstruction of the theropods groups from Açu Formation, Potiguar Basin. On in the left there is an abelisaurid dinosaur; in the center, a group of megaraptorans slaughtering a titanosaur; on the right a carcharodontosaurid awakens from its sleep; in the top center, a maniraptoran just watches. Drawing by Luciano da Silva Vidal.