THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Recombinant canine IgE Fc and an IgE Fc-TRAIL fusion protein bind to neoplastic canine mast cells

Citation for published version:

Elders, RC, Holder, A, Smith, KC, Baines, SJ & Catchpole, B 2014, 'Recombinant canine IgE Fc and an IgE Fc-TRAIL fusion protein bind to neoplastic canine mast cells', *Veterinary Immunology and Immunopathology*, vol. 159, no. 1-2, pp. 29-40. <https://doi.org/10.1016/j.vetimm.2014.02.018>

Digital Object Identifier (DOI):

[10.1016/j.vetimm.2014.02.018](https://doi.org/10.1016/j.vetimm.2014.02.018)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Veterinary Immunology and Immunopathology

Publisher Rights Statement:

This is the final accepted version of the article. The publisher version is available here: <http://www.sciencedirect.com/science/article/pii/S0165242714000543>

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Recombinant canine IgE Fc and an IgE Fc-TRAIL fusion protein bind to neoplastic canine mast cells

R.C. Elders^{a,*}, A. Holder^b, K. C. Smith^b, S.J. Baines^c, B. Catchpole^b

^a *Hospital for Small Animals, Royal (Dick) School of Veterinary Studies; University of Edinburgh; Easter Bush; Midlothian; EH25 9RG; UK*

^b *Department of Pathology and Infectious Diseases, Royal Veterinary College; Hawkshead Lane; North Mymms; Hatfield; Hertfordshire; AL9 7TA; UK*

^c *Willows Veterinary Centre and Referral Service; Highlands Road; Solihull; West Midlands; B90 4NH; UK*

* Corresponding author

Tel: +44 1638 751 000

Fax: +44 1638 555 600

E-mail address: richard.elders@aht.org.uk

Keywords: mast cell tumor; IgE; immunoconjugate; immunotherapy; TRAIL

Abbreviations: PCR, polymerase chain reaction; cMCT, canine mast cell tumor.

Abstract

Screening for expression of the high affinity receptor for IgE by reverse transcriptase PCR, revealed that almost all canine mast cell tumors expressed FcεRIα mRNA, supporting the rationale for developing anti-neoplastic treatments based on molecules that could target this receptor. Use of cytotoxic cytokines to trigger an apoptotic signal is one strategy for inducing cell death in malignant mast cells. The coding sequences for canine IgE and tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) were identified through genome analyses. Selected regions of the coding sequences for these genes were cloned and compared to the predicted genome sequences. The Fc region of canine IgE, death domain of canine TRAIL and an IgE Fc : TRAIL fusion construct were generated and epitope-tagged proteins expressed, using a eukaryotic expression system. Specific binding of recombinant canine IgE Fc-containing proteins to recombinant human FcεRIα and to a canine mast cell tumor line expressing FcεRIα (C2), but not one failing to express FcεRIα (MCLA), was demonstrated. Specific binding of the IgE:TRAIL fusion protein was not abrogated by the TRAIL moiety. These results are proof of principle that canine IgE targeting to FcεRIα can be used as a platform for selective delivery of therapies to FcεRIα-expressing cells, potentially enhancing their therapeutic index and efficacy.

1. Introduction

Canine mast cell tumors (cMCTs) are the most common skin malignancy in dogs (Brodey 1970; Finnie and Bostock, 1979; Bostock 1986; Dobson et al., 2002), likely arising from neoplastic transformation of resident tissue mast cells or their progenitors. Breed, location, stage, gross appearance, size, presence of paraneoplastic syndromes and several proliferation markers have been variably associated with prognosis, however the most consistent predictor of cMCT recurrence, metastasis and survival is histopathological grade (Bostock, 1973; Patnaik et al., 1984; O' Keefe, 1990; Gerritsen et al., 1998; Mullins et al., 2006). Although dogs with a localised cMCT are often cured by local therapy (surgery and/or radiation therapy), those with an inoperable primary mass or confirmed disseminated disease usually die of their disease (reviewed by Welle et al., 2008; Blackwood et al., 2012). The use of both local (e.g. surgery and radiation therapy) and systemic therapies (e.g. chemotherapy, receptor tyrosine kinase inhibitors) is limited by the potential for adverse effects, some of which can be idiosyncratic, cumulative and/or permanent in nature. Thus, there is a need to develop novel therapies for cMCTs with improved efficacy and higher therapeutic indices.

Mast cells avidly and specifically bind to IgE via the high-affinity IgE receptor, FcεRI. Cross-linkage of surface IgE by antigen leads to mast cell degranulation and inflammation, an important defence against parasite infestation. In addition to disease caused by malignant transformation, mast cells are also involved in type I hypersensitivity reactions in allergic disease. Additional beneficial and pathophysiological roles for mast cells are emerging, although many of these can be undertaken by other, more numerous cells (reviewed by Rao & Brown, 2008), and mast cells are not absolutely required for survival, as demonstrated by genetically-modified mouse strains lacking these cells (C57BL/6-KITW^{sh}-/W^{sh}-, Grimaldeston et al., 2005). However, other genetic mutations associated with mast cell deficiency can lead to significant developmental problems in cells of other lineages. WBB6F₁-W/W^v mice have reduced erythrocyte, granulocyte, platelet and mast cell

numbers (Kitamura et al., 1978), and although such mice are more likely than wild type mice to die in a model of acute septic peritonitis involving cecal ligation and puncture, their survival can be enhanced solely by reconstitution with cultured wild type mast cells (Echtenacher et al., 1996).

FcεRI is constitutively expressed by mast cells and basophils during early differentiation, but lower level, inducible expression has also been found on mammalian eosinophils, monocytes, platelets and dendritic cells (Thompson et al., 1990; Rottem et al; 1992; Joseph et al., 1997; Kinet 1999; Kita et al., 1999; Seminario et al., 1999). The affinity and avidity of IgE for FcεRI exceeds that of other immunoglobulin/Fc receptor interactions by several orders of magnitude (Ravetch and Kinet, 1991; Maenaka et al., 2001; McDonnell *et al.*, 2001; Wan et al., 2002). Previous work, evaluating the role of each IgE domain in binding to its high affinity receptor, has shown that C_Hε3 is critical for binding to FcεRI, but C_Hε2 and C_Hε4 are also required for high affinity and avidity binding (Keown et al., 1997; Garman et al., 2000 & 2001; McDonnell et al., 2001; Wurzburg and Jardetzky, 2002; Hunter et al., 2008). In a previous study, a partially dimerized C_Hε2-4 recombinant canine protein was used to raise an anti-canine IgE antibody which detected native canine IgE, suggesting that the quaternary structure of the recombinant C_Hε2-4 protein was similar to native canine IgE, despite the lack of a C_Hε1 domain (Ledin et al., 2006). Hunter et al. (2008) reported an alternative cell-based binding assay using RBL-2H3 cells stably transfected with rcFcεRIα and elegantly evaluated the role of each IgE heavy chain domain in binding to the receptor.

IgE has been used as a delivery platform for targeting murine and human mast cells, when linked to Pseudomonas toxin or pro-apoptotic Bcl-2 protein family members. These fusion proteins were capable of binding to and inducing apoptosis in mast cells *in vitro* and *in vivo*. (Fishman and Lorberboum-Galski, 1997; Fishman et al., 2000; Belostotsky and Lorberboum-Galski, 2001) A rmlgE₃₀₁₋₄₇₃-Pseudomonas exotoxin A chimeric protein was highly cytotoxic to both malignant murine mast cell lines and bone marrow-derived mast cells, while having no effect on other cell lines of

haematopoietic lineage (Fishman and Lorberboum-Galski, 1997), demonstrating its specificity for FcεRI-expressing cells.

Tumor necrosis factor-related apoptosis-inducing ligand (TRAIL, TNFSF10) is a type II membrane protein, an extracellular fragment of which can be cleaved to generate soluble TRAIL (sTRAIL; amino acids 37-281) (Wiley et al., 1995; Pitti et al., 1996; Mariani and Krammer, 1998). In humans and rodents, healthy cells display a repertoire of death and decoy TRAIL receptors, whereas neoplastic/transformed cells preferentially express death receptors over decoy receptors (Sheridan et al., 1997; Simonet et al., 1997; Pan et al., 1998; LeBlanc and Ashkenazi, 2003). Therefore, neoplastic cells are generally more susceptible than healthy cells to TRAIL-mediated apoptosis, making this a tempting therapeutic strategy (LeBlanc and Ashkenazi, 2003). Although the canine TRAIL receptor repertoire is poorly characterized and there are no TRAIL receptor genes identifiable in the syntenic region of the canine genome, compared to human and mouse, we and others have demonstrated that canine neoplastic mast cells are susceptible to the apoptotic effects of TRAIL (Rong et al., 2008; Elders et al., 2009).

The aim of the current study was to generate a recombinant canine IgE Fc protein that was capable of binding to the FcεRI receptor expressed on canine mast cells. The intention was that this recombinant canine IgE Fc molecule could then be modified to allow specific targeting approaches to be evaluated for treatment of canine mast cell tumors. Since we had previously shown that a canine MCT line (C2) was susceptible to TRAIL-mediated apoptosis, a canine IgE-TRAIL fusion protein was planned in the first instance to test the concept that this targeting strategy was feasible.

2. Materials and methods

2.1 Case recruitment and sample collection

Biopsy samples were obtained from dogs undergoing surgical resection of a mass, suspected or cytologically confirmed as cMCT. Tissue was stored in RNA^{later}® (Qiagen, Crawley, UK) at -20 °C prior to molecular analysis. All tumors had histopathology or cytology performed and, where available, representative slides were reviewed by a single pathologist (K.C.S.) using a published grading scheme (Patnaik et al., 1984). In one patient, tumor tissue was disaggregated and cultured to generate a novel cMCT line, designated MCLA, which demonstrated metachromatic granule staining with Toluidine blue and was positive for expression of chymase and tryptase (Elders, 2009). Residual canine EDTA blood, following completion of processing of a diagnostic sample, was used to provide canine genomic DNA for IgE cloning. Similarly, residual lymph node tissue, following diagnostic processing of a lymphoma sample, was used to provide cDNA for TRAIL cloning.

2.2 Cells and cell culture

Chinese Hamster ovary (CHO) cells (obtained from ECACC), C2 canine mastocytoma cells (a generous gift from Dr. Birgit Helm, University of Sheffield; permitted by the originator, Prof. W. Gold, University of California) (Lazarus et al., 1986) or MCLA cells were propagated at 37 °C, 5% CO₂, in 75 cm² flasks in Eagle's minimal essential medium, supplemented with 5% foetal bovine serum (FBS), 1% non-essential amino acids, 50 µg/ml gentamicin (all Sigma) and 1% L-glutamine (Invitrogen, Paisley, UK) (culture medium). For experiments, cells were dissociated using Accutase™ (PAA Laboratories, Hampshire, UK) and cultured in phenol red-free minimal essential medium (Invitrogen), supplemented with 10% FBS, 1% non-essential amino acids, 1% L-glutamine and 50 µg/ml gentamicin.

2.3 Nucleic acid extraction and polymerase chain reaction

Tumor tissue or cultured cells were homogenized in Lysis Solution for total RNA (Sigma, Poole, UK) and RNA extracted using the GenElute™ Mammalian Total RNA Miniprep Kit (Sigma). On-column (RNase-free DNase set, Qiagen) and in-solution (Turbo DNase™, Ambion, Austin, USA) digestion

steps were performed to remove contaminant genomic DNA. Reverse transcription of mRNA into complementary DNA (cDNA) was performed using oligo(dT)₁₅ primer and ImProm-II reverse transcriptase (Promega, Southampton, UK).

Polymerase chain reaction was used to amplify genes of interest using specific primers (Table 1). Each 25 µl reaction contained 1 µl both sense and anti-sense primers (final concentration 10 pmol/µl) and cDNA or gDNA (1 µl) as template. Reactions also contained 1× Hi-Spec Additive, 1× NH₄ buffer, MgCl₂ (final concentration 2.0-3.0 mM), dNTP (final concentration 10 mM) and 0.5 IU Immolase™ DNA polymerase per reaction (all from Bioline, London, UK), made up to 25 µl with water. Reactions were heated to 95 °C for 10 mins, followed by 20 - 35 cycles at 94 °C for 40 s, 55 – 62.5 °C for 30 s, and 72 °C for 1 min; with a final extension step at 72 °C for 7 mins using a G Storm thermocycler (Gene Technologies Ltd, Essex, UK). For PCR products that were to be cloned, a proof-reading DNA polymerase (Easy-A® High-Fidelity PCR Cloning Kit, Stratagene, Amsterdam, The Netherlands) was used in place of Immolase.

Horizontal gel electrophoresis was used to separate PCR products, using 1-2% agarose (Bioline) gels containing 1× SafeView Nucleic acid stain™ (NBS Biologicals Huntingdon, UK). Amplicons were visualized under 590 nm ultra-violet light, using the ImageMaster® VDS Gel Documentation System (Pharmacia Biotech, Uppsala, Sweden).

2.4 Cloning of inserts into plasmid DNA vectors

PCR amplicons were extracted (GenElute™ gel extraction kit; Sigma), cloned into the pSC-A® vector (Stratagene) and transformed into *E. coli* (Solopack™, Stratagene). Plasmid DNA was then isolated using the GenElute™ Plasmid Miniprep Kit (Sigma) and sequencing to confirm the integrity of the inserts (Geneservice, Cambridge UK). Donor (recombinant pSC-A) and recipient (pSecTagA®; Invitrogen) plasmid DNA were digested using the indicated restriction enzymes to allow directional subcloning. Following purification, insert DNA was ligated into digested recipient vector (Quick-Stick™ DNA Ligase, Bioline) and *E. coli* transformed. Endotoxin-free plasmid DNA was prepared from

E. coli cultures using GenElute™ Endotoxin-free Plasmid Midiprep Kit (Sigma), according to the manufacturer's instructions.

2.5 Transfection of Chinese hamster ovary cells

Cells for transfection were seeded at 1×10^5 cells per well into 24-well plates in a 500 μ l volume of culture medium containing 5% FBS. When 90-95% confluent, the medium was replaced with FBS and antibiotic-free medium. Cells were transfected with 800 ng endotoxin-free recombinant plasmid DNA using Lipofectamine™ 2000 (Invitrogen) according to the manufacturer's instructions. Plates were incubated for 6 h then the plasmid-containing medium was replaced with medium containing 10% FBS but lacking antibiotics, and after a further 18 h this medium was replaced with culture medium. After 72 h incubation, the supernatant, containing recombinant protein, was recovered, centrifuged ($13.8 \times g$, 5 mins), filtered (0.22 μ m filters, Millex® GV syringe filter unit, Sigma), and used for experiments.

2.6 Detection of recombinant proteins

Pre-blocked, nickel coated ELISA strips (Ni-NTA HiSorb®, Qiagen) were used to capture polyhistidine-tagged recombinant proteins from transfected CHO supernatants (100 μ l/well). After 2 h incubation, wells were washed with PBS supplemented with 0.1% Tween 20 (Sigma). A caprine anti-c-myc:HRP conjugate (A190-104P; Universal Biologicals, Cambridge, UK), diluted 1:5,000 in phenol red-free MEM/10% FBS was added for 2 h, followed by six washes. One hundred microliters of substrate (Supersensitive TMB Liquid Substrate for ELISA; Sigma) was added per well and incubated for 20 min. Twenty five microliters of 0.5 M sulphuric acid (SLS, Nottingham, UK) was added to stop the reaction, and the optical density (O.D.) at 540 nm subtracted from that at 450 nm to generate absorbance values per well. The ELISA was modified by substituting the anti-c-myc antibody with either a polyclonal caprine anti-canine IgE:HRP conjugate (1:10,000 dilution of AHP946P, Abd Serotec Ltd., Oxford, UK), or a polyclonal caprine anti-human TRAIL antibody (1:100

dilution of AF375, AbD Serotec) which was in turn detected using a polyclonal anti-caprine IgG:HRP conjugate (1:10,000 dilution of OBT1500P, AbD Serotec). Samples of transfected CHO supernatants, containing recombinant protein, were also submitted to HESKA (Fribourg, Switzerland) for analysis using their validated Allercept® canine IgE ELISA (Foster et al., 2003).

2.7 Western blotting

Polyhistidine-tagged proteins were purified from CHO supernatants using the MagneHis™ Protein Purification System (Promega). Purified recombinant proteins (32.5 µl) were added to 5 µl of 500 mM dithiothreitol (DTT) reducing agent and 12.5 µl NuPAGE® LDS sample buffer (both Invitrogen) and incubated at 70 °C for 10 mins. Proteins were separated by PAGE using 4 - 20% gels and sodium dodecyl sulphate (SDS) running buffer (both PAGEgel.com, California, USA), in the X-cell SureLock™ Mini-cell (Invitrogen) at 200 V constant for 35 mins. Proteins were then transferred to nitrocellulose membranes under reducing conditions in the X-Cell II™ Blot Module (Invitrogen) and transfer buffer (PAGEgel.com) at 30 V constant for 1 h. Membranes were rinsed, blocked overnight at 4 °C in 50 ml PBS/5% Marvel, then incubated with murine anti-His:HRP conjugate (MCA1396P, AbD Serotec, diluted 1:2500 in 25 ml PBS/5% Marvel™/0.1% Tween 20) at room temperature for 1 h. After washing, immunoreactivity was detected by chemiluminescence using the ECL™ Western blotting analysis system (GE Healthcare, Chalfont St Giles, UK) and film (Kodak X-Omat AR™, Kodak, Harrow, UK).

2.8 Assessment of FcεRIα receptor-binding activity of recombinant proteins by flow cytometry

C2 (FcεRIα positive mastocytoma line) and MCLA (FcεRIα negative mastocytoma line) cells (2×10^5 cells in a 100 µl volume of culture medium) were incubated at 4 °C for 30 mins. Cells were then centrifuged at 1200 ×g for 10 mins, resuspended and incubated at 4 °C for 2 h in 200 µl of each sterile-filtered CHO cell supernatant (transfected with either IgE₁₀₀₋₄₂₇, TRAIL₁₁₄₋₂₈₂, or IgE₁₀₀₋₄₂₇:TRAIL₁₁₄₋₂₈₂ fusion construct). Cells resuspended in supernatant from mock transfected CHO cells

acted as negative controls. Cells resuspended in PBS/0.5% canine IgE heterohybridoma supernatant (Bethyl Laboratories) were used as positive controls for IgE binding.

Cells were centrifuged and supernatants aspirated. Cell pellets were resuspended in 200 μ l PBS alone or containing caprine anti-canine IgE:FITC (1:200 of AHP946F, AbD Serotec) or murine anti-c-myc:FITC (1:200 of MCA2200F, AbD Serotec) and incubated for 20 mins. Following repeat centrifugation, cell pellets were resuspended in 200 μ l PBS for flow cytometric analysis, (FACSaria, BD Biosciences, Erembodegem, Belgium) counting up to 20,000 events. Data were analysed using FlowJo (Tree Star Inc., Oregon, USA)

3. Results

3.1 Analysis of Fc ϵ R1 α mRNA expression in canine mast cell tumors and selected mastocytoma cell lines

The canine Fc ϵ R1 α coding sequence (762 bp), consisting of 5 exons, is located on chromosome 38 (GenBank Accession# NM_00110766.1, Goitsuka et al., 1999). Screening primers, designed to amplify a 273bp fragment, were used to demonstrate that 23 of the 25 canine MCT biopsy samples were positive for Fc ϵ R1 α mRNA expression (Fig. 1A), suggesting that this represents a suitable target for potential immunotherapeutic strategies. The C2 mastocytoma cell line but not the newly established MCLA mast cell tumor line, expressed Fc ϵ R1 α mRNA (Fig. 1B).

3.2 Generation of recombinant canine IgE Fc and IgE Fc-TRAIL fusion constructs

The canine IgE heavy chain gene, encoding V ϵ H and C ϵ H1-4, is located on chromosome 8 and consists of 5 exons (GenBank Accession# XM_548007.1). Using genomic DNA as template, primers were designed at the start of exon 3 (encoding C ϵ H2) and the end of exon 5 (encoding C ϵ H4), to generate a PCR amplicon spanning the C ϵ H2-4 coding region, but which contained additional intronic sequence. This was cloned into the pSecTagA vector and following sequence verification, plasmid

DNA was transfected into Chinese hamster ovary (CHO) cells. Isolation of cDNA from the transfected cells and further PCR analysis revealed three amplicons (Fig. 2), the smallest of which was consistent with splicing out of both introns. This was confirmed by sequencing, and this canine C_Hε2-4 coding region was subsequently cloned into the pSecTagA vector between the *HindIII* and *EcoRI* sites (pSecTagA/rcIgE₁₀₀₋₄₂₇).

The canine TRAIL gene, consisting of 5 exons is located on chromosome 34 (GenBank Accession# NM_001130836.1). Using cDNA prepared from a canine lymphoma biopsy as template, canine TRAIL-specific primers were used to generate an amplicon encoding the predicted C-terminal TNF-like death domain (amino acids 114-282), which was cloned into pSecTagA (pSecTagA/rcTRAIL₁₁₄₋₂₈₂). In addition, a fusion construct was generated by subcloning the TRAIL sequence, between *EcoRI* and *XhoI* sites, downstream of and in-frame with the IgE Fc fragment to generate a fusion construct (pSecTagA/rcIgE₁₀₀₋₄₂₇ : rcTRAIL₁₁₄₋₂₈₂).

3.3 Expression of recombinant canine IgE Fc and IgE Fc-TRAIL fusion proteins

Chinese hamster ovary cells were transfected with pSecTagA containing, rcIgE₁₀₀₋₄₂₇, rcTRAIL₁₁₄₋₂₈₂, or rcIgE₁₀₀₋₄₂₇ : rcTRAIL₁₁₄₋₂₈₂ fusion constructs. Polyhistidine and c-myc epitope-tagged recombinant proteins were detected by ELISA in the supernatants of CHO cells transfected with the three constructs (Fig. 3A), while none was found in cell lysates (data not shown). This ELISA was modified by substitution of the anti-c-myc detection antibody with either an anti-canine IgE conjugate (Figure 3B), or an anti-human TRAIL antibody (Figure 3C). This confirmed expression of the relevant epitope-tagged recombinant proteins containing canine IgE Fc and/or TRAIL protein domains.

Western blotting of recombinant proteins, purified from supernatant, revealed polyhistidine-tagged proteins, somewhat heavier than the predicted weights of the translated

sequences (rcIgE₁₀₀₋₄₂₇: 42.3kDa; rcIgE₁₀₀₋₄₂₇:TRAIL₁₁₄₋₂₈₂: 61.6kDa; rcTRAIL₁₁₄₋₂₈₂: 26kDa), possibly consistent with glycosylation and/or inefficient cleavage of the signal peptide (Fig. 4).

3.4 Binding of recombinant canine IgE Fc and IgE Fc-TRAIL fusion proteins to FcεR1α

Using the Allerpcept system, it was demonstrated that rcIgE₁₀₀₋₄₂₇ and rcIgE₁₀₀₋₄₂₇:TRAIL₁₁₄₋₂₈₂ were able bind to recombinant FcεR1α, whereas rcTRAIL₁₁₄₋₂₈₂ was unable to do so (Fig. 5). By flow cytometric analysis, C2 cells (FcεR1α positive) bound monoclonal canine IgE as well as rcIgE₁₀₀₋₄₂₇ (Figure 6A), whereas MCLA cells (FcεR1α negative) failed to demonstrate any binding to monoclonal IgE or rcIgE₁₀₀₋₄₂₇ (Figure 6B). Competitive inhibition of binding of rcIgE₁₀₀₋₄₂₇ to C2 cells (detected with anti-myc antibody) was demonstrated by co-incubation with monoclonal IgE, with MFI values falling from 64.4 in the absence of the monoclonal antibody to 27.5 in the presence of the monoclonal antibody. In addition to binding to rcIgE₁₀₀₋₄₂₇, C2 cells were able to bind to the rcIgE₁₀₀₋₄₂₇:TRAIL₁₁₄₋₂₈₂ fusion protein but not rcTRAIL₁₁₄₋₂₈₂ (Figure 6C).

4. Discussion

The features of the canine patients and cMCT recruited for this study are similar to previous reports (Sfiligoi et al., 2005; Mullins et al., 2006; Newman et al., 2007; Thamm and Vail, 2007). Although a retrospective study of archived tumor tissue would have produced a much larger cohort, we have found RNA extracted from formalin-fixed material to be of relatively poor quality, compared to tissue specifically preserved for molecular analysis (Stell, 2008). FcεR1α mRNA expression was consistent throughout the biopsy samples assessed in the study, which contained both well-differentiated and poorly-differentiated tumors. Although mRNA expression does not necessarily indicate expression of functional protein, consistent FcεR1α mRNA expression suggests that development of IgE-based therapies might be worthwhile. The logical next step would be to

analyse a sample cohort composed primarily of high grade, poorly-differentiated or metastatic MCTs where IgE-based immunotherapeutics could be particularly valuable, given that the majority of high grade and disseminated tumors are refractory to current anti-cancer treatments.

Established from a cross-breed dog with tumor recurrence, C2 mastocytoma cells are highly differentiated, expressing surface FcεRI capable of binding canine IgE (Lazarus et al., 1986, Brazis et al., 2002; Hunter et al., 2009). The MCLA mastocytoma cell line was established from a Labrador retriever with a cytologically diagnosed, metastatic MCT. Cells demonstrated typical morphology, although they failed to express FcεRIα. Expression of FcεRIα might have been absent at the time of biopsy, consistent with a neoplasm derived from an early mast cell precursor cell (Thompson et al., 1990; Rottem et al., 1992), de-differentiation and loss of FcεRIα expression in a tumor arising from a later precursor or a mature mast cell, or might have been downregulated during establishment of the cell line, in the absence of IgE which stimulates expression of the receptor (Furuichi et al., 1985; Hsu and MacGlashan, 1996; Yamaguchi et al., 1997; Kubo et al., 2001). There were no detectable KIT exon 11 mutations in this cell line, which also lacked KIT mRNA expression (data not shown). This might also be evidence of an early mast cell precursor lineage, although canine mast cell lines lacking KIT expression and KIT mutation have been reported which are also independent of stem cell factor for their propagation (Ohmori et al., 2008). However, it is difficult to reconcile the lack of FcεRI and KIT mRNA expression with the characteristic morphological appearance of this cell line and the expression of chymase and tryptase mRNA (data not shown), which would suggest more advanced differentiation.

Native IgE (with dimerized heavy chains) binds to the FcεRIα chain on the cell surface with a prolonged half-life, compared to other types of Fc receptor, whereas IgE heavy chain monomers are rapidly internalized (Menon et al., 1986). In the current study, constructs were designed to allow dimerisation of the heavy chain fragments of rclgE₁₀₀₋₄₂₇ as this might facilitate long-term cell surface binding for detection and in the case of rclgE₁₀₀₋₄₂₇:TRAIL₁₁₄₋₂₈₂ would also increase the local

310 concentration of TRAIL death domain moieties on the cell surface for signalling through relevant
311 TRAIL receptors. However, formation of inappropriate intra-molecular disulphide bonds or
312 intermolecular disulphide bonds between rclgE₁₀₀₋₄₂₇ proteins, might have resulted in an altered
313 tertiary and/or quaternary structure. There are two unpaired cysteine residues within the fusion
314 protein (IgE_{100CYS} and TRAIL_{230CYS}) that could potentially lead to misfolding. Cysteine₂₃₀ in human
315 TRAIL is important for trimerisation, with mutation of this amino acid abrogating its apoptosis-
316 inducing effect (Seol and Billiar, 2000; Trabzuni et al., 2000; Kelley et al., 2001). An IgE construct
317 lacking the N-terminal cysteine was generated (pSecTagA/IgE₉₉₋₄₂₇) although this failed to express
318 recombinant protein when transfected into CHO cells (data not shown).

319 An ELISA exploiting the polyhistidine epitope tag for capture and c-myc tag for detection
320 consistently demonstrated recombinant proteins secreted into the supernatant of transfected CHO
321 cells, although the relative immunoreactivity varied between experiments, likely reflecting
322 differences in transfection efficiencies in different wells. Substituting the c-myc antibody with anti-
323 canine IgE or anti-TRAIL antibodies revealed the presence of IgE and/or TRAIL epitopes in the
324 recombinant proteins, suggesting adequate levels of protein folding. Furthermore, a modified
325 Allercept system demonstrated binding of epitope-tagged rclgE₁₀₀₋₄₂₇ to a recombinant FcεRIα
326 conjugate, usually employed to detect serum IgE in dogs suffering from allergic disease. Similar
327 binding of the rclgE₁₀₀₋₄₂₇:TRAIL₁₁₄₋₂₈₂ fusion protein was also demonstrated using this assay,
328 confirming that addition of the TRAIL component did not interfere with the ability of the IgE Fc to
329 bind to its cognate receptor.

330 Binding of recombinant IgE Fc constructs to canine FcεRI was further investigated by flow
331 cytometry, taking advantage of the availability of the novel FcεRI negative MCLA mast cell line as a
332 control for the FcεRI positive C2 mastocytoma line. Specific binding of recombinant IgE₁₀₀₋₄₂₇ to C2
333 cells was demonstrated, which could be inhibited by monoclonal canine IgE. These results provide
334 strong supportive evidence for specific binding of rclgE₁₀₀₋₄₂₇ to canine FcεRI. The C2 cellular binding

experiments were conducted at 4 °C and it might be the case that at 37 °C rclgE₁₀₀₋₄₂₇ proteins could possibly be endocytosed and internalized. This could have been detected through incubating cells for varying periods at 4 °C or 37 °C and comparing the binding of recombinant proteins on the cell surface by anti-c-myc/anti-clgE staining. Alternatives could have included assessing binding at 37 °C in the presence or absence of a metabolic inhibitor (e.g. sodium azide) to prevent internalisation, or the use of confocal microscopy.

The authors have previously demonstrated the preferential efficacy of rhTRAIL₁₁₄₋₂₈₁ in inducing apoptosis in C2 cells compared to non-neoplastic MDCK cells (Elders et al., 2009). Herein, rcTRAIL₁₁₄₋₂₈₂ was selected as a prototype cytotoxic agent that was added to the IgE delivery system, as this would potentially combine selective targeting and selective killing of malignant cells of this lineage. This particular TRAIL fragment is similar to the one shown to induce apoptosis in several canine neoplastic cell lines (Rong et al., 2008), although the rcTRAIL protein used by Rong et al., (2008) is slightly bigger than the current protein (by 3 amino acids at the N-terminus) and the current construct expressed C-terminal c-myc and polyhistidine tags.

Although receptor-binding studies demonstrated that rclgE₁₀₀₋₄₂₇:TRAIL₁₁₄₋₂₈₂ fusion protein was able to bind to C2 cells, subsequent cell viability assays failed to demonstrate any increased levels of apoptosis when C2 cells were cultured in the presence of the recombinant fusion protein (data not shown). Reasons for the lack of biological activity of the fusion protein include inadequate concentration, interference from other components in the cell supernatant or mis-folding of the recombinant protein. Furthermore, whereas IgE monomers form dimers, TRAIL functions exclusively as a trimer (Trabzuni et al., 2000). Thus, the fusion protein as designed would not have an intrinsic apoptosis-inducing effect. Further experiments combining IgE Fc-TRAIL fusion protein with TRAIL monomers, attempting to enhance C-terminal trimerisation also failed to demonstrate any biological effect (data not shown).

5. Conclusions

The specific binding of recombinant canine IgE Fc and a TRAIL fusion protein to canine mast cells expressing FcεRI is proof-of-principle that an IgE-targeted approach is a feasible strategy for mast cell-directed immunotherapeutics in dogs. TRAIL targeted to mast cells via IgE offered the possibility of induction of apoptosis specifically in neoplastic mast cells; however, this proved inefficacious *in vitro*. Future work aims to generate alternative IgE-Fc fusion proteins that can be utilized for antibody-directed enzyme prodrug therapy (ADEPT). As other cells seem capable of performing the majority of the beneficial functions of mast cells *in vivo*, the therapeutic index for such IgE-based immunotherapeutics is likely to be high for cMCT-bearing patients, provided that adequate endo-and ecto-parasite control is implemented during therapy, with likely repopulation of the tissues with mast cells from bone-marrow derived precursors after treatment is complete.

Conflict of interest statement

None of the authors of this paper has a financial or personal relationship with other people or organisations that could inappropriately influence or bias the content of the paper.

Acknowledgements

The authors would like to thank The Biotechnology and Biological Sciences Research Council and Merial Animal Health for their generous funding of this PhD studentship. We would also like to thank HESKA AG, Fribourg, Switzerland, especially Dr D. Wassom and Dr M. Derer, for generating the Allercept-based ELISA data.

Role of the funding source

382 The study design, collection, analysis and interpretation of data, the writing of this manuscript and
383 the decision to publish were made independently of the above sponsors.

References

- Belostotsky, R., Lorberboum-Galski, H., 2001, Apoptosis-inducing human-origin Fcepsilon-Bak chimeric proteins for targeted elimination of mast cells and basophils: a new approach for allergy treatment. *J Immunol* 167, 4719-4728.
- Blackwood, L., Murphy, S., Buracco, P., De Vos, J.P., De Fornel-Thibaud, P., Hirschberger, J., Kessler, M., Pastor, J., Ponce, F., Savary-Bataille, K., Argyle, D.J., European consensus document on mast cell tumours in dogs and cats. *Vet Comp Oncol* 10, e1-e29.
- Bostock, D.E., 1973, The prognosis following surgical removal of mastocytomas in dogs. *J Small Anim Pract* 14, 27-41.
- Bostock, D.E., 1986, Neoplasms of the skin and subcutaneous tissues in dogs and cats. *Br Vet J* 142, 1-19.
- Brazis, P., Torres, R., Queralt, M., de Mora, F., Ferrer, L., Puigdemont, A., 2002, Evaluation of cell-surface IgE receptors on the canine mastocytoma cell line C2 maintained in continuous culture. *Am J Vet Res* 63, 763-766.
- Brodey, R.S., 1970, Canine and feline neoplasia. *Adv Vet Sci Comp Med* 14, 309-354.
- Ding, J., Ichikawa, Y., Ishikawa, T., Shimada, H., 2004, Effect of formalin on extraction of mRNA from a formalin-fixed sample: a basic investigation. *Scand J Clin Lab Invest* 64, 229-235.
- Dobson, J.M., Samuel, S., Milstein, H., Rogers, K., Wood, J.L., 2002, Canine neoplasia in the UK: estimates of incidence rates from a population of insured dogs. *J Small Anim Pract* 43, 240-246.
- Echtenacher, B., Männel, D.N., Hültner, L., 1996. Critical protective role of mast cells in a model of acute septic peritonitis. *Nature* 381, 75-77.
- Elders, R.C., Baines, S.J., Catchpole, B., 2009, Susceptibility of the C2 canine mastocytoma cell line to the effects of tumor necrosis factor-related apoptosis-inducing ligand (TRAIL). *Vet Immunol Immunopathol* 130, 11-16.
- Finnie, J.W., Bostock, D.E., 1979, Skin neoplasia in dogs. *Aust Vet J* 55, 602-604.

410 Fishman, A., Lorberboum-Galski, H., 1997, Targeted elimination of cells expressing the high-affinity
 411 receptor for IgE (Fc epsilon RI) by a Pseudomonas exotoxin-based chimeric protein. *Eur J*
 412 *Immunol* 27, 486-494.

413 Fishman, A., Prus, D., Belostotsky, R., Lorberboum-Galski, H., 2000, Targeted Fc2'-3-PE40 chimeric
 414 protein abolishes passive cutaneous anaphylaxis in mice. *Clin Exp Immunol* 119, 398-403.

415 Foster, A.P., Littlewood, J.D., Webb, P., Wood, J.L., Rogers, K., Shaw, S.E., 2003, Comparison of
 416 intradermal and serum testing for allergen-specific IgE using a Fcepsilon RIalpha-based assay
 417 in atopic dogs in the UK. *Vet Immunol Immunopathol* 93, 51-60.

418 Furuichi, K., Rivera, J., Triche, T., Isersky, C., 1985, The fate of IgE bound to rat basophilic leukemia
 419 cells. IV. Functional association between the receptors for IgE. *J Immunol* 134, 1766-1773.

420 Garman, S.C., Sechi, S., Kinet, J.P., Jardetzky, T.S., 2001, The analysis of the human high affinity IgE
 421 receptor Fc epsilon Ri alpha from multiple crystal forms. *J Mol Biol* 311, 1049-1062.

422 Garman, S.C., Wurzburg, B.A., Tarchevskaya, S.S., Kinet, J.P., Jardetzky, T.S., 2000, Structure of the Fc
 423 fragment of human IgE bound to its high-affinity receptor Fc epsilonRI alpha. *Nature* 406, 259-
 424 266.

425 Gerritsen, R.J., Teske, E., Kraus, J.S., Rutteman, G.R., 1998, Multi-agent chemotherapy for mast cell
 426 tumours in the dog. *Vet Q* 20, 28-31.

427 Goitsuka, R., Hayashi, N., Nagase, M., Sasaki, N., Ra, C., Tsujimoto, H., Hasegawa, A., 1999, Molecular
 428 cloning of cDNAs encoding dog high-affinity IgE receptor alpha, beta, and gamma chains.
 429 *Immunogenetics* 49, 580-582.

430 Grimaldeston, M.A., Chen, C.C., Piliponsky, A.M., Tsai, M., Tam, S.Y., Galli, S.J., 2005, Mast cell-
 431 deficient W-sash c-kit mutant Kit W-sh/W-sh mice as a model for investigating mast cell
 432 biology in vivo. *Am J Pathol* 167, 835-848.

433 Hsu, C., MacGlashan, D., Jr., 1996, IgE antibody up-regulates high affinity IgE binding on murine bone
 434 marrow-derived mast cells. *Immunol Lett* 52, 129-134.

435 Hunter, M.J., Iodice, M.W., Pathak, A.K., Street, M.A., Helm, B.A., 2009, Characterization of an early
 436 signaling defect following Fc epsilonRI activation in the canine mastocytoma cell line, C2. *Mol*
 437 *Immunol* 46, 1260-1265.

438 Hunter, M.J., Vratimos, A.P., Housden, J.E., Helm, B.A., 2008, Generation of canine-human Fc IgE
 439 chimeric antibodies for the determination of the canine IgE domain of interaction with Fc
 440 epsilon RI alpha. *Mol Immunol* 45, 2262-2268.

441 Joseph, M., Gounni, A.S., Kusnierz, J.P., Vorng, H., Sarfati, M., Kinet, J.P., Tonnel, A.B., Capron, A.,
 442 Capron, M., 1997, Expression and functions of the high-affinity IgE receptor on human
 443 platelets and megakaryocyte precursors. *Eur J Immunol* 27, 2212-2218.

444 Kelley, S.K., Harris, L.A., Xie, D., Deforge, L., Totpal, K., Bussiere, J., Fox, J.A., 2001, Preclinical studies
 445 to predict the disposition of Apo2L/tumor necrosis factor-related apoptosis-inducing ligand in
 446 humans: characterization of in vivo efficacy, pharmacokinetics, and safety. *J Pharmacol Exp*
 447 *Ther* 299, 31-38.

448 Keown, M.B., Ghirlando, R., Mackay, G.A., Sutton, B.J., Gould, H.J., 1997, Basis of the 1:1
 449 stoichiometry of the high affinity receptor Fc epsilon RI-IgE complex. *Eur Biophys J* 25, 471-
 450 476.

451 Kinet, J.P., 1999, The high-affinity IgE receptor (Fc epsilon RI): from physiology to pathology. *Annu*
 452 *Rev Immunol* 17, 931-972.

453 Kita, H., Kaneko, M., Bartemes, K.R., Weiler, D.A., Schimming, A.W., Reed, C.E., Gleich, G.J., 1999,
 454 Does IgE bind to and activate eosinophils from patients with allergy? *J Immunol* 162, 6901-
 455 6911.

456 Kitamura, Y., Go, S., Hantanka, K., 1978, Decrease of mast cells in W/W^v mice and their increase by
 457 bone marrow transplantation. *Blood* 52, 447-452.

458 Kubo, S., Matsuoka, K., Taya, C., Kitamura, F., Takai, T., Yonekawa, H., Karasuyama, H., 2001, Drastic
 459 up-regulation of FcepsilonRI on mast cells is induced by IgE binding through stabilization and
 460 accumulation of FcepsilonRI on the cell surface. *J Immunol* 167, 3427-3434.

461 Lazarus, S.C., DeVinney, R., McCabe, L.J., Finkbeiner, W.E., Elias, D.J., Gold, W.M., 1986, Isolated
 462 canine mastocytoma cells: propagation and characterization of two cell lines. *Am J Physiol*
 463 251, C935-944.
 464 LeBlanc, H.N., Ashkenazi, A., 2003, Apo2L/TRAIL and its death and decoy receptors. *Cell Death Differ*
 465 10, 66-75.
 466 Ledin, A., Bergvall, K., Hillbertz, N.S., Hansson, H., Andersson, G., Hedhammar, A., Hellman, L., 2006,
 467 Generation of therapeutic antibody responses against IgE in dogs, an animal species with
 468 exceptionally high plasma IgE levels. *Vaccine* 24, 66-74.
 469 Maenaka, K., van der Merwe, P.A., Stuart, D.I., Jones, E.Y., Sonderrmann, P., 2001, The human low
 470 affinity Fcγ receptors IIa, IIb, and III bind IgG with fast kinetics and distinct
 471 thermodynamic properties. *J Biol Chem* 276, 44898-44904.
 472 Mariani, S.M., Krammer, P.H., 1998, Differential regulation of TRAIL and CD95 ligand in transformed
 473 cells of the T and B lymphocyte lineage. *Eur J Immunol* 28, 973-982.
 474 McDonnell, J.M., Calvert, R., Beavil, R.L., Beavil, A.J., Henry, A.J., Sutton, B.J., Gould, H.J., Cowburn,
 475 D., 2001, The structure of the IgE Cε2 domain and its role in stabilizing the complex with
 476 its high-affinity receptor FcεRIα. *Nat Struct Biol* 8, 437-441.
 477 Menon, A.K., Holowka, D., Webb, W.W., Baird, B., 1986, Cross-linking of receptor-bound IgE to
 478 aggregates larger than dimers leads to rapid immobilization. *J Cell Biol* 102, 541-550.
 479 Mullins, M.N., Dernell, W.S., Withrow, S.J., Ehrhart, E.J., Thamm, D.H., Lana, S.E., 2006, Evaluation of
 480 prognostic factors associated with outcome in dogs with multiple cutaneous mast cell tumors
 481 treated with surgery with and without adjuvant treatment: 54 cases (1998-2004). *J Am Vet*
 482 *Med Assoc* 228, 91-95.
 483 Newman, S.J., Mrkonjich, L., Walker, K.K., Rohrbach, B.W., 2007, Canine subcutaneous mast cell
 484 tumour: diagnosis and prognosis. *J Comp Pathol* 136, 231-239.

485 Ohmori, K., Kawarai, S., Yasuda, N., Tanaka, A., Matsuda, H., Nishimura, R., Sasaki, N., Tsujimoto, H.,
 486 Masuda, K., 2008, Identification of c-kit mutations-independent neoplastic cell proliferation of
 487 canine mast cells. *Vet Immunol Immunopathol* 126, 43-53.
 488 O'Keefe, D.A., 1990, Canine mast cell tumors. *Vet Clin North Am Small Anim Pract* 20, 1105-1115.
 489 Pan, G., Ni, J., Yu, G., Wei, Y.F., Dixit, V.M., 1998, TRUNDD, a new member of the TRAIL receptor
 490 family that antagonizes TRAIL signalling. *FEBS Lett* 424, 41-45.
 491 Patnaik, A.K., Ehler, W.J., MacEwen, E.G., 1984, Canine cutaneous mast cell tumor: morphologic
 492 grading and survival time in 83 dogs. *Vet Pathol* 21, 469-474.
 493 Pitti, R.M., Marsters, S.A., Ruppert, S., Donahue, C.J., Moore, A., Ashkenazi, A., 1996, Induction of
 494 apoptosis by Apo-2 ligand, a new member of the tumor necrosis factor cytokine family. *J Biol*
 495 *Chem* 271, 12687-12690.
 496 Rao, K.N., Brown, M.A., 2008, Mast cells: multifaceted immune cells with diverse roles in health and
 497 disease. *Ann N Y Acad Sci* 1143, 83-104.
 498 Ravetch, J.V., Kinet, J.P., 1991, Fc receptors. *Annu Rev Immunol* 9, 457-492.
 499 Rong, S., Cai, J.H., Andrews, J., 2008, Cloning and apoptosis-inducing activities of canine and feline
 500 TRAIL. *Mol Cancer Ther* 7, 2181-2191.
 501 Rottem, M., Barbieri, S., Kinet, J.P., Metcalfe, D.D., 1992, Kinetics of the appearance of Fc epsilon RI-
 502 bearing cells in interleukin-3-dependent mouse bone marrow cultures: correlation with
 503 histamine content and mast cell maturation. *Blood* 79, 972-980.
 504 Seminario, M.C., Saini, S.S., MacGlashan, D.W., Jr., Bochner, B.S., 1999, Intracellular expression and
 505 release of Fc epsilon RI alpha by human eosinophils. *J Immunol* 162, 6893-6900.
 506 Seol, D.W., Billiar, T.R., 2000, Cysteine 230 modulates tumor necrosis factor-related apoptosis-
 507 inducing ligand activity. *Cancer Res* 60, 3152-3154.
 508 Sfiligoi, G., Rassnick, K.M., Scarlett, J.M., Northrup, N.C., Gieger, T.L., 2005, Outcome of dogs with
 509 mast cell tumors in the inguinal or perineal region versus other cutaneous locations: 124 cases
 510 (1990-2001). *J Am Vet Med Assoc* 226, 1368-1374.

511 Sheridan, J.P., Marsters, S.A., Pitti, R.M., Gurney, A., Skubatch, M., Baldwin, D., Ramakrishnan, L.,
 512 Gray, C.L., Baker, K., Wood, W.I., Goddard, A.D., Godowski, P., Ashkenazi, A., 1997, Control of
 513 TRAIL-induced apoptosis by a family of signaling and decoy receptors. *Science* 277, 818-821.
 514 Simonet, W.S., Lacey, D.L., Dunstan, C.R., Kelley, M., Chang, M.S., Luthy, R., Nguyen, H.Q., Wooden,
 515 S., Bennett, L., Boone, T., Shimamoto, G., DeRose, M., Elliott, R., Colombero, A., Tan, H.L.,
 516 Trail, G., Sullivan, J., Davy, E., Bucay, N., Renshaw-Gegg, L., Hughes, T.M., Hill, D., Pattison, W.,
 517 Campbell, P., Sander, S., Van, G., Tarpley, J., Derby, P., Lee, R., Boyle, W.J., 1997,
 518 Osteoprotegerin: a novel secreted protein involved in the regulation of bone density. *Cell* 89,
 519 309-319.
 520 Stell, A.J., 2008, Development of a DNA vaccine for canine malignant melanoma. Ph.D. thesis, Royal
 521 Veterinary College, University of London.
 522 Thamm, D.H., Vail, D.M., 2007, Mast cell tumors. In: Withrow, S.J., Vail, D.M. (Eds), Withrow and
 523 MacEwen's Small Animal Clinical Oncology, 3rd edn, Saunders Elsevier, St Louis, pp. 402–424.
 524 Thompson, H.L., Metcalfe, D.D., Kinet, J.P., 1990, Early expression of high-affinity receptor for
 525 immunoglobulin E (Fc epsilon RI) during differentiation of mouse mast cells and human
 526 basophils. *J Clin Invest* 85, 1227-1233.
 527 Trabzuni, D., Famulski, K.S., Ahmad, M., 2000, Functional analysis of tumour necrosis factor-alpha-
 528 related apoptosis-inducing ligand (TRAIL): cysteine-230 plays a critical role in the
 529 homotrimerization and biological activity of this novel tumoricidal cytokine. *Biochem J* 350 Pt
 530 2, 505-510.
 531 Wan, T., Beavil, R.L., Fabiane, S.M., Beavil, A.J., Sohi, M.K., Keown, M., Young, R.J., Henry, A.J.,
 532 Owens, R.J., Gould, H.J., Sutton, B.J., 2002, The crystal structure of IgE Fc reveals an
 533 asymmetrically bent conformation. *Nat Immunol* 3, 681-686.
 534 Welle, M.M., Bley, C.R., Howard, J., Rufenacht, S., 2008, Canine mast cell tumours: a review of the
 535 pathogenesis, clinical features, pathology and treatment. *Vet Dermatol* 19, 321-339.

Wiley, S.R., Schooley, K., Smolak, P.J., Din, W.S., Huang, C.P., Nicholl, J.K., Sutherland, G.R., Smith, T.D., Rauch, C., Smith, C.A., et al., 1995, Identification and characterization of a new member of the TNF family that induces apoptosis. *Immunity* 3, 673-682.

Wurzberg, B.A., Jardetzky, T.S., 2002, Structural insights into the interactions between human IgE and its high affinity receptor FcεRI. *Mol Immunol* 38, 1063-1072.

Yamaguchi, M., Lantz, C.S., Oettgen, H.C., Katona, I.M., Fleming, T., Miyajima, I., Kinet, J.P., Galli, S.J., 1997, IgE enhances mouse mast cell Fc(ε)RI expression in vitro and in vivo: evidence for a novel amplification mechanism in IgE-dependent reactions. *J Exp Med* 185, 663-672.

Fig. 1. Screening of canine MCT biopsies and mastocytoma cell lines for expression of FcεRIα mRNA. PCR was used to assess FcεRIα mRNA expression (273 bp amplicon) in (A) cMCT biopsies (B-Z) and (B) mastocytoma cell lines C2 and MCLA (MC). PCR products were separated by agarose gel electrophoresis and visualized under UV illumination. La: 100 bp molecular weight ladder. H2O: water negative control.

Fig. 2. PCR screening for canine IgE mRNA in control and transfected CHO cells. PCR was used to amplify the C_Hε2-4 region of canine IgE from cDNA prepared from CHO cells transfected with the pSecTagA vector containing this region of the genomic sequence. PCR products were separated by agarose gel electrophoresis and visualized under UV illumination. L = 100 bp molecular weight ladder. H = water negative control; G = genomic DNA positive control; UC = untransfected CHO cDNA; UN = untransfected CHO NRT; TN = transfected CHO NRT; TC = transfected CHO cDNA, which produced amplicons with neither (largest fragment), one or both (smallest fragment) introns excised.

Fig. 3. Detection of recombinant proteins in transfected CHO supernatant. Untransfected (Mock) and pSecTagA/rcIgE₁₀₀₋₄₂₇ (IgE), pSecTagA/rcIgE₁₀₀₋₄₂₇ : rcTRAIL₁₁₄₋₂₈₂ (IgE/TRAIL), pSecTagA/rcTRAIL₁₁₄₋₂₈₂ (TRAIL) transfected CHO supernatants were harvested 90 h post transfection. Supernatants were applied to wells of Ni-NTA HisSorb plates and recombinant epitope-tagged proteins detected using by ELISA using anti-c-myc (A) anti-canine IgE (B) or anti-TRAIL (C) antibody conjugates. A polyhistidine-tagged recombinant human TRAIL protein was used as a positive control in (C). Results are show as the mean of triplicate wells ± SEM, corrected by subtracting the mean value from wells

where medium only were applied. Experiments were repeated, with demonstration of recombinant protein expressed from each construct, although the relative levels of recombinant protein immunoreactivity varied between transfections. O.D. = optical density.

Fig. 4. Detection of recombinant proteins in transfected CHO supernatants by Western blotting. Recombinant protein, purified from CHO supernatant using MagneHis beads, were separated under reducing conditions by SDS PAGE and immunoblotted. Recombinant protein was detected using anti-polyhistidine:HRP conjugate and enhanced chemiluminescence with 5 and 30 min exposure of autoradiography film. **The arrow indicates histidine-tagged recombinant canine TRAIL after 5 minutes exposure, which is more apparent after 30 min exposure.** Lad = Histidine-tagged molecular weight ladder; U = unpurified, untransfected CHO supernatant; I = purified IgE₁₀₀₋₄₂₇; F = purified IgE₁₀₀₋₄₂₇:TRAIL₁₁₄₋₂₈₂; T = purified TRAIL₁₁₄₋₂₈₂.

Fig. 5. Binding of rclgE₁₀₀₋₄₂₇ and rclgE₁₀₀₋₄₂₇:TRAIL₁₁₄₋₂₈₂ in the Allercept assay. Recombinant proteins within transfected CHO supernatants were captured using HisSorb strips (A) or Maxisorb ELISA plates coated with anti-canine IgE antibody (B) and their ability to bind to the FcεRIα chain was assessed using the Allercept system. IgE = pSecTagA/IgE₁₀₀₋₄₂₇-transfected CHO supernatant; IgE:TRAIL = pSecTagA/IgE₁₀₀₋₄₂₇ : TRAIL₁₁₄₋₂₈₂-transfected CHO supernatant; TRAIL = pSecTagA/TRAIL₁₁₄₋₂₈₂-transfected CHO supernatant.

Fig. 6. Binding of recombinant proteins to C2 and MCLA mastocytoma cells. C2 (A and C) or MCLA cells (B) were incubated with mock-transfected CHO supernatant (medium) or supernatant containing rclgE₁₀₀₋₄₂₇ (rclgE), rclgE₁₀₀₋₄₂₇:TRAIL₁₁₄₋₂₈₂ (rclgE/TRAIL) or rcTRAIL₁₁₄₋₂₈₂ (rcTRAIL), followed by labelling with polyclonal anti-canine IgE:FITC (panel B) or polyclonal anti-c-myc:FITC (panel C) or in the absence of a secondary antibody (panel A) and analysed by flow cytometry. The graphs show histogram overlays of cells incubated with the various recombinant proteins. The tables show mean fluorescence intensity (MFI) data for each incubation condition.

Figure 1A

Figure 1B

602 **Figure 2**

603

604

605

Figure 3A

Figure 3B

614 **Figure 3C**

615
616
617

Figure 4

Figure 5

627 **Figure 6A**

628 **C2 cells**

629

630

631

632 **Figure 6B**

633 **MCLA cells**

		Medium	Monoclonal cIgE	rclgE
A	None	20.04	29.74	19.54
B	cIgE:FITC	19.26	36.63	20.42
C	c-myc:FITC	18.23	27.55	20.12

634

635

636

Figure 6C

C2 cells

		Medium	rclgE	rclgE/TRAIL	rcTRAIL
A	None	34.14	29.85	29.57	29.69
B	clgE:FITC	85.59	670.5	474.23	55.02
C	c-myc:FITC	29.22	227.84	145.75	37.22

Table 1. Primers used in study

Target gene	Primer sequence (5'-3')	Size (bp)	Genbank #
FCERIA	F: AGTGGGGAGTACAGGTGTCG R: GCCTGAGCAGGAATAGTTGC	273	NM_00110766.1
IGHE	F: <u>AAGCTT</u> ATGTGCCTAAACTTCATTCCG R: TTTACCGGGGGTTTTGGACAC	988(cDNA) 1152(gDNA)	XM_548007.1
TRAIL	F: CGAGGTTCTCAGAGAGTAGCT R: <u>CTCGAG</u> GCAGCGTATTTTGCCGATTA	522	NM_001130836.1

F = forward/sense primer; R = reverse/antisense primer. Restriction enzyme sites (*Hind*III: AAGCTT and *Xho*I: CTCGAG) are shown underlined.

645 **Table 2.** Details of MCT-bearing patients.

Identifier	Age	Breed	Sex	Tumor site	Differentiation review	Location review
B	10y11m	Crossbred (Lurcher)	FN	Tail	Intermediate	Skin/SQ
C	7y6m	German shepherd dog	FN	Visceral	Poor	Skin
D	9y5m	Crossbred	FN	Thigh	Intermediate	SQ
E	8y9m	Staffordshire bull terrier	ME	Scrotum	Intermediate	Skin
F	9y1m	Rhodesian ridgeback	ME	Stifle	Well	Skin
G	8y11m	Labrador retriever	FN	Interdigital	Well	Skin
H	9y1m	Staffordshire bull terrier	MN	Muzzle	(Cytology only)	
I	11y	Crossbred (collie-type)	FN	Ventral abdomen	Poor	Skin
J	8y11m	Staffordshire bull terrier	MN	Thigh	Intermediate	SQ
K	6y4m	Labrador retriever	MN	Thigh	Intermediate	Skin
L	11y3m	Crossbred	FN	Ventral abdomen	(Cytology only)	
M	7y	Golden retriever	FE	Flank	Intermediate	Skin
N	8y1m	Labrador retriever	FN	Thigh	Intermediate	Skin/SQ
O	11y10m	Shih Tzu	MN	Prepuce	Intermediate	SQ
P	8y2m	Labrador retriever	FE	Thigh	Intermediate	Skin
Q	11y1m	Labrador retriever	MN	Scrotum	(Cytology only)	
R	9y1m	Staffordshire bull terrier	ME	Thigh	Intermediate	SQ
S	12y7m	Labrador retriever	ME	Ventral abdomen	(Cytology only)	
T	11y	Labrador retriever	MN	Prepuce	Intermediate	SQ
U	9y10m	Staffordshire bull terrier	FN	Thorax	Intermediate	SQ
V	9y5m	Labrador retriever	MN	Pinna	Intermediate	Skin
W	10y	Labrador retriever	FN	Flank	Intermediate	SQ
X	6y	Jack Russell terrier	MN	Thigh	Intermediate	Skin
Y	5y	Boxer	FN	Antebrachium	Intermediate	SQ
Z	8y4m	Staffordshire bull terrier	FN	Calf	Poor	Connective Tissue

646

647 FE: female entire; FN: female neutered; LTF: histopathology slides lost to follow-up; ME:

648 male entire; MN: male neutered; SQ: subcutaneous.

649

650

651