THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Recent advances in the regulation of plant immunity by S-nitrosylation

Citation for published version:

Lubega, J, Umbreen, S & Loake, GJ 2020, 'Recent advances in the regulation of plant immunity by S-nitrosylation', *Journal of Experimental Botany*. <https://doi.org/10.1093/jxb/eraa454>

Digital Object Identifier (DOI):

[10.1093/jxb/eraa454](https://doi.org/10.1093/jxb/eraa454)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Experimental Botany

Publisher Rights Statement:

This is a pre-copyedited, author-produced version of an article accepted for publication in Journal of Experimental Botany following peer review. The version of record [insert complete citation information here] is available online at: Jibril Lubega, Saima Umbreen, Gary J Loake, Recent Advances in the Regulation of Plant Immunity by S-Nitrosylation, Journal of Experimental Botany, , eraa454, <https://doi.org/10.1093/jxb/eraa454>

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

RECENT ADVANCES IN THE REGULATION OF PLANT IMMUNITY BY S-NITROSYLATION

Jibril Lubega¹, Saima Umbreen¹ and Gary J Loake^{1,2,3}

¹Institute of Molecular Plant Sciences, School of Biological Sciences, University of Edinburgh, Edinburgh, EH9 3BF, UK

²Centre for Synthetic and Systems Biology, School of Biological Sciences, University of Edinburgh, The King's Buildings, Edinburgh EH9 3BF, UK

³Corresponding author: Email: gloake@ed.ac.uk, Tel: 0131 650 5332, Fax: 0131 650 5392

Key words: S-nitrosylation, nitric oxide, plant immunity, SUMOylation, Zinc finger proteins, effector, autophagy.

Word Count: ≈ 3,887

Highlights

Herein we provide an update on recent developments of S-nitrosylation in plant immune function which includes: both how this redox-based post-translational modification controls key aspects of immunity and also how S-nitrosylation can disable aspects of pathogen virulence.

Abstract

S-nitrosylation, the addition of a nitric oxide (NO) moiety to a reactive protein cysteine (Cys) thiol, to form a protein S-nitrosothiol (SNO), is emerging as a key regulatory post-translational modification (PTM) to control the plant immune response. NO also S-nitrosylates the antioxidant tripeptide, glutathione (GSH), to form S-nitrosoglutathione (GSNO), both a storage reservoir of NO bioactivity and a natural NO donor. GSNO and by extension, S-nitrosylation, is controlled by GSNO reductase1 (GSNOR1). The emerging data suggests that GSNOR1 itself is a target of NO-mediated S-nitrosylation, which subsequently controls its selective autophagy, regulating cellular protein SNO levels. Recent findings also suggest that S-nitrosylation may be deployed by pathogen-challenged host cells to counteract the effect of delivered microbial effector proteins, that promote pathogenesis and by the pathogens themselves to augment virulence. Significantly, it also appears that S-nitrosylation may regulate plant immune functions by controlling SUMOylation, a peptide-based PTM. In this context, global SUMOylation is regulated by S-nitrosylation of SUMO conjugating enzyme, SCE1 at Cys 139. This redox-based PTM has also been shown to control the function of a key zinc finger transcriptional regulator during the establishment of plant immunity and the selective degradation of GSNOR1 by autophagy. Here, we provide an update of these recent advances.

60 Introduction

61 A key feature of the plant immune response is the associated striking changes in cellular redox
62 status resulting from elevated levels of reactive oxygen ~~species~~intermediates (ROS~~Is~~) and
63 reactive nitrogen species (RNS) (Couturier *et al.*, 2013; Skelly and Loake, 2013; Yu *et al.*,
64 2014). RNS comprise predominately nitric oxide (NO), the NO radical (NO $\cdot$) and other
65 derivatives including, peroxynitrite (ONOO $^-$), dinitrogen trioxide (N $_2$ O $_3$), and nitrogen dioxide
66 (NO $_2$) (Corcoran and Cotter, 2013; Grant and Loake, 2000; Couturier *et al.*, 2013). These
67 small, redox-active molecules, function in redox signalling networks helping to orchestrate the
68 plant response to a plethora of potentially stresses (Yu *et al.*, 2012; Couturier *et al.*, 2013;
69 Lisjak *et al.*, 2013; Turkan, 2017).

70

71 NO is thought to be produced majorly from the reduction of nitrates~~s~~/ nitrite by NADPH and
72 pH dependent nitrate reductase (NR) (Rockel *et al.*, 2002; Bolwell and Daudi, 2009; Sanz *et al.*
73 *et al.*, 2015; Astier *et al.*, 2018). However, other potential sources include: mitochondria, through
74 the electron transport chain, peroxisomes and chloroplasts (Kolbert *et al.*, 2019). In
75 *Arabidopsis*, NR is located in the cytosol and encoded by two structural genes *NIA1* and *NIA2*,
76 with *NIA2* exhibiting the highest observed nitrate reductase activity (Wilkinson and Crawford,
77 1991; Frederickson Matika and Loake, 2014). NO bioavailability is tightly regulated by
78 scavenging mechanisms (Chamizo-Ampudia *et al.*, 2017). Thus, NO reacts with freely
79 available intracellular glutathione to form S-nitrosogluthathione (GSNO) that acts as both a
80 storage reservoir and a natural donor of NO. GSNO is metabolized by a cytosolic enzyme S-
81 nitrosogluthathione reductase 1 (GSNOR1) to glutathione disulphide and ammonia (Feechan
82 *et al.*, 2005~~a~~; Yun *et al.*, 2011, 2016). NO can also be scavenged by ROS ~~reactive-oxygen~~
83 ~~species~~ ~~(ROS)~~ such as superoxides (O $_2^{\cdot-}$) to form ONOO $^-$ (Romero-Puertas *et al.*, 2007;
84 Gaupels *et al.*, 2011; Begara-Morales *et al.*, 2014). NO can also be consumed by pPlant
85 hemoglobins (Hb) which ~~These~~ are ubiquitous proteins conserved across biological
86 kingdoms ~~(Gardner, 2012)~~ and have ~~also~~ been shown to scavenge NO by catalyzing the
87 dioxygenation of NO to form nitrates~~s~~ (Gardner, 2012; Chamizo-Ampudia *et al.*, 2017).

88

89 Generated NO functions majorly through S-nitrosylation, a prototypic, redox-based post-
90 translational modification (PTM). This process involves the addition of an NO moiety to a rare,
91 highly reactive cysteine (Cys) thiol to form an S-nitrosothiol (SNO) in a fashion akin to more
92 established PTMs, such as phosphorylation (Spadaro *et al.*, 2010; Friso and van Wijk, 2015;
93 Gupta *et al.*, 2020). Subsequently, S-nitrosylation may regulate protein localization, protein
94 function, protein-protein interactions and protein stability in an allosteric-like fashion following
95 associated conformational changes (Astier *et al.*, 2012b; Vanzo *et al.*, 2014). To date, a
96 number of plant proteins integral to plant immunity have been found to be S-nitrosylated,
97 including: NADPH oxidase (Yun *et al.*, 2011, 2016), NPR1 (Tada *et al.*, 2008; Spoel and
98 Loake, 2011), TGA1 (Lindermayr *et al.*, 2010), AtSABP3 (Wang *et al.*, 2009) and
99 Peroxyredoxin (Romero-Puertas 2007).

100

101 In this review, we highlight recent advances associated with protein S-nitrosylation during plant
102 immunity. We discuss host-driven S-nitrosylation targeting pathogen derived effector proteins
103 delivered to the inside of plant cells, subsequently disabling their function(s) (Ling *et al.*, 2017).
104 Further, we outline the ability of S-nitrosylation to control the transcriptional reprogramming of
105 plant gene expression during the immune response (Cui *et al.*, 2018). We also review the role
106 of S-nitrosylation in promoting autophagy, by destabilizing immune-related proteins leading to
107 lysosome degradation promoting pathogen susceptibility (Zhan *et al.*, 2018). Lastly, we
108 discuss the regulation of SUMOylation by S-nitrosylation (Skelly *et al.*, 2019) during the plant
109 immune response.

110

111 **S-nitrosylation exploited by the host and pathogen during immuno responses**

112 Plant pathogens are invading microorganisms aimed to derive nutrients from the host to
113 support their growth and development (Davidsson *et al.*, 2013; Snelders *et al.*, 2018). To
114 achieve this pathogens secretes effectors which are the proteinaceous molecules released to
115 circumvent the host defence machinery (Kamoun, 2007). They are grouped majorly into two

categories, the apoplastic effectors that act outside the cell and the cytoplasmic effectors that act inside the host cell (Asai and Shirasu, 2015). In fungi and oomycetes, effectors are produced and released by haustoria, which also function as intracellular feeding structures (Wang *et al.*, 2017). In bacterial pathogens, effector proteins are secreted via six secretory systems (Setti *et al.*, 2014). For instance, the type two secretory system (T2SS) also known as sec dependent system release the apoplastic effectors, cell wall degrading enzymes and other hydrolytic enzymes such as cellulases, xylanases, amylases and proteases (Setti *et al.*, 2014; Pfeilmeier *et al.*, 2016). Whereas the type three secretory system (T3SS) is the molecular machinery that delivers cytoplasmic effectors into host cells. This system is encoded by *Hypersensitive Response* and *Pathogenicity (HRP)* genes (Setti *et al.*, 2014). The transferred effectors promote pathogenicity on a susceptible host and a hypersensitive response on a resistant host by interfering with key cellular processes (Buttner and He, 2009).

Although the molecular mechanisms underpinning the release of bacterial plant pathogen effectors into the apoplast or cytosol has been largely elucidated, the identification of their possible host protein targets is still continuing at pace (Park *et al.*, 2012; Hann *et al.*, 2014; Liu *et al.*, 2016; Wang *et al.*, 2017). As previously stated, effector-triggered immunity (ETI) results in a conspicuous nitrosative burst, although the source(s) of NO still remain controversial (Schlicht and Kombrink, 2013). During the onset of ETI in *Arabidopsis* the bacterial effector protein HopAI1 (HRP-dependent outer protein A1) of *Pseudomonas syringae* pv *tomato* DC3000 (*Pst*DC3000) is S-nitrosylated inhibiting the phosphothreonine lyase activity of this effector (Ling *et al.*, 2017). HopAI1 is known to enhances *Pst* strain 0288-9 virulence in tomato plants. In a similar fashion, the transgenic expression of HopAI1 in *Arabidopsis* increases susceptibility to *Pst*DC3000 by compromising ETI (Zhang *et al.*, 2007).

HopAI1 targets and suppresses mitogen activated protein kinases (MAPKs) activated by exposure to pathogen-associated molecular patterns (PAMPS) such as flagellin 22 (Bolwell

and Daudi, 2009). MAPKs are involved in signal transduction mediated by phosphorylation of proteins leading to activation of defence genes that orchestrate the development of a plethora of immune-related functions (Taj *et al.*, 2010). The canonical pathway for MAPK signaling is a phosphorylation cascade supported by associated mitogen activated protein kinase kinase kinase (MAPKKK), MAP kinase kinase (MAPKK) through to MAPK function (Taj *et al.*, 2010). Finally the activated MAPK is transported to the nucleus where it typically phosphorylates one or more target transcription factors, supporting transcriptional reprogramming (Taj *et al.*, 2010). The HopAI1 effector is thought to act by removing a phosphate group from phosphothreonine required for HopAI1 activity (Zhang *et al.*, 2007). S-nitrosylation of HopAI1 occurs at Cys 138, a non-catalytic residue, restoring MAPK signalling and consequently, disarming a key pathogen infection strategy (Ling *et al.*, 2017). In addition, overexpression of HopAI1 Cys138S (Cys is replaced with Serine), an S-nitrosylation insensitive mutant, triggered autoimmunity and failed to promote the anticipated pathogen susceptibility (Ling *et al.*, 2017).

On the other hand, S-nitrosylation can be utilized by the pathogen to enhance virulence. For instance, cell division control protein 48 (CDC48) also known as p97 or ~~Valslin~~ containing protein (VCP) is an evolutionally conserved ATPase of AAA family present in yeast, plants and animals (Baek *et al.*, 2013; Bodnar *et al.*, 2018). CDC48 is involved in a number of growth and development processes including vesicle trafficking, mitochondrial and proteasomal degradation of ubiquitinated proteins (Taylor and Rutter, 2011; Barthelme and Sauer, 2013). In *Arabidopsis*, it localizes across a number of subcellular components including the plasma membrane, cytoplasm, endoplasmic reticulum and nucleus (Aker *et al.*, 2006; Gallois *et al.*, 2013). Previous studies in *Drosophila* and *Arabidopsis* have shown that CDC48 undergoes S-nitrosylation at conserved cysteine sites inhibiting the ATP hydrolysis activity (Noguchi *et al.*, 2005; Fares *et al.*, 2011). Recently a similar event has been elucidated in *Nicotiana tabacum* during interaction with *Phytophthora cryptogea*, an oomycete pathogen infecting several ornamental plants (Rosnoblet *et al.*, 2017). The pathogen produces several halo proteins known as elicitors that act as avirulence factors in *Phytophthora cryptogea*-tobacco

interactions and function as *Phytophthora* pathogenicity factors on other plants (Panabières and Le Berre, 1999). The *c*Gryptogein protein is one of the elicitors produced from *Phytophthora cryptogea* during interaction with tobacco and has been shown to induce the S-nitrosylation of NtCDC48 at Cys 526 *in vivo* in *Nicotiana* cells compromising immune responses (Astier *et al.*, 2012a; Rosnoblelet *et al.*, 2017). Thus, S-nitrosylation may be exploited by both hosts and pathogens to promote either immunity or virulence, respectively.

S-nitrosylation regulates the zinc finger transcription factor, SRG1

By applying global expression profiling techniques, Cui *et al.*, (2018) uncovered a gene termed, *S-nitrosothiol regulated gene 1* (*SRG1*), which belonged to C2H2 zinc finger containing transcriptional factor (TF) family that was exquisitely sensitive to transcriptional regulation by NO. Further, accrual of *SRG1* transcripts also occurred following pathogen challenge. The C2H2 zinc finger domain is ubiquitous in higher eukaryotes. This domain consists of 28-30 amino acid residues arranged in β and α sheets. Structure stabilization is achieved by two conserved Cys of the β sheet and two conserved histidine at the C-terminus of the α sheet coordinating a zinc atom (Fedotova *et al.*, 2017). Transgenic *Arabidopsis* plants over expressing *SRG1* have increased hypersensitive response like cell death, increased ROS accumulation, increased *Pathogenesis Related 1* (*PR1*) gene expression and enhanced resistance to *Pst*DC3000 bacteria. In addition loss-of-function mutants of *SRG1* have increased susceptibility to *Pst*DC3000 (Cui *et al.*, 2018). These results suggests that *SRG1* is a positive regulator of plant immunity. However, *SRG1* possess an ETHYLENE RESPONSE FACTOR associated amphiphilic repression (EAR) domain located at the C-terminus and this enables an interaction with the corepressor, TOPLESS, to form a transcriptional repressor complex (Martin-arevalillo *et al.*, 2017; Cui *et al.*, 2018). The transcriptional repression activity of *SRG1* was validated *in vivo* in *Arabidopsis* protoplasts where *SRG1* was found to reduce the expression of *Luciferase* (*LUC*) reporter gene to $\approx 50\%$ (Cui *et al.*, 2018). Hence, *SRG1* appears to function as a transcription repressor. To function as an activator of plant immunity, *SRG1* therefore presumably must repress the activity of one or more repressors.

SRG1 exhibits reduced repressor activity in the *Arabidopsis thaliana* *S-nitrosoglutathione reductase* (*atgsnor1-3*) mutant relative to wild-type Col-0. The *atgsnor1-3* line has increased levels of *S-nitrosoglutathione* (GSNO), a natural NO donor. Thus, GSNOR1 controls the global bioavailability of GSNO (Feechan *et al.*, 2005a; Umbreen *et al.*, 2018, 2019). This suggests that the repressor activity of SRG1 is blunted by GSNO accumulation. Indeed, *S-nitrosylation* of SRG1 at Cys 87 was found to attenuate binding of this transcriptional repressor at the DNA motif AG/CT. This negatively regulated plant immunity (Fukushima *et al.*, 2012; Cui *et al.*, 2018). *S-nitrosylation* of SRG1 is thought to drive the abrogation of the zinc finger structure required for the functional activity of zinc finger proteins (Chasapis *et al.*, 2010). In mammalian systems, it is well established that NO can regulate the C2H2 zinc finger structure (Kroncke and Carlberg, 2000; Sha and Marshall, 2012), but the underpinning molecular mechanism has not been elucidated.

Collectively, these data suggest a model whereby following pathogen recognition and the activation of a nitrosative burst, leading to NO accrual, *SRG1* transcription is activated. Subsequently, SRG1 binds to the target promoter of one or more immune repressors possessing an AG/CT motif, suppressing the function of the cognate gene(s). This contributes to the activation of plant immunity. Subsequently, as the NO concentration increases during the later stages of the defence response, SRG1 becomes *S-nitrosylated* at Cys87 inhibiting its DNA binding activity and by extension, relieving the suppression of one or more immune repressors. The expression of these repressors may then contribute to the attenuation of the transient immune response. Thus, SRG1 function enables both the activation and subsequent repression of plant immunity and this switch is regulated by the *S-nitrosylation* of this C2H2 zinc finger transcription factor.

S-nitrosylation targets GSNOR for autophagy

Autophagy is a major catabolic route in eukaryotes degrading not only proteins but also entire organelle complexes (Klionsky and Codogno, 2013). The autophagy process is initiated when the lipid bilayer phagophore contributed by the endoplasmic reticulum and / or *trans*-Golgi apparatus engulf the intracellular cytoplasmic cargo leading to the formation of an autophagosome. The autophagosome fuses with the vacuole degrading the delivered materials. The products of degradation are then recycled for the construction of macromolecules (Glick *et al.*, 2010; Sieńko *et al.*, 2020; Stefaniak *et al.*, 2020). A number of autophagy related protein genes (ATG) play significant roles in autophagosome formation, but the two ubiquitination-like system of ATG8 lipidation and the ATG12 protein conjugation system are especially well investigated (Yoshimoto *et al.*, 2010; Yoshimoto and Ohsumi, 2018). In the ATG8 lipidation system, the C-terminal extension of ATG8 is processed by a Cys protease, ATG4, exposing the glycine residue. The truncated ATG8 is activated by an E1 enzyme, ATG7, via the formation of a thioester bond between the glycine residue of ATG8 and the Cys residue of ATG7. ATG8 is then transferred to a Cys residue of an E2 enzyme ATG3 and finally conjugated to a phospholipid phosphatidylethanolamine (PE) head group, leading to the formation of an autophagosome (Ichimura *et al.*, 2000). In the ATG12 conjugation system, ATG12 is activated by ATG7 and transferred to the Cys residue of an E2 like enzyme, ATG10. ATG12 is finally conjugated on to ATG5, via isopeptide bond formation between the lysine residue of ATG5 and the glycine residue of ATG12. The ATG12-ATG5 complex is subsequently able to mediate autophagosome formation (Mizushima *et al.*, 1998; Yoshimoto *et al.*, 2010; Yoshimoto and Ohsumi, 2018).

S-nitrosylation has been shown to control the availability of GSNOR1 by targeting it to autophagy pathway (Zhan *et al.*, 2018). GSNOR1 is a master regulator of intracellular levels of NO and proteome wide S-nitrosylation (Feechan *et al.*, 2005^b). Previous reports have shown that GSNOR1 can be S-nitrosylated constraining its de-nitrosylation functions achieved by scavenging GSNO (Frunghillo *et al.*, 2014). Recently, additional data has emerged,

confirming the S-nitrosylation of this key redox regulatory enzyme (Zhan *et al.*, 2018). S-nitrosylation of GSNOR1 has been shown to occur predominantly at Cys 10 and 370. *In vitro* experiments in the presence of NO donors and overexpression of GSNOR1 in *Arabidopsis NO overexpressor1 (nox1)* mutants, which exhibit elevated levels of NO, resulted in the instability of GSNOR1 (Frunghillo *et al.*, 2014). The degradation of GSNOR1 is attributed to structural modification mediated by S-nitrosylation. Autophagy related lysosomes were demonstrated as the final destination of S-nitrosylated GSNOR1. Proteasome inhibitors did not enhance the accumulation of S-nitrosylated GSNOR1, while autophagy inhibitors significantly elevated the levels of this enzyme. Thus, S-nitrosylated GSNOR1 is targeted to the autophagy pathway rather than the proteasome pathway. Moreover, elevated levels of GSNOR1 were detected in autophagosomes and in *atg2*, *atg5*, *atg7* and *atg10* mutant plants (Zhan *et al.*, 2018).

Selective autophagy is induced by the intermolecular β sheet interaction between ATG8 and the ATG8 interacting motif (AIM) of the substrate (Liu and Bassham, 2012; Michaeli *et al.*, 2016). The AIM motif in GSNOR1 is located in a β sheet between residues 152-155. The motif is located in a pocket close to Cys 10. S-nitrosylation of GSNOR1 occurs at Cys 10 and Cys 370, respectively. Though the functional role of SNO formation at Cys 370 is not elucidated, the data suggests that The S-nitrosylation of GSNOR1 at Cys 10 introduces a conformation change in protein the structure promoting the interaction of ATG8 and GSNOR1 (Zhan *et al.*, 2018). This Such interaction results in to GSNOR1 being directed into Though the functional role of Cys 370 is not elucidated, the data suggests that S-nitrosylation of GSNOR1 at Cys 10 increases interactions with the the autophagy pathway leading to GSNOR lysosome degradation. Although autophagic negative regulation of GSNOR1 mediated by S-nitrosylation could enhance the adaptability of plants to hypoxia (Zhan *et al.*, 2018), the loss of GSNOR1 in *Arabidopsis* is associated with significant immune defects (Feechan *et al.*, 2005a; Chen *et al.*, 2009; Yu *et al.*, 2012; Arasimowicz and Floryszak, 2016). Overall this information suggests that excessive autophagy-dependent degradation of GSNOR1-SNO would promote the

virulence of both adapted and non-pathogens (Figure 1). Perhaps the function of this cellular GSNOR degradation is to help maintain a supply of “fresh” GSNOR1, ensuring effective GSNOR1 activity (Feechan *et al.*, 2005a; Chen *et al.*, 2009; Yu *et al.*, 2012; Arasimowicz and Floryszak, 2016; Zhan *et al.*, 2018).

S-Nitrosylation regulates SUMOylation during plant immunity

SUMOylation, a reversible post-translational modification, occurring via the covalent attachment of the small ubiquitin-like modifier (SUMO) to target proteins through an ATP dependent reaction involving activation (E1), conjugation (E2) and ligation (E3) enzymes. SUMOylation is biochemically similar to ubiquitination but functionally distinct. SUMO is conserved from yeast to humans and the SUMO pathway is essential for the survival of eukaryotic cells (Nacerddine *et al.*, 2005; Saracco *et al.*, 2007; Zhao, 2018). In animals, SUMOylation has been studied extensively and shown to regulate numerous biological functions including various human diseases such as mental disorders, strokes, cancers, neurodegenerative and heart diseases (Coppola *et al.*, 2009; Mun *et al.*, 2016; Anderson *et al.*, 2017; Liu *et al.*, 2017; Thomas and Yang, 2017).

In plants, SUMOylation has been reported to regulate immunity via the deployment of host defence responses. In tomato, T-SUMO (Homologue of human SUMO1) protein interacts with EIX (ethylene inducing xylanase) of the fungus *Trichoderma viridae* which resulted in rapid induction of defence responses by production of ethylene (Hanania *et al.*, 1999). Pathogen triggered cell death is suppressed when T-SUMO was over-expressed and anti-sense lines displayed reverse effects, suggesting SUMO negatively regulates plant immunity in tomato (Hanania *et al.*, 1999). *Arabidopsis thaliana* SAP and MIZ1 (AtSIZ1) currently the only E3 ligase identified in *Arabidopsis*, was reported to downregulate plant immunity, as an *AtSIZ1* loss-of-function mutation produced higher levels of salicylic acid (SA) with increased expression of *PR1* and hence increased resistance to *Pst*DC3000 (Lee *et al.*, 2007). Similarly, a knock-down mutant for SUMO1/2 which has reduced levels of global SUMOylation showed

a 10-fold increase in *PR1* gene expression and increased resistance to *Pst*DC3000 (van den Burg *et al.*, 2010). Interestingly, the master regulator of SA signalling, NPR1 (NONEXPRESSOR OF PATHOGENESIS RELATED 1) acts as a SUMO substrate. NPR1 stability and binding with TFs is mediated by its SUMOylation, which is required for plant immunity (Saleh *et al.*, 2015). Interestingly, recent studies have suggested that S-nitrosylation regulates global 1/2 SUMOylation, controlling plant immunity, by directly modifying the SUMO E2 enzyme, SUMO Conjugating Enzyme 1 (SCE1), at Cys 139 (Skelly *et al.*, 2019; Gupta *et al.*, 2020). SUMO1/2 SUMOylation is thought to contribute to the negative regulation of plant immunity, maintaining the repression of plant defence responses in the absence of attempted pathogen infection (Skelly *et al.*, 2019). However, following pathogen challenge, a nitrosative burst leads to increased levels of NO accumulation, driving S-nitrosylation of SCE1 at Cys139. SNO formation at this Cys residue disables the E2 activity of SCE1 and thereby decreases global SUMOylation levels enabling the synthesis of the immune activator, SA and the activation of SA-dependent immune responses (Skelly *et al.*, 2019). Correspondingly, SCE1 Cys139S mutants, which are insensitive to S-nitrosylation, showed higher levels of SUMO conjugates and increased disease susceptibility against *Pst*DC3000. These plant lines exhibited reduced levels of endogenous SA accumulation following pathogen challenge but respond to SA in a similar fashion to wild-type plants: i.e. strong induction of SA-dependent genes upon exogenous SA application (Skelly *et al.*, 2019). Thus, S-nitrosylation of SCE1 at Cys139 may enable SA accumulation, which subsequently supports the deployment of SA-dependent defence responses (Figure 2).

Importantly, SUMOylation was previously thought to be regulated at the local level with the SUMO modification of individual substrates controlled. These findings establish a parallel and complementary paradigm, suggesting that SUMOylation can also be regulated at a global level, by the control of SCE1 by S-nitrosylation. Significantly, the function of the human homologue of SCE1, UBC9, was also found to be compromised by S-nitrosylation at the

evolutionary conserved Cys139 residue. Therefore, SNO formation at Cys139 of SCE1/UBC9 may regulate SUMOylation across phylogenetic kingdoms.

Future perspectives

Microbial pathogens deliver a plethora of effector proteins to the inside of plant cells to support pathogenesis. The work of Ling and coworkers establishes for the first time that challenged plant host cells can deploy S-nitrosylation to disable the function of effectors. Many pathogen effector proteins are rich in Cys residues and presumably, some of these must be solvent exposed, potentially providing target sites for S-nitrosylation. It will therefore be informative to determine if other effectors are disabled by this redox-based PTM. In a similar fashion, are other proteins, in addition to GSNOR, targeted by SNO formation for autophagy? Critically, are autophagy mutants disrupted in (S)NO homeostasis? Finally, it will be informative to explore if S-nitrosylation regulates other aspects of SUMOylation. It is becoming increasingly clear that S-nitrosylation is a key regulator of a plethora of molecular features underpinning the establishment of plant immunity.

Author Contributions: All the authors, J.L., S.U. and G.J.L., contributed equally to the writing and editing of this paper. All authors have read and approved the final version of the paper.

Acknowledgement: Jibril Lubega would like to thank Islamic Development Bank for funding his Doctoral studies at The University of Edinburgh. Research on S-nitrosylation within the Loake lab is supported by the BBSRC.

References

- Aker J, Borst JW, Karlova R, de Vries S.** 2006. The *Arabidopsis thaliana* AAA protein CDC48A interacts in vivo with the somatic embryogenesis receptor-like kinase 1 receptor at the plasma membrane. *Journal of Structural Biology* **156**, 62–71.
- Anderson DB, Zanella CA, Henley JM, Cimarosti H.** 2017. Sumoylation: Implications for neurodegenerative diseases (VG Wilson, Ed.). *Advances in Experimental Medicine and Biology* **963**, 261–281.
- Arasimowicz-Jelonek M, Floryszak-Wieczorek J.** 2016. Nitric oxide in the offensive strategy of fungal and Oomycete plant pathogens. *Frontiers in Plant Science* **7**, doi: 10.3389.
- Asai S, Shirasu K.** 2015. Plant cells under siege: Plant immune system versus pathogen effectors. *Current Opinion in Plant Biology* **28**, 1–8.
- Astier J, Besson-Bard A, Lamotte O, Bertoldo J, Bourque S, Terenzi H, Wendehenne D.** 2012a. Nitric oxide inhibits the ATPase activity of the chaperone-like AAA + ATPase CDC48, a target for S-nitrosylation in cryptogeiin signalling in tobacco cells. *Biochemical Journal* **447**, 249–260.
- Astier J, Gross I, Durner J.** 2018. Nitric oxide production in plants: An update. *Journal of Experimental Botany* **69**, 3401–3411.
- Astier J, Kulik A, Koen E, Besson-Bard A, Bourque S, Jeandroz S, Lamotte O, Wendehenne D.** 2012b. Protein S-nitrosylation: What's going on in plants? *Free Radical Biology and Medicine* **53**, 1101–1110.
- Baek GH, Cheng H, Choe V, Bao X, Shao J, Luo S, Rao H.** 2013. Cdc48: A Swiss Army Knife of Cell Biology. *Journal of Amino Acids* **2013**, 183421. doi: 10.1155/2013/183421. ~~4–12.~~
- Barthelme D, Sauer RT.** 2013. Bipartite determinants mediate an evolutionarily conserved interaction between Cdc48 and the 20S peptidase. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 3327–3332.
- Begara-Morales JC, Sánchez-Calvo B, Chaki M, Valderrama R, Mata-Pérez C, López-Jaramillo J, Padilla MN, Carreras A, Corpas FJ, Barroso JB.** 2014. Dual regulation of

cytosolic ascorbate peroxidase (APX) by tyrosine nitration and S-nitrosylation. *Journal of Experimental Botany* **65**, 527–538.

Bodnar NO, Kim KH, Ji Z, Wales TE, Svetlov V, Nudler E, Engen JR, Walz T, Rapoport TA. 2018. Structure of the cdc48 atpase with its ubiquitin-binding cofactor ufd1–npl4. *Nature Structural and Molecular Biology* **25**, 616–622.

Bolwell GP, Daudi A. 2009. Reactive oxygen species in plant-pathogen interactions. In: Rio L., Puppo A. (eds) Reactive Oxygen Species in Plant Signaling. Signaling and Communication in Plants, Springer, Berlin, Heidelberg. <https://doi.org/10.1>.

~~**Bolwell GP, Daudi A.** 2009. Reactive oxygen species in plant-pathogen interactions. [Signaling and Communications in plants](#), 113–133.~~

van den Burg HA, Kini RK, Schuurink RC, Takken FLW. 2010. Arabidopsis small ubiquitin-like modifier paralogs have distinct functions in development and defense. *The Plant Cell* **22**, 1998–2016.

Buttner D, He SY. 2009. Type III protein secretion in plant pathogenic bacteria. *Plant Physiology* **150**, 1656–1664.

Chamizo-Ampudia A, Sanz-Luque E, Llamas A, Galvan A, Fernandez E. 2017. Nitrate reductase regulates plant nitric oxide homeostasis. *Trends in Plant Science* **22**, 163–174.

Chasapis CT, Loutsidou AK, Orkoulas MG, Spyroulias GA. 2010. Zinc binding properties of engineered ring finger domain of arkadia E₃ ubiquitin ligase. *Bioinorganic Chemistry and Applications*, **2010**, 323152. doi: 10/1155/2010/323152. ~~4–7.~~

Chen R, Sun S, Wang C, Li Y, Liang Y, An F, Li C. 2009. The Arabidopsis PARAQUAT RESISTANT2 gene encodes an S-nitrosoglutathione reductase that is a key regulator of cell death. *Cell Research* **19**, 1377–1387.

Coppola D, Parikh V, Boulware D, Blanck G. 2009. Substantially reduced expression of PIAS1 is associated with colon cancer development. *Journal of Cancer Research and Clinical Oncology* **135**, 1287–1291.

Corcoran A, Cotter TG. 2013. Redox regulation of protein kinases. *FEBS Journal* **280**, 1944–1965.

Couturier J, Chibani K, Jacquot J-P, Rouhier N. 2013. Cysteine-based redox regulation and signaling in plants. *Frontiers in plant science* **4**, 105.

Cui B, Pan Q, Clarke D, Villarreal MO, Umbreen S, Yuan B, Shan W, Jiang J, Loake GJ. 2018. S-nitrosylation of the zinc finger protein SRG1 regulates plant immunity. *Nature Communications* **9**, 4226. <https://doi.org/10.1038/s41467-018-06578-3>.

Davidsson PR, Kariola T, Niemi O, ~~Tapio~~ Palva ~~IE~~. 2013. Pathogenicity of and plant immunity to soft rot pectobacteria. *Frontiers in Plant Science* **4**, [191](#)–[13](#).

Fares A, Rossignol M, Peltier JB. 2011. Proteomics investigation of endogenous S-nitrosylation in Arabidopsis. *Biochemical and Biophysical Research Communications* **416**, 331–336.

Fedotova AA, Bonchuk AN, Mogila VA, Georgiev PG. 2017. C2H2 zinc finger proteins : The largest but poorly explored family of higher eukaryotic transcription factors. *Reviews* **9**, 47–58.

~~**Feechan A, Kwon E, Yun BW, Wang Y, Pallas JA, Loake GJ.** 2005a. A central role for S-nitrosothiols in plant disease resistance. *PNAS* **102**, 1–6.~~

Feechan A, Kwon E, Yun B-W, Wang Y, Pallas JA, Loake GJ. 2005**b**. A central role for S-nitrosothiols in plant disease resistance. *Proceedings of the National Academy of Sciences* **102**, 8054–8059.

Frederickson Matika DE, Loake GJ. 2014. Redox regulation in plant immune function. *Antioxidants & redox signaling* **21**, 1373–88.

Friso G, van Wijk KJ. 2015. Post translational protein modifications in plant metabolism. *Plant Physiology* **169**, 1469–1487.

Frungillo L, Skelly MJ, Loake GJ, Spoel SH, Salgado I. 2014. S-nitrosothiols regulate nitric oxide production and storage in plants through the nitrogen assimilation pathway. *Nature Communications* **5**, [5401](#). doi: [10.1038/ncomms6401](https://doi.org/10.1038/ncomms6401). [4](#)–[10](#).

Fukushima S, Yoshida M, Takatsuji H. 2012. Role of linkers between zinc fingers in spacing

recognition by plant TFIIIA-type zinc-finger proteins. Journal of Amino acids [2012](#), [848037](#).doi:10.1155.

Gallois JL, Drouaud J, Lécureuil A, Guyon-Debast A, Bonhomme S, Guerche P. 2013. Functional characterization of the plant ubiquitin regulatory X (UBX) domain-containing protein AtPUX7 in Arabidopsis thaliana. Gene **526**, 299–308.

Gardner PR. 2012. Hemoglobin : A nitric-oxide dioxygenase. Scientifica [2012](#), [683729](#). doi: [10.6064/2012/683729](#). [683729](#).

Gaupels F, Spiazzi-Vandelle E, Yang D, Delledonne M. 2011. Detection of peroxynitrite accumulation in Arabidopsis thaliana during the hypersensitive defense response. Nitric Oxide **25**, 222–228.

Glick D, Barth S, Macleod KF. 2010. Autophagy : cellular and molecular mechanisms. Journal of Pathology **221**, [3-12](#). [3-12](#).

Grant JJ, Loake GJ. 2000. Update on signaling role of reactive oxygen intermediates and cognate redox signaling in disease resistance. Plant molecular biology **124**, 21–29.

Gupta KJ, Kolbert Z, Durner J, Lindermayr C, Corpas FJ, Brouquisse R, Barroso JB, Umbreen S, Palma M, Hancock JT, Petrivalsky M, Wendehenne D, Loake GJ. 2020. Regulating the regulator: nitric oxide control of post-translational modifications. New Phytologist [227](#), [1319-1325](#). doi:10.1111.

Hanania U, Furman-Matarasso N, Ron M, Avni A. 1999. Isolation of a novel SUMO protein from tomato that suppresses EIX-induced cell death. The Plant Journal **19**, 533–541.

Hann DR, Domínguez-Ferreras A, Motyka V, Dobrev PI, Schornack S, Jehle A, Felix G, Chinchilla D, Rathjen JP, Boller T. 2014. The Pseudomonas type III effector HopQ1 activates cytokinin signaling and interferes with plant innate immunity. New Phytologist **201**, 585–598.

Ichimura Y, Kirisako T, Takao T, Satomi Y, Shimonishi Y, Ishihara N, Mizushima N, Tanida I, Kominami E, Ohsumi M, Noda T, Ohsumi Y. 2000. A ubiquitin-like system mediates protein lipidation. Nature **408**, 488-492.

Kamoun S. 2007. Groovy times: filamentous pathogen effectors revealed. Current Opinion in

Plant Biology **10**, 358–365.

Klionsky DJ, Codogno P. 2013. The mechanism and physiological function of macroautophagy. *Journal of Innate Immunity* **5**, 427–433.

Kolbert Z, Barroso JB, Brouquisse R, Corpas FJ, Durner J, Gupta KJ, Lindermayr C, Loake GJ, Palma JM, Petrivalsky M, Wendehenne D, Hancock JT. 2019. Celebrating forty years of plant nitric oxide research. *Nitric Oxide* **1**, 53-70.

Kroncke KD, Carlberg C. 2000. Inactivation of zinc finger transcription factors provides a mechanism for a gene regulatory role of nitric oxide. *FASEB Journal* **14**, 166–173.

Lee J, Nam J, Park HC, Na G, Miura K, Jin JB, Yoo CY, Baek D, Kim DH, Jeong JC, Kim D, Lee SY, Salt DE, Mengiste T, Gong Q, Ma S, Bohnert HJ, Kwak SS, Bressan RA, Hasegawa PM, Yun DJ. 2007. Salicylic acid-mediated innate immunity in *Arabidopsis* is regulated by SIZ1 SUMO E3 ligase. *The Plant Journal* **49**, 79–90.

Lindermayr C, Sell S, Muller B, Leister D, Durner J. 2010. Redox regulation of the NPR1-TGA1 system of *Arabidopsis thaliana* by nitric oxide **22**, 2894-2907.

Ling T, Bellin D, Vandelle E, Imanifard Z, Delledonne M. 2017. Host-mediated S-nitrosylation disarms the bacterial effector HopAI1 to re-establish immunity. *The plant cell* **29**, 2871–2881.

Lisjak M, Teklic T, Wilson ID, Whiteman M, Hancock JT. 2013. Hydrogen sulfide: Environmental factor or signalling molecule? *Plant, Cell and Environment* **36**, 1607–1616.

Liu Y, Bassham DC. 2012. Autophagy: Pathways for Self-Eating in Plant Cells. *Annual Review of Plant Biology* **63**, 215–237.

Liu C, Pedersen C, Schultz-Larsen T, Aguilar GB, Madriz-Ordeñana K, Hovmøller MS, Thordal-Christensen H. 2016. The stripe rust fungal effector PEC6 suppresses pattern-triggered immunity in a host species-independent manner and interacts with adenosine kinases. *New Phytologist* **213**, 1556. [doi: 10.1111/nph.14362](https://doi.org/10.1111/nph.14362).

Liu Y, Zhao D, Qiu F, Zhang LL, Liu SK, Li YY, Liu MT, Wu D, Wang JX, Ding XQ, Liu YX, Dong CJ, Shao XQ, Yang BF, Chu WF. 2017. Manipulating PML SUMOylation via silencing UBC9 and RNF4 regulates cardiac fibrosis. *Molecular Therapy* **25**, 666–678.

- Martin-arevalillo R, Nanao MH, Larrieu A, Vinos-poyo T, Mast D.** 2017. Structure of the Arabidopsis TOPLESS corepressor provides insight into the evolution of transcriptional repression. *Proceedings of the National Academy of Sciences* **114**, 8107–8112.
- Michaeli S, Galili G, Genschik P, Fernie AR, Avin-Wittenberg T.** 2016. Autophagy in plants - what's new on the Menu? *Trends in Plant Science* **21**, 134–144.
- Mizushima N, Noda T, Yoshimori T, Tanaka Y, Ishii T, George DM, Klionsky JD, Ohsumi M, Ohsumi Y.** 1998. A protein conjugation system essential for autophagy. *Nature* **395**, 395–398.
- Mun M-J, Kim J-H, Choi J-Y, Kim M-S, Jang W-C, Lee JJ, Eun YL, Kwak S-J, Kim KW, Lee SB.** 2016. Polymorphisms of small ubiquitin-related modifier genes are associated with risk of Alzheimer's disease in Korean: A case-control study. *Journal of the Neurological Sciences* **364**, 122–127.
- Nacerddine K, Bhaumik M, Cohen-tannoudji M, Babinet C, Pandolfi PP, Dejean A.** 2005. The SUMO pathway is essential for nuclear integrity and chromosome segregation in mice. *Developmental cell* **9**, 769–779.
- Noguchi M, Takata T, Kimura Y, Manno A, Murakami K, Koike M, Ohizumi H, Hori S, Kakizuka A.** 2005. ATPase activity of p97/valosin-containing protein is regulated by oxidative modification of the evolutionally conserved cysteine 522 residue in walker a motif. *Journal of Biological Chemistry* **280**, 41332–41341.
- Panabières F, Le Berre JY.** 1999. A family of repeated DNA in the genome of the oomycete plant pathogen *Phytophthora cryptogea*. *Current Genetics* **36**, 105–112.
- Park C-H, Chen S, Shirsekar G, Zhou B, Khang CH, Songkumarn P, Afzal AJ, Ning Y, Wang R, Bellizzi M, Valent B, Wang GL.** 2012. The *Magnaporthe oryzae* effector AvrPiz-t targets the RING E3 ubiquitin ligase APIP6 to suppress pathogen-associated molecular pattern-triggered immunity in rice. *The Plant Cell* **24**, 4748–4762.
- Pfeilmeier S, Caly DL, Malone JG.** 2016. Bacterial pathogenesis of plants: future challenges from a microbial perspective. *Molecular plant pathology* **17**, 1298–1313.
- Rockel P, Strube F, Rockel A, Wildt J, Kaiser WM.** 2002. Regulation of nitric oxide (NO)

production by plant nitrate reductase in vivo and in vitro. *Journal of Experimental Botany* **53**, 103–110.

Romero-Puertas MC, Laxa M, Matte A, Zaninotto F, Finkemeier I, Jones AME, Perazzolli M, Vandelle E, Dietz K-J, Delledonne M. 2007. S-Nitrosylation of peroxiredoxin II E promotes peroxynitrite-mediated tyrosine nitration. *The Plant Cell Online* **19**, 4120–4130.

Rosnoblet C, Bègue H, Blanchard C, Pichereaux C, Besson-Bard A, Aimé S, Wendehenne D. 2017. Functional characterization of the chaperon-like protein Cdc48 in cryptogein-induced immune response in tobacco. *Plant Cell and Environment* **40**, 491–508.

Saleh A, Withers J, Mohan R, Marqués J, Gu Y, Yan S, Zavaliev R, Nomoto M, Tada Y, Dong X. 2015. Post translational modifications of the master transcriptional regulator NPR1 enable dynamic but tight control of plant immune responses. *Cell Host and Microbe* **18**, 169–182.

Sanz L, Albertos P, Mateos I, Sánchez-Vicente I, Lechón T, Fernández-Marcos M, Lorenzo O. 2015. Nitric oxide (NO) and phytohormones crosstalk during early plant development. *Journal of Experimental Botany* **66**, 2857–2868.

Saracco SA, Miller MJ, Kurepa J, Vierstra RD. 2007. Genetic analysis of SUMOylation in *Arabidopsis*: conjugation of SUMO1 and SUMO2 to nuclear proteins is essential. *Plant Physiology* **145**, 119–134.

Schlicht M, Kombrink E. 2013. The role of nitric oxide in the interaction of *Arabidopsis thaliana* with the biotrophic fungi, *Golovinomyces orontii* and *Erysiphe pisi*. *Frontiers in Plant Science* **4**, [351. doi.3389/fpls.2013.00351](https://doi.org/10.3389/fpls.2013.00351). 4–12.

Setti B, Bencheikh M, Henni JE. 2014. Virulence strategies of phytopathogenic bacteria and their role in plant disease pathogenesis. *African Journal of Microbiology Research* **8**, 2809–2815.

Sha Y, Marshall HE. 2012. S-nitrosylation in the regulation of gene transcription. *Biochim Biophys Acta* **1820**, 701–711.

Sieńko K, Poormassalehgoo A, Yamada K, Goto-Yamada S. 2020. Microautophagy in Plants: Consideration of Its Molecular Mechanism. *Cells* **9**, 887.

Skelly MJ, Loake GJ. 2013. Synthesis of redox-active molecules and their signaling functions during the expression of plant disease resistance. *Antioxidants & redox signaling* **19**, 990–997.

Skelly MJ, Malik SI, Le Bihan T, Bo Y, Jiang J, Spoel SH, Loake GJ. 2019. A role for S-nitrosylation of the SUMO-conjugating enzyme SCE1 in plant immunity. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 17090–17095.

Snelders NC, Kettles GJ, Rudd JJ, Thomma BPHJ. 2018. Plant pathogen effector proteins as manipulators of host microbiomes? *Molecular Plant Pathology* **19**, 257–259.

Spadaro D, Yun BW, Spoel SH, Chu C, Wang YQ, Loake GJ. 2010. The redox switch: Dynamic regulation of protein function by cysteine modifications. *Physiologia Plantarum* **138**, 360–371.

Spoel SH, Loake GJ. 2011. Redox-based protein modifications: The missing link in plant immune signalling. *Current Opinion in Plant Biology* **14**, 358–364.

Stefaniak S, Wojtyła Ł, Pietrowska-Borek M, Borek S. 2020. Completing Autophagy: Formation and Degradation of the Autophagic Body and Metabolite Salvage in Plants. *International journal of molecular sciences* **21**, 2205. [doi: 10.3390/ijms21062205](https://doi.org/10.3390/ijms21062205).

Taj G, Agarwal P, Grant M, Kumar A. 2010. MAPK machinery in plants. Recognition and response to different stresses through multiple signal transduction pathways. *Plant Signaling & Behavior* **5**, 1370–1378.

Taylor EB, Rutter J. 2011. Mitochondrial quality control by the ubiquitin-proteasome system. *Biochemical Society Transactions* **39**, 1509–1513.

Thomas SN, Yang AJ. 2017. Mass spectrometry analysis of lysine posttranslational modifications of Tau protein from Alzheimer's disease brain (C Smet-Nocca, Ed.). *Methods in Molecular Biology* **1523**, 161–177.

Turkan I. 2017. Emerging roles for ROS and RNS – versatile molecules in plants. *Journal of Experimental Botany* **68**, 4413–4416. [doi:10.1093](https://doi.org/10.1093).

- Umbreen S, Lubega J, Cui B, Pan Q, Jiang J, Loake GJ.** 2018. Specificity in nitric oxide signalling. *Journal of Experimental Botany* **69**, 3439–3448. doi:10.1093.
- Umbreen S, Lubega J, Loake GJ.** 2019. Sulfur: The heart of nitric oxide-dependent redox signalling. *Journal of Experimental Botany* **70**, 4279–4286. doi:10.1093.
- Vanzo E, Ghirardo A, Merl-Pham J, Lindermayr C, Heller W, Hauck SM, Durner J, Schnitzler JP.** 2014. S-Nitroso-proteome in poplar leaves in response to acute ozone stress. *PLoS ONE* **9**, e106886. doi:10.1371/journal.pone.0106886.
- Wang S, Boevink PC, Welsh L, Zhang R, Whisson SC, Birch PRJ.** 2017. Delivery of cytoplasmic and apoplastic effectors from *Phytophthora infestans* haustoria by distinct secretion pathways. *New Phytologist* **216**, 205–215.
- Wang YQ, Feechan A, Yun BW, Shafiei R, Hofmann A, Taylor P, Xue P, Yang FQ, Xie ZS, Pallas JA, Chu CC, Loake GJ.** 2009. S-nitrosylation of AtSABP3 antagonizes the expression of plant immunity. *Journal of Biological Chemistry* **284**, 2131–2137.
- Wilkinson JQ, Crawford NM.** 1991. Identification of the Arabidopsis CHL3 gene as the nitrate reductase structural gene NIA2. *The Plant cell* **3**, 461–471.
- Yoshimoto K, Ohsumi Y.** 2018. Unveiling the molecular mechanisms of plant autophagy—from autophagosomes to vacuoles in plants. *Plant and Cell Physiology* **59**, 1337–1344.
- Yoshimoto K, Takano Y, Sakai Y.** 2010. Autophagy in plants and phytopathogens. *FEBS Letters* **584**, 1350–1358.
- Yu M, Lamattina L, Spoel SH, Loake GJ.** 2014. Nitric oxide function in plant biology: A redox cue in deconvolution. *New Phytologist* **202**, 1142–1156.
- Yu M, Yun BW, Spoel SH, Loake GJ.** 2012. A sleigh ride through the SNO: Regulation of plant immune function by protein S-nitrosylation. *Current Opinion in Plant Biology* **15**, 424–430.
- Yun B-W, Feechan A, Yin M, Saidi NBB, Le-Bihan T, Yu M, Moore JW, Kang JG, Kwon E, Spoel SH, Pallas J, Loake GJ.** 2011. S-nitrosylation of NADPH oxidase regulates cell death in plant immunity. *Nature* **478**, 264–268.
- Yun B, Skelly MJ, Yin M, Yu M, Mun B, Lee S, Hussain A, Spoel SH, Loake GJ.** 2016.

Nitric oxide and S-nitrosoglutathione function additively during plant immunity. *New Phytologist* **211**, 516–526.

Yun BW, Spoel SH, Loake GJ. 2012. Synthesis of and signalling by small, redox active molecules in the plant immune response. *Biochimica et Biophysica Acta - General Subjects* **1820**, 770–776.

Zhan N, Wang C, Chen L, Yang H, Feng J, Gong X, Ren B, Wu R, Mu J, Li Y, Liu Z, Zhou Y, Peng J, Wang K, Huang X, Xiao S, Zuo, J. 2018. S-Nitrosylation targets GSNO reductase for selective autophagy during hypoxia responses in plants. *Molecular Cell* **71**, 142-154.

Zhang J, Shao F, Li Y, Cui H, Chen L, Li H, Zou Y, Long C, Lan L, Chai J, Chen S, Tang X, Zhou JM. 2007. A *Pseudomonas syringae* effector inactivates MAPKs to suppress PAMP-induced immunity in plants. *Cell Host and Microbe* **1**, 175–185.

Zhao X. 2018. SUMO-mediated regulation of nuclear functions and signaling processes. *Molecular cell* **71**, 409–418.

Figure legends

Figure 1. Conformational changes in GSNOR1 by S-nitrosylation mediates its autophagy. Generated NO reacts with glutathione (GSH) to form GSNO. GSNOR1 reduces GSNO to oxidized GSH (GSSG) and ammonia. GSSG is further reduced to GSH by GSH reductase (GR). Changes in GSNOR1 activity regulates protein-SNO by modulating GSNO levels. **(B)** Free NO S-nitrosylates GSNOR1 at Cys10. The S-nitrosylation of GSNOR1 results in conformation changes which enhances the interaction with autophagy-related protein 8 (ATG8) through its ATG8 interacting motif (AIM) leading to autophagosome formation. The autophagy machinery targets GSNOR1 to lysosome degradation.

Figure 2. S-nitrosylation regulates SUMOylation during plant immunity. (A) In the absence of pathogen challenge and the associated nitrosative burst, SUMO conjugating enzyme 1 (SCE1) is not S-nitrosylated and consequently SCE1 drives SUMOylation resulting in the negative regulation of plant immunity. (B) Pathogen challenge results in the nitrosative burst leading to higher levels of NO, which results in S-nitrosylation of SCE1 at Cys 139, reducing ~~SUMO conjugating enzyme 1~~ (SCE1) activity and by extension SUMOylation, contributing to the release of plant immunity suppression.