

THE UNIVERSITY of EDINBURGH

Edinburgh Research Explorer

Uncertainty in below-ground carbon biomass for major land covers in Southeast Asia

Citation for published version:

Yuen, JQ, Ziegler, AD, Webb, EL & Ryan, CM 2013, 'Uncertainty in below-ground carbon biomass for major land covers in Southeast Asia', Forest Ecology and Management, vol. 310, pp. 915-926. https://doi.org/10.1016/j.foreco.2013.09.042

Digital Object Identifier (DOI):

10.1016/j.foreco.2013.09.042

Link:

Link to publication record in Edinburgh Research Explorer

Document Version: Early version, also known as pre-print

Published In: Forest Ecology and Management

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Uncertainty in Below-ground Carbon Biomass for Major Land Covers in Southeast Asia

Jia Qi Yuen¹, Alan D Ziegler^{1*}, Edward L Webb², Casey Ryan³

¹Department of Geography, National University of Singapore, ²Department of Biological Sciences, National University of Singapore ³School of Geosciences, University of Edinburgh

* Block AS02-03-01, 1 Arts Link Kent Ridge, Singapore 107568 (tel: +65 65166640; geoadz@nus.edu.sg).

10 Abstract

1 2

3 4

5

6

7 8

9

11 Owing to difficulties associated with measuring root biomass accurately in space and time, below-ground root biomass is 12 often calculated indirectly from above-ground biomass measurements via general allometric equations. Of concern is 13 that general equations may not provide accurate site-specific calculations for accurate carbon stock assessments. This 14 review comparing more than 100 root-related studies conducted in SE Asia shows highly variable and uncertain below-15 ground woody carbon (BGC) biomass estimates for many vegetation types associated with on-going land-use changes 16 throughout the region. Most BGC data exist for Indonesia, Malaysia, and Thailand; only a few studies have been 17 conducted for Brunei, Cambodia, Lao PDR, Myanmar, Papua New Guinea, Philippines, Singapore, Timor Leste and 18 Vietnam. While substantial data exist for a variety of forests and timber-plantations, little work has focused on key 19 transition land-covers including rubber, oil palm, swidden fallows, agroforests, grasslands, and croplands. Mangroves 20 (12-219 Mg C ha⁻¹), peat forests (11-71 Mg C ha⁻¹) and other forest types (11-74 Mg C ha⁻¹) have the highest BGC values. 21 The limited data for rubber plantations (5-32 Mg C ha⁻¹), oil palm plantations (4-22 Mg C ha⁻¹), swidden fallows (3-16 22 Mg C ha⁻¹), non-swidden agroforestry (3-16 Mg C ha⁻¹) indicate modest differences in the amount of BGC for several 23 land covers that are at the heart of ongoing debates regarding the human and environmental impacts of agricultural 24 intensification. The paucity of data currently in existence for the region highlights the need for additional field 25 investigations-following-accepted protocols, of root biomass to facilitate efforts to improve carbon stock estimates. 26 Government agencies, private enterprises, and development agencies could help lead the way in developing a better 27 forest carbon database by teaming with researchers to assess total ecosystem carbon stocks prior to vegetation being 28 removed for construction, mining, or stand rotations. 29

Keywords: carbon stocks, land-cover change in the tropics, mangroves, oil palm, peat forest, root shoot ratio, rubber plantations, shifting agriculture

33 1. Introduction

34 Roots contribute up to half of the carbon cycled annually in forests; and they may account for approximately one-third 35 of global annual net primary production (Vogt et al., 1996; Jackson et al., 1997). Coarse roots provide structural support 36 and access to deep water (e.g. tap roots), while fine roots absorb water and nutrients from the surrounding soils (Jackson 37 et al., 1997; Körner, 1994; Schulze, 1983; Shi et al., 2008). Although the two root sizes are functionally different and 38 turnover at different rates, they are both important stores of biomass carbon (Clark et al., 2001; Rasse et al., 2005). 39 Carbon sequestered in roots during root growth and maintenance is eventually transferred to the soil when they turnover or die (Albrecht et al., 2004; Chalermchatwilai et al., 2011; Srivastava et al., 1986). As root-derived carbon has a long 40 41 residence time (cf. Abiven et al., 2005; Rasse et al., 2005; Sanaullah et al., 2005), below-ground woody carbon biomass 42 (BGC) is an important component of the terrestrial carbon budget. When soils are tilled, organic matter previously 43 protected from microbial action is decomposed rapidly because of changes in water, air, and temperature conditions; and 44 the breakdown of soil aggregates accelerates erosion (Sundermeier et al., 2012). Erosion, tillage, and other activities that 45 overturn and expose the soil can lead to important losses of below ground carbon. In addition, biomass burning is a major source of terrestrial carbon transfer to the atmosphere in gas form (Quéré et al., 2009). Much attention is 46 47 currently focused on reducing the loss of terrestrial carbon in both above- and below-ground stores following land-cover 48 conversion, particularly in tropical regions (Ziegler et al., 2012).

49 Forest carbon conservation in developing countries is suggested as an effective means of reducing greenhouse 50 gas emissions (Stern, 2007). For example, the United Nations Framework Convention on Climate Change (UNFCCC) 51 program for Reducing Emissions from Deforestation and forest Degradation (REDD+) is designed to 52 preserve/increase the storage of terrestrial carbon, meanwhile fostering beneficial ecosystem services and promoting 53 human livelihoods (UNFCC, 2010; 2011). Under REDD+, developing countries would receive payments from 54 industrialized nations for achieving long-term reductions in deforestation and/or replacing some land-use activities with 55 others that sequester more carbon (UNFCC, 2010; 2011). Approximately US\$4 billion was pledged for REDD+ 56 programs between 2010 and 2012 (Ballesteros et al., 2011). As of September 2013, Southeast Asia hosted number of

early REDD+ type projects (Table 1): Indonesia (44 projects), Cambodia (four projects), Malaysia (one project),
Vietnam (seven projects), Thailand (one project), Papua New Guinea (four projects), the Philippines (four projects), and
Lao PDR (one project). Several countries in the region have also started national-level preparations to engage with a
future REDD+ mechanism (CIFOR, 2011; FCPF, 2011); for example, of the 16 countries globally that have established
UN-REDD national programs, seven of those are located in the Asia-Pacific region (Cambodia, Indonesia, Papua New
Guinea, the Philippines, Solomon Islands, Sri Lanka; Vietnam, UN-REDD Programme, 2009).

63 Ideally, eligibility for financial remuneration by REDD+ requires participating countries to have accurate 64 estimates of carbon stocks and emissions associated with all important land cover transitions (Brown, 2002; UNFCCC, 65 2009). While above-ground carbon of various land covers is frequently measured, and new techniques are emerging to make AGC calculations more reliable (cf. Gibbs et al., 2007; Tollefson, 2009), much less work has addressed estimating 66 67 below-ground woody carbon biomass (cf. Mokany et al., 2006; Vogt et al., 1996; Ziegler et al., 2012). In a recent meta-68 analysis based on more than 250 studies, we found great uncertainty in total ecosystem carbon for several major land 69 covers that are related to important land-use transitions in SE Asia (Ziegler et al., 2012). Some of this uncertainty 70 stemmed from our calculation of BGC from a limited number of root:shoot ratio (RSR) data readily available in the 71 literature. Herein, we improve upon these carbon stock estimates by reviewing relevant studies/papers of below ground 72 root biomass from the SE Asia region. In addition to providing a summary of BGC estimates and root:shoot ratios for 73 vegetation types that are commonly associated with on-going and projected land-cover change, we also assess data 74 availability and quality, as they affect the accuracy of carbon accounting for land-cover change scenarios. 75

76 2. Review of below-ground woody carbon biomass

77 The countries considered in this review are Brunei, Cambodia, southern China, Indonesia, Lao PDR, Malaysia 78 (Peninsular and Insular combined), Myanmar, Papua New Guinea, the Phillippines, Singapore, Thailand, Timor Leste, 79 and Vietnam. Many of these countries are affected by on-going and drastic land-cover conversions, including forest 80 conversion to permanent cropping systems and/or plantations (e.g., rubber, oil palm), transitions from swidden 81 agriculture (shifting-agriculture) to more permanent agriculture types, logging, wetland forest (mangroves, peat forest) 82 degradation or conversion, afforestation/reforestation, and abandonment of marginal lands. We focused on the 83 following eleven major land covers related to important land-cover/land-use (LCLU) transitions now taking place in the 84 region (Fox et al., 2012; Ziegler et al., 2012; van Vliet et al., 2012): forest (FOR), logged over forest (LOF), mangrove 85 (MAN), peat forest (PF), orchard and tree-plantation (OTP), non-swidden agroforest (AGF), rubber plantation (RP), 86 swidden fallows of any length (SF), oil palm plantation (OP), grassland, pasture or shrub land (GPS) and permanent 87 cropland (PC). For each land cover, we collated literature-reported estimates of root carbon biomass and root:shoot 88 ratios (RSR). Except for Indonesia (the country with the most data), and low end-members Brunei, Laos, and Myanmar 89 (countries with only one forest study), no correlation existed between data availability and the number of 90 proposed/ongoing REDD+ type projects (Table 1). Some countries with substantial REDD+ activity (excluding 91 Indonesia) had limited data-e.g., Vietnam, Papua New Guinea, Cambodia, and the Philippines.

92 Characteristics of individual land-covers appearing in the 11 categories, and notes regarding the biomass 93 determinations for each study, are listed by country in Table S1. Biomass values for more than 300 sites/plots were 94 found throughout the 12 SE Asia countries (Table S1). Only a handful of the studies reviewed reported carbon values, 95 therefore, most of the BGC values we refer to were converted by us from biomass estimates by multiplying by 50% (cf. 96 Smith et al., 2010). Owing to insufficient data, we were not able to separate land-cover classes according to climatic 97 regimes or geographical variables that may have affected plant physiology. Because of a lack of standardization of 98 vegetation classification nomenclature (cf. Maxwell, 2004), a variety of vegetation types were lumped into some common 99 land-cover classes. For example, forests combined both evergreen and deciduous lowland forest types. In addition 100 savannah forests and ambiguous forest types were also placed in this class. We did however separate mangroves and peat 101 forests because of their known high soil organic carbon (SOC) contents. The orchard and tree plantations group 102 included a range of timber and fruit-bearing trees: e.g., Acacia sp., Eucalyptus sp., teak, cocoa, cinnamon, mango, and 103 longan. Because of insufficient data, we also use a general swidden fallow group, as opposed to splitting into short-, 104 medium-, and long-fallow classes (Zieger et al. 2012). The permanent croplands category included a range of crops, 105 including corn, cassava, and rice.

106 Great variation existed within each category, which often lumped together a variety of species, of both young 107 and old ages. In the sub-sections below, we report our strategy for determining ranges of plausible values for below-108 ground biomass and root:shoot ratios. Basically we excluded cases where values result from the sampling of only fine 109 and/or shallow roots, or where very shallow soil depths were sampled. The ranges also exclude outliers associated with 110 very young vegetation that are likely not representative of the mature land-cover class. In a few instances we eliminated 111 outliers that were extreme compared with the rest of the data. Several values were also excluded due to a lack of 112 information on sampling protocols. The final adjusted ranges represent the best-available estimates of below-ground 113 woody carbon and root:shoot ratios for mature facies of each land-cover group, based on empirical research. We do

114 caution that in some site-specific instances the true carbon stock values could be outside our summary ranges. For both 115 variables we report median and mean values that may be useful for preliminary estimates of BGC. These values are not

115 variables we report median and mean values that may be useful for preliminary estimates of BGC. These values are not 116 true mathematical medians and means, as they are determined from the entire population of minimum, maximum, mean,

and median values in adjusted ranges. In addition, we report the midpoints values of each range. We also compute RSRs

118 by examining scatterplots of AGC versus BGC (Figure 2).

119120 <u>Mangrove Forest</u>

121 More below-ground biomass data were available for mangroves than any other land-cover in the region (Table 1): 115 122 values were determined for sites/plots mostly located in Vietnam (55), Thailand (27), and Indonesia (23). Biomass data 123 were reported for a range of species, as well as a range of ages (e.g., from 1-year to mature stands). The BGC values had 124 the highest range of all land covers, as well as the highest values: <1 to 255 Mg C ha-1 (Figure 1a; Table S1). Most of the 125 low values (< 6 Mg C ha⁻¹) were associated with stands < 8 years of age. Some low outliers were determined solely from 126 soil cores. The highest BGC values (> 200 Mg C ha⁻¹) were attributed to the mangroves in Ranong, Thailand. In their 127 assessment, Komiyama et al (1987) considered several root classes (from < 2mm to > 50 mm), and estimated biomass 128 from soil blocks, a root density model determined from excavated trenches and published allometric equations. Several 129 BGC values at these sites are > 110 Mg C ha⁻¹, higher than the maximum BGC values associated in most other locations. 130 Despite the extensive work performed in the biomass calculations at the Ranong site (trench excavations), we considered 131 the highest value as an outlier. Therefore, our adjusted range of BGC values for mangroves is 12-219 Mg C ha-1 (Figure 132 1a). The reported range of RSR was 0.02-5.60 (Table S1). The high value was for a high intertidal zone inhabited by 3 133 year old Ceriops decandra (Griff) Ding Hou. The unusually high RSR was due to the inclusion of dead roots (Alongi and 134 Dixon, 2000). In most places RSR did not exceed 0.55. Our adjusted range of RSRs for this land cover class is 0.11-0.95, for which the median and midpoints are 0.40 and 0.53, respectively (Table 2; Figure 1b). The RSR derived from fitting a 135 136 line through the AGC and BGC data is 0.40 (Figure 2).

137 138 <u>Forest</u>

139 Of the approximately 61 BGC values for forests, most were determined from biomass estimates made in Malaysia (18), 140 southern China (12), Cambodia (10), and Thailand (9). Forest BGC values ranged from 1-90 Mg C ha⁻¹ (Table S1). The 141 highest forest value was associated with a lowland evergreen rainforest in Sabah, Malaysia, for which coarse roots (> 20 142 mm) were sampled (Sim and Nykvist, 1990). The next highest BGC values (74 Mg C ha⁻¹) were based on allometric 143 equations determined from root excavation for two forest types in Xishuangbanna (China) by Zheng et al. (2000; 2006). 144 Across all sites, many values were determined from published allometric equations (i.e., not determined from in situ 145 measurements). Most of the low outliers we excluded in this class were determined in studies that only considered fine 146 roots. In other cases we excluded values for which we could not assess the sampling protocol. Thus, from the range of 147 original values, we identify the adjusted range of BGC for this class to be 11-74 Mg C ha⁻¹ (Figure 1a). The 148 corresponding range for RSRs was 0.08-0.35 (Table 2, Figure 1b). In comparison, summary studies for the tropics report 149 RSR ranges of 0.09 to 0.34 (Cairns et al., 1997; GOFC-GOLD, 2009; Jackson et al., 1996). The median for this range is 150 0.17; and the midpoint is 0.22. From Figure 2, the graph-derived RSR is 0.18.

151 152 <u>Peat forests</u>

153 Peat forest BGC ranged from 3-71 Mg C ha⁻¹, for nine sites/plots in Indonesia and Malaysia (Table S1). A minimum 154 BGC value of 2.8 Mg C ha-1 was determined from small and fine roots via metal coring tubes in 3-m peat deposits in 155 Sumatra (Brady, 1997). The highest BGC value (71 Mg C ha-1) was determined similarly for 12-m peat deposits (Brady, 156 1997). The minimum value at the site was 15 Mg C ha⁻¹, demonstrating great variability. The BGC values associated 157 with the lone Malaysian study in Sarawak allowed for a comparatively high range of values (29-45 Mg h⁻¹), for which the 158 biomass estimates were derived from a published allometric equation (cf. Verwer and van der Meer, 2010; van der Meer 159 and Verwer, 2011). Our adjusted range for this class is 11-71 Mg C ha-1 (Figure 1a). This range is slightly lower than the 160 general forest category, despite these forests having thick organic layers and high soil organic carbon contents (Jaenick et 161 al., 2008; Page et al., 1999; Wibisono et al., 2011; Warren at al., 2012). From the original range of 0.06 to 0.23 we 162 determined an adjusted range of 0.08-0.23. The median of the handful of values in this range is 0.14; and the midpoint is 163 0.16 (Table 2; Figure 1b). Based on the comparison of AGC and BGC, our graph derived estimate of RSR is 0.17. The 164 low value for peat forests could be due to limited research on this land cover.

165

166 Other tree plantations

167 Other tree plantation BGC values range from 1-49 Mg C ha⁻¹ (Table S1). The lowest values were associated with young

- 168 plantations, including *Acacia* sp. and cashew, in Indonesia and Malaysia. Values < 3 Mg C ha⁻¹ were determined with
- established root:shoot ratios for longan and mixed fruit plantations in Vietnam (Zemek, 2009). The highest BGC value
- 170 was determined for mature (17-22 years) teak plantations in northern Thailand (Hiratsuka et al., 2005)—but the site had

a very high RSR of 1.11. Only two other BGC values exceeded 20 Mg C ha⁻¹. They were associated with a coconut cassava plantation (20 Mg ha⁻¹) and a mixed orchard in the Khlong Yai sub-watershed in Thailand (24 Mg C ha⁻¹). For

both, an arbitrary root:shoot ratio of 30% was applied (Gnanavelrajah et al., 2008). Only a handful of the studies

174 performed sampling to measure root biomass *in situ* (e.g., Nykvist et al., 1996; Yamada et al., 2000; Miyakuni et al., 2004;

175 Hiratsuka et al., 2005; Syahrinudin, 2005; Heriansyah et al., 2007). From the original range, we determine an adjusted

176 range of 5-33 Mg C ha⁻¹ (Table 2; Figure 1a). Root:shoot ratios for this class ranged from 0.07 to 1.11 (Table S1). Again,

177 the high value was for a mature teak plantation in Thailand (Hiratsuka et al 2005). Both the second highest value (0.57)

- and the lowest value was determined for a young (4 year) *Acacia* sp. plantation in Sabah, Malaysis (Nykvist et al., 1996).After removing outliers, we derive an adjusted range of 0.11 to 0.39. The high end of this range is associated with 10-
- 175 After femoving outliers, we derive an adjusted range of 0.11 to 0.59. The high end of this range is associated with 10-180 year cashew and 9-year cocoa plantations. The median and midpoint of our adjusted range was 0.21 and 0.25 (Table 2; 181 Σ^{-} 11) Σ^{+} Σ^{-} Σ^{-} 2)
- Figure 1b). The graph-derived value is 0.23 (from Figure 2).

183 Logged-over forest

184 Data existed in seven countries from which the range for BGC could be estimated for logged-over forests (Table 1): 1 to 185 33 Mg C ha-1 (Table S1). The low value was for a disturbed forest in Sulawesi (Leuschner et al., 2009; Leuschner et al., 186 2006; Harteveld et al., 2007); and the high value was associated with an artificial tropical forest in Xishuangbanna, China 187 (Tang et al., 2003). The former study investigated fine roots < 2 mm; the later study excavated the roots of all forest 188 plants. The next highest BGC values, 26 and 22 Mg ha⁻¹, were from a secondary forest in Singapore (Ngo et al., 2013) 189 and a logged dipterocarp forest in Sabah (Pinard and Putz, 1996). Both studies used direct sampling methods. At the low 190 end, BGC values of 4-5 Mg h⁻¹ were associated with secondary forests at least 5-10 years of age in some sites in 191 Xishuangbanna, China (Shi et al, 2001; Tang et al., 1998). Thus, we selected the value of 5 Mg C ha-1 for the low end of 192 our adjusted range of 5-26 Mg C ha-1 (Figure 1a). Most studies including both fine and coarse roots tended to yield 193 RSRs ranging from 0.09 to 0.36; and we adopt this as our adjusted range (Table 2; Figure 1b). The median and midpoint 194 of this range of values are 0.18 and 0.21, respectively. The graph-derived RSR is 0.16 (Figure 2). 195

196 <u>Rubber Plantations</u>

197 Below-ground biomass data for rubber plantations existed for only six sites in Cambodia, China, and Thailand (Table 1). 198 In Thailand, a BGC value of 31 Mg C ha-1 was associated with determinations made from above-ground biomass using 199 an arbitrary root:shoot ratio of 30% (Gnanavelrajah et al., 2008). Through sampling, BGC values ranging from 5-32 Mg 200 C ha⁻¹ were derived for three rubber clones of stand ages 10-50 years (Mizoue, 2009). Cheng et al. (2007) performed 201 field sampling in a 30-year-old rubber stand on Hainan Island (China), leading to BGC estimates of 17 Mg C ha⁻¹. 202 Obtained by excavation of coarse and small roots, the biomass values of Tang et al. (2009) suggest BGC ranges of 7-13 203 Mg C ha-1 and 13-16 Mg C ha-1 for rubber stands 13-19 and 25-47 years, respectively. From these data we identify an 204 adjusted range of 5-32 Mg C ha⁻¹ (Figure 1a). The corresponding adjusted range for RSR is 0.10-0.30, for which the 205 midpoint, median, and graph-derived values are all 0.20. 206

207 Oil Palm Plantations

208 A narrow range of low oil palm BGC values could be estimated for 15 plots/sites in Malaysia and Indonesia (Table S1): 209 2-22 Mg C ha-1. The maximum value was based on field sampling (cores to 5-m depth, excavation of the trunk) in a 30-210 year-old plantation in Sumatra (Syahrinudin, 2005). The lowest values were determined in a 50-cm soil pit in peat soil (> 211 1 m) in Sumatra, for coarse, live and dead roots (Persch et al., 2011). Several values ranging from 3-8 Mg C ha⁻¹ were 212 associated with several plantations 9-16 years of age (Henson and Chai, 1997; Henson and Dolmat, 2003). From these data we identify an adjusted range of 4-22 Mg C ha-1 for mature oil palm (Figure 1a). Reported plantation oil palm RSRs 213 214 ranged from 0.18 to 0.41 (Table S1). Several sites in Malaysia occupied the low end of this range: 0.18-0.19 for 16-23 year-old oil palm in Johor and Perak (Khalid et al., 1999; Henson and Dolmat, 2003). High values ranging from 0.39 to 215 216 0.41 were determined for stands in both Malaysia and Indonesia (Henson and Chai, 1997); Syahrinudin, 2005). Here we 217 use the original range of 0.18 to 0.41 (Table 2; Figure 1b). The median and midpoints are 0.22 and 0.30. Plots of AGC 218 and BGC suggest a RSR of 0.30 (Figure 2).

219

220 <u>Swidden Fallows</u>

221 Swidden fallows BGC values ranged from 3-16 Mg C ha⁻¹ for 10 site/plots in China, Indonesia, and Malaysia (Table S1).

222 Trends were not straightforward in this category, which included young, intermediate, and long fallows. For example,

the lowest value 3 Mg C ha⁻¹ associated with a 12.5-year secondary forest, was similar to that of much younger (1-3 years)

sites in central Kalimantan: BGC = 3 versus 3-4 Mg C ha⁻¹ (Koopmans and Andriesse, 1982 in Kenzo et al., 2010;

- Brearley, 2011). The highest BGC value (16 Mg C ha⁻¹) was for a 6-year bamboo site in west Java (Christanty et al.,
- (1996). We adopt the original minimums and maximums to define the range (Figure 1a): 3-16 Mg C ha⁻¹. The swidden
 fallow root:shoot ratio values ranged from 0.12-1.86 (Table 2). The highest RSR for was elevated by the presence of

bamboo—a grass that is sometimes found in swidden fallows (Christanty et al., 1996; Nikolic et al., 2008; Rerkasem et al.,

2009; Schmidt-Vogt, 2001). Another site in Indonesia with bamboo had a RSR value of 0.69 (Christany et al., 1996). The
 low value in the range (0.12) was associated with the 12.5 year-old secondary forest in Sarawak. We consider a realistic

range for this category is 0.12 to 0.36, unless it contains bamboo and could be greater than one (Table 2; Figure 1b).

The median of our adjusted range is 0.25; the midpoint is 0.24. The plot of AGC versus BGC shows much scatter, but

- the patterns support a value of 0.26. Here we recognize that BGC of certain fallows may in some cases be more
- accurately represented by the range of values for other groups: e.g., GPS (for short fallows), LOF (for long fallows). In
- this case the values could either be higher or lower. We also emphasize again that the presence of bamboo will elevate
- both BGC and the RSR.237

238 Agro-forestry

239 All ten non-swidden agroforestry BGC values originate from Indonesia; and they ranged from 0.04-16 Mg C ha⁻¹ (Table 240 1, S1). The lowest values were associated with the first year of cropping of bamboo talun-kebun agroforestry system 241 (Christanty et al., 1996). The highest (16 Mg C ha⁻¹) was reported for a Javanese home garden, featuring trees mixed 242 annuals and shrubs (Jensen, 1993). Values for other mature land-covers had BGC values of 7-9 Mg C ha⁻¹ (Roshetko et 243 al., 2002; Smiley and Kroschel, 2008). In general there were few land covers with mature trees in this group, thus the 244 high value of our adjusted range of 3-16 Mg C ha⁻¹ may be low (Figure 1a). The derived RSRs for the reviewed studies 245 ranged from 0.01-2.15. Both the lowest and highest values were determined for crops planted along with bamboo in a 246 Talun-Kebun agroforestry system in West Java (Christanty et al., 1996). If we eliminate these extremes, the range of 247 values reported elsewhere was 0.25-0.49 (Table 2; Figure 1b). The corresponding median for this range is 0.36; and the 248 midpoint is 0.37. The plots of AGC and BGC suggest a RSR value of 0.33 (despite one obvious outlier; Figure 2). Like 249 the swidden fallow class, the presence of bamboo would increase BGC and the RSR.

250

251 Grasslands, pastures, and shrublands

252 Grasslands BGC could be derived for five sites/plots in Thailand and Indonesia. Values ranged from 1-4 Mg C ha⁻¹ 253 (Table S1). The highest value was for Imperata grasslands in East Kalimantan in Indonesia (Syahrinudin, 2005). The low 254 value was for shrubs in Sumatra (Solichin et al., 2011); for which, the original biomass values were determined from the 255 allometric equation from Cairns et al. (1997). Values in Thailand (2-3 Mg C ha⁻¹) were associated with unburned, semi-256 natural humid grasslands. These biomass data were determined via soil cores (5 cm diameter) taken down to a depth of 257 only 15 cm (Kamnalrut and Evenson, 1992). These limited data support an adjusted range of 2-4 Mg C ha⁻¹ (Figure 1a), 258 for which the corresponding adjusted RSR range is 0.48-1.92. The median and midpoints for RSR are 1.11 and 1.20. 259 The graph-derived estimate of the RSR is 0.78. We note that the RSR could be either very low or very high for this 260 diverse land-cover category, depending on the species composition of the land cover-however, too few data exist to 261 make an accurate assessment.

262263 <u>Permanent cropland</u>

264 Eight values of BGC for permanent crops could be derived from two locations in Thailand and one in Vietnam (range = 265 1-5 Mg C ha⁻¹; Table S1). Of these, only the value (2 Mg ha⁻¹) from a mixed agriculture site in northern Thailand was 266 based on in situ sampling (Pibumrung et al., 2008). While this site included a mixture of rice paddy, corn fields, fallows, 267 and orchards, we have chosen to include them in this vegetation class, rather than AGF. The lone site in Vietnam had a 268 BGC value of 1 for a banana plantation (Zemek, 2009). Most of the BGC values of the PC class were calculated for the 269 Khlong Yai subwatershed site in Thailand from an arbitrary root:shoot ratio of 30% by Gnanavelrajah et al., (2008). The 270 highest value (5 Mg C ha-1) was associated with sugar cane. These limited data support a BGC range of 1-5 Mg C ha-1 271 (Figure 1a). Again, seven of the eight RSR values associated with permanent agriculture (0.26-0.31) were literature values 272 used to calculate below-ground biomass. The only value determined by sampling (0.31) was associated with a mixed 273 agriculture site in northern Thailand (Pibumrung et al., 2008). Thus, we use the original range of 0.26-0.31 for the RSRs 274 in this class. The median is 0.30; the midpoint, 0.29. The plots in Figure 2 suggest a value of about 0.30, but this is an 275 artifact of nearly all the AGC and BGC values being derived from RSRs, not field sampling.

276

277 3. Data limitations and uncertainty

We considered all reported data in our effort to develop plausible ranges of below-ground carbon and root:shoot ratios, despite potential flaws and differences in collection methods. In order to increase the pool of available BGC data, we

- have included studies that were undertaken for a variety of purposes, not solely biomass estimations. Adjusting the
- ranges by eliminating distinct outliers, in part, addresses the issue of underestimating total root biomass by some studies.
- Admittedly, the minimum values for many of the ranges still do appear too low for mature stands. If this is the case,
- using the medians, means, or range midpoints as a typical value for a particular class would also be low. The high
- 284 frequency of use of general allometric equations-including root:shoot ratios-is of concern for generating this

summary because these data may not be truly representative of mature vegetation at the particular study site.
Amalgamating land covers that contain myriad different plant species into the various operational categories (e.g., forests, OTP) also created uncertainty. However, a more important source of uncertainty was the non-standardization of sampling methods. Here we discuss some of the limitations we encountered during the review.

289 The methods employed in any one case often depended on geographical variables affecting accessibility, as well as the 290 type of vegetation considered. In general, soil cores were used for determining fine root biomass; and published 291 allometric equations were often used to determine total root biomass (fine + coarse roots). The most popular method 292 applied in forests was allometric relationships, followed by soil cores, soil pits and root excavation. Similarly, BGC from 293 orchards, tree and rubber plantations were mostly derived from indirect methods, whereas biomass data from logged 294 forest, oil palm, agroforest, peat forest, swidden fallows and grasslands were determined largely from direct methods 295 such as cores and pits. The few available permanent crop BGC data were determined from indirect methods; all but one 296 was determined by root:shoot ratios. Mangrove values were determined from a fairly equal mix of direct and indirect 297 methods. Again, of concern was the dependence on pre-existing allometric relationships rather than the determination 298 of new site-specific ones (eg. Komiyama et al., 2005). Granted, it is extremely difficult to perform destructive sampling 299 to make these determinations, but the point we are making is that failing to do so introduces uncertainty in the 300 determination of below ground biomass and the associated carbon.

301 Soil coring was a popular method that provided estimates for small localized points (Komiyama et al., 2000; 302 Oliveira et al., 2000). A general limitation with coring is that sampling of coarse roots with small cores is practically 303 impossible. It is also difficult to obtain samples near the base of the tree where root density may be highest. Soil 304 compaction during coring may also skew results (cf. Makkonen and Helmisaari, 1999; Park et al., 2007). For both coring 305 and excavation methods, samples are also difficult to extract in wet, sandy soils or stony soils. Total root excavation is 306 the best method for measuring large and deep vertical roots; however, very deep tap roots extending down many meters 307 and/or anchored into bedrock are often not sampled sufficiently. Nevertheless, roots will inevitably be lost during the 308 excavation process due to accidental breakage. The great need for manpower and/or machinery to facilitate root 309 removal is likely one of the reasons that only about 10% of the data were derived from root extraction methods.

310 Studies using multiple methods arguably provide more accurate biomass estimates. Pinard and Putz (1996), for 311 example, used soil monoliths (pits) to study coarse roots (>5 mm diameter) and soil cores to measure fine roots (<5 mm 312 diameter) in logged over forests in Sabah, Malaysia. Sim and Nykvist (1990) combined results from two different 313 methods to derive a single carbon biomass value for coarse roots > 20 mm in a lowland evergreen forest. Specifically, roots were cut 3 cm from a central stump for trees with a DBH > 19 cm. For smaller trees, $50 \times 50 \times 50$ cm sample pits 314 315 were excavated to study the root systems in detail. Elsewhere, Nguyen (2009) collected soil cores and soil blocks to 316 quantify belowground root carbon in mangroves. With these exceptions, most studies use only one direct method (i.e. 317 cores, pits or root excavation) to quantify coarse or fine root biomass, thereby failing to counterbalance the 318 shortcomings of any one method (cf. Hertel et al., 2009; Kitayama and Aiba, 2002; Hendricks et al., 2006; Leuscher et al., 319 2009). The importance of choosing the one method over another was demonstrated by Park et al. (2007), who found 320 that fine root biomass determined from soil cores was 27% higher than that determined from soil pits.

321 Root biomass estimates in the reviewed studies included either or both fine and coarse roots, but there was no 322 standardization for separating the two. In general, two-to-five millimeters was commonly used, although it varied with 323 vegetation type (Kenzo et al., 2010; Pinard and Putz, 1996; Stokes et al., 2009; Zobel, 2005). Arguably, roots should be 324 defined based on functionality, but the lack of knowledge of root ontogeny and morphology often prevents this (Pierret 325 et al., 2005). Some studies defined root vitality classes by either visual or physical criteria, including color, tensile strength, 326 flexibility, and chemistry. However, classification of roots into live, dead, or unknown classes was inconsistent. Lastly, 327 the definition of what constituted root biomass varied, with some researchers combining shrub and herb root biomass 328 together with that of trees. Others calculated only the biomass of the primary species (e.g., trees in a forest association). 329 Again, we point out that the motivation of the reviewed studies was not always for biomass determinations.

330 Although, rarely mentioned, processing errors may produce underestimates when roots are sampled directly. 331 Root breakage and loss-particularly of fine roots-inevitably occurs during sampling or washing (Clark et al., 2001; 332 Subedi et al., 2006). The amount of care and time spent to extract roots from soils is often a key determinant in 333 influencing root recoveries (Pierret et al., 2005). Also, there was no standardized or best way to recover roots from soils 334 - some studies separated the roots by hand (eg. Ngo et al., 2013) while others washed roots through various size sieves 335 (range: 65 µm to 5 mm) (cf. Syahrinudin, 2005; Pibumrung et al., 2008). Floatation was also suggested (Oliveira et al., 336 2000). For the floatation method, small roots may remain stuck to clayey soil particles despite repeated washing. Lastly, 337 excess lag time between root storage and processing may result in root death or loss of dry weight via decomposition, 338 thereby affecting the root biomass estimates (Oliveira et al., 2000).

Many of the case studies we reviewed arguably sampled too small an area to capture the spatial resolution of
 BGC in the stand. For example, the number of replicates in the reviewed studies for forest ranged from 2 cores within a
 30-60 ha plot (Pinard and Putz, 1996) to 160 cores within one 8-ha landform unit (Pibumrung et al., 2008). Figure 3a

342 shows the great variability in root biomass at depths down to 1 m for four replicate cores collected in a secondary forest. 343 In addition, root biomass estimated within the upper 2 m of two duplicate soil pits in the same secondary forest differed 344 by almost three-fold (7 versus 18 Mg C ha⁻¹; Figure 3b). These examples demonstrate the need for sufficient sample 345 replicates to capture the spatial variability. With respect to temporal variability, most of the reviewed studies were one-346 time snapshots, without replication. Thus, seasonal, environmental and age-related variations in root biomass were not 347 captured. In addition, few of the studies sampled below one meter, despite evidence that rooting depths of tropical trees, 348 shrubs and herbaceous plants can reach or exceed depths of 7.0, 5.0 and 2.5 m respectively (Canadell et al., 2006). The 349 importance of deep sampling is demonstrated in Figure 3b: 10-20% of the total root biomass, mostly fine roots, 350 occurred below the depth of 1 m in bamboo and secondary evergreen forests (Figure 3).

Finally, using different means of deriving carbon fractions adds to uncertainty. Most biomass carbon values
were calculated as one-half measured root biomass. In cases where carbon content was determined analytically (e.g, with
carbon-nitrogen analyzer or Walkley-Black method) values ranging from 0.37-0.53 have been determined (cf. Brearley,
2011; Hertel et al., 2009, 2009b; Kenzo et al., 2010; Leuschner et al., 2009; Pimbumrung et al., 2008).

355 These limitations, which were not always clearly elaborated upon in the reviewed case studies, contribute 356 uncertainty in the BGC and the RSRs we identify for each land cover. Specifically, the sampling related 357 limitations/difficulties complicate the development of accurate allometric relationships for estimating BGC from AGC. 358 Caution is needed when using generalized relationships because they may not be representative of the characteristics of 359 any one specific study plot, especially in heterogenous facies (Chave et al., 2005; Komiyama et al., 2008). Rarely can the 360 wealth of below-ground biomass for all vegetation be calculated from tree-based allometric relationships. In comparsion, 361 the use of indirect methods in homogeneous landscapes, such as tree-based rubber and oil palm plantations, may be 362 more reliable. However, one must also consider that allometric relationships may change with the age of vegetation and 363 changing resource availability (Hütsch et al., 2002; Laclau, 2003; Shipley and Meziane, 2002; van Noordwijk et al., 2004; 364 Wilson, 1998). 365

366 4. Total vegetation carbon stocks

The ranges of BGC we have determined for some land covers are slightly different from those we reported in the prior meta-analysis of total ecosystem carbon (Ziegler et al., 2012), particularly, those for grasslands and non-swidden agroforests categories. Again, partial motivation of this review was to improve upon our prior assessment of carbon stock changes related to land-cover conversion. In our new estimation of TEC changes (Table 3), minimum and maximum values of AGC for most land covers were adopted from the prior assessment (Ziegler et al., 2012). The AGC values for mangrove and peat forests were determined from new data presented in Table S1, for which obvious outliers were removed (as per Ziegler et al., 2012).

374 The mangrove and peat forest calculations were complicated because soil profiles for these vegetation types 375 can have very high organic contents extending down several meters. We estimate the range of SOC for mangrove 376 forests and peat forest as 225-675 Mg C ha-1 and 537-1612 Mg C ha-1, respectively. These values are estimates for a soil 377 profile 2 m deep, in agreement with the idealized soil profiles for which the other land-cover SOC values are estimated 378 (described in detail in Ziegler et al., 2012). The mangrove value is estimated as 3-fold that of forest-an assumption 379 based on the data of Donato et al. (2012) showing mangrove SOC ranging from 13-15%, compared with values for 380 upland soils that ranging from 0.4-5.5%. For the peat forest, the maximum value of 1612 Mg C ha⁻¹ (for 2 m) is within 381 the range of 1425 Mg C ha-1 (2.67 m depth) to 7889 Mg C ha-1 (12.07 m depth) reported by Warren et al (2012). It is 382 estimated by a linear regression equation, determined by fitting the data in that study: 412.85 * 2 (m depth) + 786.41 (R² 383 = 0.85; n =10). The minimum peat forest SOC value is about 2.4-fold that of the mangrove minimum (based on ratio 384 of peat maximum to mangrove maximum)

385 The rationale of these new TEC estimates is to explore the plausible impacts of changes from one land cover 386 to another in the region. Such comparisons are often the starting points in any proposal related to REDD+ or other 387 carbon accounting endeavors. Ideally such studies would collect site-specific data, but this is not always the case. In 388 absence of other more detailed data, it is evident from the values in Table 3 that conversion among several land covers 389 could result in ambiguous outcomes in carbon stocks. The improved estimates of BGC give us more confidence in 390 claiming that many projected land-cover/land-use transitions would produce uncertain or potentially neutral carbon 391 outcomes: e.g. (a) transitions between short-fallow swidden systems and permanent croplands; (b) land-cover changes 392 between/among long-fallow swidden, other agroforestry systems, and possibly rubber; and (c) land-cover changes 393 between/among intermediate-fallow swiddening, grasslands, pastures, shrub lands, and oil palm plantations. These 394 uncertainties are important to stress because many of these transitions are currently at the heart of REDD+ debates (cf. 395 Ziegler et al. 2012). Also apparent in the estimates is the importance of forests, particularly peat and mangrove forests, 396 as carbon sinks (cf. Donato et al., 2012). Conversions of these land covers almost invariably results in losses of carbon 397 stocks.

399 5. Towards a better carbon database

400 Our analysis revealed that careful attention should be given to sampling to appropriate depths, obtaining sufficient 401 replicates, and using appropriate sampling intervals to capture accurately heterogeneous root distribution, both vertically 402 and laterally (cf. Moore and McCabe, 1999). Detailed information should also be presented to provide clarity and aid in 403 interpretations. When applicable, roots should not be amalgamated into a single category, but partitioned into live and 404 dead roots. Information on the distribution of root lengths and depths for species should also be recorded. Where 405 indirect methods are to be used, information should be validated against direct methods as a form of quality control. In 406 most cases, multiple methods should be used to substantiate common calculations and provide the most accurate 407 calculation for particular root types (e.g., coarse versus fine). Long-term sampling to account for temporal variability in 408 root dynamics at a given site should be considered. In general, time-average carbon stocks are preferential for comparing 409 differences between land uses. These observations are in general agreement with published outlines of appropriate 410 sampling protocols for determining BGC accurately (Vogt et al., 1998; Oliveira et al., 2000; Qureshi et al., 2012).

411 In most cases, the use of general allometric equations or RSRs is not as desirable as using one determined 412 specifically for a location (cf. Chave et al., 2005; Kauffman and Donato, 2012). An important shortcoming of deriving such equations is that below-ground biomass components must be determined from direct sampling using destructive 413 414 methods (Chave et al., 2004; Levillain et al., 2011; Mokany et al., 2006). Destructive sampling in forests, logging 415 concessions, orchards, and plantations, however, is often not possible. In the latter three cases, root biomass 416 determinations can be conducted between rotations. This would require researchers to develop working relationships 417 with loggers and plantation owners (Laclau, 2003). Research in natural forests is complicated because most now fall 418 under the protection or jurisdiction of government forest and/or conservation departments. Government conservation 419 agencies should therefore be drawn into the forest carbon documentation process, whereby opportunistic events can be 420 used to bolster the forest carbon inventory. One means of generating new data on protected trees is to align research 421 with planned construction projects. In most countries road network, power line, and rural expansion requires the 422 removal of trees for which limited or no BGC data exist. Allowing researchers to extract trees prior to removal would 423 provide an avenue to bolster the forest carbon database. Likewise, partnerships with the mining industry to allow 424 sampling of deep rooted vegetation in open-cut mines would be useful as well. 425

426 6. Conclusions

427 Uncertainty in below-ground root carbon stocks for eleven major land covers in twelve countries across SE Asia results 428 in part from the limited amount of research that has been conducted to date and methodological inconsistencies 429 between existing studies. Limited data exists for rubber and oil palm despite their importance as cash-generating crops in 430 the region (Edwards et al., 2012; Fox et al., 2012; Koh and Wilcove, 2008; Ziegler et al., 2009; Ziegler et al., 2012). 431 Furthermore, the paucity of data for swidden fallows is surprising given the historical criminalization of slash-and-burn 432 agriculture and the uncertain role of this land cover in the future (cf. Mertz, 2009; Ziegler et al., 2011, 2012). The paucity 433 of data strengthens our claim that great uncertainty exists with regard to carbon outcomes of transitions between many 434 land covers. One implication of this uncertainty for policy-making is that reliable estimates of how any land-use changes 435 will affect below-ground root carbon simply cannot be made with existing (published) above- or below-ground meta-436 analysis data (Ziegler et al., 2012). Some land-cover transitions are less affected by uncertainty: e.g., transitions to/from 437 mature forests and other tree-based plantations to low biomass crops. However, the outcomes of changes in below-438 ground carbon regarding many non-forest changes are ambiguous. Furthermore, the time scale over which the BGC is 439 lost to the atmosphere is very poorly understood. Root biomass may persist for decades, but breakdown rates and the 440 fate of the root carbon are rarely studied.

441 New carbon stock assessment programs must include complementary site-specific, direct measurements of 442 both above- and below-ground carbon. Unfortunately, new BGC measurements will require using destructive methods 443 that quantify all biomass components (live or dead, coarse or fine) over time and space. Reliance on allometric 444 relationships already determined for the region may not provide reliable estimates, although the inclusion of data 445 determined outside SE Asia may be useful. Government agencies, private enterprises, and development agencies could 446 play a role in developing a better forest carbon database by teaming with researchers to assess AGC and BGC prior to 447 trees being removed for construction (road, dam, power lines), mining, or stand rotations (forestry, plantations). Achieving greater certainty in terrestrial carbon stock, while challenging, will allow improved assessments of stock losses 448 449 associated with the rapid landscape changes now taking place in the region. This is particularly true at forest frontier 450 areas where rapid conversion from traditional land covers to high value plantations (oil palm, rubber) is occurring at 451 unprecedented rates (Ziegler et al., 2009; Fox et al., 2012).

452

Acknowledgements: Support was provided by NASA grants NNG04GH59G and NNX08AL90G, APN grant
 ARCP2008-01CMY, and NUS grant R-109-000-134-112. This research is part of the project entitled Impacts of
 Reducing Emissions from Deforestation and Forest Degradation and Enhancing Carbon Stocks (I-REDD+). I-

- 457 Reducing Emissions from Derofestation and Porest Degradation and Emianeing Carbon Stocks (P-REDD+). 1-458 REDD+ is funded by the European Community's Seventh Framework Research Programme. More information can be
- 459 found on the web site: <u>http://www.i-redd.eu</u>.

460461 References

- Abiven, S., Recous, S., Reyes, V., Oliver, R., 2005. Mineralisation of C and N from root, stem and leaf residues in soil
 and role of their biochemical quality. *Biology and Fertility of Soils 42*, 119-128.
- 464 Albrecht, A., Cadisch, G., Blanchart, E., Sitompul, S.M., Vanlauwe, B., 2004. Below-ground inputs: Relationships with
- soil quality, soil C storage and soil structure, in: van Noordwijk, M., Cadisch, G., Ong, C.K. (Eds.), *Below-ground interactions in tropical agroecosystems: Concepts and models with multiple plant components.* CABI Publishing, Oxfordshire, pp.83-107.
- Alongi, D.M., Dixon, P., 2000. Mangrove primary production an above- and below-ground biomass in Sawi Bay,
 Southern Thailand. Phuket Marine Biological Center Special Publication 22, 31-38.
- 469

482

- Ballesteros, A., Polycarp, C., Stasio, K., Chessin, E., Easton, C., 2011. Summary of developed country fast-start climate finance pledges. Available at: <u>http://pdf.wri.org/climate_finance_pledges_2010-11-24.pdf</u> (accessed 17 July 2012).
- Brady, M.A., 1997. Organic matter dynamics of coastal peat deposits in Sumatra. Unpubl. PhD thesis. University of British
 Columbia, Vancouver.
- Brearley, F.Q., 2011. Below-ground secondary succession in tropical forests of Borneo. *Journal of Tropical Ecology 27*, 413-420.
 477
- Brown, S., 2002. Measuring carbon in forests: current status and future challenges. *Environmental Pollution 16*, 363-372.
- 480 Cairns, M.A., Brown, S., Helmer, E.H., Baumgardner, G.A., 1997. Root biomass allocation in the world's upland forests.
 481 Oecologia 111, 1-11.
- 483 Canadell, J., Jackson, R.B., Ehleringer, J.R., Mooney, H.A., Sala, O.E., Schulze, E.-D., 1996. Maximum rooting depth of
 484 vegetation types at the global scale. *Oecologia 108*, 583-595.
 485
- 486 Chalermchatwilai, B., Poungparn, S., Patanaponpaiboon, P., 2011. Distribution of fine-root necromass in a secondary
 487 mangrove forest in Trat province, Eastern Thailand. *ScienceAsia 37*, 1-5.
 488
- Chave, J., Condit, R., Aguilar, S., Hernandez, A., Lao, S., Perez, R., 2004. Error propagation and scaling for tropical forest biomass estimates. *Philosophical Transactions of the Royal Society of Biological Sciences 359*, 409-420.
- Chave, J., Andalo, C., Brown, S., Cairns, M.A., Chambers, J.Q., Eamus, D., Fölster, H., Fromard, F., HiguchI, N., Kira,
 T., Lescure, J.-P., Nelson, B.W., Ogawa, H., Puig, H., Riéra, B., Yamakura, T., 2005. Tree allometry and improved
 estimation of carbon stocks and balance in tropical forests. *Ecosystem Ecology* 145, 87-99.
- Cheng C.-M., Wang R.-S., Jiang J.-S., 2007. Variation of soil fertility and carbon sequestration by planting Hevea
 brasiliensis in Hainan Island, China. *Journal of Environmental Sciences 19*, 348-352.
- Christanty, L., Mailly, D., Kimmins, J.P., 1996. "Without bamboo, the land dies": Biomass, litterfall and soil organic
 matter dynamics of a Javanese bamboo talun-kebun system. *Forest Ecology and Management 87*, 75-88.
- 502 CIFOR, 2011. Global Database of REDD+ and Other Forest Carbon Projects. Available at:
- 503 <u>http://www.forestsclimatechange.org/redd-map/</u> (accessed 1 October 2011).
 504
- Clark, D.A., Brown, S., Kicklighter, D.W., Chambers, J.Q., Thomlinson, J. R., Jian, N., 2001. Measuring net primary
 production in forests: concepts and field methods. *Ecological Applications 11(2)*, 356-370.

- 507 Donato, D.C., Kauffman, J.B., Mackenzie, R.A., Ainsworth, A., Pfleeger, A.Z., 2012. Whole-island carbon stocks in the
 508 tropical Pacific: Implications for mangrove conservation and upland restoration. *Journal of Environmental Management 97*,
 509 89-96.
- 510

522

528

533

511 Edwards, D.P., Koh, L.P., Laurance, W.F., 2012. Indonesia's REDD+ pact: Saving imperilled forests or business as
512 usual? *Biological Conservation 151*, 41-44.
513

- 514 FCPF, 2011. Forest Carbon Partnership Facility REDD+ Country Participants Readiness Proposals. Available
 515 at: <u>http://www.forestcarbonpartnership.org/fcp/node/203</u> (accessed 1 October 2011).
 516
- Fox, F.M., Vogler, J.B., Sen, O.L., Giambelluca, T.W., Ziegler, A.D., 2012. Simulating land-cover change in Montane
 Mainland Southeast Asia. Environmental Management 49(5), 968-979.
- Gibbs, H.K., Brown, S., Niles, J.O., Foley, J.A., 2007. Monitoring and estimating tropical forest carbon stock: making
 REDD a reality. *Environmental Research Letters 2*, 1-13.
- 523 Gnanavelrajah, N., Shrestha, R. P., Schmidt-Vogt, D., Samarakoon, L., 2008. Carbon stock assessment and soil carbon 524 management in agricultural land-uses in Thailand. *Land Degradation and Development 19*, 242-256.
- 525 GOFC-GOLD, 2009. Reducing greenhouse gas emissions from deforestation and degradation in developing countries: a sourcebook of
- 526 methods and procedures for monitoring, measuring and reporting, GOFC-GOLD Report Version COP 14-2. GOFC-GOLD Project
- 527 Office, Natural Resources Canada, Alberta.
- Harteveld, M., Hertel, D., Wiens, M., Leuschner, C., 2007. Spatial and temporal variability of fine roots abundance and
 growth in tropical moist forests and agroforestry systems Sulawesi, Indonesia. *Ecotropica 13*, 111-120.
- Hendricks, J.J., Hendrick, R.L., Wilson, C.A., Mitchell, R.J., Pecot, S.D., Guo, D., 2006. Assessing the patterns and controls of fine root dynamics: an emphirical test and methodological review. *Journal of Ecology 94*, 40-57.
- Henson, I.E., Chai, S.H., 1997. Analysis of oil palm productivity. II. Biomass, distribution, productivity and turnover of
 the root system. *Elaeis Vol. 9, No. 2, December 1997*, 78-92.
- Henson, I.E., Dolmat, M.T., 2003. Physiological analysis of an oil palm density trial on a peat soil. *Journal of Oil Palm Research Vol. 15, No. 2, December 2003*, 1-27.
- 538 Heriansyah, I., Miyakuni, K., Kato, T., Kiyono, Y., Kanazawa, Y., 2007. Growth characteristics and biomass
- accumulations of acacia mangium under different management practices in Indonesia. *Journal of Tropical Forest Science 194*,
 226-235.
- Hertel, D., Harteveld, M.A., Leuschner, C., 2009. Conversion of a tropical forest into agroforest alters the fine root-related carbon flux to the soil. *Soil Biology and Biochemistry 41*, 481-490.
- Hiratsuka, M., Chingchai, V., Kantinan, P., Sirirat, J., Sato, A., Nakayama, Y., Matsunami, C., Osumi, Y., Morikawa, Y.,
 2005. Tree biomass and soil carbon in 17- and 22 year old stands of teak Tectona grandis L.f. in northern Thailand. *Tropics Vol.* 144, 377-382.
- 547
- 548 Hütsch, B.W., Augustin, J., Merbach, W., 2002. Plant rhizodeposition an important source for carbon turnover in soils.
 549 *Journal of Plant Nutrition and Soil Science 165*, 397-407.
- 550
- 551 Jackson, R.B., Canadell, J., Ehleringer, J.R., Mooney, H.A., Sala, O.E., Schulze, E.D., 1996. A global analysis of root
- distributions for terrestrial biomes. Oecologia 108, 389-411.
- 553

- Jackson, R.B., Mooney, H.A., Schulze, E.-D., 1997. A global budget for fine root biomass, surface area and nutrient contents. *Proceedings of the National Academy of Sciences of the United States of America 94*, 7362-7366.
- Jaenicke, J., Rieley, J.O., Mott, C., Kimman, P., Siegert, F., 2008. Determination of the amount of carbon stored in
 Indonesian peatlands. *Geoderma 147*, 151-158.
- Jensen, M.,1993. Soil conditions, vegetation structure and biomass of a Javanese home garden. *Agroforestry Systems 24*, 171-186.
- Kamnalrut, A., Evenson, J.P., 1992. Monsoon grassland in Thailand, in: Long, S.P., Jones, M.B., Roberts, M.J. (Eds.),
 Primary productivity of grass ecosystems of the tropics and sub-tropics. Chapman Hall, London, pp. 100-126.
- Kauffman, J.B., Donato, D.C., 2012. Protocols for the measurement, monitoring and reporting of structure, biomass and carbon stocks in
 mangrove forests. Working paper 86. CIFOR, Bogor.
- Kenzo, T., Ichie, T., Hattori, D., Kendawang, J.J., Sakurai, K., Ninomiya, I., 2010. Changes in above- and belowground biomass in early successional tropical secondary forests after shifting cultivation in Sarawak, Malaysia. *Forest Ecology and Management 260*, 875-882.

589

- 571 Khalid, H., Zin, Z.Z., Anderson, J.M., 1999. Quantification of oil palm biomass and nutrient value in a mature
 572 plantation. II. Below-ground biomass. *Journal of Oil Palm Research 11, No. 2,* 63-71.
 573
- 574 Kitayama, K., Aiba, S.-I., 2002. Ecosystem structure and productivity of tropical rain forests along altitudinal gradiens
 575 with contrasting soil phosphorus pools on Mount Kinabalu, Borneo. *Journal of Ecology 90*, 37-151.
 576
- 577 Koh, L.P., Wilcove, D.S., 2008. Is oil palm agriculture really destroying tropical biodiversity? Conservation letters 1, 60578 64.
 579
- 580 Komiyama, A., Havanond, S., Srisawatt, W., Mochida, Y., Fujimoto, K., Ohnishi, T., Ishihara, S., Miyaki, T., 2000. Top/
 root biomass ratio of a secondary mangrove (Ceriops tagal (Perr.) C.B. Rob.) forest. *Forest Ecology and Management 139*, 127-134.
 583
- 584 Komiyama, A., Ogino, K., Aksornkoae, S., Sabhasri, S., 1987. Root biomass of a mangrove forest in southern Thailand.
 585 1. Estimation by the trench method and the zonal structure of root biomass. *Journal of Tropical Ecology 3*, 97-108.
 586
- 587 Komiyama, A., Ong, J.E., Poungparn, S., 2008. Allometry, biomass and productivity of mangrove forests: A review.
 588 *Aquatic Botany 89*, 128-137.
- Komiyama, A., Poungparn, S., Kato, S., 2005. Common allometric equations for estimating the tree weight of mangroves. Journal of Tropical Ecology 21, 471-477.
- Koopmans, T.T., Andriesse, J.P., 1982. Baseline study monitoring project of nutrient cycling in shifting cultivation. Department of
 Agricultural Research Internal Report BO 82-6. Koninklijk Instituut voor de Tropen, Amsterdam.
- Körner, C., 1994. Biomass fractionation in plants: a reconsideration of definitions based on plant functions, in: Roy, J.,
 Garnier, E. (Eds.), *A whole plant perspective on carbon-nitrogen interactions*. Academic Publishing, The Hague, pp. 173-185.
- Laclau, P., 2003. Root biomass and carbon storage of ponderosa pine in a northwest Patagonia plantation. *Forest Ecology and Management 173*, 353-360.
- Leuschner, C., Harteveld, M., Hertel, D., 2009. Consequences of increasing forest use intensity for biomass, morphology
 and growth of fine roots in a tropical moist forest on Sulawesi, Indonesia. *Agriculture, Ecosystems and Environment 129*, 474 481.
- Leuschner, C., Wiens, M., Harteveld, M., Hertel, D., Tjitrosemito, S., 2006. Patterns of fine root mass and distribution
 along a disturbance gradient in a tropical montane forest, Central Sulawesi Indonesia. *Plant and Soil 283*, 163-174.

- Levillain, J., M'bou, A.T., Deleporte, P., SAint-André, L., Jourdan, C., 2011. Is the simple auger coring method reliable
 for below-ground standing biomass estimation in Eucalyptus plantations. *Annals of Botany 108*, 221-230.
- Makkonen, K., Helmisaari, H.-S., 1999. Assessing fine-root biomass ad production in a Scots pine stand comparison of
 soil core and root ingrowth methods. *Plant and Soil 210*, 43-50.
- 612

Maxwell, J.F., 2004. A synopsis of the vegetation of Thailand. *The Natural History Journal of Chulalongkorn University* 4, 1929.

- Mertz, O., 2009. Trends in shifting cultivation and the REDD mechanism. Current Opinion in Environmental
 Sustainability 1, 156-160.
- 618619 Miyakuni, K., Heriansyah, I., Heriyanto, N.M., Kiyono, Y., 2004. Allometric biomass equations, biomass expansion
- factors and root-to-shoot ratios of planted Acacia mangium Willd. forests in West Java, Indonesia. *Journal of Forest Dlaming* 10 (0.76)
- **621** *Planning 10*, 69-76.
- Mizoue, N., Kakada, K., Muthavy, P., Gyokusen, K., Koga, S., Shigematsu, A., Yoshida, S., 2009. Multiple functions of
 rubber plantations as forest and wood resources project progress report. 120th Forest Congress in Kyoto, Session T21:
 Sustainable production from manmade forests in the tropics. 28 March, 2009. Kyoto.
- Mokany, K., Raison, R.J., Prokushkin, A.S., 2006. Critical Analysis of root:shoot ratios in terrestrial biomes. *Global Change Biology 12*, 84-96.
- Moore, D., Mccabe, G.P., 1999. *Introduction to the practice of statistics. 3rd edition.* W H Freeman and Co., New York.
- Ngo, K.M., Turner, B.L., Muller-Landau, H.C., Davis, S.J., Larjavaara, M., Hassan, N.F.B.N., Lum, S., 2013. Carbon
 stocks in primary and secondary tropical forests in Singapore. *Forest Ecology and Management 296*, 81-89.
- Nguyen, T.K., Ninomiya, I., Nguyen, T.L., Nguyen, H.T., Tuan, M.S., Hong, P.N., 2009. Belowground carbon
 accumulation in young Kandelia candel (L.) Blanco plantations in Thai Binh River Mouth, Northern Vietnam. *International Journal of Ecology and Development 12(9)*, 107-117.
- Nikolic, N., Schultze-Kraft, R., Nicolic, M., Böcker, R., Holz, I., 2008. Land degradation on barren hills: a case study in northeast Vietnam. *Environnemental Management* 42, 19-36.
- 640 Nykvist, N., Sim, B.L., 1996. Effects of tractor logging and burning on biomass production and nutrient accumulation
 641 in Acacia Mangium plantations in Sabah, Malaysia. *Journal of Tropical Forest Science 92*, 161-183.
- 642 Oliveira, M.D., van NoordwijK, M., Gaze, S.R., Brouwer, G., Bona, S., Mosca, G., Hairiah, K., 2000. Auger sampling,
- 643 ingrowth cores and pinboard methods, in: Smit, A.L., Bengough, A.G., Engels, C., van Noordwijk, M., Pellerin, S., van de Geijn, S.C. (Eds.), *Root methods: A handbook*. Springer-Verlag, Berlin, pp. 175-193.
 645
- Page, S.E., Rieley, J.O., Shotyk, W., Weiss, D., 1999. Interdependence of peat and vegetation in a tropical peat swamp forest. *Proceedings of the Royal Society B 354*, 1-13.
- Park, B.B., Yanai, R.D., Vadeboncoeur, M.A., Hamburg, S.P., 2007. Estimating root biomass in rocky soils using pits, cores and allometric equations. *Soil Science Society of America Journal* 71, 206-213.
- 651

636

- Persch, S., Hergoualc'h, K., Laumonier, Y., Verchot, L., 2011. Carbon stock in coarse root biomass in different land-use
 systems on tropical peat. Workshop on Tropical Wetland Ecosystems of Indonesia: Science needs to address climate change adaptation
- 654 and mitigation. Watilan Convention Center, Sanur Beach Hotel, Bali.
- Pibumrung, P., Gajaseni, N., Popan, A., 2008. Profiles of carbon stocks in forest, reforestation and agricultural land,
 northern Thailand. *Journal of Forestry Research 19(1)*, 11-18.
- 657

- Pierret, A., Moran, C.J., Doussan, C., 2005. Conventional detection methodology is limiting our ability to understand the
 roles and functions of fine roots. *New Phytologist 166*, 967-980.
- 660 Pinard, M.A., Putz, F.E., 1996. Retaining forest biomass by reducing logging damage. *Biotropica 28(3)*, 278-295.
- 662 Quéré, C.L., Raupach, M.R., Canadell, J.G., Marland, G., et al., 2009. Trends in the sources and sinks of carbon dioxide.
- 663 *Nature Geoscience advance online publication* 1-6.
- 664 Qureshi, A., Pariva, Badola, R., Hussain, S.A., 2012. A review of protocols used for assessment of carbon stock in
 665 forested landscape. *Environmental Science and Policy 16*, 81-89.
 666
- Rasse, D.P., Rumpel, C., Dignac, M.-F., 2005. Is soil carbon mostly root carbon? Mechanisms for specific stabilisation. *Plant and Soil 269*, 341-356.
- 669

681

690

692

704

- 670 Rerkasem, K., Lawrence, D., Padoch, C., Schmidt-Vogt, D., Ziegler, A.D., 2009. Consequences of swidden transitions
 671 for crop and fallow biodiversity. *Human Ecology* 37, 347-360.
 672
- 673 Roshetko, J.M., Delaney, M., Hairiah, K., Purnomosidhi, P., 2002. Carbon stocks in Indonesian homegarden systems:
 674 Can smallholder systems be targeted for increased carbon storage? *American Journal of Alternative Agriculture 172*, 138-148.
- 675 Sanaullah, M., Chabbi, A., Leifeld, J., Bardoux, G., Billou, D., Rumpel, C., 2011. Decomposition and stabilization of root
 676 litter in top- and subsoil horizons: what is the difference? *Plant Soil 338*, 127-141.
- 677 Schmidt-Vogt, D., 2001. Secondary forests in swidden agriculture in the highlands of Thailand. *Journal of Tropical Forest Science* 13(4), 748–767.
- 680 Schulze, E.D., 1983. Root-shoot interactions and plant life forms. Netherlands Journal of Agricultural Science 4, 291-303.
- Shi, J., Tang, J., Zhang, G., Bai, K., 2001. A preliminary study on the biomass of plant community in Swidden land after
 slash and burn in Xishuangbanna. *Chinese Journal of Ecology 205*, 12-15 (in Chinese with English abstract).
- Shi, J., Wang, Z., Yu, S., Quan, X., Sun, Y., Jia, S., Mei, L., 2008. Estimation of fine root production, mortality and turnover with Minirhizotron in Larix gmelinii and Fraxinus mandshurica plantations. *Frontiers of Biology in China 3(4)*, 496-506.
 687
- Shipley, B. ,Meziane, D., 2002. The balanced-growth hypothesis and the allometry of leaf and root biomass allocation.
 Functional Ecology 16, 326-331.
- 691 Sim, B. L., Nykvist, N., 1990. Impact of forest harvesting and replanting. Journal of Tropical Forest Science 3(3), 251-284.
- 693 Smiley, G.L., Kroschel, J., 2008. Temporal change in carbon stocks of cocoa-gliricidia agroforests in Central Sulawesi,
 694 Indonesia. *Agroforestry Systems 73*, 219-231.
- 695 Smith, P., Lanigan, G., Kutsch, W.L., Buchmann, N., Eugster, W., Aubinet, M., Ceschia, E., Béziat, P., Yeluripati, J.B.,
- 696 Osborne, B., Moors, E.J., Brut, A., Wattenbach, M., Saunders, M., Jones, M., 2010. Measurements necessary for 697 assessing the net ecosystem carbon budget of croplands. *Agriculture, Ecosystems and Environment 139*, 302–315.
- 698
 699 Solichin, L.M., Steinmann, K.H., 2011. Tier 3 biomass assessment for baseline emission in Merang Peat Swamp Forest.
 700 Workshop on Tropical Wetland Ecosystems of Indonesia: Science needs to address climate change adaptation and mitigation. Watilan
 701 Convention Center, Sanur Beach Hotel, Bali.
- 702 Srivastava, S.K., Singh, K.P., Upadhyay, R.S., 1986. Fine root growth dynamics in teak (Tectona grandis Linn. F.).
 703 Canadian Journal of Forest Research 16(6), 1360-1364.
- 705 Stern, N., 2007. The Economics of Climate Change: The Stern Review, Cambridge University Press, Cambridge.

709

712

715

- Stokes, A., Atger, C., Bengough, A.G., Fourcaud, T., Sidle, R.C., 2009. Desirable plant root traits for protecting natural
 and engineered slopes against landslides. *Plant and Soil 324*, 1-30.
- Subedi, K.D., Ma, B.L., Liang, B.C., 2006. New method to estimate root biomass in soil through root-derived carbon. *Soil Biology and Biochemistry 38*, 2212-2218.
- Sundermeier A., Reeder, R., Lal, R., 2012. Soil Carbon Sequestration Fundamentals. www paper AEX-510-05. Available
 at: <u>http://ohioline.osu.edu/aex-fact/0510.html</u> (accessed 5 October 2012).
- Syahrinudin, 2005. The potential of oil palm and forest plantations for carbon sequestration on degraded land in Indonesia. Ecology and
 Development Series No. 28. Cuviller Verlag Göttingen.
- 718 Tang, J., Zhang, J., Song, Q., Cao, M., Feng, Z., 1998. A preliminary study on the biomass of secondary tropical forest in
 719 Xishuangbanna. *Acta Phytoecologica Sinica 226*, 489-498 (in Chinese with English abstract).
- Tang, J., Zhang, J., Song, Q., Huang, Z., Li, Z., Wang, L., Zeng, R., 2003. Biomass and net primary productivity of
 artificial tropical rainforest in Xishuangbanna. *Chinese Journal of Applied Ecology* 14 1, 1-6 (in Chinese with English abstract).
- Tang, J.-W., Pang, J.-P., Chen, M.-Y., Guo, X.-M., Zeng, R., 2009. Biomass and its estimation model of rubber
 plantations in Xishuangbanna, Southwest China . *Chinese Journal of Ecology 28(10)*, 1942-1948 (in Chinese with English
 abstract).
- 725

727

735

- 726 Tollefson, J., 2009. Climate: Counting carbon in the amazon. Nature 461, 1048-1052.
- 728 UNFCCC, 2009. Cost of implementing methodologies and monitoring systems relating to estimates of emissions from deforestation and forest
 729 degradation, the assessment of carbon stocks and greenhouse gas emissions from changes in forest cover, and the enhancement of forest carbon
 730 stocks. United Nations.
- 732 UNFCCC, 2010. United Nations Framework Convention on Climate Change Ad Hoc Working Group on Long-term
 733 Cooperative Action under the Convention Draft Decision [-/CP.16]. Available
- 734 at: http://unfccc.int/meetings/cancun_nov_2010/meeting/6266.php (accessed 1 October 2011).
- 736 UNFCCC, 2011. United Nations Framework Convention on Climate Change (2011) Outcome of the Ad HocWorking
 737 Group on Long-term Cooperative Action Under the Convention (Draft Decision [-/CP.17]). Available at:
 738 <u>http://unfccc.int/meetings/durban_nov_2011/meeting/6245.php</u> (accessed 1 October 2011).
- 740 UN-REDD Programme, 2009. UN-REDD Programme Partner Countries. Available at: http://www.un-
- 741 <u>redd.org/Partner_Countries/tabid/102663/Default.aspx</u> (accessed 17 September 2013).
- 742 van der Meer, P.J., Verwer, C.C., 2011. Towards a reference carbon value for peat swamp forest in Southeast Asia based
- on historical inventory data. Workshop on Tropical Wetland Ecosystems of Indonesia: Science needs to address climate change
 adaptation and mitigation. Watilan Convention Center, Sanur Beach Hotel, Bali.
- van Noordwijk, M., Rahayu, S., Williams, S.E., Hairiah, K., Khasanah, N., Schroth, G., 2004. Crop and tree root-system
- dynamics, in: van Noordwijk, M., Cadisch, G., Ong, C.K. (Eds.), *Below-ground interactions in tropical agroecosystems: Concepts and models with multiple plant components.* CABI Publishing, Oxfordshire, pp.83-107.
- 748 van Vliet, N., Mertz, O., Heineman, A., Langanke, T., Pascual, U., Schmook, B., Adams, C., Schmidt-Vogt, D., Messerli,
- P., Leisz, S., Castella, J-C., Jørgensen, L., Birch-Thomsen, T., Hett, C., Bech-Bruun, T., Ickowitz, A., Vu, Kim Chi,
 Yasuyuki, K., Fox, J., Padoch, C., Dressler, W., Ziegler, A.D., 2012. Trends, drivers and impacts of changes in swidden
- 751 agriculture on tropical forest frontiers: a global assessment. Global Environmental Change 22(2), 418-429.
- 752
- Verwer, C.C., van der Meer, P.J., 2010. Carbon pools in tropical peat forests Towards a reference value for forest biomass carbon in
 relatively undisturbed peat swamp forests in Southeast Asia. Alterra Report 2108. Alterra Wageningen UR, Wageningen.

- Vogt, K.A., Vogt, D.J., Bloomfield, J., 1998. Analysis of some direct and indirect methods for estimating root biomass
 and production of forests at an ecosystem level. *Plant and Soil 200*, 71-89.
- 757

Vogt, K.A., Vogt, D.J., Palmiotto, P.A., Boon, P., O'Hara, J., Asbjornsen, H., 1996. Review of root dynamics in forest
eccosystems grouped by climate, climatic forest type and species. *Plant and Soil 187*, 159-219.

- 760
 761 Warren, M.W., Kauffman, J.B., Murdiyarso, D., Anshari, G., Hergoualc'h, K., Kurnianto, S., Purbopuspito, J.,
 762 Gusmayanti, E., Afifudin, M., Rahajoe, J., Alhamd, L., Limin, S., Iswandi, A., 2012. A cost-efficient method to assess
 763 carbon stocks in tropical peat soil. *Biogeosciences 9*, 4477-4485.
- Wibisono, I.T., Silber, T., Lubis, I.R., Rais, D.S., Suryadiputra, N., Silvius, M., Tol, S., Joosten, H., 2011. Peatlands in Indonesia's National REDD+ strategy. Responding to the public consultation of the draft National REDD+ Strategy of Indonesia issued on August 18, 2011. Wetlands International, Bogor.
- 768

764

- Wilson, J.B., 1988. A review of evidence on the control of shoot: root ratio, in relation to models. *Annals of Botany 61*, 433-449.
- 771
- Yamada, M., Hanamizu, K., Daido, T., Tange, T., Morikawa, Y., 2000. Carbon stock in fast-growing tree species 4
 Acacia mangium man made forest in Madang, Papua New Guinea and general discussion. *Tropical Forestry No* 49, 20-33
- 774 (in Japanese).
- Zemek, O.J., 2009. Biomass and carbon stocks inventory of perennial vegetation in the Chieng Khoi Watershed, NW Vietnam. M.Sc.
 Thesis. University of Hohenheim, Stuttgart.
- Zheng, Z., Feng, Z.-L., Cao, M., Liu, H.-M., Liu, L.-H., 2000. Biomass and net primary production of primary tropical
 wet seasonal rainforest in Xishuangbanna. *Acta Phytoecological Sinica 242*, 197-203 (in Chinese with English abstract).
- Zheng, Z., Feng, Z., Cao, M., Li, Z., Zhang, J., 2006. Forest structure and biomass of a tropical seasonal rainforest in
 Xishuangbanna, Southwest China. *BIOTROPICA 38(3), 318-327*.
- 781 Ziegler, A.D., Fox, J.M., Xu, J., 2009. The rubber juggernaut. *Science* 324, 1024-1025.
- Ziegler, A.D., Fox, J.M., Webb, E.L., Padoch, C., Leisz, S., Cramb, R.A., Mertz, O., Bruun, T.T., Tran, D.V., 2011.
 Recognizing Contemporary Roles of Swidden Agriculture in Transforming Landscapes of Southeast Asia. *Conservation Biology* 25, 846-848.
- Ziegler, A.D., Phelps, J., Yuen, J.Q., Webb, E.L., Lawrence, D., Fox, J.M., Bruun T.B., Leisz, S.J., Ryan, C., Dressler, W.,
 Mertz, O., Pascual, U., Padoch, C., Koh, L.P., 2012. Carbon outcomes of major land-cover transitions in SE Asia: great
 uncertainties and REDD+ policy implications. *Global Change Biology 18, 3087-3099*.
- Zobel, R.W., 2005. Tertiary root systems, in: Zobel, R.W., Wright, S.F. (Eds.), Roots and soil management: Interactions between
 roots and the soil. American Society of Agronomy, Madison, pp. 35-56.
- 793

790

782