

# THE UNIVERSITY of EDINBURGH

# Edinburgh Research Explorer

## Above- and Belowground Carbon Stocks in a Miombo Woodland Landscape of Mozambique

#### Citation for published version:

Ryan, CM, Williams, M & Grace, J 2011, 'Above- and Belowground Carbon Stocks in a Miombo Woodland Landscape of Mozambique', Biotropica, vol. 43, no. 4, pp. 423-432. https://doi.org/10.1111/j.1744-7429.2010.00713.x

#### **Digital Object Identifier (DOI):**

10.1111/j.1744-7429.2010.00713.x

#### Link:

Link to publication record in Edinburgh Research Explorer

**Document Version:** Early version, also known as pre-print

**Published In:** Biotropica

#### **General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



LRH: Ryan, Williams and Grace

RRH: Carbon Stocks in a Miombo woodland

# 5 Above and Below Ground Carbon Stocks in a Miombo Woodland Landscape of Mozambique

Casey M Ryan<sup>1</sup>, Mathew Williams and John Grace

School of GeoSciences

Crew Building

10 University of Edinburgh

Edinburgh EH9 3JN

UK

Received \_\_\_\_; revision accepted \_\_\_\_.

<sup>&</sup>lt;sup>1</sup> Corresponding author. e-mail: casey.ryan@ed.ac.uk, Tel: +44 (0)77 3646 8828, Fax: +44 (0)131 662 0478

#### ABSTRACT

Quantifying ecosystem carbon stocks is vital for understanding the relationship between changes in land use and carbon dioxide emissions. Here we estimate carbon stocks in an area of miombo woodland in Mozambique, by identifying the major carbon stocks and their

- 5 variability. Data on the biomass of tree stems and roots, saplings, and soil carbon stocks are reported and compared to other savannas systems around the globe. A new allometric relationship between stem diameter and tree stem and root biomass is presented, based on the destructive harvest of 29 trees. These allometrics are combined with an inventory of 12,733 trees on 58 plots over an area of 27 ha.
- 10 Ecosystems carbon stocks totalled 110 tC/ha, with 76 tC/ha in the soil carbon pool (to 50 cm depth), 21.2 tC/ha in tree stem biomass, 8.5 tC/ha in tree coarse root biomass and 3.6 tC/ha in total sapling biomass. Plot-level tree root:stem ratio varied from 0.27–0.58, with a mean of 0.42, slightly higher than the mean reported for 18 other savanna sites with comparable above-ground biomass (R:S = 0.35). Tree biomass (stem + root) ranged from
- 3.1-86.5 tC/ha, but the mean (32.1 tC/ha) was well constrained (95% CI 28.4 36.6). In contrast, soil carbon stocks were almost uniformly distributed and varied from 32-133 tC/ha. Soil carbon stocks are thus the major uncertainty in the carbon storage of these woodlands. Soil texture explained 53% of the variation in soil carbon content, but only 13% of the variation in woody carbon stocks. The history of disturbance (fire, elephants,
- 20 logging/charcoal production, and shifting cultivation) is likely to decouple changes in woody carbon stocks from soil carbon stocks, mediated by tree-grass interactions.

Key words: biomass, carbon stocks, miombo, Nhambita, soil, stem, root

MIOMBO WOODLANDS ARE CHARACTERISTIC SAVANNA WOODLANDS IN SUB TROPICAL SOUTHERN AFRICA, dominated by species of the genera *Brachystegia*, *Julbernardia* and *Isoberlinia*. They occupy 11 African countries: Angola, Burundi, Botswana, Democratic Republic of the Congo, Malawi, Mozambique, Namibia, South Africa, Tanzania, Zambia

- 5 and Zimbabwe. Miombo woodlands occurs on nutrient poor soils and in areas with a distinct wet and dry seasons (annual rainfall of 700-1400 mm). They are characterised by frequent fires in the dry season (Frost 1996; Ryan and Williams 2010). The range of biophysical ecosystem goods and services provided by these woodlands is extensive, ranging from the provision of food, fuel, medicine and construction materials to larger-
- scale carbon and water management services. More than 100 million people live in, or obtain resources from, miombo, and a high proportion (up to 50% in some areas) of rural income is dependant on the woodlands, particularly among the poorest of the poor (Campbell *et al.* 2007). These woodlands formerly covered 2.7 million km<sup>2</sup> (Frost 1996), but are currently being lost and degraded to meet agricultural and energy needs (Brouwer
- 15 and Falcão 2004). Rural land use in the miombo region has for centuries been dominated by subsistence agriculture, in which woodland is cleared, by burning and felling, to grow staple crops such as maize and sorghum for a number of years before abandonment (Frost 1996; Williams *et al.* 2008).

Rapidly increasing atmospheric CO<sub>2</sub> concentrations have prompted increasing
attention to the preservation of carbon stocks in tropical ecosystems. The development of a system of payments to reduce emissions from deforestation and degradation (REDD, (Angelsen *et al.* 2009)), relies on knowledge of the magnitude of such carbon stocks.

Indeed Houghton (2005), show that knowledge of the carbon density (tC/ha) of tropical land cover is the major uncertainty in the estimation of carbon emissions from land use change. Here we address the need for carbon density information, reporting ecosystem carbon stocks for a site in the relatively understudied, but wide-spread and unique, miombo

- savanna woodlands. There is a high degree of uncertainty in carbon density estimates for savannas, particularly in the below-ground component of biomass, which in some savannas are reported to exceed the above-ground biomass (Grace *et al.* 2006). Furthermore, in savanna woodlands, soil carbon stocks typically exceed woody carbon stocks, and when the woodland is cleared, loss of soil carbon can be a significant flux (Walker and Desanker
  2004). Thus there is a need for: (1) estimates of carbon densities in all the major carbon
- pools of miombo woodlands; (2) an assessment of the spatial variability of these stocks; and (3) a quantification of the correlation between the different carbon pools, which might allow for efficient sampling.

This paper seeks to address these needs by providing estimates of ecosystem carbon stocks in miombo woodlands for a part of Africa where such data have not been collected previously. The possible existence of a large below-ground biomass pool is investigated and the root:stem biomass ratio is placed in the context of a global data set. We also examine the links between soil and woody carbon stocks and relate these to soil textural properties. The focus here is the woodlands of Sofala Province, in central Mozambique.

20 This paper is novel in that it presents a stem diameter-root biomass allometric for the miombo ecosystem, and quantifies all the major ecosystem carbon stocks at a single miombo site.

#### METHODS

SITE DESCRIPTION.– The Nhambita area (Fig. S1 & S2) is in the south of Gorongosa District in the province of Sofala in central Mozambique (18.979°S, 34.176°E). The area is partly inside the Gorongosa National Park, and occupies the western flank of the Rift Valley, the

- 5 Báruè Midlands (Tinley 1977). Nhambita falls into the Köppen Aw climate classification (Tropical Savanna) with a hot wet summer and cooler dry winter. The nearest source of meteorological data is a weather station at Chitengo, 25 km to the east of Nhambita. Here, mean annual rainfall (Oct 1-Sept 30) for the years 1956-1969 and 1998-2007 was 850±269 mm (mean ±SD). The rainfall is highly seasonal, with 82 percent falling in the five months
- between November and March. May-September receive less than 20 mm/mo, on average.
   Most of the study area is on free-draining sandy soils derived from metamorphic and
   migmatitic gneiss and granite (Tinley 1977). The soils in the south east of Nhambita are
   more hydromorphic in nature. On the sandy soils, termite (*Macrotermes*) mounds provide
   'islands' of loamy fertile soils and occur at a density of around 1 per hectare (Tinley 1977).
- 15 The vegetation consists of open woodlands, mainly dominated by species of the Caesalpinioideae, which includes the miombo species. In the less well drained areas, species from the genera *Acacia* and *Combretum* dominate. The trees form an open canopy surrounded by a continuous grass layer. The grass senesces in the dry season and provides the fuel for the frequent wildfires.
- 20 CARBON STOCK ASSESSMENT.– The major carbon pools of the ecosystem were quantified (tree stem and root biomass; sapling stem and root biomass; and soil carbon from 0-50 cm depth). To estimate vegetation (tree and sapling) carbon stocks we used the standard

approach (Chave *et al.* 2004) of a tree inventory recording stem diameter ( $D_{BH}$ ) and an allometric relationship to convert diameter to estimates of woody mass. Previous work has shown that the choice of allometric equation can strongly influence the calculation of tree biomass in miombo (Williams *et al.* 2008), a widespread problem (Chave *et al.* 2005). Thus

5 we developed a new site-specific allometric equation for both above- and below-ground biomass.

TREE STEM AND ROOT BIOMASS.– Stem and root biomass are calculated based on the inventory of 58 plots (total area 27.2 ha) and a new allometric model that relates diameter to stem and root biomass. Our approach was to compile as much inventory data as possible,

- 10 even if it was collected for different purposes. As such we used data from 58 plots recorded between 2004 and 2007:
  - Fifteen square permanent sample plots (PSPs) of 1 ha, randomly located with respect to the network of tracks and roads in the study area. We established the plots 250 m from, and perpendicular to, 15 randomly chosen points on the road and track network ('road markers').
  - Thirty square plots of 0.25 ha (Williams *et al.* 2008), also perpendicular to the 15 road markers (above), but at different distances from the road marker (randomly selected from 200, 450, 700 and 950 m).
  - Eight triangular 0.28 ha plots subsequently used in the fire experiment of Ryan and Williams (2010), near the centre of the study area.

20

15

• Five circular plots of 0.5 ha sampled away from the road network on a randomly chosen path through the study area.

Appendix 1 contains a map of the plot locations. On each plot all live, standing stems >5 cm D<sub>BH</sub> (diameter at breast height, 1.3 m) were inventoried.

5 DESTRUCTIVE HARVEST.– To estimate biomass from  $D_{BH}$ , allometric models of stem, root and total (stem+root) biomass were parameterised from destructively sampled stems in the study area. The allometric models are of the form:

#### $\log(B) = \alpha \cdot \log(D_{BH}) + \beta$

Where *B* is biomass of the stem or root, and *D* is the stem diameter.  $\alpha$  and  $\beta$  are 10 parameters to be estimated with linear least squares (Matlab, Mathworks Inc). log indicates the natural logarithm.

In October 2007, 29 trees were destructively harvested. Target trees were chosen from the dominant four species in the inventory. The  $D_{BH}$  range of the target trees exceeded the inter-quartile range of each species in the inventory. At five randomly located points on the road network, we located trees that matched the  $D_{BH}$  and species criteria as closely as possible. These trees were felled and weighed, separating trunk and branch biomass. Thereafter, multiple subsamples were taken from the felled trees from both the trunk (cubes ~10x10x10 cm) and branches (cylinders ~10 cm long). These subsamples were weighed and then dried in an oven (~80 ° C until constant mass) to determine their dry mass fraction (DMF, defined as dry mass/wet mass). Three subsamples from 19 of the trees were returned to Edinburgh for %C analysis (Carbo-Erba/400 automated CN analyser). Biomass ( $B_{dest}$ , kgC) was calculated as the product of wet mass, DMF and %C.

Twenty three of the destructively harvested trees were excavated to provide data on coarse root biomass. For each sampled stem, the area under the canopy was excavated to approximately 1 m depth using a mechanical digger. The excavated soil was sifted by hand and large root (>2 cm diameter) biomass was collected, weighed and then subsampled for moisture content and %C using the same method as for the aboveground parts (see above). Any large roots that extended deeper than 1 m were excavated individually, although it was often impossible to follow them to their very end; we excavated beyond 2 m in most cases, 10 and until the remaining roots had a diameter < 5 cm. Large primary roots (>10 cm

diameter) were weighed and sub sampled separately from smaller roots.

5

We parameterised separate allometric models for stem, root and total biomass, using the log-transformed variables to reduce heteroskedasticity. Values were back-transformed directly without the application of correction factors (Sprugel 1983).

- 15 SAPLING STEM AND ROOT BIOMASS. – Saplings, defined as stems <5 cm  $D_{BH}$  but greater than 0.3 cm at 10 cm above ground level  $(D_{10})$ , were measured and recorded on the eight fire experiment plots in 2007. Allometric models of sapling stem, root and total biomass were parameterised with data from a destructive sample of 45 saplings. The harvested saplings were located in the areas that had to be cleared to excavate the large trees (see above).
- Power fits did not give satisfactory results, so 2<sup>nd</sup> order polynomials were fitted to the data, 20 to relate  $D_{10}$  to wet biomass. Because the saplings typically had 3 or 4 similar sized stems

growing from a single root stock, we calculate an 'effective'  $D_{10}e$  of each organism, defined as:

$$D_{10}e = 2 \cdot \sqrt{\frac{\sum_{i=1}^{s} \pi \left(\frac{D_{10}}{2}\right)^2}{\pi}}$$
(1)

Where  $D_{10}e$  is the effective diameter of a multiple stemmed sapling and  $D_{10}$  is the diameter at 10 cm above the ground of each of the of the *s* stems.

5

Ten of the saplings were dried (~80°C for 24 hrs) and reweighed to determine dry mass fraction. We measured %C on subsamples of seven saplings using the same equipment as for the large trees (above).

SOIL PROPERTIES.- Soil was sampled on 13 of the permanent sample plots (PSPs). On each

10 PSP, nine sampling points were established on a 20 m grid. At five of the sampling points small pits were dug, and soil sampled at depths of 0, 5, 15, 25 and 50 cm. At the remaining sampling points only surface soil was sampled.

For elemental analysis (%C and %N) all samples were dried, sieved (2 mm sieve) and ball milled to produce a fine flour. Percentage C and N was determined using the same

15 equipment as the woody samples (above). For particle size analysis, samples at similar depths were blended to produce one sample per depth per plot. These blended samples were sieved (< 2mm), and then air dried. To de-flocculate the sample prior to particle size analysis, each sample was left overnight in a sonic bath, in a solution of 4 percent sodium hexametaphosphate. It was then analysed for percent particle size by volume using a Beckman Coulter LS 230 laser diffraction system. We report the percentage (by volume) of clay (d< 0.002 mm diameter), silt (0.002<d<0.05 mm) and sand (0.05 mm<d<2 mm), where d is the particle diameter.

To estimate bulk density, three replicate samples were extracted using a sharpened steel cylinder of known volume (209 cm<sup>3</sup>) from the depth profile at the centre of each plot at 2.5 and 30 cm depth. Dry soil bulk density was calculated from dry weight (samples were placed in an oven at 40 °C for 24 hours). Total soil C stock was determined by stepwise integration of the profile data of soil C content from 0-50 cm.

STATISTICAL TESTS.- The distribution of plot-level carbon stocks was tested for goodness of

- 10 fit to the normal and Weibull distributions, using the Jarque-Bera (J-B) and Anderson-Darling test, respectively. Having established an appropriate distribution, we report the estimated population mean and its 95% confidence interval, estimated by maximum likelihood using Matlab. Unless otherwise stated, ± indicates the standard error of the mean. For piecewise curve fitting we used the Shape Language Modeling tool (D'Errico
- 15 2009). We also assessed the degree of spatial dependence of our sampled plots by plotting and inspecting an experimental semi-variogram. All curve fitting was done in Matlab using the curve fitting toolbox (v2.1). We report the adjusted r<sup>2</sup> and the root mean squared error (RMSE) of each fit.

TERMINOLOGY. – Throughout this paper we use *B* to denote tree biomass, *SB* for sapling

20 biomass, *R*:*S* for the root:stem ratio. Subscript 's' for stem, 'r' for root, 't' for  $_{s+r}$ .  $D_{BH}$ indicates diameter at 1.3 m in cm,  $D_{10}$  diameter at 10 cm above the ground, in mm. All stocks are in tC/ha on a dry weight basis. We refer to the attributes of the destructively

harvested stems with the subscript ' $_{dest}$ '. log indicates a natural log. All species names follow Coates Palgrave (2002).

#### RESULTS

TREE STEM AND ROOT BIOMASS. – The 29 sampled trees ranged in  $D_{BH}$  from 5-73 cm. As

- well as the 20 target trees of the five dominant species (*Brachystegia boehmii* Taub., *Brachystegia spiciformis* Benth., *Combretum adenogonium* Steud. ex A.Rich., *Diplorhynchus condylocarpon* (Muell.Arg.) Pichon), we included nine other trees including two further species (*Crossopteryx febrifuga* (Afzel. ex G.Don) Benth. and *Pericopsis angolensis* (Baker) Meeuwen) that were excavated to allow access to the target species. The
- 10 sampled stems had wet mass from 8 -10,309 kg (mean 1,043 kg). The subsamples had dry mass fraction (DMF, fresh mass/dry mass) that averaged 0.65±0.01 for the trunk and 0.59±0.02 for the branches. Trunk subsample moisture content was significantly different from the branch subsamples (two tailed t-test, P <0.0001), so dry mass was calculated separately for branch and trunk mass, using stem specific moisture content values. This</p>
- 15 calculation yielded dry mass estimates of 4 6,954 kg dry matter (mean 674 kg). Percent C values were not different (two tailed t-test, P = 0.366) between trunk and branch subsamples, so the mean (47%±0.20) was used for all conversions to C mass. Final values of stem biomass ranged from 2-3268 kg C (mean 320 kg C).

Root mass for the 23 excavated trees varied from 1-760 kg C, with a mean of 123

20 kg C.  $D_{BH}$  of the excavated trees ranged from 5-72 cm. Similar to stem values, root carbon content also averaged 47±0.23 percent. Dry mass fraction averaged 0.59±0.01.

We pooled all destructive data from all species to produce one allometric model for the site, after preliminary analysis showed no significant difference (no separation of 95% CIs) in the model parameters for a fit of  $D_{BH,dest}$  and  $B_{dest}$  for the different species.

Natural log-transformed allometric models of diameter (cm) vs stem, root and total
woody biomass (kgC) fitted well (Fig. 1, upper panel; Table 1). The stem and root allometric models showed that root:stem decreases with stem size (Fig. 1, lower panel; Table 1).

INVENTORIES.— The allometric equations were applied to the inventory data to estimate stem, root and total woody biomass at a plot level. Plot total tree biomass  $(B_t)$  was not

normally or log-normally distributed, but the Anderson-Darling test showed that our sample followed a Weibull distribution. *B<sub>t</sub>* varied from 3.1–86.5 tC/ha, with a mean of 32.1 (95% CI 28.4, 36.6). *B<sub>s</sub>* ranged from 1.9-60.9 tC/ha with a mean of 21.2 (18.7, 24.3). *B<sub>r</sub>* ranged from 1.1-17.0 with a mean of 8.5 (7.7-9.6). Plot-level R:S was normally distributed and varied from 0.27–0.58, mean 0.42 (0.41-0.44) and was weakly but significantly correlated with *B<sub>t</sub>* (*R*:*S* = -0.00238 *B<sub>t</sub>* + 0.499, *r<sup>2</sup>* = 0.45, *P*<0.0001). *B<sub>r</sub>* was well correlated with *B<sub>s</sub>* (*B<sub>r</sub>* = 0.87 *B<sub>s</sub><sup>0.76</sup>*, *r<sup>2</sup>* = 0.94, RMSE = 0.87 tC/ha, *N* = 58).

Plot basal area was a good predictor of plot total woody carbon stock (Figure 3):

$$B_t = 3.972 \cdot BA \tag{2}$$

 $r^2 = 0.76$ , RMSE = 7.82 tC/ha, N = 58

Where:  $B_t$  is total (above and below ground) tree carbon stock (tC/ha) and *BA* is tree basal area (m<sup>2</sup>/ha).

SAPLING BIOMASS.– The allometric equations for sapling stem and root biomass were fitted with  $2^{nd}$  order polynomials and had  $r^2$  from 0.65-0.75 (Table 2). Dry mass fraction was

5 0.61±0.05, based on moisture measurements of the 10 saplings that were dried. %C was
44.6±0.3, based on 7 subsamples.

Using these allometric equations, and the dry mass and %C data to calculate standing stock of sapling biomass, yielded the following plot level data:  $SB_r$ , ranged from 0.7-5.1 tC/ha, with a mean of 2.5 tC/ha (95%CI 1.0-3.9).  $SB_s$  ranged from 0.3-2.4 tC/ha,

10 with a mean of 1.1 tC/ha (0.49-1.8);  $SB_t$  ranged from 1.0-7.5 with a mean of 3.6 tC/ha (1.5-5.6). Plot level sapling root:stem averaged 2.16 (2.13-2.19).

SOIL CARBON AND NITROGEN.– Across the 13 PSPs with soil data, soil carbon content was ~3 percent at the surface (Fig 4 c), decreasing to ~ 1 percent at 15 cm depth, and remained around this value at greater depths. Nitrogen content followed a similar profile (Fig 4 d)

15 with the C:N ratio decreasing from 17.2 at the surface to 14.5 at a depth of 0.5 m.

Soil carbon stocks (Fig 5) averaged  $8.0\pm0.4$  tC/ha ( $\pm 1$  SEM) from 0-2.5 cm, 17.7 $\pm 2.0$  from 2.5-10 cm, 16.0 $\pm 2.4$  from 10-20 cm and 34.5 $\pm 5.4$  tC/ha from 20-50 cm. Summing the 0-50 cm profile, soil C stocks averaged 76.3 $\pm 9.9$  tC/ha. However, the distribution of the 13 plot-level soil C stock values was almost uniform and ranged from

20 32-133 tC/ha (Fig 2), so that estimating the population mean is problematic. The mean bulk density at the surface was  $1.2\pm0.03$  t/m<sup>3</sup>, and  $1.5\pm0.04$  t/m<sup>3</sup> at 30 cm.

SOIL TEXTURE.– Soil texture at plot level varied from 3-9 percent clay, 20-48 percent silt and 46-76 percent sand, with a profile mean of 6, 33 and 60 percent, respectively. There was very little variation in soil texture with depth: mean silt content decreased down the profile from 34 to 28 percent and clay content increased from 5.6-6.7 percent (Fig 4a & b).

Soil texture was correlated with plot level soil carbon stock (Fig 6, Table 3). There was no correlation between plot level tree biomass ( $B_t$ ) and soil carbon stock ( $C_s$ ) (P = 0.5  $r^2 = 0.05$ , N = 13), and total tree biomass was only weakly negatively correlated with clay content (P = 0.16,  $r^2 = 0.10$ ; see Table 3).

#### DISCUSSION

5

- OVERVIEW AND COMPARISON WITH OTHER SITES. At our site, ecosystem carbon stocks totalled 110±13 tC/ha (Table 4). Soil carbon dominated the total, contributing 76.2 tC/ha (70%). Woody biomass totalled 33.3 tC/ha (30%), dominated by tree biomass (29.7 tC/ha). The division between soil carbon and total carbon stocks in Nhambita (70% in soil) is similar to the 60 percent reported in Walker and Desanker (2004) for Malawian miombo.
- Above-ground woody biomass in Nhambita (21.2 tC/ha) can be compared to Frost's (1996) "average for dry miombo" in Zimbabwe and Zambia of 26 tC/ha (dry mass is converted to C mass assuming 47 %C throughout this discussion). Comparing this study to that of Ribeiro *et al* (2008a) in Niassa Reserve in miombo woodland in Northern Mozambique (annual rainfall 900-1200 mm), our woodlands falls into their 'Medium
- 20 density woodland' class for which  $B_s$  was 19.5 tC/ha. Williams' (2007) summary of the data in Post (1982) and Gibbs (2006), found a "live biomass" of 28.2 tC/ha (our data: 29.7)

at the latitude of Nhambita (interpolating their data points with a spline), and soil carbon stocks of 86.6 tC/ha (to 1 m depth; our data 76.2 tC/ha, to 50 cm).

Frost (1996) developed a relationship between annual rainfall and aboveground woody biomass in miombo: using a value for our site of 850 mm/yr, and Frost's equation (Y=0.14 X - 56.21; Y is Mg/ha of dry matter; X is rainfall in mm), our biomass data (21.2 tC/ha) are lower that the equation's prediction (29.5 tC/ha).

5

Plot level root:stem ratios in this study varied from 0.28-0.58, with a mean of 0.42. Root biomass estimates in miombo are rare, but the average of root:stem in this study (0.42) can be compared to Chidumayo's (1997) values of 0.53 and 0.47 for two wet

- miombo plots in Zambia (calculated in Frost 1996). Outside of the miombo region, we found comparable data on B<sub>s</sub> and R:S for 18 sites in 11 countries in Latin America and Africa (Sarmiento and Vera 1979; Murphy and Lugo 1986; Castellanos *et al.* 1991; Woomer 1993; Malimbwi *et al.* 1994; Frost 1996; De Castro and Kauffman 1998; Tiessen *et al.* 1998; Lilienfein *et al.* 2001; Jaramillo *et al.* 2003). This summary shows that R:S falls
- 15 rapidly as  $B_s$  increases, but is very variable, up to values of  $B_s$  of around 10 tC/ha (Fig. 7), thereafter R:S levels off at between 0.3 and 0.5. It can be modelled with a piecewise linear equation, fitted with SLM (Fig. 7):

Where 
$$B_s < 11$$
: R:S = -0.30  $B_s + 3.62$ 

Where  $B_s \ge 11$ : R:S = 0.35 (3)

20 Where *R*:*S* is the root:stem ratio,  $B_s$  is above ground biomass (tC/ha).

For the data from this study, a simple linear model can be fitted:

$$R:S = -0.0046 \cdot B_s + 0.498 \tag{4}$$

 $N = 60, r^2 = 0.46, RMSE = 0.041.$ 

The mean R:S found in this study (0.42) was slightly higher than that found at sites with comparable above-ground biomass (R:S = 0.35), according to our review of the literature (Fig 7). Also, our plot-level R:S data showed a different relationship with aboveground biomass: the Nhambita plots with low  $B_s$  had much lower R:S than comparable sites in the literature (Fig. 7). This suggests a difference, probably in stem size distributions, between low biomass miombo plots, and other savannas with low biomass. Some of our low  $B_s$  plots are dominated by a few, large trees, and thus have a lower *R:S* than if they were composed

of many small trees which tend to have a high individual R:S.

The  $B_s$  vs R:S relationship can be compared to the IPCC LULCF Good Practice Guidelines (Penman *et al.* 2003) default figure for 'Woodland/savanna' of 0.48±0.19 SD, which is based on Australian and Indian data. This default figure appears to underestimate

15 root biomass for low biomass savannas in the neotropics and West Africa, but is similar to values from this study, and for savannas with  $B_S > 11$  tC/ha.

THE RELATIONSHIP BETWEEN CARBON STOCKS IN MIOMBO WOODLAND.— This study has shown that soil and woody carbon stocks are the dominant carbon pools in miombo woodland, but that the size of each carbon pool was not correlated. Soil carbon is highly

variable (32-133 tC/ha; Fig 2), and soil texture explained around half of this variation (Fig
6; Table 3) as is common in tropical soils (Feller and Beare 1997). Woody biomass was

less variable (3.1-86.5 tC/ha), but only weakly correlated with soil texture. This decoupling of the soil and woody carbon stocks is expected, because although the dynamics of wood and soil carbon are clearly linked, there are many other controls on pool size. We summarise the key processes below.

5 Woody biomass in miombo shows both regional trends with rainfall (see above, Frost 1996) and smaller scale trends with topography (Chidumayo 1997). The topographic trends relate to the catenal patterns, with low woody biomass in dambos (bottom lands with impeded drainage (von der Heyden 2004)) and higher biomass on interfluves (Chidumayo 1997). This catenal pattern may account for the weak negative correlation between clay

10 content and woody biomass found in this study. However, this simple dambo-interfluve pattern is far from universal, and we have frequently observed higher levels of woody biomass along river lines, compared to interfluves, primarily on deeper, more fertile alluvial soil.

The spatial pattern of woody biomass described above is subject to frequent and 15 widespread disturbances (Frost 1996) that reduce biomass: primarily clearance for agriculture (Williams *et al.* 2008), charcoal production (Brouwer and Falcão 2004; Falcão 2008) and fire (Ryan and Williams 2010). Elephant activity can also reduce tree populations significantly (Guy 1989; Ribeiro *et al.* 2008b), as can rare frosts (Tafangenyasha 1997), and the interactions of all these effects. Clearance for agriculture is

20 the start of a cycle whereby woody biomass is reduced nearly to zero for the duration of the agricultural activity, but then recovers to pre-disturbance levels within around 30 years of abandonment (Williams *et al.* 2008), creating a mosaic of land cover. The regular fires in

the miombo region can, if too frequent or intense, cause mortality of large and small trees and prevent regeneration (Ryan and Williams 2010), thus leading to a gradual degradation of the woodlands until no woody tree biomass is present (Furley *et al.* 2008).

Soil carbon is also generally reduced after clearance for agriculture (Walker and
Desanker 2004). Indeed Walker and Desanker (2004) report that upon conversion to
agricultural land, miombo woodland loses around 47 percent of its soil C (to 1.5 m depth).
Using the same proportion, conversion to agriculture at our site would thus result in a loss
of 36 tC/ha from the soil, slightly larger than that if all the tree biomass were to be lost
(29.7 tC/ha). Soil carbon is also lower on plots subjected to frequent fires, but the effect of

different fire regimes is similar in magnitude to the influence of soil texture (Bird *et al.*2000).

It is important to note that there is no *a priori* reason to think that low woody cover leads to lower inputs to the soil carbon pool. Grass productivity in miombo is comparable to tree productivity and because low woody biomass tends to encourage high grass biomass

15 (Frost 1996), the input to the soil could be similar under high and low woody cover. However, evidence from Bird *et al* (2000) suggests that soil carbon stocks are higher beneath trees than in adjacent open areas of woodland, which argues for the importance of carbon inputs from trees (or animals that defecate beneath them). However, the lack of a correlation between woody and soil carbon stocks in this study indicates that many other

#### 20 factors influence this relationship at larger scales.

The correlation between soil texture and soil carbon stocks shown here (Fig. 6) offers the most promising route to mapping soil carbon stocks and dealing with their

variability on a range of spatial scales. In contrast, tree below-ground biomass is very well correlated with above-ground biomass, and  $B_r$  can easily be estimated using the *R*:*S* reported here (where only plot level data is available) or our allometric equations (if stem level data is available).

- 5 LIMITATIONS.- In summarising ecosystem C stocks, several caveats must be mentioned. Firstly our plots are located on a transition from miombo woodland to the plains of the Rift valley, where a change in soil type produces a more diverse and complex woodland formation. Secondly, our plots are not evenly or randomly distributed across the landscape, but instead follow the network of small tracks and roads that provide access, and are
- clumped together. To address this, we examined a variogram of B<sub>p</sub> (Fig. S3). The variogram showed no relationship between separation distance and the variance. This suggests that the variance between closely located plots (<2 km apart) is no different from more widely spaced plots (>10 km) and supports the use of the closely spaced plots as independent data. Furthermore, if we use the mean of each set of plots perpendicular to the same road marker as a single value, then our result for above-ground woody biomass is 21.0 (95% CI 18.1-

23.8; N = 28), very similar to the mean of the 58 plots: 21.2 (18.7-24.3).

20

A further caveat is that the use of allometric models, even site specific ones, can introduce significant biases depending on their form, and how the heteroskedasticity of the destructive data is dealt with (Brown *et al.* 1989; Parresol 1999). Our estimates are bound to be subject to such biases, a subject we will address in a further paper. We also note that soils were only sampled at 13 of the plots rather than at all 58. Finally, we did not measure the more transient carbon pools such as fine roots, tree leaves and grass biomass, which

contribute to the total ecosystem carbon stocks. The major missing pool is likely to be grass biomass.

### CONCLUSIONS

At a site in the miombo woodlands of Mozambique, carbon stocks are dominated by soil

- 5 carbon: 76 tC/ha of the total 110 tC/ha are in the surface 50 cm of soil. However soil carbon is very variable across the landscape, and has a broad and almost uniform distribution, rather than converging around a mean. This requires further investigation at various spatial scales if estimates of carbon stocks in this ecosystem are to be better constrained. Soil texture is a reasonable predictor of soil carbon stocks, but soil C was not
- 10 obviously correlated to woody C stocks.

Woody tree carbon averaged  $21.3\pm1.4$  tC/ha above-ground and  $8.6\pm0.5$  tC/ha below-ground, based on allometric relationships. Sapling total biomass averages  $3.6\pm0.9$  tC/ha. Coarse root biomass is a major carbon pool in these ecosystems and is proportionally more important at low values of above-ground biomass because of shifts in root:stem ratio of trees as they grow.

#### ACKNOWLEDGEMENTS

15

The root excavation would not have been possible without the loan of a digger and front end loader from Paul Roche of Harrop-Allin & Sons, and the hard work of many people from the Nhambita community. Gary Goss felled the trees and managed much of the field

20 work. Thanks to Roberto Zolho, then of Gorongosa National Park, for allowing us to work in the Park area, Piet van Zyl, Antonio Serra and Alastair MacCrimmon of Envirotrade for their extensive assistance and support, the Envirotrade forestry technicians 'Joey' Albasine Adolfo Mucavele, Afonso Jornal, Ramaio Saimone, Neto Moulinho, Joao 'Deus' Eduardo, Salomaõ 'Baba' Nhangue, Zito Lindo, Madal and Ernesto for all their hard work in the field, and Ria van Zyl for her hospitality and the use of her oven. We thank the two

5 reviewers for their helpful comments.

#### LITERATURE CITED

Research.

15

35

- ANGELSEN, A., S. BROWN, C. LOISEL, L. PESKETT, C. STRECK and D. ZARIN (2009). Reducing Emissions from Deforestation and Forest Degradation (REDD): An Options Assessment Report
- 5 BIRD, M. I., E. M. VEENENDAAL, C. MOYO, J. LLOYD and P. FROST (2000). "Effect of fire and soil texture on soil carbon in a sub-humid savanna (Matopos, Zimbabwe)." Geoderma 94(1): 71-90.

BROUWER, R. and M. P. FALCÃO (2004). "Wood fuel consumption in Maputo, Mozambique." Biomass and Bioenergy 27(3): 233-245.

- 10 BROWN, S., R. GILLESPIE and E. LUGO (1989). "Biomass Estimation Methods for Tropical Forests with Applications to Forest Inventory Data." Forest Science 35: 881-902.
  - CAMPBELL, B. M., A. ANGELSEN, A. CUNNINGHAM, Y. KATERERE, A. SITOE and S. WUNDER (2007). Miombo woodlands - opportunities and barriers to sustainable forest management, Center for International Forestry

CASTELLANOS, J., M. MAASS and J. KUMMEROW (1991). "Root biomass of a dry deciduous tropical forest in Mexico." <u>Plant and Soil</u> V131(2): 225-228.

- 20 CHAVE, J., C. ANDALO, S. BROWN, M. A. CAIRNS, J. Q. CHAMBERS, D. EAMUS, H. FÖLSTER, F. FROMARD, N. HIGUCHI, T. KIRA, J. P. LESCURE, B. W. NELSON, H. OGAWA, H. PUIG, B. RIÉRA and T. YAMAKURA (2005). "Tree allometry and improved estimation of carbon stocks and balance in tropical forests." Oecologia V145(1): 87-99.
- 25 CHAVE, J., R. CONDIT, S. AGUILAR, A. HERNANDEZ, S. LAO and R. PEREZ (2004). "Error propagation and scaling for tropical forest biomass estimates." <u>Philosophical Transactions of the Royal Society of</u> <u>London Series B-Biological Sciences</u> 359(1443): 409-420.
- CHIDUMAYO, E. N. (1997). <u>Miombo ecology and management : an introduction</u>. 30 London, IT Publications in association with the Stockholm Environment Institute.
  - COATES PALGRAVE, K., R. B. DRUMMOND, E. J. MOLL and M. COATES PALGRAVE (2002). <u>Trees of southern Africa</u>. Cape Town, Struik Publishers.
  - D'ERRICO, J. (2009). SLM Shape Language Modeling for Matlab. http://www.mathworks.com/matlabcentral/fileexchange/24443.

DE CASTRO, E. A. and J. B. KAUFFMAN (1998). "Ecosystem structure in the Brazilian Cerrado: a vegetation gradient of aboveground biomass, root mass and consumption by fire." <u>Journal of Tropical Ecology</u> 14: 263-283.

- 40 FALCÃO, M. P. (2008). <u>Charcoal production and use in Mozambique, Malawi,</u> <u>Tanzania, and Zambia: historical overview, present situation and</u> <u>outlook</u>. The Conference on Charcoal and Communities in Africa, INBAR.
- 45 FELLER, C. and M. H. BEARE (1997). "Physical control of soil organic matter dynamics in the tropics." Geoderma 79(1-4): 69-116.

- FROST, P. (1996). The ecology of Miombo woodlands. The Miombo in transition : woodlands and welfare in Africa. B. M. Campbell. Bogor, Indonesia, Center for International Forestry Research: 11-55.
- 5 FURLEY, P. A., R. M. REES, C. M. RYAN and G. SAIZ (2008). "Savanna burning and the assessment of long-term fire experiments with particular reference to Zimbabwe." <u>Progress in Physical Geography</u> 32(6): 611-634.
- GIBBS, H. K. (2006). Olson's Major World Ecosytem Complexes Ranked by Carbon in Live Vegetation: An Updated Database Using the GLC2000 Land Cover Product. . Oak Ridge, Tennessee, Carbon Dioxide Information Center, Oak Ridge National Laboratory.

- GRACE, J., J. S. JOSÉ, P. MEIR, H. S. MIRANDA and R. A. MONTES (2006). "Productivity and carbon fluxes of tropical savannas." <u>Journal of</u> Biogeography 33(3): 387-400.
- GUY, P. R. (1989). "The Influence of Elephants and Fire on a *Brachystegia-Julbernardia* Woodland in Zimbabwe." Journal of Tropical Ecology 5(2): 215-226.
- HOUGHTON, R. A. (2005). "Aboveground forest biomass and the global carbon 20 balance." Global Change Biology 11(6): 945-958.
  - JARAMILLO, V. J., J. B. KAUFFMAN, L. RENTERÍA-RODRÍGUEZ, D. L. CUMMINGS and L. J. ELLINGSON (2003). "Biomass, Carbon, and Nitrogen Pools in Mexican Tropical Dry Forest Landscapes." Ecosystems V6(7): 609-629.
- LILIENFEIN, J., W. WILCKE, R. ZIMMERMANN, P. GERSTBERGER, G. M. ARAÚJO and W. ZECH (2001). "Nutrient storage in soil and biomass of native Brazilian Cerrado." Journal of Plant Nutrition and Soil Science 164(5): 487-495.
- MALIMBWI, R. E., B. SOLBERG and E. LUOGA (1994). "Estimation of biomass and volume in miombo woodland at Kitulangalo Forest Reserve, Tanzania."
   Journal of Tropical Forest Science 7(2): 230-242.
  - MURPHY, P. G. and A. E. LUGO (1986). "Ecology of Tropical Dry Forest." Annual Review of Ecology and Systematics 17: 67-88.
  - PARRESOL, B. R. (1999). "Assessing Tree and Stand Biomass: A Review with Examples and Critical Comparisons." Forest Science 45: 573-593.
- PENMAN, J., M. GYTARSKY, T. HIRAISHI, T. KRUG, D. KRUGER, R. PIPATTI, L. BUENDIA, K. MIWA, T. NGARA, K. TANABE and F. WAGNER (2003). <u>Good Practice</u> <u>Guidance for Land Use</u>, <u>Land-Use Change and Forestry and Definitions</u> <u>and Methodological Options to Inventory Emissions from Direct</u> <u>Human-induced Degradation of Forests and Devegetation of Other</u> <u>Vegetation Types</u>, Institute for Global Environmental Strategies (IGES) for the Intergovernmental Panel on Climate Change.
  - POST, W. M., W. R. EMANUEL, P. J. ZINKE and A. G. STANGENBERGER (1982). "Soil carbon pools and world life zones." Nature 298(5870): 156-159.
- RIBEIRO, N. S., S. S. SAATCHI, H. H. SHUGART and R. A. WASHINGTON-ALLEN (2008a).
  45 "Aboveground biomass and leaf area index (LAI) mapping for Niassa Reserve, northern Mozambique." <u>Journal of Geophysical Research-</u> Biogeosciences 113.

- RIBEIRO, N. S., H. H. SHUGART and R. WASHINGTON-ALLEN (2008b). "The effects of fire and elephants on species composition and structure of the Niassa Reserve, northern Mozambique." <u>Forest Ecology and Management</u> 255(5-6): 1626-1636.
- 5 RYAN, C. M. and M. WILLIAMS (2010). "How does fire intensity and frequency affect miombo woodland tree populations and biomass? ." <u>Ecological</u> Applications In press.
  - SARMIENTO, G. and M. VERA (1979). "Composición, estructura, biomasa y producción de diferentes sabanas en los Llanos de Venezuela." Boletín de la Sociedad Venezolana de Ciencias Naturales 136: 5-41.
  - SPRUGEL, D. G. (1983). "Correcting for Bias in Log-Transformed Allometric Equations." Ecology 64(1): 209-210.
    - TAFANGENYASHA, C. (1997). "Tree loss in the Gonarezhou National Park (Zimbabwe) between 1970 and 1983." Journal of Environmental Management 49(3): 355-366.
    - TIESSEN, H., C. FELLER, E. V. S. B. SAMPAIO and P. GARIN (1998). "Carbon Sequestration and Turnover in Semiarid Savannas and Dry Forest." Climatic Change 40(1): 105-117.
- TINLEY, K. L. (1977). Framework of the Gorongosa Ecosystem. <u>Faculty of</u> Science. Pretoria, University of Pretoria: 184.
  - VON DER HEYDEN, C. J. (2004). "The hydrology and hydrogeology of dambos: a review." Progress In Physical Geography 28(4): 544-564.
  - WALKER, S. M. and P. V. DESANKER (2004). "The impact of land use on soil carbon in Miombo Woodlands of Malawi." Forest Ecology And Management 203(1-3): 345-360.
  - WILLIAMS, M., C. M. RYAN, R. M. REES, E. SAMBANE, J. FERNANDO and J. GRACE (2008). "Carbon sequestration and biodiversity of re-growing miombo woodlands in Mozambique." <u>Forest Ecology and Management</u> 254(2): 145-155.
- 30 WOOMER, P. L. (1993). "The impact of cultivation on carbon fluxes in woody savannas of Southern Africa." <u>Water, Air, & Soil Pollution</u> 70(1): 403-412.

35

10

15

### TABLES

TABLE 1. *Tree biomass allometric equations*.  $B_{dest}$  is the destructively sampled tree stem biomass (*s*), tree coarse root biomass (*r*), and total tree biomass (*t*, i.e. stem + root), all in kgC.  $D_{BH,dest}$  is diameter at breast height in cm. Log is the natural log.  $R:S_{dest}$  is the root stem ratio.

Equation	$\mathbb{R}^2$	RMSE	Ν
$\log(B_{dest,s}) = 2.601 \log(D_{BH,dest}) - 3.629$	0.93	0.52 log(kgC)	29
$\log(B_{dest,r}) = 2.262 \log(D_{BH,dest}) - 3.370$	0.94	0.43 log(kgC)	23
$\log(B_{dest,t}) = 2.545 \log(D_{BH,dest}) - 3.018$	0.98	0.30 log(kgC)	23
$R:S_{dest} = -0.2671 \cdot \log(D_{BH,dest}) + 1.334$	0.36	0.27	23

TABLE 2. Sapling biomass allometric equations.  $SB_{dest}$  is wet sapling biomass in kg,  $D_{10}e$  is the effective diameter (mm) of the harvested saplings at 10 cm above ground (see Eq 1).

Equation	R <sup>2</sup>	RMSE	Ν
$SB_{s,dest} = 0.0007645 \cdot D_{10}e^2 + 0.004645 \cdot D_{10}e + 0.03876$	0.65	0.47 kg	45
$SB_{r,dest} = 0.001784 \cdot D_{10}e^2 + 0.0001413 \cdot D_{10}e + 0.15839$	0.72	0.84 kg	45
$SB_{t,dest} = 0.0025485 \cdot D_{10}e^2 + 0.0047865 \cdot D_{10}e + 0.19715$	0.75	1.1 kg	45

TABLE 3. *Relationships between soil texture and carbon stocks*.  $C_s$  is soil carbon stock (0-50 cm) in tC/ha,  $B_t$  is tree carbon stock (above- and below-ground, tC/ha), K and Z are clay and sand contents in percent by volume.

Equation	r <sup>2</sup>	RMSE (tC/ha)	Р	Ν
$C_s = 12.98 \cdot K + 0.2$	0.53	25.5	0.004	13
$C_s = -2.70 \cdot Z + 243$	0.51	24.9	0.004	13
$B_t = -3.70 \cdot K + 54.3$	0.10	16.8	0.16	13
$B_t = 0.81 \cdot Z - 17.2$	0.13	16.6	0.13	13

TABLE 4. Summary of ecosystem carbon stocks in the Nhambita area. All values in tC/ha.  $\pm$  indicates thestandard error of the mean.

Carbon pool		Carbon stock (tC/ha)			
saplings	stem	1.1±0.3			
	roots	2.5±0.6			
	sapling total		3.6±0.9		
trees	stem	21.2±1.4			
	roots	8.5±0.5			
	tree total		29.7±2.1		
soil	0-2.5 cm	8.0±0.4			
	2.5-10 cm	17.7±2.0			
	10-20 cm	16.0±2.4			
	20-50 cm	34.5±5.4			
	soil total		76.3±9.9		
Grand total				110±13	

#### **FIGURE CAPTIONS**

FIGURE 1. A) The allometric models of stem, root and total biomass and the destructively harvested data. The models are fitted to the log of  $D_{BH}$  and  $B_{dest}$ , but are shown back transformed. B) The allometry of the root:stem ratio of the destructively harvested stems.

5 FIGURE 2. Histograms of plot-level stem, root and total woody biomass carbon stocks on the 58 plots, and soil carbon stocks (to 50 cm depth) on 13 plots, as well as plot-level root:stem on the 58 plots. Note change in X-axis values for soil carbon stocks.

FIGURE 3. Plot-level woody carbon stocks regressed against plot basal area. The dashed line shows the 95% prediction interval for a new observation.

10 FIGURE 4. Mean depth profiles of soil texture and elemental concentrations for the 13 plots. The black line shows the mean and the grey area the full range of the observations.

FIGURE 5. Soil carbon stocks at various depths on thirteen of the 1 ha permanent sample plots.

FIGURE 6. Soil texture regressed against soil carbon stock. All data are plot means (N=13) averaged down the profile (0-50 cm depth) in the case of texture, and summed down the profile in the case of carbon stock.

15 Thick lines shows the line of best fit, and the dotted lines show the 95% predication intervals.

FIGURE 7. Root:stem ratios as a function of aboveground biomass, across the seasonally dry tropics. Definitions of stem and root biomass vary according to the source, but are 'coarse' roots (often >2 cm diameter) and trees (mainly >5 cm DBH).

# FIGURES



Figure 1.



Figure 2.



Figure 3.



Figure 4.



Figure 5.



Figure 6. Soil texture regressed against soil carbon stock. All data are plot means (n=13) averaged down the profile (0-50 cm depth) in the case of texture and summed down the profile in the case of carbon stock. Thick lines shows the line of best fit, and the dotted lines show the 95% predication

intervals.



Figure 7.

# **APPENDIX 1**



# Figure S1. Study Area Location

Figure S2. Plot locations



Figure S3. Variogram of the above ground biomass of the sample plots Separation distance of the plots is plotted against the mean variance between them, in 1 km bins.

