THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Thyroid stimulating hormone, free thyroxine and cognitive ability in old age

Citation for published version:

Booth, T, Deary, IJ & Starr, JM 2013, 'Thyroid stimulating hormone, free thyroxine and cognitive ability in old age: The Lothian birth cohort study 1936', *Psychoneuroendocrinology*, vol. 38, no. 4, pp. 597-601.
<https://doi.org/10.1016/j.psyneuen.2012.07.018>

Digital Object Identifier (DOI):

[10.1016/j.psyneuen.2012.07.018](https://doi.org/10.1016/j.psyneuen.2012.07.018)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Psychoneuroendocrinology

Publisher Rights Statement:

© Thyroid stimulating hormone, free thyroxine and cognitive ability in old age : The Lothian birth cohort study 1936. / Booth, Thomas; Deary, Ian J; Starr, John M.
In: *Psychoneuroendocrinology*, Vol. 38, No. 4, 04.2013, p. 597-601.
Research output: Contribution to journal › Article

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Thyroid Stimulating Hormone, Free Thyroxine and Cognitive Ability in Old Age:

The Lothian Birth Cohort Study 1936.

Tom Booth^{1,2}, Ian, J. Deary^{1,2} & John, M. Starr^{1,3}.

¹Centre for Cognitive Ageing and Cognitive Epidemiology, The University of Edinburgh,
EH8 9JZ, Edinburgh, UK.

²Department of Psychology, University of Edinburgh, EH8 9JZ, Edinburgh, UK.

³ Alzheimer Scotland Dementia Research Centre, The University of Edinburgh, Edinburgh,
EH8 9JZ, UK.

FINAL ACCEPTED VERSION BEFORE COPY EDITING

DO NOT CITE

Corresponding Author:

Tom Booth, Centre for Cognitive Ageing and Cognitive Epidemiology, Department of
Psychology, The University of Edinburgh, 7 George Square, EH8 9JZ, UK. Email:

tom.booth@ed.ac.uk ; Phone: (+44) 01316508405.

Abstract

The current study investigated the associations between thyroid stimulating hormone (TSH), free thyroxine (T₄) and cognitive ability (general ability, memory and processing speed), in a large age homogenous sample (n=659) of generally healthy euthyroid older adults.

Associations were considered both at baseline (Mean Age Wave 1=69.5 years; SD=0.8 years) and approximately 3 years later (Mean Age Wave 2=72.5 years; SD=0.7 years). Results indicated mean level decreases across waves in both TSH ($t=10.99, p<0.001$) and T₄ ($t=34.55, p<0.001$). There were no significant associations between TSH and T₄ with any of the cognitive variables at either wave. There was no suggestion of non-linear associations.

The lack of associations supports suggestions that the effects of thyroid hormones on cognition may be restricted to older individuals above a given threshold, and/or those with levels of thyroid hormones within the range defining clinical thyroid disorder.

Keywords: Thyroid Stimulating Hormone; TSH; Free Thyroxine; T₄; Cognitive Ability; Memory; Processing Speed.

1. Introduction

The maintenance of cognitive abilities into later life is a key determinant of a wide variety of positive life outcomes and successful ageing (Deary, 2012). Thyroid function is mooted to be important for the development and retention of cognitive function (Begin et al., 2008), with thyroid dysfunction, most commonly clinical and subclinical hypothyroidism, in later life being associated with both dementia and, more specifically, Alzheimer disease (Smith et al., 2002; Liesbeth et al., 2012). As a result, there is growing interest in whether thyroid hormones at both normal and clinical levels, are associated with cognitive performance across the life span. However, research has failed to show consistent associations between cognitive abilities and thyroid function in samples of generally healthy elderly subjects with subclinical thyroid dysfunction.

For example, St. John et al. (2009) found no association between TSH levels and cognitive ability in a sample of 489 men and women (mean age = 60.5 years). Roberts et al. (2007) found no significant differences in Mini-Mental State Exam (MMSE) or Middlesex Elderly Assessment of Mental State (MEAMS) scores between individuals with subclinical thyroid disorders and euthyroid from a large cross-sectional sample (Birmingham Elderly Thyroid Study; n=5685, age = >65 years). Higher TSH was associated with higher MMSE scores, but not to MEAMS score. Using a subsample of the same cohort in a randomized controlled trial, Parle et al (2010) found no significant differences between the placebo group (n=42, mean age 74.2 years) and a group receiving T₄ hormone treatment (n=52, mean age 73.5 years) in any cognitive measures. Similarly, in a follow up study of 599 older individuals (85 through 89 years), Gussekloo et al (2004) found no associations between TSH, T₃ or T₄ and cognitive functioning. Hogervorst et al. (2008) found that after controlling for the presence thyroid disease, stroke and dementia, higher normal T₄ was associated with lower MMSE scores at baseline, and a greater risk of decline across 2 years, in a sample of 1047 older adults (mean

age at baseline 73.6 years). Conversely, using the NHANES III sample, Beydoun et al. (2012) found that higher levels of thyroxine, TSH were associated with improved cognitive performance on a math test, and a story recall test, in the 60 to 90 years age group (n=5989 and 5878 respectively).

Begin et al. (2008) reporting on five previously published studies of euthyroid samples concluded that, from the mixed results; that it is unclear whether Thyroid Stimulating Hormone (TSH), tri-iodothyronine (T₃) or free thyroxine (T₄) is the most relevant associate of cognitive function; and that the only consistent cognitive association was with memory impairment across all biomarkers. A similar conclusions is reached from the nine studies reviewed by Hogervorst et al (2008; see Table 1, p.1014), in which four studies found some degree of significant association between memory and thyroid hormone levels in older samples (age range across studies 49 to 89).

In totality, results from prior studies have been inconsistent, and provide little evidence for associations between thyroid hormones and cognitive ability in generally healthy older adults. Samuels (2010), in reviewing the general lack of associations found in studies of subclinical hypothyroidism, noted that: a) few studies report on the largely more sensitive T₄ measurements (although for impaired thyroid function TSH might be preferred); and b) most studies rely on cognitive assessments designed to identify gross impairment (e.g. MMSE or MEAMS). Such measures may be insensitive to changes in cognitive function within the normal range of abilities.

In the current study we investigate the associations between TSH, T₄ and major domains of cognitive ability assessed using widely validated psychometric tools, in a large age-homogeneous sample of generally healthy older adults.

2. Methods

2.1 Sample

The current sample is drawn from the Lothian Birth Cohort Study 1936 (LBC1936), a longitudinal study of ageing. Full information on the recruitment and testing of participants has been previously published in study protocols and profile (Deary et al., 2007; Deary et al., 2011). In short, surviving members of the Scottish Mental Health Survey 1947 (SMS1947) resident in Edinburgh and the surrounding Lothian area, were recruited between June 2004 and November 2006.

In total, 1091 participants took part in wave 1 of testing where they completed a wide array of psychometric tests, a physical examination entailing a number of tests of physical functioning (6 metre walk, stand-sit test, grip strength, blood pressure etc.). Blood samples were also taken as part of the physical examination. At wave 2, 866 participants returned approximately three years later and repeated largely the same array of cognitive, psychometric, and physical tests. In the present study, we included participants who (n indicating the numbers who meet inclusion criteria at each step);

Completed both waves 1 and 2 (n=866).

Had complete data for TSH and T₄ (n=810).

Did not report any thyroid disease at both waves 1 and 2 (n=711).

Scored ≥ 24 on the Mini-Mental State Exam (Folstein et al., 1975), as lower scores are often considered to be a marker of cognitive impairment (n=701).

Had TSH levels within the range of 0.2 to 4.5 mU/l at both waves 1 and 2 (n=660).

Had T₄ levels with the range of 9 to 21 pmol/L waves 1 and 2 (n=659).

Applying these criteria resulted in a final sample of 659 (male=371; female=288; Mean age Wave 1=69.5 years, SD=0.8 years; Mean age wave 2=72.5 years, SD=0.7 years) participants for the current analyses.

2.2 Ethical Considerations

Ethical permission for the LBC1936 study protocol was obtained from the Multi-Centre Research Ethics Committee for Scotland and the Lothian Research Ethics Committee. All research was carried out in compliance with the Helsinki Declaration.

2.3 TSH and T₄

Blood samples were taken at both waves 1 and 2. All analyses were conducted in the same laboratory (NHS Lothian –Clinical Biochemistry & Haematology, Western General Hospital, UK) using a two-step immunoassay. Target precision of both the TSH and T₄ assays were coefficient of variability <10%.

For TSH, the laboratory reference range was 0.2 to 4.5 mU/l, with coefficients of variability ranging from 3.0 to 3.5%. For T₄, the laboratory reference range was 9 to 21 pmol/L, with coefficients of variability ranging from 5.1 to 8.9%.

In the current study we use the laboratory reference range as a selection criterion. This reference range has been used in other published work on different samples (Patterson, Lonie and Starr, 2010).

2.4 Cognitive Ability Tests

In the current study we used 14 cognitive ability subscale scores from 12 individual cognitive tests. General cognitive ability (g) was measured using 6 subtest scores of the WAIS-III^{UK} (Wechsler, 1998a; Block Design, Matrix Reasoning, Digit Span Backward, Digit Symbol, Symbol Search and Letter-Number Sequencing). Processing Speed was measured using two timed WAIS-III^{UK} subtest scores (Digit Symbol and Symbol Search), a simple and choice reaction time task, and an inspection time task. Lastly, Memory was measured using five subtest scores from the WMS-III^{UK} (Wechsler, 1998b; Logical Memory Immediate and delayed recall, Verbal Paired Associates immediate and delayed recall and Spatial Span total score) and two scores from the WAIS-III^{UK}, Digit Span Backward and Letter-Number Sequencing.

2.5 Statistical Analysis

Means, standard deviations, skew and kurtosis were computed for all raw data. Pearson's correlations between individual subtest scores, TSH and T₄ at both waves 1 and 2 were computed. Next, we used structural equation modelling (SEM) to estimate the cross-lagged correlations between three latent cognitive ability variables (*g*, processing speed, and memory) and TSH and T₄ measures at both waves 1 and 2. The latent factors represent the shared associations between individual cognitive ability subtest scores hypothesized to measure the same major domain of cognitive ability. SEM provides highly reliable estimates of broad cognitive domains by taking account of measurement error present in each subtest. All models were estimated in Mplus 6.0 using full information maximum likelihood estimation. Full details of the structural models can be found in Supplementary Material A. Lastly, we produced scatterplots with fitted loess curves of latent variable estimates in order to consider non-linear effects (see Supplementary Material B).

2.6 Covariates

All models were estimated both with and without the inclusion of health covariates. Participants' smoking status (former/current/never), alcohol consumption (drink/not drink), history (yes/no) of diabetes, hypertension, high cholesterol, cardiovascular disease, circulatory problems and stroke, were gathered from a medical interview. In addition we included the depression subscale score from the self-report Hospital Anxiety and Depression scale (HADS; Zigmond and Snaith, 1983).

In order to assess potential effects of BMI and medications known to effect thyroid hormone levels we re-ran all models excluding firstly participants with BMI less than 20 (*n*=18), and secondly those taking any of the following medications; hydrocortisone, prednisolone, growth hormone, octreotide, somatostatin, morphine, dihydrocodeine, tramadol, dopamine, L-dopa, madopar, sinemet, pimozide, phenotolamine, thioridazine,

methysergide, cyproheptadine, iodine, lithium, ropinerole, pramipexole, cabergoline, iodide or amiodarone (n=24).

3. Results

3.1 Descriptive Statistics

Full descriptive statistics are presented in Table 1. All variables were approximately normally distributed with no values for skew outside of ± 1.99 . Across the waves, there were significant mean level decreases in both TSH ($t=10.99$, $p<0.001$) and T₄ ($t=34.55$, $p<0.001$).

(Insert Table 1 about here)

Univariate correlations between individual cognitive subtests and thyroid hormones at waves 1 and 2 yielded few significant results (for full table see Supplementary Material C). Digit Span Backwards (-0.09), Letter-Number Sequencing (-0.08) and Simple Reaction Time (0.12) were significantly associated with TSH levels at wave 1, but not at wave 2.

Conversely, Block Design (-0.10) and Simple Reaction Time (0.10) were significantly associated with T₄ levels at wave 2, but not wave 1. In totality, few systematic associations were present in the current sample.

Estimates of the associations between TSH or T₄ levels and g, processing speed and memory derived from cross-lagged latent variable structural models are presented in Table 2. All structural models showed excellent fit to the data (see Supplementary Material A). No significant associations were found between either TSH or T₄ and any of the cognitive measures.

(Insert Table 2 about here)

All changes to parameter estimates removing cases with low BMI were at the second decimal place, with all values remaining non-significant. Similarly, all changes in parameter estimates were at the second decimal place in the models excluding participants currently taking potentially confounding medications (see Supplementary Material D for raw results). In this

model, the association between T₄ at wave 1 and Memory at wave 2 was significant (0.10, $p < 0.05$). However, given the large number of parameters estimated we suggest this is simply a Type 1 error.

Scatterplots were produced plotting the latent factor scores of each cognitive ability factor against TSH and T₄ at both waves. The scatterplots provided no indication of non-linear associations (see Supplementary Material B).

4. Discussion

In the current euthyroid sample, we found no significant associations between thyroid function as measured by TSH and T₄ and three major domains of cognitive ability (g, memory and processing speed). Further, we found no evidence of non-linear associations. These conclusions hold true both with and without controlling for a large number of health covariates, as well as when analyses were re-run excluding those on medications and with low BMI. Notably, both TSH and T₄ levels decreased over the three year follow-up period consistent with a fall in thyrotropin-releasing hormone (TRH), although we did not measure this directly. Although there are several cross-sectional studies that have found lower TRH levels in older participants, there is a paucity of longitudinal data (Leitol et al., 2002).

These findings support suggestions from previous studies in which the authors have suggested that thyroid-cognition associations may only be present at clinical levels of thyroid dysfunction (Begin et al. 2008; St. John et al. 2009). The association between thyroid hormone levels and memory ability was identified as perhaps the most consistent finding from past research (e.g. Begin et al. 2008; Hogervorst et al. 2008). Again in the current study, we found no support for this association.

However, the current study offers a number of important extensions to prior studies. Firstly, we use a multiple cognitive tests to identify latent cognitive ability factors of g, processing speed, and memory, and thus, provide robust estimates from structural equation models.

Secondly, we were able to estimate the associations in the same large sample (n=659) at baseline and with an approximate three year follow up. Given this sample size, the current study was powered to identify correlations of approximately 0.11 (n=659; 80% power; $p < .05$), which is a small effect size. As a result, we consider the current findings to be highly robust.

Lastly, the cohort design and narrow age range of the current sample provided almost eliminated the effect of chronological age, which otherwise could have been a major confounder of any effects of thyroid hormone levels on cognition. A number of authors have suggested that the major impact of thyroid levels on cognition occurs above a key threshold of ~80 years of age (van den Beld et al., 2005; Begin et al., 2008). If such a threshold exists, then the current sample would be too young for the major effects of thyroid function to be impacting on cognitive performance. However, given the consistency of the current results with prior findings, it is suggested they are indicative of a lack of association within the normal ranges of TSH and T₄.

References

- Begin, M.E., Langlois, M.F., Lorrain, D., Cunnane, S.C., 2008. Thyroid Function and Cognition during Aging. *Curr Gerontol Geriatr Res*. doi:10.1155/2008/474868.
- Beydoun, M.A., Beydoun, H.A., Shroff, M.R., Kitner-Triolo, M.H., Zonderman, A.B. 2012. Serum leptin, thyroxine and thyroid-stimulating hormone levels interact to affect cognitive function among US adults: evidence from a large representative survey. *Neurobiol Aging*, 33, 1730-1743.
- Deary, I.J., 2012. Human Intelligence. *Annu Rev Psychol*, 63, 453-482.
- Deary, I.J., Gow, A.J., Taylor, M.D., et al., 2007. The Lothian birth cohort 1936: a study to examine influences on cognitive ageing from age 11 to age 70 and beyond. *BMC Geriatr*, Dec. 5; 7:28. doi: 10.1186/1471-2318-7-28
- Deary, I.J., Gow, A.J., Pattie, A., Starr, J.M., 2011. Cohort profile: The Lothian Birth Cohorts of 1921 and 1936. *Int J Epidemiol*. doi:10.1093/ije/dyr197
- Folstein, M.F., Folstein, S.E., McHugh, P.R., 1975. Mini-mental state. A practical method for grading cognitive state of patients for clinicians. *J Psychiatr Res*. 12, 189-198.
- Gussekloo, J., van Exel, E., de Craen, A.J.M., et al. 2004. Thyroid Status, Disability and Cognitive Function, and Survival in Old Age. *JAMA*, 292, 2591-2599.
- Hogervorst, E., Huppert, F., Matthews, F.E., Brayne, C. 2008. Thyroid function and cognitive decline in the MRC Cognitive Function and Ageing Study. *Psychoneuroendocrinology*, 33, 1013-1022.
- Leitol, H., Behrends, J., Brabant, G., 2002. The thyroid axis in ageing. *Novartis Found Symp*, 242, 193-201
- Liesbeth, A.D.M., van Osch, Horgervorst, E., Combrinck, M., Smith, A.D. 2004. Low thyroid-stimulating hormone as an independent risk factor for Alzheimer disease. *Neurology*, 62, 1967-1971.

- Parle, J., Roberts, L., Wilson, S., et al. 2010. A Randomized Controlled Trial of the Effect of Thyroxine Replacement on Cognitive Function in Community-Living Elderly Subjects with Subclinical Hypothyroidism: The Birmingham Elderly Thyroid Study. *J Clin Endocrinol Metab*, 95, 3623-3632.
- Patterson, M., Lonie, J.A., Starr, J.M. 2010. Thyroid function, cognition, functional independence and behavioural and psychological symptoms of dementia in Alzheimers disease. *Int J Geriatr Psychiatry*, 25:1196-1199
- Roberts, L.M., Pattison, H., Roalfe, A., et al. 2006. Is subclinical thyroid dysfunction in the elderly associated with depression or cognitive dysfunction? *Ann Intern Med*. 145, 573-581.
- Samuels, M.H. 2010. Cognitive Function in Subclinical Hypothyroidism. *J Clin Endocrinol Metab*. 95, 3611-3613.
- Smith, J.W., Evans, A.T., Costall, B., Smythe, J.W. 2002. Thyroid hormones, brain function and cognition: a brief review. *Neurosci Biobehav Rev*. 26, 45-60.
- St. John, J.A., Henderson, V.W., Gatto, N.M., McCleary, C.A., Spencer, C.A., Hodis, H.N., Mack, W.J. 2009. Mildly Elevated TSH and Cognition in Middle-Aged and Older Adults. *Thyroid*, 19, 111-117.
- van den Beld, A.W., Visser, T.J., Feelders, R.A., Grobbee, D.E., Lamberts, S.W.J. 2005. Thyroid hormone concentrations, disease, physical function, and mortality in elderly men. *J Clin Endocrinol Metab*. 90, 6403-6409.
- Wechsler D. WAIS-III^{UK} administration and scoring manual. London: Psychological Corporation; 1998a.
- Wechsler D. WMS-III^{UK} administration and scoring manual. London: Psychological Corporation; 1998b.
- Zigmond, A.S. & Snaith, R.P. 1983. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand*, 67, 361-370.

Table 1: Means, Standard Deviations, Skew and Kurtosis for all variables

		Wave 1				Wave 2			
		Mean	SD	Skew	Kurt.	Mean	SD	Skew	Kurt.
<i>WAIS-III^{UK}</i>									
	Digit Symbol	57.87	12.83	0.10	-0.02	56.74	12.17	0.08	-0.19
	Digit Span Backwards	7.86	2.28	0.45	-0.24	7.86	2.23	0.33	-0.09
	Block Design	34.87	10.09	0.19	-0.21	34.15	10.14	0.43	0.09
	Letter-Number Sequencing	11.15	3.06	0.12	-0.07	11.02	3.02	0.35	0.45
	Matrix Reasoning	14.02	5.05	-0.13	-0.96	13.36	5.00	-0.10	-1.02
	Symbol Search	25.23	6.33	-0.04	0.86	24.97	5.95	-0.32	0.90
<i>WMS-III^{UK}</i>									
	Logical Memory Immediate	45.05	9.90	-0.34	-0.11	46.12	10.06	-0.38	0.14
	Logical Memory Delayed	28.17	7.63	-0.36	-0.07	29.17	7.76	-0.53	0.30
	Verbal Paired Associates Immediate	2.60	2.17	0.82	-0.17	2.77	2.23	0.65	-0.55
	Verbal Paired Associates Delayed	6.31	1.93	-1.03	0.18	6.39	2.02	-1.33	0.99
	Spatial Span	14.88	2.82	0.05	-0.23	14.77	2.70	-0.10	-0.03
<i>Speed Tests</i>									
	Simple Reaction Time	0.27	0.05	1.99	6.57	0.27	0.05	1.50	3.06
	Choice Reaction Time	0.63	0.08	1.02	2.81	0.65	0.09	0.82	1.35
	Inspection Time	112.99	11.23	-0.90	2.70	111.91	11.76	-1.06	2.58
<i>Thyroid Hormones</i>									
	TSH	1.82	0.88	0.82	0.26	1.57	0.76	1.02	1.18
	T ₄	15.29	2.04	0.12	0.05	12.65	1.58	0.22	-0.12

Table 2:

Cross-lagged Correlations between Cognitive Ability Latent Variables, TSH and T₄

	1	2	3	4	5	6	7	8	9	10
1. TSH Wave 1	-	0.76	-	-	-0.06	-0.06	-0.05	-0.07	-0.08	-0.07
2. TSH Wave 2	0.76	-	-	-	-0.02	0.00	-0.04	-0.06	-0.03	-0.01
3. T ₄ Wave 1	-	-	-	0.47	0.02	0.01	0.05	0.05	-0.01	0.07
4. T ₄ Wave 2	-	-	0.43	-	-0.01	-0.01	-0.05	-0.02	0.04	0.02
5. g Wave 1	-0.06	-0.04	0.07	-0.05	-	0.99	-	-	-	-
6. g Wave 2	-0.05	-0.01	0.04	-0.06	0.98	-	-	-	-	-
7. Processing Speed Wave 1	-0.05	-0.06	0.05	0.05	-	-	-	0.98	-	-
8. Processing Speed Wave 2	-0.07	-0.07	-0.04	-0.03	-	-	0.98	-	-	-
9. Memory Wave 1	-0.07	-0.05	0.04	0.02	-	-	-	-	-	0.89
10. Memory Wave 2	-0.07	-0.03	0.09	0.02	-	-	-	-	0.86	-

Note: All estimates shown in bold type are the wave 1 to wave 2 stability coefficients, all significant at $p < 0.001$. All other estimates are the associations between TSH, T₄ and cognition, all non-significant ($p > 0.05$). Estimates below the diagonal are uncorrected. Estimates above the diagonal are corrected for sex, age, health variables and depression scores.

Supplementary Material A

Model Specification

The basic cross-lagged model is shown in Supplementary Figure 1. The cross-lagged model comprises six primary correlations, namely, two stability coefficients (Paths A), two contemporaneous associations within time points (Paths B) and two cross-lagged associations across time points (T1/T2; Paths C). In the current models, we modelled latent estimates for the cognitive ability factors at both waves, loaded by individual sub test scores (Paths D). In addition, we included correlation residuals (Paths E) between sub-test scores from the same testing procedure (e.g. logical memory immediate and delayed recall), and between the same sub-tests across waves 1 and 2.

Figure S1:

Cross-lagged Model Design

Model Fit

Model fit was assessed based on standard cut-off criteria (Hu & Bentler, 1998; 1999; Schermelleh-Engel, Moosbrugger, & Muller, 2003) for a number of commonly used model fit indices. We adopted cut-off's of ≥ 0.95 for the comparative fit index (CFI) and Tucker-Lewis index (TLI), values ≤ 0.06 for the root-mean-square error of approximation (RMSEA) and finally values of ≤ 0.05 for the standardised root mean square residual (SRMR). As can be seen from the results in Table S2, all models met the criteria for model fit, indicating that the parameters in the model represent a good fit to the data and can be reliably interpreted.

Table S2: Model fit statistics for the 12 cross-lagged models.

	χ^2	df	Sig.	RMSEA	SRMR	CFI	TLI
g & TSH	83.88	55	$p<0.05$	0.028	0.024	0.99	0.99
g & T ₄	73.63	55	$p=0.05$	0.023	0.023	1.00	0.99
g & TSH with covariates	438.85	305	$p<0.05$	0.026	0.034	0.97	0.97
g & T ₄ with covariates	422.38	305	$p<0.05$	0.024	0.033	0.98	0.97
Processing Speed & TSH	70.25	41	$p<0.05$	0.033	0.029	0.99	0.99
Processing Speed & T ₄	62.47	41	$p<0.05$	0.028	0.026	0.99	0.99
Processing Speed & TSH with covariates	332.78	225	$p<0.05$	0.027	0.035	0.97	0.97
Processing Speed & T ₄ with covariates	313.65	225	$p<0.05$	0.025	0.033	0.98	0.97
Memory & TSH	200.53	81	$p<0.05$	0.047	0.052	0.98	0.97
Memory & T ₄	181.33	81	$p<0.05$	0.043	0.048	0.98	0.97
Memory & TSH with covariates	576.00	373	$p<0.05$	0.029	0.037	0.97	0.96
Memory & T ₄ with covariates	550.06	373	$p<0.05$	0.027	0.036	0.97	0.96

Model Parameter Estimates

Tables S3, S4 and S5 display the unstandardized and standardized parameter estimates for the two wave measurement models for each of cognitive ability latent constructs. In all cases the estimates were taken from the TSH models without covariates. Measurement parameters showed negligible changes (second decimal place) in both the T₄ models, with and without covariates.

In all tables, the following abbreviations are used; <-> = Correlation; W1 = Wave 1; W2= Wave 2; DS = Digit Symbol; SS = Symbol Search; MR = Matrix Reasoning; BD = Block Design; LN = Letter-Number Sequencing; DB = Digit Span Backward; CRT = Choice Reaction Time Mean Score; SRT = Simple Reaction Time Mean Score; IT = Inspection Time Total Score; LM1 = Logical Memory Immediate Recall; LM2 = Logical Memory Delayed Recall; VPA1 = Verbal Paired Associated Immediate Recall; VPA2 = Verbal Paired Associates Delayed Recall; SP = Spatial Span Total Score.

Table S3: Parameter Estimates for the g Measurement Model

	Unstandardized	SE	Standardized	SE	Sig
<i>Wave 1 g Loadings</i>					
DS_W1	7.368	0.568	0.574	0.037	$p<0.001$
SS_W1	4.271	0.266	0.675	0.032	$p<0.001$
MR_W1	3.19	0.222	0.632	0.036	$p<0.001$
BD_W1	6.685	0.429	0.664	0.033	$p<0.001$
LN_W1	1.821	0.129	0.592	0.034	$p<0.001$
DB_W1	1.136	0.098	0.498	0.037	$p<0.001$
<i>Correlated Residuals – Wave 1 Similar Tests</i>					
DS_W1 <-> SS_W1	17.147	3.032	0.349	0.046	$p<0.001$
MR_W1 <-> BD_W1	6.361	1.872	0.216	0.054	$p<0.001$
LN_W1 <-> DB_W1	1.543	0.247	0.315	0.041	$p<0.001$
<i>Wave 2 g Loadings</i>					
DS_W2	7.390	0.535	0.606	0.036	$p<0.001$
SS_W2	3.835	0.254	0.644	0.034	$p<0.001$
MR_W2	2.885	0.222	0.578	0.037	$p<0.001$
BD_W2	6.601	0.435	0.652	0.034	$p<0.001$
LN_W2	1.699	0.128	0.562	0.035	$p<0.001$
DB_W2	1.041	0.097	0.466	0.038	$p<0.001$
<i>Correlated Residuals – Wave 2 Similar Tests</i>					
DS_W2 <-> SS_W2	15.793	2.751	0.358	0.045	$p<0.001$
MR_W2 <-> BD_W2	6.41	1.847	0.205	0.051	$p<0.001$
LN_W2 <-> DB_W2	1.814	0.247	0.367	0.038	$p<0.001$
<i>Correlated Residuals – Same Tests Across Waves</i>					
DS_W1 <-> DS_W2	79.846	6.818	0.784	0.018	$p<0.001$
MR_W1 <-> MR_W2	7.041	0.984	0.441	0.04	$p<0.001$
LN_W1 <-> LN_W2	2.809	0.336	0.453	0.036	$p<0.001$
SS_W1 <-> SS_W2	8.636	1.4	0.406	0.045	$p<0.001$
BD_W1 <-> BD_W2	32.98	4.013	0.571	0.035	$p<0.001$
DB_W1 <-> DB_W2	2.066	0.199	0.53	0.031	$p<0.001$
<i>Correlates Residuals – Similar Tests Across Waves</i>					
DS_W1 <-> SS_W2	16.421	2.851	0.343	0.045	$p<0.001$
SS_W1 <-> DS_W2	14.061	2.838	0.311	0.048	$p<0.001$
MR_W1 <-> BD_W2	4.39	1.835	0.146	0.055	$p<0.01$
BD_W1 <-> MR_W2	8.66	1.837	0.282	0.048	$p<0.001$
LN_W1 <-> DB_W2	1.215	0.239	0.248	0.042	$p<0.001$
DB_W1 <-> LN_W2	1.509	0.243	0.306	0.04	$p<0.001$

Table S4: Parameter Estimates for the Processing Speed Measurement Model

	Unstandardized	SE	Standardized	SE	Sig
<i>Wave 1 Speed Loadings</i>					
DS_W1	-10.21	0.467	-0.795	0.022	$p<0.001$
SS_W1	-4.64	0.236	-0.733	0.025	$p<0.001$
CRT_W1	0.053	0.003	0.639	0.028	$p<0.001$
SRT_W1	0.016	0.002	0.323	0.039	$p<0.001$
IT_W1	-4.722	0.474	-0.42	0.037	$p<0.001$
<i>Correlated Residuals – Wave 1 Similar Tests</i>					
CRT_W1 <-> SRT_W1	0.001	0.000	0.299	0.039	$p<0.001$
<i>Wave 2 Speed Loadings</i>					
DS_W2	9.899	0.439	0.81	0.021	$p<0.001$
SS_W2	4.45	0.222	0.745	0.024	$p<0.001$
CRT_W2	-0.056	0.003	-0.651	0.027	$p<0.001$
SRT_W2	-0.015	0.002	-0.31	0.04	$p<0.001$
IT_W2	6.212	0.474	0.528	0.033	$p<0.001$
<i>Correlated Residuals – Wave 2 Similar Tests</i>					
CRT_W2 <-> SRT_W2	0.001	0.000	0.375	0.036	$p<0.001$
<i>Correlated Residuals – Same Tests Across Waves</i>					
DS_W1 <-> DS_W2	34.946	4.566	0.627	0.037	$p<0.001$
SS_W1 <-> SS_W2	4.685	1.037	0.273	0.049	$p<0.001$
CRT_W1 <-> CRT_W2	0.002	0.000	0.59	0.029	$p<0.001$
SRT_W1 <-> SRT_W2	0.001	0.000	0.472	0.031	$p<0.001$
IT_W1 <-> IT_W2	56.46	4.958	0.553	0.029	$p<0.001$
<i>Correlates Residuals – Similar Tests Across Waves</i>					
SRT_W1 <-> CRT_W2	0.001	0.000	0.201	0.041	$p<0.001$
CRT_W1 <-> SRT_W2	0.001	0.000	0.251	0.04	$p<0.001$

Table S5: Parameter Estimates for the Memory Measurement Model

	Unstandardized	SE	Standardized	SE	Sig
<i>Wave 1 Memory Loadings</i>					
LM1_W1	6.122	0.476	0.616	0.041	$p<0.001$
LM2_W1	4.998	0.36	0.653	0.039	$p<0.001$
VPA1_W1	1.031	0.101	0.473	0.041	$p<0.001$
VPA2_W1	1.151	0.088	0.589	0.038	$p<0.001$
LN_W1	1.572	0.149	0.516	0.044	$p<0.001$
SP_W1	1.012	0.135	0.358	0.045	$p<0.001$
DB_W1	0.914	0.109	0.405	0.045	$p<0.001$
<i>Correlated Residuals – Wave 1 Similar Tests</i>					
LM1_W1 <-> LM2_W1	35.253	3.784	0.776	0.022	$p<0.001$
VPA1_W1 <-> VPA2_W1	1.259	0.172	0.415	0.04	$p<0.001$
LN_W1 <-> DB_W1	2.043	0.29	0.379	0.04	$p<0.001$
<i>Wave 2 Memory Loadings</i>					
LM1_W2	7.49	0.458	0.743	0.036	$p<0.001$
LM2_W2	5.802	0.354	0.746	0.036	$p<0.001$
VPA1_W2	1.136	0.099	0.508	0.038	$p<0.001$
VPA2_W2	1.214	0.089	0.595	0.036	$p<0.001$
LN_W2	1.473	0.141	0.49	0.042	$p<0.001$
SP_W2	0.943	0.125	0.349	0.044	$p<0.001$
DB_W2	0.829	0.103	0.374	0.043	$p<0.001$
<i>Correlated Residuals – Wave 1 Similar Tests</i>					
LM1_W2 <-> LM2_W2	24.68	3.979	0.707	0.036	$p<0.001$
VPA1_W2 <-> VPA2_W2	1.217	0.172	0.385	0.04	$p<0.001$
LN_W2 <-> DB_W2	2.278	0.273	0.422	0.036	$p<0.001$
<i>Correlated Residuals – Same Tests Across Waves</i>					
LM1_W1 <-> LM1_W2	27.366	4.216	0.518	0.043	$p<0.001$
LM2_W1 <-> LM2_W2	14.669	2.497	0.489	0.046	$p<0.001$
VPA1_W1 <-> VPA1_W2	1.933	0.192	0.522	0.032	$p<0.001$
VPA2_W1 <-> VPA2_W2	1.425	0.154	0.55	0.034	$p<0.001$
LN_W1 <-> LN_W2	3.657	0.385	0.534	0.032	$p<0.001$
SP_W1 <-> SP_W2	3.421	0.32	0.512	0.03	$p<0.001$
DB_W1 <-> DB_W2	2.443	0.215	0.575	0.028	$p<0.001$
<i>Correlates Residuals – Similar Tests Across Waves</i>					
LM1_W1 <-> LM2_W2	18.166	3.203	0.449	0.048	$p<0.001$
LM2_W1 <-> LM1_W2	17.598	3.2	0.449	0.049	$p<0.001$
VPA1_W1 <-> VPA2_W2	0.88	0.16	0.279	0.043	$p<0.001$
VPA2_W1 <-> VPA1_W2	1.006	0.159	0.33	0.042	$p<0.001$
LN_W1 <-> DB_W2	1.795	0.264	0.334	0.039	$p<0.001$
DB_W1 <-> LN_W2	2.062	0.271	0.381	0.038	$p<0.001$

Supplementary Material B

In order to investigate whether the lack of significant associations was due to the linearity assumptions of correlation coefficients, we produced scatter-plots of the latent factor scores for the cognitive ability variables derived from the structural equation models, against TSH and T₄ levels at both waves 1 and 2 (Total 24 plots). Next we fitted loess curves to the scatterplots. Loess curves are generated by locally fitting points along the x-axis, with the curve created with these points are joined (Jacoby, 1997).

For illustrative purposes, figures S2 to S25 display all plots. As can be seen from these plots, no consistent and significant non-linearity is present in the current data.

Figure S2:

Figure S3:

Figure S4:

Figure S5:

Figure S6:

Figure S7:

Figure S8:

Figure S9:

Figure S10:

Figure S11:

Figure S12:

Figure S13:

Figure S14:

Figure S15:

Figure S16:

Figure S17:

Figure S18:

Figure S19:

Figure S20:

Figure S21:

Figure S22:

Figure S23:

Figure S24:

Figure S25:

Supplementary Material C

Table S6:

Pearsons Correlations between TSH, T₄ and Individual Cognitive Ability Subscale Scores.

		Wave 1				Wave 2			
		Raw		Partial		Raw		Partial	
		TSH	T ₄	TSH	T ₄	TSH	T ₄	TSH	T ₄
<i>WAIS-III^{UK}</i>									
	Digit Symbol	-0.03	0.07	-0.03	-0.03	-0.06	0.00	-0.03	0.00
	Digit Span Backwards	-0.09*	0.01	-0.10*	-0.01	0.05	0.02	0.06	0.03
	Block Design	0.02	0.06	0.01	0.04	0.02	-0.10*	0.02	-0.04
	Letter-Number Sequencing	-0.08*	0.04	-0.09*	0.02	0.01	0.02	0.01	0.03
	Matrix Reasoning	-0.03	0.05	-0.03	0.03	0.00	-0.04	0.00	0.01
	Symbol Search	-0.03	0.05	-0.03	-0.01	-0.05	-0.02	-0.04	0.01
<i>WMS-III^{UK}</i>									
	Logical Memory Immediate	-0.06	0.03	-0.06	-0.01	-0.07	0.01	-0.07	0.01
	Logical Memory Delayed	-0.06	0.01	-0.06	-0.03	-0.05	0.03	-0.04	0.03
	Verbal Paired Associates Immediate	0.07	-0.01	0.08	-0.02	0.06	0.03	0.07	0.02
	Verbal Paired Associates Delayed	0.05	0.03	0.05	-0.01	0.04	0.01	0.06	-0.01
	Spatial Span	-0.07	0.01	-0.07	-0.03	-0.03	-0.05	-0.03	-0.01
<i>Speed Tests</i>									
	Simple Reaction Time	0.12**	-0.01	0.13**	0.00	0.07	0.10*	0.07	0.08*
	Choice Reaction Time	0.02	0.03	0.03	0.05	0.07	0.03	0.06	0.01
	Inspection Time	-0.06	0.03	-0.06	0.01	-0.05	-0.04	-0.05	0.00

* $p < 0.05$; ** $p < 0.01$; Raw correlations are uncorrected. Partial correlations are controlling for age, sex, HADS score and health covariates

Supplementary Material D

Table S7:

Cross-lagged Correlations between Cognitive Ability Latent Variables, TSH and T₄ in Split Samples by BMI and Drug Use

	1	2	3	4	5	6	7	8	9	10
1. TSH Wave 1	-	0.76***	-	-	-0.07	-0.06	0.07	0.09	-0.09	-0.09
2. TSH Wave 2	0.77***	-	-	-	-0.04	-0.01	0.08	0.08	-0.05	-0.02
3. T ₄ Wave 1	-	-	-	0.43***	0.08	0.05	-0.05	-0.06	0.05	0.10*
4. T ₄ Wave 2	-	-	0.43***	-	-0.05	-0.05	0.04	0.02	0.03	0.04
5. g Wave 1	-0.06	-0.03	0.07	-0.05	-	0.98***	-	-	-	-
6. g Wave 2	-0.06	0.00	0.04	-0.06	0.98***	-	-	-	-	-
7. Processing Speed Wave 1	0.06	0.06	-0.04	0.04	-	-	-	0.98***	-	-
8. Processing Speed Wave 2	0.07	0.07	-0.04	0.02	-	-	0.98***	-	-	-
9. Memory Wave 1	-0.07	-0.02	0.03	0.01	-	-	-	-	-	0.86***
10. Memory Wave 2	-0.07	-0.01	0.07	0.00	-	-	-	-	0.87***	-

Note: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. All estimates shown in bold type are the wave 1 to wave 2 stability coefficients. Estimates below the diagonal are uncorrected associations having removed cases with BMI < 20. Estimates above the diagonal are uncorrected associations having removed cases taking a number of drugs known to influence thyroid hormone levels.

References

- Hu, L.T., Bentler, P.M. 1998. Fit indices in covariance structure modelling: Sensitivity to underparameterized model misspecification. *Psychol Methods*, 3, 424–453.
- Hu, L.T., Bentler, P.M. 1999. Cut-off criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Struct Equ Modelling*, 6, 1–55.
- Jacoby, W.G. 1997. *Statistical Graphics for Univariate and Bivariate Data*. Thousand Oaks, CA: Sage Publications, Inc.
- Schermelleh-Engel, K., Moosbrugger, H., Muller, H. 2003. Evaluating the Fit of Structural Equation Models: Tests of Significance and Descriptive Goodness-of-Fit Measures. *Methods of Psychological Research*, 8, 23-74.