THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Evidence from Late Cretaceous-Paleogene volcanic rocks of the Kyrenia Range, northern Cyprus for the northern, active continental margin of the Southern Neotethys

Citation for published version:

Chen, G & Robertson, AHF 2021, 'Evidence from Late Cretaceous-Paleogene volcanic rocks of the Kyrenia Range, northern Cyprus for the northern, active continental margin of the Southern Neotethys', *Lithos*, vol. 380-381, pp. 105835. <https://doi.org/10.1016/j.lithos.2020.105835>

Digital Object Identifier (DOI):

[10.1016/j.lithos.2020.105835](https://doi.org/10.1016/j.lithos.2020.105835)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Lithos

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

LITHOS

Evidence from Late Cretaceous-Paleogene volcanic rocks of the Kyrenia Range, northern Cyprus for the northern, active continental margin of the Southern Neotethys --Manuscript Draft--

Manuscript Number:	LITHOS9043R2
Article Type:	Regular Article
Keywords:	Late Cretaceous-Paleogene; Volcanic rocks; Kyrenia Range; Arc magmatism; Marginal basin formation; Southern Neotethys
Corresponding Author:	Guohui Chen Institute of Geology and Geophys, Chinese Academy of Sciences Beijing, CHINA
First Author:	Guohui Chen
Order of Authors:	Guohui Chen Alastair Robertson
Abstract:	<p>Late Cretaceous felsic and latest Cretaceous-Paleogene basaltic volcanic rocks are exposed throughout the Kyrenia Range, N Cyprus. Field mapping of the key, well-exposed western range indicates that the felsic volcanics are mainly crop out in the southerly, frontal part of the range, separated from the latest Cretaceous-Paleogene basaltic volcanics farther north by a thrust. New U-Pb zircon dating of the felsic volcanics indicates a primary age of c. 74.0 Ma (Late Campanian). The felsic volcanics are characterised by evolved compositions that were probably generated in an extensional subduction-related setting. Their relatively low Nb and Y but high Rb concentrations, together with characteristic Th/Ta ratios (6-20), suggest a mature continental arc setting. The latest Cretaceous-Paleogene basaltic volcanics have mainly within-plate chemical characteristics (in the east), although some have a subordinate subduction influence; e.g. negative Nb (in the west). Sr-Nd-Hf isotopic signatures (i.e. high positive $\epsilon\text{Nd}(t)$ and $\epsilon\text{Hf}(t)$ values) suggest derivation of the basalts from several OIB-like mantle sources, with probable involvement of a crustal (recycled) component (i.e. elevated Nb/Y ratios). Comparisons with SE Turkey, where Late Cretaceous arc-related granitic rocks are widely developed, suggest that the Kyrenia Range igneous rocks may have originated to the south of the Alanya metamorphic massif (N of Cyprus) and correlative continental units in SE Turkey. The latest Cretaceous-Paleogene basaltic volcanics in N Cyprus and SE Turkey (Maden Complex) are interpreted to represent incipient marginal basin formation, possibly in an oblique-convergent setting, prior to Miocene suturing with Arabia. In the light of alternatives, we infer genesis of the N Cyprus Late Cretaceous and Paleogene volcanic rocks related to stages in the development of the northerly active continental margin of the Southern Neotethys.</p>
Suggested Reviewers:	<p>Fatih Karaoğlu fkaraoglan@cu.edu.tr expert on arc magmatism in SE Turkey</p> <p>Osman Parlak parlak@cu.edu.tr expert on magmatism and tectonics especially in Turkey</p> <p>Pamela Kempton pkempton@ksu.edu</p>
Opposed Reviewers:	

**Evidence from Late Cretaceous-Paleogene volcanic rocks of the Kyrenia Range,
northern Cyprus for the northern, active continental margin of the Southern
Neotethys**

Guohui Chen^{1*}, Alastair H. F. Robertson¹

¹School of GeoSciences, University of Edinburgh, West Mains Road, Edinburgh EH9
3JW, UK

*Current Address: State Key Laboratory of Lithospheric Evolution, Institute of Geology
and Geophysics, Chinese Academy of Sciences, P.O. Box 9825, Beijing 100029,
China

Corresponding author: Guohui Chen (Guohui.Chen@live.cn)

Revision notes:

Dear Professor Shellnutt,

Thank you for considering the further revision of our paper. We have acted on the comments made. We have marked the main edits using track changes. Please note that we not made any significant changes to the results or discussion, only refinements in the light of the review comments.

The main further changes of the paper are:

- 1) The field descriptions are reorganized somewhat, with local description first, and then a summary;*
- 2) The section on the Late Cretaceous magmatic rocks in SE Turkey are revised beginning with units in the south, then the units in the north. More consideration is given to possible equivalents in the Kyrenia Range volcanics (see new section 5.4);*
- 3) More information is given on the HP-LT metamorphism and shoshonites related to Late Cretaceous subduction and possible collision in SE Turkey (lines 492-497) as this is relevant to the regional setting;*
- 4) The Eocene volcanics are discussed in slightly more detail, as suggested (lines 585-593, 653-664).*

In summary, we hope that you will decide that our paper is now suitable for publication in Lithos.

*Yours sincerely,
Guohui Chen and Alastair Robertson*

COMMENTS FROM EDITORS AND REVIEWERS

Editor: I requested the previous reviewers of your manuscript to evaluate the revised version however, only Reviewer #2 accepted the invitation. As you will see, Reviewer #2 is mostly satisfied with the improvements but, points out a few minor issues in the discussion that should be addressed. Furthermore, I think it is appropriate to move section 2 after section 3. It is common to have the methods after the 'geological background'. Please update the manuscript accordingly.

All the best.
Greg Shellnutt
Co-Editor-in-Chief

Comments

Line 73. The methods should come after the geological background... or in this case the "Volcanism in the Kyrenia Range.

OK. It is reordered now.

Line 75. Delete 'initially'

OK. It is deleted now.

Line 100. Should it be "Whole rock procedural blanks"?

No. It refers the total procedural blanks. So it is correct;

Line 208. Should be either "... and a felsic debris-flow deposit are exposed." Or "... and felsic debris-flow deposits are exposed."

OK. It is revised to be "a felsic debris-flow deposit are exposed" now.

Lines 534-535. Should be, "... arc was explained by northward..."

Yes. Revised as suggested.

Line 565. Should be, "... work is needed..."

Line 670. Should be, "... alternatives are suggested; ..."

OK. They are revised now.

Reviewer #2: Dear Editor and authors,

I review the revised MS entitled "Evidence from Late Cretaceous-Paleogene volcanic rocks of the Kyrenia Range, Northern Cyprus for northern, active continental margin of the Southern Neotethys" submitted by Chen and Robertson.

First of all, I want to respond one of the authors' comments in "General Response (7)": "Also, it is normal for this research community to widely cite relevant literature (it is not uncommon in Turkey for aggrieved authors to take cases to university ethics committees if they feel their work has not been adequately cited)."

As a Turkish scientist, this sentence is directly addressing myself and looks trying to provoking and manipulating me to make political review rather than a scientific review. So, I strongly suggest the authors relying on scientific comments. Here all my comments rely on nothing but my scientific vision. The rest is not the authors' business.

I suggest using the subtitle "Results and Discussion" instead "Discussion". The authors present mostly their field and analytical data in sections 5.1 and 5.2.

The Discussion section, especially sections 5.1 Genetic type/source characteristics and 5.2 Tectonic discrimination, provides further interpretation of the field and analytical data. So we think the subtitle "Discussion" here is OK and in line with use in similar papers.

The introduction, Methodology and the presentation of the results of the work seem to be alright. However, the discussion is still debatable.

Discussion

The authors still insisting on linking Kyrenia range to SE Anatolia. It is obvious that these units are continuation of each other in terms of paleogeography, however, their tectonic setting is quite different. The author should explain in detail, possibility of roll-back in a southerly located oceanic crust (Kızıldağ-Troodos oceanic crust) with an Eocene arc.

More details of the paleogeographic continuation (agreed) are now given, to help clarify the setting of the various tectonic units in SE Turkey. In addition, more detailed comparisons (paleogeographically and geochemically) between the possible equivalents of the Kyrenia Range volcanics are included, together with literature interpretations. Several additional relevant recent references are included (section 5.4). However, we have kept the additions to a necessary minimum to contain the length of this section.

If the Bitlis-Pütürge (BP) and Tauride platforms collided during late Cretaceous. The HP metamorphism of the BP and the shoshonitic magmatism during late Cretaceous located in Pertek (Elazığ) are addressing a late Cretaceous collision (see results of Oberhänsli et al, 2010, 2014; Sar et al., 2019). After this there should be no Berit Ocean in SE Anatolia and BP attached to Tauride platform acting the active margin of southerly located subduction, which produces the Kızıldağ and Troodos ophiolites.

The genesis of oceanic crust, including Kızıldağ and Troodos ophiolites, by SSZ spreading is c. 90 Ma (e.g. Mukasa and Ludden, 1987). In contrast, the geochronology indicates a tectono-metamorphic age of Santonian-Campanian for the HP-LT metamorphism of the BP microcontinents (82-80 Ma; Oberhänsli et al., 2010, 2013; Çetinkaplan et al., 2016). Therefore, the younger magmatism is likely to result from an advanced stage of northward subduction of the oceanic plate, although a more southerly origin is possible for the Kyrenia volcanics, see Line 570-577. However, we agree that the Berit ocean probably sutured during the latest Cretaceous, while S Neotethys remained open to the south until the Miocene. We make this clearer.

The Eocene arc magmatism should be a result of the southerly located subduction and the authors proposal of roll-back during Paleogene rejects the arc magmatism during Eocene.

The Eocene arc magmatism could be caused by a southerly located subduction given the fact that the Berit meta-ophiolite is unconformably overlain by the Eocene Maden Complex N of the Malatya Metamorphic (Robertson et al., 2006). If the Meydan ophiolite and cross cutting granite are the same unit (likely), then the Kyrenia Late Cretaceous volcanics should have a different origin,

maybe an arc further south. We indicated this before but now we are more explicit (lines 583-593).

The proposed model does not reject the arc magmatism during Eocene. The proposed model suggests the Eocene volcanics formed in an oblique-convergent setting, in other words, arc magmatism could possibly take place especially in local regions where remnant oceanic crust still exists. In addition, the Eocene Kyrenia rocks are mainly within-plate in affinity, rather than arc-related.

If the authors proposal explains the existing of HP metamorphism and shoshonitic magmatism during late Cretaceous and arc magmatism during Eocene, I will be happy to read.

More information about the HP-LT metamorphism is given in Line 487-497, as requested. The Late Cretaceous shoshonitic magmatism is now mentioned. However, literature review indicates that shoshonitic magmatism does not have a unique tectonic origin but may occur in a range of extensional to post-collisional subduction-related settings (see lines 551-561). Concerning the Eocene arc magmatism, there is geochemical evidence for this in places in SE Turkey but little supporting evidence for a regional-scale arc although as we note this could have subducted (see lines 583-593). However, in agreement with the reviewer, we infer an active margin rather than post-collisional setting for these volcanics in contrast to some other authors.

Figures&tables look adequate and good in drawing.

My suggestion is accepting the MS with considering the comments above. The text should be spell checked.

Regards,
Dr. Fatih KARAOĞLAN

Oberhänsli R, Koralay E, Candan O, Pourteau A, Bousquet R (2014) Late Cretaceous eclogitic high-pressure relics in the Bitlis Massif Geodin Acta 26:175-190 doi:10.1080/09853111.2013.858951

Oberhänsli R, Candan O, Bousquet R, Rimmele G, Okay A, Goff J (2010) Alpine high pressure evolution of the eastern Bitlis complex, SE Turkey. In: Sosson M, Kaymakci N, Stephenson RA, Bergerat F, Starostenko V (eds) Sedimentary Basin Tectonics from the Black Sea and Caucasus to the Arabian Platform, vol 340. vol 1. Geological Society, London, Special Publications, London, pp 461-483. doi:10.1144/sp340.20

Sar A, Ertürk MA, Rizeli ME (2019) Genesis of Late Cretaceous intra-oceanic arc intrusions in the Pertek area of Tunceli Province, eastern Turkey, and implications for the geodynamic evolution of the southern Neo-Tethys: Results of zircon U-Pb geochronology and geochemical and Sr-Nd isotopic analyses Lithos 350-351:105263 doi:10.1016/j.lithos.2019.105263

These references were included.

Abstract

Late Cretaceous felsic and latest Cretaceous-Paleogene basaltic volcanic rocks are exposed throughout the Kyrenia Range, N Cyprus. Field mapping of the key, well-exposed western range indicates that the felsic volcanics are mainly crop out in the southerly, frontal part of the range, separated from the latest Cretaceous-Paleogene basaltic volcanics farther north by a thrust. New U-Pb zircon dating of the felsic volcanics indicates a primary age of c. 74.0 Ma (Late Campanian). The felsic volcanics are characterised by evolved compositions that were probably generated in an extensional subduction-related setting. Their relatively low Nb and Y but high Rb concentrations, together with characteristic Th/Ta ratios (6-20), suggest a mature continental arc setting. The latest Cretaceous-Paleogene basaltic volcanics have mainly within-plate chemical characteristics (in the east), although some have a subordinate subduction influence; e.g. negative Nb (in the west). Sr-Nd-Hf isotopic signatures (i.e. high positive $\epsilon\text{Nd}(t)$ and $\epsilon\text{Hf}(t)$ values) suggest derivation of the basalts from several OIB-like mantle sources, with probable involvement of a crustal (recycled) component (i.e. elevated Nb/Y ratios). Comparisons with SE Turkey, where Late Cretaceous arc-related granitic rocks are widely developed, suggest that the Kyrenia Range igneous rocks may have originated to the south of the Alanya metamorphic massif (N of Cyprus) and correlative continental units in SE Turkey. The latest Cretaceous-Paleogene basaltic volcanics in N Cyprus and SE Turkey (Maden Complex) are interpreted to represent incipient marginal basin formation, possibly in an oblique-convergent setting, prior to Miocene suturing with Arabia. In the light of alternatives, we infer genesis of the N Cyprus Late Cretaceous and Paleogene volcanic rocks related to stages in the development of the northerly active continental margin of the Southern Neotethys.

Highlights

- Felsic volcanics erupted at c. 74 Ma in an active continental margin arc setting
- Some within-plate basalts (c. dated at 70-45 Ma) were affected by subduction
- The basalts are interpreted to represent incipient marginal basin formation
- The volcanism provides new evidence of subduction in the E Mediterranean region
- Northward subduction is supported by coeval arc magmatism in SE Turkey

**Evidence from Late Cretaceous-Paleogene volcanic rocks of the Kyrenia Range,
northern Cyprus for the northern, active continental margin of the Southern
Neotethys**

Guohui Chen^{1*}, Alastair H. F. Robertson¹

¹School of GeoSciences, University of Edinburgh, West Mains Road, Edinburgh EH9
3JW, UK

*Current Address: State Key Laboratory of Lithospheric Evolution, Institute of Geology
and Geophysics, Chinese Academy of Sciences, P.O. Box 9825, Beijing 100029,
China

Corresponding author: Guohui Chen (Guohui.Chen@live.cn)

Abstract

Late Cretaceous felsic and latest Cretaceous-Paleogene basaltic volcanic rocks are exposed throughout the Kyrenia Range, N Cyprus. Field mapping of the key, well-exposed western range indicates that the felsic volcanics are mainly crop out in the southerly, frontal part of the range, separated from the latest Cretaceous-Paleogene basaltic volcanics farther north by a thrust. New U-Pb zircon dating of the felsic volcanics indicates a primary age of c. 74.0 Ma (Late Campanian). The felsic volcanics are characterised by evolved compositions that were probably generated in an extensional subduction-related setting. Their relatively low Nb and Y but high Rb concentrations, together with characteristic Th/Ta ratios (6-20), suggest a mature continental arc setting. The latest Cretaceous-Paleogene basaltic volcanics have mainly within-plate chemical characteristics (in the east), although some have a subordinate subduction influence; e.g. negative Nb (in the west). Sr-Nd-Hf isotopic

signatures (i.e. high positive $\epsilon\text{Nd}(t)$ and $\epsilon\text{Hf}(t)$ values) suggest derivation of the basalts from several OIB-like mantle sources, with probable involvement of a crustal (recycled) component (i.e. elevated Nb/Y ratios). Comparisons with SE Turkey, where Late Cretaceous arc-related granitic rocks are widely developed, suggest that the Kyrenia Range igneous rocks may have originated to the south of the Alanya metamorphic massif (N of Cyprus) and correlative continental units in SE Turkey. The latest Cretaceous-Paleogene basaltic volcanics in N Cyprus and SE Turkey (Maden Complex) are interpreted to represent incipient marginal basin formation, possibly in an oblique-convergent setting, prior to Miocene suturing with Arabia. In the light of alternatives, we infer genesis of the N Cyprus Late Cretaceous and Paleogene volcanic rocks related to stages in the development of the northerly active continental margin of the Southern Neotethys.

Keywords: Late Cretaceous-Paleogene; Volcanic rocks; Kyrenia Range; Arc magmatism; Marginal basin formation; Southern Neotethys

1. Introduction

Continental margin magmatism is a key feature of subduction at convergent plate margins (Dewey, 1969; Şengör and Natal'in, 1996). Information concerning the field relations, age, petrology and geochemistry are essential to understand the growth and demise of arcs, and are critical to reconstruct paleogeography and to test alternative tectonic hypotheses (e.g. Trehu et al., 1994; Tatsumi and Kogiso, 2003; Bowman et al., 2019).

The Eastern Mediterranean region, situated between the North African-Arabian and Anatolian (Eurasian) plates, includes tectonically emplaced remnants of Neotethys that developed from the preceding Paleotethys farther north (e.g. Le Pichon, 1982; Şengör et al., 1984; Robertson and Dixon, 1984; Stampfli and Borel, 2002). Much of the arc magmatism in the north, in the Pontides, of Jurassic-Eocene age (Fig. 1), relates to long-lived northward subduction of the Northern Neotethys (İzmir-Ankara-Erzincan ocean) (Adamia et al., 1977; Dercourt et al., 1993; Ustaömer and Robertson, 1997; Rice et al., 2006). Subduction of the Southern Neotethys resulted in both continental margin and oceanic magmatism that ranges, overall, from Late Cretaceous to Paleogene (Fig. 1). Here, we focus on evidence of Late Cretaceous and Paleogene volcanism in the Kyrenia Range of northern Cyprus, representing the most westerly known occurrence of arc-type magmatism that can be related to subduction of the Southern Neotethys (Fig. 1) (Moore, 1960; Ducloz, 1972; Pearce, 1975; Baroz, 1979, 1980; Robertson and Woodcock, 1986; Huang et al., 2007; Robertson et al., 2012b).

Our specific objectives here are: (1) to understand the tectono-stratigraphy of two, contrasting, felsic and basaltic volcanic units in relation to associated sedimentary units; (2) to determine the eruptive ages of both of the volcanic units directly, using geochronology. Previously, the volcanics were dated indirectly using microfossils within interbedded pelagic carbonates (Baroz, 1979; Robertson et al., 2012b); (3) to determine whether there is any petrological or geochemical variation (including

isotopic variation) along the Kyrenia Range (c. 100 km) and, if so, the implications; (4) to infer the magmatic-tectonic setting of eruption, compared with the evidence of arc volcanism, especially in southeast Turkey; (5) to test alternative tectono-magmatic hypotheses bearing in mind that arc magmatism plays a key role in plate tectonic interpretations.

2. Volcanism in the Kyrenia Range

Rifting of the Southern Neotethys during Late Permian and Early-Middle Triassic was followed by continental break-up during the Late Triassic-Early Jurassic (Robertson and Woodcock, 1979; Şengör and Yılmaz, 1981; Robertson and Dixon, 1984; Garfunkel, 1998; Robertson et al., 2020). The Kyrenia Range then formed part of a carbonate platform that slowly subsided during Jurassic-Early Cretaceous in a passive margin setting (Robertson and Woodcock, 1986). The carbonate platform was deformed and metamorphosed under greenschist facies during the Late Cretaceous, extensionally exhumed, and then unconformably overlain by basic extrusive igneous rocks of latest Cretaceous-Paleogene age (Figs. 2-3) (Ducloz, 1972; Baroz, 1979; Robertson and Woodcock, 1986). In addition, felsic volcanics and tuffs are exposed as thrust slices at a low structural level in the south of the range (Figs. 2b, 3) (Robertson et al., 2012b). These volcanic rocks have no exposed base and are directly overlain by a much larger thrust sheet that includes the mainly basic extrusive igneous rocks.

Baroz (1979, 1980) mapped the volcanic rocks and carried out petrographic and whole-rock chemical analysis of major elements. He reported the presence of a lower stratigraphic sequence of basalt, dolerite, trachybasalt, trachyandesite, dacite and rhyolitic tuff, and interpreted this as a bimodal basic-acidic calc-alkaline suite, related to a Late Maastrichtian volcanic arc. In contrast, the Paleogene volcanic assemblage

was suggested to have erupted stratigraphically above in a post-collisional (intra-continental) strike-slip setting (Baroz, 1980). However, a thrust was later mapped between the mainly felsic and mainly basic suites (Robertson et al., 2012b), complicating this simple stratigraphy.

Chemical analysis of the Paleogene basaltic lavas, including immobile trace elements, initially suggested an alkaline, within-plate eruptive setting (Pearce, 1975). Subsequent chemical analysis, including some immobile elements, confirmed this mainly alkaline within-plate setting but also revealed some evidence of a subduction influence (e.g. negative Nb anomaly) (Robertson and Woodcock, 1986). The Paleogene lavas were inferred by these authors to have erupted in a transtensional setting along the northern, active margin of the 'Troodos ocean'. Huang et al. (2007) carried out additional chemical analysis of the Paleogene basaltic lavas, mainly from the eastern and western Kyrenia Range, with emphasis on immobile elements, and proposed a Late Cretaceous-Paleogene back-arc setting related to northward subduction.

The felsic rocks of the lowermost thrust sheet, known as the Fourkovouno (Selvilitepe)¹ Formation, up to 400 m thick, begin with marine water-lain felsic tuffs and subaqueous felsic debris-flow deposits and culminate in laterally discontinuous, thick-bedded to massive rhyolitic lava flows (Fig. 3) (Moore, 1960; Baroz, 1979; Huang et al., 2007; Robertson et al., 2012b). The felsic rocks are locally cut by small (meter-sized) basaltic/doleritic intrusions (not studied). A Late Cretaceous age (Late Campanian) was suggested for the felsic lavas, based on sparse planktic foraminifera (*Globotruncana* sp.) that were recognised in a chalky interbed near the top of the

¹ For simplicity, we use the traditional stratigraphy for the formation names (more recent Turkish equivalents are mentioned initially). However, we use current the Turkish names (e.g. for settlements); pre-existing names are mentioned initially.

117 succession (Baroz, 1979).

118 The Late Cretaceous and Paleogene basaltic igneous rocks have been divided
119 into two formations, namely the Maastrichtian Melounda (Mallıdağ) Formation and the
120 Paleocene-Middle Eocene Ayios Nikolaos (Yamaçköy) Formation (Fig. 2a).

121 The Melounda Formation, c. 300 m thick, unconformably overlies the exhumed
122 Mesozoic platform carbonates of the Trypa (Tripa) Group or, in places,
123 unmetamorphosed terrigenous turbidites of the Kiparisso Vouno (Alevkaya Tepe)
124 Member of the Melounda Formation (Baroz, 1979; Robertson and Woodcock, 1986;
125 Robertson et al., 2012b). The basaltic lavas of the Melounda Formation are commonly
126 intercalated with pinkish pelagic limestones (Fig. 2a). The succession locally includes
127 lenticular, lithoclastic breccias and finer-grained clastic facies that were reworked from
128 the underlying meta-platform carbonates (Fig. 2a) (Robertson and Woodcock, 1986;
129 Robertson et al., 2012b). Planktic foraminifera from pelagic carbonates between the
130 lavas indicate a Late Maastrichtian age (Baroz, 1979; Robertson et al., 2012b).

131 The overlying Ayios Nikolaos Formation, 300-400 m thick, comprises basaltic
132 volcanics that are variably interbedded with pelagic carbonates. Calciturbidites,
133 calcareous debris-flow deposits and carbonate-rock breccias appear above the lavas,
134 towards the top of the overall succession (Baroz, 1979; Robertson and Woodcock,
135 1986; Robertson et al., 2012b, 2014) (Fig. 2a). Planktic foraminifera from pelagic
136 carbonates that are interbedded with the lavas indicate a Late Paleocene to Mid-
137 Eocene age range (Baroz, 1979; Robertson et al., 2012b).

138 139 3. Methods

140 The Kyrenia Range is a complex thrust belt and therefore it was essential to remap
141 the western Kyrenia Range, where felsic and basaltic volcanic rocks are exposed
142 together (Moore, 1960; Baroz, 1979; Robertson et al., 2012b). In addition, sedimentary

logs were measured and correlated to produce a composite stratigraphy of the volcanogenic successions that are exposed in two superimposed thrust sheets.

Optical microscopy was carried out on the samples collected. For the felsic lithologies, 13 samples were studied from the western range. For the basaltic lithologies, we used a combination of our samples (n=29) and also some that were previously collected for paleomagnetic study (n=26) throughout the Kyrenia Range (Morris et al., 2015).

Whole-rock major and trace element concentrations of the volcanic rocks were measured by X-ray fluorescence (XRF) on fused glass beads and pressed powder pellets at the School of GeoSciences, University of Edinburgh, using the well-known methods of Fitton et al. (1998) and Fitton and Godard (2004). Accuracy and precision are typically c. 5%. For representative samples, additional trace and rare earth element (REE) analysis was carried out by inductively coupled plasma-mass spectrometry (ICP-MS) at the ACME Analytical Laboratories, Vancouver. Major element contents were determined from a LiBO₂ fusion by ICP-ES, using 5 g of sample pulp. Trace element contents were determined from a LiBO₂ fusion by ICP-MS, again using 5 g of sample pulp. Detection limits range between 0.01 and 0.04 wt% for major oxides, 0.01 and 0.1 ppm for trace and rare earth elements. The relative standard deviation for the REE is ~5% and for all other trace elements is up to 10%, with quality control using international geostandards (see <http://acmelab.com>).

In addition, Sr-Nd-Hf isotopic analysis was performed on a Neptune Plus multi-collector (MC)-ICP-MS at Wuhan SampleSolution Analytical Technology Co., Ltd., China, as reported by Li et al. (2012). Whole procedural blanks were <100 pg for Sr, <50 pg for Nd and <50 pg for Hf. ⁸⁷Sr/⁸⁶Sr, ¹⁴³Nd/¹⁴⁴Nd and ¹⁷⁶Hf/¹⁷⁷Hf ratios were normalised to ⁸⁶Sr/⁸⁸Sr = 0.1194, ¹⁴⁶Nd/¹⁴⁴Nd = 0.7219 and ¹⁷⁹Hf/¹⁷⁷Hf = 0.7325, using the exponential law. Standard analysis yielded ⁸⁷Sr/⁸⁶Sr = 0.710240 ± 11 (2SD, n = 4) for NBS987, ¹⁴³Nd/¹⁴⁴Nd = 0.512440 ± 8 (2SD, n = 8) for GSB 04-3258-2015, and

$^{176}\text{Hf}/^{177}\text{Hf} = 0.282224 \pm 6$ (2SD, $n = 6$) for Alfa Hf. In addition, USGS reference materials BCR-2 and RGM-2 were also analysed for Sr–Nd–Hf isotopes, and gave ratios of 0.705039 ± 8 and 0.704169 ± 11 for $^{87}\text{Sr}/^{86}\text{Sr}$, 0.512644 ± 6 and 0.512808 ± 9 for $^{143}\text{Nd}/^{144}\text{Nd}$ and 0.282864 ± 6 and 0.283016 ± 7 for $^{176}\text{Hf}/^{177}\text{Hf}$, which is within error of recommended values (Thirlwall, 1991; Weis et al., 2006, 2007). The analytical data for the major, trace and rare earth elements/isotope ratio of the felsic and basaltic volcanics are listed in the Supplementary Table S1, S2, respectively.

In addition, two polished thin sections of basalt (no. 02 and 21) that contain relatively fresh feldspar and pyroxene crystals were selected for analysis of major elements using a Cameca SX100 electron microprobe at the School of GeoSciences, University of Edinburgh. Details of the analytical conditions and methods are given by Hayward (2011) and by Hartley and Thordarson (2013). The accuracy of major element determinations is $<\pm 1\%$ of total value. Analytical data for feldspar and pyroxene are given in Supplementary Table S3, S4, respectively.

For dating, zircon crystals within 30–160 μm fraction were separated from crushed felsic tuffaceous rock and rhyolitic lava (no. 14–18, 14–19 and 14–20; c. 5 kg each). A Wilfley table, Frantz Isodynamic magnetic separator and a high-density solution (lithium polytungstate; 2.85 g/ml) were used to aid separation. The zircon grains were randomly handpicked under a binocular microscope, mounted in epoxy resin and polished sufficiently to expose the centre of the grains. Internal structures were studied with a scanning electron microscope using cathodoluminescence (CL) at the School of GeoSciences, University of Edinburgh. U–Pb analysis was then performed using a Cameca IMS-1270 secondary ion mass spectrometer (SIMS) at the School of GeoSciences, University of Edinburgh, using the methods reported by Kelly et al. (2008) and Ustaömer et al. (2012). Errors of the reported ages are $\pm 1\sigma$. Related geochronological plots were produced using ISOPLOT (Ludwig, 2012). Analytical results are listed in the Supplementary Table S5.

197

198 4. Results

199 4.1. Volcanic successions

200 The felsic volcanics, mainly crop out in the western part of the Kyrenia Range in
201 two thrust sheets (Figs. 3-4). The lower of these, which is relatively intact, extends from
202 Geçitköy (Panagra) eastwards to Selvilitepe (Fourkovouno) (Fig. 5a-e). The second,
203 structurally higher unit, comprises discontinuous exposures extending from north of
204 Geçitköy, to southwest of Karsiyaka (Vasileia) and west of Alevkaya Tepe (Kiparisso
205 Vouno) (Fig. 5f-j). Farther west, in the Kayalar (Orga) area (Fig. 3a), a small thrust slice
206 exposes coarse-grained felsic tuff and a felsic debris-flow deposit. Small (tens of m-
207 sized) outcrops of felsic volcanogenic rocks also occur farther east; e.g. northeast of
208 Aşağı Dikmen (Kato Dikomon), between thrust sheets of Mesozoic meta-carbonate
209 rocks (Baroz, 1979). Small exposures are also intersliced with Late Cretaceous pelagic
210 carbonates of the Melounda Formation to the north of Ergenekon (Agios Khariton) (Fig.
211 3a) (Robertson et al., 2012b).

212 The local successions are restored as a lower thrust sheet and an upper thrust
213 sheet. The succession in the lower thrust sheet, c. 90 m thick, begins with weakly-
214 bedded, white felsic tuff (beds <0.5 m thick) (Fig. 5a, c-e), followed by thick layers of
215 matrix-supported felsic breccia-conglomerate (c. 10 m thick) that we interpret as
216 debris-flow deposits (Fig. 5a, c, e). The succession passes upwards into massive,
217 vitreous, rhyolitic lava flows, individually up to 10 m thick (Figs. 5a, Supplementary
218 Figure S1). Localised normal grading; e.g. northeast of Geçitköy, indicates that the
219 succession is the right way up, stratigraphically (Fig. 5a-b). In places, steeply dipping,
220 irregularly shaped basaltic or doleritic intrusions, up to 2 m across, cut the felsic
221 layering (Figs. 5a, Supplementary Figure S1). Adjacent to these intrusions, poorly
222 consolidated tuffaceous sediments have undergone contact metamorphism to form

hard, dark grey, flint-like, recrystallised felsic rock (Robertson et al., 2012b). There is no evidence of similar intrusions in the structurally overlying basalts, which is in keeping with the mapped tectonic contact between the two units. The succession in the upper thrust sheet begins with greyish to greenish, tuffaceous matrix-supported conglomerates, including clasts of silicified rhyolite, chalk and basalt (Supplementary Figure S1). The estimated thickness ranges from c. 40 m north of Geçitköy (Fig. 5f), to c. 80 m southwest of Karsiyaka (Fig. 5g-h), to 30 m west of the Alevkaya Tepe (Fig. 5i-j).

Contrasting field relations facilitate mapping of the Late Cretaceous versus the Paleogene formations (Baroz, 1979; Robertson and Woodcock, 1986; Hakyemez et al., 2000; Robertson et al., 2012b). In the far east of the Kyrenia Range, the Karpas Peninsula (Fig. 3a), both pillowed and massive basalt are widely exposed near Balalan (Platanissos). Interpillow carbonate and lenticular interbeds of pelagic carbonate (< 1 m thick) (Figs. 3a, Supplementary Figure S2) contain Late Maastrichtian microfossils (Robertson et al., 2012b). In the eastern range, basaltic rocks of the Melounda Formation contain abundant pelagic carbonate and are overlain by massive basalts of the Ayios Nikolaos Formation; i.e. from Ağıllar (Mandres) to Mallıdağ (Melounda) (Fig. 3a) (Baroz, 1979; Robertson and Woodcock, 1986; Hakyemez et al., 2000; Robertson et al., 2012b). In the central range, pillow lava with interbedded pelagic carbonate of the Melounda Formation occurs in two settings. The first is directly above the Upper Cretaceous basal unconformity, with the meta-carbonate platform rock beneath; e.g. near Tirmen (Trypimeni). The second setting is as one (or several), small thrust sheets along the southern flank of the range; e.g. Değirmenlik (Kythrea), Boğaz (Bogaz) and southwest of Alevkaya Tepe (Figs. 3a, Supplementary Figure S2). Basalt with some interbedded pelagic carbonate of the Ayios Nikolaos Formation is exposed higher in the succession (within a shear zone), near Tirmen, Arapköy (Klepini) and İncesu (Motides) (Baroz, 1979). In the western range, basaltic lavas interbedded with pelagic

carbonates of the Melounda Formation are exposed in a thrust sheet along the southern margin of the range; e.g. Geçitköy (Figs. 3, Supplementary Figure S2). Similar basaltic lavas with pelagic carbonates and local carbonate breccias are exposed along the northern margin of the range; e.g. near Karşıyaka and farther west (Baroz, 1979; Hakyemez et al., 2000; Robertson et al., 2012b). In addition, basaltic lava interbedded with pelagic chalk of the Ayios Nikolaos Formation occurs higher in the succession (southward-younging) along the eastern side of Geçitköy road (Huang et al., 2007; Robertson et al., 2012b).

In summary, the Late Maastrichtian basalts (Melounda Formation) are typically pillowed, forming repeated flows (individually up to c. 10 m thick), reaching a maximum of > 90 m in the east (near Balalan). The lavas include pink pelagic carbonate as interstitial sediment and discontinuous layers. In contrast, the Paleogene basalts (Ayios Nikolaos Formation) are commonly massive, especially in the eastern range, where individual lava flows reach c. 45 m thick, with relatively few pelagic intercalations (Supplementary Figure S2). In the central and western ranges, thinner lava flows (<10m) are intercalated with pelagic carbonates and reddish chert (formed by diagenetic replacement). During this work, we identified geochemical differences between the Late Cretaceous-Paleogene basaltic lavas of similar age along the far-eastern, eastern, central and western Kyrenia Range (see below).

4.2. Petrography

The felsic rocks are porphyritic, composed of quartz, plagioclase and sanidine, together with subordinate biotite and rare hornblende. Large sanidine crystals (0.3-1 mm) show simple twinning, whereas plagioclase commonly has lamellar twinning (Supplementary Figure S3). Biotite laths (0.3-0.8 mm in length) are preferentially orientated parallel to flow layering. Rare quartz phenocrysts (<1%, 0.2-0.5 mm) are

commonly fragmented, probably related to quenching during eruption; i.e. explosive fragmentation (Supplementary Figure S3). The groundmass comprises microcrystalline quartz and feldspar, together with minor muscovite, biotite and rare hornblende (c. 60 μm). In some samples (e.g. nos. 14-19 and 14-20), the groundmass is cryptocrystalline, almost glassy (Supplementary Figure S3).

The basaltic rocks of both Late Cretaceous and Paleogene age generally fall into two groups in terms of mineral composition and texture. The first group (most common) is porphyritic with subhedral to euhedral clinopyroxene phenocrysts (30-40%), up to 0.6 mm long, that occur interstitially or are intergrown with subhedral feldspar (Supplementary Figure S4, Table S3). Feldspar (45-60%) forms elongate, acicular laths, up to 1 mm long. Olivine is relatively rare (<5%). The second group is ophitic, as locally observed in the western Kyrenia Range with abundant anhedral to subhedral augite phenocrysts (up to 0.6 mm in size). Most plagioclase is acicular, enclosed or surrounded by augite (Supplementary Figure S4).

Most samples are slightly to moderately altered. Amygdales and veins are infilled with secondary minerals such as calcite and zeolite. Alteration is variable, for example, pyroxene and plagioclase are heavily altered to chlorite and clay minerals (e.g. smectite/sericite). The relatively high loss on ignition values result from secondary processes, which need to be taken into account prior to rock identification and tectonic discrimination.

4.3. Whole-rock chemistry

Major-element oxides (K_2O , Na_2O and CaO) and trace elements (e.g. Rb, Sr and Ba) exhibit a relatively wide compositional range, consistent with the effects of alteration (see Supplementary Table S1-S2). Alteration can be inferred from variable LOI values that range between 0.3-5.4 wt% for the felsic volcanics and 4.09-14.57 wt% for the basaltic rocks. This is consistent with the presence of considerable amounts of

water- and/or carbonate dioxide-bearing minerals, for example chlorite and calcite, as observed petrographically (see Supplementary Figure S3, S4). Trace elements such as Ti, Zr, Y, Nb, Ta, V, Co, Th and REEs tend to be immobile during weathering and/or metamorphism below amphibole facies are, therefore, preferred for rock-type identification and tectonic discrimination (e.g. Pearce and Cann, 1973; Rollinson, 1993). For basaltic rocks, samples with high crystal contents (cumulate composition or highly porphyritic) should be discounted from geochemical discrimination (Pearce, 1996). In general, samples with immobile element $\text{Al}_2\text{O}_3 > 20 \text{ wt\%}$ (concentrated in feldspar), $\text{Sc} > 50 \text{ ppm}$ (concentrated in clinopyroxene), or $\text{Ni} > 200 \text{ ppm}$ (concentrated in olivine) are predicted to contain high amounts of cumulated minerals (Pearce, 1996). On this basis, basaltic rocks nos. 14-66, 14-50, 14-51, 14-58, 14-68, 08, 16, 25 and 28 are discounted.

The more felsic assemblage is mainly rhyolitic in the classification of Winchester and Floyd (1977) but shows a trachytic affinity in the revised Zr/Ti vs. Nb/Y (Fig. 6a) diagram of Pearce (1996). Relatively high abundances of Th ($>14 \text{ ppm}$) but low Co concentrations ($<12 \text{ ppm}$) are consistent with high-K and shoshonitic affinities (Fig. 6b). Primitive mantle-normalised multi-element plots (Fig. 6c) show variable enrichments and depletions in large-ion lithophile elements (LILEs; e.g. Cs, Rb, Ba, K and Sr) that are influenced by alteration. The samples show negative anomalies of Nb, Zr and Ti, coupled with marked positive anomalies of Th/U, Pb, Nd and probably Sm (Fig. 6c). Total REE contents of the felsic assemblage (Fourkovouno Formation) vary from 115 to 173 ppm. La ranges from 29-44 ppm, $(\text{La/Yb})_{\text{N}}=12.63\text{-}18.98$, suggestive of an alkalic affinity. The ratios $(\text{La/Sm})_{\text{N}}=6.13\text{-}7.22$ and $(\text{Gd/Yb})_{\text{N}}=1.26\text{-}1.68$ suggest a large LREE fractionation but low to moderate HREE fractionation (Fig. 6d). Pronounced negative Eu anomalies ($\text{Eu/Eu}^*=0.32\text{-}0.48$) are consistent with plagioclase crystallization, as seen petrographically (see Supplementary Figure S3). In general, the samples are comparable to upper continental crust (UCC). The basaltic rocks plot in the fields of

basalt and alkali basalt, with intermediate degrees of fractionation ($Zr/Ti=0.01-0.02$) and moderate alkalinity ($Nb/Y=0.1-1.8$) on the rock-type discrimination plot (Fig. 7a). The basalts have a calc-alkaline affinity, except for samples from Mallıdağ (no. 01), Tirmen (no. 19-51), Değirmenlik (no. 14-64) and İncesu (no. 19-67) that exhibit high-K and shoshonitic affinities (Fig. 7b). The trace element compositions of the basaltic rocks exhibit wide ranges of Sr, Ba, Rb, Th, Ta, Nb and Ce. However, Zr, Hf, Sm, Ti and Y generally comparable with enriched mid-ocean ridge basalt (E-MORB) and oceanic island basalt (OIB) (Fig. 8a-c).

The basaltic rocks are characterised by variable REE concentrations (67.2-239.4 ppm) and light REE enrichments ($(La/Yb)_N=3.18-19.35$) (Fig. 8d). Samples from the central range, at Değirmenlik and Tirmen, show much higher total REE contents and steeper REE patterns. Westwards in the Kyrenia Range, the patterns become smoother, with less pronounced enrichment in LREEs. In general, no marked Eu anomaly (~ 1) is observed.

4.4. Whole-rock Sr-Nd-Hf isotopes

The selected basaltic samples have relatively uniform whole-rock Sr-Nd-Hf isotopic characteristics. Adopting a Late Cretaceous eruption age based on the paleontological dating (Robertson et al., 2012b), the calculated initial $^{87}Sr/^{86}Sr$ ratios range from 0.705054 to 0.705701, with $\epsilon Nd(t)$ values of 2.27 to 5.76. Initial $^{176}Hf/^{177}Hf$ and $\epsilon Hf(t)$ values are 0.282934-0.283017, 7.36-10.14, respectively (see Supplementary Table S2).

4.5. Zircon U-Pb geochronology

Zircon crystals separated from the felsic rocks are typically angular to sub-angular, 30-160 μm long. The grains have variable internal textures, including banded zoning (e.g. zircon 1), concentric zoning (nos. 2, 4, 5 and 12), and minor sector zoning (nos.

10, 11; Fig. 9a). Fractures (or depressions) and some inherited cores are affected by fluid-related chemical alteration (nos. 6 and 8) (Fig. 9a). The zircon grains have high Th/U ratios (0.19-1.24), suggestive of a magmatic origin (Rubatto, 2002). Twelve zircon grains were analysed with concordance levels ranging from 99-105%. The calculated $^{206}\text{Pb}/^{238}\text{U}$ age with the generally lowest possible error was used for age determination. The much older zircon grain (no. 4) is subhedral with a patchy xenocrystic texture, suggestive of a recycled origin (Fig. 9a). The concordant or nearly concordant data (100-102%) with a young age distribution yielded weighted mean $^{206}\text{Pb}/^{238}\text{U}$ gave ages of 74.0 ± 0.6 Ma, 74.0 ± 0.4 Ma and 71.7 ± 0.7 Ma (Fig. 9b-d).

5. Discussion

5.1. Genetic type/source characteristics

For the felsic rocks, the high LREE/HREE (e.g. $(\text{La}/\text{Yb})_{\text{N}}=12.63-18.98$) and negative Eu anomalies, resemble model UCC but with depletion in Eu, Sr and P (Fig. 6c-d). The A/CNK (molar ratio of $\text{Al}_2\text{O}_3/(\text{CaO}+\text{Na}_2\text{O}+\text{K}_2\text{O})$) ratio is useful to identify the source of felsic rocks (Chappell and White, 1992) but is questionable for altered rocks like those of the Kyrenia Range. SiO_2 , P_2O_5 and Th can also be used as indices of the fractional crystallization of a felsic melt (Chappell and White, 1992), but again may be affected by alteration (specially SiO_2). In our samples, P_2O_5 decreases with increasing SiO_2 and Th (Fig. 10a-b), trends that are indicative of I (igneous)- or A (anorogenic)-type granites. The relatively low contents of Zr, Nb, Y, La, Ce, Zn and Ga further characterise these rocks as I-type (Fig. 10c-d). The abundances of Th and Co, in particular hint at a shoshonitic composition (Hastie et al., 2007).

The trace element patterns of the basaltic rocks show a wider range of variation (Fig. 8). E-MORB and OIB-type basalts were derived from relatively enriched sources,

specifically for the basalts of the Karpas Peninsula, eastern range and most of the central range. The lesser enrichment of Nb within the basalts of the central range at Ergenekon (no. 31), Arapköy (no. 14-62) and Alevkaya Tepe (no. 14-49), and the western range at Karsiyaka (no. 14) and Geçitköy (no. 10), can be explained by a relatively depleted mantle source, together with subduction modification. There is little evidence of the addition of radiogenic Sr from seawater, either to the source melt or during post-magmatic alteration. The Sr vs. Nd isotope correlation diagram (Fig. 11a) does not show a significant shift towards higher initial $^{87}\text{Sr}/^{86}\text{Sr}$ ratios as would be expected in both of these cases.

The basalts all have positive $\epsilon\text{Nd}(t)$ values (+2.27 - +5.76), consistent with derivation from an OIB-like source mantle (Fig. 11a). The samples plot directly on the terrestrial array, overlapping with the Nd isotope compositions of Etna and the Aeolian Arc, but intermediate between these settings in Hf isotopes (Fig. 11b). The samples that show trace element evidence of a subduction-modified source, based on relatively flat MORB-normalised multi-element patterns and negative Nb anomalies (no. 10, Geçitköy), plot in the Aeolian Arc-type mantle field. Conversely, samples from the Karpas Peninsular, at Balalan, plot close to the field of Etna and exhibit less subduction modification (Fig. 11b). The trace element and isotopic data, therefore, yield consistent results.

The involvement of a crustal component is also suggested by incompatible element ratio plots, on which the basalts exhibit two distinct trends. The samples with relatively strong subduction signatures plot away from the MORB array (Fig. 12a), fingerprinting a subduction-related lithospheric mantle end member. A probable lower crust input is suggested by relatively higher La/Nb and Ba/Nb ratios (e.g. Wang et al., 2004). In contrast, an OIB-like mantle component dominated most of the other basalt samples, as indicated by relatively low La/Nb, Zr/Nb but high Nb/Y ratios (Fig. 12b). The ratios of highly incompatible elements with small distribution coefficients (e.g.

Zr/Nb, Nb/Th, Nb/La) are sensitive indicators of mantle source heterogeneity (Saunders et al., 1988; Condie, 2003; Pearce and Stern, 2006). The samples from the western locations fall in the fields of an enriched component and upper continental crust (Fig. 12c), which could represent mantle mixing with continental sediments (subducted) or crustal contamination. In contrast, the samples from the easterly locations resemble those of the OIB/recycled component, and cluster around a different batch-melting trajectory (10-20%) (Fig. 12c) that is representative of recycled oceanic lithosphere (Saunders et al., 1988). We, therefore, infer that the basalts are likely to have been derived from multiple sources (i.e. two or more source components were involved).

5.2. Tectonic discrimination

On the Nb vs. Y diagram (Fig. 13a), the felsic volcanics show similarities with volcanic arc granites and syn-collisional granites. One sample from Geçitköy (no. 16-08), with relatively high Nb and Y concentrations, plots near the field of within-plate granite. On the Th/Yb vs. Ta/Yb diagram (Fig. 13b), the volcanics mainly erupted on active continental margins. An active continental margin setting is therefore inferred for the felsic volcanics.

Basalts of the Kyrenia Range (e.g. easterly locations) mostly exhibit a within-plate affinity (Fig. 8). However, some samples, especially in the west, have a subduction-related signature (e.g. negative Nb anomalies) (Fig. 8b-c) akin to volcanic arc basalts. Most samples have a within-plate affinity on the Zr/Y vs. Zr (Fig. 13c) diagram, except for two samples, from Ergenekon and Arapköy (central range), that straddle the island arc and MORB fields. The basalts overlap the MORB and back-arc basin basalt fields on the Ti vs. V plot (Fig. 13d). Clinopyroxene compositions, as shown in the ternary $\text{TiO}_2\text{-MnO-Na}_2\text{O}$ plot (see Supplementary Figure S5), indicate a dominantly within-

plate affinity. The relatively high TiO_2 contents of the clinopyroxenes point to an enriched MORB-related source (see Supplementary Figure S5). Overall, the inferred subduction contribution decreases from west to east along the range.

5.3. Global comparison of the Kyrenia Range volcanic rocks

Global comparisons (see Supplementary Figure S6) lend support to our interpretation of the Kyrenia Range volcanic rocks as being subduction-related. In particular, within the Late Cretaceous felsic rocks, the enrichment of Rb, Th and Sm versus the depletion of Ti, Ba, Nb, P compared to primitive mantle, are characteristic of volcanic arc granites (Fig. 6c) (Pearce et al., 1984).

The Kyrenia Range Maastrichtian and Paleogene basalts of within-plate affinities (e.g. Balalan, eastern range) mainly have MORB-normalised spider patterns that are similar to the Pliocene-Quaternary OIB-type basalts of the Tyrrhenian Sea, W Mediterranean (Peccerillo, 2017). The MORB-normalised patterns of these basalts suggest a within-plate eruptive setting, with a small sub-crustal lithosphere metasomatic influence (Kastens et al. 1988; Peccerillo, 2017).

Back-arc basalts of the Early Cretaceous Sarmiento Complex, southern Chile (Saunders et al., 1979) and the Miocene Japan Sea (Tamaki et al., 1990) resemble a minority of the Kyrenia Range basalts (c. 10%; e.g. Geçitköy, western range), although with lower abundances of K, Rb, Ba, Th, Ta and Nb. The relative depletion of these elements is characteristic of eruption in a relatively wide, evolved back-arc basin (e.g. Bransfield Strait; Weaver et al., 1979). However, another possible setting is eruption in a back-arc basin that was influenced by oblique extension (e.g. Guaymas pull-apart basin; Brune et al., 2012). The relatively flat MORB-normalised patterns, akin to E-MORB, of both the Bransfield Strait and Guaymas back-arc basin basalts (e.g. Saunders et al., 1982; Keller et al., 2002) point to substantial involvement of depleted

458 MORB sources.

459

460 5.4. Comparison with the adjacent Tethyan region

461 Late Cretaceous and Paleogene arc-related rocks; i.e. similar in age of those in
462 the Kyrenia Range are critical to any regional magmatic-tectonic interpretation.

463 In western Cyprus, a Late Cretaceous arc-derived volcanogenic succession
464 (Kannaviou Formation) overlies the Troodos ophiolite. The volcanoclastic sediments
465 were derived from intermediate-felsic eruptions, c. 6 Ma before the felsic eruptions in
466 the Kyrenia Range (c. 80 vs. 74 Ma). Grain size and textural evidence suggest that,
467 depending on prevailing wind direction and strength, the eruptive centers are likely to
468 have been <100 km away (Chen and Robertson, 2019). The centers for the Kannaviou
469 Formation were located to the south of the Kyrenia Range or its lateral extension, areas
470 that are not now exposed. The Kyrenia Range was affected by south-directed thrusting
471 during Mid-Eocene and Late Miocene-earliest Pliocene (Baroz, 1979; Robertson et al.,
472 2012b, 2014; McCay and Robertson, 2013; Robertson and Kinnaird, 2016), such that
473 the distal (southerly) edge of the inferred over-riding active margin (see below) could
474 have been subducted, structurally over-ridden or eroded. In other words, additional arc
475 units could have existed to the south of their present location in the Kyrenia Range.

476 To the northeast (c. 200 km), the Misis-Andırın Range in southern Turkey is an
477 extension of the Kyrenia lineament (Robertson et al., 2004). The outcrop, known as
478 the Misis-Andırın Complex, includes massive basalts, pillow lavas, lava breccias and
479 volcanoclastic debris-flows. These lithologies were previously inferred to be Miocene
480 (Floyd et al., 1991, 1992). However, inter-lava sediments there have been dated as
481 Campanian-Maastrichtian (Robertson et al., 2004), similar in age to the oldest basaltic
482 rocks in the Kyrenia Range. The Miocene sediments are structurally intercalated. The
483 basalts of the Misis-Andırın Complex are tholeiitic, with mildly enriched, subduction-

related characteristics (e.g. moderate La/Nb ratios), and are interpreted as of back-arc basin origin (Floyd et al., 1991, 1992), which is consistent with our interpretation of the Kyrenia Range basalts (Fig. 14a-b, Supplementary Figure S6).

The Kyrenia Range is backed to the north by the Alanya Massif (Fig. 1), which has been interpreted as a continental fragment, separate from the Tauride carbonate platform to the north (Robertson et al., 2012a; Çetinkaplan et al., 2016). The Alanya Massif is represented by the Late Precambrian-Mesozoic, mainly meta-sedimentary rocks. High pressure-low temperature (HP-LT) metamorphics document Late Cretaceous (82-80 Ma) subduction/exhumation (Oberhänsli et al., 2010, 2013). This is inferred to have taken place along the northern, active margin of the Southern Neotethys (Çetinkaplan et al., 2016; Robertson and Parlak, 2020). The felsic volcanics (Fourkovouno Formation) in the Kyrenia Range could have erupted along the southern (trailing) margin of the Alanya microcontinent where they could have escaped metamorphism. However, they could also represent a separate more southerly unit.

Late Cretaceous continental margin arc-type magmatic rocks also occur extensively farther east, in SE Turkey, as two separate belts.

The most southerly, known Late Cretaceous arc rocks in SE Turkey are exposed in the Helete area, part of a zone of frontal thrust sheets (Yıldırım, 2015; Nurlu et al., 2016) (Fig. 1). Two thrust assemblages are exposed, both cut by granites that have been radiometrically dated at 93-83 Ma (Cenomanian-Campanian) (Nurlu et al., 2016). The lower of the assemblages, the Helete volcanics, comprises arc-related basaltic andesite, andesite, dacite, rhyolite and common tuffaceous rocks. The upper of the two thrust assemblages is a dismembered Late Cretaceous supra-subduction zone-type ophiolite (Meydan ophiolite) (Yıldırım, 2015; Nurlu et al., 2016). The cross-cutting intrusive rocks sealed the thrust sheets during the Late Cretaceous. The granitic rocks are interpreted as a calc-alkaline, subduction-related suite that formed by mixing of mantle and crustal sources, as suggested by Sr-Nd-Pb isotopic data (Nurlu et al.,

2016). The granitic rocks exhibit relatively flat chondrite-normalised REE patterns, similar to E-MORB (Nurlu et al., 2016) (Supplementary Figure S6). In contrast, the chondrite-normalised REE patterns of the Kyrenia Range felsic rocks are similar to UCC (Fig. 6c), suggesting a more continentally-influenced setting.

The second belt of arc-related magmatism is located up to c. 100 km farther north and north-east, in the Keban (Malatya), Göksun (Kahramanmaraş) and Baskil (Elazığ) regions (Fig. 1) (Yazgan and Chessex, 1991; Parlak, 2006; Rızaoğlu et al., 2009; Karaoğlu et al., 2013, 2016). Arc-granitic rocks (Baskil intrusives) cut the Late Cretaceous Göksun ophiolite and equivalents (i.e. Berit meta-ophiolite (=North Berit ophiolite), Kömürhan ophiolite and İspendere ophiolite) that are inferred to have formed above a northward-dipping subduction zone (Parlak, 2006). Similar arc granites also cut the structurally over-riding Malayta metamorphic unit, which is interpreted as the southern part of the Tauride microcontinent. The Göksun ophiolite (and equivalents) and the Malayta metamorphic (Tauride) units were juxtaposed along an active continental margin when the arc-granitic rocks were intruded (Robertson et al., 2006; Rızaoğlu et al., 2009; Karaoğlu et al., 2013, 2016).

The extrusive rocks of the Göksun and related ophiolites can also be broadly correlated with widespread assemblages of basic igneous rocks and volcanogenic sediments in the Malatya-Elazığ region, known as the Yüksekova Complex (Perincek and Özkaya, 1981; Aktaş and Robertson, 1984). The basalts have a tholeiitic to tholeiitic-transitional character, variable Zr/Y (1.5-6) ratios and mantle-array-aligned Sr-Nd isotopes (Ural et al., 2015). These basic volcanics were derived from a depleted mantle source without a significant crustal contribution; i.e. from an intra-oceanic setting within the Southern Neotethys (Ural et al., 2015).

The arc rocks that cut both the Göksun (and related) ophiolites and the Malatya Metamorphics (Baskil intrusives) in the northern belt are mainly hornblende-biotite granodiorites and 'normal' granites, dated radiometrically at 88-82 Ma (Santonian-

Campanian) (Parlak, 2006; Rızaoğlu et al., 2009; Karaoğlu et al., 2016); i.e. up to c. 5 Ma younger than Helete area granites in the south. These southerly granitic rocks have I-type, calc-alkaline arc affinities, with both mantle and crustal-derivation features (Nurlu et al., 2016). The Baskil intrusions of the northern belt are relatively enriched in LREE (Parlak, 2006; Rızaoğlu et al., 2009) compared to the Helete granites of the south, which suggests a more crustally influenced source during magma genesis for the former. Similarly, the Kyrenia Range basalts have relatively high Zr/Y (2.8-7.7) and Nb/Y (Figs. 12b, 13c), suggestive of a magmatic contribution from continental crust or subducted continentally derived sediments (Zindler and Hart, 1986; Pearce, 1996).

The northerly arc has been explained by northward subduction of Mesozoic oceanic basin (Berit ocean) that was located between two continental units; i.e. the Bitlis and Pütürge massifs in the south and the Tauride continent represented by the Malatya Metamorphics in the north (Robertson et al., 2012a; Karaoğlu et al., 2013, 2016; Barrier et al., 2018). The Late Cretaceous arc-related intrusive magmatism of the northern belt in SE Turkey (Baskil intrusives) evolved towards collisional, including shoshonitic compositions (74-72 Ma), which are interpreted to indicate continental and post-collisional settings (Kuşçu et al., 2013; Ertürk et al., 2018; Sar et al., 2019). The shoshonitic compositions of some of the granitic rocks, together with the evidence of HP/LT metamorphism, are consistent with the collision of the Tauride carbonate platform (i.e. Malatya Metamorphics) to the north with the Bitlis and Pütürge continental units to the south. However, the origins of shoshonites remain controversial because the necessary partial melting of mantle and interaction with subduction-related fluids can take place in a variety of pre-, syn- and post-collisional settings (e.g. Campbell et al., 2014).

Several authors have correlated the Late Cretaceous Göksun and related ophiolites in the north with the Meydan ophiolite (Berit meta-ophiolite of Yılmaz et al., 1993) in the south. If correct, the granitic intrusives in both belts originated as different

parts of a single arc lineament that straddled the northerly active continental margin of the oceanic basin. The southerly arc and ophiolitic rocks (Helete-Meydan) were emplaced southwards during the latest Cretaceous, although further southward thrusting took place, mainly during the Eocene and Miocene (Perinçek and Kozlu, 1984; Yılmaz et al., 1993).

In many reconstructions, oceanic crust still remained between the Bitlis-Pütürge continental units and Arabia until the Miocene (Yılmaz, 1993; Robertson et al., 2012a; Barrier et al., 2018; van Hinsbergen et al., 2020). The Bitlis and Pütürge massifs have been generally correlated with the Alanya metamorphic massif to the north of Cyprus (with or without a direct continuation) (Çetinkaplan et al., 2016). If correct, the Late Cretaceous Kyrenia Range felsic arc volcanism (and that of the Kannaviou Formation) could have been located to south of the Alanya continent, with a possible eastward continuation to south of the Bitlis and Pütürge continental units.

Paleogene magmatic rocks are also present in SE Turkey. Eocene volcanic rocks and associated minor intrusions, known as the Maden Complex, unconformably overlie and cut the Pütürge and Bitlis metamorphosed continental units (Fig. 1) (Hempton, 1985; Yazgan and Chessex, 1991; Aktaş and Robertson, 1984, 1990; Yılmaz, 1993; Elmas and Yılmaz, 2003; Robertson et al., 2006; Ertürk et al., 2018). The volcanics show enrichment in LILEs and relative depletion in Nb, Ta and Ti compared to MORB (Ertürk et al., 2018), similar to a minority of the subduction-related volcanics in the Kyrenia Range (see Supplementary Figure S6). The Eocene Maden Complex magmatism is widely proposed to relate to back-arc rifting of the (Eurasian) active continental margin during the later stages of northward subduction of the Southern Neotethys, in keeping with the widespread view that collision with Arabia was delayed until the Miocene (Aktaş and Robertson, 1984, 1990; Yılmaz, 1993; Yılmaz et al., 1993; Yiğitbaş and Yılmaz, 1996; Robertson et al., 2006, 2007; van Hinsbergen et al., 2020). Oblique convergence, proposed by several authors (Aktaş and Robertson, 1990;

Elmas and Yılmaz, 2003), could have resulted in along-strike variation in the geochemical signatures of extension versus subduction in the Kyrenia Range, although more geochemical work is needed, regionally and globally to test this hypothesis.

Farther north, Eocene granitic rocks locally cut the continental Malatya metamorphic unit (Doğanşehir region) (Perinçek and Kozlu, 1984; Karaoğlu et al., 2013, 2016). However, no extrusive equivalents are known and, as noted above, the magmatic rocks in this lineament are unlikely to correlate directly with the Kyrenia Range.

During the Miocene, the active continental margin bordering the remnant Southern Neotethys collided with the Arabian passive continental margin to the south. In both the Kyrenia Range and SE Turkey, the Late Cretaceous and Paleogene magmatic rocks are exposed close the thrust front (Eurasia), implying that at least tens of kms of fore-arc crust have been lost (subducted or overridden), potentially removing additional frontal arc-related crust, both Late Cretaceous and Paleogene. Thus, the Kyrenia, Kannaviou (W Cyprus) and Maden Complex magmatism could all represent surviving products of a regional active continental margin.

5.5. Tectono-magmatic hypotheses

In the light of the regional setting, the felsic volcanics (Fourkovouno Formation, c. 74 Ma), with continental margin arc affinities, are interpreted to have resulted from northward subduction beneath continental crust, possibly the Alanya continental unit to the north (Çetinkaplan et al., 2016; Robertson et al., 2020). The felsic volcanism is characteristic of relatively advanced subduction, suggesting that earlier eruptive centers existed along the active continental margin. The early volcanism is likely to be represented by the Campanian (c. 80 Ma) arc-derived volcanogenic sediments

(Kannaviou Formation) in western Cyprus (Robertson, 1977; Gilbert and Robertson, 2013; Chen and Robertson, 2019). The underlying mantle wedge (sub-continental lithosphere) was metasomatically enriched during the Late Cretaceous subduction, with implications for the subsequent basaltic volcanism.

For the Maastrichtian to Late Eocene basaltic volcanics, there are three alternative tectono-magmatic models. The first model envisages a marginal basin setting related to northward subduction. Compared to typical back-arc basin basalts (Fig. 14a-b), the Kyrenia Range basalts have relatively high Th/Nb but low La/Nb ratios. The high abundances of Th are likely to reflect a continental crust influence (Floyd et al., 1991). The variation in La/Nb ratios of the basalts could reflect the influence of a within-plate mantle source (i.e. easterly basalts) and/or the arc maturity; i.e. the arc-like western basalts could have erupted closer to the distal (oceanward) edge of the convergent margin. However, a problem with this model is the absence of evidence for a Paleogene volcanic arc to the south. There is no evidence of arc-derived fallout tuff or volcanoclastic sediments in the Maastrichtian-Paleogene basaltic succession, in contrast with, for example, the SW Pacific Mariana and Tonga marginal basins (Clift, 1994; Bryan et al., 2004).

The second model involves an extensional (or transtensional) setting. In this interpretation, the Late Maastrichtian and Paleogene 'enriched' within-plate volcanism was triggered by strike-slip (or transtension) along the active continental margin. One option is that the volcanism was linked to the anticlockwise paleorotation of the Troodos ophiolite to the south during Late Campanian-Early Eocene (Clube et al., 1985; Clube and Robertson, 1986; Morris et al., 2006, 2015; Hodgson et al., 2010). More generally, oblique convergence of the African plate with respect to Eurasia could have resulted in segmentation of the South Neotethyan active margin into subducting segments with arc magmatism, and also strike-slip segments without arc magmatism (Aktaş and Robertson, 1990). A modern example is the Andaman Sea where oblique

convergence between the Asian and the Australian plates resulted in segmented arc magmatism (e.g. Curray et al., 2005). However, there is limited information from different modern settings to test this hypothesis.

The third, preferred, scenario is that both the felsic and the basaltic volcanism are related to northward subduction during stages in the closure of the Mesozoic ocean basin to the south. The Late Cretaceous magmatism can then be explained by regional northward subduction (nearly orthogonal) resulting in continental margin arc volcanism. Slab-derived and residual slab components (fluid, melt) were released into the upper mantle lithosphere (e.g. Pearce, 1983; Harangi et al., 2007). The Late Maastrichtian and Paleogene basaltic volcanics then erupted in an extensional (or transtensional) setting, perhaps related to slab rollback and incipient marginal basin formation. Several different mantle sources are likely to have contributed along the Kyrenia Range. The variable subduction-related signatures were inherited from the previously modified mantle lithosphere in this interpretation. Examples of subduction-influenced settings that were affected by later extension-related melting include the Rio Grande Rift, west USA (e.g. Riecker, 1979) and the Neogene north Aegean-west Anatolia volcanic province (e.g. Şengör and Yılmaz, 1981; Ersoy and Palmer, 2013). In summary, the available evidence is consistent with Late Maastrichtian-Paleogene incipient marginal basin formation along the northern, active margin of an ocean basin (Southern Neotethys) that did not close until the Miocene.

5.6. Testing alternative tectonic models

Finally, the Late Cretaceous-Paleogene volcanism in N Cyprus and SE Turkey can be used to test alternative plate tectonic models, including those by Robertson et al. (2012a), Karaoğlu et al. (2013, 2016), Maffione et al. (2017), Barrier et al. (2018), McPhee and van Hinsbergen (2019) and van Hinsbergen et al. (2020), as summarised in Figure 15.

In Reconstruction 1 (Fig. 15a-c) (Robertson et al., 2012a), the Southern Neotethys rifted in the Triassic, while Paleotethys evolved into the Northern Neotethys (İzmir-Ankara-Erzincan ocean). The Kyrenia Range is restored to the northern passive margin of the Southern Neotethys. Both intra-oceanic and continental margin subduction zones (northward-dipping) were active in the Southern Neotethys during the Late Cretaceous. The continental margin arc volcanism was constructed on, or to the south of one, or more, continental blocks (microcontinents), represented by the Kyrenia-Alanya-Pütürge-Bitlis lineament in the south and the Malatya-Keban lineament farther north. This reconstruction is compatible with our new evidence.

Reconstruction 2 (Fig. 15d-f) (Barrier et al., 2018) has many common features with (1), in particular the arc volcanism along the northern margin of the Southern Neotethys. However, the Kyrenia, Malatya and Pütürge-Bitlis massifs are restored as a single elongate continental unit rather than as microcontinents, as in (1). Also, little or no oceanic crust is inferred between the southerly continental unit and the Tauride carbonate platform to the north (i.e. Göksun and related ophiolites). This is difficult to reconcile with the Late Cretaceous and Eocene granitic magmatism cutting the Mesozoic Malatya (Keban) platform unless large-scale strike-slip and terrane duplication has taken place for which there is little supporting evidence.

In Reconstruction 3 (Fig. 15g-h) (Maffione et al., 2017; van Hinsbergen et al., 2020), the Kyrenia Range is restored to the southern passive margin of the Mesozoic Tethyan ocean. Late Cretaceous ophiolites that are presently located in several c. E-W, sub-parallel belts (i.e. N and S of the Bitlis and Pütürge massifs) originated above a single subduction zone that originated within Neotethys, dipping generally to the east or northeast (Stampfli and Borel, 2002; Moix et al., 2008). The subduction zone rolled back generally westwards until it collided with the Tauride and Arabian continents. In the south, the subduction zone broke through the oceanic gap between the African and Tauride continents, 'invaded' and replaced the Southern Neotethys with Late

Cretaceous supra-subduction zone fore-arc crust. Fore-arc ophiolites were then obducted generally northwards and southwards over the opposing continents (van Hinsbergen et al., 2020). Two alternatives are suggested; in one, the Bitlis and Pütürge continental units are correlated with Arabia (Maffione et al., 2017), whereas in the other these units are correlated with the southern margin of the Tauride continent (van Hinsbergen et al., 2020). The Kyrenia Range (Trypa Group) is placed on the N African continental margin in both alternatives. The metamorphism of the Mesozoic carbonate platform in the Kyrenia Range (Trypa Group) resulted from northeastward underthrusting/subduction beneath the obducted forearc Troodos ophiolite (van Hinsbergen et al., 2020). However, there are several problematic aspects: (1) seismic tomography in the Eastern Mediterranean supports generally northward subduction (e.g. Hafkenscheid et al., 2006), without evidence of a relict southward-dipping slab, as implied by the inferred northward emplacement of some Late Cretaceous ophiolites in S Turkey; (2) it does not allow for the tens to hundreds of km of subduction required to produce the Late Cretaceous arc magmatism in N Cyprus and SE Turkey. The Malatya metamorphic unit, cut by Late Cretaceous granites, is instead restored to the lower, downgoing plate (i.e. beneath the south margin of the Tauride continent) implying northward subduction; (3) the model (van Hinsbergen et al., 2020) does not allow for any Late Cretaceous continental arc-related magmatism in northern Cyprus, SE Turkey or Iran (e.g. Agard et al., 2011); (4) in this model, the proposed Late Eocene-Miocene (post 45 Ma) northward subduction is too young to explain the Late Cretaceous-Paleogene magmatic rocks in the Kyrenia Range and SE Turkey; (5) arc magmatism of Oligocene-Miocene age in S Turkey is absent, although this would be expected if significant northward subduction took place during this time, as in model 3.

723 6. Conclusions

- 724 ● Latest Cretaceous mainly felsic volcanic rocks and latest Cretaceous-Paleogene
725 mainly basaltic volcanic rocks are exposed throughout the Kyrenia Range,
726 separated by a thrust.
- 727 ● Mapping and logging of the key, well-exposed western Kyrenia Range segment
728 shows that the Late Cretaceous felsic arc volcanics occur as variably
729 dismembered thrust sheets in the southerly, frontal part of thrust belt.
- 730 ● The primary eruptive age of the felsic arc volcanics (Fourkovouno Formation) is c.
731 74.0 Ma (Late Campanian), based on U-Pb dating of zircons.
- 732 ● The latest Cretaceous-Paleogene basalts vary geochemically along the Kyrenia
733 Range (E-W). The volcanics in the east resemble normal rift products. In contrast,
734 the volcanics in the west have a subordinate subduction-related signature (e.g.
735 negative Nb anomaly).
- 736 ● Sr-Nd-Hf isotopic data for the latest Cretaceous-Paleogene basalts suggest that
737 they were derived from several OIB-like mantle sources, with the involvement of a
738 crustally-derived (recycled) component.
- 739 ● The Kyrenia Range felsic volcanics show some geochemical similarities with the
740 late Cretaceous arc-related rocks in S Turkey (Helete granite; 93-83 Ma). These
741 felsic volcanics are interpreted to represent a fragment of a mature magmatic arc
742 that was active along a regional-scale active continental margin. Regional
743 reconstructions suggest that the Helete granites and related ophiolitic rocks were
744 emplaced southwards from an oceanic basin (Berit ocean) to the north.
- 745 ● The latest Cretaceous-Paleogene volcanics in the Kyrenia Range have
746 geochemical similarities with the Middle Eocene (c. 47 Ma) basaltic rocks in S
747 Turkey (Maden Complex), suggesting an along-strike continuation of the same
748 active continental margin.

● The latest Cretaceous-Paleogene basaltic volcanics in the Kyrenia Range are interpreted as the products of an incipient marginal basin that developed during northward subduction, possibly related to oblique convergence, prior to collision with Arabia during the Miocene.

● In the light of alternative regional tectonic models, the Late Cretaceous and Paleogene magmatic rocks of N Cyprus are interpreted to represent stages in the development of the northerly active continental margin of the Southern Neotethys.

Acknowledgements

We thank Richard Hinton, Steffen Kutterolf and Dick Kroon for scientific discussion. Mike Hall is thanked for thin section and polished blocks preparation. Nick Odling kindly assisted with the XRF analysis. Antony Morris is thanked for providing the samples of basalt from drill cores that were originally collected for paleomagnetic studies. Rory McKavney kindly assisted with the fieldwork in northern Cyprus. The first author gratefully acknowledges the receipt of a joint studentship of the Principal's Career Development PhD Scholarship and Edinburgh Global Research Scholarship. The authors are grateful for financial support via the Natural Environment Research Council Ion Microprobe Facility (to A.H.F. Robertson) to carry out the secondary ion mass spectrometry U-Pb dating of detrital zircons. Fieldwork and geochemical analysis were aided by financial support from the International Association of Sedimentologists [Postgraduate Grant Scheme], the Mineralogical Society of Great Britain and Ireland [Postgraduate Student Bursary Awards], and the Edinburgh Geological Society [Clough Fund], all to the first author. Additional financial support was provided by the John Dixon Memorial Fund. Osman Parlak is thanked for discussion of SE Turkey geology. The manuscript benefitted from comments by Pamela Kempton, Fatih Karaoğlu and the editor, Greg Shellnutt.

775 References

- 776 Adamia, S.A., Lordkipanidze, M., Zakariadze, G., 1977. Evolution of an active
777 continental margin as exemplified by the Alpine history of the Caucasus.
778 Tectonophysics 40, 183-199.
- 779 Agard, P., Omrani, J., Jolivet, L., Whitechurch, H., Vrielynck, B., Spakman, W., Monié,
780 P., Meyer, B., Wortel, R., 2011. Zagros orogeny: a subduction-dominated process.
781 Geological Magazine 148, 692-725.
- 782 Aktaş, G., Robertson, A.H.F., 1984. The Maden Complex, SE Turkey: evolution of a
783 Neotethyan active margin, in: Dixon, J.E., Robertson, A.H.F. (Eds.), The
784 Geological Evolution of the Eastern Mediterranean. Geological Society of London,
785 Special Publications 17, pp. 375-402.
- 786 Aktaş, G., Robertson, A.H.F., 1990. Tectonic evolution of the Tethys suture zone in SE
787 Turkey: evidence from the petrology and geochemistry of Late Cretaceous and
788 Middle Eocene extrusives, in: Malpas, J., Moores, E.M., Panayiotou, A.,
789 Xenophontos, C. (Eds.), Ophiolites-Oceanic Crustal Analogues. Proceedings of
790 the International Symposium 'Troodos 1987'. Cyprus Geological Survey
791 Department, Nicosia, pp. 311-329.
- 792 Baroz, F., 1979. Etude géologique dans le Pentadaktylos et la Mesaoria (Chypre
793 Septentrionale). Doctor of Science Thesis (Published). Université de Nancy,
794 France (434 pp).
- 795 Baroz, F., 1980. Volcanism and continent-island arc collision in the Pentadaktylos
796 range, Cyprus, in: Panayiotou, A. (Ed.), Ophiolites: Proceedings of the
797 International Ophiolite Symposium. Cyprus Ministry of Agriculture and Natural
798 Resources, Geology Survey Department, Nicosia, Cyprus, pp. 73-85.
- 799 Barrier, E., Vrielynck, B., Brouillet, J.F., Brunet, M.F., 2018. Paleotectonic
800 Reconstruction of the Central Tethyan Realm. Tectono-Sedimentary-Palinspastic
801 Maps from Late Permian to Pliocene. Atlas of 20 maps (scale 1/15000000).

802 CCGM/CGMW, Paris.

803 Beccaluva, L., Macciotta, G., Piccardo, G., Zeda, O., 1989. Clinopyroxene composition
804 of ophiolite basalts as petrogenetic indicator. *Chemical Geology* 77, 165-182.

805 Bowman, N., Van Otterloo, J., Cairns, C., Taylor, D., Cas, R., 2019. Complex evolution
806 of volcanic arcs: The lithofacies and palaeogeography of the Cambrian Stavelo
807 Arc, Delamerian Fold Belt, Western Victoria. *Journal of Volcanology and
808 Geothermal Research* 373, 120-132.

809 Brune, S., Popov, A.A., Sobolev, S.V., 2012. Modeling suggests that oblique extension
810 facilitates rifting and continental break-up. *Journal of Geophysical Research: Solid
811 Earth* 117. doi: 10.1029/2011JB008860

812 Bryan, S., Cook, A., Evans, J., Colls, P., Wells, M., Lawrence, M., Jell, J., Greig, A.,
813 Leslie, R., 2004. Pumice rafting and faunal dispersion during 2001–2002 in the
814 Southwest Pacific: record of a dacitic submarine explosive eruption from Tonga.
815 *Earth and Planetary Science Letters* 227, 135-154.

816 Campbell, I.H., Stepanov, A.S., Liang, H.-Y., Allen, C.M., Norman, M.D., Zhang, Y.-Q.,
817 Xie, Y.-W., 2014. The origin of shoshonites: new insights from the Tertiary high-
818 potassium intrusions of eastern Tibet. *Contributions to Mineralogy and Petrology*
819 167. doi: 10.1007/s00410-014-0983-9

820 Çetinkaplan, M., Pourteau, A., Candan, O., Koralay, O.E., Oberhänsli, R., Okay, A.I.,
821 Chen, F., Kozlu, H., Şengün, F., 2016. P–T–t evolution of eclogite/blueschist facies
822 metamorphism in Alanya Massif: time and space relations with HP event in Bitlis
823 Massif, Turkey. *International Journal of Earth Sciences* 105, 247-281.

824 Chappell, B.W., White, A.J.R., 1992. I- and S-type granites in the Lachlan Fold Belt.
825 *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*
826 83, 1-26.

827 Chen, G., Robertson, A.H.F., 2019. Provenance and magmatic-tectonic setting of
828 Campanian-aged volcanoclastic sandstones of the Kannaviou Formation in
829 western Cyprus: Evidence for a South-Neotethyan continental margin volcanic arc.

830 Sedimentary Geology 388, 114-138.

831 Chen, S.-s., Liu, J.-q., Chen, S.-s., Guo, Z.-f., Sun, C.-q., 2015. Variations in the
832 geochemical structure of the mantle wedge beneath the northeast Asian marginal
833 region from pre- to post-opening of the Japan Sea. *Lithos* 224-225, 324-341.

834 Clift, P., 1994. Controls on the Sedimentary and Subsidence History of an Active Plate
835 Margin: An Example from the Tonga Arc (Southwest Pacific), in: Hawkins, J.,
836 Parson, L., Allan, J. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific*
837 *Results*. Ocean Drilling Program 135, College Station, TX, pp. 173–189.

838 Clube, T.M.M., Creer, K.M., Robertson, A.H.F., 1985. Palaeorotation of the Troodos
839 microplate, Cyprus. *Nature* 317, 522-525.

840 Clube, T.M.M., Robertson, A.H.F., 1986. The palaeorotation of the Troodos microplate,
841 Cyprus, in the Late Mesozoic-Early Cenozoic plate tectonic framework of the
842 Eastern Mediterranean. *Surveys in Geophysics* 8, 375-437.

843 Condie, K.C., 2003. Incompatible element ratios in oceanic basalts and komatiites:
844 tracking deep mantle sources and continental growth rates with time.
845 *Geochemistry, Geophysics, Geosystems* 4, 1005. doi:10.1029/2002GC000333

846 Condie, K.C., 2005. High field strength element ratios in Archean basalts: a window to
847 evolving sources of mantle plumes? *Lithos* 79, 491-504.

848 Curray, J.R., 2005. Tectonics and history of the Andaman Sea region. *Journal of Asian*
849 *Earth Sciences* 25, 187-232.

850 Dercourt, J., Ricou, L.E., Vrielynck, B., 1993. *Atlas of Peri-Tethys Palaeogeographical*
851 *Maps*. CCGM/CGMW, Paris (268 pp).

852 Dewey, J.F., 1969. Evolution of the Appalachian/Caledonian orogen. *Nature* 222, 124-
853 129.

854 Ducloz, C., 1972. The geology of the Bellapais-Kythrea area of the Central Kyrenia
855 Range. *Bulletin of the Geological Survey Department* 6, Nicosia, Cyprus (75 pp).

856 Elmas, A., Yilmaz, Y., 2003. Development of an oblique subduction zone-tectonic
857 evolution of the Tethys suture zone in southeast Turkey. *International Geology*

858 Review 45, 827-840.

859 Ersoy, E.Y., Palmer, M.R., 2013. Eocene-Quaternary magmatic activity in the Aegean:
860 implications for mantle metasomatism and magma genesis in an evolving orogeny.
861 *Lithos* 180, 5-24.

862 Erturk, M.A., Beyarslan, M., Chung, S.L., Lin, T.H., 2018. Eocene magmatism (Maden
863 Complex) in the Southeast Anatolian Orogenic Belt: Magma genesis and tectonic
864 implications. *Geoscience Frontiers* 9, 1829-1847.

865 Fitton, J.G., Godard, M., 2004. Origin and evolution of magmas on the Ontong Java
866 Plateau, in: Fitton, J.G., Mahoney, J.J., Wallace, P.J., Saunders, A.D. (Eds.),
867 Origin and Evolution of the Ontong Java Plateau. Geological Society of London,
868 Special Publications 229, pp. 151-178.

869 Fitton, J.G., Saunders, A.D., Larsen, L.M., Hardarson, B.S., Norry, M.J., 1998. Volcanic
870 rocks from the southeast Greenland margin at 63°N: composition, petrogenesis
871 and mantle sources, in: Saunders, A.D., Larsen, H.C., Wise, S.W. (Eds.),
872 Proceedings of the Ocean Drilling Program, Scientific Results. Ocean Drilling
873 Program 152, College Station, TX, pp. 331-350.

874 Floyd, P., Kelling, G., Gökçen, S., Gökçen, N., 1991. Geochemistry and tectonic
875 environment of basaltic rocks from the Misis ophiolitic mélange, south Turkey.
876 *Chemical Geology* 89, 263-280.

877 Floyd, P., Kelling, G., Gökçen, S., Gökçen, N., 1992. Arc-related origin of volcanoclastic
878 sequences in the Misis Complex, Southern Turkey. *The Journal of Geology* 100,
879 221-230.

880 Garfunkel, Z., 1998. Constrains on the origin and history of the Eastern Mediterranean
881 basin. *Tectonophysics* 298, 5-35.

882 Gilbert, M.F., Robertson, A.H.F., 2013. Field relations, geochemistry and origin of the
883 Upper Cretaceous volcanoclastic Kannaviou Formation in western Cyprus:
884 evidence of a southerly Neotethyan volcanic arc, in: Robertson, A.H.F., Parlak, O.,
885 Ünlügenç, U.C. (Eds.), *Geological Development of Anatolia and the Easternmost*

886 Mediterranean Region. Geological Society of London, Special Publications 372,
887 pp. 273-298.

888 Gorton, M.P., Schandl, E.S., 2000. From continents to island arcs: a geochemical index
889 of tectonic setting for arc-related and within-plate felsic to intermediate volcanic
890 rocks. *The Canadian Mineralogist* 38, 1065-1073.

891 Hafkenscheid, E., Wortel, M., Spakman, W., 2006. Subduction history of the Tethyan
892 region derived from seismic tomography and tectonic reconstructions. *Journal of*
893 *Geophysical Research: Solid Earth* 111, B08401. doi: 10.1029/2005JB003791

894 Hakyemez, Y., Turhan, N., Sönmez, I., Sümengen, M., 2000. Kuzey Kıbrıs Türk
895 Cumhuriyeti'nin Jeolojisi (Geology of the Turkish Republic of Northern Cyprus).
896 Genel Müdürlüğü Jeoloji Etütleri Diaresi, Maden Tektik ve Arama, Ankara (44 pp).

897 Harangi, S., Downes, H., Thirlwall, M., Gmeling, K., 2007. Geochemistry, Petrogenesis
898 and Geodynamic Relationships of Miocene Calc-alkaline Volcanic Rocks in the
899 Western Carpathian Arc, Eastern Central Europe. *Journal of Petrology* 48, 2261-
900 2287.

901 Hartley, M.E., Thordarson, T., 2013. The 1874-1876 volcano-tectonic episode at Askja,
902 North Iceland: Lateral flow revisited. *Geochemistry, Geophysics, Geosystems* 14,
903 2286-2309.

904 Hastie, A.R., Kerr, A.C., Pearce, J.A., Mitchell, S.F., 2007. Classification of Altered
905 Volcanic Island Arc Rocks using Immobile Trace Elements: Development of the
906 Th–Co Discrimination Diagram. *Journal of Petrology* 48, 2341-2357.

907 Hayward, C., 2011. High spatial resolution electron probe microanalysis of tephra and
908 melt inclusions without beam-induced chemical modification. *The Holocene* 22,
909 119-125.

910 Hempton, M.R., 1985. Structure and deformation history of the Bitlis suture near Lake
911 Hazar, southeastern Turkey. *Geological Society of America Bulletin* 96, 233-243.

912 Hodgson, E., Morris, A., Anderson, M., Robertson, A., 2010. First palaeomagnetic
913 results from the Kyrenia Range terrane of northern Cyprus and their implication

914 for the regional plate tectonic evolution of the eastern Mediterranean, EGU
 915 General Assembly Conference Abstracts, Vienna, Austria, p. 6449.

916 Hu, Z., Gao, S., 2008. Upper crustal abundances of trace elements: a revision and
 917 update. *Chemical Geology* 253, 205-221.

918 Huang, K., Malpas, J., Xenophontos, C., 2007. Geological studies of igneous rocks
 919 and their relationships along the Kyrenia Range, in: Moumani, K., Shawabkeh, K.,
 920 Al-Malabeh, A., Abdelghafoor, M. (Eds.), 6th International Congress of Eastern
 921 Mediterranean Geology, Amman, Jordan, p. 53.

922 Karaoğlu, F., Parlak, O., Hejl, E., Neubauer, F., Kloetzli, U., 2016. The temporal
 923 evolution of the active margin along the Southeast Anatolian Orogenic Belt (SE
 924 Turkey): Evidence from U–Pb, Ar–Ar and fission track chronology. *Gondwana*
 925 *Research* 33, 190-208.

926 Karaoğlu, F., Parlak, O., Robertson, A., Thöni, M., Klötzli, U., Koller, F., Okay, A.İ.,
 927 2013. Evidence of Eocene high-temperature/high-pressure metamorphism of
 928 ophiolitic rocks and granitoid intrusion related to Neotethyan subduction
 929 processes (Doğanşehir area, SE Anatolia), in: Robertson, A.H.F., Parlak, O.,
 930 Ünlügenç, U.C. (Eds.), *Geological Development of Anatolia and the Easternmost*
 931 *Mediterranean Region*. Geological Society of London, Special Publications 372,
 932 pp. 249-272.

933 Keller, R.A., Fisk, M.R., Smellie, J.L., Strelin, J.A., Lawver, L.A., 2002. Geochemistry
 934 of back arc basin volcanism in Bransfield Strait, Antarctica: Subducted
 935 contributions and along-axis variations. *Journal of Geophysical Research: Solid*
 936 *Earth* 107. doi: 10.1029/2001JB000444

937 Kelly, N., Hinton, R., Harley, S., Appleby, S., 2008. New SIMS U–Pb zircon ages from
 938 the Langavat Belt, South Harris, NW Scotland: implications for the Lewisian
 939 terrane model. *Journal of the Geological Society* 165, 967-981.

940 Kempton, P.D., Downes, H., Lustrino, M., 2018. Pb and Hf isotope evidence for mantle

941 enrichment processes and melt interactions in the lower crust and lithospheric
 942 mantle in Miocene orogenic volcanic rocks from Monte Arcuentu (Sardinia, Italy).
 943 *Geosphere* 14, 926-950.

944 Kuşcu, İ., Tosdal, R.M., Gencalioğlu-Kuşcu, G., Friedman, R., Ullrich, T.D., 2013. Late
 945 Cretaceous to Middle Eocene Magmatism and Metallogeny of a Portion of the
 946 Southeastern Anatolian Orogenic Belt, East-Central Turkey. *Economic Geology*
 947 108, 641-666.

948 Le Pichon, X., 1982. Land-locked oceanic basins and continental collision: the Eastern
 949 Mediterranean as a case example, in: Hsü, K.J. (Ed.), *Mountain building*
 950 *processes*. Academic Press, New York, pp. 201-211.

951 Li, C.-F., Li, X.-H., Li, Q.-L., Guo, J.-H., Li, X.-H., Yang, Y.-H., 2012. Rapid and precise
 952 determination of Sr and Nd isotopic ratios in geological samples from the same
 953 filament loading by thermal ionization mass spectrometry employing a single-step
 954 separation scheme. *Analytica Chimica Acta* 727, 54-60.

955 Ludwig, K.R., 2012. Users manual for Isoplot 3.75. A geochronological toolkit for
 956 Microsoft Excel. Berkeley Geochronology Centre, Special Publication No. 5,
 957 Berkeley.

958 Maffione, M., van Hinsbergen, D.J., de Gelder, G.I., van der Goes, F.C., Morris, A.,
 959 2017. Kinematics of Late Cretaceous subduction initiation in the Neo-Tethys
 960 Ocean reconstructed from ophiolites of Turkey, Cyprus, and Syria. *Journal of*
 961 *Geophysical Research: Solid Earth* 122, 3953-3976.

962 McCay, G.A., Robertson, A.H.F., 2013. Upper Miocene–Pleistocene deformation of the
 963 Girne (Kyrenia) Range and Dar Dere (Ovgos) lineaments, northern Cyprus: role
 964 in collision and tectonic escape in the easternmost Mediterranean region, in:
 965 Robertson, A.H.F., Parlak, O., Ünlügenç, U.C. (Eds.), *Geological Development of*
 966 *Anatolia and the Easternmost Mediterranean Region*. Geological Society of
 967 London, Special Publications 372, pp. 421-445.

968 McCay, G.A., Robertson, A.H.F., Kroon, D., Raffi, I., Ellam, R.M., Necdet, M., 2013.
 969 Stratigraphy of Cretaceous to Lower Pliocene sediments in the northern part of
 970 Cyprus based on comparative $^{87}\text{Sr}/^{86}\text{Sr}$ isotopic, nannofossil and planktonic
 971 foraminiferal dating. *Geological Magazine* 150, 333-359.

972 McPhee, P.J., van Hinsbergen, D.J., 2019. Tectonic reconstruction of Cyprus reveals
 973 Late Miocene continental collision of Africa and Anatolia. *Gondwana Research* 68,
 974 158-173.

975 Moix, P., Beccaleto, L., Kozur, H.W., Hochard, C., Rosselet, F., Stampfli, G.M., 2008.
 976 A new classification of the Turkish terranes and sutures and its implication for the
 977 paleotectonic history of the region. *Tectonophysics* 451, 7-39.

978 Moore, T.A., 1960. The geology and mineral resources of the Astromeritis-Kormakiti
 979 area. Geological Survey Department, Memoir 6, Nicosia, Cyprus (96 pp).

980 Morris, A., Anderson, M.W., Inwood, J., Robertson, A.H., 2006. Palaeomagnetic
 981 insights into the evolution of Neotethyan oceanic crust in the eastern
 982 Mediterranean, in: Robertson, A.H.F., Mountrakis, D. (Eds.), *Tectonic
 983 Development of the Eastern Mediterranean Region*. Geological Society of London,
 984 Special Publications 260, pp. 351-372.

985 Morris, A., Robertson, A.H.F., Anderson, M.W., Hodgson, E., 2015. Did the Kyrenia
 986 Range of northern Cyprus rotate with the Troodos–Hatay microplate during the
 987 tectonic evolution of the eastern Mediterranean? *International Journal of Earth
 988 Sciences* 105, 399-415.

989 Nisbet, E.G. Pearce, J.A., 1977. Clinopyroxene composition in mafic lavas from
 990 different tectonic settings. *Contributions to Mineralogy and Petrology* 63, 149-160.

991 Nurlu, N., Parlak, O., Robertson, A.H.F., von Quadt, A., 2016. Implications of Late
 992 Cretaceous U–Pb zircon ages of granitic intrusions cutting ophiolitic and
 993 volcanogenic rocks for the assembly of the Tauride allochthon in SE Anatolia
 994 (Helete area, Kahramanmaraş Region, SE Turkey). *International Journal of Earth
 995 Sciences* 105, 283-314.

996 Oberhänsli, R., Candan, O., Bousquet, R., Rimmerle, G., Okay, A., Goff, J., 2010. Alpine
 997 high pressure evolution of the eastern Bitlis complex, SE Turkey, in: Sosson, M.,
 998 Kaymakci, N., Stephenson, R.A., Bergerat, F., Starostenko, V. (Eds.),
 999 Sedimentary Basin Tectonics from the Black Sea and Caucasus to the Arabian
 1000 Platform. Geological Society of London, Special Publications 340, pp. 461-483.

1001 Oberhänsli, R., Koralay, E., Candan, O., Pourteau, A., Bousquet, R., 2013. Late
 1002 Cretaceous eclogitic high-pressure relics in the Bitlis Massif. *Geodinamica Acta*
 1003 26, 175-190.

1004 Parlak, O., 2006. Geodynamic significance of granitoid magmatism in the southeast
 1005 Anatolian orogen: geochemical and geochronological evidence from Göksun–Afşin
 1006 (Kahramanmaraş, Turkey) region. *International Journal of Earth Sciences* 95, 609-
 1007 627.

1008 Pearce, J.A., 1975. Basalt geochemistry used to investigate past tectonic
 1009 environments on Cyprus. *Tectonophysics* 25, 41-67.

1010 Pearce, J.A., 1983. Role of the sub-continental lithosphere in magma genesis at active
 1011 continental margins, in: Hawkersworth, C.J., Norry, M.J. (Eds.), *Continental*
 1012 *Basalts and Mantle Xenoliths*. Shiva, Cheshire, UK, pp. 230-249.

1013 Pearce, J.A., 1996. A user's guide to basalt discrimination diagrams, in: Wyman, D.A.
 1014 (Ed.), *Trace Element Geochemistry of Volcanic Rocks: Applications for Massive*
 1015 *Sulphide Exploration*. Geological Association of Canada, Short Course Notes 12,
 1016 pp. 79-113.

1017 Pearce, J.A., Cann, J., 1973. Tectonic setting of basic volcanic rocks determined using
 1018 trace element analyses. *Earth and Planetary Science Letters* 19, 290-300.

1019 Pearce, J.A., Harris, N.B., Tindle, A.G., 1984. Trace element discrimination diagrams
 1020 for the tectonic interpretation of granitic rocks. *Journal of Petrology* 25, 956-983.

1021 Pearce, J.A., Norry, M.J., 1979. Petrogenetic implications of Ti, Zr, Y, and Nb variations
 1022 in volcanic rocks. *Contributions to Mineralogy and Petrology* 69, 33-47.

1023 Peccerillo, A., 2017. *Cenozoic Volcanism in the Tyrrhenian Sea Region*. Springer

1024 International Publishing, Cham (399 pp).

1025 Perinçek, D., Kozlu, H., 1984. Stratigraphy and structural relations of the units in the
1026 Afşin-Elbistan-Doğanşehir region (Eastern Taurus), in: Tekeli, O., Göncüoğlu, M.C.
1027 (Eds.), *Geology of the Taurus Belt: Proceedings of the International Symposium*.
1028 Maden Tetkik ve Arama Enstitüsü, Ankara, pp. 181-198.

1029 Perinçek, D., Özkaya, İ., 1981. Tectonic evolution of the northern margin of Arabian
1030 plate. *Bulletin of Institute of Earth Science, Hacettepe University* 8, 91-101.

1031 Rice, S.P., Robertson, A.H.F., Ustaömer, T., 2006. Late Cretaceous-Early Cenozoic
1032 tectonic evolution of the Eurasian active margin in the Central and Eastern
1033 Pontides, northern Turkey, in: Robertson, A.H.F., Mountrakis, D. (Eds.), *Tectonic
1034 Development of the Eastern Mediterranean Region*. Geological Society of London,
1035 Special Publications 260, pp. 413-445.

1036 Riecker, R.E., 1979. Rio Grande Rift: Tectonics and Magmatism. American
1037 Geophysical Union, Washington, DC (438 pp).

1038 Rızaoğlu, T., Parlak, O., Höck, V., Koller, F., Hames, W.E., Billor, Z., 2009. Andean-
1039 type active margin formation in the eastern Taurides: Geochemical and
1040 geochronological evidence from the Baskil granitoid (Elazığ, SE Turkey).
1041 *Tectonophysics* 473, 188-207.

1042 Robertson, A.H.F., 1977. The Kannaviou Formation, Cyprus: volcanoclastic
1043 sedimentation of a probable Late Cretaceous volcanic arc. *Journal of the
1044 Geological Society* 134, 269-292.

1045 Robertson, A.H.F., Dixon, J.E., 1984. Introduction: aspects of the geological evolution
1046 of the Eastern Mediterranean, in: Dixon, J.E., Robertson, A.H.F. (Eds.), *The
1047 Geological Evolution of the Eastern Mediterranean*. Geological Society of London,
1048 Special Publications 17, pp. 1-74.

1049 Robertson, A.H.F., Kinnaird, T.C., 2016. Structural development of the central Kyrenia
1050 Range (north Cyprus) in its regional setting in the eastern Mediterranean region.
1051 *International Journal of Earth Sciences* 105, 417-437.

1052 Robertson, A.H.F., Parlak, O., 2020. Late Cretaceous-Palaeocene subduction-
 1053 collision-exhumation of a microcontinent along the northern, active margin of
 1054 South Neotethys: evidence from the Alanya Massif and the adjacent Antalya
 1055 Complex (S Turkey). *Journal of Asian Earth Sciences*, 104467. doi:
 1056 10.1016/j.jseaes.2020.104467

1057 Robertson, A.H.F., Parlak, O., Kinnaird, T.C., Taslı, K., Dumitrica, P., 2020. Cambrian-
 1058 Eocene pre-rift, pulsed rift, passive margin and emplacement processes along the
 1059 northern margin of the Southern Neotethys: Evidence from the Antalya Complex
 1060 in the Alanya Window (S Turkey). *Journal of Asian Earth Sciences: X* 3, 100026.
 1061 doi: 10.1016/j.jaesx.2020.100026

1062 Robertson, A.H.F., Parlak, O., Rizaoglu, T., Ünlügenç, Ü., İnan, N., Taslı, K., Ustaömer,
 1063 T., 2007. Tectonic evolution of the South Tethyan ocean: evidence from the
 1064 Eastern Taurus Mountains (Elazığ region, SE Turkey), in: Ries, A.C., Butler,
 1065 R.W.H., Graham, R.H. (Eds.), *Deformation of the Continental Crust: The Legacy*
 1066 *of Mike Coward*. Geological Society of London, Special Publications 272, pp. 231-
 1067 270.

1068 Robertson, A.H.F., Parlak, O., Ustaömer, T., 2012a. Overview of the Palaeozoic-
 1069 Neogene evolution of Neotethys in the Eastern Mediterranean region (southern
 1070 Turkey, Cyprus, Syria). *Petroleum Geoscience* 18, 381-404.

1071 Robertson, A.H.F., Taslı, K., İnan, N., 2012b. Evidence from the Kyrenia Range, Cyprus,
 1072 of the northerly active margin of the Southern Neotethys during Late Cretaceous–
 1073 Early Cenozoic time. *Geological Magazine* 149, 264-290.

1074 Robertson, A.H.F., Ünlügenç, Ü.C., İnan, N., Taslı, K., 2004. The Misis-Andırın
 1075 Complex: a Mid-Tertiary melange related to late-stage subduction of the Southern
 1076 Neotethys in S Turkey. *Journal of Asian Earth Sciences* 22, 413-453.

1077 Robertson, A.H.F., Ustaömer, T., Parlak, O., Ünlügenç, U.C., Taşlı, K., İnan, N., 2006.
 1078 The Berit transect of the Tauride thrust belt, S Turkey: Late Cretaceous-Early
 1079 Cenozoic accretionary/collisional processes related to closure of the Southern

1080 Neotethys. *Journal of Asian Earth Sciences* 27, 108-145.

1081 Robertson, A.H.F., Woodcock, N.H., 1979. Mamonia Complex, southwest Cyprus:
 1082 Evolution and emplacement of a Mesozoic continental margin. *Geological Society*
 1083 *of America Bulletin* 90, 651-665.

1084 Robertson, A.H.F., Woodcock, N.H., 1986. The role of the Kyrenia Range Lineament,
 1085 Cyprus, in the geological evolution of the eastern Mediterranean area.
 1086 *Philosophical Transactions of the Royal Society of London. Series A,*
 1087 *Mathematical and Physical Sciences* 317, 141-177.

1088 Rollinson, H.R., 1993. *Using Geochemical Data: Evaluation, Presentation,*
 1089 *Interpretation.* Routledge, London (384 pp).

1090 Rubatto, D., 2002. Zircon trace element geochemistry: partitioning with garnet and the
 1091 link between U–Pb ages and metamorphism. *Chemical Geology* 184, 123-138.

1092 Rudnick, R.L., Gao, S., 2003. Composition of the continental crust, in: Holland, H.D.,
 1093 Turekian, K.K. (Eds.), *Treatise on geochemistry.* Elsevier-Pergamon 3, Oxford, pp.
 1094 1-64.

1095 Sar, A., Ertürk, M.A., Rizeli, M.E., 2019. Genesis of Late Cretaceous intra-oceanic arc
 1096 intrusions in the Pertek area of Tunceli Province, eastern Turkey, and implications
 1097 for the geodynamic evolution of the southern Neo-Tethys: Results of zircon U–Pb
 1098 geochronology and geochemical and Sr–Nd isotopic analyses. *Lithos* 350-351,
 1099 105263. doi: 10.1016/j.lithos.2019.105263

1100 Saunders, A.D., Fornari, D.J., Joron, J., Tarney, J., Treuil, M., 1982. Geochemistry of
 1101 basic igneous rocks, Gulf of California, Deep Sea Drilling Project Leg 64, in:
 1102 Curray, J.R., Moore, D.G., Kelts, K. (Eds.), *Initial Reports of the Deep Sea Drilling*
 1103 *Project. Ocean Drilling Program* 64, College Station, TX, pp. 595-642.

1104 Saunders, A.D., Tarney, J., Stern, C.R., Dalziel, I.W., 1979. Geochemistry of Mesozoic
 1105 marginal basin floor igneous rocks from southern Chile. *Geological Society of*
 1106 *America Bulletin* 90, 237-258.

1107 Şengör, A.M.C., Natal'in, B.A., 1996. Turkic-type orogeny and its role in the making of

1108 the continental crust. *Annual Review of Earth and Planetary Sciences* 24, 263-
 1109 337.

1110 Şengör, A.M.C., Yılmaz, Y., 1981. Tethyan evolution of Turkey: a plate tectonic
 1111 approach. *Tectonophysics* 75, 181-241.

1112 Şengör, A.M.C., Yılmaz, Y., Sungurlu, O., 1984. Tectonics of the Mediterranean
 1113 Cimmerides: nature and evolution of the western termination of Palaeo-Tethys, in:
 1114 Dixon, J.E., Robertson, A.H.F. (Eds.), *The Geological Evolution of the Eastern*
 1115 *Mediterranean*. Geological Society of London, Special Publications 17, pp. 77-112.

1116 Shervais, J.W., 1982. Ti-V plots and the petrogenesis of modern and ophiolitic lavas.
 1117 *Earth and Planetary Science Letters* 59, 101-118.

1118 Stampfli, G.M., Borel, G., 2002. A plate tectonic model for the Paleozoic and Mesozoic
 1119 constrained by dynamic plate boundaries and restored synthetic oceanic
 1120 isochrons. *Earth and Planetary Science Letters* 196, 17-33.

1121 Sun, S.-s., McDonough, W.F., 1989. Chemical and isotopic systematics of oceanic
 1122 basalts: implications for mantle composition and processes, in: Saunders, A.D.,
 1123 Norry, M.J. (Eds.), *Magmatism in the Ocean Basins*. Geological Society of London,
 1124 Special Publications 42, pp. 313-345.

1125 Tamaki, K., Pisciotto, K., Allan, J., Alexandrovich, J.M., Barnes, D.A., Boggs, S.,
 1126 Brumsack, H., Brunner, C.A., Cramp, A., Jolivet, L., Kawka, O.E., Koizumi, I.,
 1127 Kuramoto, S.i., Langseth, M.G., McEvoy, J., Meredith, J.A., Mertz, K.A., Murray,
 1128 R.W., Nobes, D.C., Rahman, A., Schaar, R., Stewart, K.P., Tada, R., Thy, P.,
 1129 Vigliotti, L., White, L.D., Wipperfurth, J., Yamashita, S., 1990. *Proceedings of the*
 1130 *Ocean Drilling Program, Initial Reports vol. 127*. Ocean Drilling Program.

1131 Tatsumi, Y., Kogiso, T., 2003. The subduction factory: its role in the evolution of the
 1132 Earth's crust and mantle, in: Larter, R.D., Leat, P.T. (Eds.), *Intra-Oceanic*
 1133 *Subduction Systems: Tectonic and Magmatic Processes*. Geological Society of
 1134 London, Special Publications 219, pp. 55-80.

1135 Thirlwall, M., 1991. Long-term reproducibility of multicollector Sr and Nd isotope ratio

1136 analysis. *Chemical Geology* 94, 85-104.

1137 Trehu, A., Asudeh, I., Brocher, T., Luetgert, J., Mooney, W., Nabelek, J., Nakamura, Y.,
 1138 1994. Crustal architecture of the Cascadia forearc. *Science* 266, 237-243.

1139 Ural, M., Arslan, M., Göncüoğlu, M., Tekin, U., Kürüm, S., 2015. Late Cretaceous arc
 1140 and back-arc formation within the southern Neotethys: Whole-rock, trace element
 1141 and Sr-Nd-Pb isotopic data from basaltic rocks of the Yüksekova Complex
 1142 (Malatya- Elazığ, SE Turkey). *Ofioliti* 40, 52-72.

1143 Ustaömer, P.A., Ustaömer, T., Robertson, A.H.F., 2012. Ion probe U-Pb dating of the
 1144 Central Sakarya basement: a peri-Gondwana terrane intruded by late Lower
 1145 Carboniferous subduction/collision-related granitic rocks. *Turkish Journal of Earth
 1146 Sciences* 21, 905-932.

1147 Ustaömer, T., Robertson, A.H.F., 1997. Tectonic-sedimentary evolution of the North-
 1148 Tethyan active margin in the Central Pontides of Northern Turkey, in: Robinson,
 1149 A.G. (Ed.), *Regional and Petroleum Geology of the Black Sea and Surrounding
 1150 Region*. AAPG Memoirs 68, pp. 255-290.

1151 van Hinsbergen, D.J., Torsvik, T.H., Schmid, S.M., Mañenco, L.C., Maffione, M., Vissers,
 1152 R.L., Gürer, D., Spakman, W., 2020. Orogenic architecture of the Mediterranean
 1153 region and kinematic reconstruction of its tectonic evolution since the Triassic.
 1154 *Gondwana Research* 81, 79-229.

1155 Wang, Y., Fan, W., Zhang, Y., Guo, F., Zhang, H., Peng, T., 2004. Geochemical,
 1156 $^{40}\text{Ar}/^{39}\text{Ar}$ geochronological and Sr-Nd isotopic constraints on the origin of
 1157 Paleoproterozoic mafic dikes from the southern Taihang Mountains and
 1158 implications for the ca. 1800Ma event of the North China Craton. *Precambrian
 1159 Research* 135, 55-77.

1160 Weaver, S.D., Saunders, A.D., Pankhurst, R.J., Tarney, J., 1979. A geochemical study
 1161 of magmatism associated with the initial stages of back-arc spreading.
 1162 *Contributions to Mineralogy and Petrology* 68, 151-169.

1163 Weis, D., Kieffer, B., Hanano, D., Nobre Silva, I., Barling, J., Pretorius, W., Maerschalk,

1164 C., Mattielli, N., 2007. Hf isotope compositions of US Geological Survey reference
1165 materials. *Geochemistry, Geophysics, Geosystems* 8, 57-77.

1166 Weis, D., Kieffer, B., Maerschalk, C., Barling, J., De Jong, J., Williams, G.A., Hanano,
1167 D., Pretorius, W., Mattielli, N., Scoates, J.S., 2006. High-precision isotopic
1168 characterization of USGS reference materials by TIMS and MC-ICP-MS.
1169 *Geochemistry, Geophysics, Geosystems* 7, Q08006. doi:
1170 10.1029/2006GC001283

1171 Whalen, J.B., Currie, K.L., Chappell, B.W., 1987. A-type granites: geochemical
1172 characteristics, discrimination and petrogenesis. *Contributions to Mineralogy and*
1173 *Petrology* 95, 407-419.

1174 Winchester, J., Floyd, P., 1977. Geochemical discrimination of different magma series
1175 and their differentiation products using immobile elements. *Chemical Geology* 20,
1176 325-343.

1177 Yazgan, E., Chessex, R., 1991. Geology and Tectonic Evolution of the Southeastern
1178 Taurides in the Region of Malatya. *Turkish Association of Petroleum Geologists* 3,
1179 1-42.

1180 Yiğitbaş, E., Yılmaz, Y., 1996. Post-Late Cretaceous strike-slip tectonics and its
1181 implications for the Southeast Anatolian orogen, Turkey. *International Geology*
1182 *Review* 38, 818-831.

1183 Yıldırım, E., 2015. Geochemistry, petrography and tectonic significance of the ophiolitic
1184 rocks, felsic intrusions and Eocene volcanic rocks of an imbrication zone (Helete
1185 area, Southeast Turkey). *Journal of African Earth Sciences* 107, 89-107.

1186 Yılmaz, Y., 1993. New evidence and model on the evolution of the southeast Anatolian
1187 orogen. *Geological Society of America Bulletin* 105, 251-271.

1188 Yılmaz, Y., Yiğitbaş, E., Genç, Ş.C., 1993. Ophiolitic and metamorphic assemblages of
1189 southeast Anatolia and their significance in the geological evolution of the
1190 orogenic belt. *Tectonics* 12, 1280-1297.

1191 Zhang, Y., Wang, Y., Geng, H., Zhang, Y., Fan, W., Zhong, H., 2013. Early

1192 Neoproterozoic (~850Ma) back-arc basin in the Central Jiangnan Orogen
1193 (Eastern South China): Geochronological and petrogenetic constraints from meta-
1194 basalts. Precambrian Research 231, 325-342.
1195 Zindler, A., Hart, S., 1986. Chemical geodynamics. Annual Review of Earth and
1196 Planetary Sciences 14, 493-571.
1197

1198 Listing of Figures

1199 Fig. 1 Outline tectonic map of the Eastern Mediterranean region, including the main
1200 volcanic units mentioned in the text. The tectonic framework is modified after
1201 Robertson et al. (2012a). Note the location of the Kyrenia Range in northern Cyprus
1202 (red box). Jurassic, Cretaceous and Eocene igneous rocks are shown in blue, red and
1203 yellow, respectively (see the text for literature sources).

1204 Fig. 2 Successions shown in two main thrust sheets exposed in the Kyrenia Range,
1205 northern Cyprus. (a) Simplified log of the succession in the largest thrust sheet
1206 including the latest Cretaceous-Paleogene volcanics (Melounda Formation); (b)
1207 Restored log of the small, dismembered frontal thrust sheets that expose the Late
1208 Cretaceous felsic volcanics (Fourkovouno Formation) (data from Baroz, 1979;
1209 Robertson et al., 2012b and this study).

1210 Fig. 3 (a) Outline geological map of the Kyrenia Range (modified from Robertson et al.,
1211 2012b). Locations from which basalts were collected are numbered on the map, with
1212 small yellow boxes. Karpas Peninsula: 1-Balalan (Platanissos); Eastern Range: 2-
1213 Ağıllar (Mandres), 3-Çınarlı (Platani), 4-Mallıdağ (Melounda); Central Range: 5-Tirmen
1214 (Trypimeni), 6-Ergenekon (Agios Khariton), 7-Değirmenlik (Kythrea), 8-Arapköy
1215 (Klepini), 9-Beylerbeyi (Bellapais), 10-Boğaz (Bogaz), 11-Pınarbaşı (Krini), 12-İncesu
1216 (Motides), 13-Alevkaya Tepe (Kiparisso Vouno); Western Range, 14-Karşıyaka
1217 (Vasileia), 15-Geçitköy (Panagra); (b) Geological map of the western Kyrenia Range,
1218 northern Cyprus made during this study (based on mapping by Baroz, 1979). Locations
1219 of cross sections (AA', BB' and CC') are indicated. Note that the two contrasting, mainly
1220 felsic and mainly basaltic volcanic units are separated by a thrust fault (Robertson et
1221 al., 2012b; this study; see Fig. 2).

1222 Fig. 4 (a) A-A' cross section of Geçitköy that is dominated by a south-verging,
1223 recumbent anticline (Late Miocene-earliest Pliocene); (b) B-B' cross-section of
1224 Selvilitepe showing southward imbrication of basalt/pelagic chalk and felsic

1225 volcanogenic rocks; (c) C-C' cross-section showing imbricated felsic volcanogenic
1226 rocks.

1227 Fig. 5 Measured stratigraphic logs of partial successions in the felsic volcanics and
1228 structurally associated units. The structural lower unit (logs a-e) is relatively intact,
1229 whereas the higher unit (logs f-j) is characterised by the thrust intercalations with the
1230 Trypa Group. Fig. 6 (a) Zr/Ti vs. Nb/Y diagram (after Pearce, 1996) and (b) Th vs. Co
1231 diagram (after Hastie et al., 2007) for the Kyrenia Range felsic rocks; (c) Primitive
1232 mantle-normalised multi-element spider diagram and (d) chondrite-normalised rare
1233 earth element (REE) patterns of the felsic volcanics with reference data for upper
1234 continental crust (UCC) and lower continental crust (LCC). Normalising values of
1235 primitive mantle and chondrite after Sun and McDonough (1989), upper continental
1236 crust (UCC) and lower continental crust (LCC) data are from Rudnick and Gao (2003),
1237 and Hu and Gao (2008). The Chile volcanic arc granite data are after Pearce et al.
1238 (1984).

1239 Fig. 7 (a) Zr/Ti vs. Nb/Y diagram (after Pearce, 1996) and (b) Th vs. Co diagram (after
1240 Hastie et al., 2007) for the basaltic volcanics. Red symbols: Karpas Peninsula and
1241 eastern range; green symbols: central range; blue symbols: western range. Paleogene
1242 basalt samples are indicated by thick magenta-outlined symbols.

1243 Fig. 8 Mid-ocean ridge basalt (MORB)-normalised multi-element spider diagram for
1244 basalts from the Karpas Peninsula and the eastern (a), central (b) and western (c)
1245 Kyrenia Range. MORB-normalised data are from Pearce et al. (1983); (d) Chondrite-
1246 normalised REE patterns of selected basalts. Chondrite, OIB and E-MORB data are
1247 from Sun and McDonough (1989). Paleogene basalt samples are indicated by solid
1248 black lines and magenta-outlined symbols.

1249 Fig. 9 (a) Cathodoluminescence images of zircon grains separated from the samples
1250 of Geçitköy. Red circles mark inconsistent analyses that were omitted from the

1251 weighted mean $^{206}\text{Pb}/^{238}\text{U}$ age calculation. Locations of the ion probe analysis spots
1252 and the corresponding ages ($^{206}\text{Pb}/^{238}\text{U} \pm 1\sigma$) are indicated. Scale bar = 20 μm ; (b-d)
1253 Wetherill Concordia diagram for the zircons analyses from sample 14-18 (b), 14-19 (c)
1254 and 14-20 (d).

1255 Fig. 10 (a-b) Plots of P_2O_5 vs. SiO_2 and Th, respectively, show that all of the samples
1256 follow the I-type granite trend; (c) Zr vs. $10000 \times \text{Ga}/\text{Al}$, and (d) Nb vs. $10000 \times \text{Ga}/\text{Al}$
1257 discrimination diagrams of Whalen et al. (1987), showing the I-, S (sedimentary)- and
1258 M (depleted mantle source)-type nature of the Late Cretaceous felsic volcanics.

1259 Fig. 11 (a) $\epsilon\text{Nd}(\text{t})$ vs. $\epsilon\text{Sr}(\text{t})$ diagram for the basaltic rocks analysed. The samples are
1260 mainly Late Cretaceous ($n=4$); Paleogene basalt samples ($n=2$) are indicated by
1261 magenta-outlined symbols. Comparative compositions are from Zindler and Hart
1262 (1986); (b) $\epsilon\text{Hf}(\text{t})$ vs. $\epsilon\text{Nd}(\text{t})$ plot showing rocks for Kyrenia Range, northern Cyprus
1263 relative to other volcanic rocks of the western Mediterranean (after Kempton et al.,
1264 2018).

1265 Fig. 12 (a) Ba/Nb versus La/Nb diagram (after Zhang et al., 2013); (b) Nb/Y versus
1266 Zr/Y diagram (after Condie, 2005), arrows in (b) indicate the effects of batch melting
1267 (F) and the probable effect of fluids derived from subduction (SUB); (c) Zr/Nb vs. Nb/Th
1268 diagram (modified after Condie, 2003). Abbreviations: PM, primitive mantle; DM,
1269 shallow depleted mantle; HIMU, high μ (U/Pb) source; EMI and EMII, enriched
1270 mantle sources; ARC, arc-related basalt; N-MORB, normal ocean ridge basalt; OIB,
1271 oceanic island basalt; DEP, deep depleted mantle; EN, enriched component; REC,
1272 recycled component; UCC, upper upper continental crust; LCC, lower continental crust;
1273 BM, batch melting trajectory with percent melting noted. Numbers on mixing lines are
1274 percents.

1275 Fig. 13 (a) Nb vs. Y diagram (Pearce et al., 1984) and (b) Th/Yb vs. Ta/Yb (Gorton and
1276 Schandl, 2000) for the felsic volcanics; (c) Zr/Y vs. Zr diagram (after Pearce and Norry,
1277 1979) and (d) V vs. Ti discrimination diagram (after Shervais, 1982) for the basaltic

1278 rocks. Abbreviations: WPG, within-plate granite; VAG, volcanic arc granite; syn-COLG,
1279 syn-collisional granite; ORG, ocean ridge granite; WPB, within-plate basalt; MORB,
1280 mid-ocean ridge basalt; BABB, back-arc basin basalt; ARC, arc-related basalt; OFB,
1281 oceanic floor basalt.

1282 Fig. 14 (a) Th/Nb vs. La/Nb and (b) La/Nb versus Y diagrams (after Floyd et al., 1991).

1283 Abbreviations: IAT, island arc tholeiite; BABB, back-arc basin basalt; OFB, oceanic
1284 flood basalt; FAPB, fore-arc platform basalt; T/E-MORB, T-type/enriched-mid-ocean
1285 ridge basalt.

1286 Fig. 15 Alternative tectonic models of Tethys in the Eastern Mediterranean region. (a-
1287 c) Northward subduction with continental fragments rifted from Gondwana (Robertson
1288 et al., 2012a); (d-f) Northward subduction and marginal basin formation (Barrier et al.,
1289 2018); (g-h) Genesis of Late Cretaceous supra-subduction ophiolites at a single
1290 subduction zone to the NE followed by roll-back of segments including the Southern
1291 Neotethys (Maffione et al., 2017; McPhee and van Hinsbergen, 2019; van Hinsbergen
1292 et al., 2020). Based on discussion (see text), model 1 is generally favoured.

1293

1294 Listing of Supplementary Figure

1295 Supplementary Figure S1 Field photographs of the felsic volcanics. (a) Exposure in
1296 Geçitköy. Samples 14-18 and 14-19 are from this location; (b) Irregular-shaped
1297 doleritic intrusion (hammer for scale); (c) Localised occurrence of greyish to greenish,
1298 rhyolitic debris-flow unit, north of Geçitköy; (d) Greenish rhyolitic debris-flow unit with
1299 angular rhyolitic clasts (pen for scale).

1300 Supplementary Figure S2 Field photographs of the basaltic volcanics. (a) Pillow lavas
1301 and interstitial pink pelagic carbonate; roadcut c. 200 m north of Balalan; (b) basalt
1302 interbedded with pelagic carbonate; thrust sheet above Miocene siliciclastic sediment;
1303 c. 800 m northwest of Değirmenlik; (c) Basaltic volcanics intercalated with pelagic
1304 carbonates and small slices of meta-platform carbonates of the Trypa Group along the
1305 southern flank of the range, 800 m north of Boğaz; (d) Basalt-pelagic carbonate
1306 intercalation, south of Karşıyaka.

1307 Supplementary Figure S3 Photomicrographs of felsic volcanics. (a) Subhedral
1308 sanidine (Sa), plagioclase (Pl) and biotite (Bt), in a groundmass of devitrified glassy
1309 volcanic shards (cross-polarised light); (b) felsic glass groundmass (Vg) with
1310 microcrystalline quartz, irregular-shaped quartz (Q) and feldspar (Pl) (cross-polarised
1311 light). Sample number is indicated in the bottom-left corner.

1312 Supplementary Figure S4 Photomicrographs of basalts. (a) Intersertal basalt with
1313 euhedral granular augite (Px), elongate plagioclase laths (Pl) and opaque grains (Op)
1314 (cross-polarised light); (b) Basalt with intersertal texture, in which randomly-oriented
1315 plagioclase generally enclose pyroxenes (plane-polarised light); (c) Porphyritic basalt
1316 with euhedral plagioclase and strongly altered olivine phenocrysts (cross-polarised
1317 light); (d) Ophitic basalt with plagioclase phenocrysts, augite occurs as an inclusion
1318 within individual plagioclase crystals (cross-polarised light). Sample number is
1319 indicated in the bottom-left corner. Samples no. 21, 14-51 and 14-69 are from the
1320 Melounda Formation, whereas sample no. 19-51 is from the Ayios Nikolaos Formation.

Supplementary Figure S5 Tectonic discrimination plots. (a) TiO_2 - MnO - Na_2O ternary diagram (after Nisbet and Pearce, 1977) for clinopyroxenes in basalt; (b) TiO_2 - SiO_2 - Na_2O diagram (after Beccaluva et al., 1989) for clinopyroxenes in basalt. Abbreviations: VAB, volcanic arc basalt; OFB, ocean-floor basalt; WPA, within-plate alkali basalt; WPT, within-plate tholeiitic basalt; BON, boninite; IAT, island-arc tholeiite; WPB, within-plate basalt.

Supplementary Figure S6 (a) Chondrite-normalised REE comparison patterns of the Kyrenia Range felsic rocks and the Helete granites in SE Turkey (data after Nurlu et al., 2016). Chondrite data after Sun and McDonough (1989); (b)-(c) MORB-normalised multi-element spider diagrams for basalts. For comparison, the coloured fields show the composition variations of basalts from the Kyrenia Range (this study), the Misis Complex, S Turkey (Floyd et al., 1991), the Maden Complex, S Turkey (Erturk et al., 2018), the Japan Sea (Chen et al., 2015), the Sarmiento Complex, Chile (Saunders et al., 1979), the Bransfield Strait, Antarctic (Keller et al., 2002), the Guaymas Basin, Caribbean (Saunders et al., 1982) and the Tyrrhenian Sea, Italy (Peccerillo, 2017). MORB-normalised data are from Pearce et al. (1983). N-MORB, E-MORB and OIB data are from Sun and McDonough (1989). Specifically, two representative basalts of different affinities (within-plate vs. volcanic arc) from the Kyrenia Range are indicated.

1348 Listing of Supplementary Tables

1349 Supplementary Table S1. Major element oxides, trace and rare earth elements for the
1350 Late Cretaceous Fourkovouno (Selvilitepe) Formation.

1351 Supplementary Table S2. Major element oxides, trace and rare earth element analyses
1352 for the latest Cretaceous-Paleogene basaltic volcanics.

1353 Supplementary Table S3. Electron microprobe analyses of feldspar in the latest
1354 Cretaceous-Paleogene basaltic volcanics.

1355 Supplementary Table S4. Electron microprobe analyses of pyroxene in the latest
1356 Cretaceous-Paleogene basaltic volcanics.

1357 Supplementary Table S5. SIMS zircon U-Pb analyses of the zircon grains separated
1358 from the Late Cretaceous Fourkovouno (Selvilitepe) Formation.

**Evidence from Late Cretaceous-Paleogene volcanic rocks of the Kyrenia Range,
northern Cyprus for the northern, active continental margin of the Southern
Neotethys**

Guohui Chen^{1*}, Alastair H. F. Robertson¹

¹School of GeoSciences, University of Edinburgh, West Mains Road, Edinburgh EH9
3JW, UK

*Current Address: State Key Laboratory of Lithospheric Evolution, Institute of Geology
and Geophysics, Chinese Academy of Sciences, P.O. Box 9825, Beijing 100029,
China

Corresponding author: Guohui Chen (Guohui.Chen@live.cn)

Abstract

Late Cretaceous felsic and latest Cretaceous-Paleogene basaltic volcanic rocks are
exposed throughout the Kyrenia Range, N Cyprus. Field mapping of the key, well-
exposed western range indicates that the felsic volcanics are mainly crop outexposed
in the southerly, frontal part of the range, separated from the latest Cretaceous-
Paleogene basaltic volcanics farther north by a thrust. New U-Pb zircon dating of the
felsic volcanics indicates a primary age of c. 74.0 Ma (Late Campanian). The felsic
volcanics are characterised by evolved high potassium and shoshonitic compositions
that were probably generated in an extensional subduction-related setting. Their
relatively low Nb and Y but high Rb concentrations, together with characteristic Th/Ta
ratios (6-20), suggest a mature continental arc setting. The latest Cretaceous-
Paleogene basaltic volcanics have mainly within-plate chemical characteristics (in the
east), although some have a subordinate subduction influence; e.g. negative Nb (in

Formatted: Font color: Auto

Formatted: Font color: Auto

the west). Sr-Nd-Hf isotopic signatures (i.e. high positive $\epsilon\text{Nd}(t)$ and $\epsilon\text{Hf}(t)$ values) suggest derivation of the basalts from several OIB-like mantle sources, with probable involvement of a crustal (recycled) component (i.e. elevated Nb/Y ratios). Comparisons with SE Turkey, where Late Cretaceous arc-related granitic rocks are widely developed, suggest that the Kyrenia Range igneous rocks may have originated to the south of the Alanya metamorphic massif (N of Cyprus) and correlative continental units in SE Turkey. ~~The Kyrenia Range Late Cretaceous felsic volcanics are comparable with rare Late Cretaceous granitic rocks in the front of the over-riding Tauride thrust belt in SE Turkey (Hölele unit). They are also comparable with Late Cretaceous arc-related magmatic rocks higher in the regional tectono-stratigraphy in SE Turkey, although these may relate to a different continental margin arc. We propose that the Late Cretaceous felsic volcanics in N Cyprus record mature arc magmatism, as in SE Turkey. In contrast,~~ the latest Cretaceous-Paleogene basaltic volcanics in N Cyprus and SE Turkey (Maden Complex) are interpreted to represent incipient marginal basin formation, possibly in an oblique-convergent setting, prior to Miocene suturing with Arabia. In the light of alternatives, we infer genesis of the N Cyprus Late Cretaceous and Paleogene volcanic rocks related to stages in the development of the northerly active continental margin of the Southern Neotethys.

Keywords: Late Cretaceous-Paleogene; Volcanic rocks; Kyrenia Range; Arc magmatism; Marginal basin formation; Southern Neotethys

1. Introduction

Continental margin magmatism is a key feature of subduction at convergent plate margins (Dewey, 1969; Şengör and Natal'in, 1996). Information concerning the field relations, age, petrology and geochemistry are essential to understand the growth and demise of arcs, and are critical to reconstruct paleogeography and to test alternative tectonic hypotheses (e.g. Trehu et al., 1994; Tatsumi and Kogiso, 2003; Bowman et al., 2019).

The Eastern Mediterranean region, situated between the North African-Arabian and Anatolian (Eurasian) plates, includes tectonically emplaced remnants of Neotethys that developed from the preceding ~~larger~~ Paleotethys farther north (e.g. Le Pichon, 1982; Şengör et al., 1984; Robertson and Dixon, 1984; Stampfli and Borel, 2002). Much of the arc magmatism in the north, in the Pontides, of Jurassic-Eocene age (Fig. 1), relates to long-lived northward subduction of the Northern Neotethys (İzmir-Ankara-Erzincan ocean) (Adamia et al., 1977; Dercourt et al., 1993; Ustaömer and Robertson, 1997; Rice et al., 2006). Subduction of the Southern Neotethys ~~also took place during the Late Cretaceous (95-90 Ma), resulting in~~ both continental margin and oceanic magmatism that ranges, overall, from Late Cretaceous to Paleogene (Fig. 1). Here, we focus on evidence of Late Cretaceous and Paleogene volcanism in the Kyrenia Range of northern Cyprus, representing the most westerly known occurrence of arc-type magmatism that can be related to subduction of the Southern Neotethys (Fig. 1) (Moore, 1960; Ducloz, 1972; Pearce, 1975; Baroz, 1979, 1980; Robertson and Woodcock, 1986; Huang et al., 2007; Robertson et al., 2012b).

Our specific objectives here are: (1) to understand the tectono-stratigraphy of two, contrasting, felsic and basaltic volcanic units in relation to associated sedimentary units; (2) to determine ~~directly~~ the eruptive ages of both of the volcanic units directly, using geochronology. Previously, the volcanics ~~These were previously~~ dated indirectly using microfossils within interbedded pelagic carbonates (Baroz, 1979; Robertson et

al., 2012b); (3) to ~~investigate~~determine whether there is any petrological or geochemical variation (including isotopic variation) along the Kyrenia Range (c. 100 km) and, if so, the implications; (4) to infer the magmatic-tectonic setting of eruption, compared with the evidence of arc volcanism ~~globally, and~~ especially in southeast Turkey; (5) to test alternative tectono-magmatic hypotheses bearing in mind that arc magmatism ~~plays~~has a key role in plate tectonic interpretations.

~~2. Methods~~

~~The Kyrenia Range is a complex thrust belt, and therefore it was essential initially, to remap the western Kyrenia Range, where felsic and basaltic volcanic rocks are exposed together (Moore, 1960; Baroz, 1979; Robertson et al., 2012b). In addition, sedimentary logs were measured and correlated to produce a composite stratigraphy of the volcanogenic successions that are exposed in two superimposed thrust sheets.~~

~~Optical microscopy was carried out on the samples collected. For the felsic lithologies, 13 samples were studied from the western range. For the basaltic lithologies, we used a combination of new samples (n=29) and also samples that were previously collected for paleomagnetic study (n=26) (Morris et al., 2015) throughout the Kyrenia Range.~~

~~Whole-rock major and trace element concentrations of the volcanic rocks were measured by X-ray fluorescence (XRF) on fused glass beads and pressed powder pellets at the School of GeoSciences, University of Edinburgh, using the well-known methods of Fitton et al. (1998) and Fitton and Godard (2004). Accuracy and precision are typically c. 5%. For representative samples, additional trace and rare earth element (REE) analysis was carried out by inductively coupled plasma-mass spectrometry (ICP-MS) at the ACME Analytical Laboratories, Vancouver. Major element contents were determined from a LiBO₂ fusion by ICP-ES by using 5 g of sample pulp. Trace~~

element contents were determined from a LiBO₂ fusion by ICP-MS by using 5 g of sample pulp. Detection limits range between 0.01 and 0.04 wt% for major oxides, 0.01 and 0.1 ppm for trace and rare earth elements. The relative standard deviation for the REE is 5% and for all other trace elements is up to 10%, with quality control using international geostandards (see <http://aemelab.com>).

In addition, Sr-Nd-Hf isotopic analysis was performed on a Neptune Plus multi-collector (MC)-ICP-MS at Wuhan SampleSolution Analytical Technology Co., Ltd., China, as reported by Li et al. (2012). Whole procedural blanks were <100 pg for Sr, <50 pg for Nd and <50 pg for Hf. ⁸⁷Sr/⁸⁶Sr, ¹⁴³Nd/¹⁴⁴Nd and ¹⁷⁶Hf/¹⁷⁷Hf ratios were normalised to ⁸⁶Sr/⁸⁸Sr = 0.1194, ¹⁴⁶Nd/¹⁴⁴Nd = 0.7219 and ¹⁷⁹Hf/¹⁷⁷Hf = 0.7325, using the exponential law. Standard analysis yielded ⁸⁷Sr/⁸⁶Sr = 0.710240 ± 11 (2SD, n = 4) for NBS987, ¹⁴³Nd/¹⁴⁴Nd = 0.512440 ± 8 (2SD, n = 8) for GSB-04-3258-2015, and ¹⁷⁶Hf/¹⁷⁷Hf = 0.282224 ± 6 (2SD, n = 6) for Alfa-Hf. In addition, USGS reference materials BCR-2 and RGM-2 were also analysed for Sr-Nd-Hf isotopes, and gave ratios of 0.705039 ± 8 and 0.704169 ± 11 for ⁸⁷Sr/⁸⁶Sr, 0.512644 ± 6 and 0.512808 ± 9 for ¹⁴³Nd/¹⁴⁴Nd and 0.282864 ± 6 and 0.283016 ± 7 for ¹⁷⁶Hf/¹⁷⁷Hf, which is within error of recommended values (Thirlwall, 1991; Weis et al., 2006, 2007). The analytical data for the major, trace and rare earth elements/isotope ratio of the felsic and basaltic volcanics are listed in the Supplementary Table S1, S2, respectively.

Furthermore, two polished thin sections of basalt (no. 02 and 21) that contain relatively fresh feldspar and pyroxene crystals were selected for analysis of major elements using a Cameca SX100 electron microprobe at the School of GeoSciences, University of Edinburgh. Details of the analytical conditions and methods are given by Hayward (2011) and by Hartley and Thordarson (2013). The accuracy of major element determinations is <± 1% of total value. Analytical data for feldspar and pyroxene are given in Supplementary Table S3, S4, respectively.

~~Zircon crystals within 30–160 µm fraction were separated from crushed felsic tuffaceous rock and rhyolitic lava (no. 14-18, 14-19 and 14-20; c. 5 kg each). A Wilfley table, Frantz Isodynamic magnetic separator and a high density solution (lithium polytungstate; 2.85 g/ml) were used to aid separation. Zircon grains were randomly handpicked under a binocular microscope, mounted in epoxy resin and polished sufficiently to expose the center of the grains. Internal structures were studied with a scanning electron microscope using cathodoluminescence (CL) at the School of GeoSciences, University of Edinburgh. U-Pb analysis was then performed using a Cameca IMS-1270 secondary ion mass spectrometer (SIMS) at the School of GeoSciences, University of Edinburgh, using the methods reported by Kelly et al. (2008) and Ustaömer et al. (2012). Errors of the reported ages are $\pm 1\sigma$. Related geochronological plots were produced using ISOPLOT (Ludwig, 2012). Analytical results are listed in the Supplementary Table S5.~~

~~32.~~ Volcanism in the Kyrenia Range

Rifting of the Southern Neotethys during Late Permian and Early-Middle Triassic was followed by continental break-up during the Late Triassic-Early Jurassic (Robertson and Woodcock, 1979; Şengör and Yılmaz, 1981; Robertson and Dixon, 1984; Garfunkel, 1998; Robertson et al., 2020). The Kyrenia Range then formed part of a carbonate platform that slowly subsided during Jurassic ~~and~~ Early Cretaceous in a along a passive margin ~~of the Southern Neotethys setting~~ (Robertson and Woodcock, 1986). The carbonate platform was deformed and metamorphosed under greenschist facies during the Late Cretaceous, extensionally exhumed, and then unconformably overlain by basic extrusive igneous rocks of latest Cretaceous ~~and~~ Paleogene age (Figs. 2-3) (Ducloz, 1972; Baroz, 1979; Robertson and Woodcock, 1986). In addition, felsic volcanics and tuffs are exposed as thrust slices at a low structural level in the south of the range (Figs. 2b, 3) (Robertson et al., 2012b). These volcanic rocks have

no exposed base and are directly overlain by ~~the a~~ much larger thrust sheet that includes the mainly basic extrusive igneous rocks.

Baroz (1979, 1980) mapped the volcanic rocks and carried out petrographic and whole-rock chemical analysis of major elements. He reported the presence of a lower stratigraphic sequence of basalt, dolerite, trachybasalt, trachyandesite, dacite and rhyolitic tuff, and interpreted this as a bimodal basic-acidic calc-alkaline suite, related to a Late Maastrichtian volcanic arc. In contrast, the Paleogene volcanic assemblage was suggested to have erupted stratigraphically above in a post-collisional (intra-continental) strike-slip setting (Baroz, 1980). However, a thrust was later mapped between the mainly felsic and mainly basic suites (Robertson et al., 2012b), complicating this simple stratigraphy.

Chemical analysis of the Paleogene basaltic lavas, including immobile trace elements, initially suggested an alkaline, within-plate eruptive setting (Pearce, 1975). Subsequent chemical analysis, including some immobile elements, confirmed ~~this a~~ mainly alkaline within-plate setting but also revealed some evidence of a subduction influence (e.g. negative Nb anomaly) (Robertson and Woodcock, 1986). The Paleogene lavas were ~~at that time~~ inferred by these authors to have erupted in a transtensional setting along the northern, active margin of the 'Troodos ocean'. Huang et al. (2007) carried out additional chemical analysis of the Paleogene basaltic lavas, mainly from the eastern and western Kyrenia Range, with emphasis on immobile elements, and proposed a Late Cretaceous-Paleogene back-arc setting related to northward subduction.

The felsic rocks of the lower most thrust sheet, known as the Fourkovouno (Selvilitepe)¹ Formation, ~~(up to 400 m thick),~~ begin with marine water-lain felsic tuffs

¹ For simplicity, ~~w~~We use the traditional stratigraphy for the formation names (more recent Turkish equivalents are mentioned initially). However, we use current the ~~the~~ Turkish names

and subaqueous felsic debris-flow deposits and culminate in laterally discontinuous, thick-bedded to massive rhyolitic lava flows (Fig. 3) (Moore, 1960; Baroz, 1979; Huang et al., 2007; Robertson et al., 2012b). The felsic rocks are locally cut by small (meter-sized) basaltic/doleritic intrusions (~~that were~~ not studied). A Late Cretaceous age (Late Campanian) was ~~previously~~ suggested for the felsic lavas, based on sparse planktic foraminifera (*Globotruncana* sp.) that were recognised in a chalky interbed near the top of the succession (Baroz, 1979).

The Late Cretaceous and Paleogene basaltic igneous rocks have been divided into ~~three-two~~ formations, namely the Maastrichtian Melounda (Mallıdağ) Formation ~~and~~ the Paleocene-Middle Eocene Ayios Nikolaos (Yamaçköy) Formation (Fig. 2a).

The Melounda Formation, c. 300 m thick, unconformably overlies the exhumed Mesozoic platform carbonates of the Trypa (Tripa) Group or, in places, unmetamorphosed terrigenous turbidites of the Kiparisso Vouno (Alevkaya Tepe) Member of the Melounda Formation (Baroz, 1979; Robertson and Woodcock, 1986; Robertson et al., 2012b). The basaltic lavas of the Melounda Formation are commonly intercalated with pinkish pelagic limestones (Fig. 2a). The succession locally includes lenticular, lithoclastic breccias and finer-grained clastic facies that were reworked from the underlying meta-platform carbonates (Fig. 2a) (Robertson and Woodcock, 1986; Robertson et al., 2012b). Planktic foraminifera from ~~the~~ pelagic carbonates between the lavas indicate a Late Maastrichtian age (Baroz, 1979; Robertson et al., 2012b).

The overlying Ayios Nikolaos Formation, 300-400 m thick, comprises basaltic volcanics that are variably interbedded with pelagic carbonates. Calciturbidites, calcareous debris-flow deposits and carbonate-rock breccias appear above the lavas, towards the top of the overall succession (Baroz, 1979; Robertson and Woodcock,

(e.g. ~~for~~that are currently in use (e.g. ~~for~~ settlements); pre-existing names are mentioned only initially).

1986; Robertson et al., 2012b, 2014) (Fig. 2a). Planktic foraminifera from pelagic carbonates that are interbedded with the lavas indicate a ~~maximum~~ Late Paleocene to Mid-Eocene age range (Baroz, 1979; Robertson et al., 2012b).

3. Methods

The Kyrenia Range is a complex thrust belt and therefore it was essential to remap the western Kyrenia Range, where felsic and basaltic volcanic rocks are exposed together (Moore, 1960; Baroz, 1979; Robertson et al., 2012b). In addition, sedimentary logs were measured and correlated to produce a composite stratigraphy of the volcanogenic successions that are exposed in two superimposed thrust sheets.

Optical microscopy was carried out on the samples collected. For the felsic lithologies, 13 samples were studied from the western range. For the basaltic lithologies, we used a combination of our samples (n=29) and also some that were previously collected for paleomagnetic study (n=26) throughout the Kyrenia Range (Morris et al., 2015).

Whole-rock major and trace element concentrations of the volcanic rocks were measured by X-ray fluorescence (XRF) on fused glass beads and pressed powder pellets at the School of GeoSciences, University of Edinburgh, using the well-known methods of Fitton et al. (1998) and Fitton and Godard (2004). Accuracy and precision are typically c. 5%. For representative samples, additional trace and rare earth element (REE) analysis was carried out by inductively coupled plasma-mass spectrometry (ICP-MS) at the ACME Analytical Laboratories, Vancouver. Major element contents were determined from a LiBO₂ fusion by ICP-ES, using 5 g of sample pulp. Trace element contents were determined from a LiBO₂ fusion by ICP-MS, again using 5 g of sample pulp. Detection limits range between 0.01 and 0.04 wt% for major oxides, 0.01 and 0.1 ppm for trace and rare earth elements. The relative standard deviation for the

Formatted: Indent: First line: 2 ch

REE is ~5% and for all other trace elements is up to 10%, with quality control using international geostandards (see <http://acmelab.com>).

In addition, Sr-Nd-Hf isotopic analysis was performed on a Neptune Plus multi-collector (MC)-ICP-MS at Wuhan SampleSolution Analytical Technology Co., Ltd., China, as reported by Li et al. (2012). Whole procedural blanks were <100 pg for Sr, <50 pg for Nd and <50 pg for Hf. $^{87}\text{Sr}/^{86}\text{Sr}$, $^{143}\text{Nd}/^{144}\text{Nd}$ and $^{176}\text{Hf}/^{177}\text{Hf}$ ratios were normalised to $^{86}\text{Sr}/^{88}\text{Sr} = 0.1194$, $^{146}\text{Nd}/^{144}\text{Nd} = 0.7219$ and $^{179}\text{Hf}/^{177}\text{Hf} = 0.7325$, using the exponential law. Standard analysis yielded $^{87}\text{Sr}/^{86}\text{Sr} = 0.710240 \pm 11$ (2SD, $n = 4$) for NBS987, $^{143}\text{Nd}/^{144}\text{Nd} = 0.512440 \pm 8$ (2SD, $n = 8$) for GSB 04-3258-2015, and $^{176}\text{Hf}/^{177}\text{Hf} = 0.282224 \pm 6$ (2SD, $n = 6$) for Alfa Hf. In addition, USGS reference materials BCR-2 and RGM-2 were also analysed for Sr–Nd–Hf isotopes, and gave ratios of 0.705039 ± 8 and 0.704169 ± 11 for $^{87}\text{Sr}/^{86}\text{Sr}$, 0.512644 ± 6 and 0.512808 ± 9 for $^{143}\text{Nd}/^{144}\text{Nd}$ and 0.282864 ± 6 and 0.283016 ± 7 for $^{176}\text{Hf}/^{177}\text{Hf}$, which is within error of recommended values (Thirlwall, 1991; Weis et al., 2006, 2007). The analytical data for the major, trace and rare earth elements/isotope ratio of the felsic and basaltic volcanics are listed in the Supplementary Table S1, S2, respectively.

In addition, two polished thin sections of basalt (no. 02 and 21) that contain relatively fresh feldspar and pyroxene crystals were selected for analysis of major elements using a Cameca SX100 electron microprobe at the School of GeoSciences, University of Edinburgh. Details of the analytical conditions and methods are given by Hayward (2011) and by Hartley and Thordarson (2013). The accuracy of major element determinations is $<\pm 1\%$ of total value. Analytical data for feldspar and pyroxene are given in Supplementary Table S3, S4, respectively.

For dating, zircon crystals within 30–160 μm fraction were separated from crushed felsic tuffaceous rock and rhyolitic lava (no. 14-18, 14-19 and 14-20; c. 5 kg each). A Wilfley table, Frantz Isodynamic magnetic separator and a high-density solution

(lithium polytungstate; 2.85 g/ml) were used to aid separation. The zircon grains were randomly handpicked under a binocular microscope, mounted in epoxy resin and polished sufficiently to expose the centre of the grains. Internal structures were studied with a scanning electron microscope using cathodoluminescence (CL) at the School of GeoSciences, University of Edinburgh. U-Pb analysis was then performed using a Cameca IMS-1270 secondary ion mass spectrometer (SIMS) at the School of GeoSciences, University of Edinburgh, using the methods reported by Kelly et al. (2008) and Ustaömer et al. (2012). Errors of the reported ages are $\pm 1\sigma$. Related geochronological plots were produced using ISOPLOT (Ludwig, 2012). Analytical results are listed in the Supplementary Table S5.

4. Results

4.1. ~~Restored~~ Volcanic successions

The felsic volcanics, mainly crop out in the western part of the Kyrenia Range in two thrust sheets (Figs. 3-4). The lower of these, which is relatively intact, extends from Geçitköy (Panagra) eastwards to Selvitepe (Fourkovouno) (Fig. 5a-e). The second, structurally higher unit, comprises discontinuous exposures extending from north of Geçitköy, to southwest of Karsıyaka (Vasileia) and west of Alevkaya Tepe (Kiparisso Vouno) (Fig. 5f-j). Farther west, in the Kayalar (Orga) area (Fig. 3a), a small thrust slice exposes ~~of~~ coarse-grained felsic tuff and a felsic debris-flow ~~deposits-depositare~~ exposed. Small (tens of m-sized) outcrops of felsic volcanogenic rocks also occur farther east; e.g. northeast of Aşağı Dikmen (Kato Dikomon), between thrust sheets of ~~the~~ Mesozoic meta-carbonate rocks (Baroz, 1979). Small exposures are also intersliced with Late Cretaceous pelagic carbonates of the Melounda Formation to the north of Ergenekon (Agios Khariton) (Fig. 3a) (Robertson et al., 2012b).

The ~~restored local~~ successions are restored as a in the lower thrust sheet and an
upper thrust sheet. The succession in the lower thrust sheet, c. 90 m thick, begins with
weakly-bedded, white felsic tuff (beds <0.5 m thick) (Fig. 5a, c-e), followed by thick
layers of matrix-supported felsic breccia-conglomerate (c. 10 m thick) that ~~weare~~
~~interpreted~~ as debris-flow deposits (Fig. 5a, c, e). The succession passes upwards into
massive, vitreous, rhyolitic lava flows, individually up to 10 m thick (Figs. 5a,
Supplementary Figure S1). Localised normal grading; e.g. northeast of Geçitköy,
indicates that the succession is ~~stratigraphically~~ the right way up stratigraphically (Fig.
5a-b). In places, steeply dipping, irregularly shaped basaltic or doleritic intrusions, up
to 2 m ~~thick across~~, cut the felsic layering (Figs. 5a, Supplementary Figure S1). ~~In~~
~~contact with~~ Adjacent to these intrusions, poorly consolidated tuffaceous sediments
have undergone contact metamorphism to form hard, dark grey, flint-like, recrystallised
felsic rock (Robertson et al., 2012b). There is no evidence of similar intrusions in the
structurally overlying basalts, which is in keeping with the mapped tectonic contact
between the two units.

The succession in the upper thrust sheet begins with greyish to greenish,
tuffaceous matrix-supported conglomerates, including clasts of silicified rhyolite, chalk
and basalt (Supplementary Figure S1). The estimated thickness ranges from c. 40 m
north of Geçitköy (Fig. 5f), to c. 80 m southwest of Karsiyaka (Fig. 5g-h), to 30 m west
of the Alevkaya Tepe (Fig. 5i-j).

Contrasting field relations facilitate mapping of the Late Cretaceous versus the
Paleogene formations (Baroz, 1979; Robertson and Woodcock, 1986; Hakyemez et
al., 2000; Robertson et al., 2012b). ~~The Late Maastrichtian basalts (Melounda~~
~~Formation) are typically pillowed, forming repeated flows, up to c. 10 m thick. They~~
~~include pink pelagic carbonate as interstitial sediment and discontinuous layers (up to~~
~~40s cm thick), extending laterally for up to c. 10 m. In contrast, the Paleogene basalts~~

~~(Ayios Nikolaos Formation) are commonly massive, especially in the eastern range, where individual lava flows reach 10s of m thick, with relatively few pelagic intercalations (Supplementary Figure S2). However, in the central and western ranges, the lava flows are mainly thinner (<10m) and are intercalated with pelagic carbonates that contain abundant reddish chert (formed by diagenetic replacement). The contrasting features facilitate mapping of the two formations (Baroz, 1979; Robertson and Woodcock, 1986; Hakyemez et al., 2000; Robertson et al., 2012b).~~

~~During this work, it was found that there are geochemical differences between lavas of similar age along the far eastern, eastern, central and western Kyrenia Range (see below). We, therefore, give some more specific information on the lavas in these areas.~~

In the far east of the Kyrenia Range, the Karpas Peninsula (Fig. 3a), both pillowed and massive basalt are widely exposed near Balalan (Platanissos). ~~The lavas include interpillow carbonate and lenticular interbeds of pelagic carbonate (< 1 m thick) (Figs. 3a, Supplementary Figure S2) contain with~~ Late Maastrichtian microfossils (Robertson et al., 2012b).~~.-~~

In the eastern range, basaltic rocks of the Melounda Formation contain abundant pelagic carbonate and are overlain by massive basalts of the Ayios Nikolaos Formation; i.e. from Ağıllar (Mandres) to Mallıdağ (Melounda) (Fig. 3a) (Baroz, 1979; Robertson and Woodcock, 1986; Hakyemez et al., 2000; Robertson et al., 2012b).~~.-~~

In the central range, pillow lava with interbedded pelagic carbonate of the Melounda Formation occurs in two settings. The first is directly above the Upper Cretaceous basal unconformity, with the ~~underlying~~ meta-carbonate platform rock ~~beneath~~; e.g. near Tirmen (Trypimeni). The second setting is as one (or several), small thrust sheets along the southern flank of the range; e.g. Değirmenlik (Kythrea), Boğaz (Bogaz) and southwest of Alevkaya Tepe (Figs. 3a, Supplementary Figure S2). Basalt

Formatted: Indent: First line: 2.5 ch

with some interbedded pelagic carbonate of the Ayios Nikolaos Formation is exposed higher in the succession (within a shear zone), near Tirmen, Arapköy (Klepini) and İncesu (Motides) (Baroz, 1979).

In the western range, basaltic lavas interbedded with pelagic carbonates of the Melounda Formation are exposed in a thrust sheet along the southern margin of the range; e.g. Geçitköy (Figs. 3, Supplementary Figure S2). Similar basaltic lavas with pelagic carbonates and local carbonate breccias are exposed along the northern margin of the range; e.g. near Karşıyaka and farther west (Baroz, 1979; Hakyemez et al., 2000; Robertson et al., 2012b). In addition, basaltic lava interbedded with pelagic chalk of the Ayios Nikolaos Formation occurs higher in the succession (southward-younging) along the eastern side of Geçitköy road (Huang et al., 2007; Robertson et al., 2012b).

In summary, the Late Maastrichtian basalts (Melounda Formation) are typically pillowed, forming repeated flows (individually up to c. 10 m thick), reaching a maximum of > 90 m in the east (near Balalan). The lavas include pink pelagic carbonate as interstitial sediment and discontinuous layers. In contrast, the Paleogene basalts (Ayios Nikolaos Formation) are commonly massive, especially in the eastern range, where individual lava flows reach c. 45 m thick, with relatively few pelagic intercalations (Supplementary Figure S2). In the central and western ranges, thinner lava flows (<10m) are intercalated with pelagic carbonates and reddish chert (formed by diagenetic replacement). During this work, we identified geochemical differences between the Late Cretaceous-Paleogene basaltic lavas of similar age along the far-eastern, eastern, central and western Kyrenia Range (see below).

4.2. Petrography

The felsic rocks are porphyritic, composed of quartz, plagioclase and sanidine,

together with subordinate biotite and rare hornblende. Large sanidine crystals (0.3-1 mm) show simple twinning, whereas plagioclase commonly has lamellar twinning (Supplementary Figure S3). Biotite laths (0.3-0.8 mm in length) are preferentially orientated parallel to flow layering. Rare quartz phenocrysts (<1%, 0.2-0.5 mm) are commonly fragmented, probably related to quenching during eruption; i.e. explosive fragmentation (Supplementary Figure S3). The groundmass comprises microcrystalline quartz and feldspar, together with minor muscovite, biotite and rare hornblende (c. 60 μ m). In some samples (e.g. nos. 14-19 and 14-20), the groundmass is cryptocrystalline, almost glassy (Supplementary Figure S3).

The basaltic rocks of both Late Cretaceous and Paleogene age generally fall into two groups in terms of mineral composition and texture. The first group (most common) is porphyritic with subhedral to euhedral clinopyroxene phenocrysts (30-40%), up to 0.6 mm long, that occur interstitially or are intergrown with subhedral feldspar (Supplementary Figure S4, Table S3). Feldspar (45-60%) forms elongate, acicular laths, up to 1 mm long. Olivine is relatively rare (<5%). The second group is ophitic, as locally observed in the western Kyrenia Range with abundant anhedral to subhedral augite phenocrysts (up to 0.6 mm in size). Most plagioclase is acicular, enclosed or surrounded by augite (Supplementary Figure S4).

Most samples are slightly to moderately altered. Amygdales and veins are infilled with secondary minerals such as calcite and zeolite. Alteration is variable, for example, pyroxene and plagioclase are heavily altered to chlorite and clay minerals (e.g. smectite/sericite). The relatively high loss on ignition values result from secondary processes, which need to be taken into account prior to rock identification and tectonic discrimination.

4.3. Whole-rock chemistry

Major-element oxides (K_2O , Na_2O and CaO) and trace elements (e.g. Rb, Sr and

384 Ba) exhibit a relatively wide compositional range, consistent with the effects of
385 alteration (see Supplementary Table S1-S2). Alteration can be inferred from variable
386 LOI values that range between 0.3-5.4 wt% for the felsic volcanics and 4.09-14.57 wt%
387 for the basaltic rocks. This is consistent with the presence of considerable amounts of
388 water- and/or carbonate dioxide-bearing minerals, for example chlorite and calcite, as
389 observed petrographically (see Supplementary Figure S3, S4). Trace elements such
390 as Ti, Zr, Y, Nb, Ta, V, Co, Th and REEs tend to be immobile during weathering and/or
391 metamorphism below amphibole facies are, therefore, preferred for rock-type
392 identification and tectonic discrimination (e.g. Pearce and Cann, 1973; Rollinson,
393 1993). For basaltic rocks, samples with high crystal contents (cumulate composition or
394 highly porphyritic) should be discounted from geochemical discrimination (Pearce,
395 1996). In general, samples with immobile element $\text{Al}_2\text{O}_3 > 20$ wt% (concentrated in
396 feldspar), Sc > 50 ppm (concentrated in clinopyroxene), or Ni > 200 ppm (concentrated
397 in olivine) are predicted to contain high amounts of cumulated minerals (Pearce, 1996).
398 On this basis, basaltic rocks nos. 14-66, 14-50, 14-51, 14-58, 14-68, 08, 16, 25 and 28
399 are discounted.

400 The more felsic assemblage is mainly rhyolitic in the classification of Winchester
401 and Floyd (1977) but shows a trachytic affinity in the revised Zr/Ti vs. Nb/Y (Fig. 6a)
402 diagram of Pearce (1996). Relatively high abundances of Th (>14 ppm) but low Co
403 concentrations (<12 ppm) are consistent with high-K and shoshonitic affinities (Fig. 6b).
404 Primitive mantle-normalised multi-element plots (Fig. 6c) show variable enrichments
405 and depletions in large-ion lithophile elements (LILEs; e.g. Cs, Rb, Ba, K and Sr) that
406 are influenced by alteration. The samples show negative anomalies of Nb, Zr and Ti,
407 coupled with marked positive anomalies of Th/U, Pb, Nd and probably Sm (Fig. 6c).
408 Total REE contents of the felsic assemblage (Fourkovouno Formation) vary from 115
409 to 173 ppm. La ranges from 29-44 ppm, $(\text{La}/\text{Yb})_{\text{N}}=12.63\text{-}18.98$, suggestive of an alkalic
410 affinity. The ratios $(\text{La}/\text{Sm})_{\text{N}}=6.13\text{-}7.22$ and $(\text{Gd}/\text{Yb})_{\text{N}}=1.26\text{-}1.68$ suggest a large LREE

fractionation but low to moderate HREE fractionation (Fig. 6d). Pronounced negative Eu anomalies ($\text{Eu}/\text{Eu}^*=0.32\text{-}0.48$) are consistent with plagioclase crystallization, as seen petrographically (see Supplementary Figure S3). In general, the samples are comparable to upper continental crust (UCC). The basaltic rocks plot in the fields of basalt and alkali basalt, with intermediate degrees of fractionation ($\text{Zr}/\text{Ti}=0.01\text{-}0.02$) and moderate alkalinity ($\text{Nb}/\text{Y}=0.1\text{-}1.8$) on the rock-type discrimination plot (Fig. 7a). The basalts have a calc-alkaline affinity, except for samples from Mallıdağ (no. 01), Tirmen (no. 19-51), Değirmenlik (no. 14-64) and İncesu (no. 19-67) that exhibit high-K and shoshonitic affinities (Fig. 7b). The trace element compositions of the basaltic rocks exhibit wide ranges of Sr, Ba, Rb, Th, Ta, Nb and Ce. However, Zr, Hf, Sm, Ti and Y generally comparable with enriched mid-ocean ridge basalt (E-MORB) and oceanic island basalt (OIB) (Fig. 8a-c).

The basaltic rocks are characterised by variable REE concentrations (67.2-239.4 ppm) and light REE enrichments ($(\text{La}/\text{Yb})_N=3.18\text{-}19.35$) (Fig. 8d). Samples from the central range, at Değirmenlik and Tirmen, show much higher total REE contents and steeper REE patterns. Westwards in the Kyrenia Range, the patterns become smoother, with less pronounced enrichment in LREEs. In general, no marked Eu anomaly (~ 1) is observed.

4.4. Whole-rock Sr-Nd-Hf isotopes

The selected basaltic samples have relatively uniform whole-rock Sr-Nd-Hf isotopic characteristics. Adopting a Late Cretaceous eruption age based on the paleontological dating (Robertson et al., 2012b), the calculated initial $^{87}\text{Sr}/^{86}\text{Sr}$ ratios range from 0.705054 to 0.705701, with $\epsilon\text{Nd}(t)$ values of 2.27 to 5.76. Initial $^{176}\text{Hf}/^{177}\text{Hf}$ and $\epsilon\text{Hf}(t)$ values are 0.282934-0.283017, 7.36-10.14, respectively (see Supplementary Table S2).

4.5. Zircon U-Pb geochronology

Zircon crystals separated from the felsic rocks are typically angular to sub-angular, 30-160 μm long. The grains have variable internal textures, including banded zoning (e.g. zircon 1), concentric zoning (nos. 2, 4, 5 and 12), and minor sector zoning (nos. 10, 11; Fig. 9a). Fractures (or depressions) and some inherited cores are affected by fluid-related chemical alteration (nos. 6 and 8) (Fig. 9a). The zircon grains have high Th/U ratios (0.19-1.24), suggestive of a magmatic origin (Rubatto, 2002). Twelve zircon grains were analysed with concordance levels ranging from 99-105%. The calculated $^{206}\text{Pb}/^{238}\text{U}$ age with the generally lowest possible error was used for age determination. The much older zircon grain (no. 4) is subhedral with a patchy xenocrystic texture, suggestive of a recycled origin (Fig. 9a). The concordant or nearly concordant data (100-102%) with a young age distribution yielded weighted mean $^{206}\text{Pb}/^{238}\text{U}$ gave ages of 74.0 ± 0.6 Ma, 74.0 ± 0.4 Ma and 71.7 ± 0.7 Ma (Fig. 9b-d).

5. Discussion

5.1. Genetic type/source characteristics

For the felsic rocks, the high LREE/HREE (e.g. $(\text{La}/\text{Yb})_{\text{N}}=12.63-18.98$) and negative Eu anomalies, resemble model UCC but with depletion in Eu, Sr and P (Fig. 6c-d). The A/CNK (molar ratio of $\text{Al}_2\text{O}_3/(\text{CaO}+\text{Na}_2\text{O}+\text{K}_2\text{O})$) ratio is useful to identify the source of felsic rocks (Chappell and White, 1992) but is questionable for altered rocks like those of the Kyrenia Range. SiO_2 , P_2O_5 and Th can also be used as indices of the fractional crystallization of a felsic melt (Chappell and White, 1992), but again may be affected by alteration (specially SiO_2). In our samples, P_2O_5 decreases with increasing SiO_2 and Th (Fig. 10a-b), trends that are indicative of I (igneous)- or A (anorogenic)-type granites. The relatively low contents of Zr, Nb, Y, La, Ce, Zn and Ga further

characterise these rocks as I-type (Fig. 10c-d). [The abundances of Th and Co, in particular hint at a shoshonitic composition \(Hastie et al., 2007\).](#)

The trace element patterns of the basaltic rocks show a wider range of variation (Fig. 8). E-MORB and OIB-type basalts were derived from relatively enriched sources, specifically for the basalts of the Karpas Peninsula, eastern range and most of the central range. The lesser enrichment of Nb within the basalts of the central range at Ergenekon (no. 31), Arapköy (no. 14-62) and Alevkaya Tepe (no. 14-49), and the western range at Karsiyaka (no. 14) and Geçitköy (no. 10), can be explained by a relatively depleted mantle source, together with subduction modification. There is little evidence of the addition of radiogenic Sr from seawater, either to the source melt or during post-magmatic alteration. The Sr vs. Nd isotope correlation diagram (Fig. 11a) does not show a significant shift towards higher initial $^{87}\text{Sr}/^{86}\text{Sr}$ ratios as would be expected in both of these cases.

The basalts all have positive $\epsilon\text{Nd}(t)$ values (+2.27 - +5.76), consistent with derivation from an OIB-like source mantle (Fig. 11a). The samples plot directly on the terrestrial array, overlapping with the Nd isotope compositions of Etna and the Aeolian Arc, but intermediate between these settings in Hf isotopes (Fig. 11b). The samples that show trace element evidence of a subduction-modified source, based on relatively flat MORB-normalised multi-element patterns and negative Nb anomalies (no. 10, Geçitköy), plot in the Aeolian Arc-type mantle [field](#). Conversely, samples from the Karpas Peninsular, at Balalan, plot close to the field of Etna and exhibit less subduction modification (Fig. 11b). The trace element and isotopic data, therefore, yield consistent results.

The involvement of a crustal component is also suggested by incompatible element ratio plots, on which the basalts exhibit two distinct trends. The samples with relatively strong subduction signatures plot away from the MORB array (Fig. 12a), fingerprinting a subduction-related lithospheric mantle end member. A probable lower

crust input is suggested by relatively higher La/Nb and Ba/Nb ratios (e.g. Wang et al., 2004). In contrast, an OIB-like mantle component dominated most of the other basalt samples, as indicated by relatively low La/Nb, Zr/Nb but high Nb/Y ratios (Fig. 12b). The ratios of highly incompatible elements with small distribution coefficients (e.g. Zr/Nb, Nb/Th, Nb/La) are sensitive indicators of mantle source heterogeneity (Saunders et al., 1988; Condie, 2003; Pearce and Stern, 2006). The samples from the western locations fall in the fields of an enriched component and upper continental crust (Fig. 12c), which could represent mantle mixing with continental sediments (subducted) or crustal contamination. In contrast, the samples from the easterly locations resemble those of the OIB/recycled component, and cluster around a different batch-melting trajectory (10-20%) (Fig. 12c) that is representative of recycled oceanic lithosphere (Saunders et al., 1988). We, therefore, infer that the basalts are likely to have been derived from multiple sources (i.e. two or more source components were involved).

5.2. Tectonic discrimination

On the Nb vs. Y diagram (Fig. 13a), the felsic volcanics show similarities with volcanic arc granites and syn-collisional granites. One sample from Geçitköy (no. 16-08), with relatively high Nb and Y concentrations, plots near the field of within-plate granite. On the Th/Yb vs. Ta/Yb diagram (Fig. 13b), the volcanics mainly erupted on active continental margins. An active continental margin setting is therefore inferred for the felsic volcanics.

Basalts of the Kyrenia Range (e.g. easterly locations) mostly exhibit a within-plate affinity (Fig. 8). However, some samples, especially in the west, have a subduction-related signature (e.g. negative Nb anomalies) (Fig. 8b-c) akin to volcanic arc basalts. Most samples have a within-plate affinity on the Zr/Y vs. Zr (Fig. 13c) diagram, except

for two samples, from Ergenekon and Arapköy (central range), that straddle the island arc and MORB fields. The basalts overlap the MORB and back-arc basin basalt fields on the Ti vs. V plot (Fig. 13d). Clinopyroxene compositions, as shown in the ternary $\text{TiO}_2\text{-MnO-Na}_2\text{O}$ plot (see Supplementary Figure S5), indicate a dominantly within-plate affinity. The relatively high TiO_2 contents of the clinopyroxenes point to an enriched MORB-related source (see Supplementary Figure S5). Overall, the inferred subduction contribution decreases from west to east along the range.

5.3. Global comparison of the Kyrenia Range volcanic rocks

Global ~~comparison (see Supplementary Figure S6) lend~~ comparisons (see Supplementary Figure S6) lend -support to our interpretation of the Kyrenia Range volcanic rocks as being subduction-related. In particular,

~~Within the Late Cretaceous felsic rocks of the Kyrenia Range,~~ the enrichment of Rb, Th and Sm ~~and versus~~ the depletion of Ti, Ba, Nb, P compared to primitive mantle, are characteristic of volcanic arc granites (Fig. 6c) (Pearce et al., 1984).

The Kyrenia Range Maastrichtian and Paleogene basalts of within-plate affinities (e.g. Balalan, eastern range) mainly ~~have~~ MORB-normalised spider patterns that are similar to the Pliocene-Quaternary OIB-type basalts of the Tyrrhenian Sea, W Mediterranean (Peccerillo, 2017). The MORB-normalised patterns of these basalts suggest a within-plate eruptive setting, with a small sub-crustal lithosphere metasomatic influence (Kastens et al. 1988; Peccerillo, 2017).

Back-arc basalts of the Early Cretaceous Sarmiento Complex, southern Chile (Saunders et al., 1979) and the Miocene Japan Sea (Tamaki et al., 1990) resemble a minority of the Kyrenia Range basalts (c. 10%; e.g. Geçitköy, western range), ~~but~~ although with lower abundances of K, Rb, Ba, Th, Ta and Nb. The relative ~~depletions~~

~~depletion of these elements is characteristic of~~ ~~are likely to be representative of~~
eruption in a relatively wide, evolved back-arc basin (e.g. Bransfield Strait; Weaver et
al., 1979). However, another possible setting is eruption in a back-arc basin that was
influenced by oblique extension (e.g. Guaymas pull-apart basin; Brune et al., 2012).
The relatively flat MORB-normalised patterns, akin to E-MORB, of both the Bransfield
Strait and Guaymas back-arc basin basalts (e.g. Saunders et al., 1982; Keller et al.,
2002) point to substantial involvement of depleted MORB sources.

5.4. Comparison with the adjacent Tethyan region

Late Cretaceous and Paleogene arc-related rocks; i.e. similar in age of those in
the Kyrenia Range are critical to any regional magmatic-tectonic interpretation.

In western Cyprus, ~~there is a~~ Late Cretaceous arc-derived volcanogenic
succession (Kannaviou Formation) ~~that~~ overlies the Troodos ophiolite. The
volcaniclastic sediments ~~there~~ were derived from intermediate-felsic ~~composition~~
eruptions, c. 6 Ma before the felsic eruptions in the Kyrenia Range (c. 80 vs. 74 Ma).
Grain size and textural evidence suggest that, depending on prevailing wind direction
and strength, the eruptive centers are likely to have been <100 km away (Chen and
Robertson, 2019). The centers ~~for the Kannaviou Formation were~~ ~~could have been~~
located to the south of the Kyrenia Range or its lateral extension, areas that are not
now exposed. The Kyrenia Range was affected by south-directed thrusting during Mid-
Eocene and Late Miocene-earliest Pliocene (Baroz, 1979; Robertson et al., 2012b,
2014; McCay and Robertson, 2013; Robertson and Kinnaird, 2016), such that the distal
(southerly) edge of the inferred over-riding active margin (see below) could have been
subducted, structurally over-ridden or eroded. In other words, additional arc units could
have existed to the south of their present location in the Kyrenia Range.

To the northeast (c. 200 km), the Misis-Andırın Range in southern Turkey is an

extension of the Kyrenia lineament (Robertson et al., 2004). The outcrop, known as the Misis-Andırın Complex, includes massive basalts, pillow lavas, lava breccias and volcanoclastic debris-flows. These lithologies were previously inferred to be Miocene (Floyd et al., 1991, 1992). However, inter-lava sediments there have been dated as Campanian-Maastrichtian (Robertson et al., 2004), similar in age to the oldest basaltic rocks in the Kyrenia Range. The Miocene sediments are structurally intercalated. The basalts of the Misis-Andırın Complex are tholeiitic, with mildly enriched, subduction-related characteristics (e.g. moderate La/Nb ratios), and are interpreted as of back-arc basin origin (Floyd et al., 1991, 1992), which is consistent with our interpretation of the Kyrenia Range basalts ~~(Floyd et al., 1991, 1992)~~ (Fig. 14a-b, Supplementary Figure S6).

The Kyrenia Range is backed to the north by the Alanya Massif (Fig. 1), which has been interpreted as a continental fragment, separate from the Tauride carbonate platform to the north (Robertson et al., 2012a; Çetinkaplan et al., 2016). The Alanya Massif is represented by the Late Precambrian-Mesozoic, mainly meta-sedimentary rocks. High pressure-low temperature (HP-LT) metamorphics document Late Cretaceous (82-80 Ma) subduction/exhumation (Oberhänsli et al., 2010, 2013). This is inferred to have taken place along the northern, active margin of the Southern Neotethys (Çetinkaplan et al., 2016; Robertson and Parlak, 2020). The felsic volcanics (Fourkovouno Formation) in the Kyrenia Range could have erupted along the southern (trailing) margin of the Alanya microcontinent where they could have escaped metamorphism. However, they could also represent a separate more southerly unit.

Late Cretaceous continental margin arc-type magmatic rocks also occur extensively farther east, in SE Turkey, as two separate belts. ~~The Kyrenia Range volcanics are most comparable with volcanogenic units that occur within an extension of the same (or similar) tectonic lineament; i.e. close to the leading edge of the over-riding Tauride (Eurasian) continent.~~

The most southerly, known Late Cretaceous arc rocks in SE Turkey are exposed in the Helete area, part of a zone of frontal thrust sheets (Yıldırım, 2015; Nurlu et al., 2016) (Fig. 1). Two thrust assemblages are exposed, both cut by granites that have been ~~Arc-related volcanogenic units there comprise basaltic andesite, andesite, dacite, rhyolite and common tuffaceous rocks. Cross-cutting granites have been~~ radiometrically dated at 93-83 Ma (Cenomanian-Campanian) (Nurlu et al., 2016). The lower of the assemblages, the Helete volcanics, comprises arc-related basaltic andesite, andesite, dacite, rhyolite and common tuffaceous rocks. The upper of the two thrust assemblages is a dismembered Late Cretaceous supra-subduction zone-type ophiolite (Meydan ophiolite) (Yıldırım, 2015; Nurlu et al., 2016). The cross-cutting intrusive rocks sealed the thrust sheets during the Late Cretaceous. The granitic rocks ~~The intrusive rocks~~ are interpreted as a calc-alkaline, subduction-related suite that formed by mixing of mantle and crustal sources, as suggested by Sr-Nd-Pb isotopic data (Nurlu et al., 2016). The granitic rocks exhibit relatively flat chondrite-normalised REE patterns, similar to E-MORB (Nurlu et al., 2016) (Supplementary Figure S6). In contrast, the chondrite-normalised REE patterns of the Kyrenia Range felsic rocks are similar to UCC (Fig. 6c), suggesting a more ~~continentally-influenced proximal (northerly)~~ setting.

The second belt of arc-related magmatism is located up to c. 100 km ~~located~~ farther north and north-east, in the Keban (Malatya), Göksun (Kahramanmaraş) and Baskil (Elazığ) regions (Fig. 1) (Yazgan and Chessex, 1991; Parlak, 2006; Rızaoğlu et al., 2009; Karaoğlu et al., 2013, 2016). ~~(c. 100 km), where a~~ Arc-granitic rocks (Baskil intrusives) cut both the Late Cretaceous Göksun ophiolite and equivalents (i.e. Berit meta-ophiolite (=North Berit ophiolite), Kömürhan ophiolite and İspendere ophiolite) that are inferred to have formed above a northward-dipping subduction zone (Parlak, 2006). Similar ~~the arc granites also cut the~~ structurally over-riding Malayta metamorphic unit, ~~which that~~ is interpreted as the southern part of the Tauride microcontinent. The

Göksun ophiolite (and equivalents) and the Malatya metamorphic (Tauride) units were juxtaposed along an active continental margin when the arc-granitic rocks were intruded (Robertson et al., 2006; Rızaoğlu et al., 2009; Karaoğlu et al., 2013, 2016).

The extrusive rocks of the Göksun and related ophiolites can also be broadly correlated with widespread assemblages of basic igneous rocks and volcanogenic sediments in the Malatya-Elazığ region, known as the Yüksekova Complex (Perincek and Özkaya, 1981; Aktaş and Robertson, 1984). The basalts have a tholeiitic to tholeiitic-transitional character, variable Zr/Y (1.5-6) ratios and mantle-array-aligned Sr-Nd isotopes (Ural et al., 2015). These basic volcanics were derived from a depleted mantle source without a significant crustal contribution; i.e. from an intra-oceanic setting within the Southern Neotethys (Ural et al., 2015).

The arc rocks that cut both the Göksun (and related) ophiolites and the Malatya Metamorphics (Baskil intrusives) in the northern belt are mainly hornblende-biotite granodiorites and 'normal' granites, dated radiometrically at 88-82 Ma (Santonian-Campanian) (Parlak, 2006; Rızaoğlu et al., 2009; Karaoğlu et al., 2016); i.e. up to c. 5 Ma younger than Helete area granites in the south. These southerly granitic rocks have I-type, calc-alkaline arc affinities, with both mantle and crustal-derivation features (Nurlu et al., 2016). The Baskil intrusions of the northern belt are relatively enriched in LREE (Parlak, 2006; Rızaoğlu et al., 2009) compared to the Helete granites of the south, which suggests a more crustally influenced source during magma genesis for the former. Similarly, the Kyrenia Range basalts have relatively high Zr/Y (2.8-7.7) and Nb/Y (Figs. 12b, 13c), suggestive of a magmatic contribution from continental crust or subducted continentally derived sediments (Zindler and Hart, 1986; Pearce, 1996).

The northerly arc has been explained by northward subduction of Mesozoic oceanic basin (Berit ocean) that was located between two continental units; i.e. the Bitlis and Pütürge massifs in the south and the Tauride continent represented by the Malatya Metamorphics in the north (Robertson et al., 2012a; Karaoğlu et al., 2013,

2016; Barrier et al., 2018). The Late Cretaceous arc-related intrusive magmatism of the northern belt in SE Turkey (Baskil intrusives) evolved towards collisional, including shoshonitic compositions (74-72 Ma), which are interpreted to indicate continental and post-collisional settings (Kuşcu et al., 2013; Erturk et al., 2018; Sar et al., 2019). The shoshonitic compositions of some of the granitic rocks, together with the evidence of HP/LT metamorphism, are consistent with the collision of the Tauride carbonate platform (i.e. Malatya Metamorphics) to the north with the Bitlis and Pütürge continental units to the south. However, the origins of shoshonites remain controversial because the necessary partial melting of mantle and interaction with subduction-related fluids can take place in a variety of pre-, syn- and post-collisional settings (e.g. Campbell et al., 2014).

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Several authors have correlated the Late Cretaceous Göksun and related ophiolites in the north with the Meydan ophiolite (Berit meta-ophiolite of Yılmaz et al., 1993) in the south. If correct, the granitic intrusives in both belts originated as different parts of a single arc lineament that straddled the northerly active continental margin of the oceanic basin. The southerly arc and ophiolitic rocks (Helete-Meydan) were emplaced southwards during the latest Cretaceous, although further southward thrusting took place, mainly during the Eocene and Miocene (Perinçek and Kozlu, 1984; Yılmaz et al., 1993).

Formatted: Font color: Purple

In many reconstructions, oceanic crust still remained between the Bitlis-Pütürge continental units and Arabia until the Miocene (Yılmaz, 1993; Robertson et al., 2012a; Barrier et al., 2018; van Hinsbergen et al., 2020). The Bitlis and Pütürge massifs have been generally correlated with the Alanya metamorphic massif to the north of Cyprus (with or without a direct continuation) (Çetinkaplan et al., 2016). If correct, the Late Cretaceous Kyrenia Range felsic arc volcanism (and that of the Kannaviou Formation) could have been located to south of the Alanya continent, with a possible eastward continuation to south of the Bitlis and Pütürge continental units. (Parlak, 2006;

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Robertson et al., 2006). Examples occur in the Keban (Malatya), Göksun (Kahramanmaraş) and Baskil (Elazığ) regions (Fig. 1) (Yazgan and Chessex, 1991; Parlak, 2006; Rızaoğlu et al., 2009; Karaoğlu et al., 2013, 2016). The are rocks are mainly hornblende biotite granodiorites and 'normal' granites that are radiometrically dated as 88–82 Ma (Santonian–Campanian) (Rızaoğlu et al., 2009; Karaoğlu et al., 2016). The granitic rocks have I-type, calc-alkaline arc affinities, with both mantle and crustal derivation features (Parlak, 2006; Rızaoğlu et al., 2009; Karaoğlu et al., 2013, 2016). The intrusions are relatively enriched in LREE (Parlak, 2006; Rızaoğlu et al., 2009) compared to the Holote granites, which suggests a more crustally influenced source during magma genesis. The above granitic rocks are located in a structurally higher position than the Kyrenia and Holote arc rocks (i.e. above the Pütürge metamorphic massif) and are likely to have a different region. The northerly arc have been explained by northward subduction of a small oceanic basin (Berit ocean) that was located between two continental units; i.e. Bitlis–Pütürge in the south and the Tauride continent (Malatya) in the north (Robertson et al., 2012a; Karaoğlu et al., 2013, 2016; Barrier et al., 2018).

All of the above extrusive and intrusive felsic arc rocks differ from the Late Cretaceous ophiolitic and oceanic arc rocks in the region. These include a widely exposed assemblage (Malatya–Elazığ region) of basic igneous rocks and volcanogenic sediments, known as the Yüksekova Complex (Perincek and Özkaya, 1981; Aktaş and Robertson, 1984). The basalts have a tholeiitic to tholeiitic transitional character, variable Zr/Y (1.5–6) ratios and mantle array aligned Sr–Nd isotopes (Ural et al., 2015). These basic volcanics were derived from a depleted mantle source without a significant crustal contribution; i.e. from an intra-oceanic setting within the Southern Neotethys (Ural et al., 2015). In contrast, the Kyrenia Range basalts have higher Zr/Y (2.8–7.7) and Nb/Y (Figs. 12b, 13c), suggestive of a magmatic contribution from continental crust or subducted continentally derived sediments (Zindler and Hart, 1986; Pearce, 1996).

Formatted: Font color: Red

Paleogene magmatic rocks are also present in SE Turkey. Eocene volcanic rocks and associated minor intrusions, known as the Maden Complex, unconformably overlie and cut the Pütürge and Bitlis ~~continental metamorphic units~~metamorphosed continental units (Fig. 1) (Hempton, 1985; Yazgan and Chessex, 1991; Aktaş and Robertson, 1984, 1990; Yılmaz, 1993; Elmas and Yılmaz, 2003; Robertson et al., 2006; Erturk et al., 2018). The volcanics show enrichment in LILEs and relative depletion in Nb, Ta and Ti compared to MORB (Erturk et al., 2018), similar to a minority of the subduction-related volcanics in the Kyrenia Range (see Supplementary Figure S6). The Eocene Maden Complex magmatism is widely proposed to relate to back-arc rifting of the (Eurasian) active continental margin during the later stages of northward subduction of the Southern Neotethys, in keeping with the widespread view that collision with Arabia was delayed until the Miocene (Aktaş and Robertson, 1984, 1990; Yılmaz, 1993; Yılmaz et al., 1993; Yiğitbaş and Yılmaz, 1996; Robertson et al., 2006, 2007; van Hinsbergen et al., 2020). Oblique convergence, ~~as~~ proposed by several authors (Aktaş and Robertson, 1990; Elmas and Yılmaz, 2003), could have resulted in along-strike variation in the geochemical signatures of extension versus subduction in the Kyrenia Range, although more geochemical work ~~is~~ needed, regionally and globally to test this ~~possibility~~hypothesis.

Farther north, Eocene granitic rocks locally cut the continental Malatya metamorphic unit (Doğanşehir region) (Perinçek and Kozlu, 1984; Karaoğlu et al., 2013, 2016). However, no extrusive equivalents are known and, as noted above, the magmatic rocks in this lineament are unlikely to correlate directly with the Kyrenia Range.

During the Miocene, the active continental margin bordering the remnant Southern Neotethys collided with the Arabian passive continental margin to the south. In both the Kyrenia Range and SE Turkey, the Late Cretaceous and Paleogene magmatic rocks are exposed close the thrust front (Eurasia), implying that at least tens of kms of

fore-arc crust have been lost (subducted or overridden), potentially removing additional frontal arc-related crust, both Late Cretaceous and Paleogene. Thus, the Kyrenia, Kannaviou (W Cyprus) and ~~Helete-Maden Complex~~ magmatism could all represent surviving products of a regional ~~Late Cretaceous~~active continental margin-~~are~~.

5.5. Tectono-magmatic hypotheses

In the light of the regional setting, ~~t~~The felsic volcanics (Fourkovouno Formation, c. 74 Ma), with continental margin arc affinities, are interpreted to have resulted from northward subduction beneath continental crust, possibly the Alanya continental unit to the north (Çetinkaplan et al., 2016; Robertson et al., 2020), of the Taurides. ~~The~~ felsic volcanism is characteristic of relatively advanced subduction, suggesting that earlier eruptive centers existed along the active continental margin. The early volcanism is likely to be represented by the Campanian (c. 80 Ma) arc-derived volcanogenic sediments (Kannaviou Formation) in western Cyprus (Robertson, 1977; Gilbert and Robertson, 2013; Chen and Robertson, 2019). The underlying mantle wedge (sub-continental lithosphere) was metasomatically enriched during the Late Cretaceous subduction, with implications for the subsequent basaltic volcanism.

For the Maastrichtian to Late Eocene basaltic volcanics, there are three alternative tectono-magmatic models. The first model envisages a marginal basin setting related to northward subduction. ~~The Southern Neotethys remained partly open between the Taurides and the Troodos ophiolite during the Late Cretaceous to Mid-Eocene. Subduction of the remnant ocean gave rise to the basaltic volcanism, including the Misis-Andırın Range.~~ Compared to typical back-arc basin basalts (Fig. 14a-b), the ~~basalts of the Kyrenia Range~~ basalts (and the Misis-Andırın Range) have relatively high Th/Nb but low La/Nb ratios. The high abundances of Th are likely to reflect a continental crust ~~influence~~input (Floyd et al., 1991). The variation in La/Nb ratios of the

basalts could reflect the influence of a within-plate mantle source (i.e. easterly basalts) and/or the arc maturity; (i.e. the arc-like western basalts could have erupted closer to the distal (oceanward) edge of the convergent margin). However, a problem with this model is the absence of evidence for a Paleogene volcanic arc to the south. There is no evidence of arc-derived fallout tuff or volcanoclastic sediments in the Maastrichtian-Paleogene basaltic succession, in contrast with, for example, the SW Pacific Mariana and Tonga marginal basins (Clift, 1994; Bryan et al., 2004).

The second model involves an extensional (or transtensional) setting. In this interpretation, the Late Maastrichtian and Paleogene 'enriched' within-plate volcanism was triggered by strike-slip (or transtension) along the ~~South Neotethyan~~ active continental margin. One option is that the volcanism was linked to the anticlockwise paleorotation of the Troodos ophiolite to the south during Late Campanian-Early Eocene (Clube et al., 1985; Clube and Robertson, 1986; Morris et al., 2006, 2015; Hodgson et al., 2010). More generally, oblique convergence of the African plate with respect to Eurasia could have resulted in segmentation of the South Neotethyan active margin into subducting segments with arc magmatism, and also strike-slip segments without arc magmatism (Aktaş and Robertson, 1990). A modern example is the Andaman Sea where oblique convergence between the Asian and the Australian plates resulted in segmented arc magmatism (e.g. Curray et al., 2005). However, there is limited information from different modern settings to test this hypothesis.

The third, preferred, scenario is that both the felsic and the basaltic volcanism are related to northward subduction during stages in the closure of the ~~Southern Neotethys~~ Mesozoic ocean basin to the south. The Late Cretaceous magmatism can then be explained by regional northward subduction (nearly orthogonal) resulting in continental margin arc volcanism. Slab-derived and residual slab components (fluid, melt) were released into the upper mantle lithosphere (e.g. Pearce, 1983; Harangi et al., 2007). The Late Maastrichtian and Paleogene basaltic volcanics then erupted in

an extensional (or transtensional) setting, perhaps related to slab rollback and incipient marginal basin formation. Several different mantle sources are likely to have contributed along the Kyrenia Range. The variable subduction-related signatures were inherited from the previously modified mantle lithosphere in this interpretation. Examples of subduction-influenced settings that were affected by later extension-related melting include the Rio Grande Rift, west USA (e.g. Riecker, 1979) and the Neogene north Aegean-west Anatolia volcanic province (e.g. Şengör and Yılmaz, 1981; Ersoy and Palmer, 2013). In summary, the available evidence is consistent with Late Maastrichtian-Paleogene incipient marginal basin formation along the northern, active margin of ~~the Southern Neotethys~~ an ocean basin (Southern Neotethys) that did not close until the Miocene.

5.6. Testing alternative tectonic models

Finally, the Late Cretaceous-Paleogene volcanism in N Cyprus and SE Turkey can be used to test alternative plate tectonic models, including those by Robertson et al. (2012a), Karaoğlu et al. (2013, 2016), Maffione et al. (2017), Barrier et al. (2018), McPhee and van Hinsbergen (2019) and van Hinsbergen et al. (2020), as summarised in Figure 15.

In Reconstruction 1 (Fig. 15a-c) (Robertson et al., 2012a), the Southern Neotethys rifted in the Triassic, while Paleotethys evolved into the Northern Neotethys (İzmir-Ankara-Erzincan ocean). The Kyrenia Range is restored to the northern passive margin of the Southern Neotethys. Both intra-oceanic and continental margin subduction zones (northward-dipping) were active in the Southern Neotethys during the Late Cretaceous. The continental margin arc volcanism was constructed on, or to the south of one, or more, continental blocks (microcontinents), represented by the Kyrenia-Alanya-Pütürge-Bitlis lineament in the south and the Malatya-Keban lineament farther north. This reconstruction is compatible with our new evidence, ~~while~~

prompting questions concerning the implications of two subparallel arc magmatic belts (outside the present scope).

Reconstruction 2 (Fig. 15d-f) (Barrier et al., 2018) has many common features with (1), in particular the arc volcanism along the northern margin of the Southern Neotethys. However, the Kyrenia, Malatya and Pütürge-Bitlis massifs are restored as a single elongate continental unit rather than as microcontinents, as in (1). Also, little or no oceanic crust is inferred between the southerly continental unit and the Tauride carbonate platform to the north (i.e. Göksun and related ophiolites). This is difficult to reconcile with the Late Cretaceous and Eocene granitic magmatism cutting the Mesozoic Malatya (Keban) platform unless large-scale strike-slip and terrane duplication has taken place for which there is little supporting evidence.

In Reconstruction 3 (Fig. 15g-h) (Maffione et al., 2017; van Hinsbergen et al., 2020), the Kyrenia Range is restored to the southern passive margin of the Mesozoic Tethyan ocean. Late Cretaceous ophiolites that are presently located in several c. E-W, sub-parallel belts (i.e. N and S of the Bitlis and Pütürge massifs) originated above a single subduction zone that originated within Neotethys, dipping generally to the east or northeast (Stampfli and Borel, 2002; Moix et al., 2008). The subduction zone rolled back generally westwards until it collided with the Tauride and Arabian continents. In the south, the subduction zone broke through the oceanic gap between the African and Tauride continents, 'invaded' and replaced the Southern Neotethys with Late Cretaceous supra-subduction zone fore-arc crust. Fore-arc ophiolites were then obducted generally northwards and southwards over the opposing continents (van Hinsbergen et al., 2020). Two alternatives ~~have been~~are suggested; in one, the Bitlis and Pütürge continental units are correlated with Arabia (Maffione et al., 2017), whereas in the other these units are correlated with the southern margin of the Tauride ~~continents-continent~~ (van Hinsbergen et al., 2020). The Kyrenia Range (Trypa Group) is placed on the N African continental margin in both alternatives. The metamorphism

of the Mesozoic carbonate platform in the Kyrenia Range (Trypa Group) resulted from northeastward underthrusting/subduction beneath the obducted forearc Troodos ophiolite (van Hinsbergen et al., 2020). However, there are several problematic aspects: (1) seismic tomography in the Eastern Mediterranean supports generally northward subduction (e.g. Hafkenscheid et al., 2006), without evidence of a relict southward-dipping slab, as implied by the inferred northward emplacement of some Late Cretaceous ophiolites in S Turkey (~~it is however unclear for how long such a trace would be preserved~~); (2) it does not allow for the tens to hundreds of km of subduction required to produce the Late Cretaceous arc magmatism in N Cyprus and SE Turkey. Specifically, ~~the magmatically crosscut~~ Malatya metamorphic unit, cut by Late Cretaceous granites, is instead restored to the lower, downgoing plate (i.e. beneath the south margin of the Tauride continents) ~~implying-suggesting~~ northward subduction; (3) the model (van Hinsbergen et al., 2020) does not allow for any Late Cretaceous continental arc-related magmatism in northern Cyprus, SE Turkey or Iran (e.g. Agard et al., 2011); (4) in this model, the proposed Late Eocene-Miocene (post 45 Ma) northward subduction is ~~also~~ too young to explain the Late Cretaceous-Paleogene magmatic rocks in the Kyrenia Range and SE Turkey; (5) arc magmatism of Oligocene-Miocene age in S Turkey is absent, although this would be expected if significant northward subduction took place during this time, as in model 3.

6. Conclusions

- Latest Cretaceous mainly felsic volcanic rocks and latest Cretaceous-Paleogene mainly basaltic volcanic rocks are exposed throughout the Kyrenia Range, separated by a thrust.
- Mapping and logging of the key, well-exposed ~~segment of the~~ western Kyrenia Range segment, N Cyprus shows that the Late Cretaceous felsic arc volcanics

occur as variably dismembered thrust sheets in the southerly, frontal part of thrust belt.

- The primary eruptive age of the felsic arc volcanics (Fourkovouno Formation) is ~~determined as~~ c. 74.0 Ma (Late Campanian), based on U-Pb dating of zircons.

- The latest Cretaceous-Paleogene basalts vary geochemically along the Kyrenia Range (E-W). The volcanics in the east resemble normal rift products. In contrast, the volcanics in the west have a subordinate subduction-related signature (e.g. negative Nb anomaly).

- Sr-Nd-Hf isotopic data for the latest Cretaceous-Paleogene basalts suggest that they were derived from several OIB-like mantle sources, ~~probably~~ with the involvement of a crustally ly-derived (recycled) component.

- The Kyrenia Range felsic volcanics show some geochemical similarities with the late Cretaceous (~~93-83 Ma~~) arc-related rocks in S Turkey (Helete granite; 93-83 Ma). These felsic volcanics are interpreted to represent a fragment of a mature magmatic arc that was active along a regional-scale active continental margin. Regional reconstructions suggest that the Helete granites and related ophiolitic rocks were emplaced southwards from an oceanic basin (Berit ocean) to the north.

- The latest Cretaceous-Paleogene volcanics in the Kyrenia Range have geochemical similarities with the Middle Eocene (c. 47 Ma) basaltic rocks in S Turkey (Maden Complex), suggesting an along-strike continuation of the same active continental margin.-

- The latest Cretaceous-Paleogene basaltic volcanics in the Kyrenia Range are interpreted as the products of an incipient marginal basin that developed during northward subduction, possibly related to oblique convergence, prior to collision with Arabia during the Miocene.-

- 884 ● In the light of alternative regional ~~Having taken account of a range of~~ tectonic
885 models, the Late Cretaceous and Paleogene magmatic rocks ~~in Kyrenia Range,~~
886 of N Cyprus are interpreted to represent stages in the development of the northerly
887 active continental margin of the Southern Neotethys.

888
889 Acknowledgements

890 We thank Richard Hinton, Steffen Kutterolf and Dick Kroon for scientific discussion.
891 Mike Hall is thanked for thin section and polished blocks preparation. Nick Odling kindly
892 assisted with the XRF analysis. Antony Morris is thanked for providing the samples of
893 basalt from drill cores that were originally collected for paleomagnetic studies. Rory
894 McKavney kindly assisted with the fieldwork in northern Cyprus. The first author
895 gratefully acknowledges the receipt of a joint studentship of the Principal's Career
896 Development PhD Scholarship and Edinburgh Global Research Scholarship. The
897 authors are grateful for financial support via the Natural Environment Research Council
898 Ion Microprobe Facility (to A.H.F. Robertson) to carry out the secondary ion mass
899 spectrometry U-Pb dating of detrital zircons. Fieldwork and geochemical analysis were
900 aided by financial support from the International Association of Sedimentologists
901 [Postgraduate Grant Scheme], the Mineralogical Society of Great Britain and Ireland
902 [Postgraduate Student Bursary Awards], and the Edinburgh Geological Society
903 [Clough Fund], all to the first author. Additional financial support was provided by the
904 John Dixon Memorial Fund. Osman Parlak is thanked for discussion of SE Turkey
905 geology. The manuscript benefitted from comments by Pamela Kempton, Fatih
906 Karaoğlu and the editor, Greg Shellnutt.

907
Formatted: Justified, Don't add space between paragraphs of the same style, No widow/orphan control

908 References

- 909 Adamia, S.A., Lordkipanidze, M., Zakariadze, G., 1977. Evolution of an active
910 continental margin as exemplified by the Alpine history of the Caucasus.
911 Tectonophysics 40, 183-199.
- 912 Agard, P., Omrani, J., Jolivet, L., Whitechurch, H., Vrielynck, B., Spakman, W., Monié,
913 P., Meyer, B., Wortel, R., 2011. Zagros orogeny: a subduction-dominated process.
914 Geological Magazine 148, 692-725.
- 915 Aktaş, G., Robertson, A.H.F., 1984. The Maden Complex, SE Turkey: evolution of a
916 Neotethyan active margin, in: Dixon, J.E., Robertson, A.H.F. (Eds.), The
917 Geological Evolution of the Eastern Mediterranean. Geological Society of London,
918 Special Publications 17, pp. 375-402.
- 919 Aktaş, G., Robertson, A.H.F., 1990. Tectonic evolution of the Tethys suture zone in SE
920 Turkey: evidence from the petrology and geochemistry of Late Cretaceous and
921 Middle Eocene extrusives, in: Malpas, J., Moores, E.M., Panayiotou, A.,
922 Xenophontos, C. (Eds.), Ophiolites-Oceanic Crustal Analogues. Proceedings of
923 the International Symposium 'Troodos 1987'. Cyprus Geological Survey
924 Department, Nicosia, pp. 311-329.
- 925 Baroz, F., 1979. Etude géologique dans le Pentadaktylos et la Mesaoria (Chypre
926 Septentrionale). Doctor of Science Thesis (Published). Université de Nancy,
927 France (434 pp).
- 928 Baroz, F., 1980. Volcanism and continent-island arc collision in the Pentadaktylos
929 range, Cyprus, in: Panayiotou, A. (Ed.), Ophiolites: Proceedings of the
930 International Ophiolite Symposium. Cyprus Ministry of Agriculture and Natural
931 Resources, Geology Survey Department, Nicosia, Cyprus, pp. 73-85.
- 932 Barrier, E., Vrielynck, B., Brouillet, J.F., Brunet, M.F., 2018. Paleotectonic
933 Reconstruction of the Central Tethyan Realm. Tectono-Sedimentary-Palinspastic
934 Maps from Late Permian to Pliocene. Atlas of 20 maps (scale 1/15000000).
935 CCGM/CGMW, Paris.

936 Beccaluva, L., Macciotta, G., Piccardo, G., Zeda, O., 1989. Clinopyroxene composition
 937 of ophiolite basalts as petrogenetic indicator. *Chemical Geology* 77, 165-182.
 938 Bowman, N., Van Otterloo, J., Cairns, C., Taylor, D., Cas, R., 2019. Complex evolution
 939 of volcanic arcs: The lithofacies and palaeogeography of the Cambrian Staveland
 940 Arc, Delamerian Fold Belt, Western Victoria. *Journal of Volcanology and*
 941 *Geothermal Research* 373, 120-132.
 942 Brune, S., Popov, A.A., Sobolev, S.V., 2012. Modeling suggests that oblique extension
 943 facilitates rifting and continental break-up. *Journal of Geophysical Research: Solid*
 944 *Earth* 117. doi: 10.1029/2011JB008860
 945 Bryan, S., Cook, A., Evans, J., Colls, P., Wells, M., Lawrence, M., Jell, J., Greig, A.,
 946 Leslie, R., 2004. Pumice rafting and faunal dispersion during 2001–2002 in the
 947 Southwest Pacific: record of a dacitic submarine explosive eruption from Tonga.
 948 *Earth and Planetary Science Letters* 227, 135-154.
 949 [Campbell, I.H., Stepanov, A.S., Liang, H.-Y., Allen, C.M., Norman, M.D., Zhang, Y.-Q.,](#)
 950 [Xie, Y.-W., 2014. The origin of shoshonites: new insights from the Tertiary high-](#)
 951 [potassium intrusions of eastern Tibet. *Contributions to Mineralogy and Petrology*](#)
 952 [167. doi: 10.1007/s00410-014-0983-9](#)
 953 Çetinkaplan, M., Pourteau, A., Candan, O., Koralay, O.E., Oberhänsli, R., Okay, A.I.,
 954 Chen, F., Kozlu, H., Şengün, F., 2016. P–T–t evolution of eclogite/blueschist facies
 955 metamorphism in Alanya Massif: time and space relations with HP event in Bitlis
 956 Massif, Turkey. *International Journal of Earth Sciences* 105, 247-281.
 957 Chappell, B.W., White, A.J.R., 1992. I- and S-type granites in the Lachlan Fold Belt.
 958 *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*
 959 83, 1-26.
 960 Chen, G., Robertson, A.H.F., 2019. Provenance and magmatic-tectonic setting of
 961 Campanian-aged volcanoclastic sandstones of the Kannaviou Formation in
 962 western Cyprus: Evidence for a South-Neotethyan continental margin volcanic arc.
 963 *Sedimentary Geology* 388, 114-138.

964 Chen, S.-s., Liu, J.-q., Chen, S.-s., Guo, Z.-f., Sun, C.-q., 2015. Variations in the
 965 geochemical structure of the mantle wedge beneath the northeast Asian marginal
 966 region from pre- to post-opening of the Japan Sea. *Lithos* 224-225, 324-341.

967 Clift, P., 1994. Controls on the Sedimentary and Subsidence History of an Active Plate
 968 Margin: An Example from the Tonga Arc (Southwest Pacific), in: Hawkins, J.,
 969 Parson, L., Allan, J. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific*
 970 *Results. Ocean Drilling Program 135*, College Station, TX, pp. 173–189.

971 Clube, T.M.M., Creer, K.M., Robertson, A.H.F., 1985. Palaeorotation of the Troodos
 972 microplate, Cyprus. *Nature* 317, 522-525.

973 Clube, T.M.M., Robertson, A.H.F., 1986. The palaeorotation of the Troodos microplate,
 974 Cyprus, in the Late Mesozoic-Early Cenozoic plate tectonic framework of the
 975 Eastern Mediterranean. *Surveys in Geophysics* 8, 375-437.

976 Condie, K.C., 2003. Incompatible element ratios in oceanic basalts and komatiites:
 977 tracking deep mantle sources and continental growth rates with time.
 978 *Geochemistry, Geophysics, Geosystems* 4, 1005. doi:10.1029/2002GC000333

979 Condie, K.C., 2005. High field strength element ratios in Archean basalts: a window to
 980 evolving sources of mantle plumes? *Lithos* 79, 491-504.

981 Curray, J.R., 2005. Tectonics and history of the Andaman Sea region. *Journal of Asian*
 982 *Earth Sciences* 25, 187-232.

983 Dercourt, J., Ricou, L.E., Vrielynck, B., 1993. *Atlas of Peri-Tethys Palaeogeographical*
 984 *Maps. CCGM/CGMW, Paris* (268 pp).

985 Dewey, J.F., 1969. Evolution of the Appalachian/Caledonian orogen. *Nature* 222, 124-
 986 129.

987 Ducloz, C., 1972. The geology of the Bellapais-Kythrea area of the Central Kyrenia
 988 Range. *Bulletin of the Geological Survey Department* 6, Nicosia, Cyprus (75 pp).

989 Elmas, A., Yilmaz, Y., 2003. Development of an oblique subduction zone-tectonic
 990 evolution of the Tethys suture zone in southeast Turkey. *International Geology*
 991 *Review* 45, 827-840.

992 Ersoy, E.Y., Palmer, M.R., 2013. Eocene-Quaternary magmatic activity in the Aegean:
 993 implications for mantle metasomatism and magma genesis in an evolving orogeny.
 994 *Lithos* 180, 5-24.

995 Erturk, M.A., Beyarslan, M., Chung, S.L., Lin, T.H., 2018. Eocene magmatism (Maden
 996 Complex) in the Southeast Anatolian Orogenic Belt: Magma genesis and tectonic
 997 implications. *Geoscience Frontiers* 9, 1829-1847.

998 Fitton, J.G., Godard, M., 2004. Origin and evolution of magmas on the Ontong Java
 999 Plateau, in: Fitton, J.G., Mahoney, J.J., Wallace, P.J., Saunders, A.D. (Eds.),
 1000 Origin and Evolution of the Ontong Java Plateau. Geological Society of London,
 1001 Special Publications 229, pp. 151-178.

1002 Fitton, J.G., Saunders, A.D., Larsen, L.M., Hardarson, B.S., Norry, M.J., 1998. Volcanic
 1003 rocks from the southeast Greenland margin at 63°N: composition, petrogenesis
 1004 and mantle sources, in: Saunders, A.D., Larsen, H.C., Wise, S.W. (Eds.),
 1005 Proceedings of the Ocean Drilling Program, Scientific Results. Ocean Drilling
 1006 Program 152, College Station, TX, pp. 331-350.

1007 Floyd, P., Kelling, G., Gökçen, S., Gökçen, N., 1991. Geochemistry and tectonic
 1008 environment of basaltic rocks from the Misis ophiolitic mélange, south Turkey.
 1009 *Chemical Geology* 89, 263-280.

1010 Floyd, P., Kelling, G., Gökçen, S., Gökçen, N., 1992. Arc-related origin of volcanoclastic
 1011 sequences in the Misis Complex, Southern Turkey. *The Journal of Geology* 100,
 1012 221-230.

1013 Garfunkel, Z., 1998. Constrains on the origin and history of the Eastern Mediterranean
 1014 basin. *Tectonophysics* 298, 5-35.

1015 Gilbert, M.F., Robertson, A.H.F., 2013. Field relations, geochemistry and origin of the
 1016 Upper Cretaceous volcanoclastic Kannaviou Formation in western Cyprus:
 1017 evidence of a southerly Neotethyan volcanic arc, in: Robertson, A.H.F., Parlak, O.,
 1018 Ünlügenç, U.C. (Eds.), Geological Development of Anatolia and the Easternmost
 1019 Mediterranean Region. Geological Society of London, Special Publications 372,

1020 pp. 273-298.

1021 Gorton, M.P., Schandl, E.S., 2000. From continents to island arcs: a geochemical index
1022 of tectonic setting for arc-related and within-plate felsic to intermediate volcanic
1023 rocks. *The Canadian Mineralogist* 38, 1065-1073.

1024 Hafkenscheid, E., Wortel, M., Spakman, W., 2006. Subduction history of the Tethyan
1025 region derived from seismic tomography and tectonic reconstructions. *Journal of*
1026 *Geophysical Research: Solid Earth* 111, B08401. doi: 10.1029/2005JB003791

1027 Hakyemez, Y., Turhan, N., Sönmez, I., Sümengen, M., 2000. Kuzey Kıbrıs Türk
1028 Cumhuriyeti'nin Jeolojisi (Geology of the Turkish Republic of Northern Cyprus).
1029 Genel Müdürlüğü Jeoloji Etütleri Diaresi, Maden Tektik ve Arama, Ankara (44 pp).

1030 Harangi, S., Downes, H., Thirlwall, M., Gmeling, K., 2007. Geochemistry, Petrogenesis
1031 and Geodynamic Relationships of Miocene Calc-alkaline Volcanic Rocks in the
1032 Western Carpathian Arc, Eastern Central Europe. *Journal of Petrology* 48, 2261-
1033 2287.

1034 Hartley, M.E., Thordarson, T., 2013. The 1874-1876 volcano-tectonic episode at Askja,
1035 North Iceland: Lateral flow revisited. *Geochemistry, Geophysics, Geosystems* 14,
1036 2286-2309.

1037 Hastie, A.R., Kerr, A.C., Pearce, J.A., Mitchell, S.F., 2007. Classification of Altered
1038 Volcanic Island Arc Rocks using Immobile Trace Elements: Development of the
1039 Th–Co Discrimination Diagram. *Journal of Petrology* 48, 2341-2357.

1040 Hayward, C., 2011. High spatial resolution electron probe microanalysis of tephra and
1041 melt inclusions without beam-induced chemical modification. *The Holocene* 22,
1042 119-125.

1043 Hempton, M.R., 1985. Structure and deformation history of the Bitlis suture near Lake
1044 Hazar, southeastern Turkey. *Geological Society of America Bulletin* 96, 233-243.

1045 Hodgson, E., Morris, A., Anderson, M., Robertson, A., 2010. First palaeomagnetic
1046 results from the Kyrenia Range terrane of northern Cyprus and their implication
1047 for the regional plate tectonic evolution of the eastern Mediterranean, EGU

1048 General Assembly Conference Abstracts, Vienna, Austria, p. 6449.

1049 Hu, Z., Gao, S., 2008. Upper crustal abundances of trace elements: a revision and
1050 update. *Chemical Geology* 253, 205-221.

1051 Huang, K., Malpas, J., Xenophontos, C., 2007. Geological studies of igneous rocks
1052 and their relationships along the Kyrenia Range, in: Moumani, K., Shawabkeh, K.,
1053 Al-Malabeh, A., Abdelghafoor, M. (Eds.), 6th International Congress of Eastern
1054 Mediterranean Geology, Amman, Jordan, p. 53.

1055 Karaoğlu, F., Parlak, O., Hejl, E., Neubauer, F., Kloetzli, U., 2016. The temporal
1056 evolution of the active margin along the Southeast Anatolian Orogenic Belt (SE
1057 Turkey): Evidence from U–Pb, Ar–Ar and fission track chronology. *Gondwana
1058 Research* 33, 190-208.

1059 Karaoğlu, F., Parlak, O., Robertson, A., Thöni, M., Klötzli, U., Koller, F., Okay, A.İ.,
1060 2013. Evidence of Eocene high-temperature/high-pressure metamorphism of
1061 ophiolitic rocks and granitoid intrusion related to Neotethyan subduction
1062 processes (Doğanşehir area, SE Anatolia), in: Robertson, A.H.F., Parlak, O.,
1063 Ünlügenç, U.C. (Eds.), *Geological Development of Anatolia and the Easternmost
1064 Mediterranean Region*. Geological Society of London, Special Publications 372,
1065 pp. 249-272.

1066 Keller, R.A., Fisk, M.R., Smellie, J.L., Strelin, J.A., Lawver, L.A., 2002. Geochemistry
1067 of back arc basin volcanism in Bransfield Strait, Antarctica: Subducted
1068 contributions and along-axis variations. *Journal of Geophysical Research: Solid
1069 Earth* 107. doi: 10.1029/2001JB000444

1070 Kelly, N., Hinton, R., Harley, S., Appleby, S., 2008. New SIMS U–Pb zircon ages from
1071 the Langavat Belt, South Harris, NW Scotland: implications for the Lewisian
1072 terrane model. *Journal of the Geological Society* 165, 967-981.

1073 Kempton, P.D., Downes, H., Lustrino, M., 2018. Pb and Hf isotope evidence for mantle
1074 enrichment processes and melt interactions in the lower crust and lithospheric

1075 mantle in Miocene orogenic volcanic rocks from Monte Arcuentu (Sardinia, Italy).
1076 Geosphere 14, 926-950.

1077 [Kuşcu, İ., Tosdal, R.M., Gencalioglu-Kuşcu, G., Friedman, R., Ullrich, T.D., 2013. Late](#)
1078 [Cretaceous to Middle Eocene Magmatism and Metallogeny of a Portion of the](#)
1079 [Southeastern Anatolian Orogenic Belt, East-Central Turkey. Economic Geology](#)
1080 [108, 641-666.](#)

1081 Le Pichon, X., 1982. Land-locked oceanic basins and continental collision: the Eastern
1082 Mediterranean as a case example, in: Hsü, K.J. (Ed.), Mountain building
1083 processes. Academic Press, New York, pp. 201-211.

1084 Li, C.-F., Li, X.-H., Li, Q.-L., Guo, J.-H., Li, X.-H., Yang, Y.-H., 2012. Rapid and precise
1085 determination of Sr and Nd isotopic ratios in geological samples from the same
1086 filament loading by thermal ionization mass spectrometry employing a single-step
1087 separation scheme. Analytica Chimica Acta 727, 54-60.

1088 Ludwig, K.R., 2012. Users manual for Isoplot 3.75. A geochronological toolkit for
1089 Microsoft Excel. Berkeley Geochronology Centre, Special Publication No. 5,
1090 Berkeley.

1091 Maffione, M., van Hinsbergen, D.J., de Gelder, G.I., van der Goes, F.C., Morris, A.,
1092 2017. Kinematics of Late Cretaceous subduction initiation in the Neo-Tethys
1093 Ocean reconstructed from ophiolites of Turkey, Cyprus, and Syria. Journal of
1094 Geophysical Research: Solid Earth 122, 3953-3976.

1095 McCay, G.A., Robertson, A.H.F., 2013. Upper Miocene–Pleistocene deformation of the
1096 Girne (Kyrenia) Range and Dar Dere (Ovgos) lineaments, northern Cyprus: role
1097 in collision and tectonic escape in the easternmost Mediterranean region, in:
1098 Robertson, A.H.F., Parlak, O., Ünlügenç, U.C. (Eds.), Geological Development of
1099 Anatolia and the Easternmost Mediterranean Region. Geological Society of
1100 London, Special Publications 372, pp. 421-445.

1101 McCay, G.A., Robertson, A.H.F., Kroon, D., Raffi, I., Ellam, R.M., Necdet, M., 2013.

1102 Stratigraphy of Cretaceous to Lower Pliocene sediments in the northern part of
 1103 Cyprus based on comparative $^{87}\text{Sr}/^{86}\text{Sr}$ isotopic, nannofossil and planktonic
 1104 foraminiferal dating. *Geological Magazine* 150, 333-359.

1105 McPhee, P.J., van Hinsbergen, D.J., 2019. Tectonic reconstruction of Cyprus reveals
 1106 Late Miocene continental collision of Africa and Anatolia. *Gondwana Research* 68,
 1107 158-173.

1108 Moix, P., Beccaleto, L., Kozur, H.W., Hochard, C., Rosselet, F., Stampfli, G.M., 2008.
 1109 A new classification of the Turkish terranes and sutures and its implication for the
 1110 paleotectonic history of the region. *Tectonophysics* 451, 7-39.

1111 Moore, T.A., 1960. The geology and mineral resources of the Astromeritis-Kormakiti
 1112 area. Geological Survey Department, Memoir 6, Nicosia, Cyprus (96 pp).

1113 Morris, A., Anderson, M.W., Inwood, J., Robertson, A.H., 2006. Palaeomagnetic
 1114 insights into the evolution of Neotethyan oceanic crust in the eastern
 1115 Mediterranean, in: Robertson, A.H.F., Mountrakis, D. (Eds.), *Tectonic*
 1116 *Development of the Eastern Mediterranean Region*. Geological Society of London,
 1117 Special Publications 260, pp. 351-372.

1118 Morris, A., Robertson, A.H.F., Anderson, M.W., Hodgson, E., 2015. Did the Kyrenia
 1119 Range of northern Cyprus rotate with the Troodos–Hatay microplate during the
 1120 tectonic evolution of the eastern Mediterranean? *International Journal of Earth*
 1121 *Sciences* 105, 399-415.

1122 Nisbet, E.G. Pearce, J.A., 1977. Clinopyroxene composition in mafic lavas from
 1123 different tectonic settings. *Contributions to Mineralogy and Petrology* 63, 149-160.

1124 Nurlu, N., Parlak, O., Robertson, A.H.F., von Quadt, A., 2016. Implications of Late
 1125 Cretaceous U–Pb zircon ages of granitic intrusions cutting ophiolitic and
 1126 volcanogenic rocks for the assembly of the Tauride allochthon in SE Anatolia
 1127 (Helete area, Kahramanmaraş Region, SE Turkey). *International Journal of Earth*
 1128 *Sciences* 105, 283-314.

1129 [Oberhänsli, R., Candan, O., Bousquet, R., Rimmele, G., Okay, A., Goff, J., 2010. Alpine](#)

1130 [high pressure evolution of the eastern Bitlis complex, SE Turkey, in: Sosson, M.,](#)
 1131 [Kaymakci, N., Stephenson, R.A., Bergerat, F., Starostenko, V. \(Eds.\),](#)
 1132 [Sedimentary Basin Tectonics from the Black Sea and Caucasus to the Arabian](#)
 1133 [Platform. Geological Society of London, Special Publications 340, pp. 461-483.](#)
 1134 [Oberhänsli, R., Koralay, E., Candan, O., Pourteau, A., Bousquet, R., 2013. Late](#)
 1135 [Cretaceous eclogitic high-pressure relics in the Bitlis Massif. *Geodinamica Acta*](#)
 1136 [26, 175-190.](#)
 1137 Parlak, O., 2006. Geodynamic significance of granitoid magmatism in the southeast
 1138 Anatolian orogen: geochemical and geochronological evidence from Göksun–Afşin
 1139 (Kahramanmaraş, Turkey) region. *International Journal of Earth Sciences* 95, 609-
 1140 627.
 1141 Pearce, J.A., 1975. Basalt geochemistry used to investigate past tectonic
 1142 environments on Cyprus. *Tectonophysics* 25, 41-67.
 1143 Pearce, J.A., 1983. Role of the sub-continental lithosphere in magma genesis at active
 1144 continental margins, in: Hawkersworth, C.J., Norry, M.J. (Eds.), *Continental*
 1145 *Basalts and Mantle Xenoliths*. Shiva, Cheshire, UK, pp. 230-249.
 1146 Pearce, J.A., 1996. A user's guide to basalt discrimination diagrams, in: Wyman, D.A.
 1147 (Ed.), *Trace Element Geochemistry of Volcanic Rocks: Applications for Massive*
 1148 *Sulphide Exploration*. Geological Association of Canada, Short Course Notes 12,
 1149 pp. 79-113.
 1150 Pearce, J.A., Cann, J., 1973. Tectonic setting of basic volcanic rocks determined using
 1151 trace element analyses. *Earth and Planetary Science Letters* 19, 290-300.
 1152 Pearce, J.A., Harris, N.B., Tindle, A.G., 1984. Trace element discrimination diagrams
 1153 for the tectonic interpretation of granitic rocks. *Journal of Petrology* 25, 956-983.
 1154 Pearce, J.A., Norry, M.J., 1979. Petrogenetic implications of Ti, Zr, Y, and Nb variations
 1155 in volcanic rocks. *Contributions to Mineralogy and Petrology* 69, 33-47.
 1156 Peccerillo, A., 2017. *Cenozoic Volcanism in the Tyrrhenian Sea Region*. Springer
 1157 International Publishing, Cham (399 pp).

1158 Perinçek, D., Kozlu, H., 1984. Stratigraphy and structural relations of the units in the
1159 Afşin-Elbistan-Doğanşehir region (Eastern Taurus), in: Tekeli, O., Göncüoğlu, M.C.
1160 (Eds.), *Geology of the Taurus Belt: Proceedings of the International Symposium*.
1161 Maden Tetkik ve Arama Enstitüsü, Ankara, pp. 181-198.

1162 Perinçek, D., Özkaya, İ., 1981. Tectonic evolution of the northern margin of Arabian
1163 plate. *Bulletin of Institute of Earth Science, Hacettepe University* 8, 91-101.

1164 Rice, S.P., Robertson, A.H.F., Ustaömer, T., 2006. Late Cretaceous-Early Cenozoic
1165 tectonic evolution of the Eurasian active margin in the Central and Eastern
1166 Pontides, northern Turkey, in: Robertson, A.H.F., Mountrakis, D. (Eds.), *Tectonic
1167 Development of the Eastern Mediterranean Region*. Geological Society of London,
1168 Special Publications 260, pp. 413-445.

1169 Riecker, R.E., 1979. Rio Grande Rift: Tectonics and Magmatism. *American
1170 Geophysical Union, Washington, DC* (438 pp).

1171 Rızaoğlu, T., Parlak, O., Höck, V., Koller, F., Hames, W.E., Billor, Z., 2009. Andean-
1172 type active margin formation in the eastern Taurides: Geochemical and
1173 geochronological evidence from the Baskil granitoid (Elazığ, SE Turkey).
1174 *Tectonophysics* 473, 188-207.

1175 Robertson, A.H.F., 1977. The Kannaviou Formation, Cyprus: volcanoclastic
1176 sedimentation of a probable Late Cretaceous volcanic arc. *Journal of the
1177 Geological Society* 134, 269-292.

1178 Robertson, A.H.F., Dixon, J.E., 1984. Introduction: aspects of the geological evolution
1179 of the Eastern Mediterranean, in: Dixon, J.E., Robertson, A.H.F. (Eds.), *The
1180 Geological Evolution of the Eastern Mediterranean*. Geological Society of London,
1181 Special Publications 17, pp. 1-74.

1182 Robertson, A.H.F., Kinnaird, T.C., 2016. Structural development of the central Kyrenia
1183 Range (north Cyprus) in its regional setting in the eastern Mediterranean region.
1184 *International Journal of Earth Sciences* 105, 417-437.

1185 Robertson, A.H.F., Parlak, O., 2020. Late Cretaceous-Palaeocene subduction-

1186 collision-exhumation of a microcontinent along the northern, active margin of
 1187 South Neotethys: evidence from the Alanya Massif and the adjacent Antalya
 1188 Complex (S Turkey). *Journal of Asian Earth Sciences*, 104467. doi:
 1189 10.1016/j.jseaes.2020.104467
 1190 Robertson, A.H.F., Parlak, O., Kinnaird, T.C., Taslı, K., Dumitrica, P., 2020. Cambrian-
 1191 Eocene pre-rift, pulsed rift, passive margin and emplacement processes along the
 1192 northern margin of the Southern Neotethys: Evidence from the Antalya Complex
 1193 in the Alanya Window (S Turkey). *Journal of Asian Earth Sciences*: X 3, 100026.
 1194 doi: 10.1016/j.jaesx.2020.100026
 1195 Robertson, A.H.F., Parlak, O., Rızaoğlu, T., Ünlügenç, Ü., İnan, N., Taslı, K., Ustaömer,
 1196 T., 2007. Tectonic evolution of the South Tethyan ocean: evidence from the
 1197 Eastern Taurus Mountains (Elazığ region, SE Turkey), in: Ries, A.C., Butler,
 1198 R.W.H., Graham, R.H. (Eds.), *Deformation of the Continental Crust: The Legacy*
 1199 *of Mike Coward*. Geological Society of London, Special Publications 272, pp. 231-
 1200 270.
 1201 Robertson, A.H.F., Parlak, O., Ustaömer, T., 2012a. Overview of the Palaeozoic-
 1202 Neogene evolution of Neotethys in the Eastern Mediterranean region (southern
 1203 Turkey, Cyprus, Syria). *Petroleum Geoscience* 18, 381-404.
 1204 Robertson, A.H.F., Taslı, K., İnan, N., 2012b. Evidence from the Kyrenia Range, Cyprus,
 1205 of the northerly active margin of the Southern Neotethys during Late Cretaceous–
 1206 Early Cenozoic time. *Geological Magazine* 149, 264-290.
 1207 Robertson, A.H.F., Ünlügenç, Ü.C., İnan, N., Taslı, K., 2004. The Misis-Andırın
 1208 Complex: a Mid-Tertiary melange related to late-stage subduction of the Southern
 1209 Neotethys in S Turkey. *Journal of Asian Earth Sciences* 22, 413-453.
 1210 Robertson, A.H.F., Ustaömer, T., Parlak, O., Ünlügenç, Ü.C., Taşlı, K., İnan, N., 2006.
 1211 The Berit transect of the Tauride thrust belt, S Turkey: Late Cretaceous-Early
 1212 Cenozoic accretionary/collisional processes related to closure of the Southern
 1213 Neotethys. *Journal of Asian Earth Sciences* 27, 108-145.

1214 Robertson, A.H.F., Woodcock, N.H., 1979. Mamonia Complex, southwest Cyprus:
 1215 Evolution and emplacement of a Mesozoic continental margin. Geological Society
 1216 of America Bulletin 90, 651-665.

1217 Robertson, A.H.F., Woodcock, N.H., 1986. The role of the Kyrenia Range Lineament,
 1218 Cyprus, in the geological evolution of the eastern Mediterranean area.
 1219 Philosophical Transactions of the Royal Society of London. Series A,
 1220 Mathematical and Physical Sciences 317, 141-177.

1221 Rollinson, H.R., 1993. Using Geochemical Data: Evaluation, Presentation,
 1222 Interpretation. Routledge, London (384 pp).

1223 Rubatto, D., 2002. Zircon trace element geochemistry: partitioning with garnet and the
 1224 link between U–Pb ages and metamorphism. Chemical Geology 184, 123-138.

1225 Rudnick, R.L., Gao, S., 2003. Composition of the continental crust, in: Holland, H.D.,
 1226 Turekian, K.K. (Eds.), Treatise on geochemistry. Elsevier-Pergamon 3, Oxford, pp.
 1227 1-64.

1228 [Sar, A., Ertürk, M.A., Rizeli, M.E., 2019. Genesis of Late Cretaceous intra-oceanic arc](#)
 1229 [intrusions in the Pertek area of Tunceli Province, eastern Turkey, and implications](#)
 1230 [for the geodynamic evolution of the southern Neo-Tethys: Results of zircon U–Pb](#)
 1231 [geochronology and geochemical and Sr–Nd isotopic analyses. Lithos 350-351,](#)
 1232 [105263. doi: 10.1016/j.lithos.2019.105263](#)

1233 Saunders, A.D., Fornari, D.J., Joron, J., Tarney, J., Treuil, M., 1982. Geochemistry of
 1234 basic igneous rocks, Gulf of California, Deep Sea Drilling Project Leg 64, in:
 1235 Curray, J.R., Moore, D.G., Kelts, K. (Eds.), Initial Reports of the Deep Sea Drilling
 1236 Project. Ocean Drilling Program 64, College Station, TX, pp. 595-642.

1237 Saunders, A.D., Tarney, J., Stern, C.R., Dalziel, I.W., 1979. Geochemistry of Mesozoic
 1238 marginal basin floor igneous rocks from southern Chile. Geological Society of
 1239 America Bulletin 90, 237-258.

1240 Şengör, A.M.C., Natal'in, B.A., 1996. Turkic-type orogeny and its role in the making of
 1241 the continental crust. Annual Review of Earth and Planetary Sciences 24, 263-

1242 337.

1243 Şengör, A.M.C., Yılmaz, Y., 1981. Tethyan evolution of Turkey: a plate tectonic
1244 approach. *Tectonophysics* 75, 181-241.

1245 Şengör, A.M.C., Yılmaz, Y., Sungurlu, O., 1984. Tectonics of the Mediterranean
1246 Cimmerides: nature and evolution of the western termination of Palaeo-Tethys, in:
1247 Dixon, J.E., Robertson, A.H.F. (Eds.), *The Geological Evolution of the Eastern*
1248 *Mediterranean*. Geological Society of London, Special Publications 17, pp. 77-112.

1249 Shervais, J.W., 1982. Ti-V plots and the petrogenesis of modern and ophiolitic lavas.
1250 *Earth and Planetary Science Letters* 59, 101-118.

1251 Stampfli, G.M., Borel, G., 2002. A plate tectonic model for the Paleozoic and Mesozoic
1252 constrained by dynamic plate boundaries and restored synthetic oceanic
1253 isochrons. *Earth and Planetary Science Letters* 196, 17-33.

1254 Sun, S.-s., McDonough, W.F., 1989. Chemical and isotopic systematics of oceanic
1255 basalts: implications for mantle composition and processes, in: Saunders, A.D.,
1256 Norry, M.J. (Eds.), *Magmatism in the Ocean Basins*. Geological Society of London,
1257 Special Publications 42, pp. 313-345.

1258 Tamaki, K., Pisciotto, K., Allan, J., Alexandrovich, J.M., Barnes, D.A., Boggs, S.,
1259 Brumsack, H., Brunner, C.A., Cramp, A., Jolivet, L., Kawka, O.E., Koizumi, I.,
1260 Kuramoto, S.i., Langseth, M.G., McEvoy, J., Meredith, J.A., Mertz, K.A., Murray,
1261 R.W., Nobes, D.C., Rahman, A., Schaar, R., Stewart, K.P., Tada, R., Thy, P.,
1262 Vigliotti, L., White, L.D., Wipperfurth, J., Yamashita, S., 1990. *Proceedings of the*
1263 *Ocean Drilling Program, Initial Reports vol. 127*. Ocean Drilling Program.

1264 Tatsumi, Y., Kogiso, T., 2003. The subduction factory: its role in the evolution of the
1265 Earth's crust and mantle, in: Larter, R.D., Leat, P.T. (Eds.), *Intra-Oceanic*
1266 *Subduction Systems: Tectonic and Magmatic Processes*. Geological Society of
1267 London, Special Publications 219, pp. 55-80.

1268 Thirlwall, M., 1991. Long-term reproducibility of multicollector Sr and Nd isotope ratio
1269 analysis. *Chemical Geology* 94, 85-104.

1270 Trehu, A., Asudeh, I., Brocher, T., Luetgert, J., Mooney, W., Nabelek, J., Nakamura, Y.,
1271 1994. Crustal architecture of the Cascadia forearc. *Science* 266, 237-243.

1272 Ural, M., Arslan, M., Göncüoğlu, M., Tekin, U., Kürüm, S., 2015. Late Cretaceous arc
1273 and back-arc formation within the southern Neotethys: Whole-rock, trace element
1274 and Sr-Nd-Pb isotopic data from basaltic rocks of the Yüksekova Complex
1275 (Malatya- Elazığ, SE Turkey). *Ofioliti* 40, 52-72.

1276 Ustaömer, P.A., Ustaömer, T., Robertson, A.H.F., 2012. Ion probe U-Pb dating of the
1277 Central Sakarya basement: a peri-Gondwana terrane intruded by late Lower
1278 Carboniferous subduction/collision-related granitic rocks. *Turkish Journal of Earth
1279 Sciences* 21, 905-932.

1280 Ustaömer, T., Robertson, A.H.F., 1997. Tectonic-sedimentary evolution of the North-
1281 Tethyan active margin in the Central Pontides of Northern Turkey, in: Robinson,
1282 A.G. (Ed.), *Regional and Petroleum Geology of the Black Sea and Surrounding
1283 Region*. AAPG Memoirs 68, pp. 255-290.

1284 van Hinsbergen, D.J., Torsvik, T.H., Schmid, S.M., Mañenco, L.C., Maffione, M., Vissers,
1285 R.L., Gürer, D., Spakman, W., 2020. Orogenic architecture of the Mediterranean
1286 region and kinematic reconstruction of its tectonic evolution since the Triassic.
1287 *Gondwana Research* 81, 79-229.

1288 Wang, Y., Fan, W., Zhang, Y., Guo, F., Zhang, H., Peng, T., 2004. Geochemical,
1289 $^{40}\text{Ar}/^{39}\text{Ar}$ geochronological and Sr-Nd isotopic constraints on the origin of
1290 Paleoproterozoic mafic dikes from the southern Taihang Mountains and
1291 implications for the ca. 1800Ma event of the North China Craton. *Precambrian
1292 Research* 135, 55-77.

1293 Weaver, S.D., Saunders, A.D., Pankhurst, R.J., Tarney, J., 1979. A geochemical study
1294 of magmatism associated with the initial stages of back-arc spreading.
1295 *Contributions to Mineralogy and Petrology* 68, 151-169.

1296 Weis, D., Kieffer, B., Hanano, D., Nobre Silva, I., Barling, J., Pretorius, W., Maerschalk,
1297 C., Mattielli, N., 2007. Hf isotope compositions of US Geological Survey reference

1298 materials. *Geochemistry, Geophysics, Geosystems* 8, 57-77.

1299 Weis, D., Kieffer, B., Maerschalk, C., Barling, J., De Jong, J., Williams, G.A., Hanano,
1300 D., Pretorius, W., Mattioli, N., Scoates, J.S., 2006. High-precision isotopic
1301 characterization of USGS reference materials by TIMS and MC-ICP-MS.
1302 *Geochemistry, Geophysics, Geosystems* 7, Q08006. doi:
1303 10.1029/2006GC001283

1304 Whalen, J.B., Currie, K.L., Chappell, B.W., 1987. A-type granites: geochemical
1305 characteristics, discrimination and petrogenesis. *Contributions to Mineralogy and
1306 Petrology* 95, 407-419.

1307 Winchester, J., Floyd, P., 1977. Geochemical discrimination of different magma series
1308 and their differentiation products using immobile elements. *Chemical Geology* 20,
1309 325-343.

1310 Yazgan, E., Chessex, R., 1991. Geology and Tectonic Evolution of the Southeastern
1311 Taurides in the Region of Malatya. *Turkish Association of Petroleum Geologists* 3,
1312 1-42.

1313 Yiğitbaş, E., Yılmaz, Y., 1996. Post-Late Cretaceous strike-slip tectonics and its
1314 implications for the Southeast Anatolian orogen, Turkey. *International Geology
1315 Review* 38, 818-831.

1316 Yıldırım, E., 2015. Geochemistry, petrography and tectonic significance of the ophiolitic
1317 rocks, felsic intrusions and Eocene volcanic rocks of an imbrication zone (Helete
1318 area, Southeast Turkey). *Journal of African Earth Sciences* 107, 89-107.

1319 Yılmaz, Y., 1993. New evidence and model on the evolution of the southeast Anatolian
1320 orogen. *Geological Society of America Bulletin* 105, 251-271.

1321 [Yılmaz, Y., Yiğitbaş, E., Genç, Ş.C., 1993. Ophiolitic and metamorphic assemblages of
1322 southeast Anatolia and their significance in the geological evolution of the
1323 orogenic belt. *Tectonics* 12, 1280-1297.](#)

1324 Zhang, Y., Wang, Y., Geng, H., Zhang, Y., Fan, W., Zhong, H., 2013. Early
1325 Neoproterozoic (~850Ma) back-arc basin in the Central Jiangnan Orogen

1326 (Eastern South China): Geochronological and petrogenetic constraints from meta-
1327 basalts. *Precambrian Research* 231, 325-342.
1328 Zindler, A., Hart, S., 1986. Chemical geodynamics. *Annual Review of Earth and*
1329 *Planetary Sciences* 14, 493-571.
1330

1331 Listing of Figures

1332 Fig. 1 Outline tectonic map of the Eastern Mediterranean region, including the main
1333 volcanic units mentioned in the text. The tectonic framework is modified after
1334 Robertson et al. (2012a). Note the location of the Kyrenia Range in northern Cyprus
1335 (red box). Jurassic, Cretaceous and Eocene igneous rocks are shown in blue, red and
1336 yellow, respectively (see the text for literature sources).

1337 Fig. 2 Successions shown in two main thrust sheets exposed in the Kyrenia Range,
1338 northern Cyprus. (a) Simplified log of the succession in the largest thrust sheet
1339 including the latest Cretaceous-Paleogene volcanics (Melounda Formation); (b)
1340 Restored log of the small, dismembered frontal thrust sheets that expose the Late
1341 Cretaceous felsic volcanics (Fourkovouno Formation) (data from Baroz, 1979;
1342 Robertson et al., 2012b and this study).

1343 Fig. 3 (a) Outline geological map of the Kyrenia Range (modified from Robertson et al.,
1344 2012b). Locations from which basalts were collected are numbered on the map, with
1345 small yellow boxes. Karpas Peninsula: 1-Balalan (Platanissos); Eastern Range: 2-
1346 Ağıllar (Mandres), 3-Çınarlı (Platani), 4-Mallıdağ (Melounda); Central Range: 5-Tirmen
1347 (Trypimeni), 6-Ergenekon (Agios Khariton), 7-Değirmenlik (Kythrea), 8-Arapköy
1348 (Klepini), 9-Beylerbeyi (Bellapais), 10-Boğaz (Bogaz), 11-Pınarbaşı (Krini), 12-İncesu
1349 (Motides), 13-Alevkaya Tepe (Kiparisso Vouno); Western Range, 14-Karşıyaka
1350 (Vasileia), 15-Geçitköy (Panagra); (b) Geological map of the western Kyrenia Range,
1351 northern Cyprus made during this study (based on mapping by Baroz, 1979). Locations
1352 of cross sections (AA', BB' and CC') are indicated. Note that the two contrasting, mainly
1353 felsic and mainly basaltic volcanic units are separated by a thrust fault (Robertson et
1354 al., 2012b; this study; see Fig. 2).

1355 Fig. 4 (a) A-A' cross section of Geçitköy that is dominated by a south-verging,
1356 recumbent anticline (Late Miocene-earliest Pliocene); (b) B-B' cross-section of
1357 Selvilitepe showing southward imbrication of basalt/pelagic chalk and felsic

1358 volcanogenic rocks; (c) C-C' cross-section showing imbricated felsic volcanogenic
1359 rocks.

1360 Fig. 5 Measured stratigraphic logs of partial successions in the felsic volcanics and
1361 structurally associated units. The structural lower unit (logs a-e) is relatively intact,
1362 whereas the higher unit (logs f-j) is characterised by the thrust intercalations with the
1363 Trypa Group. Fig. 6 (a) Zr/Ti vs. Nb/Y diagram (after Pearce, 1996) and (b) Th vs. Co
1364 diagram (after Hastie et al., 2007) for the Kyrenia Range felsic rocks; (c) Primitive
1365 mantle-normalised multi-element spider diagram and (d) chondrite-normalised rare
1366 earth element (REE) patterns of the felsic volcanics with reference data for upper
1367 continental crust (UCC) and lower continental crust (LCC). Normalising values of
1368 primitive mantle and chondrite after Sun and McDonough (1989), upper continental
1369 crust (UCC) and lower continental crust (LCC) data are from Rudnick and Gao (2003),
1370 and Hu and Gao (2008). The Chile volcanic arc granite data are after Pearce et al.
1371 (1984).

1372 Fig. 7 (a) Zr/Ti vs. Nb/Y diagram (after Pearce, 1996) and (b) Th vs. Co diagram (after
1373 Hastie et al., 2007) for the basaltic volcanics. Red symbols: Karpas Peninsula and
1374 eastern range; green symbols: central range; blue symbols: western range. Paleogene
1375 basalt samples are indicated by thick magenta-outlined symbols.

1376 Fig. 8 Mid-ocean ridge basalt (MORB)-normalised multi-element spider diagram for
1377 basalts from the Karpas Peninsula and the eastern (a), central (b) and western (c)
1378 Kyrenia Range. MORB-normalised data are from Pearce et al. (1983); (d) Chondrite-
1379 normalised REE patterns of selected basalts. Chondrite, OIB and E-MORB data are
1380 from Sun and McDonough (1989). Paleogene basalt samples are indicated by solid
1381 black lines and magenta-outlined symbols.

1382 Fig. 9 (a) Cathodoluminescence images of zircon grains separated from the samples
1383 of Geçitköy. Red circles mark inconsistent analyses that were omitted from the

1384 weighted mean $^{206}\text{Pb}/^{238}\text{U}$ age calculation. Locations of the ion probe analysis spots
1385 and the corresponding ages ($^{206}\text{Pb}/^{238}\text{U} \pm 1\sigma$) are indicated. Scale bar = 20 μm ; (b-d)
1386 Wetherill Concordia diagram for the zircons analyses from sample 14-18 (b), 14-19 (c)
1387 and 14-20 (d).

1388 Fig. 10 (a-b) Plots of P_2O_5 vs. SiO_2 and Th, respectively, show that all of the samples
1389 follow the I-type granite trend; (c) Zr vs. $10000 \times \text{Ga}/\text{Al}$, and (d) Nb vs. $10000 \times \text{Ga}/\text{Al}$
1390 discrimination diagrams of Whalen et al. (1987), showing the I-, S (sedimentary)- and
1391 M (depleted mantle source)-type nature of the Late Cretaceous felsic volcanics.

1392 Fig. 11 (a) $\epsilon\text{Nd}(t)$ vs. $\epsilon\text{Sr}(t)$ diagram for the basaltic rocks analysed. The samples are
1393 mainly Late Cretaceous ($n=4$); Paleogene basalt samples ($n=2$) are indicated by
1394 magenta-outlined symbols. Comparative compositions are from Zindler and Hart
1395 (1986); (b) $\epsilon\text{Hf}(t)$ vs. $\epsilon\text{Nd}(t)$ plot showing rocks for Kyrenia Range, northern Cyprus
1396 relative to other volcanic rocks of the western Mediterranean (after Kempton et al.,
1397 2018).

1398 Fig. 12 (a) Ba/Nb versus La/Nb diagram (after Zhang et al., 2013); (b) Nb/Y versus
1399 Zr/Y diagram (after Condie, 2005), arrows in (b) indicate the effects of batch melting
1400 (F) and the probable effect of fluids derived from subduction (SUB); (c) Zr/Nb vs. Nb/Th
1401 diagram (modified after Condie, 2003). Abbreviations: PM, primitive mantle; DM,
1402 shallow depleted mantle; HIMU, high $m\mu$ (U/Pb) source; EMI and EMII, enriched
1403 mantle sources; ARC, arc-related basalt; N-MORB, normal ocean ridge basalt; OIB,
1404 oceanic island basalt; DEP, deep depleted mantle; EN, enriched component; REC,
1405 recycled component; UCC, upper upper continental crust; LCC, lower continental crust;
1406 BM, batch melting trajectory with percent melting noted. Numbers on mixing lines are
1407 percents.

1408 Fig. 13 (a) Nb vs. Y diagram (Pearce et al., 1984) and (b) Th/Yb vs. Ta/Yb (Gorton and
1409 Schandl, 2000) for the felsic volcanics; (c) Zr/Y vs. Zr diagram (after Pearce and Norry,
1410 1979) and (d) V vs. Ti discrimination diagram (after Shervais, 1982) for the basaltic

1411 rocks. Abbreviations: WPG, within-plate granite; VAG, volcanic arc granite; syn-COLG,
1412 syn-collisional granite; ORG, ocean ridge granite; WPB, within-plate basalt; MORB,
1413 mid-ocean ridge basalt; BABB, back-arc basin basalt; ARC, arc-related basalt; OFB,
1414 oceanic floor basalt.

1415 Fig. 14 (a) Th/Nb vs. La/Nb and (b) La/Nb versus Y diagrams (after Floyd et al., 1991).
1416 Abbreviations: IAT, island arc tholeiite; BABB, back-arc basin basalt; OFB, oceanic
1417 flood basalt; FAPB, fore-arc platform basalt; T/E-MORB, T-type/enriched-mid-ocean
1418 ridge basalt.

1419 Fig. 15 Alternative tectonic models of Tethys in the Eastern Mediterranean region. (a-
1420 c) Northward subduction with continental fragments rifted from Gondwana (Robertson
1421 et al., 2012a); (d-f) Northward subduction and marginal basin formation (Barrier et al.,
1422 2018); (g-h) Genesis of Late Cretaceous supra-subduction ophiolites at a single
1423 subduction zone to the NE followed by roll-back of segments including the Southern
1424 Neotethys (Maffione et al., 2017; McPhee and van Hinsbergen, 2019; van Hinsbergen
1425 et al., 2020). Based on discussion (see text), model **a-1** is generally favoured.

1426

1427 Listing of Supplementary Figure

1428 Supplementary Figure S1 Field photographs of the felsic volcanics. (a) Exposure in
1429 Geçitköy. Samples 14-18 and 14-19 are from this location; (b) Irregular-shaped
1430 doleritic intrusion (hammer for scale); (c) Localised occurrence of greyish to greenish,
1431 rhyolitic debris-flow unit, north of Geçitköy; (d) Greenish rhyolitic debris-flow unit with
1432 angular rhyolitic clasts (pen for scale).

1433 Supplementary Figure S2 Field photographs of the basaltic volcanics. (a) Pillow lavas
1434 and interstitial pink pelagic carbonate; roadcut c. 200 m north of Balalan; (b) basalt
1435 interbedded with pelagic carbonate; thrust sheet above Miocene siliciclastic sediment;
1436 c. 800 m northwest of Değirmenlik; (c) Basaltic volcanics intercalated with pelagic
1437 carbonates and small slices of meta-platform carbonates of the Trypa Group along the
1438 southern flank of the range, 800 m north of Boğaz; (d) Basalt-pelagic carbonate
1439 intercalation, south of Karşıyaka.

1440 Supplementary Figure S3 Photomicrographs of felsic volcanics. (a) Subhedral
1441 sanidine (Sa), plagioclase (Pl) and biotite (Bt), in a groundmass of devitrified glassy
1442 volcanic shards (cross-polarised light); (b) felsic glass groundmass (Vg) with
1443 microcrystalline quartz, irregular-shaped quartz (Q) and feldspar (Pl) (cross-polarised
1444 light). Sample number is indicated in the bottom-left corner.

1445 Supplementary Figure S4 Photomicrographs of basalts. (a) Intersertal basalt with
1446 euhedral granular augite (Px), elongate plagioclase laths (Pl) and opaque grains (Op)
1447 (cross-polarised light); (b) Basalt with intersertal texture, in which randomly-oriented
1448 plagioclase generally enclose pyroxenes (plane-polarised light); (c) Porphyritic basalt
1449 with euhedral plagioclase and strongly altered olivine phenocrysts (cross-polarised
1450 light); (d) Ophitic basalt with plagioclase phenocrysts, augite occurs as an inclusion
1451 within individual plagioclase crystals (cross-polarised light). Sample number is
1452 indicated in the bottom-left corner. Samples no. 21, 14-51 and 14-69 are from the
1453 Melounda Formation, whereas sample no. 19-51 is from the Ayios Nikolaos Formation.

1454 Supplementary Figure S5 Tectonic discrimination plots. (a) TiO_2 - MnO - Na_2O ternary
1455 diagram (after Nisbet and Pearce, 1977) for clinopyroxenes in basalt; (b) TiO_2 - SiO_2 -
1456 Na_2O diagram (after Beccaluva et al., 1989) for clinopyroxenes in basalt. Abbreviations:
1457 VAB, volcanic arc basalt; OFB, ocean-floor basalt; WPA, within-plate alkali basalt;
1458 WPT, within-plate tholeiitic basalt; BON, boninite; IAT, island-arc tholeiite; WPB,
1459 within-plate basalt.

1460 Supplementary Figure S6 (a) Chondrite-normalised REE comparison patterns of the
1461 Kyrenia Range felsic rocks and the Helete granites in SE Turkey (data after Nurlu et
1462 al., 2016). Chondrite data after Sun and McDonough (1989); (b)-(c) MORB-normalised
1463 multi-element spider diagrams for basalts. For comparison, the coloured fields show
1464 the composition variations of basalts from the Kyrenia Range (this study), the Misis
1465 Complex, S Turkey (Floyd et al., 1991), the Maden Complex, S Turkey (Erturk et al.,
1466 2018), the Japan Sea (Chen et al., 2015), the Sarmiento Complex, Chile (Saunders et
1467 al., 1979), the Bransfield Strait, Antarctic (Keller et al., 2002), the Guaymas Basin,
1468 Caribbean (Saunders et al., 1982) and the Tyrrhenian Sea, Italy (Peccherillo, 2017).
1469 MORB-normalised data are from Pearce et al. (1983). N-MORB, E-MORB and OIB
1470 data are from Sun and McDonough (1989). Specifically, two representative basalts of
1471 different affinities (within-plate vs. volcanic arc) from the Kyrenia Range are indicated.

1472

1473

1474

1475

1476

1477

1478

1479

1481 Listing of Supplementary Tables

1482 Supplementary Table S1. Major element oxides, trace and rare earth elements for the

1483 Late Cretaceous Fourkovouno (Selvilitepe) Formation.

1484 Supplementary Table S2. Major element oxides, trace and rare earth element analyses

1485 for the latest Cretaceous-Paleogene basaltic volcanics.

1486 Supplementary Table S3. Electron microprobe analyses of feldspar in the latest

1487 Cretaceous-Paleogene basaltic volcanics.

1488 Supplementary Table S4. Electron microprobe analyses of pyroxene in the latest

1489 Cretaceous-Paleogene basaltic volcanics.

1490 Supplementary Table S5. SIMS zircon U-Pb analyses of the zircon grains separated

1491 from the Late Cretaceous Fourkovouno (Selvilitepe) Formation.

Figure 1

[Click here to access/download;Figure;Fig. 1.jpg](#)

Figure 2

[Click here to access/download;Figure;Fig. 2.jpg](#)

Figure 3

[Click here to access/download;Figure;Fig. 3.jpg](#)

Figure 4

[Click here to access/download;Figure;Fig. 4.jpg](#)

Figure 5

Figure 6

[Click here to access/download;Figure;Fig. 6.jpg](#)

Figure 7

Figure 8

[Click here to access/download;Figure;Fig. 8.jpg](#)

Figure 9

[Click here to access/download;Figure;Fig. 9.jpg](#)

a

b

c

d

Figure 10

Figure 13

[Click here to access/download;Figure;Fig. 13.jpg](#)

Figure 15

[Click here to access/download;Figure;Fig. 15.jpg](#)

[Click here to access/download](#)

**Supplementary material/Appendix (Files for online
publication only)**

Supplementary Figure S1.jpg

[Click here to access/download](#)

**Supplementary material/Appendix (Files for online
publication only)**

Supplementary Figure S2.jpg

[Click here to access/download](#)

**Supplementary material/Appendix (Files for online
publication only)**

Supplementary Figure S3.jpg

[Click here to access/download](#)

**Supplementary material/Appendix (Files for online
publication only)**

Supplementary Figure S4.jpg

[Click here to access/download](#)

**Supplementary material/Appendix (Files for online
publication only)**

Supplementary Figure S5.jpg

[Click here to access/download](#)

**Supplementary material/Appendix (Files for online
publication only)**

Supplementary Figure S6.jpg

[Click here to access/download](#)

**Supplementary material/Appendix (Files for online
publication only)**

Supplementary Table S1.xlsx

[Click here to access/download](#)

**Supplementary material/Appendix (Files for online
publication only)**

Supplementary Table S2.xlsx

[Click here to access/download](#)

**Supplementary material/Appendix (Files for online
publication only)**

Supplementary Table S3.xlsx

[Click here to access/download](#)

Supplementary material/Appendix (Files for online publication only)

Supplementary Table S4.xlsx

[Click here to access/download](#)

Supplementary material/Appendix (Files for online publication only)

Supplementary Table S5.xlsx

[Click here to access/download](#)

RDM Data Profile XML

LITHOS9043_DataProfile.xml

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

--