THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

First and most northern occurrence of a thalattosuchian crocodylomorph from the Jurassic of the Isle of Skye, Scotland

Citation for published version:

Kean, K, Foffa, D, Johnson, M, Young, M, Greitens, G & Brusatte, S 2020, 'First and most northern occurrence of a thalattosuchian crocodylomorph from the Jurassic of the Isle of Skye, Scotland', *Scottish Journal of Geology*, vol. 56, no. 2. <https://doi.org/10.1144/sjg2020-013>

Digital Object Identifier (DOI):

[10.1144/sjg2020-013](https://doi.org/10.1144/sjg2020-013)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Scottish Journal of Geology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

First and most northern occurrence of a thalattosuchian crocodylomorph from the Jurassic of the Isle of Skye, Scotland

Kim J. Kean^{1*}, Davide Foffa², Michela M. Johnson, Mark T. Young¹, Gert Greitens, Stephen L. Brusatte¹

¹School of Geosciences, University of Edinburgh, Grant Institute, The King's Buildings, James Hutton Road, Edinburgh EH9 3FE, UK

² National Museums Scotland, Chambers St, Edinburgh EH1 1JF, UK

**Corresponding author (email: kean.kim.j@gmail.com)*

5982 Words, 122 References, 1 Table, 7 Figures

Abbreviated Title: First Thalattosuchian from Skye, Scotland

Abstract:

The Jurassic was a key interval for the evolution of dinosaurs, crocodylomorphs, and many other vertebrate groups. In recent years, new vertebrate fossils have emerged from the Early-Middle Jurassic of the Isle of Skye, Scotland; however, much more is known about Skye's dinosaur fauna than its crocodylomorphs. Here we report new crocodylomorph material collected from Jurassic marine deposits at Prince Charlie's Cave on the north east coast of Skye. The specimen is a small cobble containing postcranial elements from an individual that is considerably larger in size than previous crocodylomorphs described from Skye. Based on features of the vertebrae and osteoderms, the specimen is assigned to Thalattosuchia, an extinct clade of semiaquatic/pelagic crocodylomorphs. Specifically, the sub-circular and bean-shaped pit ornamentation on the dorsal surface of the osteoderms in alternating rows suggest affinities with the semi-aquatic lineage Teleosauroidea. Although the ornamentation pattern on the osteoderms is most similar to *Macrospondylus* ("Steneosaurus") *bollensis*, we conservatively assign the specimen to Teleosauroidea indeterminate. Regardless of its precise affinities and fragmentary nature, the specimen is the first thalattosuchian discovered in Scotland and is the most northerly reported Jurassic thalattosuchian globally, adding to our understanding of the palaeobiogeography and evolution of this group.

Introduction

Crocodylomorpha is a highly diverse clade of archosaurian reptiles with a long evolutionary history that spans over 200 million years from the Late Triassic to the present (Irmis et al., 2013). While modern crocodylomorph diversity is limited to semiaquatic ambush predators, extinct species experimented with an enormous variety of niches and modes of life ranging from small terrestrial cursorial insectivores and armoured herbivores (Crush, 1984; Buckley et al., 2000; O'Connor et al., 2010), to large terrestrial apex predators (Zanno et al., 2015; Godoy et al., 2014), and even some fully marine forms (Andrews, 1913; Young et al., 2010; Schwab et al., 2020).

Thalattosuchians were the only crocodylomorph clade, and the only archosaurs, that transitioned to a fully marine pelagic lifestyle. This transition occurred during the Jurassic, which therefore represents a key time in the evolutionary history of the group. Thalattosuchia is divided into two subgroups: Metriorhynchoidea, which includes the fully aquatic Metriorhynchidae (Young et al., 2010), and the primarily semi-aquatic Teleosauroidea (Johnson et al., 2015, 2017, 2018, 2020a,b,c; Foffa et al., 2019). The fossil record of Thalattosuchia spans from the Early Jurassic to the Early Cretaceous; however, it is patchy and strongly influenced by a few high-diversity European localities (Lagerstätten effect) in the Early, late-Middle, and Late Jurassic (Benson et al., 2010). Only a few million years after their initial appearance, both thalattosuchian lineages had achieved near-global distribution by the Middle Jurassic, with species known from Sub-Boreal, Continental and Tethyan Europe (e.g. Andrews, 1913; Cau and Fanti, 2011; Parrilla-Bel et al., 2013; Chiarenza et al., 2015), North America (Buffetaut, 1979; Wilberg, 2015), South America (Gasparini et al., 2000, 2005; Fernández and Gasparini, 2000; Fernández et al., 2011; Herrera et al., 2013a; Herrera, 2015), Siberia (Efimov, 1988), India (Phansalkar et al., 1994), and Africa (Fara et al., 2002; Haddoumi et al., 2016; Jouve et al., 2016; Dridi and Johnson, 2020). However, unlike other contemporaneous marine reptile groups, neither thalattosuchian lineage apparently inhabited high latitudes (Delsett et al., 2015; Zverkov et al., 2015a, b; Roberts et al., 2020), perhaps due to physiological constraints (Mannion et al. 2015; Séon et al., 2020). To date, teleosauroids and metriorhynchoids from the Middle-Late Jurassic of Yorkshire, England, represent the northernmost published occurrence of the clade (Benton and Taylor, 1984; Foffa et al., 2018a).

Here we report new crocodylomorph material (NMS G.2020.11.1) from the Jurassic of the Isle of Skye, Scotland. This new specimen is the sixth crocodylomorph taxon (Table 1) (Evans and Waldman, 1996; Wills et al., 2014; Young et al., 2016; Yi et al., 2017) from the increasingly diverse tetrapod assemblage of the Jurassic lagoonal/coastal deposits of Skye (Clark, 1993; Anquetin, 2010; Brusatte et al., 2015; Young et al., 2016; dePolo et al., 2018, 2020; Panciroli et al., 2017, 2018, 2019). Extensive comparisons with contemporaneous terrestrial and marine tetrapods demonstrate that this specimen shares affinities with thalattosuchian teleosauroids. Therefore, it represents the first known occurrence of thalattosuchian crocodylomorphs from the Isle of Skye (and Scotland), the most northern occurrence of the group globally, and provides new insights on the palaeogeography and early evolution of teleosauroids. Note that throughout the text, we refer to the recently published teleosauroid taxonomy provided by Johnson et al. (2020c), with species previously referred to as “*Steneosaurus*” originally indicated in brackets.

Institutional Abbreviations:

CAMSM – Sedgwick Museum of Earth Sciences, Cambridge, UK; IVPP – Institution of Palaeontology and Palaeoanthropology, Beijing, China; MNHN – Muséum National d’Histoire Naturelle, Paris, France; NHMUK – Natural History Museum, London, UK; NMS – National Museum of Scotland, Edinburgh, UK; PETMG – Peterborough Museum and Art Gallery, Peterborough, UK; SMNS – Staatliches Museum für Naturkunde Stuttgart, Stuttgart, Germany; SNHM – Staatliches Naturhistorisches Museum Braunschweig, Braunschweig, Germany.

1. Geological Background

The Inner Hebrides boasts one of the most complete fossiliferous Middle Jurassic sequences in the world, and the Isle of Skye, in particular, has become a globally important site for Jurassic research (Clark, 1993, 2001, 2018; Morton and Hudson, 1995; Anquetin, 2010; Wills et al., 2014; Brusatte et al., 2015; Young et al., 2016; Yi et al., 2017; dePolo et al., 2018, 2020; Panciroli et al., 2017, 2018, 2019). The specimen described here was found in 2018 by Gert Greitens as a loose cobble at Prince Charlie’s Cave, north of Portree (Fig. 1). As the cobble was not found in situ, it is difficult to determine its exact age and stratigraphic origin. Nevertheless, we can derive some hints of its provenance by comparing the matrix surrounding the specimen with the lithology of the geological formations exposed in the area. Fortunately, the Jurassic succession of Prince Charlie’s Cave is

one of the most complete of the Hebrides Basin, and spans from the uppermost Pliensbachian (~190.8 – 182.7 Ma - Early Jurassic) to the Bajocian (~170.3 – 168.3 Ma - Middle Jurassic), as described by Morton (2004). Here we summarise the main lithological features of each unit and invite the reader to refer to Morton (2004) for a more detailed account.

Surrounding the cave are almost continuous sea cliffs of the Scalpay Sandstone Formation (Pliensbachian-Toarcian). Stratigraphically the Scalpay Sandstone Formation is overlain by the Portree Shale Formation (Toarcian, ~182.7 – 174.1Ma), which is thicker and more complete at Prince Charlie's Cave than elsewhere in the Hebrides Basin, and is itself overlain by the Raasay Ironstone Formation (Toarcian) and Dun Caan Shale Member (Toarcian). The Scalpay Sandstone Formation consists of a grey, fine-medium grained sandstone that has been extensively bioturbated, and contains marine invertebrates such as crinoids, bivalves, brachiopods, and ammonites. At its base, the Portree Shale Formation is a claystone which becomes silty and sandy, and contains belemnites. Moving up the beds are light-dark grey claystones containing ammonites, and the top of the succession is defined by scattered iron-ooliths and abundant belemnites. The Raasay Ironstone Formation is divided into three parts; a basal section of crystalline calcareous ironstones, or ferruginous limestones; a middle section of claystone with two bands of ironstone nodules; and a top section consisting of a light-medium grey oolitic ironstone. Both ammonites and belemnites are found throughout the Raasay Formation, with belemnites being particularly abundant in the middle section. Overlaying the Raasay Formation is the dark-grey micaceous shale of the Dun Caan Shale Member which has no ammonites (Morton, 2004; Morton and Hudson, 1995).

NMS G.2020.11.1 is a cobble of claystone with a fine-grained clay matrix that is well-cemented and grey in colour. This may indicate that the cobble belongs to the Portree Shale Formation, suggesting a lower Toarcian age (~182.7 Ma) for the specimen, which would place it in the Early Jurassic. With that said, because the specimen was found loose and water worn, it has certainly been reworked, and it therefore could have derived from nearly any Jurassic-aged unit along the east coast of the Trotternish Peninsula of Skye, so we conservatively consider it Early to Middle Jurassic in age here.

2. Systematic Palaeontology

Specimen. *NMS G.2020.11.1*: a small cobble containing eight complete and partial osteoderms, three incomplete vertebrae, and additional rib fragments. The osteoderms are fragmented and due to the low contrast between matrix and bones in the μ CT-scans, diagnostic features can only be observed clearly on two of the osteoderms. The vertebrae are also incomplete; the neural spine is present in only one of the vertebrae, which has a single complete transverse process. In the remaining vertebrae the neural spine is missing, and the transverse processes are either absent or incomplete. There are no chevrons or cervical ribs preserved.

Locality. Collected by G. G. at Prince Charlie's Cave, north of Portree (the cobble was not found in situ).

Preparation and Scanning. As the majority of the fossil material in NMS G.2020.11.1 is embedded in the matrix, the specimen was subjected to X-ray computed tomography (CT) scanning by Dr. Thomas Davies using a Nikon XT H 225 μ CT scanner at the Palaeobiology Lab of the University of Bristol. Digital segmentation of the fossils from the matrix was conducted by Kim Kean at the University of Edinburgh, School of GeoSciences, using Materialise Mimics 20.0.

3. Description

NMS G.2020.11.1 is a 14 cm long, 8 cm wide, and 4 cm tall cobble (Fig. 2). It contains at least eighteen bones: three vertebrae, eight osteoderms, and seven ribs, plus other fragmentary unidentified elements, all associated in a fine grained, grey, well-cemented, clay matrix. The bones are well preserved but slightly damaged. The cobble has been rounded, likely by attrition and hydraulic action while it was in the North Atlantic Ocean. The bones that can be seen on the surface of the cobble have been damaged by these erosional processes, in particular the vertebrae, which are all incomplete as a result.

3.1 Vertebrae

Three incomplete vertebrae are preserved in NMS G.2020.11.1 (Fig. 3). They are all partially exposed on the surface of the cobble and weathered to varying degrees. The neural spine and a prezygapophysis are preserved on one of the three vertebrae (Fig. 3A). The postzygapophyses are missing in all vertebrae, and only two have preserved transverse processes. The transverse

processes (Fig. 3A and B) are long and wide, with rounded distal ends. They are broadened anteroposteriorly and show the diapophyseal and parapophyseal rib attachments. The neural spine is partially preserved and has been tilted approximately 35° laterally relative to the centrum; it is rectangular in lateral view, thin, and taller than the centrum at approximately 3.44 cm tall (Fig. 3A). The single preserved prezygapophysis is ellipsoidal in shape and extends slightly anterior to the articular surface of the centrum. The centra of all three vertebrae are spool-shaped, anteroposteriorly elongated, taller than wide, and amphicoelous. In ventral and lateral views, the centra show the typical hourglass figure of crocodylomorph vertebrae (Fig. 3) (Andrews, 1913; Young et al., 2013; Johnson et al., 2017). The exact position of the vertebrae along the spinal column is unclear, but the absence of rib attachment on the centra rules out a cervical position. Similarly, the absence of sacral ribs, or caudal chevrons/haemal arches indicate that these centra do not come from the sacral or caudal regions. Instead, the long, spool-shaped centra, and the elongated transverse processes with separated rib attachments, demonstrate that these are vertebrae from the middle-posterior dorsal section of the vertebral column (Andrews, 1913; Romer, 1956). As the neural arch is largely missing in the third preserved vertebra (Fig. 3C), it is possible that the neurocentral suture was open, which would indicate that this individual was not yet fully mature (Brochu, 1996; Herrera et al., 2013b), although its exact ontogenetic stage is unclear.

3.2 Osteoderms

As the bones in NMS G.2020.11.1 are disarticulated and have not maintained their in-vivo association, it is impossible to say with confidence if the osteoderms originated from the dorsal or ventral series. The osteoderms are rectangular- or triangular-shaped with a flat, pitted dorsal surface, and a slightly convex and smooth ventral surface. There are eight osteoderms in the block in total, five of which could be confidently segmented due to low contrast in the μ CT data (Figs. 4, 5). The most complete osteoderms are shown in Fig. 4. The osteoderm in Fig. 4A is rectangular in shape; it is approximately 5 cm in length and 3 cm in width. It is ornamented with well separated bean-shaped, sub-circular, and sub-triangular pits that are arranged into alternate rows. The osteoderm shown in Fig. 4B is comparatively thin and irregular in shape. It is approximately 4 cm in length and ranges in width from 6 to 2 cm as it tapers to a point. The unfragmented edges of the osteoderm suggest that its irregular shape is not due to erosion or wear. The pits ornamenting the surface are circular and sub-triangular and decrease in size as they radiate out from the centre.

The osteoderms depicted in Fig. 5 are incomplete, and around 3 cm in width and 1.5-3 cm in length. The pits are bean-shaped and sub-circular, and arranged into alternating rows, similar to the pattern observed in Fig. 4A. The difference in shape and pit pattern of Fig. 4B could be reflective of the osteoderms belonging to different areas of the body (Vickaryous and Hall, 2008). A keel is absent on all the osteoderms observed, however, we do not consider this a reliable diagnostic character as ornamentation on the osteoderms of crocodylomorphs is formed by the resorption and redeposition of cortical bone. Therefore, pits and ridges are routinely altered, eroded, or filled during ontogeny (Clarac et al., 2015). In addition, the prominence of a keel can vary greatly within individuals of a single species. Given that a keel is absent in NMS G.2020.11.1, we have focused on comparisons between the pit ornamentation of sacral osteoderms amongst teleosauroids, which are generally diagnostic.

3.3 Other fragments

There are several ribs and unidentifiable bone fragments in NMS G.2020.11.1. In cross-section the ribs are semi-circular. The most complete fragment as shown in Fig. 5D is 16 cm long and approximately 1 cm in diameter in the rib body, and has a similar morphology to the thoracic ribs of the Middle Jurassic teleosauroid from Tunisia described by Dridi and Johnson (2020). The rib has a break towards the distal end of the rib body, but when complete it was likely uniformly curved distally. The proximal end of the rib is double headed with a tuberculum and capitulum. The capitulum would have been in contact with the parapophysis of the transverse process. It would appear that the tuberculum is rounded, however this, along with finer details of the rib's morphology, are difficult to discern due to the low contrast between it and the surrounding matrix in the μ CT-scan.

4. Crocodylomorph Affinity of NMS G.2020.11.1

The Inner Hebrides were home to diverse terrestrial and marine faunas during the Jurassic. Since the earliest discoveries by Hugh Miller (1858), an ever increasing variety of amphibians (Evans and Waldman, 1996), lepidosauromorphs (Waldman and Evans, 1994), turtles (Anquetin, 2010), dinosaurs (Clark, 2018; dePolo et al., 2018, 2020), ichthyosaurs (Brusatte et al., 2015), plesiosaurs (Clark, 1993), crocodylomorphs (Evans and Waldman, 1996; Wills et al., 2014; Young et al., 2016;

Yi et al., 2017), and mammaliamorphs (Waldman and Savage, 1972; Panciroli et al., 2017, 2018, 2019) have been reported, represented both by body and trace fossils.

Although NMS G.2020.11.1 is largely incomplete and poorly preserved, it still yields a suite of vertebral and osteodermal characters that distinguishes it from contemporaneous taxa previously reported on the Isle of Skye. The following vertebral and osteodermal characters suggest a crocodylomorph affinity for NMS G.2020.11.1:

i) Long hourglass-shape and amphicoelous vertebral centra (Fig. 3) (Andrews, 1913; Young et al., 2013; Johnson et al., 2017). Turtles also have vertebrae with an hourglass centrum (Joyce et al., 2009; Pérez-García, 2012). However, as in *Eileanchelys waldmani* from Skye, the vertebrae of turtles differ from those of crocodylomorphs and NMS G.2020.11.1 in being platycoelous (Joyce et al., 2009; Anquetin, 2010; Pérez-García, 2012). An affinity with dinosaurs can also be ruled out based on the morphology of the vertebrae in NMS G.2020.11.1. Although the dorsal vertebrae of ornithischians are typically longer than they are tall as is the case with NMS G.2020.11.1, they do not have a ventral surface as strongly concave as those in crocodylomorphs and NMS G.2020.11.1 (Galton, 1985). The vertebrae of saurischians are similarly distinct; the dorsals of sauropods, for example, are typically opisthocoelous and ornamented by deep pleurocoelous fossae (Romer, 1956; Wilson et al., 2011; Wilson, 2012). Pleurocoels (pneumatic fossae and foramina) are also seen in the dorsals of theropods, whose centra tend to be platycoelous (Romer, 1956; Rauhut, 2005). Furthermore, saurischians typically have a series of thin laminae on their dorsal vertebrae, but these are absent in NMS G.2020.11.1 (Wilson et al., 2011; Wilson, 2012). Amongst marine reptiles, the vertebrae of ichthyosaurs are amphicoelous, but are very short and their centra are not hourglass-shaped (McGowan and Motani, 2003). The vertebrae of plesiosaurs do not possess an “hourglass” profile, and the transverse processes of their dorsal vertebrae are rod-like, not plate-like as in crocodylomorphs and NMS G.2020.11.1. The transverse processes also diverge from the centra at an angle from the sagittal plane in plesiosaurs, whereas they are perpendicular to the centra in crocodylomorphs and NMS G.2020.11.1 (Andrews, 1910, 1913).

ii) Double rib attachment (diapophyses and parapophyses) on the transverse process (Fig. 3). Double-headed ribs are not exclusive to crocodylomorphs, and are also seen in ichthyosaurs and dinosaurs, whilst marine reptiles such as plesiosaurs and turtles have single-headed ribs (McGowan and Motani, 2003; Huh et al., 2011; Kim et al., 2011). However, the morphology of the rib attachments differs among these groups. In ichthyosaurs the diapophyses and parapophyses are located on the lateral surfaces of the centra of the dorsal vertebrae (McGowan and Motani, 2003). In the anterior dorsals of saurischian dinosaurs the parapophysis lies on the centrum and migrates to the neural arch in the middle and posterior dorsal vertebrae (Wilson et al., 2011). In the dorsal vertebrae of crocodylomorphs the parapophyses and diapophyses are located on the transverse process (Andrews, 1913), as is the case in NMS G.2020.11.1.

iii) Presence and shape of osteoderms (Figs. 4, 5). Amongst species in the Jurassic terrestrial and marine faunas of Skye, an integumentary skeleton of dermal bone is present in turtles, lepidosauromorphs, and archosaurs (Waldman and Evans, 1994; Vickaryous and Sire, 2009; Anquetin, 2010; Wills et al., 2014). However, the carapace of turtles and lepidosaurian scales differ substantially from the thick, disc-shaped osteoderms of archosaurs (Vickaryous and Sire, 2009). Amongst Early-Middle Jurassic archosaurs, osteoderms can be found in the vast majority of pseudosuchians (the larger group of crocodylomorphs and close relatives) and some ornithischian dinosaurs (e.g. stegosaurs and ankylosaurs). Osteoderms are also present in non-archosaurian archosauriformes such as the Triassic Proterochampsia and Doswelliidae (Cerdeña et al., 2015; Ezcurra 2016). Some of these groups can be readily excluded as potential identities for NMS G.2020.11.1. Non-crocodylomorph pseudosuchians and non-archosaurian archosauromorphs did not survive into the Jurassic. The osteoderms of stegosaurs take the form of unornamented and enlarged dermal plates that are organised in rows along the dorsolateral body surface (de Buffrenil et al., 1986; Main et al., 2005; Maidment et al., 2015). Ankylosaur osteoderms are often characterised by a keel and a dendritic pattern of vascularised furrows that radiate away from the keel and randomly organised pits on the dorsal surface (Burns, 2008). By contrast the pitted osteoderms present in NMS G.2020.11.1 resembles those of crocodylomorphs in shape, ornamentation, size, and association with the vertebral column (Burns et al., 2013; Schwarz et al., 2017).

Overall, the size and combination of vertebral and osteodermal characters demonstrates that NMS G.2020.11.1 can be referred to a relatively large-bodied crocodylomorph. As several crocodylomorph taxa have been reported from the Jurassic of Skye (Table 1) over the last two decades (Evans and Waldman, 1996; Wills et al., 2014; Young et al., 2016; Yi et al., 2017), the next section aims to clarify whether NMS G.2020.11.1 belongs to any of these taxa. Note that all previous Skye crocodylomorphs come from Middle Jurassic rocks, whereas the provenance of NMS G.2020.11.1 cannot be substantiated beyond the Jurassic generally (as discussed above).

4.1 Taxonomic comparisons with other Early and Middle Jurassic crocodylomorphs

Here we compare NMS G.2020.11.1 to Early and Middle Jurassic crocodylomorph clades found on Skye, as well as worldwide. We have divided them into non-thalattosuchian crocodylomorphs, namely Goniopholididae, Atoposauridae and Pholidosauridae, and thalattosuchian crocodylomorphs.

4.1.1 Non-thalattosuchian crocodylomorphs

The crocodylomorph material known from Skye (Table 1) includes an incomplete dentary of a hylaeochampsid-like taxon (Yi et al., 2017), a partial lower jaw from the atoposaurid *Theriosuchus* (Young et al., 2016), a disarticulated incomplete postcranial skeleton from a goniopholidid (Evans and Waldman, 1996), a left pubis from an indeterminate mesoeucrocodylian (Wills et al., 2014), and a tooth and disarticulated osteoderms from an indeterminate neosuchian (Wills et al., 2014). These specimens demonstrate that Skye hosted a variety of small-sized neosuchians in nearshore environments during the Middle Jurassic. However, none of these groups share the same vertebral and osteodermal characteristics as NMS G.2020.11.1, and were significantly smaller in body size than NMS G.2020.11.1.

Indeterminate neosuchian (Wills et al., 2014) - Two previous crocodylomorph specimens containing osteoderms have been reported from the Jurassic of Skye: NHMUK PV R 36713 (assigned by Wills et al. (2014) to *Neosuchia* indet.) and G.1992.47.6 (assigned to Goniopholididae, discussed below). NHMUK PV R 36713 was found in the Kilmaluag Formation on the southwest Elgol Peninsula of Skye and contains thirteen partial osteoderms. Some of the more complete osteoderms show an overall quadrilateral structure ornamented by a straight

longitudinal keel and circular to oval pits (Fig. 6A) (Wills et al., 2014). The largest of the osteoderms is approximately 20 mm wide (Wills et al., 2014), which is significantly smaller than the osteoderms of NMS G.2020.11.1, which are up to 50 mm in width. Most pits ornamenting the osteoderms in NHMUK PV R 36713 are circular, with a few that are oval in shape (Fig. 6A) (Wills et al., 2014). The pits on NMS G.2020.11.1 are more varied in shape, with some that are circular, oval, and bean-shaped (Figs. 4, 5).

Goniopholididae - The dorsal osteoderms of Goniopholididae are plate-like, quadrilateral structures which are wider than long. The paramedial osteoderms of *Goniopholis*, for example, have a heavily ornamented dorsal surface, lack a keel, and are sub-rectangular and wider than long (Andrade et al., 2012; Wills et al., 2014). In contrast, cervical dorsolateral osteoderms from *Sunosuchus* described by Averianov (2000) are longer than wide and have a high longitudinal keel with oval-shaped pits arranged in alternating rows ornamenting the dorsal surface. Typically, the pits ornamenting the dorsal surface of Goniopholididae osteoderms are oval to circular in shape, while the anterior margin of the dorsal surface tends to be smooth (Fig. 6B) (Averianov, 2000; Kuzmin et al., 2013; Puértolas-Pascual et al., 2015; Puértolas-Pascual and Mateus, 2020). A partial postcranial skeleton with osteoderms discovered in the Kilmaluag Formation of Skye was assigned to Goniopholididae by Evans and Waldman (1996). This specimen (G.1992.47.6) is currently being held in the NMS collections and has significantly smaller osteoderms than NMS G.2020.11.1. The osteoderms in NMS G.2020.11.1 are also not as densely pitted as that of Goniopholididae indet. shown in Fig. 6B, nor are the pits as large. Whilst the pits are arranged in regular rows in both *Sunosuchus* and NMS G.2020.11.1, the pits in NMS G.2020.11.1 vary more in morphology than those in *Sunosuchus*.

***Theriosuchus* and Atoposauridae** - The *Theriosuchus* material previously described from Skye by Young et al. (2016) does not include osteoderms. Therefore, we will be making direct comparisons between NMS G.2020.11.1 and taxa closely related to *Theriosuchus*. In addition to *Theriosuchus* sp., atoposaurid taxa from the Jurassic include *Alligatorellus* sp., *Alligatorium* sp., *Atoposaurus* sp., and *Knoetschkesuchus langenbergensis* (Schwarz-Wings et al., 2011; Tennant and Mannion, 2014; Schwarz et al., 2017). The ventral osteoderms of these Late Jurassic atoposaurids tend to be quadrilateral in shape and ornamented with a shallow keel and circular pits

(Schwarz-Wings et al., 2011; Tennant and Mannion, 2014; Schwarz et al., 2017). NMS G.2020.11.1 can be excluded from Atoposauridae in part because of its larger size. Atoposaurids were characteristically very small in size, rarely reaching more than 1 m in overall body length (Schwarz and Salisbury, 2005; Schwarz et al., 2017). Given that NMS G.2020.11.1 shows single vertebrae 3-4 cm in length, the animal would have been significantly larger. In addition, the osteoderms of NMS G.2020.11.1 do not show the same circular pitting as observed in atoposaurid taxa.

Pholidosauridae - Pholidosaurids were common in freshwater and marginal marine environments during the Late Jurassic and Early Cretaceous (Martin and Buffetaut, 2012). Their dorsal osteoderms are typically rectangular in shape and are densely ornamented by small, deep, circular and subcircular pits (Jouve and Jalil, 2020). With the thoracic osteoderms being highly elongated mediolaterally (see Jouve and Jalil, 2020). This does not match the shape or ornamentation observed in NMS G.2020.11.1.

4.1.2 NMS G.2020.11.1 within Thalattosuchia

Despite the diversity of crocodylomorphs on Skye, no evidence of thalattosuchians has previously been recovered from the island, even though they are one of the most diverse crocodylomorph clades of the Mesozoic (Pierce and Benton, 2006; Young et al., 2010, 2014; Young and Steel, 2014; Johnson et al., 2017; Foffa et al., 2019; Johnson et al., 2020c). This is puzzling for two reasons: firstly, there are abundant marine deposits on Skye, and secondly, thalattosuchians were relatively diverse in the Early and Middle Jurassic with multiple taxa found in the lagerstätten of England (Andrews, 1910, 1913; Pierce and Benton, 2006; Brusatte et al., 2016; Foffa et al., 2018a,b). Specifically, three teleosauroid taxa (*Plagiophthalmosuchus* [“*Steneosaurus*”] *gracilirostris*, *Macrospondylus* [“*Steneosaurus*”] *bollensis*, and *Mystriosaurus*) (Westphal, 1961; Walkden et al., 1987; Sachs et al., 2019), and one basal metriorhynchoid (*Pelagosaurus typus*) (Pierce and Benton, 2006) are known from the Lower Jurassic of Southern England and Yorkshire; and at least four teleosauroids (*Neosteneosaurus* [“*Steneosaurus*”] *edwardsi*, *Charitomenosuchus* [“*Steneosaurus*”] *leedsi*, *Mycterosuchus nasutus*, *Lemmysuchus obtusidens*) (Andrews 1913; Johnson et al., 2015, 2017; Foffa et al., 2018a,b), and seven metriorhynchids (*Ieldraan melkshamensis*, *Suchodus durobrivensis*, “*Metriorhynchus*” *brachyrhynchus*, *Tyrannoneustes*

lythrodektikos, *Dakosaurus*–like specimens, *Metriorhynchus superciliosus*, *Gracilineustes leedsi*) (Andrews, 1913; Young and Andrade, 2009; Young et al., 2012; Foffa and Young, 2014; Foffa et al., 2017, 2018a,b, 2019), are known from the late Middle Jurassic of the Oxford Clay Formation. The following paragraphs will be used to compare NMS G.2020.11.1 to worldwide Early and Middle Jurassic teleosauroids and basal metriorhynchoids, with emphasis on taxa from England. Due to the limited variation of vertebrae in teleosauroids we largely focus on differences in osteoderm shape, thickness, and ornamentation. NMS G.2020.11.1 will not be compared with metriorhynchids as they do not possess osteoderms (Young et al., 2013).

Basal Metriorhynchoids - Early Jurassic metriorhynchoids such as *Pelagosaurus typus* (Fig. 6C) and *Magyarosuchus fitosi* (Fig. 6D) had osteoderms, unlike derived metriorhynchoids of the family Metriorhynchidae. However, the osteoderms of all known basal metriorhynchoids differ from those in NMS G.2020.11.1; for instance, the paramedial dorsal osteoderms of *Pelagosaurus typus*, which are also sub-rectangular, have a very low keel, and a honey-comb pattern of pits (Pierce and Benton, 2006), and are more densely ornamented (e.g. more numerous pits) than NMS G.2020.11.1 (Fig. 6C). The dorsal osteoderms of *Magyarosuchus fitosi*, on the other hand, are large, rectangular to slightly rounded in shape, and are ornamented with an anteroposterior dorsal keel and irregularly shaped pits (Fig. 6D) (Ősi et al., 2018). The pits range from circular, ellipsoid, bean-shaped, triangular to quadrangular morphologies, and show variations in size from small to very large on the dorsal surface (Ősi et al., 2018). Whilst the pits ornamenting *M. fitosi* are predominantly triangular in shape, NMS G.2020.11.1 is largely adorned by sub-circular and bean-shaped pits. In addition, the disparity in size of pits is not as large in NMS G.2020.11.1 as it is in *M. fitosi*.

Early Jurassic Teleosauroids - Specific teleosauroids that are known from the Early Jurassic include *Pl. gracilirostris* (NHMUK PV OR 14792) (Fig. 7A), *Mystriosaurus laurillardii* (Fig. 7B), *Platysuchus multiscrobiculatus* (SMNS 9930) (Fig. 7C), and *Ma. bollensis* (SMNS 51563) (Fig. 7D). The Chinese teleosauroid (IVPP RV 10098), which was previously referred to as *Peipehsuchus teleorhinus* (Li, 1993), may also be from the Early Jurassic. However, the provenance of IVPP RV 10098 and the holotype (IVPP RV 48001) is somewhat uncertain, and so the Chinese teleosauroid is considered to be Early-Middle Jurassic in age (Martin et al., 2018).

Osteoderms are known from all but *Mystriosaurus laurillardi* and the Chinese teleosauroid. However, one particular specimen located at the Staatliches Naturhistorisches Museum (SNHM) in Braunschweig, Germany (SNHM-IG-008-R), which was assigned to “*Steneosaurus*” af. *bollensis* by Wincierz (1967) contains partial osteoderms, and is currently thought to belong to *Mystriosaurus* sp. Comparisons will therefore be made between SNHM-IG-008-R and NMS G.2020.11.1. It is important to note that a formal description is currently in preparation by MMJ and colleagues, and is beyond the scope of this paper.

The osteoderms of *Pl. gracilirostris* are unfortunately poorly preserved, making direct comparisons difficult. The osteoderms appear to have small, moderately spaced pits, and is ornamented by a very weak keel (Fig. 7A). In contrast, the osteoderms of *Ma. bollensis* are ornamented by numerous, relatively close pits which are semi-ellipsoidal in shape (Fig. 7D). However, there is great variety in keel size within *Ma. bollensis*, whilst some have prominent keels, other individuals of comparable size can have weak keels. *Platysuchus multiscrobiculatus* has a weak keel on the osteoderms, with small, circular, numerous pits that are densely packed together (Fig. 7C). The pits on the dorsal osteoderms of *Mystriosaurus* sp. are closely packed together, ellipsoidal in shape, and are arranged in a subtle ‘starburst’ pattern (Fig. 7B).

The shape of the pits in NMS G.2020.11.1 are not dissimilar to those of *Ma. bollensis*, however unlike the osteoderms of *Ma. bollensis*, *P. multiscrobiculatus*, and *Mystriosaurus* sp., those of NMS G.2020.11.1 do not have densely packed pits or a keel. The pits on NMS G.2020.11.1 are not arranged in any kind of ‘starburst’ pattern either, as is the case in *Mystriosaurus* sp.

Middle Jurassic Teleosauroids - Teleosauroid taxa from the Bathonian include *Teleosaurus cadomensis* (Fig. 7E), *Yvridiosuchus boutilieri*, *Deslongchampsina larteti*, *Seldsienean* (“*Steneosaurus*”) *megistorhynchus*, *Clovesuurdameredeor* (“*Steneosaurus*”) *stephani*, and *Andrianavoay* (“*Steneosaurus*”) *baroni* (Fig. 7F); however, osteoderms are currently only known from *T. cadomensis* (NMHUK PV R 119a) and *A. baroni* (NHMUK PV R 1999). The dorsal osteoderms of *T. cadomensis* are similar to those of *Platysuchus*, in which they are ornamented by a weak keel and numerous, small, circular, densely packed pits (Fig. 7E) (Johnson et al., 2020a). Only a small fragment of dorsal osteoderm is preserved for *A. baroni* (Fig. 7F). It shows well

separated, moderately sized, and subcircular pits which resembles the pit ornamentation of NMS G.2020.11.1. However, it is currently unclear where this osteoderm may have been along the body; it is also important to note that *A. baroni* (NHMUK PV R 1999) is currently only known from Madagascar (Newton, 1893).

More well-known, Callovian teleosauroids include *C. leedsi* (NHMUK PV R 3806) (Fig. 7G), *N. edwardsi* (PETMG R178) (Fig. 7H), *Lemmysuchus obtusidens* (NHMUK PV R 3168) (Fig. 7I), and *Mycterosuchus nasutus* (NHMUK PV R 2617) (Fig. 7J), all of which have well preserved, multiple dorsal osteoderms. Those of *C. leedsi* are ornamented by a shallow keel and small-to-large sub-circular and elliptical pits, arranged in a semi-circular pattern (Fig. 7G). In *N. edwardsi* (Fig. 7H), the pits are few, subcircular and exceptionally large (especially in the middle of the osteoderm). In *L. obtusidens* (Fig. 7I), the osteoderms display distinctive elongated pits that merge together (notably along the lateral margin) as well as irregularly shaped, relatively close pits that are arranged in a ‘starburst’ pattern. The pits on the osteoderms of *My. nasutus* are well separated and are largely irregular in shape, some of which are bean-shaped. The osteoderms in this taxon possess a relatively well developed keel (Fig. 7J). This pit pattern is not dissimilar to that of NMS G.2020.11.1, which also has sub-circular and some bean-shaped pits arranged into alternate rows, with the smallest pits on the edges of the osteoderm. However, among Teleosauroidea, the dorsal osteoderms of *My. nasutus* are unusually large and thick (DF. pers. obs. on NHMUK PV R 2617 and CAMSM J. 1420), which are features not seen in NMS G.2020.11.1.

In summary, NMS G.2020.11.1 was recovered from marine beds, and is significantly larger than any crocodylomorph material previously discovered on Skye. Its vertebral morphology and osteoderm ornamentation exclude an affinity between NMS G.2020.11.1 and any other crocodylomorph clade previously described from the Isle of Skye. Instead, the osteoderms of NMS G.2020.11.1 most closely resemble those of the Early Jurassic thalattosuchian teleosauroid *Ma. bollensis*, indicating that it is most likely a teleosauroid, which is concordant with its large body size, Jurassic age, and aquatic provenance. However, we acknowledge that NMS G.2020.11.1 is composed of fragmentary material which are of limited diagnostic utility. Therefore, we do not attempt to assign NMS G.2020.11.1 to a specific taxon and instead conservatively consider it Teleosauroidea indeterminate.

5. Discussion

Research over the last two decades has shown that Skye was home to a diverse crocodylomorph fauna of small-bodied, non-pelagic, neosuchians such as goniopholidids, atoposaurids and a hylaeochampsid-like taxon in the Jurassic (Evans and Waldman, 1996; Young et al., 2016; Yi et al., 2017). Not only is NMS G.2020.11.1 the first evidence of larger semi-aquatic crocodylomorphs on Skye during the Jurassic, it is also the most northerly thalattosuchian discovered to date. Previously the most northern thalattosuchians were recovered from the Blue Lias of Yorkshire on the east coast of England (Benton and Taylor, 1984; Foffa et al., 2018a).

The distribution of thalattosuchians appears to be more restricted than that of other marine reptiles. Marine pseudosuchians, including thalattosuchians, inhabited waters between palaeolatitudes of $\sim 44^{\circ}\text{N}$ and $\sim 38^{\circ}\text{S}$ (Mannion et al., 2015). With a palaeolatitude of $\sim 40^{\circ}\text{N}$, Scotland was within the upper limits of this hypothesised habitable range for thalattosuchians (Nunn et al., 2009). Other marine tetrapods (e.g. ichthyosaurs and plesiosaurs) had a wider latitudinal range between $\sim 69^{\circ}\text{N}$ and 42°S (Mannion et al., 2015); and unlike thalattosuchians, pelagic sauropterygians and ichthyosaurs have been discovered much further north of Scotland, in Svalbard and Siberia (Delsett et al., 2015; Rogov et al., 2018; Zverkov et al., 2015a,b, 2018, 2019a,b). One possible explanation for this discrepancy is that plesiosaurs and ichthyosaurs were fully homeothermic endotherms, possibly being able to maintain their core temperatures between $32\text{--}40^{\circ}\text{C}$ (Bernard et al., 2010; Motani, 2010; Séon et al., 2020).

While Jurassic thalattosuchians do not show evidence for homeothermic endothermy, with large-bodied teleosauroids potentially being ‘gigantothermic’ (like modern saltwater crocodiles), metriorhynchids may have been endothermic but poorly homeothermic (see Séon et al., 2020). Young et al. (2010) hypothesized that the closure of the external mandibular fenestrae in metriorhynchids was due to them no longer requiring mouth-gape basking to thermoregulate, a behaviour which teleosauroids likely utilized to raise their body temperatures, similar to extant crocodylians (Séon et al., 2020). This may have restricted teleosauroids to warmer, more hospitable climates and rendered more pole-ward habitats unfeasible. However, among extant crocodylians the American and Chinese alligators (*Alligator mississippiensis* and *Alligator*

sinensis) are able to inhabit temperate climates, and have the greatest tolerance to cold environments of living crocodylomorphs, surviving by brumating in the winter, either underwater or in burrows (e.g. Lang, 1987).

There is growing evidence that sympleiomorphically, crocodylomorphs had some form of homeothermic and/or endothermic thermophysiology (Seymour et al., 2004; Summers, 2005; de Ricqlès et al., 2008; Farmer and Sanders, 2010; Legendre et al., 2016). It has been hypothesised that thalattosuchians reverted to poikilothermic ectothermy when they became aquatic ambush predators during the Early Jurassic (Seymour et al., 2004; Mannion et al., 2015). Séon et al. (2020) suggested that endothermy may have been ‘reactivated’ as metriorhynchids became better adapted to an active pelagic lifestyle. Because of this difference in thermoregulatory strategy, metriorhynchoids were likely less restricted by palaeolatitude than teleosauroids (Séon et al., 2020). In recent years, several major disparities in the ecological niches and modes of life between metriorhynchoids and teleosauroids have been uncovered using skeletal, neuroanatomical, and geochemical data (e.g. Foffa et al., 2018b; Schwab et al., 2020; Séon et al., 2020).

6. Conclusion

Here we report the discovery of new crocodylomorph material recovered from Jurassic marine deposits on the Isle of Skye. NMS G.2020.11.1 is a small cobble containing partial vertebrae, osteoderms, and ribs, from an animal larger than previous Jurassic crocodylomorphs described from Skye. We establish that the material can be assigned to an indeterminate teleosauroid thalattosuchian crocodylomorph based on vertebral and osteodermal characters. The pattern of the pitting ornamenting the dorsal surface of the osteoderms, for instance, most closely resembles that of the teleosauroid *Macrospondylus bollensis*. NMS G.2020.11.1 is consequently the first reported thalattosuchian from Scotland, and the most northerly thalattosuchian material recovered to date worldwide. NMS G.2020.11.1 is therefore an important specimen despite its fragmentary nature, as it reveals further information on the palaeobiogeography of thalattosuchians during a poorly sampled period of time. It suggests that other thalattosuchians may be found in Scotland in the future, including other teleosauroids, and metriorhynchoids which were probably better adapted for more northerly waters in the Jurassic.

7. Acknowledgments

We are grateful to Stig Walsh for facilitating the borrowing of the specimen from the NMS collection. Thanks to Dr Thomas Davies at the University of Bristol who conducted the μ CT scanning in August 2019. We thank Dugald Ross, Mark Wilkinson, and Neil Clark for their collaboration in studying Skye vertebrates, and for discussion and fieldwork collaboration. We also thank Atilla Ösi for providing photos of osteoderms of Goniopholididae indet., and Sven Sachs for providing information regarding the *Mystriosaurus* sp. specimen SNHM-IG-008-R. We would also like to thank our reviewers, Neil Clark and Eric Wilberg, for their kind and constructive comments, and Romesh Niranjana Palamakumbura for handling the manuscript as editor. The PalAlba team's work on the Isle of Skye is funded by the National Geographic Society (GEFNE185-16), Derek and Maureen Moss, the Edinburgh Zoo, and the Edinburgh Geological Society. This is PalAlba publication Number 10.

8. References

- Andrade, M.B., Edmonds, R., Benton, M.J. and Schouten, R. 2012. A new Berriasian species of *Goniopholis* (Mesoeucrocodylia, Neosuchia) from England, and a review of the genus. *Zoological Journal of the Linnean Society*. Vol. 163 (suppl_1), pp. S66–S108
- Andrews, C.W. 1910. A descriptive catalogue of the marine reptiles of the Oxford Clay, Part One *British Museum (Natural History), London*
- Andrews, C.W. 1913. A descriptive catalogue of the marine reptiles of the Oxford Clay, Part Two. *British Museum (Natural History), London*
- Anquetin, J. 2010. The anatomy of the basal turtle *Eileanchelys waldmani* from the Middle Jurassic of the Isle of Skye, Scotland. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*. Vol 101, pp. 67–98
- Averianov, A.O. 2000. *Sunosuchus* sp. (Crocodylomorpha, Goniopholididae) from the Middle Jurassic of Kirghisia. *Journal of Vertebrate Paleontology*. Vol. 20 (4), pp. 776–779

- Benson, R.B.J., Butler, R.J., Lindgren, J., and Smith, A.S. 2010. Mesozoic marine tetrapod diversity: mass extinctions and temporal heterogeneity in geological megabiases affecting vertebrates. *Proceedings of the Royal Society B*. Vol. 277, pp. 829-834
- Benton, M.J. and Taylor, M.A. 1984. Marine reptiles from the Upper Lias (Lower Toarcian, Lower Jurassic) of the Yorkshire Coast. *Proceedings of the Yorkshire Geological Society*. Vol. 44 (4), pp. 399–429
- Bernard, A., Lécuyer, C., Vincent, P., Amiot, R., Bardet, N., Buffetaut, E., Cuny, G., Fourel, F., Martineau, F., Mazin, J-M. and Prieur, A. 2010. Regulation of Body Temperatures by some Mesozoic Marine Reptiles. *Science*. Vol. 328 (5984), pp. 1379-1382
- Brochu, C.A. 1996. Closure of neurocentral sutures during crocodilian ontogeny: implications for maturity assessment in fossil archosaurs. *Journal of Vertebrate Paleontology*. Vol. 16, pp. 49–62
- Brusatte, S.L., Young, M.T., Challands, T.J., Clark, N.D.L., Fischer, V., Fraser, N.C., Liston, J.J., MacFadyen, C.C.J., Ross, D.A., Walsh, S. and Wilkinson, M. 2015. Ichthyosaurs from the Jurassic of Skye, Scotland. *Scottish Journal of Geology*. Vol. 51(1), DOI: 10.1144/sjg2014-018
- Brusatte, S.L., Muir, A., Young, M.T., Walsh, S. Steel, L. and Witmer, L.M. 2016. The braincase and neurosensory anatomy of an Early Jurassic marine crocodylomorph: implications for crocodylian sinus evolution and sensory transitions. *The Anatomical Record*. Vol. 299, pp. 1511–1530
- Buckley, G.A., Brochu, C.A., Krause, D.W. and Pol, D. 2000. A pug-nosed crocodyliiform from the Late Cretaceous of Madagascar. *Nature*. Vol. 405, pp. 941–944
- Buffetaut, E.A. 1979. Jurassic Marine Crocodilians (Mesosuchia: Teleosauridae) from Central Oregon: First Record in North America. *Journal of Paleontology*. Vol. 53 (1), pp. 210–215

- 585
586 Burns, M.E. 2008. Taxonomic utility of ankylosaur (Dinosauria, Ornithischia) osteoderms:
587 *Glyptodontoelta mimus* Ford, 2000: a test case. *Journal of Vertebrate Paleontology*. Vol. 28
588 (4), pp. 1102–1109
589
- 590 Burns, M.E., Vickaryous, M.K. and Currie, P.J. 2013. Histological variability in fossil and
591 recent alligatoroid osteoderms: systematic and functional implications. *Journal of*
592 *Morphology*. Vol. 274 (6), pp. 676–686
593
- 594 Cau, A. and Fanti, F. 2011. The oldest known metriorhynchid crocodilian from the Middle
595 Jurassic of North-eastern Italy: *Neptunidraco ammoniticus* gen. et sp. nov. *Gondwana*
596 *Research*. Vol. 19, pp. 550–565
597
- 598 Cerda, I.A., Desojo, J.B., Trotteyn, M.J. and Scheyer, T.M. 2015. Osteoderm histology of
599 *Proterochampsia* and Doswelliidae (Reptilia: Archosauriformes) and their evolutionary and
600 paleobiological implications. *Journal of Morphology*. Vol. 276, pp. 385–402
601
- 602 Chiarenza, A.A., Foffa, D., Young, M.T., Insacco, G., Cau, A. Carnevale, G. and Catanzariti,
603 R. 2015. The youngest record of metriorhynchid crocodylomorphs, with implications for the
604 extinction of Thalattosuchia. *Cretaceous Research*. Vol. 56, pp. 698–616
605
- 606 Clarac, F., Souter, T., Cornette, R., Cubo, J. and de Buffrénil, V. 2015. A Quantitative
607 Assessment of Bone Area increase due to Ornamentation in the Crocodylia. *Journal of*
608 *Morphology*. Vol. 276, pp. 1183–1192
609
- 610 Clark, N.D.L., 1993. A new occurrence of Scottish plesiosaurian remains from the Isle of Skye.
611 *Scottish Journal of Geology*. Vol. 29, pp. 197–199
612
- 613 Clark, N.D.L. 2001. A thyreophoran dinosaur from the Early Bajocian (Middle Jurassic) of the
614 Isle of Skye, Scotland. *Scottish Journal of Geology*. Vol. 37 (1), pp. 19–26
615

- Clark, N.D.L. 2018. Review of the Dinosaur Remains from the Middle Jurassic of Scotland, UK. *Geosciences*. Vol. 8 (2), 53
- Crush, P.J. 1984. A late upper Triassic sphenosuchid crocodilian from Wales. *Palaeontology*. Vol. 27 (1), pp.131–157
- de Buffrenil, V., Farlow, J.O. and de Ricqlès, A. 1986. Growth and function of *Stegosaurus* plates: evidence from one histology. *Paleobiology*. Vol. 12 (4), pp. 459–473
- Delsett, L.L., Novis, L.K., Roberts, A.J., Koevoets, M.J., Hammer, Ø, Druckenmiller, P.S. and Hurum, J.H. 2015. The Slottsmøya marine reptile Lagerstätte: Depositional environments, taphonomy and diagenesis. *Geological Society Special Publication*. Vol. 434 (1), pp. 165–188
- dePolo, P.E., Brusatte, S.L., Challands, T.J., Foffa, D., Ross, D.A., Wilkinson, M. and Hongyu, Y. 2018. A sauropod-dominated tracksite from Rubha nam Brathairean (Brothers' Point), Isle of Skye, Scotland. *Scottish Journal of Geology*. Vol. 54 (1), pp. 1–12
- dePolo, P.E., Brusatte, S.L., Challands, T.J., Foffa, D., Wilkinson, M., Clark, N.D.L., Hoad, J., DaCosta Pereira, P.V.L.G., Ross, D.A. and Wade, T.J. 2020. Novel track morphotypes from new tracksites indicate increased Middle Jurassic dinosaur diversity on the Isle of Skye, Scotland. *PLoS ONE*. Vol. 15 (3): e0229640
- de Ricqlès, A., Padian, K., Knoll, F. and Horner, J.R. 2008. On the origin of high growth rates in archosaurs and their ancient relatives: complementary histological studies on Triassic archosauriforms and the problem of a 'phylogenetic signal' in bone histology. *Annales de Paléontologie*. Vol. 94, pp. 57–76.
- Dridi, J and Johnson, M.M. 2020. On a longirostrine crocodylomorph (Thalattosuchia) from the Middle Jurassic of Tunisia. *Geobios*. Vol. 56, pp. 95-106

Efimov, M.B. 1988. The fossil crocodiles and champsosaurides of Mongolia and the USSR. *Trudy Somestnaya Sovetsko-Mongolskya Paleontologicheskaya Expeditsiya*. Vol. 36, pp. 1-108 [in Russian]

Evans, S. E. and Waldman, M. 1996. Small reptiles and amphibians from the Middle Jurassic of Skye, Scotland. *Museum of Northern Arizona Bulletin*. Vol. 60, pp. 219–26

Ezcurra, M.D. 2016. The phylogenetic relationships of basal archosauromorphs, with an emphasis on the systematics of proterosuchian archosauriforms. *PeerJ*. Vol. 4(4): e1778.

Fara, E., Ouaja, M., Buffetaut, E. and Srarfi, D. 2002. First occurrences of thalattosuchian crocodiles in the Middle and Upper Jurassic of Tunisia. *Neues Jahrbuch für Geologie und Paläontologie*. Vol. 8, pp. 465–476

Farmer, C.G. and Sanders, K. 2010. Unidirectional airflow in the lungs of alligators. *Science*. Vol. 327, pp. 338–340

Fernández, M. and Gasparini, Z.B. 2000. Salt glands in a Tithonian metriorhynchid crocodyliform and their physiological significance. *Lethaia*. Vol. 33, pp. 269–276

Fernández, M., Carabajak, A.P., Gasparini, Z. and Chong, G. 2011. A metriorhynchid crocodyliform braincase from northern Chile. *Journal of Vertebrate Paleontology*. Vol. 31 (2), pp. 369-377

Foffa, D. and Young, M.T. 2014. The cranial osteology of *Tyrannoneustes lythrodectikos* (Crocodylomorpha: Metriorhynchidae) from the Middle Jurassic of Europe. *PeerJ*. 2: e608, DOI: <https://doi.org/10.7717/peerj.608>

Foffa, D., Young, M.T., Brusatte, S.L., Graham, M.R. and Steel, L. 2017. A new metriorhynchid crocodylomorph from the Oxford Clay Formation (Middle Jurassic) of

England, with implications for the origin and diversification of Geosaurini. *Journal of Systematic Palaeontology*. DOI: 10.1080/14772019.2017.1367730

Foffa, D., Young, M.T. and Brusatte, S.L. 2018a. Filling the Corallian gap: New information on Late Jurassic marine reptile faunas from England. *Acta Palaeontologica Polonica*. Vol. 63 (2), pp. 287–313

Foffa, D. Young, M.T., Stubbs, T.L., Dexter, K.G. and Brusatte, S.L. 2018b. The long-term ecology and evolution of marine reptiles in a Jurassic seaway. *Nature Ecology and Evolution*. Vol. 2, pp. 1548–1555

Foffa, D., Johnson, M.M., Young, M.T., Steel, L. and Brusatte, S.L. 2019. Revision of the Late Jurassic deep-water teleosauroid crocodylomorph *Teleosaurus megarhinus* Hulke, 1871 and evidence of pelagic adaptations in Teleosaroidea. *PeerJ*. Vol. 7, e6646

Galton, P.M. 1985. British Plated Dinosaurs (Ornithischia: Stegosauridae). *Journal of Vertebrate Paleontology*. Vol. 5 (3), pp. 211–254

Gasparini, Z., Vignaud, P. and Chong, G. 2000. The Jurassic Thalattosuchia (Crocodyliformes) of Chile: a paleobiogeographic approach. *Bulletin Société géologique de France*. Vol. 171 (6), pp. 657–664

Gasparini, Z., Cichowolski, M. and Lazo, D.G. 2005. First record of *Metriorhynchus* (Reptilia: Crocodyliformes) in the Bathonian (Middle Jurassic) of the Eastern Pacific. *Journal of Paleontology*. Vol. 79 (4), pp. 801–805

Godoy, P.L., Montefeltro, F.C., Norell, M.A. and Langer, M.C. 2014. An Additional baurusuchid from the Cretaceous of Brazil with evidence of interspecific predation among Crocodyliformes. *PLoS ONE*. Vol. 9 (5), e97138

Haddoumi, H., Allain, R., Meslouh, S., Metais, G., Monbaron, M., Pons, D., Rae, J-C., Vullo, R., Zouhri, S. and Geerbrant, E. 2016. Guelb el Ahmar (Bathonian, Anoual Syncline, eastern Morocco): First continental flora and fauna including mammals from the Middle Jurassic of Africa. *Gondwana Research*. Vol. 29 (1), pp. 290–319

Herrera, Y., Gasparini, Z. and Fernández, M.S. 2013a. A new Patagonian species of *Cricosaurus* (Crocodyliformes, Thalattosuchia): first evidence of *Cricosaurus* in Middle-Upper Tithonian lithographic limestone from Gondwana. *Palaeontology*. Vol. 56 (3), pp. 663–678

Herrera, Y., Fernández, M.S. and Gasparini, Z. 2013b. Postcranial skeleton of *Cricosaurus araucanensis* (Crocodyliformes: Thalattosuchia): morphology and palaeobiological insights. *Alcheringa: An Australasian Journal of Palaeontology*. Vol. 37 (3), pp. 285–298

Herrera, Y. 2015. Metriorhynchidae (Crocodylomorpha: Thalattosuchia) from Upper Jurassic-Lower Cretaceous of Neuquén Basin (Argentina), with comments on the natural casts of the brain. *Publicación Electrónica de la Asociación Paleontológica Argentina*. Vol. 15 (1), pp. 159–171

Huh, M., Lee, D-G., Kim, J-K., Lim-J-D. and Godefroit, P. 2011. A new basal ornithomimid dinosaur from the Upper Cretaceous. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*. Vol. 259, pp. 1–24

Irmis, R.B., Nesbitt, S.J. and Sues, H-D. 2013. Early Crocodylomorpha. *Geological Society, London, Special Publications*. Vol. 379, pp. 275–302, DOI: <https://doi.org/10.1144/SP379.24>

Johnson, M.M., Young, M.T., Steel, L. and Lepage, Y. 2015. *Steneosaurus edwardsi* (Thalattosuchia: Teleosauridae), the largest known crocodylomorph of the Middle Jurassic. *Biological Journal of the Linnean Society*. Vol. 115 (4), pp. 911–918

Johnson, M.M., Young, M.T., Steel, L., Foffa, D., Smith, S.A., Hua, S., Havlik, P., Howlett, E.A. and Dyke, G. 2017. Re-description of ‘*Steneosaurus*’ *obtusidens* Andrews, 1909, an unusual microphagous teleosaurids crocodylomorph from the Middle Jurassic of England. *Zoological Journal of the Linnean Society*. Vol.182 (2), pp. 385-418

Johnson, M.M., Young, M.T., Brusatte, S.L., Thuy, B. and Weis, R. 2018. A catalogue of teleosaurids (Crocodylomorpha: Thalattosuchia) from the Toarcian and Bajocian (Jurassic) of southern Luxembourg. *Historical Biology*. Vol. 31 (9), pp. 1179-1194, DOI: 10.1080/08912963.2018.1427090

Johnson, M.M., Young, M.T. and Brusatte, S.L. 2020a. Re-description of two contemporaneous mesorostrine teleosauroids (Crocodylomorpha: Thalattosuchia) from the Bathonian of England and insights into the early evolution of Machimosaurini. *Zoological Journal of the Linnean Society*. Vol. 189 (2), pp. 449-482

Johnson, M.M., Young, M.T. and Brusatte, S.L. 2020b. Emptying the wastebasket: a historical and taxonomic revision of the Jurassic crocodylomorph *Steneosaurus*. *Zoological Journal of the Linnean Society*. Vol. 189 (2), pp. 428-448

Johnson, M.M., Young, M.T., and Brusatte, S.L. 2020c. The phylogenetics of Teleosauroidea (Crocodylomorpha, Thalattosuchia) and implications for their ecology and evolution. *PeerJ* 8: e9808 DOI 10.7717/peerj.9808

Jouve, S., Mennecart, B., Douteau, J. and Jalil, N-E. 2016. The oldest durophagous Teleosauroid (Crocodylomorpha, Thalattosuchia) from the Lower Bathonian of Central High Atlas, Morocco. *Palaeontology*. Vol. 59 (6), pp. 863–876

Jouve, S. and Jalil, N-E. 2020. Paleocene resurrection of a crocodylomorph taxon: Biotic crises, climatic and sea level fluctuations. *Gondwana Research*. Vol. 85, pp. 1-18

- Joyce, W.G., Lucas, S.G., Scheyer, T.M., Heckert, A.B. and Hunt, A.P. 2009. A thin-shelled reptile from the Late Triassic of North America and the origin of the turtle shell. *Proceedings. Biological sciences*. Vol. 276 (1656), pp. 507–513
- Kim, J-K., Huh, M., Lee, S-G. and Kim, Y-J. 2011. Preliminary study on dinosaur rib microstructure by applying correlative microscopy techniques. *Geosciences Journal*. Vol. 15 (3), pp. 225–235
- Kuzmin, I.T., Skutschas, P.P., Grigorieva, O.I. and Krasnolutskaa, S.A. 2013. Goniopholidid Crocodylomorph from the Middle Jurassic Berezovsk Quarry Locality (Western Siberia, Russia). *Proceedings of the Zoological Institute RAS*. Vol. 317 (4), pp. 452–458
- Lang, J.W. 1987. Crocodilian Behaviour: Implications for Management. Wildlife Management: Crocodiles and Alligators. In: Webbs, G.J.W., Manolis, S.C., Whitehead, P.J, ed. *Wildlife Management: Crocodiles and Alligators*. Sydney: Surrey Beatty. pp. 273-294
- Legendre, L.J., Guénard, G., Botha-Brink, J and, Cubo, J. 2016. Palaeohistological evidence for ancestral high metabolic rate in archosaurs. *Systematic Biology*. Vol. 65, pp. 989–996, DOI: [doi:10.1093/sysbio/syw033](https://doi.org/10.1093/sysbio/syw033)
- Li J. 1993. A new specimen of *Peipehsuchus teleorhinus* from Ziliujing Formation of Daxion, Sichuan. *Vertebrata PalAsiatica*. Vol. 31, pp. 85–94.
- Maidment, S.C.R., Brassey, C. and Barrett, P.M. 2015. The Postcranial Skeleton of an Exceptionally Complete Individual of the Plated Dinosaur *Stegosaurus stenops* (Dinosauria: Thyreophora) from the Upper Jurassic Morrison Formation of Wyoming, U.S.A. *PLoS ONE*. Vol. 10 (10): e0138352
- Main, R.P., de Ricqlès, A., Horner, J.R. and Padian, K. 2005. The evolution and function of thyreophoran dinosaur scutes: implications for plate function in stegosaurs. *Paleobiology*. Vol. 31 (2), pp. 291–314

- Mannion, P.D., Benson, R.B.J., Carrano, M.T., Tennant, J.P., Judd, J. and Butler, R.J. 2015. Climate constrains the evolutionary history and biodiversity of crocodilians. *Nature Communications*. Vol. 6, 8438
- Martin, J.E. and Buffetaut, E. 2012. The maxillary depression of Pholidosauridae: an anatomical study. *Journal of Vertebrate Paleontology*. Vol. 32 (6), pp. 1442–1446
- Martin, J.E., Suteethorn, S., Lauprasert, K., Tong, H., Buffetaut, E., Liard, R., Salaviale, C., Deesri, U., Suteethorn, V., and Claude, J. 2018. A new freshwater teleosaurids from the Jurassic of northeastern Thailand. *Journal of Vertebrate Paleontology*. Vol. 38 (6), DOI: 10.1080/02724634.2018.1549059
- McGowan, C. and Motani, R. 2003. Handbook of Paleoherpetology, Part 8. Ichthyopterygia. Verlag Der Friedrich Pfeil. München
- Miller, H. 1858. The Cruise of the Betsey; or, a summer ramble among the fossiliferous deposits of the Hebrides with Rambles of a Geologist; or, ten thousand miles over the fossiliferous deposits of Scotland. Edinburgh: Thomas Constable and Co. Hamilton, Adams, and Co., London
- Morton, N. and Hudson, J.D. 1995. Field guide to the Jurassic of the isles of Raasay and Skye, Inner Hebrides, NW Scotland. In: Taylor, P.D. ed. 1995. *Field Geology of the British Jurassic*. London: Geological Society, London, pp. 209-280
- Morton, N. 2004. Chapter 8: The Hebrides Basin. In Simms, M. J., Childlaw, N., Page, K. N. and Morton, N. *British Lower Jurassic Stratigraphy*. GCR Vol. 30 (8), GCR ID: 249
- Motani, R. 2010. Warm-Blooded “Sea Dragons”?. *Science*. Vol. 328 (5984), pp. 1361-1362

Newton, R.B. 1893. On the discovery of a secondary reptile in Madagascar: *Steneosaurus baroni* (n. sp.); with a reference to some post Tertiary vertebrate remains from the same country recently acquired by the British Museum (Natural History). *The Geological Magazine*. Vol. 10 (5), pp. 193-198

Nunn, E.V., Price, G.D., Hart, M.B., Page, K.N. and Leng, M.J. 2009. Isotopic signals from Callovian-Kimmeridgian (Middle-Upper Jurassic) belemnites and bulk organic carbon, Staffin Bay, Isle of Skye, Scotland. *Journal of the Geological Society, London*. Vol. 166, pp. 633–641

O'Connor, P.M., Sertich, J.J.W., Stevens, N.J., Roberts, E.M., Gottfried, M.D., Hieronymus, T.L., Jinnah, Z.A., Ridgely, R., Ngasala, S.E. and Temba, J. 2010. The evolution of mammal-like crocodyliforms in the Cretaceous Period of Gondwana. *Nature*. Vol. 466 pp. 748–751

Ősi, A., Young, M.T., Galácz, A. and Rabi, M. 2018. A new large-bodied thalattosuchian crocodyliform from the Lower Jurassic (Toarcian) of Hungary, with further evidence of the mosaic acquisition of marine adaptations in Metriorhynchoidea. *PeerJ*. Vol. 6, e4668

Panciroli, E., Walsh, S., Fraser, N.C., Brusatte, S.L. and Corfe, I. 2017. A reassessment of the postcanine dentition and systematics of the tritylodontid *Stereognathus* (Cynodontia, Tritylodontidae, Mammaliaformes) from the Middle Jurassic of the United Kingdom. *Journal of Vertebrate Paleontology*. Vol. 37 (5), e1351448, DOI:[10.1080/02724634.2017.1351448](https://doi.org/10.1080/02724634.2017.1351448)

Panciroli, E., Benson, R.B.J. and Butler, R.J. 2018. New partial dentaries of amphitheriid mammal *Palaeoxonodon ooliticus* from Scotland, and posterior dentary morphology in early cladotherians. *Acta Palaeontologica Polonica*. Vol. 63 (2), pp. 197-206

Panciroli, E., Schultz, J.A. and Luo, Z-X. 2019. Morphology of the petrosal and stapes of *Borealestes* (Mammaliaformes, Docodonta) from the Middle Jurassic of Skye, Scotland. *Papers in Palaeontology*. Vol. 5 (1), pp. 139–156

- Parrilla-Bel, J., Young, M.T., Moreno-Azanza and M., Canudo, J.I. 2013. The first metriorhynchid crocodylomorph from the Middle Jurassic of Spain, with implications for evolution of the subclade Rhacheosaurini. *PLoS ONE*. Vol. 8 (1), e54275
- Pérez-García, A. 2012. The European Lower Cretaceous *Chitrasephalus dumonii* (Testudines: Cryptodira) and the diversity of a poorly known lineage of turtles. *Acta Palaeontologica Polonica*. Vol. 57, pp. 575–588
- Phansalkar, V.G., Sudha, G. and Khadkikar, A.S. 1994. Giant marine reptilian skulls from the Jurassic of Kachchh, Gujarat. *Current Science*. Vol. 67, pp. 460-461
- Pierce, S.E. and Benton, M.J. 2006. *Pelagosaurus typus* Bronn, 1841 (Mesoeucrocodylia: Thalattosuchia) from the Upper Lias (Toarcian Lower Jurassic) of Somerset, England. *Journal of Vertebrate Paleontology*. Vol. 26 (3), pp. 621–635
- Puértolas-Pascual, E., Canudo, J.I. and Sender, L.M. 2015. New material from a huge specimen of *Anteophthalmosuchus* cf. *esuchae* (Goniopholididae) from the Albian of Andorra (Teruel, Spain): phylogenetic implications. *Journal of Iberian Geology*. Vol. 41 (1), pp. 41–56
- Puértolas-Pascual, E. and Mateus, O. 2020. A three-dimensional skeleton of Goniopholididae from the Late Jurassic of Portugal: implications for the Crocodylomorpha bracing system. *Zoological Journal of the Linnean Society*. Vol. 189 (2), pp. 521-548
- Rauhut, O.W.M. 2005. Osteology and relationships of a new theropod dinosaur from the Middle Jurassic of Patagonia. *Palaeontology*. Vol. 48 (1), pp. 87–110
- Roberts, A.J., Druckenmiller, P.S., Cordonnier, B., Delsett, L.L. and Hurum, J.H. 2020. A new plesiosaurian from the Jurassic-Cretaceous transitional interval of the Slottsmøya Member (Volgian), with insights into the cranial anatomy of cryptoclidids using computed tomography. *PeerJ*. Vol. 8: e8652

889 Rogov, M.A., Zverkov, N.G., Zakharov, V.A. and Arkangelsky, M.S. 2018. Marine Reptiles
890 and Climates of the Jurassic and Cretaceous of Siberia. Stratigraphy and Geological
891 Correlation. Vol. 27 (4), pp. 398-423

892
893 Romer, A.S. 1956. Osteology of the Reptiles. Chicago: The University of Chicago Press.

894
895 Sachs, S., Johnson, M.M., Young, M.T. and Abel, P. 2019. The mystery of *Mystriosaurus*:
896 redescribing the poorly known Early Jurassic teleosaurids thalattosuchians *Mystriosaurus*
897 *laurillardii* and *Steneosaurus brevior*. *Acta Palaeontologica Polonica*. Vol. 64 (3), pp. 565–
898 579

899
900 Schwab., J.A., Young, M.T., Neenan, J.M., Walsh, S.A., Witmer, L.M., Herrera, Y., Allain, R.
901 Brochu, C.A., Choiniere, J.N., Clark, J.M., Dollman, K.N., Etches, S., Fritsch, G., Gignac,
902 P.M., Alexander, R., Sachs, S., Turner, A.H., Vignaud, P. Wilberg, E.W., Xu, X., Zanno, L.E.
903 and Brusatte, S.L. 2020. Inner ear sensory system changes as extinct crocodylomorphs
904 transitioned from land to water. *PNAS*. Vol. 117 (19), pp. 10422–10428

905
906 Schwarz, D. and Salisbury, S.W. 2005. A new species of *Theriosuchus* (Atoposauridae,
907 Crocodylomorpha) from the Late Jurassic (Kimmeridgian) of Guimarota, Portugal. *Geobios*.
908 Vol. 38 (6), pp. 779–802

909
910 Schwarz, D., Raddatz, M. and Wings, O. 2017. *Knoetschkesuchus langenbergensis* gen. nov.
911 sp. nov., a new atoposaurid crocodyliiform from the Upper Jurassic Langenberg Quarry (Lower
912 Saxony, northwestern Germany), and its relationships to *Theriosuchus*. *PLoS ONE*. Vol. 12
913 (2): e0160617

914
915 Schwarz-Wings, D., Klein, N., Neumann, C. and Resch, U. 2011. A new partial skeleton of
916 *Alligatorellus* (Crocodyliiformes) associated with echinoids from the Late Jurassic (Tithonian)
917 lithographic limestone of Kelheim, Germany. *Fossil Record*. Vol. 14 (2), 195–205

918

- Séon, N., Amiot, R., Martin, J.E., Young, M.T., Middleton, H. Fourrel, F., Picot, L., Valentin, X. and Lécuyer, C. 2020. Thermophysiologicals of Jurassic marine crocodylomorphs inferred from the oxygen isotope composition of their tooth apatite. *Philosophical Transactions of the Royal Society B*. Vol. 375: 20190139, DOI: <https://doi.org/10.1098/rstb.2019.0139>
- Seymour, R.S., Bennet-Stamper, C.L., Johnston, S.D., Carrier, D.R. and Grigg, G.C. 2004. Evidence for Endothermic Ancestors of Crocodiles at the Stem of Archosaur Evolution. *Physiological and Biochemical Zoology*. Vol. 77 (6), pp. 1051-1067
- Summers, A.P. 2005. Evolution: warm-hearted crocs. *Nature*. Vol. 434, pp. 833–834
- Tennant, J.P. and Manion, P.D. 2014. Revision of the Late Jurassic crocodyliform *Alligatorellus*, and evidence for allopatric speciation driving high diversity in western European atoposaurids. *PeerJ*. Vol. 2, e599
- Vickaryous, M.K. and Hall, B.K. 2008. Development of the dermal skeleton in *Alligator mississippiensis* (Archosauria, Crocodylia) with comments on the homology of osteoderms. *Journal of Morphology*. Vol. 269 (4), pp. 398-422
- Vickaryous, M.K. and Sire, J-Y. 2009. The integumentary skeleton of tetrapods: origin, evolution, and development. *Journal of Anatomy*. Vol. 214, pp. 441–464
- Waldman, M. and Savage, R.J. 1972. The first Jurassic mammal from Scotland. *Journal of the Geological Society, London*. Vol. 128, pp. 119–125
- Waldman, M. and Evans, S.E. 1994. Lepidosauromorph reptiles from the Middle Jurassic of Skye. *Zoological Journal of the Linnean Society*. Vol. 112 (1–2), pp. 135–150
- Walkden, G.M., Fraser, N.C. and Muir, J. 1987. A new specimen of *Steneosaurus* (Mesosuchia, Crocodylia) from the Toarcian of the Yorkshire coast. *Proceedings of the Yorkshire Geological Society*. Vol. 46, pp. 279–287

Westphal, F. 1961. Zur Systematik der Deutschen und Englischen Lias-Krokodilier. *Neues Jahrbuch Geologie und Paläontologie*. Vol. 113, pp. 207–218

Wilberg, E.W. 2015. A new metriorhynchoid (Crocodylomorpha, Thalattosuchia) from the Middle Jurassic of Oregon and the evolutionary timing of marine adaptations in thalattosuchian crocodylomorphs. *Journal of Vertebrate Paleontology*. Vol. 35 (2), e902846

Wills, S., Barrett, P.M. and Walker, A. 2014. New dinosaur and crocodylomorph material from the Middle Jurassic (Bathonian) Kilmaluag Formation, Skye, Scotland. *Scottish Journal of Geology*. Vol. 50 (2), pp. 183–190

Wilson, J.A., D’Emic, M.D., Ikejiri, T., Moacdieh, E.M. and Whitlock, J.A. 2011. A nomenclature for vertebral fossae in sauropods and other saurischian dinosaurs. *PLoS One*. Vol. 6 (2), e17114

Wilson, J.A., 2012. New vertebral laminae and patterns of serial variation in vertebral laminae of sauropod dinosaurs. *Contributions from the Museum of Paleontology, University of Michigan*. Vol. 32 (7), pp. 91–110

Wincierz, J. 1967. Ein *Steneosaurus* Fund aus dem nordwestdeutschen Oberen Lias. *Paläontologische Zeitschrift*. Vol. 41, pp. 60-72

Yi, H., Tennant, J.P., Young, M.T., Challands, T.J., Foffa, D., Hudson, J.D., Ross, D.A. and Brusatte, S.L. 2017. An unusual small-bodied crocodyliform from the Middle Jurassic of Scotland, UK, and potential evidence for an early diversification of advanced neosuchians. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*. Vol. 107, pp. 1–12

979 Young, M.T. and Andrade, M.B. 2009. What is *Geosaurus*? Redescription of *Geosaurus*
980 *giganteus* (Thalattosuchia: Metriorhynchidae) from the Upper Jurassic of Bayern, Germany.
981 *Zoological Journal of the Linnean Society*. Vol. 157 (3), pp. 551–585

982
983 Young, M.T., Brusatte, S.L. Ruta, M. and Andrade, M.B. 2010. The evolution of
984 Metriorhynchoidea (Mesoeucrocodylia, Thalattosuchia): an integrated approach using
985 geometric morphometrics, analysis of disparity, and biomechanics. *Zoological Journal of the*
986 *Linnean Society*. Vol. 158 (4), pp. 801–859

987
988 Young, M.T., Brusatte, S.L., Andrade, M.B., Desojo, J.B., Beatty, B.L., Steel, L., Fernández,
989 M.S. Sakamoto, M., Ignacio Ruiz-Omeñaca, J. and Schoch, R.R. 2012. The cranial osteology
990 and feeding ecology of the metriorhynchid crocodylomorph genera *Dakosaurus* and
991 *Plesiosuchus* from the Late Jurassic of Europe. *PLoS ONE*. Vol. 7(9), e44985, DOI:
992 <https://doi.org/10.1371/journal.pone.0044985>

993
994 Young, M.T., Andrade, M.B., Brusatte, S.L., Sakamoto, M. and Liston, J. 2013. The oldest
995 known metriorhynchid super-predator: a new genus and species from the Middle Jurassic of
996 England, with implications for serration and mandibular evolution in predacious clades.
997 *Journal of Systematic Paleontology*. Vol. 11 (4), pp. 475–513

998
999 Young, M.T. and Steel, L. 2014a. Evidence for the teleosaurids crocodylomorph genus
1000 *Machimosaurus* in the Kimmeridge Clay Formation (Late Jurassic) of England. *Historical*
1001 *Biology*. Vol.26 (4), pp. 472–479

1002
1003 Young MT, Hua S, Steel L, Foffa D, Brusatte SL, Thüring S, Mateus O, Ruiz- Omeñaca JI,
1004 Havlik P, Lepage Y and Andrade MB. 2014b. Revision of the Late Jurassic teleosaurid genus
1005 *Machimosaurus* (Crocodylomorpha, Thalattosuchia). *Royal Society Open Science*. 1:140222

1006
1007 Young, M.T., Tennant, J.P., Brusatte, S.L., Challands, T.J., Fraser, N.C., Clark, N.D.L. and
1008 Ross, D.A. 2016. The first definitive Middle Jurassic atoposaurid (Crocodylomorpha,

Neosuchia), and a discussion on the genus *Theriosuchus*. *Zoological Journal of the Linnean Society*. Vol. 176, pp. 443–462

Zanno, L.E., Drymala, S., Nesbitt, S.J. and Schneider, V.P. 2015. Early crocodylomorph increases top tier predator diversity during rise of dinosaurs. *Scientific Reports*. Vol. 5, 9276

Zverkov, N.G., Arkhangelsky, M.S., Pardo Pérez, J.M. and Beznosov, P.A. 2015a. On the Upper Jurassic ichthyosaur remains from the Russia. *Proceedings of the Zoological Institute RAS*. Vol. 319 (1), pp. 81–97

Zverkov, N.G., Arkhangelsky, M.S. and Stenshin, I.M. 2015b. A review of Russian Upper Jurassic ichthyosaurs with an intermedium/humeral contact. Reassessing *Grendelius* McGowan, 1976. *Proceedings of the Zoological Institute RAS*. Vol. 319 (4), pp. 558–588

Zverkov, N.G., Fischer, V., Madzia, D. and Benson, R.B.J. 2018. Increased pliosaurid dental disparity across the Jurassic-Cretaceous transition. *Palaeontology*. Vol. 61 (6), pp. 825–846

Zverkov, N.G. and Efimov, V.M. 2019a. Revision of *Undorosaurus*, a mysterious Late Jurassic ichthyosaur of the Boreal Realm. *Journal of Systematic Palaeontology*. Vol. 17 (14), pp. 963–993

Zverkov, N.G. and Prilepskaya, N.E. 2019b. A prevalence of *Arthropterygius* (Ichthyosauria: Ophthalmosauridae) in the Late Jurassic – earliest Cretaceous of the Boreal Realm. *PeerJ* Vol. 7, e6799

TABLES

Table 1: Summary of crocodylomorph specimens from the Middle Jurassic of Skye reported in the literature

FIGURE CAPTIONS

Figure 1: Map of Skye, Scotland, showing the geographical and geological context for the crocodylomorph specimen described here. Cross indicating location of Prince Charlie's Cave. Maps adapted from Brusatte et al. (2015) and Morton (2004).

Figure 2: (a) 3D model and (b) photograph of a cobble containing vertebrae, osteoderms, and ribs of a Teleosauroida indet. from the Jurassic of Skye, Scotland. Scale bar 5cm.

Figure 3: Teleosauroida indet. Vertebrae from the Jurassic of Skye, Scotland. A1–C6 series of three unassociated dorsal vertebrae from the Jurassic (NMS G.2020.11.1) in anterior, left lateral, posterior, right lateral, ventral, and dorsal views. Scale bar = 2cm
cent. = centrum; nc. = neural canal; ns. = neural spine; prz.= prezygapophyses; tp. = transverse process

Figure 4: Isolated Teleosauroida indet. osteoderms from the Jurassic of Skye, Scotland (NMS G.2020.11.1) in dorsal, ventral, left lateral, right lateral, anterior, posterior views. Scale bar = 2cm.

Figure 5: Isolated Teleosauroida indet. osteoderm fragments and rib from the Jurassic of Skye, Scotland (NMS G.2020.11.1). A–C osteoderms in dorsal and ventral views. D – rib in dorsal, ventral, left and right lateral views. Scale bar = 2cm.

Figure 6: Comparative plate of non-teleosauroid Jurassic crocodylomorph dorsal osteoderms. A – *Neosuchia* indet. (Wills et al., 2014) scale bar = 2cm; B – *Goniopholididae* indet. (Kuzmin et al., 2013) scale bar = 2cm; C – *Pelagosaurus typus* (MNHN.F RJN 463) scale bar = 2cm; D - *Magyarosuchus fitosi* (Ösi et al., 2018) scale bar = 5cm.

Figure 7: Comparative plate of Jurassic thalattosuchians dorsal osteoderms. A – *Plagiophthalmosuchus gracilirostris* (NHMUK PV OR 14792) scale bar = 2cm; B – *Mystriosaurus* sp. (SNHM-IG-008-R) scale bar = 2cm; C – *Platysuchus multiscrobiculatus* (SMNS 9930) scale bar = 5cm; D – *Macrospondylus bollensis* (SMNS 51984) scale bar = 2cm; E – *Teleosaurus* sp. (NHMUK PV R 119a) scale bar = 2cm; F – *Andrianavoay baroni* (NHMUK PV R 1999) scale bar = 2cm; G – *Charitomenosuchus leedsii* (NHMUK PV R 3806) scale bar = 2cm; H – *Neosteneosaurus edwardsi* (PETMG R178) scale bar = 5cm; I – *Lemmysuchus obtusidens* (NHMUK PV R 3168) scale bar = 2cm; J – *Mycterosuchus nasutus* (NHMUK PV R 2617) scale bar = 5cm (A-D = Early Jurassic; E-J = Middle Jurassic (E-F = Bathonian; G-J = Callovian)). Note that the positioning of (F) is unknown.

1086

1087

1088

Age	Geology	Collection	Material	Reference	Taxon
Bathonian	Kilmaluag Formation	NMS G. 1992.47.6	Disarticulate d partial postcranial skeleton	Evans and Waldman (1996)	Goniopholididae
Bathonian	Duntulum Formation	NMS G.2016.21.1	Incomplete left dentary and incomplete splenial	Yi et al. (2017)	Neosuchia cf. Hylaeochampsidae
Late Bajocian-Bathonian	Valtos Sandstone Formation	NMS G.2014.52	Anterior dentary	Young et al. (2016)	Theriosuchus
Bathonian	Kilmaluag Formation	NMS G.1992.47.5 1	Left pubis	Wills et al. (2014)	Indeterminate mesoeucrocodylian
Bathonian	Kilmaluag Formation	NHMUK R36713	Disarticulate d osteoderms and single tooth	Wills et al (2014)	Neosuchia indet.