THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Prenatal tobacco exposure and self-regulation in early childhood: Implications for developmental psychopathology

Citation for published version:

Wiebe, SA, Clark, CAC, De Jong, DM, Chevalier, N, Espy, KA & Wakschlag, LS 2015, 'Prenatal tobacco exposure and self-regulation in early childhood: Implications for developmental psychopathology', *Development and Psychopathology*, vol. 27, no. 2, pp. 397-409.
<https://doi.org/10.1017/S095457941500005X>

Digital Object Identifier (DOI):

[10.1017/S095457941500005X](https://doi.org/10.1017/S095457941500005X)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Development and Psychopathology

Publisher Rights Statement:

© Wiebe, S. A., Clark, C. A., De Jong, D. M., Chevalier, N., Espy, K. A., & Wakschlag, L. S. (2014). Prenatal tobacco exposure and self-regulation in early childhood: Implications for developmental psychopathology. *Development and Psychopathology*.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Prenatal tobacco exposure and self-regulation in early childhood: Implications for developmental
psychopathology

Sandra A. Wiebe

Department of Psychology and Centre for Neuroscience, University of Alberta

Caron A. C. Clark

Department of Psychology, University of Arizona

Desiree M. de Jong

Department of Psychology, University of Massachusetts Amherst

Nicolas Chevalier

Department of Psychology, University of Edinburgh

Kimberly Andrews Espy

Department of Psychology, University of Arizona, and Developmental Cognitive Neuroscience

Laboratory, University of Nebraska-Lincoln

Lauren Wakschlag

Department of Medical Social Sciences, Feinberg School of Medicine, and Institute for Policy

Research, Northwestern University

Acknowledgments

This work was supported by NIDA grants R21DA024769 to Sandra Wiebe, R01DA014661 to Kimberly Andrews Espy, and R01 DA023653 to Kimberly Andrews Espy and Lauren Wakschlag. We gratefully acknowledge the members of the Developmental Cognitive Neuroscience Laboratory for assistance with data collection and coding, and the families who made this research possible. Correspondence regarding this paper may be addressed to Sandra Wiebe, Department of Psychology, University of Alberta, P217 Biological Sciences Building, Edmonton, AB, Canada T6G 2E9, telephone (780) 492-2237, fax (780) 492-1768, electronic mail sandra.wiebe@ualberta.ca.

Prenatal tobacco exposure and self-regulation in early childhood: Implications for developmental psychopathology

Abstract

Prenatal tobacco exposure (PTE) has a well-documented association with disruptive behavior in childhood, but the neurocognitive effects of exposure that underlie this link are not sufficiently understood. The present study was designed to address this gap, through longitudinal follow-up in early childhood of a prospectively-enrolled cohort with well-characterized prenatal exposure. Three-year-old children ($n = 151$) were assessed using a developmentally sensitive battery capturing both cognitive and motivational aspects of self-regulation. PTE was related to motivational self-regulation, where children had to delay approach to attractive rewards, but not cognitive self-regulation, where children had to hold information in mind and inhibit prepotent motor responses. Furthermore, PTE predicted motivational self-regulation more strongly in boys than girls, and when propensity scores were covaried to control for confounding risk factors, the effect of PTE on motivational self-regulation was significant only in boys. These findings suggest that PTE's impact on neurodevelopment may be greater in boys than girls, perhaps reflecting vulnerability in neural circuits that subserve reward sensitivity and emotion regulation, and may also help to explain why PTE is more consistently related to disruptive behavior disorders than attention problems.

Prenatal tobacco exposure and self-regulation in early childhood: Implications for
developmental psychopathology

The impact of early experience on development is a central theme in the field of developmental psychopathology. Its role in canalizing behavior patterns was highlighted in the last Special Issue on “Neural Plasticity, Sensitive Periods, and Psychopathology” (Cicchetti & Tucker, 1994a, 1994b; Courchesne, Chisum, & Townsend, 1994; Fox, Calkins, & Bell, 1994). In the intervening two decades, there has been increasing recognition that the prenatal environment is a potent influence on adaptive and maladaptive behavior patterns (Glover, O’Connor, & O’Donnell, 2010; Schuetze, Eiden, Colder, Gray, & Huestis, 2011). One prenatal influence with documented long-term effects on children’s health is smoking during pregnancy (DiFranza, Aligne, & Weitzman, 2004). Although there is now general awareness of the risks of smoking during pregnancy, prenatal tobacco exposure (PTE) is still common. In 2010, 23.2% of women in the US smoked cigarettes immediately prior to or during pregnancy, and 10.7% of women smoked throughout pregnancy (Tong et al., 2013). Rates are even higher in some groups, particularly among pregnant women who are younger, more financially disadvantaged, less educated, and White or Native American.

PTE has been linked to higher levels of externalizing behavior across developmental periods, including atypical externalizing trajectories in early childhood (Wakschlag, Leventhal, Pine, Pickett, & Carter, 2006), and clinical patterns of disruptive behavior (i.e., oppositional defiant disorder (ODD) and conduct disorder (CD) (Cornelius, Goldschmidt, DeGenna, & Day, 2007; Huijbregts, Warren, de Sonnevile, & Swaab-Barneveld, 2008; Monuteaux, Blacker, Biederman, Fitzmaurice, & Buka, 2006; Orlebeke, Knol, & Verhulst, 1999; Robinson et al., 2010; Wakschlag & Hans, 2002; Wakschlag et al., 2011). There is also mixed support for an

association between PTE and attention deficit hyperactivity disorder (ADHD), where some studies have found an association (Cornelius et al., 2007; Froehlich et al., 2009; Keyes, Davey Smith, & Susser, 2014; Kotimaa et al., 2003; Nomura, Marks, & Halperin, 2010; Orlebeke et al., 1999; Robinson et al., 2010), and others have not (Ball et al., 2010; Day, Richardson, Goldschmidt, & Cornelius, 2000; Huijbregts, Séguin, Zoccolillo, Boivin, & Tremblay, 2007), or have found associations only in children with comorbid ODD (Wakschlag, Pickett, Kasza, & Loeber, 2006) or dopaminergic risk alleles (Becker, El-Faddagh, Schmidt, Esser, & Laucht, 2008; Neuman et al., 2007).

However, the underlying neurodevelopmental mechanisms that drive these behavioral problems are as yet unclear. Nigg and Casey (2005) suggested that the common thread among externalizing behavior disorders is impaired self-regulation, reflecting dysfunction in neural circuits including frontal cortex and striatum; other theories have similarly emphasized the role of early self-regulatory difficulties in the etiology of ADHD (Barkley, 1997; Castellanos, Sonuga-Barke, Milham, & Tannock, 2006), ODD (Matthys, Vanderschuren, & Schutter, 2013) and antisocial behavior (Moffitt, 1993). The goal of the present study was to examine the construct of self-regulation in young children with PTE, as a potential marker of a developmental trajectory at risk for elevated externalizing behavior. In light of sex differences in vulnerability to pre- and perinatal risk factors (Elsmén, Steen, & Hellström-Westas, 2004) and in risk for externalizing behavior (Card, Stucky, Sawalani, & Little, 2008; Gershon, 2002), we explored sex as a moderator.

The Emergence of Self-Regulation

The last two decades have been marked by substantial theoretical and methodological advances in our understanding of the early development of self-regulation (Carlson, 2005; Espy, Kaufmann, Glisky, & McDiarmid, 2001; Garon, Bryson, & Smith, 2008; Wiebe et al., 2011; Willoughby, Blair, Wirth, Greenberg, & Family Life Project Investigators, 2010). Unfolding self-regulation in infancy and early childhood can be characterized as a gradual shift from exogenous to endogenous control of behavior. In infancy, behavior is largely driven by environmental stimuli, where external stimuli capture the infant's attention and elicit prepotent responses (e.g., approach, distress). Children become increasingly capable of volitional action in the environment, for example choosing to direct their attention to less salient stimuli or select alternative responses to a situation (e.g., delaying gratification, shifting attention from a stressor), and thereby develop the capacity to adhere to caregiver demands and social norms (Kopp, 1982). This exogenous-endogenous transition is supported by changes in prefrontal circuitry, in particular refinement of long-range connections between prefrontal cortex and other cortical and subcortical areas (Huttenlocher & Dabholkar, 1997; Thatcher, 1994).

Recent models of self-regulation include cognitive, emotional, and motivational processes as mutually interactive and reciprocal (e.g., Bell & Deater-Deckard, 2007; Gross & Thompson, 2007; Posner & Rothbart, 2000). There is evidence for a meaningful distinction between contexts that primarily tax children's self-regulation capabilities due to cognitive load (e.g., holding information in mind, overcoming automatic responses; often termed "cool EF") relative to motivational load (e.g., regulating responses to a salient reward or distressing situation; "hot EF") (Hongwanishkul, Happaney, Lee, & Zelazo, 2005; Willoughby, Kupersmidt, Voegler-Lee, & Bryant, 2011). Neurologically, the ability to self-regulate in the face of cognitive

load appears to be based in dorsolateral prefrontal cortex (DLPFC), whereas motivationally loaded tasks place demands on orbitofrontal cortex (OFC) (Zelazo, Carlson, & Kesek, 2008).

Theoretical links between childhood self-regulation and externalizing outcomes, mentioned earlier, are supported by empirical evidence. There are group differences between children with ADHD and typically-developing controls in response inhibition (Berlin, Bohlin, Nyberg, & Janols, 2004; Oosterlaan, Logan, & Sergeant, 1998; Schoemaker et al., 2012; Schoemaker, Mulder, Deković, & Matthys, 2013) and delay tolerance (Bitsakou, Psychogiou, Thompson, & Sonuga-Barke, 2009); the former construct is encompassed by cognitive self-regulation, whereas the latter falls within the rubric of motivational self-regulation. Aggressive and disruptive behavior is also associated with self-regulation problems (Raaijmakers et al., 2008; Schoemaker et al., 2012). In community samples, cognitive self-regulation difficulties have likewise been found to predict higher levels of externalizing behavior (Berlin & Bohlin, 2002; Riggs, Blair, & Greenberg, 2003). In one of the few studies to directly compare cognitive and motivational self-regulation, Willoughby and colleagues (2011) found that motivational self-regulation uniquely predicted inattentive and hyperactive behavior, whereas cognitive self-regulation uniquely predicted academic outcomes. Examining distinct cognitive and motivational components of self-regulation in a PTE sample at risk for externalizing behavior problems may allow for more precise specification of the neurodevelopmental basis of these difficulties.

PTE and Emergent Self-Regulation

Based on evidence from animal studies, in which exposure can be experimentally manipulated, PTE is believed to impact dopaminergic, serotonergic, and noradrenergic systems (Dwyer, McQuown, & Leslie, 2009; Slotkin, Pinkerton, Tate, & Seidler, 2006). These systems in

turn support the networks involved in adaptive control of cognition, emotion, and behavioral responses, including the anterior cingulate, and specific regions in ventrolateral, dorsolateral, and orbitofrontal cortex (Nigg & Casey, 2005; Zelazo & Carlson, 2012). Neuroimaging findings in humans likewise show that PTE is related to abnormalities in neural systems involved in self-regulation, including anterior cingulate regions involved in response inhibition, and orbitofrontal regions involved in motivational self-regulation (Liu et al., 2013; Lotfipour et al., 2009; El Marroun et al., 2014). Therefore, a causal pathway linking PTE, disordered self-regulation, and externalizing behavior is plausible.

PTE is associated with neurobehavioral deficits in infancy, but it is unclear to what extent these deficits reflect problems with self-regulation. Although neonates and infants with PTE show problems with attention and distractibility (Espy et al., 2011; G. A. Richardson, Day, & Taylor, 1989; Wiebe, Fang, Johnson, James, & Espy, 2014), these differences likely reflect bottom-up rather than top-down processes based on the developmental timeline of the emergence of top-down control. In previous work with the present sample, for example, we found no relation between PTE and working memory or information processing at 6 months (Wiebe et al., 2014). PTE neonates also show heightened irritability and difficulty soothing (Jacobson, Fein, Jacobson, Schwartz, & Dowler, 1984; Stroud et al., 2009). PTE-related differences in irritability persist through the first year (Schuetze & Eiden, 2007; Wiebe et al., 2014). In two studies that incorporated physiological measurement tapping self-regulation during presentation of a stressor, PTE infants showed greater peak cortisol response at 7 months (Schuetze, Lopez, Granger, & Eiden, 2008) and maladaptive patterns of autonomic activity at 9 months (Schuetze, Eiden, Colder, Gray, & Huestis, 2013), indicating problems with emotion regulation late in the first year of life.

In early and middle childhood, measurement of self-regulation is more straightforward, and studies have found associations between PTE and poorer performance on tasks requiring the control of attention and inhibition of prepotent responses (Cornelius, Ryan, Day, Goldschmidt, & Willford, 2001; Mezzacappa, Buckner, & Earls, 2011; Noland et al., 2005; Wiebe et al., 2009; for exceptions see Derauf et al., 2012; Huijbregts, Warren, de Sonnevile, & Swaab-Barneveld, 2008). Fewer studies have examined effects of PTE on self-regulation under emotional or motivational load, but the limited evidence supports a connection. One study found that at age 3, PTE children were less likely to wait for a delayed reward, although these differences were no longer significant by the time children were 5 (Kelsey, Hoffman, Wiebe, James, & Espy, 2014). In another study conducted with school-aged children, those with PTE evidenced greater difficulty in coping with frustration while playing a computer game with randomly interposed delays, relative to their non-exposed peers (Huijbregts et al., 2008). Taken together, these studies highlight disturbances in self-regulation among children with PTE, although it is less clear whether motivational and cognitive aspects of self-regulation are differentially vulnerable and to what extent these aspects may portend different dimensions of externalizing behavior.

Taken together with the evidence linking PTE with externalizing behavior, correlations between PTE and key behavioral outcomes are well-established. However, a straightforward causal interpretation of these associations is not warranted. Studies using genetically informed designs have suggested that putative teratologic effects may be a marker for genetic risk for problem behavior (D'Onofrio et al., 2008; D'Onofrio, Van Hulle, Goodnight, Rathouz, & Lahey, 2012; Maughan, Taylor, Caspi, & Moffitt, 2004; Silberg et al., 2003), as smoking during pregnancy is in itself a maternal problem behavior (Eiden et al., 2011; McGrath et al., 2012). One shortcoming of these studies is their relatively crude exposure measurement, which has

often been retrospective, brief, and has relied solely on maternal self-report, which is subject to biases and non-disclosure. These studies have also typically lacked a developmental framework. An alternative approach to addressing potential confounding has been the development of statistical methods that can deal with these potential biases and confounding risks. One approach that is gaining in popularity is propensity scores (McCaffrey, Ridgeway, & Morral, 2004; Rosenbaum & Rubin, 1984). To estimate propensity scores, risk factors related to selection bias in observational studies are identified and combined into a single metric using logistic regression or generalized boosted models better suited for non-normally distributed variables. In the case of PTE, propensity scores are then incorporated into statistical models to adjust exposure differences via matching or as a covariate (Ellis, Berg-Nielsen, Lydersen, & Wichstrøm, 2012; Fang et al., 2010; Willoughby, Greenberg, Blair, Stifter, & Family Life Investigative Group, 2007). Using these more rigorous control methods, these studies have typically found that exposure effects on externalizing behavior persist (Ellis et al., 2012; Keyes et al., 2014; Paradis, Fitzmaurice, Koenen, & Buka, 2011). Thus, whether PTE has direct teratologic effects is as yet unclear: rigorous prospective, multi-method assessment of exposure and modeling of unfolding brain:behavior relations to PTE is key to advancing understanding of this issue.

Sex Differences in PTE Vulnerability

It is well-documented that boys are more vulnerable to risk factors before and at birth—for example, boys are more likely to be born preterm or low birth weight, and preterm-born males face greater health risks than their female counterparts (Elsmén et al., 2004; Moe & Slinning, 2001). As such, it makes sense to consider potential sex differences in vulnerability to PTE. There is some evidence that boys are more vulnerable: PTE's effect on birthweight is

greater in boys than girls (Tayie & Powell, 2012), and sex has been found to moderate PTE's effects on attention and irritability in infancy (Pickett et al., 2008; Schuetze et al., 2013, 2008; Willoughby et al., 2007; but for an exception see Wiebe et al., 2014). Several studies have found that PTE is more strongly associated with externalizing behavior in boys, in independent samples assessed in early childhood and preadolescence (Hutchinson, Pickett, Green, & Wakschlag, 2010; Wakschlag & Hans, 2002). Unfortunately, many studies have not analyzed or reported differences in PTE effects, disregarding or simply controlling for sex in their analyses.

Research in developmental psychopathology also highlights the importance of examining sex differences in pathways to externalizing behavior. First, there are large sex differences in the prevalence of externalizing behavior. For ADHD, boys outnumber girls 9:1 in clinically-referred samples, and 3:1 in community samples (Gershon, 2002). For ODD, one large study estimated prevalence was 2.4 times more common in boys than girls (Maughan, Rowe, Messer, Goodman, & Meltzer, 2004).

There are also indications that risk factors and trajectories leading to externalizing behavior differ between boys and girls. Keenan and Shaw (1997) proposed that girls' trajectories show greater discontinuity because they face greater expectations to suppress externalizing behavior and, because of their earlier development of language and prosocial skills, they are more responsive to these socialization pressures. Consistent with this hypothesis, in a longitudinal study that used behavior problems in preschool and the transition to school to predict later externalizing behavior problems, Mesman, Bongers, and Koot (2001) found greater continuity in boys' developmental trajectories, where behavior problems in preschool emerged as stronger predictors of externalizing problems in preadolescence for boys than for girls. In a study examining moderated mediation, Chang, Olson, Sameroff, and Sexton (2011) found that for

boys, hostile parenting in early childhood contributed to externalizing behavior at age 6 via self-regulation problems assessed at age 3, whereas for girls relations between parenting and externalizing behavior were not mediated by self-regulation. This evidence supports the importance of considering sex as a moderator of developmental trajectories related to PTE.

The Present Study

The overarching goal of the present study was to study the link between PTE and self-regulation in early childhood, to gain insight into how PTE confers increased risk for externalizing behavior. We addressed this question within the Midwestern Infant Development Study (MIDS) cohort. This cohort was enrolled to examine the effects of PTE, and as such participants were selected to oversample women who smoked and enroll nonsmokers from similar backgrounds (Espy et al., 2011). Prospective, repeated measurement of smoking during pregnancy was used to minimize recall bias, and maternal self-report of smoking was integrated with biomarkers (i.e., assessing levels of cotinine, a byproduct of nicotine, in biospecimens) to generate a best estimate indicator of PTE and avoid misclassification because of underreporting of smoking.

Self-regulation was assessed at age 3 years, adapting a battery of tasks previously used with this age group (Wiebe et al., 2011) to include measures of both cognitive and motivational self-regulation. Individual tasks were used as indicators of latent factors in a structural equation model (SEM). Individual self-regulation tasks are often unreliable measures of self-regulation because performance reflects not only variation in regulatory capacity, but also variation in the basic abilities required to complete the task (e.g., motor abilities, naming), known as the task

impurity problem (Miyake et al., 2000). SEM parses the common, self-regulatory variability shared across tasks from unique, task-specific variability.

We hypothesized that PTE would be related to poorer self-regulation in early childhood, and based on previous research we expected differences to be evident across cognitive and motivational domains. We also tested the hypothesis that boys would show greater vulnerability to PTE. Finally, we tested whether relations between PTE and self-regulation could be accounted for by confounding risk factors related to maternal risk for smoking during pregnancy, using a propensity score approach.

Method

Participants

Mother-child dyads included in the present study ($N = 151$ children; 72 girls, 79 boys) were recruited during pregnancy to study the effects of PTE on neonatal attention and irritability (Espy et al., 2011) and participated in a follow-up study when children were 3 years old ($M = 3$ years 7 days, $SD = 22$ days). Mothers reported that their children's racial or ethnic background was European American (52%), African American (24%), Hispanic or Latino (20%), Native American (1.3%) or more than one race (3.3%). Demographic information is summarized in Table 1.

The initial cohort included 369 mother-child dyads at two Midwestern study sites, a small city in Nebraska and a rural tri-county area in Illinois. Because cigarette smoking is associated with lower socioeconomic status, stratified enrollment procedures were used to minimize potentially confounding demographic differences between the smoking and non-smoking groups. Exclusionary criteria for mothers included binge drinking and illegal drug use, with the

exception of occasional marijuana use. Exclusionary criteria for infants were preterm birth (<35 weeks) and birth complications known to affect developmental outcome (e.g., neonatal seizures). Due to funding constraints that made a multi-site follow-up untenable, the 3-year follow-up was limited to the Nebraska site ($n = 198$). Of this sample, 14 dyads were not eligible to participate at the 3-year follow-up because the mother no longer had custody ($n = 5$), their families had moved internationally ($n = 3$), the mother or child had died ($n = 1$), the child had been diagnosed with a neurological/medical condition ($n = 1$), or the mother had requested not to be contacted for follow-up studies ($n = 4$). Twenty dyads did not participate because they declined to participate ($n = 4$), could not be located ($n = 9$) or could not be enrolled within the time window ($n = 7$). Of the 164 dyads (89%) who participated in the 3-year wave, 13 were excluded from the present analyses because they completed questionnaires and interviews but were unable to visit the lab and so did not provide self-regulation data ($n = 11$) or because propensity scores could not be calculated due to missing data on key predictor variables ($n = 2$). Thus, the analytic sample for the present study included 151 dyads. Dyads included in the final sample did not differ significantly from those not followed up or excluded in their prenatal exposure status, maternal education, race or ethnicity, or child sex.

Procedures

Prenatal recruitment has been described in detail elsewhere (Espy et al., 2011; Fang et al., 2010) and so only a brief overview is presented here. Women were recruited during pregnancy and provided written, informed consent. At 14 and 28 weeks and immediately after delivery, women completed a set of interviews and questionnaires regarding their smoking behavior and use of other substances, physical and mental health, stress levels, and social supports. Urine

samples from the mother (and, after birth, the infant) were obtained at each visit to verify self-reported smoking data.

At the 3-year follow-up, each child was tested individually in a developmental laboratory setting by a trained research assistant who was blind to prenatal exposure status. A battery of tasks that included measures of self-regulation was administered in three sessions, in a fixed order to ensure that any potential carry-over effects would be comparable, as is typical in individual differences research; tasks selected for the present analyses are described below. Adherence to experimental protocols was maintained by regular team meetings and reviews of session videorecordings. Upon completion of each follow-up wave, children received a small toy and mothers received a gift card as compensation.

Measures

Prenatal Tobacco Exposure. At 14 and 28 weeks gestation and at delivery, mothers provided a month-by-month report of the number of cigarettes smoked per day using a modified timeline-followback interview (Espy et al., 2011). Maternal urine samples and infant meconium were tested for cotinine, a metabolite of nicotine, by US Drug Testing Laboratories. Infants were classified as exposed if mothers endorsed smoking and/or if they had cotinine values over 50 ng/mL at any time during the pre- or perinatal period. In total, 56% of infants in the present sample were classified as exposed. Mean levels of exposure across pregnancy are summarized in Table 1.

Propensity scores. Propensity scores were used to statistically control for selection bias by adjusting for potentially confounding risk factors that may differ between women who do and do not smoke during pregnancy. Propensity scores reflect the estimated likelihood that a

participant will smoke during pregnancy derived statistically from maternal background variables. As previously reported in detail by Fang et al. (2010), maternal mental and physical health and demographic background variables were used to estimate the propensity scores. Propensity scores were estimated using a generalized boosted model (GBM), a non-parametric approach that is robust to situations where covariates may be non-normally distributed, their effects may be non-linear and non-additive, and multicollinearity or missing values may be present (Friedman, 2001; Imbens, 2003; McCaffrey et al., 2004). GBM was implemented using the “twang” package in R 2.8.1 (R Foundation for Statistical Computing, Vienna, Austria). All 42 variables used in propensity score estimation are reported in Fang et al. (2010). The variables that contributed most strongly to propensity score computation (and proportion of variance in propensity scores explained) included drinking in the first month of pregnancy (14.9%), education (9.8%), drinking at the last menstrual period (8.6%), age (7.7%), IQ (6.4%), and history of hyperactivity (5.8%).

3-year Self-Regulation. Tasks in the self-regulation battery placed demands on children’s ability to hold information in mind and inhibit a prepotent response (see Wiebe et al., 2011, for a detailed description of the battery). The *Big-Little Stroop* task measured children’s inhibitory control of distractor interference. Stimuli were line drawings of everyday objects containing smaller embedded pictures that either matched or conflicted in identity with the larger object. The dependent measure was accuracy on conflict trials. The *Preschool Go/No-Go* task assessed children’s ability to inhibit a prepotent motor response. Pictures of fish were displayed on a computer screen, and children were instructed “catch” them by responding within 1500 milliseconds on a button box. On less frequent “no-go” trials (25% of trials), a shark appeared and children were instructed to “let it go” by withholding their response. The dependent measure

was *d*-prime, or the standardized difference between the hit and false alarm rates (Stanislaw & Todorov, 1999). Response inhibition was also assessed with a computerized version of the *Shape School*. Children viewed red and blue cartoon shape characters and were instructed to name their colors only when characters had happy faces and to inhibit naming those with sad faces. The dependent measure was accuracy on inhibit trials. The *Delayed Alternation* task required children to hold previously rewarded locations in working memory. The hiding location of a food reward alternated between left and right sides after each correct retrieval, and trials were separated by a 10 second filled delay. The dependent measure was the proportion of correct responses. Finally, in the *Nebraska Barnyard* task, children had to remember a sequence of animal names and press corresponding colored buttons on a touch screen in the corresponding order; this task was simplified from the version in Wiebe et al. (2011) to include only four colored buttons to speed administration, as in previous studies 3-year-old children seldom advanced beyond two-item sequences. The dependent measure was a composite score reflecting the summed proportion correct responses at each span length.

Two tasks required children to regulate their behavior under conditions where reward was highly salient. Children's ability to wait for a delayed food reward was assessed in the *Snack Delay* task, in which children were instructed to stand still with their hands on a placemat marked with two handprints, without moving or talking, in front of a handful of M&M candies under a transparent cup during a four-minute delay. Performance was indexed by (1) a score indicating whether children ate the snack during the delay, and (2) a summary score reflecting children's compliance with task rules, where children's behavior was scored in each 5 second epoch (up to 3 points for standing still, keeping their hands on the mat, and remaining silent) summed across all epochs prior to children eating the snack or until the task ended at 240

seconds. Children's ability to comply with a directive in the face of temptation was assessed using the *Goody Shelf* task, administered as part of the Disruptive Behavior Diagnostic Observation Schedule (DB-DOS; Wakschlag et al., 2008). The examiner instructed the child to sit at a small table and then unveiled an appealing set of toys (e.g., flashing wand, jelly bean dispenser) on a small shelf, instructing the child that the toys were only for looking at. During the 5-minute delay, the child was given three crayons and paper with which to draw, and the examiner sat in the corner of the room. If children touched the toys, then the examiner provided series of increasingly supportive prompts (e.g., verbal reminders, moving the shelf). Each instance of touching the toys was coded for intensity (brief touches = 1; sustained touches = 2; sustained touches where child was resistant to examiner prompts = 3). The dependent measure was the total score.

For all self-regulation tasks except the Preschool Go/No-Go (which required a button-press response), a randomly-selected subset of sessions ($\geq 20\%$) were independently scored by a second coder. Inter-rater reliability was high ($M = 91\text{-}100\%$ for all tasks).

Statistical Methods

Descriptive statistics were calculated in SAS 9.2 (SAS Institute, Cary, NC). Structural equation modeling (SEM) was conducted in Mplus 7.11 (Muthén & Muthén, 2012), using full information maximum likelihood estimation to handle missing data. Model fit to the data was assessed using the chi-square (χ^2) statistic, where a non-significant χ^2 value signifies good fit. Because the χ^2 test is sensitive to small deviations from perfect fit, additional indices used for model evaluation included the root mean square error of approximation (*RMSEA*), where values less than .06 indicate good fit and values between .06 and .08 indicate acceptable fit; and the

comparative fit index (*CFI*), where values between .95 and 1.00 indicate good fit (Hu & Bentler, 1999; Kline, 2011).

SEM was used to test substantive hypotheses about PTE and self-regulation, and to assess gender as a moderator. First, the best-fitting model of self-regulation was determined using confirmatory factor analysis (CFA). Then, the relationship between PTE and self-regulation was assessed. Additional models were tested to (1) explore the effect of gender as a moderator by constraining paths to equality across boys and girls; and (2) examine whether associations were robust to confounding risk factors by adding propensity scores as a covariate.

Results

Descriptive Statistics

Children's performance on the self-regulation battery is summarized in Table 2, presented separately by prenatal exposure status and by sex. For all measures of self-regulation under motivational load, children with PTE performed significantly worse than their non-exposed peers. No measures of self-regulation under cognitive load differed between exposure groups (all $ps > 0.10$), and there were no significant sex differences in performance on individual self-regulation tasks (all $ps > 0.10$). Table 3 presents correlations among the self-regulation measures. Correlations among tasks indexing self-regulation under cognitive load were generally significant but tended to be small in magnitude. In contrast, correlations among tasks indexing self-regulation under motivational load were significant and moderate to large. Most correlations across domains of self-regulation were non-significant.

Data Reduction

Before conducting SEM analyses, several dependent measures (Goody Shelf and Snack Delay movement scores) were standardized to minimize the range of variance across indicators (Kline, 2011), and if necessary scores were reflected so that a higher score always represented better self-regulation. Next, the factor structure of the self-regulation battery was evaluated using CFA; indices of model fit and model comparisons are summarized in Table 5. Unsurprisingly given the observed pattern of correlations among tasks, the best-fitting model had two factors reflecting latent self-regulation under conditions of cognitive relative to motivational load. All factor loadings were statistically significant, and standardized factor loadings ranged from .41 to .56 for the cognitive load factor and .44 to .95 for the motivational load factor. Tests of measurement invariance supported invariance by sex and exposure at the configural, metric, and scalar levels. In analyses of invariance by sex, it was necessary to constrain the residual variance for the *Snack Delay* summary score indicator to zero in boys for model identification purposes. If metric and scalar invariance are supported, then one can validly compare the means of latent variables across groups (Kline, 2011; Vandenberg & Lance, 2000).

Does PTE Predict Self-Regulation in Early Childhood?

When self-regulation was regressed on PTE, children with PTE showed significantly poorer self-regulation under motivational load relative to non-exposed children ($b = -.60$, $SE = .19$, $p = .001$), but the groups did not differ in self-regulatory performance on tasks tapping cognitive load ($b = -.21$, $SE = .21$, $p = .33$). Latent variables are scaled in standard deviation units, so children with PTE performed over half a standard deviation below their non-exposed peers. This model showed good fit to the data, $\chi^2(25) = 31.90$, $p = .16$, $RMSEA = 0.043$, $CFI =$

0.967. The pattern of significant results was unchanged when propensity scores were added to the model as a covariate. Model fit remained adequate with the exception of a significant model chi-square test, $\chi^2(31) = 45.94, p = .04, RMSEA = 0.056, CFI = 0.933$.

Does Sex Moderate the Effect of PTE?

Multiple-group models were tested to examine whether PTE's relation to self-regulation differed between boys and girls. Based on measurement invariance testing, the residual variance for the *Snack Delay* summary score indicator was fixed to zero for boys in all models. PTE was a significant predictor of self-regulation under motivational load in boys ($b = -.95, SE = .22, p < .001$) and in girls ($b = -.44, SE = .22, p = .045$); PTE's effect was significantly greater in boys, $\chi^2(1) = 5.49, p = .02$. PTE's effect on self-regulation under cognitive load was non-significant ($b = -.16, SE = .22, p = .46$) and equivalent in boys and girls, $\chi^2(1) = 0.52, p = .47$. A model in which PTE's effect on cognitively-loaded self-regulation was constrained to be equal across sexes and its effect on motivationally-loaded self-regulation was free to vary showed good fit to the data, $\chi^2(68) = 78.46, p = .18, RMSEA = 0.045, CFI = 0.954$. When the propensity score covariate was added to this model, the pattern of findings changed, as depicted in Figure 1. For boys, PTE's effect on motivationally-loaded self-regulation was robust ($b = -1.61, SE = .39, p < .001$), such that PTE boys performed over 1.5 standard deviation units poorer below non-exposed boys after confounding risks were controlled. In girls, this effect was no longer significant ($b = -.56, SE = .43, p = .19$). PTE's effect on cognitively-loaded self-regulation remained insignificant ($b = .18, SE = .36, p = .61$). For this model, fit statistics indicated marginally acceptable fit to the data, $\chi^2(80) = 103.48, p = .04, RMSEA = 0.062, CFI = 0.903$.

Discussion

In this study, we examined whether smoking during pregnancy was related to self-regulation in early childhood. We predicted that children with PTE would show lower levels of self-regulation in both cognitive and motivational domains; this prediction was only partially supported, as PTE-related differences were present only in the motivational domain. We also predicted that the relation between PTE and self-regulation would be stronger in boys than in girls. This hypothesis was supported, as PTE's impact on motivational self-regulation was significantly greater for boys than for girls. Furthermore, when propensity scores were added as a covariate to determine whether these results could be explained by other risk factors that were confounded with smoking during pregnancy, PTE no longer significantly predicted motivational self-regulation in girls, whereas the effect in boys was unchanged.

Our finding that motivational but not cognitive self-regulation was related to PTE was unexpected, but is in fact consistent with a previous study by Huijbregts and colleagues (2008). Their study had a relatively small sample, assessed exposure via maternal retrospective self-report, and included only a single task indexing cognitive and motivational self-regulation; however, as in the present study, they found that PTE was related to problems with motivational self-regulation—in their study, a computerized measure of delay aversion—but not cognitive self-regulation, assessed via a sustained attention task. The present study replicates these earlier findings in a larger, prospective sample, with more sophisticated measurement of both prenatal exposure and childhood self-regulation, and controlling for confounding risks.

The dissociation between cognitive and emotional self-regulation observed in the present study may also contribute to our understanding of the pattern of associations between PTE and externalizing disorders in the literature. As reviewed earlier, research has found that PTE has a

robust association with disruptive behavior disorders (DBDs) such as ODD and CD, but inconsistent relations with ADHD. For example, Nigg and Breslau (2007) found that PTE directly predicted ODD, which in turn accounted for PTE's relation with CD. In contrast, PTE's relation with ADHD was confounded by other risk factors such as maternal psychopathology and socioeconomic status, and was also mediated by birth weight. The motivational underpinnings of DBDs are well-established (Matthys et al., 2013; White et al., 2013, 2014). For ADHD, motivational and emotional factors are less recognized as a core deficit, but they are implicated in a subset of individuals with the disorder. Notably, 25-45% of children with ADHD exhibit emotion dysregulation as well (Shaw, Stringaris, Nigg, & Leibenluft, 2014). In Sonuga-Barke's dual-pathway model of ADHD (e.g., Castellanos et al., 2006; Sonuga-Barke, Auerbach, Campbell, Daley, & Thompson, 2005), it is proposed that subgroups of children with ADHD are typified by different neurocognitive risk factors. Within this model, executive dysfunction (or disordered cognitive self-regulation) is a risk factor for inattentive symptoms, whereas problems with delay aversion (or disordered motivational self-regulation) are a risk factor for hyperactive symptoms. If, as the findings of the present study and that of Huijbregts et al. (2008) suggest, PTE has a selective effect on motivational self-regulation, then heterogeneity within the ADHD population may contribute to the inconsistency of findings with respect to the association between PTE and ADHD; however, further research is needed to test this hypothesis.

The specific relation between PTE and motivational self-regulation implicates brain networks involved in processing emotion and reward, including the OFC, ACC, amygdala, and dopamine neurons in the striatum and midbrain. Although no studies to date have examined neural correlates of PTE in early childhood, studies in adolescence have revealed structural differences in these regions (Lotfipour et al., 2009; Toro et al., 2008). In animal models of PTE,

researchers have found differences in neurodevelopment and gene expression of dopaminergic systems (Muneoka, Nakatsu, Fuji, Ogawa, & Takigawa, 1999; S. A. Richardson & Tizabi, 1994; see Dwyer et al., 2009 for a review). Functionally, PTE is associated with a decreased neural response to reward in adolescent humans (Müller et al., 2013) and rats (Franke, Park, Belluzzi, & Leslie, 2008). Interestingly, this profile of reduced reward sensitivity has been proposed to result in increased reward seeking, as individuals strive to reach an optimal level of arousal, thereby increasing risk for externalizing behavior (Matthys et al., 2013; Shaw et al., 2014).

It is important to consider the present results within a developmental context. Children in the present study were assessed in a period of early childhood marked by rapid growth in self-regulation. In tests of self-regulation, between their third and fourth birthdays children become less likely to respond impulsively and inaccurately (Lemmon & Moore, 2007; Wiebe, Sheffield, & Espy, 2012; Zelazo et al., 2008). Because perseveration and impulsive responding are prevalent in the age range of the present study, it is possible that PTE effects on cognitive self-regulation may have been masked but could become apparent with development, as trajectories of exposed and non-exposed children diverge. Such a pattern of findings was observed for externalizing behavior in the second year (Wakschlag, Leventhal, et al., 2006). It is also possible that the relation between PTE and motivational self-regulation may change with development, as observed in another recent study (Kelsey et al., 2014). Longitudinal follow-up of the present sample later in development will help to clarify relations between PTE, self-regulation early in the preschool years, and outcomes in later childhood.

Our ability to detect relations between PTE and self-regulation likely benefited from our strong measurement of self-regulation. We adopted a latent variable approach that resulted in a model with good fit to the data, separating regulatory contributors to task performance from

extraneous task-specific contributors. Our self-regulation battery included a broad array of tasks tapping cognitive self-regulation, across differing presentation and response formats. However, our three measures of motivational self-regulation were drawn from only two tasks that differed in appetitive stimulus (food vs. appealing toys) but were similar in other respects (requirement to delay gratification, suppress approach and engage in a less appealing alternative behavior). So far there is very little research on the factor structure of motivational self-regulation in early childhood. However, one study that was able to capture the distinction between cognitive and motivational processes measured motivational self-regulation using tasks that, like ours, required that children wait for a reward, and found that only the motivational factor predicted externalizing behavior (Willoughby et al., 2011). Another study assessed motivational self-regulation using tasks that involved a salient reward but not delay, and found that cognitive and motivational measures converged to a single factor with modest relations to externalizing behavior (Allan & Lonigan, 2011). More work is needed in this domain both to parse the relative contributions of reward, delay, and emotion to motivational self-regulation requirements and to specify relations between facets of self-regulation and key predictors and outcomes.

We explored sex as a potential moderator of PTE risk. Consistent with our predictions and with previous studies examining a variety of pre- and perinatal risk factors (Elsmén et al., 2004; Moe & Slinning, 2001), boys emerged as more vulnerable to PTE-related deficits. Prenatally-exposed boys had significantly more difficulty coping with a delay in which they were prohibited from eating a desired treat or playing with highly salient toys. It is important to note that in the sample as a whole, girls and boys were equally likely to eat the treat prematurely or approach the forbidden toys, but in girls these behaviors were less strongly related to PTE, and when confounding risks were controlled by covarying the propensity score, no significant

relation between PTE and motivational self-regulation in girls remained. This finding may suggest that the determinants of self-regulation differ between boys and girls; it is possible that, for PTE girls, other factors such as parenting, home environment characteristics, or genetic risk factors play a larger role that overshadows the risk conferred by prenatal exposure.

Beyond the limitations already discussed, several other limitations of the present study warrant mention. The present follow-up was only able to include a subset of the original MIDS cohort; if the full cohort had been included, then the study would have had more power to detect more complex interactions. As well, although extensive background information was collected on participating women and used in calculation of the propensity scores, there may remain unmeasured confounding that was not accounted for.

The present study builds on previous research, and raises questions that need to be followed up in future studies. Our finding of a selective association between PTE and motivational self-regulation in boys should be revisited in a larger sample, so that additional questions such as the role of parenting can be addressed. It will be important to examine a broader range of outcomes to understand whether girls are developmentally “immune” from PTE impact or whether they are affected but in other ways. The inclusion of genetic information (e.g., measured genotype; genetically informed designs) would be beneficial to further advance understanding of putative teratologic effects. In the present sample, further follow-up assessments are underway and will make it possible to examine whether motivational self-regulation at age 3 mediates the pathway from PTE to later disruptive behavior.

The relation between PTE and motivational self-regulation in boys was independent of a host of confounding factors, rigorously controlled using propensity score methods. The present findings advance the current state of knowledge via methodologic rigor in assessment of both

prenatal exposures and potential confounds, and in the developmental framing. Of course, the definitive answer to the question of whether PTE causes disruptions in self-regulation requires experimental investigation. D’Onofrio, Thapar, and their colleagues have done pioneering work in the use of quasi-experimental designs for this purpose (D’Onofrio et al., 2008, 2012; Rice et al., 2009). Combining such methods with high quality exposure measurement and nuanced examination of multi-level pathways is the crucial next step for bringing clarity to the ongoing debate about the putative behavioral teratologic effects of PTE.

PTE may be a particularly salient example of an experience that canalizes developmental trajectories by altering the ontogeny of neurophysiological systems during a sensitive period of development. In keeping with the hierarchical nature of development, these early disruptions then appear to be recapitulated as children face new developmental challenges, such as the need to acquire basic self-regulation of behavioral responses during early childhood. The present findings clearly demonstrate that self-regulatory difficulties are detectable in children with PTE as early as age 3 years.

However, this pathological progression is not inevitable, considering that each child’s developmental trajectory is influenced by a complex system comprised of multiple interacting risk and protective factors. One goal of the field of developmental psychopathology is to promote protective factors (e.g., environmental enrichment) to compensate for genetic or environmental risk factors (see Cicchetti & Curtis, 2006, for further discussion). Some have argued that early childhood is a sensitive period in the development of neural structures supporting self-regulation, marked by enhanced sensitivity to environmental influences (Center on the Developing Child, 2011; Zelazo & Carlson, 2012), although of course neural plasticity is present to varying degrees throughout the lifespan (Cicchetti & Tucker, 1994a; Kramer, Bherer,

Colcombe, Dong, & Greenough, 2004). Recent studies have demonstrated that self-regulatory abilities are enhanced in children at sociodemographic risk when they have access to enriched home environments (Nelson et al., 2014) or classroom contexts (Diamond, Barnett, Thomas, & Munro, 2007). Leveraging the enhanced plasticity of self-regulation conferred by this sensitive period by implementing interventions that target self-regulatory skills may be key to altering trajectories towards antisocial behavior in children with PTE.

References

- Allan, N. P., & Lonigan, C. J. (2011). Examining the dimensionality of effortful control in preschool children and its relation to academic and socioemotional indicators. *Developmental Psychology, 47*(4), 905–915.
- Ball, S. W., Gilman, S. E., Mick, E., Fitzmaurice, G., Ganz, M. L., Seidman, L. J., & Buka, S. L. (2010). Revisiting the association between maternal smoking during pregnancy and ADHD. *Journal of Psychiatric Research, 44*(15), 1058–1062.
- Barkley, R. A. (1997). Behavioral inhibition, sustained attention, and executive functions: constructing a unifying theory of ADHD. *Psychological Bulletin, 121*(1), 65–94.
- Becker, K., El-Faddagh, M., Schmidt, M. H., Esser, G., & Laucht, M. (2008). Interaction of dopamine transporter genotype with prenatal smoke exposure on ADHD symptoms. *The Journal of Pediatrics, 152*(2), 263–269.
- Bell, M. A., & Deater-Deckard, K. (2007). Biological systems and the development of self-regulation: Integrating behavior, genetics, and psychophysiology. *Journal of Developmental and Behavioral Pediatrics, 28*(5), 409–420.
- Berlin, L., & Bohlin, G. (2002). Response inhibition, hyperactivity, and conduct problems among preschool children. *Journal of Clinical Child Psychology, 31*(2), 242–251.
- Berlin, L., Bohlin, G., Nyberg, L., & Janols, L.-O. (2004). How well do measures of inhibition and other executive functions discriminate between children with ADHD and controls? *Child Neuropsychology, 10*(1), 1–13.
- Bitsakou, P., Psychogiou, L., Thompson, M., & Sonuga-Barke, E. J. S. (2009). Delay aversion in attention deficit/hyperactivity disorder: An empirical investigation of the broader phenotype. *Neuropsychologia, 47*(2), 446–456.

- Card, N. A., Stucky, B. D., Sawalani, G. M., & Little, T. D. (2008). Direct and indirect aggression during childhood and adolescence: A meta-analytic review of gender differences, intercorrelations, and relations to maladjustment. *Child Development, 79*(5), 1185–1229.
- Carlson, S. M. (2005). Developmentally sensitive measures of executive function in preschool children. *Developmental Neuropsychology, 28*(2), 595–616.
- Castellanos, F. X., Sonuga-Barke, E. J. S., Milham, M. P., & Tannock, R. (2006). Characterizing cognition in ADHD: Beyond executive dysfunction. *Trends in Cognitive Sciences, 10*(3), 117–123.
- Center on the Developing Child (2011). *Building the Brain’s “Air Traffic Control” System: How Early Experiences Shape the Development of Executive Function* (Working Paper No. 11). Boston, MA: Harvard University.
- Chang, H., Olson, S. L., Sameroff, A. J., & Sexton, H. R. (2011). Child effortful control as a mediator of parenting practices on externalizing behavior: Evidence for a sex-differentiated pathway across the transition from preschool to school. *Journal of Abnormal Child Psychology, 39*(1), 71–81.
- Cicchetti, D., & Curtis, W. J. (2006). The developing brain and neural plasticity: Implications for normality, psychopathology, and resilience. In D. Cicchetti & D. J. Cohen (Eds.), *Developmental Psychopathology. Vol. 2: Developmental Neuroscience* (2nd ed., pp. 1–64). Hoboken, NJ: Wiley.
- Cicchetti, D., & Tucker, D. (1994a). Development and self-regulatory structures of the mind. *Development and Psychopathology, 6*(4), 533–549.

- Cicchetti, D., & Tucker, D. (1994b, Eds.). Neural plasticity, sensitive periods, and psychopathology [Special Issue]. *Development and Psychopathology*, 6(4), 531–814.
- Cornelius, M. D., Goldschmidt, L., DeGenna, N., & Day, N. L. (2007). Smoking during teenage pregnancies: effects on behavioral problems in offspring. *Nicotine & Tobacco Research*, 9(7), 739–750.
- Cornelius, M. D., Ryan, C. M., Day, N. L., Goldschmidt, L., & Willford, J. A. (2001). Prenatal tobacco effects on neuropsychological outcomes among preadolescents. *Journal of Developmental & Behavioral Pediatrics*, 22(4), 217–225.
- Courchesne, E., Chisum, H., & Townsend, J. (1994). Neural activity-dependent brain changes in development: Implications for psychopathology. *Development and Psychopathology*, 6(4), 697–722.
- D’Onofrio, B. M., Van Hulle, C. A., Goodnight, J. A., Rathouz, P. J., & Lahey, B. B. (2012). Is maternal smoking during pregnancy a causal environmental risk factor for adolescent antisocial behavior? Testing etiological theories and assumptions. *Psychological Medicine*, 42(7), 1535–1545.
- D’Onofrio, B. M., Van Hulle, C. A., Waldman, I. D., Rodgers, J. L., Harden, K. P., Rathouz, P. J., & Lahey, B. B. (2008). Smoking during pregnancy and offspring externalizing problems: An exploration of genetic and environmental confounds. *Development and Psychopathology*, 20(1), 139–164.
- Day, N. L., Richardson, G. A., Goldschmidt, L., & Cornelius, M. D. (2000). Effects of prenatal tobacco exposure on preschoolers’ behavior. *Journal of Developmental and Behavioral Pediatrics*, 21(3), 180–188.

- Derauf, C., Lester, B. M., Neyzi, N., Kekatpure, M., Gracia, L., Davis, J., Kallianpur, K., et al. (2012). Subcortical and cortical structural central nervous system changes and attention processing deficits in preschool-aged children with prenatal methamphetamine and tobacco exposure. *Developmental Neuroscience*, 34(4), 327–341.
- Diamond, A., Barnett, W. S., Thomas, J., & Munro, S. (2007). Preschool program improves cognitive control. *Science*, 318(5855), 1387–1388.
- DiFranza, J. R., Aligne, C. A., & Weitzman, M. (2004). Prenatal and postnatal environmental tobacco smoke exposure and children's health. *Pediatrics*, 113(4), 1007–1015.
- Dwyer, J. B., McQuown, S. C., & Leslie, F. M. (2009). The dynamic effects of nicotine on the developing brain. *Pharmacology & Therapeutics*, 122(2), 125–139.
- Eiden, R. D., Leonard, K. E., Colder, C. R., Homish, G. G., Schuetze, P., Gray, T. R., & Huestis, M. A. (2011). Anger, hostility, and aggression as predictors of persistent smoking during pregnancy. *Journal of Studies on Alcohol and Drugs*, 72(6), 926–932.
- Ellis, L. C., Berg-Nielsen, T. S., Lydersen, S., & Wichstrøm, L. (2012). Smoking during pregnancy and psychiatric disorders in preschoolers. *European Child & Adolescent Psychiatry*, 21(11), 635–644.
- Elsmén, E., Steen, M., & Hellström-Westas, L. (2004). Sex and gender differences in newborn infants: Why are boys at increased risk? *Journal of Men's Health and Gender*, 1(4), 303–311.
- Espy, K. A., Fang, H., Johnson, C., Stopp, C., Wiebe, S. A., & Respass, J. (2011). Prenatal tobacco exposure: Developmental outcomes in the neonatal period. *Developmental Psychology*, 47(1), 153–169.

- Espy, K. A., Kaufmann, P. M., Glisky, M. L., & McDiarmid, M. D. (2001). New procedures to assess executive functions in preschool children. *The Clinical Neuropsychologist*, 15(1), 46–58.
- Fang, H., Johnson, C., Chevalier, N., Stopp, C., Wiebe, S. A., Wakschlag, L. S., & Espy, K. A. (2010). Using propensity score modeling to minimize the influence of confounding risks related to prenatal tobacco exposure. *Nicotine & Tobacco Research*, 12(12), 1211–1219.
- Fox, N. A., Calkins, S. D., & Bell, M. A. (1994). Neural plasticity and development in the first two years of life: Evidence from cognitive and socioemotional domains of research. *Development and Psychopathology*, 6(4), 677–696.
- Franke, R. M., Park, M., Belluzzi, J. D., & Leslie, F. M. (2008). Prenatal nicotine exposure changes natural and drug-induced reinforcement in adolescent male rats. *The European Journal of Neuroscience*, 27(11), 2952–2961.
- Friedman, J. H. (2001). Stochastic gradient boosting. *Computational Statistics and Data Analysis*, 29(5), 1189–1232.
- Froehlich, T. E., Lanphear, B. P., Auinger, P., Hornung, R., Epstein, J. N., Braun, J., & Kahn, R. S. (2009). Association of tobacco and lead exposures with attention-deficit/hyperactivity disorder. *Pediatrics*, 124(6), e1054–63.
- Garon, N., Bryson, S. E., & Smith, I. M. (2008). Executive function in preschoolers: A review using an integrative framework. *Psychological Bulletin*, 134(1), 31–60.
- Gershon, J. (2002). A meta-analytic review of gender differences in ADHD. *Journal of Attention Disorders*, 5(3), 143–154.
- Glover, V., O'Connor, T. G., & O'Donnell, K. (2010). Prenatal stress and the programming of the HPA axis. *Neuroscience and Biobehavioral Reviews*, 35(1), 17–22.

- Gross, J. J., & Thompson, R. A. (2007). Emotion regulation: Conceptual foundations. In J. J. Gross (Ed.), *Handbook of Emotion Regulation* (pp. 3–24). New York, NY: Guilford Press.
- Hongwanishkul, D., Happaney, K. R., Lee, W. S. C., & Zelazo, P. D. (2005). Assessment of hot and cool executive function in young children: Age-related changes and individual differences. *Developmental Neuropsychology*, 28(2), 617–644.
- Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling*, 6(1), 1–55.
- Huijbregts, S. C. J., Séguin, J. R., Zoccolillo, M., Boivin, M., & Tremblay, R. E. (2007). Associations of maternal prenatal smoking with early childhood physical aggression, hyperactivity-impulsivity, and their co-occurrence. *Journal of Abnormal Child Psychology*, 35(2), 203–215.
- Huijbregts, S. C. J., Warren, A. J., de Sonnevile, L. M. J., & Swaab-Barneveld, H. (2008). Hot and cool forms of inhibitory control and externalizing behavior in children of mothers who smoked during pregnancy: An exploratory study. *Journal of Abnormal Child Psychology*, 36(3), 323–333.
- Hutchinson, J., Pickett, K. E., Green, J., & Wakschlag, L. S. (2010). Smoking in pregnancy and disruptive behaviour in 3-year-old boys and girls: An analysis of the UK Millennium Cohort Study. *Journal of Epidemiology and Community Health*, 64(1), 82–88.
- Huttenlocher, P. R., & Dabholkar, A. S. (1997). Regional differences in synaptogenesis in human cerebral cortex. *The Journal of Comparative Neurology*, 387(2), 167–178.
- Imbens, G. (2003). Nonparametric Estimation of Average Treatment Effects Under Exogeneity: A Review. National Bureau of Economic Research.

- Jacobson, S. W., Fein, G. G., Jacobson, J. L., Schwartz, P. M., & Dowler, J. K. (1984). Neonatal correlates of prenatal exposure to smoking, caffeine, and alcohol. *Infant Behavior and Development*, 7(3), 253–265.
- Keenan, K., & Shaw, D. (1997). Developmental and social influences on young girls' early problem behavior. *Psychological Bulletin*, 121(1), 95–113.
- Kelsey, K. M., Hoffman, L., Wiebe, S. A., James, T. D., & Espy, K. A. (2014). Impact of prenatal tobacco exposure on developmental change in motor control and delayed gratification in preschoolers. *Manuscript under review*.
- Keyes, K. M., Davey Smith, G., & Susser, E. (2014). Associations of prenatal maternal smoking with offspring hyperactivity: Causal or confounded? *Psychological Medicine*, 44(4), 857–867.
- Kline, R. B. (2011). *Principles and practice of structural equation modeling* (3rd ed.). New York, NY: Guilford Press.
- Kopp, C. B. (1982). Antecedents of self-regulation: A developmental perspective. *Developmental Psychology*, 18(2), 199–214.
- Kotimaa, A. J., Moilanen, I., Taanila, A., Ebeling, H., Smalley, S. L., McGough, J. J., Hartikainen, A.-L., et al. (2003). Maternal smoking and hyperactivity in 8-year-old children. *Journal of the American Academy of Child & Adolescent Psychiatry*, 42(7), 826–833.
- Kramer, A. F., Bherer, L., Colcombe, S. J., Dong, W., & Greenough, W. T. (2004). Environmental influences on cognitive and brain plasticity during aging. *Journal of Gerontology: Medical Sciences*, 59(9), M940–M957.
- Lemmon, K., & Moore, C. (2007). The development of prudence in the face of varying future rewards. *Developmental Science*, 10(4), 502–511.

- Liu, J., Lester, B. M., Neyzi, N., Sheinkopf, S. J., Gracia, L., Kekatpure, M., & Kosofsky, B. E. (2013). Regional brain morphometry and impulsivity in adolescents following prenatal exposure to cocaine and tobacco. *JAMA Pediatrics*, 167(4), 348–354.
- Lotfipour, S., Ferguson, E., Leonard, G., Perron, M., Pike, B., Richer, L., Seguin, J. R., et al. (2009). Orbitofrontal cortex and drug use during adolescence: Role of prenatal exposure to maternal smoking and BDNF genotype. *Archives of General Psychiatry*, 66(11), 1244–1252.
- El Marroun, H., Schmidt, M. N., Franken, I. H. A., Jaddoe, V. W. V., Hofman, A., van der Lugt, A., Verhulst, F. C., et al. (2014). Prenatal tobacco exposure and brain morphology: A prospective study in young children. *Neuropsychopharmacology*, 39(4), 792–800. Nature Publishing Group.
- Matthys, W., Vanderschuren, L. J. M. J., & Schutter, D. J. L. G. (2013). The neurobiology of oppositional defiant disorder and conduct disorder: Altered functioning in three mental domains. *Development and Psychopathology*, 25(1), 193–207.
- Maughan, B., Rowe, R., Messer, J., Goodman, R., & Meltzer, H. (2004). Conduct disorder and oppositional defiant disorder in a national sample: Developmental epidemiology. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 45(3), 609–621.
- Maughan, B., Taylor, A., Caspi, A., & Moffitt, T. E. (2004). Prenatal smoking and early childhood conduct problems: Testing genetic and environmental explanations of the association. *Archives of General Psychiatry*, 61(8), 836–843.
- McCaffrey, D., Ridgeway, G., & Morral, A. (2004). Propensity score estimation with boosted regression for evaluating causal effects in observational studies. *Psychological Methods*, 9(4), 403–425.

- McGrath, L. M., Mustanski, B., Metzger, A., Pine, D. S., Kistner-Griffin, E., Cook, E., & Wakschlag, L. S. (2012). A latent modeling approach to genotype-phenotype relationships: Maternal problem behavior clusters, prenatal smoking, and MAOA genotype. *Archives of Women's Mental Health, 15*(4), 269–282.
- Mesman, J., Bongers, I. L., & Koot, H. M. (2001). Preschool developmental pathways to preadolescent internalizing and externalizing problems. *Journal of Child Psychology & Psychiatry, & Allied Disciplines, 42*(5), 679–689.
- Mezzacappa, E., Buckner, J. C., & Earls, F. (2011). Prenatal cigarette exposure and infant learning stimulation as predictors of cognitive control in childhood. *Developmental Science, 14*(4), 881–891.
- Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., Howerter, A., & Wager, T. D. (2000). The unity and diversity of executive functions and their contributions to complex “frontal lobe” tasks: A latent variable analysis. *Cognitive Psychology, 41*(1), 49–100.
- Moe, V., & Slinning, K. (2001). Children prenatally exposed to substances: Gender-related differences in outcome from infancy to 3 years of age. *Infant Mental Health Journal, 22*(3), 334–350.
- Moffitt, T. E. (1993). The neuropsychology of conduct disorder. *Development and Psychopathology, 5*(1-2), 135–151.
- Monuteaux, M. C., Blacker, D., Biederman, J., Fitzmaurice, G., & Buka, S. L. (2006). Maternal smoking during pregnancy and offspring overt and covert conduct problems: A longitudinal study. *Journal of Child Psychology and Psychiatry, and Allied Disciplines, 47*(9), 883–890.

- Müller, K. U., Mennigen, E., Ripke, S., Banaschewski, T., Barker, G. J., Büchel, C., Conrod, P., et al. (2013). Altered reward processing in adolescents with prenatal exposure to maternal cigarette smoking. *JAMA Psychiatry*, 70(8), 847–856.
- Muneoka, K., Nakatsu, T., Fuji, J.-I., Ogawa, T., & Takigawa, M. (1999). Prenatal administration of nicotine results in dopaminergic alterations in the neocortex. *Neurotoxicology and Teratology*, 21(5), 603–609.
- Muthén, L. K., & Muthén, B. O. (2012). *Mplus User's Guide* (7th ed.). Los Angeles, CA: Muthén & Muthén.
- Nelson, J. M., Choi, H.-J., Clark, C. A. C., James, T. D., Fang, H., Wiebe, S. A., & Espy, K. A. (2014). Sociodemographic risk and early environmental factors that contribute to resilience in executive control: A factor mixture model of 3-year-olds. *Child Neuropsychology*.
- Neuman, R. J., Lobos, E., Reich, W., Henderson, C. A., Sun, L.-W., & Todd, R. D. (2007). Prenatal smoking exposure and dopaminergic genotypes interact to cause a severe ADHD subtype. *Biological Psychiatry*, 61(12), 1320–1328.
- Nigg, J. T., & Breslau, N. (2007). Prenatal smoking exposure, low birth weight, and disruptive behavior disorders. *Journal of the American Academy of Child & Adolescent Psychiatry*, 46(3), 362–369.
- Nigg, J. T., & Casey, B. J. (2005). An integrative theory of attention-deficit/ hyperactivity disorder based on the cognitive and affective neurosciences. *Development and Psychopathology*, 17(3), 785–806.
- Noland, J. S., Singer, L. T., Short, E. J., Minnes, S., Arendt, R. E., Kirchner, H. L., & Bearer, C. (2005). Prenatal drug exposure and selective attention in preschoolers. *Neurotoxicology and Teratology*, 27(3), 429–438.

- Nomura, Y., Marks, D. J., & Halperin, J. M. (2010). Prenatal exposure to maternal and paternal smoking on attention deficit hyperactivity disorders symptoms and diagnosis in offspring. *Journal of Nervous and Mental Disease, 198*(9), 672–678.
- Oosterlaan, J., Logan, G. D., & Sergeant, J. A. (1998). Response inhibition in AD/HD, CD, comorbid AD/HD + CD, anxious, and control children: a meta-analysis of studies with the stop task. *Journal of Child Psychology and Psychiatry, and Allied Disciplines, 39*(3), 411–425.
- Orlebeke, J. F., Knol, D. L., & Verhulst, F. C. (1999). Child behavior problems increased by maternal smoking during pregnancy. *Archives of Environmental Health, 54*(1), 15–19.
- Paradis, A. D., Fitzmaurice, G. M., Koenen, K. C., & Buka, S. L. (2011). Maternal smoking during pregnancy and criminal offending among adult offspring. *Journal of Epidemiology and Community Health, 65*(12), 1145–1150.
- Pickett, K. E., Wood, C., Adamson, J., D’Souza, L., Desouza, L., & Wakschlag, L. S. (2008). Meaningful differences in maternal smoking behaviour during pregnancy: Implications for infant behavioural vulnerability. *Journal of Epidemiology and Community Health, 62*(4), 318–324.
- Posner, M. I., & Rothbart, M. K. (2000). Developing mechanisms of self-regulation. *Development and Psychopathology, 12*(3), 427–441.
- Raaijmakers, M. A. J., Smidts, D. P., Sergeant, J. A., Maassen, G. H., Posthumus, J. A., van Engeland, H., & Matthys, W. (2008). Executive functions in preschool children with aggressive behavior: Impairments in inhibitory control. *Journal of Abnormal Child Psychology, 36*(7), 1097–1107.

- Rice, F., Harold, G. T., Boivin, J., Hay, D. F., van den Bree, M., & Thapar, A. (2009). Disentangling prenatal and inherited influences in humans with an experimental design. *Proceedings of the National Academy of Sciences*, 106(7), 2464–2467.
- Richardson, G. A., Day, N. L., & Taylor, P. M. (1989). The effect of prenatal alcohol, marijuana, and tobacco exposure on neonatal behavior. *Infant Behavior and Development*, 12(2), 199–209.
- Richardson, S. A., & Tizabi, Y. (1994). Hyperactivity in the offspring of nicotine-treated rats: Role of the mesolimbic and nigrostriatal dopaminergic pathways. *Pharmacology, Biochemistry, and Behavior*, 47(2), 331–337.
- Riggs, N. R., Blair, C. B., & Greenberg, M. T. (2003). Concurrent and 2-year longitudinal relations between executive function and the behavior of 1st and 2nd grade children. *Child Neuropsychology*, 9(4), 267–276.
- Robinson, M., McLean, N. J., Oddy, W. H., Mattes, E., Bulsara, M., Li, J., Zubrick, S. R., et al. (2010). Smoking cessation in pregnancy and the risk of child behavioural problems: a longitudinal prospective cohort study. *Journal of Epidemiology & Community Health*, 64(7), 622–629.
- Rosenbaum, P. R., & Rubin, D. B. (1984). Reducing bias in observational studies using subclassification on the propensity score. *Journal of the American Statistical Association*, 79(387), 516–524.
- Schoemaker, K., Bunte, T., Wiebe, S. A., Espy, K. A., Deković, M., & Matthys, W. (2012). Executive function deficits in preschool children with ADHD and DBD. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 53(2), 111–119.

- Schoemaker, K., Mulder, H., Deković, M., & Matthys, W. (2013). Executive functions in preschool children with externalizing behavior problems: A meta-analysis. *Journal of Abnormal Child Psychology*, 41(3), 457–471.
- Schuetze, P., & Eiden, R. D. (2007). The association between prenatal exposure to cigarettes and infant and maternal negative affect. *Infant Behavior and Development*, 30(3), 387–398.
- Schuetze, P., Eiden, R. D., Colder, C. R., Gray, T. R., & Huestis, M. a. (2011). Physiological regulation in cigarette exposed infants: An examination of potential moderators. *Neurotoxicology and Teratology*, 33(5), 567–574.
- Schuetze, P., Eiden, R. D., Colder, C. R., Gray, T. R., & Huestis, M. A. (2013). Physiological regulation at 9 months of age in infants prenatally exposed to cigarettes. *Infancy*, 18(2), 233–255.
- Schuetze, P., Lopez, F. A., Granger, D. A., & Eiden, R. D. (2008). The association between prenatal exposure to cigarettes and cortisol reactivity and regulation in 7-month-old infants. *Developmental Psychobiology*, 50(8), 819–834.
- Shaw, P., Stringaris, A., Nigg, J., & Leibenluft, E. (2014). Emotion dysregulation in attention deficit hyperactivity disorder. *The American Journal of Psychiatry*, 171(3), 276–293.
- Silberg, J. L., Parr, T., Neale, M. C., Rutter, M., Angold, A., & Eaves, L. J. (2003). Maternal smoking during pregnancy and risk to boys' conduct disturbance: An examination of the causal hypothesis. *Biological Psychiatry*, 53(2), 130–135.
- Slotkin, T. A., Pinkerton, K. E., Tate, C. A., & Seidler, F. J. (2006). Alterations of serotonin synaptic proteins in brain regions of neonatal Rhesus monkeys exposed to perinatal environmental tobacco smoke. *Brain Research*, 1111(1), 30–35.

- Sonuga-Barke, E. J. S., Auerbach, J. G., Campbell, S. B., Daley, D., & Thompson, M. (2005). Varieties of preschool hyperactivity: Multiple pathways from risk to disorder. *Developmental Science*, 8(2), 141–150.
- Stanislaw, H., & Todorov, N. (1999). Calculation of signal detection theory measures. *Behavior Research Methods, Instruments, & Computers*, 31(1), 137–149.
- Stroud, L. R., Paster, R. L., Goodwin, M. S., Shenassa, E., Buka, S., Niaura, R., Rosenblith, J. F., et al. (2009). Maternal smoking during pregnancy and neonatal behavior: A large-scale community study. *Pediatrics*, 123(5), e842–e848.
- Tayie, F. A., & Powell, C. (2012). Sex differences in the association between prenatal smoking and decreased birthweight, and intensive health care of the neonate. *Behavioral Medicine*, 38(4), 138–142.
- Thatcher, R. W. (1994). Psychopathology of early frontal lobe damage: Dependence on cycles of development. *Development and Psychopathology*, 6(4), 565–596.
- Tong, V. T., Dietz, P. M., Morrow, B., D’Angelo, D. V, Farr, S. L., Rockhill, K. M., & England, L. J. (2013). Trends in smoking before, during, and after pregnancy - Pregnancy Risk Assessment Monitoring System (PRAMS), United States, 40 sites, 2000-2010. *Morbidity and Mortality Weekly Report*, 62(6), 1–19.
- Toro, R., Leonard, G., Lerner, J. V, Lerner, R. M., Perron, M., Pike, G. B., Richer, L., et al. (2008). Prenatal exposure to maternal cigarette smoking and the adolescent cerebral cortex. *Neuropsychopharmacology*, 33(5), 1019–1027.
- Vandenberg, R. J., & Lance, C. E. (2000). A review and synthesis of the measurement invariance literature: Suggestions, practices, and recommendations for organizational research. *Organizational Research Methods*, 3(1), 4–69.

- Wakschlag, L. S., & Hans, S. L. (2002). Maternal smoking during pregnancy and conduct problems in high-risk youth: A developmental framework. *Development and Psychopathology, 14*(2), 351–369.
- Wakschlag, L. S., Henry, D. B., Blair, R. J. R., Dukic, V., Burns, J., & Pickett, K. E. (2011). Unpacking the association: Individual differences in the relation of prenatal exposure to cigarettes and disruptive behavior phenotypes. *Neurotoxicology and Teratology, 33*(1), 145–154.
- Wakschlag, L. S., Hill, C., Carter, A. S., Danis, B., Egger, H. L., Keenan, K., Leventhal, B. L., et al. (2008). Observational assessment of preschool disruptive behavior. Part I: Reliability of the Disruptive Behavior Diagnostic Observation Schedule (DB-DOS). *Journal of the American Academy of Child and Adolescent Psychiatry, 47*(6), 622–631.
- Wakschlag, L. S., Leventhal, B. L., Pine, D. S., Pickett, K. E., & Carter, A. S. (2006). Elucidating early mechanisms of developmental psychopathology: The case of prenatal smoking and disruptive behavior. *Child Development, 77*(4), 893–906.
- Wakschlag, L. S., Pickett, K. E., Kasza, K. E., & Loeber, R. (2006). Is prenatal smoking associated with a developmental pattern of conduct problems in young boys? *Journal of the American Academy of Child and Adolescent Psychiatry, 45*(4), 461–467.
- White, S. F., Clanton, R., Brislin, S. J., Meffert, H., Hwang, S., Sinclair, S., & Blair, R. J. R. (2014). Temporal discounting and conduct disorder in adolescents. *Journal of Personality Disorders, 28*(1), 5–18.
- White, S. F., Pope, K., Sinclair, S., Fowler, K. A., Brislin, S. J., Williams, W. C., Pine, D. S., et al. (2013). Disrupted expected value and prediction error signaling in youths with disruptive

- behavior disorders during a passive avoidance task. *The American Journal of Psychiatry*, 170(3), 315–323.
- Wiebe, S. A., Espy, K. A., Stopp, C., Respass, J., Stewart, P., Jameson, T. R., Gilbert, D. G., et al. (2009). Gene-environment interactions across development: Exploring DRD2 genotype and prenatal smoking effects on self-regulation. *Developmental Psychology*, 45(1), 31–44.
- Wiebe, S. A., Fang, H., Johnson, C., James, K. E., & Espy, K. A. (2014). Determining the impact of prenatal tobacco exposure on self-regulation at 6 months. *Developmental Psychology*.
- Wiebe, S. A., Sheffield, T. D., & Espy, K. A. (2012). Separating the fish from the sharks: A longitudinal study of preschool response inhibition. *Child Development*, 83(4), 1245–1261.
- Wiebe, S. A., Sheffield, T., Nelson, J. M., Clark, C. A. C., Chevalier, N., & Espy, K. A. (2011). The structure of executive function in 3-year-olds. *Journal of Experimental Child Psychology*, 108(3), 436–452.
- Willoughby, M. T., Blair, C. B., Wirth, R. J., Greenberg, M., & Family Life Project Investigators (2010). The measurement of executive function at age 3 years: Psychometric properties and criterion validity of a new battery of tasks. *Psychological Assessment*, 22(2), 306–317.
- Willoughby, M. T., Greenberg, M., Blair, C. B., Stifter, C., & Family Life Investigative Group (2007). Neurobehavioral consequences of prenatal exposure to smoking at 6 to 8 months of age. *Infancy*, 12(3), 273–301.
- Willoughby, M. T., Kupersmidt, J., Voegler-Lee, M., & Bryant, D. (2011). Contributions of hot and cool self-regulation to preschool disruptive behavior and academic achievement. *Developmental Neuropsychology*, 36(2), 162–180.
- Zelazo, P. D., & Carlson, S. M. (2012). Hot and cool executive function in childhood and adolescence: Development and plasticity. *Child Development Perspectives*, 6(4), 354–360.

Zelazo, P. D., Carlson, S. M., & Kesek, A. (2008). The development of executive function in childhood. In C. A. Nelson & M. Luciana (Eds.), *Handbook of Developmental Cognitive Neuroscience* (2nd ed., pp. 553–576). Cambridge, MA: MIT Press.

Table 1

Descriptive statistics for demographic and exposure variables

Measure	PTE (<i>n</i> = 81)		NE (<i>n</i> = 65)	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
Infant sex (% female)	49.4%		56.1%	
Maternal education (years)	13.6	1.54	13.9	1.67
Child PPVT standardized score	94.6	13.64	96.9	15.76
Propensity score (estimated propensity to smoke during pregnancy; 0-1)***	0.76	0.189	0.30	0.169
Self-reported smoking (cigarettes/day):				
First trimester ***	3.96	6.56	0	0
Second trimester ***	3.83	6.75	0	0
Third trimester ***	3.13	6.26	0	0

Note. PPVT = Peabody Picture Vocabulary Test; PTE = prenatally tobacco-exposed; NE = non-exposed; *** $p < .001$;

Table 2

Descriptive statistics for measures of child self-regulation, by prenatal tobacco exposure status and child sex

Construct	Range	PTE		NE		Boys		Girls	
Dependent measure		(n = 51-80)		(n = 49-66)		(n = 52-76)		(n = 48-70)	
		<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
Cognitively Loaded Self-regulation									
Big-Little Stroop conflict trial accuracy	0.0 – 1.0	0.22	0.229	0.28	0.262	0.27	0.251	0.22	0.239
Delayed Alternation accuracy	0.0 - .94	0.51	0.159	0.50	0.211	0.51	0.183	0.49	0.186
Go/no-go d-prime	-1.4 – 3.1	0.54	0.979	0.52	0.993	0.63	1.039	0.43	0.913
Nebraska Barnyard composite score	0.6 – 8.1	3.2	1.76	3.5	1.71	3.4	1.71	3.2	1.78
Shape School Inhibit accuracy	0.0 – 1.0	0.33	0.223	0.39	0.301	0.38	0.291	0.34	0.234
Motivationally Loaded Self-regulation									
Goody Shelf rule-breaking ^a	0 - 33	4.8	7.84	2.2	6.20	4.5	8.40	2.7	5.62
Snack Delay ate treat ^a	0.0 – 1.9	0.43	0.497	0.20	0.406	0.35	0.479	0.31	0.467
Snack Delay movement score ^a	3 - 117	43.7	32.58	59.9	29.05	47.2	30.88	54.3	33.09

^a significant difference between exposure groups ($p < .05$). NE = non-exposed; PTE = prenatally tobacco-exposed

Table 3

Correlations among self-regulation measures

Task	2.	3.	4.	5.	6.	7.	8.
1. Big-Little Stroop	0.25**	0.23**	0.20*	0.30**	0.03	0.13	0.30***
2. Delayed Alternation	--	0.19*	0.30***	0.14	0.15+	0.07	0.14
3. Go/No-go		--	0.34***	0.29**	0.19*	0.08	0.10
4. Nebraska Barnyard			--	0.06	0.15+	0.12	0.22*
5. Shape School				--	0.10	0.12	0.22*
6. Goody Shelf (reversed)					--	0.37***	0.41***
7. Snack Delay (ate treat; reversed)						--	0.74***
8. Snack Delay (movement score)							--

Notes: To ease interpretation, scores were transformed if necessary so that higher scores represented better self-regulation for all measures. ⁺

p < .10; * p < .05; ** p < .01; *** p < .001.

Table 4

Model fit indices for alternative CFA models of self-regulation

Model (number of factors)	χ^2	<i>df</i>	<i>p</i>	<i>RMSEA</i>	<i>CFI</i>	<i>BIC</i>	Model comparison	$\Delta\chi^2$	<i>df</i>	<i>p</i>
1. Unitary self-regulation (2)	64.13	20	<.001	0.121	0.784	4279				
2. Cognitive and motivational load (2)	28.06	19	.082	0.056	0.956	4248	1 vs. <u>2</u>	36.07	1	<.001
3. Inhibition, working memory, and motivational load (3)	25.99	17	.075	0.059	0.956	4255	<u>2</u> vs. 3	2.07	2	.356
4. Cognitive and motivational load with correlated errors (2)	26.80	18	.083	0.057	0.957	4251	<u>2</u> vs. 4	1.26	1	.262

Notes: For model comparisons, the preferred model is underlined. Where two nested models showed equivalent fit to the data, the more parsimonious model was preferred.

Figure Captions

Figure 1: Path diagram illustrating the effect of prenatal tobacco exposure on self-regulation in boys (top) and girls (bottom). Both unstandardized and standardized (in parentheses) parameters are presented; error variances and the propensity score covariate are not shown. ** $p < .01$; *** $p < .001$.

Boys:

Girls: