THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Acute restraint stress does not alter corticosteroid receptor or 11-Hydroxysteroid dehydrogenase gene expression at Hypothalamic-Pituitary-Adrenal axis regulatory sites in captive male white-crowned sparrows (*Zonotrichia leucophrys gambelii*)

Citation for published version:

Krause, JS, Pérez, JH, Reid, A, Cheah, JC, Bishop, V, Wingfield, JC & Meddle, S 2021, 'Acute restraint stress does not alter corticosteroid receptor or 11-Hydroxysteroid dehydrogenase gene expression at Hypothalamic-Pituitary-Adrenal axis regulatory sites in captive male white-crowned sparrows (*Zonotrichia leucophrys gambelii*)', *General And Comparative Endocrinology*.
<https://doi.org/10.1016/j.ygcen.2020.113701>

Digital Object Identifier (DOI):

[10.1016/j.ygcen.2020.113701](https://doi.org/10.1016/j.ygcen.2020.113701)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

General And Comparative Endocrinology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

**Acute restraint stress does not alter corticosteroid receptors or 11 β -
hydroxysteroid dehydrogenase gene expression at hypothalamic-pituitary-
adrenal axis regulatory sites in captive male white-crowned sparrows
(*Zonotrichia leucophrys gambelii*)**

**Jesse S. Krause^{1,2}, Jonathan H. Pérez^{2,4,5}, Angus M.A. Reid^{5,6}, Jeffrey Cheah¹, Valerie
Bishop⁵, John C. Wingfield², and Simone L. Meddle⁵**

**¹Department of Biology, University of Nevada, Reno, 1664 N. Virginia Street, Reno, NV
89557, USA**

**²Department of Neurobiology, Physiology and Behavior, University of California Davis,
One Shields Avenue, Davis, CA 95616, USA**

**³Department of Biology, University of South Alabama, 5871 USA Dr. N. Room 124, Mobile,
AL 36688, USA**

**⁴Institute of Biodiversity, Animal Health & Comparative Medicine, University of Glasgow,
Glasgow G12 8QQ, Scotland, UK.**

**⁵The Roslin Institute, The Royal (Dick) School of Veterinary Studies, University of
Edinburgh, Easter Bush, Midlothian, EH25 9RG, Scotland, UK**

**⁶MRC Human Genetics Unit, Institute of Genetics & Molecular Medicine, University of
Edinburgh, EH4 2XU, Scotland, UK**

Running Title: Effects of acute restraint on gene expression

21 **This Manuscript has 18 pages, 2 tables and 2 figures**

22 **Corresponding author: Jesse S. Krause**

23 **Email: jskrause@unr.edu**

24

25 **Key words: acute stress, corticosterone, hypothalamic-pituitary-adrenal (HPA) axis,**
26 **corticosteroid, bird, corticosterone binding globulin (CBG), gonadotropin inhibitory**
27 **hormone (GnIH), 11 β Hydroxysteroid dehydrogenase (11 β HSD1, 11 β HSD2),**
28 **glucocorticoid receptor, mineralocorticoid receptor**

29

Abstract

Capture-restraint is often used to investigate the acute hypothalamic-pituitary-adrenal axis (HPA) response to stress in wild and captive animals through production of glucocorticoids. Although this approach is useful for understanding changes in glucocorticoids, it overlooks potential changes in the complex regulatory systems associated with the glucocorticoid response, including genomic receptors, steroid metabolizing enzymes, carrier proteins, and downstream target proteins (e.g. gonadotropin-inhibitory hormone; GnIH). The present study in captive male white-crowned sparrows (*Zonotrichia leucophrys*) tests the hypothesis that corticosteroid receptors (mineralocorticoid - MR and glucocorticoid - GR), 11 β -hydroxysteroid dehydrogenase 1 (11 β HSD1) and 2 (11 β HSD2), corticosteroid binding globulin (CBG), and GnIH undergo rapid changes in expression to mediate the glucocorticoid response to acute stress. To determine dynamic changes in gene mRNA expression in the hippocampus, hypothalamus, pituitary gland, and liver, birds were sampled within 3 minutes of entering the room and after 10, 30, and 60 minutes of capture restraint stress in a cloth bag. Restraint stress handling increased CBG and decreased GnIH mRNA expression in the liver and hypothalamus, respectively. MR, GR, 11 β HSD1, and 11 β HSD2 mRNA expression in the brain, pituitary gland and liver did not change. No correlations were found between gene expression and baseline or stress-induced plasma corticosterone levels. No rapid changes of MR, GR, 11 β HSD1, and 11 β HSD2 mRNA expression during a standardized acute restraint protocol suggests that tissue level sensitivity may remain constant during acute stressors. However, the observed rise in CBG mRNA expression could act to facilitate transport to target tissues or buffer the rise in circulating glucocorticoids. Further studies on tissue specific sensitivity are warranted.

53

54 **Introduction**

55 In vertebrates the hypothalamic-pituitary-adrenal (HPA) axis regulates the production of
56 glucocorticoid hormones, cortisol or corticosterone, that circulate through the blood stream
57 bound to corticosteroid binding globulin (CBG) which is produced by the liver (Schoech,
58 Romero et al. 2013). In response to a challenge or stressor, the activity of the HPA axis is
59 modulated, resulting in increased glucocorticoid production, which can be several orders of
60 magnitude higher than initial capture or baseline levels (de Kloet 2014). The actions of
61 glucocorticoids are mediated through two separate classes of receptors: the high affinity
62 mineralocorticoid receptor (MR) and the low affinity glucocorticoid receptor (GR) (de Kloet
63 2014). The difference in receptor affinity allows for primary activation of the MR at low levels
64 of glucocorticoids, often referred to as ‘baseline’. GR is activated as glucocorticoid levels elevate
65 above baseline in response to a challenge or stressor, which results in behavioral and
66 physiological adjustments (Joëls, Karst et al. 2008, de Kloet 2014). The cellular levels of
67 glucocorticoids are further affected by the enzymes 11 β -hydroxysteroid dehydrogenase 1
68 (11 β HSD1) and 2 (11 β HSD2). 11 β HSD2 deactivates corticosterone by metabolizing it to a less
69 biologically active glucocorticoid, while this process is reversed by 11 β HSD1 (Seckl and
70 Chapman 1997). Thus, these enzymes act antagonistically to one another to regulate intracellular
71 corticosterone levels which has direct effects on receptor activation. In addition, there is
72 evidence that CBG can mediate glucocorticoid signaling by either facilitating the delivery of
73 steroids to their target tissues, reducing the bioavailability by binding steroids at the target tissue,
74 or controlling the amount of steroid that is free in the plasma (Hammond, Smith et al. 1991,
75 Grasa, Cabot et al. 2001, Breuner and Orchinik 2002, Schoech, Romero et al. 2013, Sivukhina

and Jirikowski 2014, Gulfo, Ledda et al. 2016, Gulfo, Castel et al. 2019). Ultimately, all these factors combine to determine the rate at which MR and/or GR are activated, thus affecting stress physiology.

The hippocampus, the paraventricular nucleus (PVN) of the hypothalamus and the anterior pituitary gland are the major control sites for HPA axis function in mammals (Dallman, Akana et al. 1992, Joëls and Kloet 2017) and birds (Bouillé and Baylé 1975, Bouillé and Baylé 1976, Hodgson, Meddle et al. 2007, Dickens, Romero et al. 2009, Krause, McGuigan et al. 2015, Smulders 2017, Cornelius, Perreau et al. 2018). Through the interactions of MR and GR, the hippocampus controls the amplitude and timing of circadian rhythms for baseline glucocorticoids (Dallman, Akana et al. 1992, Sapolsky, Romero et al. 2000, Smulders 2017). Stress-induced levels of glucocorticoids are thought to provide negative feedback at the level of the PVN and anterior pituitary gland primarily through GR (Sapolsky, Romero et al. 2000, Landys, Ramenofsky et al. 2006). Pharmacological or genetic manipulations of MR, GR, 11 β HSD1 and 11 β HSD2 at each or all levels of the HPA axis can affect baseline and/or stress-induced plasma glucocorticoids, which suggests a complex and coordinated signaling cascade (Harris, Kotelevtsev et al. 2001, Kolber, Wiczorek et al. 2008, Schmidt, Sterlemann et al. 2009, Harris, Holmes et al. 2013, Pérez, Swanson et al. 2020).

Field biologists often assess acute HPA axis function by using a standard capture restraint protocol for up to 60 minutes and taking serial blood samples to measure circulating glucocorticoids. However, it remains unknown if the corticosteroid receptors or glucocorticoid metabolizing enzymes also change within this standard 60-minute restraint stress protocol. We hypothesized that the regulatory components of the HPA axis undergo rapid changes in expression in order to potentiate glucocorticoid signaling. In order to determine the rapidity at

which gene expression of the regulatory elements of the HPA axis may change, tissues from male captive white-crowned sparrows were sampled within three minutes of entering the room and after 10, 30, and 60 minutes of restraint stress. We predicted that gene expression for MR, GR, and 11 β HSD1 would be upregulated and 11 β HSD2 downregulated in response to stress in the hippocampus, hypothalamus, pituitary gland, and liver. In addition, we quantified gene expression of two known downstream targets of glucocorticoids; CBG and gonadotropin inhibitory hormone (GnIH) (Calisi, Rizzo et al. 2008, Kirby, Geraghty et al. 2009). We predicted that hepatic CBG expression would be upregulated to buffer the increase in plasma glucocorticoids induced by restraint stress. We also predicted upregulation of gonadotropin-inhibitory hormone (GnIH), because it is elevated by increased GR activation and plays a role in suppressing reproductive pathways (Kirby, Geraghty et al. 2009, Calisi, Díaz-Muñoz et al. 2011, Ernst, Lynn et al. 2016).

Methods

Birds

A total of 35 male Gambel's white-crowned sparrows, (*Zonotrichia leucophrys gambelii*), were caught using mist nets and baited potter traps in late October through early November 2015 in the vicinity of Davis, California (38.5449° N, 121.7405° W). Birds were housed in individual cages (35 cm (w) x 40 cm (l) x 45 cm (h)) in three separate indoor aviaries at 20 $\pm$ 2°C. A diet of 60% Mazuri maintenance chow and 40% wild bird seed and water were provided *ad libitum*. Birds were exposed to naturally changing photoperiod for Davis, CA, USA (38.55° N, -121.74° E) during the acclimation and experimentation period. Increasing natural photoperiod results in the activation of physiological cascades for migration and breeding (Farner and Follett 1966). Increasing photoperiod in captive Gambel's white-crowned sparrows

does not significantly affect HPA axis activity as they do not undergo the seasonal changes in baseline and stress-induced corticosterone observed in free-living individuals (Romero and Wingfield 1999, Ramenofsky, Campion et al. 2017).

Blood and Tissue collection

The effect of restraint stress on plasma levels of glucocorticoids was determined on a mid-March photoperiod 12L:12D (lights on: 07:00 and off 19:00) on separate days from tissue collection. Approximately 60µL of blood was collected following venipuncture of the alar vein with a 26-gauge needle and collected into a heparinized glass microcapillary tubes (VWR: 15401-56). Samples were collected within 3 minutes of entering the room and 30 minutes post restraint in an opaque cloth bag. Birds were bled at 09:00 over two consecutive days. Blood was centrifuged at 15,000 *g* for 5 minutes to separate plasma from red blood cells. The plasma was aspirated with a Hamilton syringe, transferred into a labeled microcentrifuge tube, and stored at -30°C until radioimmunoassay for corticosterone.

Birds were euthanized in early July when the photoperiod was 14L:10D (lights on: 07:00 and off 21:00) to ensure they were in a fully reproductive state which was confirmed by assessment of an enlarged cloacal protuberance and large testes size. As determined at the time of euthanasia, the testes had an average length of 6.96 ± 1.28 mm and width of 5.2 ± 1.12 mm ($n=35$; mean $\pm$ S.D.) and there was no evidence of body or wing feather molt. Birds were sampled between the hours of 09:00 and 11:00 over two consecutive days and were randomly assigned to four groups on each day; a baseline sampled in under 3 minutes ($n=8$), and three other groups in which the bird was placed into a cloth bag for 10 ($n=8$), 30 ($n=9$) or 60 ($n=9$) minutes. The time points were chosen as they represent the standardized capture restraint time

intervals at which blood samples are collected in stress physiology studies. Birds were euthanized by sedation with isoflurane followed by rapid decapitation. Birds were dissected and tissues were quickly collected and fresh frozen on powdered dry ice, wrapped in aluminum foil and placed in labeled plastic bags and stored at -80°C. In the case of the liver, the left lobe was collected. Samples were later shipped on dry ice to the Roslin Institute, University of Edinburgh where they were stored at -80°C until cDNA preparation. All work was approved by University of California, Davis - Institutional Animal Care and Use Committee (IACUC) under protocol 19758, United States Fish and Wildlife Service permits - Federal MB90026B-0, and California State SC13449.

Preparation of cDNA

Whole frozen brains were warmed to -18°C in a cryostat and the hippocampus (~30mg) and hypothalamus (~80-100mg) were dissected in standardized blocks for isolation of RNA. The hippocampus from both hemispheres was isolated from the dorsal aspect of the telencephalon. Hypothalamic blocks were created by placing the brain into a brain matrix and landmarks were used according to Stokes, Leonard et al. (1974) to make a coronal cut just caudal to the Tractus septomesencephalicus (TrSM) and a rostral coronal cut at the start of the cerebellum. Tissues were homogenized using an Ultra Turrax homogenizer (IKA-Werke GmbH & Co. KG) in conjunction with 1000µL of trizol (Sigma-Aldrich T9424), except for the pituitary gland in which 300µL was used due to the small mass (<3mg). Total RNA was extracted from 250µL of tissue homogenate using Zymo Research Direct-zol 96 well plates (R2057) including the DNA digestion step per the manufacturer's instructions. Total RNA for hippocampus, hypothalamus, pituitary gland, and liver were assessed using a Nanodrop 1000 (Thermo Scientific) and found to have an average yield of 261 ± 112 , 254 ± 126 , 26 ± 10 , and 219 ± 89 ng/µL (mean±SD) with an

average 260/280 ratio of 2.05 ± 0.05 , 2.02 ± 0.02 , 2.06 ± 0.16 , and 2.08 ± 0.02 (mean $\pm$ SD), respectively. cDNA was produced by reverse transcribing 250 ng of RNA per sample using the High Capacity Reverse Transcription Kit (Applied Biosystems) according to the manufacturer's guidelines and the final volume was adjusted to 110 μ L per sample using ultra-pure (Milli-Q) water.

Quantitative polymerase chain reaction (qPCR)

All assays used Brilliant 111 Ultra-fast SYBR Green qPCR Master mix (Agilent Technologies 60083) which was read on an Agilent Mx3005p qPCR system and processed with MxPro software (Agilent Technologies) as previously described (Reid, Wilson et al. 2017). In brief, a 20 μ L total reaction volume per sample was generated by combining 10 μ L SYBR Green, 8 μ L cDNA, 0.4 μ L 20 μ M forward primer, 0.4 μ L 20 μ M reverse primer, 0.3 μ L 1/500 dilution of ROX reference dye solution, and 0.9 μ L Milli-Q H₂O. Each assay used the following thermal conditions: 50°C; 120s, 95°C; 120s, (40 cycles of 95°C; 15s, 60°C; 30s), then 95°C; 60s, 60°C; 30s, 95°C; 15s. Apparent reaction efficiencies were determined by analyses of the standard curves and ranged between 92.4 - 96.8%. The R² for standard curves ranged between 0.984-1.000.

Primers were designed using the relevant white-throated sparrow coding sequence (cds) unless white-crowned sparrow cds were available (CBG and GnIH). CBG was recently identified in *Taeniopygia guttata* after nearly 40 years of research. Primers were designed against the published because few genome sequences have been updated (Vashchenko, Das et al. 2016). Multiple primer combinations were tested for each gene using the previously mentioned thermal conditions for SYBR reactions and with FastStart PCR (Roche) using a standard PCR thermal conditions [95°C; 240s, (40 cycles of 95°C; 30s, 58°C; 30s, 72°C; 30s), then 72°C; 420s].

Products from both PCR and qPCR reactions were visualized using 2% agarose gel electrophoresis. The primer combination for each gene was selected on the amplicons that yielded highest amplifying reaction that lacked visible primer dimers or infidelity signals (Table 1). Once primers were selected, FastStart amplicons were isolated from the agarose gel using a QIAquick Gel Extraction Kit (Qiagen 28704). Amplicons were sent for sequencing and GR, MR, 11 β HSD1, 11 β HSD2, YWHAZ were found to have sequence homologies with the published white-throated sparrow (*Zonotrichia albicollis*) genome of 98, 95, 100, 99.72, and 98.82% respectively. The CBG amplicon shared 90% identity with *Taeniopygia guttata* CBG, and the GnIH amplicon shared 98% identity with the published *Zonotrichia leucophrys gambelii* GnIH amplicon. Stock amplicons were serially diluted to produce standard curves for qPCR. For each gene assay per tissue, duplicate samples from each bird were all run on a single plate. The purified cDNA was used to generate standard curves by quantification on a Nanodrop and then a serial dilution was conducted from an aliquot of the stock amplicon solution. For each gene assay per tissue, duplicate samples from each bird were all run on a single plate. All genes were normalized to the reference gene YWHAZ due to its reliability in previous avian studies (Olias, Adam et al. 2014, Reid, Wilson et al. 2017).

MxPro software compares Ct values of unknown samples to those of standards with known target concentration to determine an absolute measure of the number of target cDNA copies in each reaction volume, which represented a constant proportion of total RNA. Absolute expression values for each gene of interest were then divided by the corresponding YWHAZ absolute expression value to give a normalized relative measure of expression, with arbitrary units, for each sample. This controlled for any inconsistency in total RNA representation (e.g. difference in total RNA reverse-transcribed per sample), as well as any disproportional

representation caused by large differences in unassessed transcripts (e.g. ribosomal RNAs or tissue-specific genes under certain seasonal conditions) compared to YWHAZ basal expression. Additional retrospective assessment of YWHAZ for reference suitability found expression to be consistent across all samples for each tissue type (see results).

Quantification of plasma corticosterone

Corticosterone concentrations were determined using a previously validated radioimmunoassay for corticosterone (Krause, Pérez et al. 2017). In brief, 15µL from baseline plasma samples and 10µL from the 30-minute samples were combined with 2000CPM of tritiated corticosterone (Perkin Elmer NET399250UC). 4mL of freshly distilled dichloromethane was used to extract corticosterone from the plasma samples. Aspirated dichloromethane was placed into a water bath at 35°C and was dried using nitrogen gas. Dried extracts were reconstituted in 550µL of phosphate buffer saline with gelatin (PBSG). 100µL of reconstituted extract was added to a scintillation vial and combined with 3mL of scintillation fluid (Perkin Elmer Ultima Gold: 6013329) to determine extraction recovery percentages. Next, 200µL of reconstituted extracts were added to duplicate RIA assay tubes with 100µL of tritiated corticosterone (Perkin Elmer NET399250UC) and 100µL of antiserum (MP Biomedical 07–120016, lot 3R3-PB-20E antibody). Unbound steroid was separated from bound steroid using 500µL of dextran coated charcoal solution. Samples were then placed in a centrifuge for 10 minutes at 4°C at 3000 g to separate bound from free steroids in solution. The supernatant containing bound steroids was decanted into scintillation vials and 3.5mL of scintillation fluid were added. Mean recovery was 85.25±4.2% (mean±SD) and intra- (calculated using C.V. between duplicates) and inter-assay variations were 5.05 and 6.14%, respectively.

237

238 **Statistical Analyses**

239 Statistical analyses of the data were performed in R Statistical Analysis software with R
240 studio version 3.6.3 (R Core Development Team 2018). Gene expression data were analyzed
241 using an analysis of variance (ANOVA) with the main effect of sampling time points. The
242 residuals were checked for normal distribution using Shapiro-Wilks and normality using
243 Levene's test. If the assumptions were not met, the data were log transformed. All post hoc tests
244 were performed using the emmeans package (Lenth, Singmann et al. 2018) to perform Tukey's
245 Honestly Significant Difference (HSD) tests and generate effects size estimates based on
246 Cohen's D. Pearson's correlations were used to investigate the relationship between either
247 baseline or stress-induced corticosterone and gene expression in each of the measured tissues.

248 **Results**

249 *Effects of restraint handling stress on gene expression*

250 MR, GR, 11 β HSD1 and 11 β HSD2 mRNA expressions were not affected by restraint handling in
251 any of the tissues measured (Figs. 1&2; Table 2). However, restraint handling resulted in
252 changes in mRNA expression of GnIH in the hypothalamus and CBG in the liver (Fig. 3; Table
253 2). Hypothalamic GnIH expression was higher at the 3-minute compared to 10-minute sampling
254 points ($t=4.26$, $P=0.002$, $df=21$, Cohen's $d=2.39$) and 30 ($t=2.82$, $P=0.04$, $df=21$, Cohen's
255 $d=1.38$; Fig. 2A). CBG in the liver was higher at the 60-minute sampling point compared to the
256 3-minute ($t=3.04$, $P=0.02$, $df=23$, Cohen's $d=1.57$) and 10 minute ($t=2.82$, $P=0.045$; Fig. 2B,
257 $df=23$, Cohen's $d=1.50$) sampling points.

YWHAZ gene expression was not affected by restraint handling in the hippocampus ($F_{3,22}=0.25$, $P=0.85$), hypothalamus ($F_{3,24}=0.007$, $P=0.99$), pituitary gland ($F_{3,19}=0.04$, $P=0.98$), or liver ($F_{3,23}=2.21$, $P=0.11$).

Relationships between gene expression and circulating corticosterone

No correlations were found between circulating baseline or stress-induced glucocorticoids and any of the genes measured (Table 2).

Discussion

We tested whether regulatory elements downstream of plasma levels of corticosterone respond rapidly to the stress of a standardized restraint handling protocol. There were no changes in MR, GR, 11 β HSD1, or 11 β HSD2 mRNA expression, suggesting that with respect to standardized restraint stress protocols lasting 1 hour or less, quantification of plasma corticosteroid levels may be sufficient to characterize HPA axis activity in white-crowned sparrows.. In rock doves, *Columba livia*, transcriptomic analysis showed no changes in MR, GR, 11 β HSD1, or 11 β HSD2 mRNA expression in the hypothalamus nor pituitary gland following 30 minutes of restraint stress (Calisi, Austin et al. 2017). Our findings along with Calisi et al. 2017, may be explained by the fact that changes in gene expression, particularly for genomic receptors, may occur on a longer time scale and were likely missed in the present study given the maximum observation window of one hour. For instance, in rats, 120 minutes, but not 15, 30, or 60 minutes, of acute stress decreased hippocampal GR expression but not MR (Paskitti, McCreary et al. 2000), upregulated hypothalamic GR expression but not MR, and decreased pituitary GR expression but not MR (Karandrea, Kittas et al. 2000). In birds, MR and GR mRNA expression

vary over longer time periods, for example in response to seasonal changes (Liebl, Shimizu et al. 2013, Krause, McGuigan et al. 2015, Li, Davis et al. 2020), social stress (Cornelius, Perreau et al. 2018), early life stress (Zimmer and Spencer 2014), chronic stress (Dickens, Romero et al. 2009), selection for high HPA axis activity (Hodgson, Meddle et al. 2007), and range expansion (Liebl and Martin 2013).

Another possible mechanism for rapidly modulating tissue level availability of corticosterone is through glucocorticoid regulatory enzymes. 11 β -hydroxysteroid dehydrogenase 1 (11 β HSD1) catalyzes the reactivation of corticosterone from inactive metabolites, and 11 β -hydroxysteroid dehydrogenase 2 (11 β HSD2) catalyzes the deactivation corticosterone to inactive metabolites. In the current study, we did not detect any rapid effects of stress on the expression of 11 β HSD1 or 11 β HSD2. A study in rodents found upregulation of 11 β HSD1 mRNA in the hippocampus at 1 and 2 hours post physiological stressor, but no other brain regions involved with HPA axis signaling were investigated (Spiers, Chen et al. 2016). 11 β HSD2 is not highly expressed in the brain of mammals (unlike in birds) thus no comparative studies could be found investigating the effects of stress (Chapman, Holmes et al. 2013). In white-crowned sparrows, pharmacological blockade of 11 β HSD1 and 11 β HSD2 enzyme function during both systemic administration and infusion into the third ventricle of the brain (11 β HSD2, only) altered levels of plasma corticosteroids over the course of a standardized restraint stressor (Pérez, Swanson et al. 2020). The central administration of pharmacological antagonists to the third ventricle suggested that 11 β HSD2 was involved in regulating negative feedback control at PVN of the hypothalamus. While, systemic injections targeted the liver, kidneys, and pituitary gland suggested that 11 β HSD2 and 11 β HSD1 are important for balancing net deactivation of corticosterone by the kidney and net reactivation by the liver and/or possible negative feedback

at the pituitary gland (Pérez, Swanson et al. 2020). However, regulation of gene expression is not the only way to modulate enzyme kinetics. Both enzymes are dependent upon cofactors, NAD⁺ and NADPH, for proper functioning and thus these could potentially regulate their activity. Studies on 3 β -hydroxysteroid dehydrogenase (3 β HSD) have been shown to undergo rapid modulation, by changes of NAD⁺ cofactors, in songbirds (Pradhan, Newman et al. 2010). Our data suggest that rapid regulation of expression is not occurring, but further studies will need to elucidate if either 11 β HSD1 or 2 undergoes a rapid response to cofactors.

Binding assays in birds have verified the presence of both MR and GR in the liver (Lattin, Waldron-Francis et al. 2012), but we found no effect on liver GR or MR mRNA expression in response to restraint handling stress. We have not found any published studies to date that have investigated the effects of short-term acute stress on corticosteroid receptor gene expression in the liver of any species. This represents a serious gap in our knowledge for how the liver responds to acute stress. In house sparrows, *Passer domesticus*, a chronic stress paradigm also had no effect on MR or GR in the liver when analyzed using receptor binding assays (Lattin and Romero 2014). However in fish and rodents, treatment of hepatocytes or whole animals with glucocorticoids increased GR mRNA expression after 24 h of exposure but not at 1, 4, or 8h post treatment (Sathiyaa and Vijayan 2003, Liu, Nakagawa et al. 2005). In addition, 6 h of chronic immobilization for 1 day in rodents resulted in upregulation of 11 β HSD1 and down regulation of 11 β HSD2 (Chen 2020). Thus, changes in gene expression associated with glucocorticoid signaling appear to be context and duration dependent.

Rapid changes in the expression of downstream targets of corticosteroid were observed following restraint stress. CBG expression in the liver increased in as little as 30 minutes in response to acute restraint stress. The current literature shows mixed results for the effects of

restraint handling on circulating plasma CBG in birds; some species show a decline while others are unaffected (Breuner, Lynn et al. 2006). Gambel's white-crowned sparrows do not show a decline in circulating CBG after 1 hour of restraint (Breuner, Lynn et al. 2006). The binding capacity of CBG in songbirds has been reported to be relatively high, allowing binding of additional steroids or potentially buffering the loss of cleaved CBG (Charlier, Underhill et al. 2009, Li, Sun et al. 2017) so that a substantial increase in circulating CBG may not be necessary to regulate HPA axis activity. Based on our data, a longer time course than that used previously by Breuner et al., (2006) may be required to detect a corresponding change in circulating CBG. Alternatively, increased CBG transcript production may be keeping pace with increased CBG cleavage and clearance during a stressor. Long-term stressors, such as food removal or restriction studies, have shown that plasma CBG levels decline in white-crowned sparrows within 22 hours (Lynn, Breuner et al. 2003).

GnIH is important for reducing the activity of the hypothalamic-pituitary-gonadal (HPG) axis, but contrary to our predictions, GnIH expression was markedly reduced after 10 and 30 minutes of restraint stress. Hypothalamic GnIH peptide and mRNA typically increase with stress, and glucocorticoid receptors colocalize with GnIH neurons (Calisi, Rizzo et al. 2008, Kirby, Geraghty et al. 2009, Gojska and Belsham 2014). Nonetheless studies in captive breeding zebra finches have shown GnIH immunoreactive cells to decrease in response to restraint stress and food removal (Ernst, Lynn et al. 2016, Wilsterman, Alonge et al. 2020). Studies in European starlings and rats have shown that GnIH levels in both sexes peak following clutch completion, which may serve to suppress HPG axis function in the event that the clutch is lost the reproductive system can reactivate (Calisi, Geraghty et al. 2016). Thus, it is difficult to determine the functional significance of the observed decline in GnIH, given its transient occurrence

(Kirby, Geraghty et al. 2009, Calisi, Díaz-Muñoz et al. 2011). Based on our existing knowledge of GnIH the decrease in expression may play a functional role in altering either HPG axis activity or sexual behavior in response to acute environmental stressors.

Conclusion

This is the first study in birds to test whether or not rapid changes in corticosteroid receptors or the 11 β Hydroxysteroid dehydrogenase system occur at multiple levels within the HPA axis and at target tissues during a standardized restraint handling sampling period. Our findings suggest that neither the expression of corticosteroid receptors nor the 11 β -hydroxysteroid dehydrogenase system change over the course of a standardized restraint handling protocol at any regulatory point within HPA axis. However, hepatic mRNA expression of CBG was rapidly upregulated which may serve to buffer circulating levels of glucocorticoids and may also be important for delivering corticosterone to target tissues. Thus, genes that are regulated by glucocorticoid signaling - GnIH and CBG, undergo rapid changes. Our data suggest that tissue level sensitivity for glucocorticoids as accessed by corticosteroid receptors and 11 β HSD enzymes suggests that these elements are stable on the acute time scale. Long-term stress studies are required to understand if these regulatory elements change over longer time periods.

Acknowledgements

We would like to thank Dr. Thomas P. Hahn, Dr. Marilyn Ramenofsky and our dedicated undergraduates Ryan Swanson and Hannah Lau for assistance with blood sampling, tissue collections, laboratory work, and animal husbandry.

Funding

This work was supported by the National Science Foundation Division of Integrative Organismal Systems IOS 1558049 to John C. Wingfield. Simone L. Meddle acknowledges Roslin Institute strategic grant funding from the Biotechnology and Biological Sciences Research Council (BB/P013759/1). J.C. Wingfield would like to acknowledge the University of California, Davis Endowed Chair in Physiology.

References

- Bouillé, C. and J. D. Baylé (1975). "Influence of Septal Nuclei on Basal Pituitary-Adrenocortical Function in Birds." Neuroendocrinology **18**(4): 281-289.
- Bouillé, C. and J. D. Baylé (1976). "Comparison between Hypothalamic, Hippocampal and Septal Multiple Unit Activity and Basal Corticotropic Function in Unrestrained, Unanesthetized Resting Pigeons." Neuroendocrinology **22**(2): 164-174.
- Breuner, C. and M. Orchinik (2002). "Plasma binding proteins as mediators of corticosteroid action in vertebrates." Journal of Endocrinology **175**(1): 99-112.
- Breuner, C. W., S. E. Lynn, G. E. Julian, J. M. Cornelius, B. J. Heidinger, O. P. Love, R. S. Sprague, H. Wada and B. A. Whitman (2006). "Plasma-binding Globulins and Acute Stress Response." Horm Metab Res **38**(04): 260,268.
- Calisi, R., S. Austin, A. S. Lang and M. D. MacManes (2017). "Sex-biased transcriptomic response of the reproductive axis to stress." bioRxiv.
- Calisi, R. M., S. L. Díaz-Muñoz, J. C. Wingfield and G. E. Bentley (2011). "Social and breeding status are associated with the expression of GnIH." Genes, Brain and Behavior **10**(5): 557-564.
- Calisi, R. M., A. C. Geraghty, A. Avila, D. Kaufer, G. E. Bentley and J. C. Wingfield (2016). "Patterns of hypothalamic GnIH change over the reproductive period in starlings and rats." General and Comparative Endocrinology **237**: 140-146.
- Calisi, R. M., N. O. Rizzo and G. E. Bentley (2008). "Seasonal differences in hypothalamic EGR-1 and GnIH expression following capture-handling stress in house sparrows (*Passer domesticus*)." General and Comparative Endocrinology **157**(3): 283-287.
- Chapman, K., M. Holmes and J. Seckl (2013). 11 β -Hydroxysteroid Dehydrogenases: Intracellular Gate-Keepers of Tissue Glucocorticoid Action.
- Charlier, T. D., C. Underhill, G. L. Hammond and K. K. Soma (2009). "Effects of aggressive encounters on plasma corticosteroid-binding globulin and its ligands in white-crowned sparrows." Hormones and Behavior **56**(3): 339-347.
- Chen, H.-J. C. Y., T.; Lee, J.K.; Juliani, J.; Sernia, C.; Hill, A.F.; Lavidis, N.A.; Spiers, J.G. (2020). "Restraint Stress Alters Expression of Glucocorticoid Bioavailability Mediators, Suppresses Nrf2, and Promotes Oxidative Stress in Liver Tissue. ." Antioxidants **9**(853).
- Cornelius, J. M., G. Perreau, V. R. Bishop, J. S. Krause, R. Smith, T. P. Hahn and S. L. Meddle (2018). "Social information changes stress hormone receptor expression in the songbird brain." Hormones and Behavior **97**: 31-38.

409 Dallman, M. F., S. F. Akana, K. A. Scribner, M. J. Bradbury, C. D. Walker, A. M. Strack and C. S. Cascio
 410 (1992). "Stress, Feedback and Facilitation in the Hypothalamo-Pituitary-Adrenal Axis." Journal of
 411 Neuroendocrinology **4**(5): 517-526.
 412 de Kloet, E. R. (2014). "From receptor balance to rational glucocorticoid therapy." Endocrinology **155**(8):
 413 2754-2769.
 414 Dickens, M., L. M. Romero, N. E. Cyr, I. C. Dunn and S. L. Meddle (2009). "Chronic stress alters
 415 glucocorticoid receptor and mineralocorticoid receptor mRNA expression in the European starling
 416 (*Sturnus vulgaris*) brain." Journal of Neuroendocrinology **21**(10): 832-840.
 417 Ernst, D. K., S. E. Lynn and G. E. Bentley (2016). "Differential response of GnIH in the brain and gonads
 418 following acute stress in a songbird." General and Comparative Endocrinology **227**: 51-57.
 419 Farner, D. S. and B. K. Follett (1966). "Light and other environmental factors affecting avian
 420 reproduction." Journal of Animal Science **25**(Supplement): 90-115.
 421 Gojska, N. M. and D. D. Belsham (2014). "Glucocorticoid receptor-mediated regulation of Rfrp (GnIH)
 422 and Gpr147 (GnIH-R) synthesis in immortalized hypothalamic neurons." Molecular and Cellular
 423 Endocrinology **384**(1): 23-31.
 424 Grasa, M. M., C. Cabot, J. A. Fernández-López, X. Remesar and M. Alemany (2001). "Modulation of
 425 corticosterone availability to white adipose tissue of lean and obese Zucker rats by corticosteroid-
 426 binding globulin." Hormone and metabolic research **33**(7): 407.
 427 Gulfo, J., R. Castel, A. Ledda, M. d. M. Romero, M. Esteve and M. Grasa (2019). "Corticosteroid-Binding
 428 Globulin is expressed in the adrenal gland and its absence impairs corticosterone synthesis and
 429 secretion in a sex-dependent manner." Scientific Reports **9**(1): 14018.
 430 Gulfo, J., A. Ledda, E. Serra, C. Cabot, M. Esteve and M. Grasa (2016). "Altered lipid partitioning and
 431 glucocorticoid availability in CBG-deficient male mice with diet-induced obesity." Obesity **24**(8): 1677-
 432 1686.
 433 Hammond, G. L., C. L. Smith and D. A. Underhill (1991). "Molecular studies of corticosteroid binding
 434 globulin structure, biosynthesis and function." The Journal of Steroid Biochemistry and Molecular
 435 Biology **40**(4-6): 755-762.
 436 Harris, A. P., M. C. Holmes, E. R. de Kloet, K. E. Chapman and J. R. Seckl (2013). "Mineralocorticoid and
 437 glucocorticoid receptor balance in control of HPA axis and behaviour." Psychoneuroendocrinology **38**(5):
 438 648-658.
 439 Harris, H. J., Y. Kotelevtsev, J. J. Mullins, J. R. Seckl and M. C. Holmes (2001). "Intracellular Regeneration
 440 of Glucocorticoids by 11 β -Hydroxysteroid Dehydrogenase (11 β -HSD)-1 Plays a Key Role in Regulation of
 441 the Hypothalamic-Pituitary-Adrenal Axis: Analysis of 11 β -HSD-1-Deficient Mice." Endocrinology **142**(1):
 442 114-120.
 443 Hodgson, Z. G., S. L. Meddle, M. L. Roberts, K. L. Buchanan, M. R. Evans, R. Metzdorf, M. Gahr and S. D.
 444 Healy (2007). "Spatial ability is impaired and hippocampal mineralocorticoid receptor mRNA expression
 445 reduced in zebra finches (*Taeniopygia guttata*) selected for acute high corticosterone response to
 446 stress." Proceedings of the Royal Society B: Biological Sciences **274**(1607): 239-245.
 447 Joëls, M., H. Karst, R. DeRijk and E. R. de Kloet (2008). "The coming out of the brain mineralocorticoid
 448 receptor." Trends in Neurosciences **31**(1): 1-7.
 449 Joëls, M. and E. R. d. Kloet (2017). "30 YEARS OF THE MINERALOCORTICOID RECEPTOR: The brain
 450 mineralocorticoid receptor: a saga in three episodes." **234**(1): T49.
 451 Karandrea, D., C. Kittas and E. Kitraki (2000). "Contribution of sex and cellular context in the regulation
 452 of brain corticosteroid receptors following restraint stress." Neuroendocrinology **71**(6): 343-353.
 453 Kirby, E. D., A. C. Geraghty, T. Ubuka, G. E. Bentley and D. Kaufer (2009). "Stress increases putative
 454 gonadotropin inhibitory hormone and decreases luteinizing hormone in male rats." Proceedings of the
 455 National Academy of Sciences **106**(27): 11324-11329.

Kolber, B. J., L. Wiczorek and L. J. Muglia (2008). "Hypothalamic–pituitary–adrenal axis dysregulation and behavioral analysis of mouse mutants with altered glucocorticoid or mineralocorticoid receptor function." Stress **11**(5): 321-338.

Krause, J. S., M. A. McGuigan, V. R. Bishop, J. C. Wingfield and S. L. Meddle (2015). "Decreases in mineralocorticoid but not glucocorticoid receptor mRNA expression during the short Arctic breeding season in free-living Gambel's white-crowned sparrow (*Zonotrichia leucophrys gambelii*)."
Journal of Neuroendocrinology **27**(1): 66-75.

Krause, J. S., J. H. Pérez, S. L. Meddle and J. C. Wingfield (2017). "Effects of short-term fasting on stress physiology, body condition, and locomotor activity in wintering male white-crowned sparrows." Physiology & Behavior **177**: 282-290.

Landys, M. M., M. Ramenofsky and J. C. Wingfield (2006). "Actions of glucocorticoids at a seasonal baseline as compared to stress-related levels in the regulation of periodic life processes." General and Comparative Endocrinology **148**(2): 132-149.

Lattin, C. R. and L. M. Romero (2014). "Chronic stress alters concentrations of corticosterone receptors in a tissue-specific manner in wild house sparrows (*Passer domesticus*)."
The Journal of Experimental Biology **217**(14): 2601-2608.

Lattin, C. R., K. Waldron-Francis, J. W. Richardson, R. de Bruijn, C. M. Bauer, C. W. Breuner and L. Michael Romero (2012). "Pharmacological characterization of intracellular glucocorticoid receptors in nine tissues from house sparrow (*Passer domesticus*)."
General and Comparative Endocrinology **179**(2): 214-220.

Lenth, R., H. Singmann, J. Love, P. Buerkner and M. Herve (2018). "Emmeans: Estimated marginal means, aka least-squares means." R package version **1**(1): 3.

Li, D., J. E. Davis, G. Wang, G. Nabi, V. R. Bishop, Y. Sun, S. L. Meddle, J. C. Wingfield and F. Lei (2020). "Coping with extremes: Remarkably blunt adrenocortical responses to acute stress in two sympatric snow finches on the Qinghai-Tibet Plateau during winter relative to other seasons." General and Comparative Endocrinology: 113434.

Li, Y., Y. Sun, J. S. Krause, M. Li, X. Liu, W. Zhu, Y. Yao, Y. Wu and D. Li (2017). "Dynamic interactions between corticosterone, corticosteroid binding globulin and testosterone in response to capture stress in male breeding Eurasian tree sparrows." Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology **205**: 41-47.

Liebl, A. L. and L. B. Martin (2013). "Stress hormone receptors change as range expansion progresses in house sparrows." Biology Letters **9**(3).

Liebl, A. L., T. Shimizu and L. B. Martin (2013). "Covariation among glucocorticoid regulatory elements varies seasonally in house sparrows." General and Comparative Endocrinology **183**(0): 32-37.

Liu, Y., Y. Nakagawa, Y. Wang, R. Sakurai, P. V. Tripathi, K. Lutfy and T. C. Friedman (2005). "Increased Glucocorticoid Receptor and 11 β -Hydroxysteroid Dehydrogenase Type 1 Expression in Hepatocytes May Contribute to the Phenotype of Type 2 Diabetes in *db/db* Mice." Diabetes **54**(1): 32-40.

Lynn, S. E., C. W. Breuner and J. C. Wingfield (2003). "Short-term fasting affects locomotor activity, corticosterone, and corticosterone binding globulin in a migratory songbird." Hormones and Behavior **43**(1): 150-157.

Olias, P., I. Adam, A. Meyer, C. Scharff and A. D. Gruber (2014). "Reference genes for quantitative gene expression studies in multiple avian species." PLOS ONE **9**(6): e99678.

Paskitti, M. E., B. J. McCreary and J. P. Herman (2000). "Stress regulation of adrenocorticosteroid receptor gene transcription and mRNA expression in rat hippocampus: time-course analysis." Molecular Brain Research **80**(2): 142-152.

Pérez, J. H., R. E. Swanson, H. J. Lau, J. Cheah, V. R. Bishop, K. R. S. Snell, A. M. A. Reid, S. L. Meddle, J. C. Wingfield and J. S. Krause (2020). "Tissue-specific expression of 11 β -HSD and its effects on plasma corticosterone during the stress response." The Journal of Experimental Biology **223**(1): jeb209346.

Pradhan, D. S., A. E. M. Newman, D. W. Wacker, J. C. Wingfield, B. A. Schlinger and K. K. Soma (2010). "Aggressive interactions rapidly increase androgen synthesis in the brain during the non-breeding season." Hormones and Behavior **57**(4-5): 381-389.

R Core Development Team (2018). R: a language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria.

Ramenofsky, M., A. W. Campion, J. H. Pérez, J. S. Krause and Z. Németh (2017). "Behavioral and physiological traits of migrant and resident white-crowned sparrows: a common garden approach." The Journal of Experimental Biology **220**(7): 1330-1340.

Reid, A. M. A., P. W. Wilson, S. D. Caughey, L. M. Dixon, R. B. D'Eath, V. Sandilands, T. Boswell and I. C. Dunn (2017). "Pancreatic PYY but not PPY expression is responsive to short-term nutritional state and the pancreas constitutes the major site of PYY mRNA expression in chickens." General and Comparative Endocrinology **252**: 226-235.

Romero, L. M. and J. C. Wingfield (1999). "Alterations in hypothalamic–pituitary–adrenal function associated with captivity in Gambel's white-crowned sparrows (*Zonotrichia leucophrys gambelii*)."
Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology **122**(1): 13-20.

Sapolsky, R. M., L. M. Romero and A. U. Munck (2000). "How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions." Endocrine Reviews **21**(1): 55-89.

Sathiyaa, R. and M. M. Vijayan (2003). "Autoregulation of glucocorticoid receptor by cortisol in rainbow trout hepatocytes." American Journal of Physiology-Cell Physiology **284**(6): C1508-C1515.

Schmidt, M. V., V. Sterlemann, K. Wagner, B. Niederleitner, K. Ganea, C. Liebl, J. M. Deussing, S. Berger, G. n. Schütz, F. Holsboer and M. B. Müller (2009). "Postnatal Glucocorticoid Excess Due to Pituitary Glucocorticoid Receptor Deficiency: Differential Short- and Long-Term Consequences." Endocrinology **150**(6): 2709-2716.

Schoech, S. J., L. M. Romero, I. T. Moore and F. Bonier (2013). "Constraints, concerns and considerations about the necessity of estimating free glucocorticoid concentrations for field endocrine studies." Functional Ecology **27**(5): 1100-1106.

Seckl, J. R. and K. E. Chapman (1997). "Medical and physiological aspects of the 11beta-hydroxysteroid dehydrogenase system." European journal of biochemistry / FEBS **249**(2): 361-364.

Sivukhina, E. V. and G. F. Jirikowski (2014). "Adrenal steroids in the brain: Role of the intrinsic expression of corticosteroid-binding globulin (CBG) in the stress response." Steroids **81**: 70-73.

Smulders, T. V. (2017). "The avian hippocampal formation and the stress response." Brain, Behavior and Evolution **90**(1): 81-91.

Spiers, J. G., H.-J. C. Chen, J. S. M. Cuffe, C. Sernia and N. A. Lavidis (2016). "Acute restraint stress induces rapid changes in central redox status and protective antioxidant genes in rats." Psychoneuroendocrinology **67**: 104-112.

Stokes, T. M., C. M. Leonard and F. Nottebohm (1974). "The telencephalon, diencephalon, and mesencephalon of the canary, *Serinus canaria*, in stereotaxic coordinates." The Journal of Comparative Neurology **156**(3): 337-374.

Vashchenko, G., S. Das, K.-M. Moon, J. C. Rogalski, M. D. Taves, K. K. Soma, F. Van Petegem, L. J. Foster and G. L. Hammond (2016). "Identification of Avian Corticosteroid-binding Globulin (Serpina6) Reveals the Molecular Basis of Evolutionary Adaptations in SerpinA6 Structure and Function as a Steroid-binding Protein." Journal of Biological Chemistry **291**(21): 11300-11312.

Wilsterman, K., M. M. Alonge, X. Bao, K. A. Conner and G. E. Bentley (2020). "Food access modifies GnIH, but not CRH, cell number in the hypothalamus in a female songbird." General and Comparative Endocrinology: 113438.

Zimmer, C. and K. A. Spencer (2014). "Modifications of Glucocorticoid Receptors mRNA Expression in the Hypothalamic-Pituitary-Adrenal Axis in Response to Early-life Stress in Female Japanese Quail." Journal of Neuroendocrinology **26**(12): 853-860.

Table and Figure Legends

Table 1.

Primers that were utilized in this study.

Gene	Accession Number	Primer	Primer Seq
Corticosteron Binding Globulin (CBG)	KU180443.1	F	GCAGATCTCTCTGGCATCAC
		R	GAGGAAGGGATGGCTGAATTC
Glucocorticoid Receptor (GR)	XM_002192952.3	F	TGAAGAGCCAGTCCCTGTTTCGAG
		R	CAACCACATCATGCATAGAGTCCAGCA
Mineralocorticoid Receptor (MR)	NM_001076690.1	F	AAGAGTCGGCCAAACATCCTTGTCT
		R	AAGAAACGGGTGGTCTAAATCCCAG
Gonadotropin Inhibitory Hormone (GnIH)	AB128164.1	F	CTGCCTCGAGACATTCAGGA
		R	GACCACCACTGAAGCAAAC
11 β -Hydroxysteroid dehydrogenase 1 (11 β HSD1)	XM_005492865.1	F	GCTCATCCTCAACCACATCG
		R	CCATCTAGGGCGAACTTGGT
11 β -Hydroxysteroid dehydrogenase 2 (11 β HSD2)	XM_005490743.3	F	ATATCCAGGCCACACCAAC
		R	CACGTTGTCCCTGTTTGTAGT
Tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta (YWHAZ)	NM_001031343.1	F	GTGGAGCAATCACAACAGGC
		R	GCGTGCGTCTTTGTATGACTC

Table 2.

Analyses of variance results investigating the effects of acute capture restraint stress on gene expression and Pearson's correlations between baseline and stress-induced corticosterone and gene expression of glucocorticoid receptor (GR), mineralocorticoid receptor (MR), 11 β -hydroxysteroid dehydrogenase 1 (11 β HSD1), 11 β -hydroxysteroid dehydrogenase 2 (11 β HSD2), corticosterone binding globulin

(CBG) and gonadotropin inhibitory hormone (GnIH) mRNA in tissues including the hippocampus, hypothalamus, pituitary gland, and liver of male white crowned sparrows.

	Gene expression			Baseline CORT Correlation			Stress-induced CORT Correlation		
Mineralocorticoid receptor (MR)	DF	F	P	DF	r	P	DF	r	P
Hippocampus	3,22	0.01	0.98	24	0.28	0.10	24	0.22	0.26
Hypothalamus	3,24	0.73	0.54	2	0.24	0.21	26	0.13	0.48
Pituitary gland	3,14	3.13	0.06	16	0.05	0.81	16	0.02	0.92
Liver	3,22	0.86	0.48	24	0.27	0.17	24	0.14	0.46
Glucocorticoid receptor (GR)									
Hippocampus	3,22	0.13	0.94	24	0.22	0.27	24	0.13	0.51
Hypothalamus	3,24	0.12	0.94	26	0.06	0.73	26	0.28	0.15
Pituitary gland	3,14	1.20	0.34	15	0.11	0.66	15	0.17	0.49
Liver	3,22	0.45	0.71	24	0.02	0.88	24	0.28	0.16
11 β -Hydroxysteroid dehydrogenase 1 (11 β HSD1)									
Hippocampus	3,21	1.24	0.31	23	-0.05	0.8	23	0.08	0.68
Hypothalamus	3,23	0.23	0.87	25	-0.01	0.92	25	0.03	0.87
Pituitary gland	3,16	0.39	0.76	18	0.43	0.06	18	0.26	0.25
Liver	3,22	0.77	0.51	25	0.33	0.63	25	0.22	0.26
11 β -Hydroxysteroid dehydrogenase 2 (11 β HSD2)									
Hippocampus	3,21	1.95	0.15	23	0.01	0.93	23	-0.07	0.71
Hypothalamus	3,22	1.84	0.16	24	-0.27	0.16	24	-0.04	0.81
Pituitary gland	3,16	0.13	0.93	18	0.52	0.01	18	0.16	0.5
Liver	3,22	0.77	0.52	24	0.21	0.29	24	0.35	0.07
Corticosterone binding globulin (CBG)									
Liver	3,23	22.85	0.02	25	0.06	0.75	25	0.16	0.41
Gonadotropin-inhibitory hormone									
Hypothalamus	3,21	8.76	<0.0001	23	0.03	0.88	23	0.16	0.56

Figure 1.

Effects of capture restraint stress on the expression of glucocorticoid (GR; A, B, C, D) and mineralocorticoid (MR; E, F, G, H) receptor mRNA in the hippocampus, hypothalamus, pituitary gland, and liver from time of capture in male white-crowned sparrows. Data were normalized relative to the house keeping gene YWHAZ. Letters that are different from one another indicate a significant difference using contrasts with a Tukey's correction. Values represent means $\pm$ SEM.

Figure 2.

Effects of capture restraint stress on the expression of 11 β -hydroxysteroid dehydrogenase 1 (11 β HSD1; A, B, C, D) and 11 β -hydroxysteroid dehydrogenase 2 (11 β HSD2; E, F, G, H) mRNA in the hippocampus, hypothalamus, pituitary gland, and liver from time of capture in male white-crowned sparrows. Data were normalized relative to the house keeping gene YWHAZ. Letters that are different from one another indicate a significant difference using contrasts with a Tukey's correction. Values represent means $\pm$ SEM.

Figure 3.

585 Effects of restraint stress on the expression of (A) hypothalamic gonadotropin inhibitory hormone (GnIH)
586 and (B) liver corticosteroid binding globulin (CBG) mRNA. Data were normalized relative to the house
587 keeping gene YWHAZ. Letters that are different from one another indicate a significant difference using
588 contrasts with a Tukey's correction. Values represent means $\pm$ SEM.

589

590 Supplemental Figure 1. YWHAZ reference gene expression in hippocampus, hypothalamus, pituitary
591 gland and liver of white crowned sparrows exposed to restraint handling.