THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Tropical peatlands and their contribution to the global carbon cycle and climate change

Citation for published version:

Ribeiro, K, Pacheco, FS, Ferreira, JW, Sousaneto, ER, Hastie, A, Krieger Filho, GC, Alvalá, PC, Forti, MC & Ometto, JP 2021, 'Tropical peatlands and their contribution to the global carbon cycle and climate change', *Global Change Biology*, vol. 27, no. 3, pp. 489-505. <https://doi.org/10.1111/gcb.15408>, <https://doi.org/10.1111/gcb.v27.3>

Digital Object Identifier (DOI):

[10.1111/gcb.15408](https://doi.org/10.1111/gcb.15408)

[10.1111/gcb.v27.3](https://doi.org/10.1111/gcb.v27.3)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Global Change Biology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Title: Tropical peatlands and their contribution to the global carbon cycle and climate change

Running Title: Tropical peatlands, carbon cycle and climate

Kelly Ribeiro¹, Felipe Siqueira Pacheco¹, José Willian Ferreira¹, Eráclito Rodrigues de Sousa-Neto¹, Adam Hastie², Guenther Carlos Krieger Filho³, Plínio Carlos Alvalá¹, Maria Cristina Forti¹, Jean Pierre Ometto¹

1 – Earth System Science Center (CCST), National Institute for Space Research (INPE), Av. Astronautas, 1758, São José dos Campos, São Paulo, 12227-000, Brazil.

2 - School of GeoSciences, University of Edinburgh, EH9 3FF, Edinburgh, Scotland, UK

3- Laboratory of Thermal and Environmental Engineering, Polytechnic School of the University of São Paulo, Av. Prof. Mello Moraes, 2231, São Paulo 05508-030, Brazil

Contact Information:

Kelly Ribeiro, Earth System Science Center (CCST),
São José dos Campos, Brazil

Email: kelly.ribeiro@inpe.br

Tel: +55 12 32087926

Abstract

Peatlands are carbon-rich ecosystems that cover 185-423 million hectares of the earth's surface. The majority of the world's peatlands are in temperate and boreal zones, whereas tropical ones cover only a total area of 90-170 million hectares. However, there are still considerable uncertainties in C stock estimates as well as a lack of information about depth, bulk density and carbon accumulation rates. The incomplete data is notable especially in tropical peatlands located in South America, which are estimated to have the largest area of peatlands in the tropical zone. This paper displays the current state of knowledge surrounding tropical peatlands and their biophysical characteristics, distribution and carbon stock, role in the global climate, the impacts of direct human disturbances on carbon accumulation rates and greenhouse gas emissions. Based on the new peat extension and depth data, we estimate that tropical peatlands store 152-288 GtC, or about half of the global peatland emitted carbon. We discuss the knowledge gaps in research on distribution, depth, C stock and fluxes in these ecosystems which play an important role in the global carbon cycle and risk releasing large quantities of greenhouse gases into the atmosphere (CO_2 and CH_4) when subjected to anthropogenic interferences (e.g. drainage and deforestation). Recent studies show that although climate change has an impact on the carbon fluxes of these ecosystems, the direct anthropogenic disturbance may play a greater role. The future of these systems as carbon sinks will depend on advancing current scientific knowledge and incorporating local understanding to support policies geared toward managing and conserving peatlands in vulnerable regions, such as the Amazon where recent records show increased forest fires and deforestation.

Keywords: tropical peatlands, carbon cycling, greenhouse gas emissions, climate change, land use change

Introduction

Peatlands are a type of wetland that form when waterlogged anoxic conditions limit the decomposition and respiration of organic matter (Vitt, 2013) and creates an accumulation of peat. Like most wetlands, peatlands can have dense vegetation cover with lacustrine characteristics (presence of the water plants), sometimes influenced by river seasonality and dynamics (Lähteenoja, Flores, & Nelson, 2013), as well as seasonal or annual floods with geomorphological features where water is retained (Finlayson & Milton, 2018; Kelly et al., 2013; Warner & Rubec, 1997). Whilst there is no absolute consensus on what defines peat, most studies have settled on two criteria: soils that have both an organic matter content of at least 30% (Reiche, Gleixner, & Küsel, 2010; Sorensen, 1993), though typically with a higher threshold of 50% (Gumbricht et al., 2017) or 65% (e.g. (Dargie et al., 2017), and a minimum depth of 30cm to 40cm (Dargie et al., 2017; Page et al., 2011; Page & Baird, 2016; Dargie et al., 2017; Page et al., 2011; Page & Baird, 2016) .

These ecosystems provide unique ecosystem services, such as water storage by regulating the river's discharge, thereby benefiting ecosystems and human communities (Harenda, Lamentowicz, Samson, & Chojnicki, 2018), along with regulating water flow in hydrographic basins, including buffering floods (Joseph, 2005). Moreover, they are fertile fields for agricultural and horticultural production (Rieley et al., 2008), play an important role in sediment, nutrient and carbon (C) retention (Rieley et al., 2008), and are home to a unique biodiversity that includes a variety of endemic species (Wilson, Griffiths, & Anielski, 2001).

Peatlands cover a total area of about 185-423 million hectares throughout the world (1.2-2.8% of the earth's total land area) (Xu, Morris, Liu, & Holden, 2018). In the tropical area, zones covered by peat range from 90-170 million hectares and are located mainly in South America, Southeast Asia and Central Africa (Gumbricht et al., 2017). These system store large amounts of C (469-694 Giga tonnes of C) (Lähteenoja et al., 2012; Leifeld & Menichetti, 2018;

Page et al., 2011; Yu et al., 2010) and act as net sinks of atmospheric carbon dioxide (CO₂). However, they can also act as major sources of greenhouse gas (GHG) emissions, such as CO₂ and methane (CH₄), into the atmosphere (Leifeld & Menichetti, 2018; Roulet, 2012), due to either natural processes such as changes in autotrophic and heterotrophic respiration rates, changes in river paths, droughts and natural fires, or anthropogenic interferences including logging, drainage, deforestation, fires, and land use and land cover (LULC) changes (Hooijer et al., 2010; Leng, Ahmed, & Jalloh, 2019; Yule, Lim, & Lim, 2016).

Changes to the gross C uptake and/or release of these ecosystems can reverse whole-peatlands carbon budget and significantly alter the current and future global climate (Worrall et al., 2011; Wu & Roulet, 2014). In recent years, tropical peatlands have been receiving more attention not only because of their contribution to the global carbon budget and climate change, but also because of new estimates of larger peatland areas in the tropics (Dargie et al., 2017; Draper et al., 2014; Gumbrecht et al., 2017; Page et al., 2011; Xu et al., 2018). To understand how tropical peatlands contribute to global climate change, it is important to understand their geographical coverage, capacity to store and sequester carbon, and the main factors that drive their degradation (Yu, 2011).

In contrast to temperate peatlands, in which the relationship between climate, ecosystem dynamics and carbon (C) accumulation is well studied, the body of literature on tropical peatlands is mainly concentrated on Southeast Asia (Page, Rieley, Shotyk, & Weiss, 1999; San José et al., 2013), Peru in South America (Kelly et al., 2017; Läähteenoja et al., 2012; Roucoux et al., 2017; Sorribas et al., 2016) and, to a lesser degree, the Cuvette Centrale basin in Africa (Dargie et al., 2019, 2017). This paper presents an extensive review about tropical peatlands in terms of their biophysical conditions that promotes peat formation (e.g. temperature, rainfall, ground water, nutrient pool and substrate quality), spatial distribution and carbon stock, as well as how these ecosystems are affected under different disturbance regimes.

Moreover, the paper identifies and discusses knowledge gaps surrounding this highly threatened, yet poorly understood ecosystem in several regions of the tropical area.

Peat formation process and biophysical characteristics of tropical peatlands

In general, peat is formed when the amount of photosynthetically produced organic matter exceeds the loss of organic matter through fire, decomposition and lateral loss (Hodgkins et al., 2018). Peat formation is led by several factors s, such as hydrological dynamics (groundwater, seasonality and river dynamics), climatic characteristics (temperature and precipitation), underlying topography and geology of the area, nutrient pool, chemistry, and vegetation dynamics (Biancalani & Avagyan, 2014; Hapsari et al., 2017; Yu, 2012). Hydrological dynamics are among the main factors that regulate peatlands and control peat formation processes, predominant vegetation, nutrient content, carbon sequestration capacity, and decomposition processes (Blodau, 2002; Limpens et al., 2008). In certain peatlands, where water-saturated condition occurs all year around, peat soil profiles identified peat domes that reach depths up to 15 m (Gumbricht et al., 2017).

In many tropical peatlands the soil is seasonally flooded mostly by large rivers with high nutrient content and intense sediment deposition. These factors associated to high precipitation patterns and temperatures favor the development of flooded peatlands concomitant of dense tree coverage, with high floristic diversity and high net primary productivity (NPP) (Gillman et al., 2015) and absence of mosses (Page et al., 1999). On the contrary, in many northern peatlands, the low temperature and low nutrient inputs favors the dominance of the bryophyte genus *Sphagnum* (Clymo, 1987) and a shrub layer is usually well developed with sparse occurrence of large trees (Vitt, 2013; Ingram, 1987).

The peatland vegetation cover described above, is an important characteristic that influences the composition and the process of peat formation. Peat in tropical peatlands is mainly formed by woody material and dead branches and roots (Dommain et al., 2015; Gallego-

Sala et al., 2018) whereas most of the peat in northern peatlands is formed of Sphagnum mosses and sedges. The woody material contains high C:N and lignin:N ratios that degrade slowly (Gandois et al., 2012; Gandois et al., 2014). This promotes the release of phenolic components that inhibit decomposition (H. Wang, Richardson, & Ho, 2015). These conditions increase the aromatic content in the soil and create a reduced oxidation state in which C remains and recalcitrance is high, despite high temperatures (Hodgkins et al., 2018). Divergent from the peat formation process in northern peatlands, the low soil temperatures, freezing and the acid characteristics of the cell wall of Sphagnum species favor the reduction of C oxidative processes even with abundant labile carbohydrates (Sphagnum) (Vitt, 2013; Clymo, Kramer, & Hammerton, 1984).

“In general, there are two types of peatlands: ombrotrophic and minerotrophic (Clymo, 1987), being divided according to the origin of nutrient input in the system . Ombrotrophic peatlands are influenced exclusively by water from precipitation (no other sources) (Bourbonniere, 2009; Takada, Shimada, & Takahashi, 2016; Vitt, 2013) while minerotrophic peatlands are typically formed in depressions and floodplains and receive mineral nutrients with incoming surface or ground water (Bourbonniere, 2009; International Peatland Society (IPS), n.d.; Lahteenoja et al., 2009; Takada et al., 2016; Vitt, 2013).

At the start of the peat formation, the peatland is initially minerotrophic (Clymo, 1987). As the peat layer grows in height, the dome becomes elevated and the peatland may no longer be affected by the river that feeds into it or by the entry of groundwater, thereby obtaining water exclusively from precipitation and becoming ombrotrophic. At this stage, nutrient and mineral deposits are mainly from atmospheric dry deposition or precipitation, but large amounts of nutrients can also come from dust and air pollution (Ponette-Gonzalez et al., 2016). For instance, according to Swindles et al. (2018), the oldest Peruvian tropical peatlands discovered to date were formed in three stages: first, peat was formed in an abandoned river channel with

open water and aquatic plants; then inundated forest swamp was formed; and finally the peat dome raised as the peat accumulated.

Many peatlands in tropical region are minerotrophic having been formed from the lateral migration of rivers (Lähteenoja et al., 2013, 2012; Lähteenoja et al., 2009; Schumann & Joosten, 2008). Most of them are located in river deltas, floodplain areas, abandoned river channels and shallow oxbow lakes (dead arms) (Baker, 2014; C. B. T.-C. and R. W. Craft, 2016; Rieley et al., 2008; Rebelo, Finlayson, & Nagabhatla, 2009). However, there are examples of ombrotrophic peat bogs in the tropics in South America (S E Page et al., n.d.; Swindles et al., 2018), Southeast Asia (S E Page, Rieley, & Wüst, 2006; Wösten, Clymans, Page, Rieley, & Limin, 2008) and Africa (Dargie et al., 2017) reported in the literature. In Southeast Asia, different formation processes have been observed and most of the peat is currently ombrotrophic, with some related to ancient sea-level rise and an increase of Holocene precipitation (Dommain, Couwenberg, & Joosten, 2011). Thus, even at similar latitudes, the mechanisms of peat formation, regulation and carbon accumulation can differ between regions.”

Distribution and carbon stock of tropical peatlands

There is a lot of variation in the published data about the occurrence and distribution of tropical peatlands. Up to a few years ago, Southeast Asia (Indonesia, East Sumatra, Kalimantan, Papua New Guinea, Papua New Guinea, and Malaysia) was considered to have the largest peatland C reservoirs in the tropical area (Dargie et al., 2017; Joosten, 2009; Lähteenoja et al., 2009; Miettinen & Liew, 2010; Miettinen, Shi, & Liew, 2016; Page, Rieley, & Banks, 2011; Page et al., 2002), however large intact peatlands have recently been described in South America (Draper et al., 2014) and Africa (Dargie et al., 2017). For example, Dargie et al. (2017) used field measurements combined with remote sensing data to estimate the extent of a peat complex in the Cuvette Centrale region of the Congo Basin, the largest intact tropical peatland

to date at 14.6 (13.2-15.6) million hectares (Mha). As a result of these recent studies, estimates of tropical peatlands have been revised (see Gumbrecht et al., 2017; Xu et al., 2018) and the total area of tropical peatlands is now considered to cover 90-170 Mha, (23% to 30% of the total area covered by peatlands throughout the world). This new estimate is two to three times larger than the 56 Mha that Page et al. (2011) reported and which led to new discussions on the physical and chemical factors that define wetlands and peatlands (Figure 1).

The new estimates of total peat cover in the tropics represent a volume of about 3,850-7,268 km³ (estimated using area from Xu et al., 2018 and Gumbrecht et al., 2017, and mean depth from Gumbrecht et al., 2017), which is much higher than the previous estimate of 1,758 km³ (Page et al., 2011) (Figure 1). Considering these estimates, the largest reserves of peat are located in Brazil (area and volume of 23 Mha and 900 km³, respectively), Indonesia (14 Mha and 578 km³) and the Democratic Republic of the Congo (9 Mha and 445 km³, Figure 2). It is important to note that in Indonesia there is a longer history of fieldwork and, therefore, a relatively large database of ground-truthing points (Jaenicke et al., 2008), whereas to date there are relatively few published field data from the Congo Basin (Dargie et al., 2017), and even fewer from Brazil (Lähteenoja. 2013).

Based on the estimated volume of peat in the tropics and the average carbon content per km³ of peat (Lähteenoja et al., 2009), we estimated that peat in the tropics stores an equivalent of 152-288 GtC (Table 1), which is significantly higher than previously reported estimates of 119.2 (Leiffield & Menichetti, 2018), 104.7 (Dargie et al., 2017), 90 (Moore et al., 2013), 88.6 (Page et al., 2011) and 52 GtC (Zoltai & Martikainen, 1996). The stock of 152-288 GtC is equivalent to the amount of C emitted by burning fossil fuels at a rate of 10 GtC year⁻¹ for the next 15-30 years (Murdiyarso, Hergoualch, & Verchot, 2010; Raupach et al., 2013). C emission from fossil fuel in 2014 were 9.8 GtC, (<https://www.globalcarbonproject.org/>). In addition, the mid-range value of our estimated C stock (215 Gt) represents about 25% of the terrestrial carbon

pool in the tropics (846.3 GtC), considering both carbon above ground (374.9 GtC, phytomass) and stored in the soil (571.3 GtC, Scharlemann et al., 2014).

The main explanation for the large range in our new estimate of tropical peatland C stock (152–288 GtC) is the different methodological approaches adopted for the estimation of the area. For instance, the numerical model that Gumbrecht et al. (2017) adopted to estimate total area uses a set of factors associated with hydrological modeling, time series of vegetation, soil moisture and hydro-geomorphological data. Xu et al. (2018) considered a wide variety of sources from different authors and regions and applied criteria of relevance, spatial resolution as well as age, and combined these data sources to produce a new amalgamated global map of peatland distribution. For areas where peatland-specific datasets were not available, they estimated peatland extent based on the distribution of histosols derived from the Harmonized World Soil Database v1.2 (HWSD). Page et al., (2011) considered data from national soil inventories from different countries. Data from the latter may not be comparable given the different definitions of peat and inclusion of non-peat organic soils. The new estimates of Gumbrecht et al. (2017) and Xu et al. (2018) suggest that the extent of differing with what was previously reported of what was previously reported (Page et al., 2011).

Peatlands in the tropical zone are found in many countries, however some regions have large peatland areas and carbon stock. The South American peatlands are estimated to be located mainly in the Rio Negro Basin (Brazil) and Pastaza-Marañón Foreland Basin (PMFB, Peru) (Draper et al., 2014, Lähteenoja et al., 2013; Lähteenoja et al., 2009), however, to date there has been limited ground-truthing of the former (Lähteenoja et al., 2013) and therefore larger uncertainty associated with the extent and volume of Brazilian peatlands. The PMFB alone is estimated to represent a C stock of 3.14 (0.44–8.15) GtC with 90% of this total contained belowground. The large uncertainty reflects the need for more field-data.

231 In Africa, peatlands occur in many countries, but extensive peatlands are located in the
232 Rugezi Marsh in Rwanda, the Okavango Delta in Botswana, the Sudd catchment in Sudan and
233 in particular the Congo basin (Grundling and Grootjans, 2018). The Cuvette Centrale wetland
234 of the Congo basin is estimated to contain a C stock of 30.6 (6.3–46.8) Gt C (Bwangoy et al.,
235 2010; Dargie et al., 2017, Table 1). Again, note the large uncertainty range, which is a reflection
236 of the fact that this estimate is based on a relatively sparse set of field measurements (Dargie et
237 al., 2017). Peatlands have also been reported in southern Africa, mainly along the eastern coast
238 (Mozambique Coastal Plain) and in the central plateau (Grundling & Grobler, 2005; McWethy
239 et al., 2016).

240 The total area covered by peatlands in Southeast Asia is roughly 21 Mha (Xu et al., 2018).
241 Most of these peatlands are in Indonesia (15 Mha), Malaysia (2.2 Mha), Thailand (40 thousand
242 ha) and, to a lesser extent, Vietnam, Brunei and the Philippines. A recent estimate put the peat
243 C store in Indonesia alone at 28.1 (13.6–40.5) Gt C (Warren et al., 2017). Unlike other large
244 tropical peat reservoirs in the world that are either untouched or have had little alteration,
245 peatlands in Southeast Asia have faced intense anthropogenic disturbances since the 1970s,
246 when permission was granted to use these extensive areas for commercial purposes.

247 Due to the current large uncertainties around carbon stocks in tropical peatlands, it is
248 notorious that with the advance of knowledge in the identification of tropical peatlands (mainly
249 extension and depth) resulted in significantly higher estimates for carbon stocks in the tropical
250 zones. For South America and Africa the large uncertainty reflects the need for more field-data
251 (Dargie et al., 2017). Peatlands in relatively remote African and Amazonian regions currently
252 face low human intervention, however as anthropogenic activities, such as commercial
253 agriculture, exploitation of waters for hydropower (in Andes), forestry (including
254 deforestation), construction of impoundments, roads and ports, and gas exploration in peatlands
255 increase, so does the degradation of these ecosystems (Baker, 2014; Lahteenoja et al., 2009;

Roucoux et al., 2017). Therefore, decreasing uncertainties about area, and C stock in such remote regions is crucial to estimate the true C accumulation potential of these peatlands and to prevent future impact of human activities that peatlands may face mainly in South America and Africa.

267

268 Figure 1 – Estimated peat area (a), depth (b) and volume (c), presented by Page et al. (2011),

269 Gumbrecht et al. (2017) and Xu et al. (2018) of tropical peatlands. (d) Estimated carbon stock

270 (GtC) in tropical peatlands. Error bars are minimum and maximum estimates when available.

271 *Values estimated using peatland area from Xu et al. (2018) and mean depth from Gumbrecht

272 et al. (2017) and Page et al. (2011).

Figure 2 – Distribution of peatlands in tropical regions. Data from Xu et al., (2018)

Tropical peatland carbon accumulation, climate change and the global carbon cycle

The carbon accumulation rates in undisturbed tropical peatlands range from 24-300 gC m⁻² yr⁻¹ (Table 2), while for boreal and temperate undisturbed peatlands they are generally lower (vary from 2 to 271 gC m⁻² yr⁻¹, Olefeldt et al. 2012; Renou-Wilson et al. 2019). Although substantial variation occurs depending on peatland type, hydrology, vegetation type and peat formation (C. Craft, Washburn, & Parker, 2008; Sjögersten et al., 2014), C accumulation rates are, with a few exceptions, greater in the tropics and decrease with latitude (Sjögersten et al., 2014). Additionally, the carbon accumulation rates in undisturbed tropical peatlands are generally much higher than in intact old-growth tropical forests, commonly over mineral soils, in Africa and Amazonia (40-91 and 0-47 gC m² yr⁻¹, respectively) (Hubau et al., 2020).

The accumulation rates depend on the balance between carbon uptake by vegetation and carbon emitted to the atmosphere and lost to adjacent terrestrial or aquatic system. CO₂

emission vary greatly in tropical peatlands (250 and $13841 \text{ gC m}^{-2} \text{ yr}^{-1}$ (Table 3) and tend to be greater than in non-tropical systems ($411 \pm 128 \text{ gC m}^{-2} \text{ yr}^{-1}$) (Bubier, Bhatia, Moore, Roulet, & Lafleur, 2003; Clair, Arp, Moore, Dalva, & Meng, 2002; Crow & Wieder, 2005; Mäkiranta et al., 2009; Silvola, Alm, Ahlholm, Nykanen, & Martikainen, 1996). Estimated fluxes of CH_4 from peatlands are typically several orders of magnitude lower than those for CO_2 (Table 3). CH_4 emissions are indeed undetectable in some peatlands and an uptake from the atmosphere might occur instead (Sjögersten et al., 2014). Previous studies have estimated that undisturbed temperate and boreal environments emit moderate to high level of CH_4 (-7.1 – $2088.6 \text{ gC m}^{-2} \text{ yr}^{-1}$) (Inubushi, Furukawa, Hadi, Purnomo, & Tsuruta, 2003; Martikainen, Nykänen, Alm, & Silvola, 1995; Melling, Hatano, & Goh, 2005; Mitsch et al., 2010; Turetsky 2014), whereas CH_4 emissions from undisturbed tropical peatlands have been estimated at moderate range of -9.2 – $110.6 \text{ gC m}^{-2} \text{ yr}^{-1}$ (Table 3).

Methane formation is driven by methanogenic microorganisms activity (anaerobic decomposers) that degrades organic matter slowly in an anoxic environment (Mitsch et al., 2010). A peatland's capacity to emit less CH_4 appears to be a complex mechanism developed over several thousands of years, given that formerly human-disturbed restored peatlands in temperate systems with well-established vegetation and carbon stock have CH_4 emissions about 150% higher than older peatlands (Renou-Wilson et al., 2019). This fact suggests that to maintain low CH_4 emissions and higher carbon sequestration rates it is important to not only invest in actions that seek to recover impacted areas, but also to ensure that ecosystems are protected.

In the absence of direct human disturbance, many tropical peat deposits are actively accumulating carbon or are in steady states (Dargie et al., 2017; Fatoyinbo, 2017). However, climate change may significantly impact peatlands, and this relationship is poorly understood, particularly in the case of tropical peatlands, and thus the fate of peatlands under future change

remains uncertain (Frey & Smith, 2005; Gallego-Sala et al., 2018; Hapsari et al., 2017; Hirano et al., 2012; Hodgkins et al., 2018; Rieley et al., 2008). The effect of climate change will depend mainly on how temperature, total precipitation, sea level and frequency of extreme events will change in a specific region and how they will affect hydrology, vegetation composition and, consequently, primary production, substrate quality, decomposition process, lateral carbon fluxes and C accumulation rates of peatlands.

Some recent work, using Dynamic Global Vegetation Models (DGVMs) indicated, for Northern Hemisphere peatlands, a carbon sink twice as big than the 1861-2005 mean under two climate scenarios (defined by the RCPs 2.6 and 6.0), even though rapid climate change (under RCP8.5) might impact negatively the extent of northern peatlands, and the capacity of these areas to act as a carbon sink (Chaudhary et al, 2019; Qiu et al, 2020). As well, some models have predicted continued peat accumulation through to 2100 (Gallego-Sala et al., 2018; Spahni et al., 2013), while most models agree that there will be substantial losses over the next centuries (Avis, Weaver, & Meissner, 2011; Gallego-Sala et al., 2018; Ise et al., 2008), and some models have predicted that loss to start before 2100 (Avis, Weaver, & Meissner, 2011; Ise et al., 2008). Few pan-tropic modelling studies have been undertaken (Gallego-Sala et al., 2018; Treat et al., 2019) largely due to the sparsity of available data on tropical peatlands that is needed for model parametrization and validation. Climate models for the western Amazon predict increasing precipitation and river discharge over the century (Duffy et al., 2015; Sorribas et al., 2016; Zulkafli et al., 2016) whereas the opposite is predicted for the Eastern Amazon (Duffy et al., 2015; Sorribas et al., 2016), meaning that Brazilian peatlands are likely to be more vulnerable than those in Peru. However, a recent modelling study in the PMFB in Peru predicted that temperature increases would offset any positive effect of increased precipitation on peat accumulation by the end of the century through an increase in decomposition. i.e., Peruvian peatlands will cease peat accumulation despite increases in precipitation (Wang et al., 2018).

Likewise, in the Congo Basin there is a clear consensus that temperature will increase under all future scenarios while precipitation is predicted to increase under high emission scenarios and remain relatively unchanged under low emission scenarios (Haensler, Saeed, & Jacob, 2013) and, therefore, Congolese peatlands may also be vulnerable to future climate change. Likewise, studies in Southeast Asian peatlands, many of which are already degraded from deforestation and drainage, have been shown that additional carbon emission could also occur if dry seasons are extended or are more severe due to future climate change (Warren, et al., 2017).

Conversely, there is evidence to suggest that tropical peats may be more resistant to temperature changes. Hodgkins et al. (2018) observed that the higher aromatic content of tropical peat compared to the peat located at higher latitudes creates both a reduced oxidation state and higher recalcitrance, which prevents carbon release, even at high temperatures. In many peatlands in the northern hemisphere, deep peat has also high recalcitrance characteristics, which means that despite the expected temperature increases from climate change, the deep peat will probably remain stable, suggesting that these carbon stocks may be preserved in the face of climate change given their similar characteristics to tropical peat. Although there may only be a direct relationship between temperature and decomposition in high recalcitrant peat, it is recognized that changes in precipitation can alter the natural hydrology of these environments and enhance the degradation processes of recalcitrant peat (Chimner & Ewel, 2005).

In summary, many peatland areas are projected to stop accumulating peat by the end of the century and beyond, thus creating a positive climate feedback loop where further warming means C losses and, in turn, greater radiative forcing (Gallego-Sala et al., 2018). However, across the entire tropics, and particularly the Amazon and Congo basins, further field data is required to better parameterize and validate models so that we can improve projections of the future C balance in tropical peatlands, which at the moment remain highly uncertain.

Although climate change (such as changes to temperatures and precipitation) has an impact on the dynamics of these ecosystems, direct anthropogenic changes (LULC changes, drainage and deforestation) currently play a greater role. Therefore, understanding the impact of direct anthropogenic changes on these ecosystems can help us understand whether tropical peatlands are a net sink or net source in the global carbon cycle.

Direct human disturbances and their impacts on carbon accumulation rates and GHG emissions in tropical peatlands

Anthropogenic activities, such as logging, drainage, deforestation, fires and the conversion of native forests to agricultural lands, have been rapidly increasing in peatlands since the 1990s (Hooijer et al., 2010), particularly in developing countries, and have put these ecosystems at risk (Swindles et al., 2018). Although most of the scientific literature on the degradation processes of tropical peatlands focuses on Southeast Asia (Hapsari et al., 2017; Hirano, Jauhiainen, Inoue, & Takahashi, 2009; Hirano et al., 2012; Inubushi et al., 2003; Könönen, Jauhiainen, Laiho, Kusin, & Vasander, 2015; Rieley et al., 2008), the degradation of large areas of peat and the impacts that may alter their natural conditions have also been documented in both South America and Africa (Baker, 2014; Dargie et al., 2017; Dargie et al., 2019; Roucoux et al., 2017; Swindles et al., 2018).

In Southeast Asia, domestic and international demand for agricultural and forest products and services has put pressure on tropical peatlands and, by 2010, it was estimated that only 36% of the original peatland area in the Southeast Asia was covered by primary and secondary peat swamp forest (Miettinen, Shi, & Liew, 2012; Dohong, Aziz & Dargusch, 2017). In the Indonesian regions of Sumatra and Kalimantan, the two regions of Indonesia with the greatest impacts, only 6% were pristine peat swamp forests (Miettinen, Shi, & Liew, 2012). To meet the high demand of agricultural products, the peatlands have been subjected to deforestation (Hirano et al., 2012), widespread drainage (Fatoyinbo, 2017), and recurrent fires (Page et al.,

2002). Page et al. (2011) argues that, on the one hand, expansion of agriculture and forestry in the region has provided opportunities to industries and businesses, yet on the other hand it has also had, has also had negative environmental impacts. Between 2000 and 2010, Southeast Asia has had the highest annual rate of deforestation (rate of 2.2%) among all tropical humid regions in the world. This deforestation has resulted in the loss of 11 Mha of native forests and has led to significant changes in natural ecosystem dynamics, mainly related to carbon balance (Miettinen & Liew, 2010). Harris et al. (2013) projected land use and emissions from peatlands between 2010 and 2050 across Indonesia, Malaysia, and in Papua New Guinea and found that under the “business as usual” scenario, in which total production of oil palm will increase without peatland protection measures, the average annual CO₂ emissions would almost double between 2020 and 2050 (from 264 to 424 Tg CO₂ yr⁻¹). In contrast, restoring the peat to native forest vegetation (restoration scenario) would bring annual emissions close to zero.

In Africa, increased economic development could have a negative impact on peatlands through hydrocarbon exploration, logging, plantations and other forms of disturbance that significantly damage these ecosystems, although they are still intact today (Dargie et al., 2019). Additionally, land-use changes occur as a result of multiple complex and interacting environmental, economic and political factors, which can accelerate the negative impacts of human activities. In Cuvette Centrale region in Congo, rivers are the main transport network and there are relatively few roads. This, along with the large distance from any international port and low population densities, is among the reasons why the Congo basin peatlands have so far been spared from more severe degradation typical for Southeast Asian peatlands. Although limited in number, roads have already been constructed across some of the peatland areas of the Cuvette Centrale. No studies have yet considered the specific impacts of these roads on the peat properties, hydrology or vegetation; however, the observed swamp forest death following road construction suggests that roads could be having a negative impact on the

416 wetlands of the region (Dargie et al., 2017). The low level of human intervention in the Cuvette
417 Centrale peatlands at present suggests that there is still time to protect the peatlands in a largely
418 intact state, possibly by encouraging funding for mitigation of land-use change (Dargie et al.,
419 2019).

420 In South America, large areas of undisturbed peatlands are increasingly facing a range
421 of threats, including hydroelectricity (river damming) projects, road and railway projects (Finer
422 & Orta-Martínez, 2010; Gutiérrez-Vélez et al., 2011), ore, gas, and oil exploration, logging and
423 drainage for agriculture (Baker, 2014; Roucoux et al., 2017). Over exploitation of the palm fruit
424 (*Mauritia flexuosa* – commonly found in wetlands) is also an increasing concern (Kahn &
425 Mejia, 1990; Lilleskiv et al, 2019). In contrast with the better-known but highly degraded and
426 at-risk peatlands of Southeast Asia (Miettinen et al., 2012), many peatlands in South America
427 remain largely intact and the threat of destruction from direct human impacts is comparatively
428 low (Baker, 2014).

429 In general, the degradation process of tropical peatlands begins with the felling of natural
430 vegetation, which reduces the amount of biomass in the system (Könönen et al., 2016), and
431 promotes an increase of C oxidation rates and a reduction of soil moisture because of the
432 increased incidence of direct radiation (Dargie et al., 2019; Jauhiainen, Hooijer, & Page, 2012).
433 However to a lesser extent, a reduction in vegetation can also lead to increase in soil moisture
434 due to the decrease in transpiration (Porporato, Laio, Ridolfi, & Rodriguez-Iturbe, 2001). After
435 the deforestation process, the peatlands are artificially drained in order to reduce groundwater
436 levels to plant perennial and rotating crops (Dargie et al., 2019) are not adapted to the naturally
437 flooded environment. Next, aerial biomass crops are produced, which reduce the ecosystem
438 carbon uptake because the soil no longer has the environmental conditions of peatlands to
439 accumulate carbon, and the carbon accumulated by the crop primary production is removed
440 from the system through the harvest (Roucoux et al., 2017)

The most recent studies on GHG soil emissions from natural and impacted environments show that tropical peatlands have high CO₂ emissions in drained environments used for agricultural production and in recovering areas (Leifeld & Menichetti, 2018). Although non-impacted forests emit C through soil respiration, on average emissions are lower due to the maintenance of natural soil moisture conditions and groundwater levels.

In Indonesia, the carbon lost in peatlands after LULC changes has averaged approximately 60 Mg ha⁻¹ yr⁻¹ over 25 years of interference (Murdiyarso et al., 2010). This loss is, in part, due to the absence of vegetation in impacted and/or drained peatlands, given that in undisturbed peatlands C sequestration from vegetation cover offsets C emissions from the soil. In Southeast Asia, it is estimated that in 2006 CO₂ emissions from organic matter decomposition in drained peat soil were equivalent to 1% to 3% of all global CO₂ emissions from burning fossil fuels (~630 Mt), and that 82% of these emissions were from Indonesian peatlands (Hooijer et al., 2010). Other studies have pointed to even higher emission rates associated with peat decomposition in Indonesia, reaching about 8% of global emissions from burning fossil fuels (2000 Mt yr⁻¹ of CO₂, Rieley et al., 2008). Moreover, major events were reported in 1997 and 2015, in which widespread forest and peatland fires burned large areas of the Southeast Asia (Page et al. 2002; Huijnen et al., 2016), especially Indonesia, releasing large amounts of carbon land-based in the atmosphere, mainly in the form of CO₂, CO and CH₄. With an average emission rate of 11.3 Tg CO₂ per day during these events, emissions exceeded the European Union's (EU28) fossil fuel CO₂ release rate of 8.9 Tg CO₂ per day (Huijnen et al., 2016).

Methane fluxes also change as a result of human disturbance (Reay et al., 2018). The conversion of peatland forests to areas of intensive cultivation, along with significant inputs of nitrogen fertilizers, may alter the natural dynamics of methane and nitrous oxide emissions (Tian et al., 2015). Rice crops in Indonesia have shown very significant CH₄ emissions after

being converted from peatland forests (Table 3) because the production of CH₄ by methanogenic microorganisms is boosted by both the ever-flooded system and the use of nitrogen fertilizers (Conrad, 2002). Emissions from these crops may be about 20-fold greater than emissions from natural areas. Nitrous oxide emissions in Indonesia have been shown to increase substantially with land use change and the introduction of agricultural activities in peatlands (Oktarita et al., 2017). Nitrous oxide emissions from *Elaeis guineensis* (oil palm) monocultures in Indonesia were reported by Hadi et al. (2005) at 9.1 gC m⁻² years⁻¹, higher than those reported by Inubushi et al. (2003) in native peat forests, 1.25 gC m⁻² years⁻¹.

In addition to GHG emissions, drainage enables organic matter to be transported to adjacent watercourses in the form of dissolved organic carbon (DOC), particulate organic matter and dissolved inorganic matter. For instance, Baum et al. (2007) suggest that Indonesian rivers, particularly those receiving effluents drained from peatlands, transfer large amounts of carbon, in the form of DOC, to the oceans (21 Tg yr⁻¹) and that this accounts for approximately 10% of global riverine DOC inputs into the ocean (Rieley et al., 2008).

Roucoux et al. (2017) examined the services provided by large, intact tropical peatlands, the factors threatening them, and opportunities to conserve them, and cite that, although their contribution from tropical peatlands to climate regulation on the planet is evident, their importance is weakly articulated within existing conservation agendas, mainly because they are poorly described and mapped and are frequently unrecognized by local agencies and institutions. Fortunately, in Amazonia, Africa, and New Guinea tropical peatland ecosystems are also widespread and often much less intensively exploited. Many can be described as intact at the landscape scale; their hydrology is unaffected by human activity and their vegetation cover is not fragmented or substantially degraded.

490 Table 1 –Tropical peatland carbon stock (GtC) showing mean values and/or (range) if available.

System/Location	Land cover	Carbon stock (GtC)	Ref.
<i>Tropical Asia</i>		68.9 (66.6 – 70.4)	Page et al. (2011)
Central Kalimantan, South Sumatra and West Papua	Peat swamp forest	55 ± 10	Jaenicke et al. (2008)
Indonesia	Native Forest	23.2	Dommain et al. (2014)
Indonesia	Native vegetation and impacted areas	30	Rudiyanto et al. (2015)
Indonesia	Native vegetation and impacted areas	57.4	Page et al. (2011)
Malaysia	Native vegetation and impacted areas	9.1	Page et al. (2011)
Southeast Asia	Native vegetation and impacted areas	172	Sjögersten et al. (2014)
Southeast Asia	Native Forest	65	Dommain et al. (2011)
Southeast Asia	Native vegetation and impacted areas	20	Dommain et al. (2011), (2014)
<i>Tropical America</i>		12.7 (11.5 – 13.4)	Page et al. (2011)
Peru (Pastaza-Marañon)	Native Forest	3.14 (0.4 – 8.1)	Baker, (2014)
Peru (Pastaza-Marañon)	Native Forest	3.12 (0.8 - 9.5)	Lähteenoja et al. (2012)
<i>Tropical Africa</i>		6.9 (3.5 – 8.1)	Page et al. (2011)
Cuvette Centrale, Congo	Native Forest	30.6 (6.3 – 46.8)	Dargie et al. (2017)
<i>Global scenario</i>		(469 – 694)	Page et al. (2011); Yu et al (2010)
Tropical undisturbed		(139 – 251)	Miettinen & Liew (2010); Zoltai (1996)
Tropical disturbed		(13 – 37)	Kurnianto et al. (2015); Zoltai & Martikainen (1996)
Tropical		(152-288)	This study
Non-Tropical		(387 – 394)	Page et al. (2011); Gorham (1991); Immirzi & Maltby (1993); Gorham (1991)

491

492

493 Table 2 – Carbon accumulation rates (gC m⁻² yr⁻¹) from tropical peatlands and from non-tropical peatlands (for comparison) showing mean values
494 and/or (range) if available. Positive values are carbon accumulation and negative values are carbon loss.

System/Location	Land cover	Carbon accumulation rates (gC m ⁻² yr ⁻¹)	Ref.
<i>Tropical Asia</i>			
Central Kalimantan	Native Forest	22.3 (6.5 – 121.4)	Page et al. (2004)

Central Kalimantan	Native Forest	31.3 (16.6 – 73.2)	Dommain et al. (2011)
Central Sumatra, Indonesia	Secondary peat swamp forest	55	Hapsari et al. (2017)
Indonesia	Native Forest	72	Dommain et al. (2015)
Indonesia	Native Forest	94	Page et al. (2004)
Brunei (Borneo)	Peat swamp forest (mangrove forest)	300	Dommain et al.(2015)
Brunei (Borneo)	Peat swamp forest (<i>Shorea albida</i>)	50	Dommain et al.(2015)
Kalimantan Central	Drained peatlands and forest	85	Page et al. (2004)
Kalimantan, Indonesia	Peat swamp forest	94.3	Page et al. (2004)
Malaysia	Rain forest	(79 – 147)	Kosugi et al. (2008)
Riau, Sumatra	Native Forest	81 ± 1.4	Neuzil et al. (1997)
Sarawak, Malaysia	Undrained peat swamp forest	8.46 ± 0.51	Wong et al. (2020)
Sarawak, Malaysia	Relatively disturbed secondary peat swamp forest	4.17 ± 0.69	Wong et al. (2020)
Sarawak, Malaysia	oil palm plantation	2.19 ± 0.21	Wong et al. (2020)
Southeast Asia	Native Forest	(30 – 270)	Page et al. (2004)
Southeast Asia	Drained affected peat swamp forest	(-499 – -174)	Rieley et al. (2008)
West Kalimantan	Drained peatlands and forest	(74 – 85)	Neuzil et al. (1997)
<i>Tropical America</i>			
Amazonia	Amazonian forests without El Niño event	-100	Saleska et al. (2003)
Amazonia	Amazonian forests with El Niño event	(100 – 200)	Saleska et al. (2003)
Amazonian peatlands	Forested peatland	(26 – 195)	Lähteenoja et al. (2009)
Costa Rica	Fragments of Yolillo (<i>Raphia</i>)	(250 – 260)	Mitsch eta al. (2010)
French Guiana	Pristine tropical rain forest	-138	Bonalet et al. (2008)
Peru (Pastaza-Marañon)	Native Forest	52 ± 22 (36 – 85)	Lähteenoja et al. (2009)
Peru (Pastaza-Marañon)	Native Forest	(28 – 108)	Lähteenoja et al. (2012)
Cayambre-Coca	Peatlands in the Andes Mountains	51.1	Chimner & Ewel (2005)
<i>Tropical Africa</i>			
Cuvette Centrale, Congo	Native Forest	23.9 ± 5.8 (18.3 – 33.1)	Dargie et al. (2017)
Kenya	Tropical papyrus peatland	160	Jones and Humphries (2002)
Burundi	Buyongwe Swamp	125	Panujen (1996)
Burundi	Ndurumu Swamp	65	Panujen (1996)
Rwanda	Cyili Swamp	113	Panujen (1996)
Rwanda	Gishoma Swamp	(86-106)	Panujen (1996)
Rwanda	Mashya Bog	91	Panujen (1996)
Rwanda	Cyabaralika Swamp	33	Panujen (1996)
Rwanda	Kiguhu Swamp	31	Panujen (1996)

Tropical undisturbed
Tropical disturbed
Non-Tropical

(24 – 300)
(-499 – -174)
(-248 – 271)

Kurnianto et al. (2015); Chimner (2004)
Rieley et al. (2008)
Roulet et al. (2007); Olefeldt et al. (2012);
Renou-Wilson et al. (2019)

495

496 Table 3 – Estimates of soil emission of CO₂ and CH₄ from peatlands (gC m⁻² yr⁻¹) showing mean values and/or (range) if available. *Values of
497 CO₂ emission are from autotrophic and heterotrophic respiration and do not consider primary production. *Positive values of soil emission mean
498 carbon emission and negative mean carbon uptake.

499

System/Location	Land cover	CO ₂ *	CH ₄ **	Ref.
		(gC m ⁻² yr ⁻¹)		
Tropical Asia				
Malaysia	Forested peatland		0.02 (-0.05 – 0.10)	Melling et al. (2005)
Thailand	Forest peatland		9.81 ± 23.6 (1.7 – 110.4)	Ueda et al. (2000)
Central Kalimantan	Peat swamp forest floor	3493 ± 316	1.36 ± 0.57	Jauhiainen et al. (2004)
Indonesia	Peat swamp forest floor		1.35	Jauhiainen et al. (2004)
Indonesia	Poorly drained forest	174 ± 203		Hirano et al. (2012)
Indonesia	Drained forest	328 ± 204		Hirano et al. (2012)
Indonesia	Burnt and drained forest	499 ± 72		Hirano et al. (2012)
Indonesia	Tropical peatlands (including rice)		(4.4 – 19.3)	Hadi et al.(2005)
Kalimantan, Indonesia	Forested peatland	2777 ± 8322		Hirano et al.(2009)
Kalimantan, Indonesia	Secondary forest	4494	1.66	Hadi et al. (2005)
Kalimantan, Indonesia	Secondary forest	3460 (1603 - 35522)	4.4 (0 – 29)	Hadi et al. (2005)
Kalimantan, Indonesia	Forested peatland	3495 ± 315 (438 - 4818)	1.4 ± 5.7 (-0.09 – 3.1)	Jauhiainen et al. (2004)
Kalimantan, Indonesia	Forested peatland	4932 (692 - 13841)		Sundari et al. (2012)
Kalimantan, Indonesia	Forested peatland		9.6 ± 5.3	Inubushi et al. (1998)
Malaysia	Forested peatland	3889		Murayama and Bakar (1996)
Micronesia	Forested peatland	3469 ± 315 (2978 - 3522)		Chimner 2004
Sarawak, Malásia	Sago	(552 – 2146)		Melling et al. (2005)
Sarawak, Malásia	Oil palm	(403 – 29334)		Melling et al. (2005)
Sarawak, Malaysia	Forest ecosystem	(876 – 4669)		Melling et al. (2005)
South Kalimantan	Secondary forest	1200 ± 430	1.2 ± 0.4	Inubushi et al. (2003)
South Kalimantan	Secondary forest peatland to paddy field	(1200 – 1500)	(1.2 – 1.9)	Inubushi et al. (2003)
South Kalimantan	Changing land-use from Secondary forest to upland tended	(1000 – 2000)	(1.2 – 0.6)	Inubushi et al. (2003)

South Kalimantan	Abandoned upland crops field	990 ± 110	0.6 ± 0.7	Inubushi et al. (2003)
South Kalimantan	Abandoned paddy fields	1540 ± 290	1.9 ± 0.5	Inubushi et al. (2003)
Southeast Asia	Secondary Native Forest	3460	4.4	Hadi et al. (2005)
Southeast Asia	Secondary forest	3500	0.5	Hadi et al. (2005)
Southeast Asia	forest ecosystem	2100		Melling et al. (2005)
Southeast Asia	Lowland peatlands	250	1.09	Couwenberg et al. (2010)
Southeast Asia	Undrained peat swamp forest	3892 ± 304	1.36 ± 0.57	Rieley et al. (2008)
Southeast Asia	Drained uncultivated agricultural land	1928 ± 526	0.12 ± 0.09	Rieley et al. (2008)
Southeast Asia	Drained affected peat swamp forest	4000 ± 1091	1.3 ± 0.98	Rieley et al. (2008)
Southeast Asia	Burned areas	2900		Hadi et al. (2005)
Southeast Asia	Converting peat swamp forests into oil palm	5940		Murdiyarso et al. (2010)
Southeast Asia	Paddy field	1389	19.6	Radjagukguk (1997)
Southeast Asia	Rice-soybean rotation field	2019	2.6	Bouwman (1990)
Southeast Asia	Paddy field	1400	1.4	Hadi et al. (2005)
Southeast Asia	Rice-soybean rotation field	2000		Hadi et al. (2005)
Southeast Asia	Oil palm	1500		Melling et al. (2005)
Southeast Asia	Sago	1100		Melling et al. (2005)
Southeast Asia	Cultivation of palm oil (<i>Elaeisis guineensis</i>)	5940		Murdiyarso et al. (2010)
Southeast Asia	Rice crops (Mega Rice Project)	2178		Hadi et al. (2005)
Southeast Asia	Rice crops	1389	26.6	Hadi et al. (2005)
Southeast Asia	Agricultural Soils	2019	1.7	Hadi et al. (2005)
Southeast Asia	Horticulture	1500	1.9	Inubushi et al. (2003)
Southeast Asia	Cultures with nitrogen fertilization (<i>Acacia</i> sp e <i>Metroxylon sagu</i>)	2130	2.6	Couwenberg et al. (2010)
Southeast Asia	Clear felled recovering peat Swamp forest	3400 ± 927	2 ± 1.5	Rieley et al. (2008)
Sumatra, Indonesia	Forested peatland	3329 ± 481.8	7.8 ± 4.2	Furukawa et al. (2005)
Sumatra, Indonesia	Forested peatland	2435 ± 140	10.6 ± 11.9	Furukawa et al. (2005)
Sumatra, Indonesia	Forested peatland	3294 ± 937	6.7 ± 2.4	Furukawa et al. (2005)
Sumatra, Indonesia	Natural swamp forest drained more than 5 years	267		Jauhiainen et al. (2012)
<i>Tropical America</i>				
Bocas del Toro, Panama	Forested peatland (<i>Raphia</i> sp.)	1857 (96 – 14839)	(1.1 – 110.6)	Wright et al. (2011)
Bocas del Toro, Panama	Forested peatland (<i>Camptosperma</i> sp.)	2085 (543 – 7017)	(-7.7 – 31.8)	Wright et al. (2011)
Bocas del Toro, Panama	Open peatland (<i>Cyperus</i> sp.)	1269 (61 – 8322)	(-9.3 – 27.2)	Wright et al. (2011)
Ka’au, Hawaii	Montane swamp	1112.5 ± 412		Chimner (2004)
Mauim, Hawaii	Montane peatland	2497 ± 657		Chimner (2004)
Orinoco Llanos, Venezuela	Palm peatland	263 (149 – 473)		Bracho & San José (1990)
Brazil (Lowlands in São Paulo state)	Pastureland (dry season and wet season)	3210	(-5.2 – 4.0)	Ribeiro et al. (2018)

Brazil (Lowlands in São Paulo state)	Native forest (dry season and wet season)	2174	(-3.1 – 4.2)	Ribeiro et al. (2018)
Brazil (Lowlands in São Paulo state)	Irrigated rice crop (dry season and wet season)	2074	3.1	Ribeiro et al. (2018)
<i>Global scenario</i>				
Tropical undisturbed		(250 – 13841)	(-9.3 – 110.6)	Kurnianto et al. (2015); Chimner (2004)
Tropical disturbed		(263 – 29334)	(1.9 – 26.6)	Kurnianto et al. (2015); Zoltai & Martikainen (1996)
Non-Tropical		411 ± 128	35.1 ± 2.6 (-7.1 – 2088.6)	Lund et al.(2010)

500

501

502 Final remarks

503 Tropical peatlands are different from boreal and temperate peatlands, particularly their
504 climatic settings, peat matter formation, and vegetation coverage. They cover 90-170 Mha
505 which represents 23% to 30% of the total area covered by peatlands throughout the world.
506 Bringing together the most up-to-date estimates of peatland area, peat depth, peat volume and
507 peat carbon content, we estimate that tropical peatlands store 152-288 Gt of carbon, which is
508 significantly higher than the previously reported values. The large uncertainty in these estimates
509 is related to methodological approach and the sparse field data on depth and C content, mainly
510 in South America and Africa. Despite tropical peatlands covering a smaller area and storing
511 less carbon than non-tropical peatlands, carbon accumulation rates are greater in the tropics and
512 decrease with latitude, which gives tropical peatlands the important role of accumulating carbon
513 emitted by human activities now and in the future.

514 Climate change is a threat to peatlands, but at local and regional levels, direct human
515 interventions have played a more important role in impairing the capacity of peatlands to
516 sequester carbon. In the tropical zone, the carbon sequestration rate of peatlands in Southeast
517 Asia has changed from 79-300 (uptake) to (-499) – (-174) g m⁻² yr⁻¹ (emission) after direct
518 human interference. Integrated development and management mechanisms supported by strong
519 policies and meaningful incentives can balance this scenario and contribute to more effective
520 measures.

521 The Amazon region potentially holds the largest natural peatland across the tropics.
522 However, the factors that contribute to tropical peatland degradation are only well understood
523 for Southeast Asian peatlands. To date, very few studies have addressed the impacts of peatland
524 degradation in South America and in Africa. Thus, advancing to fill the current scientific
525 knowledge gaps and incorporating local understanding is crucial for supporting policies geared
526 toward managing and conserving undisturbed peatlands in these regions. Furthermore,

understanding and mapping peatlands in Brazil by encouraging research projects can enhance the current knowledge about the potential of these system to uptake and store carbon, and can encourage actions aimed to protect peatlands in this region. Due to the high level of conservation and the expected high capacity of carbon accumulation, the Amazon region peatlands are particularly important in the context of climate change mitigation.

Acknowledgement

We thank the financial support of the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) through the program AUXPE-PROEX (number 0711/2018) and the São Paulo Research Foundation (FAPESP, project number 15/24201-0 and Nexus Project). The authors are grateful to Dr. G. Dargie for commenting on a draft and reorganize the manuscript and to Dr. M. Miranda for her valuable help with data and references.

Data Sharing and Accessibility

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

References

- Avis, C. A., Weaver, A. J., & Meissner, K. J. (2011). Reduction in areal extent of high-latitude wetlands in response to permafrost thaw. *Nature Geoscience*, 4(7), 444–448. <https://doi.org/10.1038/ngeo1160>
- Baker, F. C. D. and K. H. R. and I. T. L. and E. T. A. M. and E. N. H. C. and O. L. and L. T. M. and E. V. S. and R. Z. and T. R. (2014). The distribution and amount of carbon in the largest peatland complex in Amazonia. *Environmental Research Letters*, 9(12), 124017. Retrieved from <http://stacks.iop.org/1748-9326/9/i=12/a=124017>
- Baum, A., Rixen, T., & Samiaji, J. (2007). Relevance of peat draining rivers in central Sumatra for the riverine input of dissolved organic carbon into the ocean. *Estuarine, Coastal and Shelf Science*, 73(3), 563–570.

- 555 <https://doi.org/http://dx.doi.org/10.1016/j.ecss.2007.02.012>
- 556 Biancalani, R., & Avagyan, A. (2014). Towards climate-responsible peatlands management.
- 557 *Mitigation of Climate Change in Agriculture Series (MICCA)*, (9).
- 558 Blodau, C. (2002). Carbon cycling in peatlands □ A review of processes and controls.
- 559 *Environmental Reviews*, 10(2), 111–134. <https://doi.org/10.1139/a02-004>
- 560 BONAL, D., BOSC, A., PONTON, S., GORET, J.-Y., BURBAN, B., GROSS, P., ...
- 561 GRANIER, A. (2008). Impact of severe dry season on net ecosystem exchange in the
- 562 Neotropical rainforest of French Guiana. *Global Change Biology*, 14(8), 1917–1933.
- 563 <https://doi.org/10.1111/j.1365-2486.2008.01610.x>
- 564 Bourbonniere, R. A. (2009). Review of water chemistry research in natural and disturbed
- 565 peatlands. *Canadian Water Resources Journal*, 34, 393+.
- 566 Bouwman, A. F. (1990). Global distribution of the major soils and land cover types. *Soils and*
- 567 *the Greenhouse Effect*, 33–59.
- 568 Bracho, R., & San José, J. (1990). Energy fluxes in a morichal (swamp palm community) at the
- 569 Orinoco Llanos, Venezuela: microclimate, water vapour and CO₂ exchange.
- 570 *Photosynthetica*, 24(3), 468–494.
- 571 Bubier, J. L., Bhatia, G., Moore, T. R., Roulet, N. T., & Lafleur, P. M. (2003). Spatial and
- 572 Temporal Variability in Growing-Season Net Ecosystem Carbon Dioxide Exchange at a
- 573 Large Peatland in Ontario, Canada. *Ecosystems*, 6(4), 353–367.
- 574 <https://doi.org/10.1007/s10021-003-0125-0>
- 575 Bwangoy, J.-R. B., Hansen, M. C., Roy, D. P., Grandi, G. De, & Justice, C. O. (2010). Wetland
- 576 mapping in the Congo Basin using optical and radar remotely sensed data and derived
- 577 topographical indices. *Remote Sensing of Environment*, 114(1), 73–86.
- 578 <https://doi.org/http://dx.doi.org/10.1016/j.rse.2009.08.004>
- 579 Chimner, R.A. (2004). Soil respiration rates of tropical peatlands in Micronesia and Hawaii.
- 580 *Wetlands*, 24(1), 51–56. Retrieved from [http://www.bioone.org/bioone/?request=get-](http://www.bioone.org/bioone/?request=get-journals-list&issn=0277-5212)
- 581 [journals-list&issn=0277-5212](http://www.bioone.org/bioone/?request=get-journals-list&issn=0277-5212)
- 582 Chimner, Rodney A, & Ewel, K. C. (2005). A Tropical Freshwater Wetland: II. Production,
- 583 Decomposition, and Peat Formation. *Wetlands Ecology and Management*, 13(6), 671–684.
- 584 <https://doi.org/10.1007/s11273-005-0965-9>
- 585 Clair, T. A., Arp, P., Moore, T. R., Dalva, M., & Meng, F.-R. (2002). Gaseous carbon dioxide
- 586 and methane, as well as dissolved organic carbon losses from a small temperate wetland
- 587 under a changing climate. *Environmental Pollution*, 116, S143–S148.
- 588 [https://doi.org/https://doi.org/10.1016/S0269-7491\(01\)00267-6](https://doi.org/https://doi.org/10.1016/S0269-7491(01)00267-6)

- Clymo, R. S. (1987). The ecology of peatlands. *Science Progress (1933-)*, 71(4 (284)), 593–614. Retrieved from <http://www.jstor.org/stable/43420701>
- Conrad, R. (2002). Conrad R.. Control of microbial methane production in wetland rice fields [Review]. *Nutr Cycl Agroecosyst* 64: 59-69. *Nutrient Cycling in Agroecosystems*, 64, 59–69. <https://doi.org/10.1023/A:1021178713988>
- COUWENBERG, J., DOMMAIN, R., & JOOSTEN, H. (2010). Greenhouse gas fluxes from tropical peatlands in south-east Asia. *Global Change Biology*, 16(6), 1715–1732. <https://doi.org/10.1111/j.1365-2486.2009.02016.x>
- Craft, C. B. T.-C. and R. W. (Ed.). (2016). *Index*. <https://doi.org/https://doi.org/10.1016/B978-0-12-407232-9.18001-7>
- Craft, C., Washburn, C., & Parker, A. (2008). *Latitudinal Trends in Organic Carbon Accumulation in Temperate Freshwater Peatlands BT - Wastewater Treatment, Plant Dynamics and Management in Constructed and Natural Wetlands* (J. Vymazal, Ed.). https://doi.org/10.1007/978-1-4020-8235-1_3
- Crow, S. E., & Wieder, R. K. (2005). SOURCES OF CO₂ EMISSION FROM A NORTHERN PEATLAND: ROOT RESPIRATION, EXUDATION, AND DECOMPOSITION. *Ecology*, 86(7), 1825–1834. <https://doi.org/10.1890/04-1575>
- Dargie, G. C., Lawson, I. T., Rayden, T. J., Miles, L., Mitchard, E. T. A., Page, S. E., ... Lewis, S. L. (2019). Congo Basin peatlands: threats and conservation priorities. *Mitigation and Adaptation Strategies for Global Change*, 24(4), 669–686. <https://doi.org/10.1007/s11027-017-9774-8>
- Dargie, G. C., Lewis, S. L., Lawson, I. T., Mitchard, E. T. A., Page, S. E., Bocko, Y. E., & Ifo, S. A. (2017). Age, extent and carbon storage of the central Congo Basin peatland complex. *Nature*, 542(7639), 86–90. Retrieved from <http://dx.doi.org/10.1038/nature21048>
- Dommain, R., Cobb, A., Joosten, H., Glaser, P., Chua, A., Gandois, L., ... Harvey, C. (2015). Forest dynamics and tip-up pools drive pulses of high carbon accumulation rates in a tropical peat dome in Borneo (Southeast Asia). *Journal of Geophysical Research: Biogeosciences*, 120. <https://doi.org/10.1002/2014JG002796>
- Dommain, R., Couwenberg, J., & Joosten, H. (2011). Development and carbon sequestration of tropical peat domes in south-east Asia: Links to post-glacial sea-level changes and Holocene climate variability. *Quaternary Science Reviews - QUATERNARY SCI REV*, 30, 999–1010. <https://doi.org/10.1016/j.quascirev.2011.01.018>
- Duffy, P. B., Brando, P., Asner, G. P., & Field, C. B. (2015). Projections of future meteorological drought and wet periods in the Amazon. *Proceedings of the National*

- 623 *Academy of Sciences*, 112(43), 13172 LP – 13177.
 624 <https://doi.org/10.1073/pnas.1421010112>
- 625 Fatoyinbo, L. (2017). Ecology: Vast peatlands found in the Congo Basin. *Nature*, 542(7639),
 626 38–39. Retrieved from <http://dx.doi.org/10.1038/542038b>
- 627 Finer, M., & Orta-Martínez, M. (2010). A second hydrocarbon boom threatens the Peruvian
 628 Amazon: trends, projections, and policy implications. *Environmental Research Letters*,
 629 5(1), 14012. <https://doi.org/10.1088/1748-9326/5/1/014012>
- 630 Finlayson, C. M., & Milton, G. R. (2018). *Peatlands BT - The Wetland Book: II: Distribution,*
 631 *Description, and Conservation* (C. M. Finlayson, G. R. Milton, R. C. Prentice, & N. C.
 632 Davidson, Eds.). https://doi.org/10.1007/978-94-007-4001-3_202
- 633 Frey, K. E., & Smith, L. C. (2005). Amplified carbon release from vast West Siberian peatlands
 634 by 2100. *Geophysical Research Letters*, 32(9), n/a-n/a.
 635 <https://doi.org/10.1029/2004GL022025>
- 636 Furukawa, Y., Inubushi, K., Ali, M., Itang, A. M., & Tsuruta, H. (2005). Effect of changing
 637 groundwater levels caused by land-use changes on greenhouse gas fluxes from tropical
 638 peat lands. *Nutrient Cycling in Agroecosystems*, 71(1), 81–91.
 639 <https://doi.org/10.1007/s10705-004-5286-5>
- 640 Gallego-Sala, A. V, Charman, D. J., Brewer, S., Page, S. E., Prentice, I. C., Friedlingstein, P.,
 641 ... Zhao, Y. (2018). Latitudinal limits to the predicted increase of the peatland carbon sink
 642 with warming. *Nature Climate Change*, 8(10), 907–913. [https://doi.org/10.1038/s41558-](https://doi.org/10.1038/s41558-018-0271-1)
 643 [018-0271-1](https://doi.org/10.1038/s41558-018-0271-1)
- 644 Gandois, L., Cobb, A., Lim, I., Kamariah, A. S., & Harvey, C. (2012). Impact of deforestation
 645 on solid and dissolved organic matter characteristics of tropical peat forests: Implications
 646 for carbon release. *Biogeochemistry*, 114, 1–17. [https://doi.org/10.1007/s10533-012-](https://doi.org/10.1007/s10533-012-9799-8)
 647 [9799-8](https://doi.org/10.1007/s10533-012-9799-8)
- 648 Gandois, L., Teisserenc, R., Cobb, A., Hei Ing, C., Lim, L., Kamariah, A. S., ... Harvey, C.
 649 (2014). Origin, composition, and transformation of dissolved organic matter in tropical
 650 peatlands. *Geochimica et Cosmochimica Acta*, 137, 35–47.
 651 <https://doi.org/10.1016/j.gca.2014.03.012>
- 652 Gillman, L. N., Wright, S. D., Cusens, J., McBride, P. D., Malhi, Y., & Whittaker, R. J. (2015).
 653 Latitude, productivity and species richness. *Global Ecology and Biogeography*, 24(1),
 654 107–117. <https://doi.org/10.1111/geb.12245>
- 655 Gorham, E. (1991). Northern Peatlands: Role in the Carbon Cycle and Probable Responses to
 656 Climatic Warming. *Ecological Applications*, 1(2), 182–195.

<https://doi.org/10.2307/1941811>

- Grundlin, P.-L., & Glober, R. (2005). Peatlands and Mires of South Africa. *Biologiezentrum Linz/Austria*, (35), 379–396. Retrieved from www.biologiezentrum.at
- Grundling, P.-L., & Grootjans, A. P. (2018). *Peatlands of Africa BT - The Wetland Book: II: Distribution, Description, and Conservation* (C. M. Finlayson, G. R. Milton, R. C. Prentice, & N. C. Davidson, Eds.). https://doi.org/10.1007/978-94-007-4001-3_112
- Gumbrecht, T., Roman-Cuesta, R. M., Verchot, L., Herold, M., Wittmann, F., Householder, E., ... Murdiyarso, D. (2017). An expert system model for mapping tropical wetlands and peatlands reveals South America as the largest contributor. *Global Change Biology*, 23(9), 3581–3599. <https://doi.org/10.1111/gcb.13689>
- Gutiérrez-Vélez, V. H., DeFries, R., Pinedo-Vásquez, M., Uriarte, M., Padoch, C., Baethgen, W., ... Lim, Y. (2011). High-yield oil palm expansion spares land at the expense of forests in the Peruvian Amazon. *Environmental Research Letters*, 6(4), 44029. <https://doi.org/10.1088/1748-9326/6/4/044029>
- Hadi, A., Inubushi, K., Furukawa, Y., Purnomo, E., Rasmadi, M., & Tsuruta, H. (2005). Greenhouse gas emissions from tropical peatlands of Kalimantan, Indonesia. *Nutrient Cycling in Agroecosystems*, 71(1), 73–80. <https://doi.org/10.1007/s10705-004-0380-2>
- Haensler, A., Saeed, F., & Jacob, D. (2013). Assessing the robustness of projected precipitation changes over central Africa on the basis of a multitude of global and regional climate projections. *Climatic Change*, 121(2), 349–363. <https://doi.org/10.1007/s10584-013-0863-8>
- Hapsari, K. A., Biagioni, S., Jennerjahn, T. C., Reimer, P. M., Saad, A., Achnopha, Y., ... Behling, H. (2017). Environmental dynamics and carbon accumulation rate of a tropical peatland in Central Sumatra, Indonesia. *Quaternary Science Reviews*, 169, 173–187. <https://doi.org/https://doi.org/10.1016/j.quascirev.2017.05.026>
- Harenda, K., Lamentowicz, M., Samson, M., & Chojnicki, B. (2018). The Role of Peatlands and Their Carbon Storage Function in the Context of Climate Change. In *GeoPlanet: Earth and Planetary Sciences* (pp. 169–187). https://doi.org/10.1007/978-3-319-71788-3_12
- Hirano, T., Jauhiainen, J., Inoue, T., & Takahashi, H. (2009). Controls on the Carbon Balance of Tropical Peatlands. *Ecosystems*, 12(6), 873–887. <https://doi.org/10.1007/s10021-008-9209-1>
- Hirano, T., Segah, H., Kusin, K., Limin, S., Takahashi, H., & Osaki, M. (2012). Effects of disturbances on the carbon balance of tropical peat swamp forests. *Global Change Biology*, 18(11), 3410–3422. <https://doi.org/10.1111/j.1365-2486.2012.02793.x>

- 691 Hodgkins, S. B., Richardson, C. J., Dommain, R., Wang, H., Glaser, P. H., Verbeke, B., ...
 692 Chanton, J. P. (2018). Tropical peatland carbon storage linked to global latitudinal trends
 693 in peat recalcitrance. *Nature Communications*, 9. [https://doi.org/10.1038/s41467-018-](https://doi.org/10.1038/s41467-018-06050-2)
 694 06050-2
- 695 Hooijer, A., Page, S., Canadell, J. G., Silvius, M., Kwadijk, J., Wosten, H., & Jauhiainen, J.
 696 (2010). *Current and future CO2 emissions from drained peatlands in Southeast Asia*. 7,
 697 1505–1514. Retrieved from <http://edepot.wur.nl/160538>
- 698 Hubau, W., Lewis, S. L., Phillips, O. L., Affum-Baffoe, K., Beeckman, H., Cuní-Sanchez, A.,
 699 ... Zemagho, L. (2020). Asynchronous carbon sink saturation in African and Amazonian
 700 tropical forests. *Nature*, 579(7797), 80–87. <https://doi.org/10.1038/s41586-020-2035-0>
- 701 Huijnen, V., Wooster, M. J., Kaiser, J. W., Gaveau, D. L. A., Flemming, J., Parrington, M., ...
 702 van Weele, M. (2016). Fire carbon emissions over maritime southeast Asia in 2015 largest
 703 since 1997. *Scientific Reports*, 6(1), 26886. <https://doi.org/10.1038/srep26886>
- 704 Immirzi, C. ., & Maltby, E. (1993). The Global Status of Peatlands and their Role in Carbon
 705 Cycling, a report for Friends of the Earth by the Wetland Ecosystem. *Oryx*, 27(2), 127.
 706 <https://doi.org/DOI: 10.1017/S0030605300020755>
- 707 INGRAM, H. A. P. (1978). SOIL LAYERS IN MIRES: FUNCTION AND TERMINOLOGY.
 708 *Journal of Soil Science*, 29(2), 224–227. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2389.1978.tb02053.x)
 709 2389.1978.tb02053.x
- 710 International Peatland Society (IPS). (n.d.). Types of peatlands. Retrieved from
 711 <https://peatlands.org/peatlands/types-of-peatlands/>
- 712 Inubushi, K., Furukawa, Y., Hadi, A., Purnomo, E., & Tsuruta, H. (2003). Seasonal changes of
 713 CO₂, CH₄ and N₂O fluxes in relation to land-use change in tropical peatlands located in
 714 coastal area of South Kalimantan. *Chemosphere*, 52(3), 603–608.
 715 [https://doi.org/http://dx.doi.org/10.1016/S0045-6535\(03\)00242-X](https://doi.org/http://dx.doi.org/10.1016/S0045-6535(03)00242-X)
- 716 Inubushi, K., Hadi, A., Okazaki, M., & Yonebayashi, K. (1998). Effect of converting wetland
 717 forest to sago palm plantations on methane gas flux and organic carbon dynamics in
 718 tropical peat soil. *Hydrological Processes*, 12(13-14), 2073–2080.
 719 [https://doi.org/10.1002/\(SICI\)1099-1085\(19981030\)12:13/14<2073::AID-](https://doi.org/10.1002/(SICI)1099-1085(19981030)12:13/14<2073::AID-HYP720>3.0.CO;2-K)
 720 HYP720>3.0.CO;2-K
- 721 Ise, T., Dunn, A. L., Wofsy, S. C., & Moorcroft, P. R. (2008). High sensitivity of peat
 722 decomposition to climate change through water-table feedback. *Nature Geoscience*, 1(11),
 723 763–766. <https://doi.org/10.1038/ngeo331>
- 724 Jaenicke, J., Rieley, J. O., Mott, C., Kimman, P., & Siegert, F. (2008). Determination of the

amount of carbon stored in Indonesian peatlands. *Geoderma*, 147(3), 151–158.
<https://doi.org/http://dx.doi.org/10.1016/j.geoderma.2008.08.008>

Jauhiainen, J., Hooijer, A., & Page, S. E. (2012). Carbon dioxide emissions from an *Acacia* plantation on peatland in Sumatra, Indonesia. *Biogeosciences*, 9(2), 617–630.
<https://doi.org/10.5194/bg-9-617-2012>

Jauhiainen, Jyrki, Jaya, A., Inoue, T., Heikkinen, J. E. P., Martikainen, P., & Vasander, H. (2004). Carbon balance in managed tropical peat in central Kalimantan, Indonesia. *Proc. 12th Int. Peat Congr*, 653–659.

Jones, M. B., & Humphries, S. W. (2002). Impacts of the C4 sedge *Cyperus papyrus* L. on carbon and water fluxes in an African wetland. *Hydrobiologia*, 488(1), 107–113.
<https://doi.org/10.1023/A:1023370329097>

Joosten, H. (2009). *The Global Peatland CO2 Picture: peatland status and drainage related emissions in all countries of the world*. (p. 35 pp.). p. 35 pp. Wageningen: Wetlands International.

Joseph, H. (2005). Peatland hydrology and carbon release: why small-scale process matters. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 363(1837), 2891–2913. <https://doi.org/10.1098/rsta.2005.1671>

Kahn, F., & Mejia, K. (1990). Palm communities in wetland forest ecosystems of Peruvian Amazonia. *Forest Ecology and Management*, 33–34, 169–179.
[https://doi.org/https://doi.org/10.1016/0378-1127\(90\)90191-D](https://doi.org/https://doi.org/10.1016/0378-1127(90)90191-D)

Kelly (Letcher), R. A., Jakeman, A. J., Barreteau, O., Borsuk, M. E., ElSawah, S., Hamilton, S. H., ... Voinov, A. A. (2013). Selecting among five common modelling approaches for integrated environmental assessment and management. *Environmental Modelling & Software*, 47, 159–181. <https://doi.org/https://doi.org/10.1016/j.envsoft.2013.05.005>

Kelly, T. J., Lawson, I. T., Roucoux, K. H., Baker, T. R., Jones, T. D., & Sanderson, N. K. (2017). The vegetation history of an Amazonian domed peatland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 468, 129–141.
<https://doi.org/https://doi.org/10.1016/j.palaeo.2016.11.039>

Könönen, M., Jauhiainen, J., Laiho, R., Kusin, K., & Vasander, H. (2015). Physical and chemical properties of tropical peat under stabilised land uses. *Mires and Peat*, 16, Article 8.

Könönen, M., Jauhiainen, J., Laiho, R., Spetz, P., Kusin, K., Limin, S., & Vasander, H. (2016). Land use increases the recalcitrance of tropical peat. *Wetlands Ecology and Management*, 24(6), 717–731. <https://doi.org/10.1007/s11273-016-9498-7>

- 759 Kosugi, Y., Takanashi, S., Ohkubo, S., Matsuo, N., Tani, M., Mitani, T., ... Nik, A. R. (2008).
760 CO₂ exchange of a tropical rainforest at Pasoh in Peninsular Malaysia. *Agricultural and*
761 *Forest Meteorology*, 148(3), 439–452.
762 <https://doi.org/https://doi.org/10.1016/j.agrformet.2007.10.007>
- 763 Kurnianto, S., Warren, M., Talbot, J., Kauffman, B., Murdiyarso, D., & Frohking, S. (2015).
764 Carbon accumulation of tropical peatlands over millennia: a modeling approach. *Global*
765 *Change Biology*, 21(1), 431–444. <https://doi.org/10.1111/gcb.12672>
- 766 Lhteenoja, O., Flores, B., & Nelson, B. (2013). Tropical Peat Accumulation in Central
767 Amazonia. *Wetlands*, 33(3), 495–503. <https://doi.org/10.1007/s13157-013-0406-0>
- 768 Lhteenoja, O., Retegui, Y. R., Rsnen, M., Torres, D. D. C., Oinonen, M., & Page, S. (2012).
769 The large Amazonian peatland carbon sink in the subsiding Pastaza-Maraon foreland
770 basin, Peru. *Global Change Biology*, 18(1), 164–178. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2486.2011.02504.x)
771 [2486.2011.02504.x](https://doi.org/10.1111/j.1365-2486.2011.02504.x)
- 772 Lhteenoja, O., Ruokolainen, K., Schulman, L., & Alvarez, J. (2009). Amazonian floodplains
773 harbour minerotrophic and ombrotrophic peatlands. *CATENA*, 79(2), 140–145.
774 <https://doi.org/https://doi.org/10.1016/j.catena.2009.06.006>
- 775 LHTEENOJA, O., RUOKOLAINEN, K., SCHULMAN, L., & OINONEN, M. (2009).
776 Amazonian peatlands: an ignored C sink and potential source. *Global Change Biology*,
777 15(9), 2311–2320. <https://doi.org/10.1111/j.1365-2486.2009.01920.x>
- 778 Leifeld, J., & Menichetti, L. (2018). The underappreciated potential of peatlands in global
779 climate change mitigation strategies. *Nature Communications*, 9(1), 1071.
780 <https://doi.org/10.1038/s41467-018-03406-6>
- 781 Leng, L. Y., Ahmed, O. H., & Jalloh, M. B. (2019). Brief review on climate change and tropical
782 peatlands. *Geoscience Frontiers*, 10(2), 373–380.
783 <https://doi.org/https://doi.org/10.1016/j.gsf.2017.12.018>
- 784 Limpens, J., Berendse, F., Blodau, C., Canadell, J. G., Freeman, C., Holden, J., ... Schaepman-
785 Strub, G. (2008). Peatlands and the carbon cycle: from local processes to global
786 implications – a synthesis. *Biogeosciences*, 5(5), 1475–1491. [https://doi.org/10.5194/bg-](https://doi.org/10.5194/bg-5-1475-2008)
787 [5-1475-2008](https://doi.org/10.5194/bg-5-1475-2008)
- 788 LUND, M., LAFLEUR, P. M., ROULET, N. T., LINDROTH, A., CHRISTENSEN, T. R.,
789 AURELA, M., ... NILSSON, M. B. (2010). Variability in exchange of CO₂ across 12
790 northern peatland and tundra sites. *Global Change Biology*, 16(9), 2436–2448.
791 <https://doi.org/10.1111/j.1365-2486.2009.02104.x>
- 792 Mkiranta, P., Laiho, R., Fritze, H., Hytnen, J., Laine, J., & Minkkinen, K. (2009). Indirect

- regulation of heterotrophic peat soil respiration by water level via microbial community structure and temperature sensitivity. *Soil Biology and Biochemistry*, 41(4), 695–703. <https://doi.org/https://doi.org/10.1016/j.soilbio.2009.01.004>
- Martikainen, P. J., Nykänen, H., Alm, J., & Silvola, J. (1995). Change in fluxes of carbon dioxide, methane and nitrous oxide due to forest drainage of mire sites of different trophy. *Plant and Soil*, 168(1), 571–577. <https://doi.org/10.1007/BF00029370>
- McWethy, D. B., Neumann, F. H., Steinbruch, F., Ryan, C. M., & Valsecchi, V. (2016). Late Quaternary vegetation development and disturbance dynamics from a peatland on Mount Gorongosa, central Mozambique. *Quaternary Science Reviews*, 137, 221–233. <https://doi.org/https://doi.org/10.1016/j.quascirev.2016.02.004>
- Melling, L., Hatano, R., & Goh, K. J. (2005). Soil CO₂ flux from three ecosystems in tropical peatland of Sarawak, Malaysia. *Tellus B: Chemical and Physical Meteorology*, 57(1), 1–11. <https://doi.org/10.3402/tellusb.v57i1.16772>
- Miettinen, J., & Liew, S. C. (2010). Status of Peatland Degradation and Development in Sumatra and Kalimantan. *AMBIO: A Journal of the Human Environment*, 39(5–6), 394–401. <https://doi.org/10.1007/s13280-010-0051-2>
- Miettinen, J., Shi, C., & Liew, S. C. (2012). Two decades of destruction in Southeast Asia's peat swamp forests. *Frontiers in Ecology and the Environment*, 10(3), 124–128. <https://doi.org/10.1890/100236>
- Miettinen, J., Shi, C., & Liew, S. C. (2016). Land cover distribution in the peatlands of Peninsular Malaysia, Sumatra and Borneo in 2015 with changes since 1990. *Global Ecology and Conservation*, 6, 67–78. <https://doi.org/http://dx.doi.org/10.1016/j.gecco.2016.02.004>
- Mitsch, W. J., Nahlik, A., Wolski, P., Bernal, B., Zhang, L., & Ramberg, L. (2010). Tropical wetlands: seasonal hydrologic pulsing, carbon sequestration, and methane emissions. *Wetlands Ecology and Management*, 18(5), 573–586. <https://doi.org/10.1007/s11273-009-9164-4>
- Moore, S., Evans, C. D., Page, S. E., Garnett, M. H., Jones, T. G., Freeman, C., ... Gauci, V. (2013). Deep instability of deforested tropical peatlands revealed by fluvial organic carbon fluxes. *Nature*, 493(7434), 660–663. Retrieved from <http://dx.doi.org/10.1038/nature11818>
- Murdiyarso, D., Hergoualch, K., & Verchot, L. (2010). *Opportunities for reducing greenhouse gas emissions in tropical peatlands* (Vol. 107, p. 19655–19660). Vol. 107, p. 19655–19660. <https://doi.org/http://dx.doi.org/10.1073/pnas.0911966107>

- Neuzil, S. G. (1997). Onset and rate of peat and carbon accumulation in four domed ombrogenous peat deposit. In R. And & S. E. Page (Eds.), *Biodiversity and Sustainability of Tropical Peatlands – Proceedings of the International Symposium on Tropical Peatlands* (pp. 55–72). Palangkaraya, Indonesia; Cardigan.
- Oktarita, S., Hergoualc'h, K., Anwar, S., & Verchot, L. V. (2017). Substantial N₂O emissions from peat decomposition and N fertilization in an oil palm plantation exacerbated by hotspots. *Environmental Research Letters*, 12(10), 104007. <https://doi.org/10.1088/1748-9326/aa80f1>
- Olefeldt, D., Roulet, N. T., Bergeron, O., Crill, P., Bäckstrand, K., & Christensen, T. R. (2012). Net carbon accumulation of a high-latitude permafrost palsamire similar to permafrost-free peatlands. *Geophysical Research Letters*, 39(3). <https://doi.org/10.1029/2011GL050355>
- PAGE, S. E., RIELEY, J. O., & BANKS, C. J. (2011). Global and regional importance of the tropical peatland carbon pool. *Global Change Biology*, 17(2), 798–818. <https://doi.org/10.1111/j.1365-2486.2010.02279.x>
- Page, S E, & Baird, A. J. (2016). Peatlands and Global Change: Response and Resilience. *Annual Review of Environment and Resources*, 41(1), 35–57. <https://doi.org/10.1146/annurev-environ-110615-085520>
- Page, S E, Rieley, J. O., Shotyk, W., & Weiss, D. (1999). Interdependence of peat and vegetation in a tropical peat swamp forest. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 354(1391), 1885–1897. <https://doi.org/10.1098/rstb.1999.0529>
- Page, S E, Rieley, J. O., & Wüst, R. (2006). Chapter 7 Lowland tropical peatlands of Southeast Asia. In I. P. Martini, A. Martínez Cortizas, & W. B. T.-D. in E. S. P. Chesworth (Eds.), *Peatlands* (Vol. 9, pp. 145–172). [https://doi.org/10.1016/S0928-2025\(06\)09007-9](https://doi.org/10.1016/S0928-2025(06)09007-9)
- Page, S E, Siegert, F., Rieley, J. O., Boehm, H. D. V, Jaya, A., Limin, S., ... Limin, S. H. (n.d.). Inception, history and development of peatlands in the Amazon Basin. *Dynamics*, 19, 625–635.
- Page, S E, Wüst, R. A. J., Weiss, D., Rieley, J. O., Shotyk, W., & Limin, S. H. (2004). A record of Late Pleistocene and Holocene carbon accumulation and climate change from an equatorial peat bog (Kalimantan, Indonesia): implications for past, present and future carbon dynamics. *Journal of Quaternary Science*, 19(7), 625–635. <https://doi.org/10.1002/jqs.884>

- Page, Susan E, Siegert, F., Rieley, J. O., Boehm, H.-D. V, Jaya, A., & Limin, S. (2002). The amount of carbon released from peat and forest fires in Indonesia during 1997. *Nature*, 420(6911), 61–65. Retrieved from <http://dx.doi.org/10.1038/nature01131>
- Pajunen, H. (1996). Mires as late Quaternary accumulation basins in Rwanda and Burundi, Central Africa. In *Geological Survey of Finland* (p. 106).
- Ponette-González, A. G., Curran, L. M., Pittman, A. M., Carlson, K. M., Steele, B. G., Ratnasari, D., ... Weathers, K. C. (2016). Biomass burning drives atmospheric nutrient redistribution within forested peatlands in Borneo. *Environmental Research Letters*, 11(8), 85003. <https://doi.org/10.1088/1748-9326/11/8/085003>
- Porporato, A., Laio, F., Ridolfi, L., & Rodriguez-Iturbe, I. (2001). Plants in water-controlled ecosystems: active role in hydrologic processes and response to water stress: III. Vegetation water stress. *Advances in Water Resources*, 24(7), 725–744. [https://doi.org/https://doi.org/10.1016/S0309-1708\(01\)00006-9](https://doi.org/https://doi.org/10.1016/S0309-1708(01)00006-9)
- Qiu, C., Zhu, D., Ciais, P., Guenet, B., & Peng, S. (2020). The role of northern peatlands in the global carbon cycle for the 21st century. *Global Ecology and Biogeography*, 29(5), 956–973. <https://doi.org/10.1111/geb.13081>
- Radjagukguk, B. (1997). Peat soil of Indonesia: Location, classification and problem for sustainability. In R. J.O. & P. S.E (Eds.), *Biodiversity and Sustainability of Tropical Peatlands* (pp. 42–54). Samara Publishing Ltd.
- Raupach, M., Gloor, M., Sarmiento, J., G. Canadell, J., Frölicher, T., Gasser, T., ... Trudinger, C. (2013). The declining uptake rate of atmospheric CO₂ by land and ocean sinks. In *Biogeosciences Discussions* (Vol. 10). <https://doi.org/10.5194/bgd-10-18407-2013>
- Reay, D., Smith, P., Christensen, T. R., James, R., & Clark, H. (2018). Methane and Global Environmental Change. *Annual Review of Environment and Resources*, 43. <https://doi.org/10.1146/annurev-environ-102017-030154>
- Rebelo, L.-M., Finlayson, C. M., & Nagabhatla, N. (2009). Remote sensing and GIS for wetland inventory, mapping and change analysis. *Journal of Environmental Management*, 90(7), 2144–2153.
- Reiche, M., Gleixner, G., & Küsel, K. (2010). Effect of peat quality on microbial greenhouse gas formation in an acidic fen. *Biogeosciences*, 7(1), 187–198.
- Renou-Wilson, F., Moser, G., Fallon, D., Farrell, C. A., Müller, C., & Wilson, D. (2019). Rewetting degraded peatlands for climate and biodiversity benefits: Results from two raised bogs. *Ecological Engineering*, 127, 547–560. <https://doi.org/https://doi.org/10.1016/j.ecoleng.2018.02.014>

- Ribeiro, K., Pacheco, F. S., Ferreira, W., Sousa Neto, E. R., Ometto, J. P., Forti, M. C., ... Alvalá, P. C. (2018). Effect of land use and land cover on greenhouse gas emissions from peatlands in the Paraíba Valley, Brazil. *AGUFM*, 2018, GC21I-1213.
- Rieley, J., Wüst, R., Jauhiainen, J., Page, S., Wösten, H., Hooijer, A., ... Stahlhut, M. (2008). Tropical peatlands: Carbon stores, carbon gas emissions and contribution to climate change processes. In *Peatlands and climate change* (pp. 148–181). University of Calgary, Canada.
- Roucoux, K. H., Lawson, I. T., Baker, T. R., Del Castillo Torres, D., Draper, F. C., Lähteenoja, O., ... Vriesendorp, C. F. (2017). Threats to intact tropical peatlands and opportunities for their conservation. *Conservation Biology*, 31(6), 1283–1292. <https://doi.org/10.1111/cobi.12925>
- Roulet, N. (2012). Peatlands, carbon storage, greenhouse gases, and the Kyoto Protocol: Prospects and significance for Canada. *Wetlands*, 20, 605–615. [https://doi.org/10.1672/0277-5212\(2000\)020\[0605:PCSGGA\]2.0.CO;2](https://doi.org/10.1672/0277-5212(2000)020[0605:PCSGGA]2.0.CO;2)
- ROULET, N. T., LAFLEUR, P. M., RICHARD, P. J. H., MOORE, T. I. M. R., HUMPHREYS, E. R., & BUBIER, J. (2007). Contemporary carbon balance and late Holocene carbon accumulation in a northern peatland. *Global Change Biology*, 13(2), 397–411. <https://doi.org/10.1111/j.1365-2486.2006.01292.x>
- Rudiyanto, Setiawan, B. I., Arief, C., Saptomo, S. K., Gunawan, A., Kuswarman, ... Indriyanto, H. (2015). Estimating Distribution of Carbon Stock in Tropical Peatland Using a Combination of an Empirical Peat Depth Model and GIS. *Procedia Environmental Sciences*, 24, 152–157. <https://doi.org/10.1016/j.proenv.2015.03.020>
- Saleska, S. R., Miller, S. D., Matross, D. M., Goulden, M. L., Wofsy, S. C., da Rocha, H. R., ... Silva, H. (2003). Carbon in Amazon forests: unexpected seasonal fluxes and disturbance-induced losses. *Science (New York, N.Y.)*, 302(5650), 1554–1557. <https://doi.org/10.1126/science.1091165>
- San José, J., Montes, R., Nikonova, N., Grace, J., Buendía, C., Davidson, E. a., ... Tayasu, I. (2013). Effects of simulated drought and nitrogen fertilizer on plant productivity and nitrous oxide (N₂O) emissions of two pastures. *Biogeosciences*, 55(1–2), 291–302. <https://doi.org/10.1007/s11104-012-1248-x>
- Scharlemann, J. P. W., Tanner, E. V. J., Hiederer, R., & Kapos, V. (2014). Global soil carbon: understanding and managing the largest terrestrial carbon pool. *Carbon Management*, 5(1), 81–91. <https://doi.org/10.4155/cmt.13.77>
- Schumann, M., & Joosten, H. (2008). *Global peatland restoration manual*. Retrieved from

- 929 http://www.imcg.net/media/download_gallery/books/gprm_01.pdf
- 930 Silvola, J., Alm, J., Ahlholm, U., Nykanen, H., & Martikainen, P. J. (1996). CO_2
- 931 Fluxes from Peat in Boreal Mires under Varying Temperature and Moisture Conditions.
- 932 *Journal of Ecology*, 84(2), 219–228. <https://doi.org/10.2307/2261357>
- 933 Sjögersten, S., Black, C., Evers, S., Hoyos-Santillan, J., Wright, E., & Turner, B. (2014).
- 934 Tropical wetlands: A missing link in the global carbon cycle? *Global Biogeochemical*
- 935 *Cycles*. <https://doi.org/10.1002/2014GB004844>
- 936 Sorensen, K. W. (1993). Indonesian peat swamp forests and their role as a carbon sink.
- 937 *Chemosphere*, 27(6), 1065–1082.
- 938 Sorribas, M. V., Paiva, R. C. D., Melack, J. M., Bravo, J. M., Jones, C., Carvalho, L., ... Costa,
- 939 M. H. (2016). Projections of climate change effects on discharge and inundation in the
- 940 Amazon basin. *Climatic Change*, 136(3), 555–570. <https://doi.org/10.1007/s10584-016->
- 941 1640-2
- 942 Spahni, R., Joos, F., Stocker, B. D., Steinacher, M., & Yu, Z. C. (2013). Transient simulations
- 943 of the carbon and nitrogen dynamics in northern peatlands: from the Last Glacial
- 944 Maximum to the 21st century. *Clim. Past*, 9(3), 1287–1308. <https://doi.org/10.5194/cp-9->
- 945 1287-2013
- 946 Swindles, G. T., Morris, P. J., Whitney, B., Galloway, J. M., Gałka, M., Gallego-Sala, A., ...
- 947 Lähteenoja, O. (2018). Ecosystem state shifts during long-term development of an
- 948 Amazonian peatland. *Global Change Biology*, 24(2), 738–757.
- 949 <https://doi.org/10.1111/gcb.13950>
- 950 Takada, M., Shimada, S., & Takahashi, H. (2016). *Tropical Peat Formation BT - Tropical*
- 951 *Peatland Ecosystems* (M. Osaki & N. Tsuji, Eds.). <https://doi.org/10.1007/978-4-431->
- 952 55681-7_8
- 953 Tian, H., Chen, G., Lu, C., Xu, X., Ren, W., Zhang, B., ... Wofsy, S. (2015). Global methane
- 954 and nitrous oxide emissions from terrestrial ecosystems due to multiple environmental
- 955 changes. *Ecosystem Health and Sustainability*, 1(1), 1–20.
- 956 <https://doi.org/10.1890/EHS14-0015.1>
- 957 Treat, C. C., Kleinen, T., Broothaerts, N., Dalton, A. S., Dommain, R., Douglas, T. A., ...
- 958 Brovkin, V. (2019). Widespread global peatland establishment and persistence over the
- 959 last 130,000 y. *Proceedings of the National Academy of Sciences*, 116(11), 4822 LP –
- 960 4827. <https://doi.org/10.1073/pnas.1813305116>
- 961 Ueda, S., Go, C.-S. U., Yoshioka, T., Yoshida* ast;, N., Wada, E., Miyajima* ast ast;, T., ...
- 962 Boonprakub, S. (2000). Dynamics of dissolved O₂, CO₂, CH₄, and N₂O in a tropical

- coastal swamp in southern Thailand. *Biogeochemistry*, 49(3), 191–215.
<https://doi.org/10.1023/A:1006393214432>
- Vitt, D. H. (2013). *Peatlands* ☆ (B. B. T.-E. of E. (Second E. Fath, Ed.).
<https://doi.org/https://doi.org/10.1016/B978-0-12-409548-9.00741-7>
- Wang, H., Richardson, C. J., & Ho, M. (2015). Dual controls on carbon loss during drought
 in peatlands. *Nature Climate Change*, 5, 584.
- Wang, S., Zhuang, Q., Lähteenoja, O., Draper, F. C., & Cadillo-Quiroz, H. (2018). Potential
 shift from a carbon sink to a source in Amazonian peatlands under a changing climate.
Proceedings of the National Academy of Sciences, 115(49), 12407 LP – 12412.
<https://doi.org/10.1073/pnas.1801317115>
- Warner, B. G., & Rubec, C. D. A. (1997). *The Canadian Wetland Classification System* (2nd
 ed.). National Wetlands Working Group.
- Warren, M., Hergoualc'h, K., Kauffman, J. B., Murdiyarso, D., & Kolka, R. (2017). An
 appraisal of Indonesia's immense peat carbon stock using national peatland maps:
 uncertainties and potential losses from conversion. *Carbon Balance and Management*,
 12(1), 12. <https://doi.org/10.1186/s13021-017-0080-2>
- Wilson, S., Griffiths, M., & Anielski, M. (2001). *The Alberta GPI Accounts: Wetlands and*
Peatlands. Retrieved from
http://www.pembina.org/reports/23_wetlands_and_peatlands.pdf
- Wong, G. X., Hirata, R., Hirano, T., Kiew, F., Aeries, E. B., Musin, K. K., ... Melling, L.
 (2020). How do land use practices affect methane emissions from tropical peat
 ecosystems? *Agricultural and Forest Meteorology*, 282–283, 107869.
<https://doi.org/https://doi.org/10.1016/j.agrformet.2019.107869>
- Worrall, F., Chapman, P., Holden, J., Evans, C., Artz, R., Smith, P., & Grayson, R. (2011). *A*
review of current evidence on carbon fluxes and greenhouse gas emissions from UK
peatlands. <https://doi.org/09638901>
- Wösten, J. H. M., Clymans, E., Page, S. E., Rieley, J. O., & Limin, S. H. (2008). Peat–water
 interrelationships in a tropical peatland ecosystem in Southeast Asia. *CATENA*, 73(2),
 212–224. <https://doi.org/https://doi.org/10.1016/j.catena.2007.07.010>
- WRIGHT, E. L., BLACK, C. R., CHEESMAN, A. W., DRAGE, T., LARGE, D., TURNER,
 B. L., & SJÖGERSTEN, S. (2011). Contribution of subsurface peat to CO₂ and CH₄
 fluxes in a neotropical peatland. *Global Change Biology*, 17(9), 2867–2881.
<https://doi.org/10.1111/j.1365-2486.2011.02448.x>
- Wu, J., & Roulet, N. T. (2014). Climate change reduces the capacity of northern peatlands to

- absorb the atmospheric carbon dioxide: The different responses of bogs and fens. *Global Biogeochemical Cycles*, 28(10), 1005–1024. <https://doi.org/10.1002/2014GB004845>
- Xu, J., Morris, P. J., Liu, J., & Holden, J. (2018). PEATMAP: Refining estimates of global peatland distribution based on a meta-analysis. *CATENA*, 160, 134–140. <https://doi.org/https://doi.org/10.1016/j.catena.2017.09.010>
- Yu, G. R., Fang, H. J., Cheng, S. L., Zhu, T. H., Wang, Y. S., Yan, J. H., ... Zhou, M. (2010). Effects of multiple environmental factors on CO₂ emission and CH₄ uptake from old-growth forest soils. *Biogeosciences*, 7(1), 395–407.
- Yu, Z. (2011). Holocene carbon flux histories of the world's peatlands: Global carbon-cycle implications. *The Holocene*, 21(5), 761–774. <https://doi.org/10.1177/0959683610386982>
- Yu, Z. C. (2012). Northern peatland carbon stocks and dynamics: a review. *Biogeosciences*, 9, 4071–4085. <https://doi.org/10.5194/bg-9-4071-2012>
- Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W., & Hunt, S. J. (2010). Global peatland dynamics since the Last Glacial Maximum. *Geophysical Research Letters*, 37(13). <https://doi.org/10.1029/2010GL043584>
- Yule, C. M., Lim, Y. Y., & Lim, T. Y. (2016). Degradation of Tropical Malaysian Peatlands Decreases Levels of Phenolics in Soil and in Leaves of *Macaranga pruinosa*. *Frontiers in Earth Science*, Vol. 4, p. 45. Retrieved from <https://www.frontiersin.org/article/10.3389/feart.2016.00045>
- Zoltai, S. C., & Martikainen, P. J. (1996). *Estimated extent of forested peatlands and their role in the global carbon cycle BT - Forest Ecosystems, Forest Management and the Global Carbon Cycle* (M. J. Apps & D. T. Price, Eds.). https://doi.org/10.1007/978-3-642-61111-7_5
- Zulkafli, Z., Buytaert, W., Manz, B., Rosas, C. V., Willems, P., Lavado-Casimiro, W., ... Santini, W. (2016). Projected increases in the annual flood pulse of the Western Amazon. *Environmental Research Letters*, 11(1), 14013. <https://doi.org/10.1088/1748-9326/11/1/014013>

Figure 1 – Estimated peat area (a), depth (b) and volume (c), presented by Page et al. (2011), Gumbricht et al. (2017) and Xu et al. (2018) of tropical peatlands. (d) Estimated carbon stock (GtC) in tropical peatlands. Error bars are minimum and maximum estimates when available. *Values estimated using peatland area from Xu et al. (2018) and mean depth from Gumbricht et al. (2017) and Page et al. (2011).

136x241mm (300 x 300 DPI)

Figure 2 – Distribution of peatlands in tropical regions. Data from Xu et al., (2018)

307x208mm (300 x 300 DPI)