

THE UNIVERSITY of EDINBURGH

Edinburgh Research Explorer

Culture and characterisation of canine mitral valve interstitial and endothelial cells

Citation for published version:

Liu, MM, Flanagan, TC, Lu, CC, French, AT, Argyle, DJ & Corcoran, BM 2015, 'Culture and characterisation of canine mitral valve interstitial and endothelial cells', *Veterinary Journal*, vol. 204, no. 1, pp. 32-39. https://doi.org/10.1016/j.tvjl.2015.01.011

Digital Object Identifier (DOI):

10.1016/j.tvjl.2015.01.011

Link:

Link to publication record in Edinburgh Research Explorer

Document Version: Peer reviewed version

Published In: Veterinary Journal

Publisher Rights Statement:

This is the author's final peer-reviewed manuscript as accepted for publication. The final published version is available here: http://www.sciencedirect.com/science/article/pii/S1090023315000295

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Elsevier Editorial System(tm) for The Veterinary Journal Manuscript Draft

Manuscript Number: YTVJL-D-14-00751R2

Title: Culture and characterisation of canine mitral valve interstitial and endothelial cells

Article Type: Original Article

Keywords: Valve interstitial cell; Valve endothelial cell; Cell culture; Canine; Myxomatous mitral valve disease.

Corresponding Author: Prof. Brendan Corcoran, MVB PhD

Corresponding Author's Institution: University of Edinburgh

First Author: Brendan Corcoran, MVB PhD

Order of Authors: Brendan Corcoran, MVB PhD; Meng Meng Liu; Tom Flangan; Chi-Chien Lu; Anne French; David Argyle

Abstract: Valve interstitial cells (VICs) have an important role in the aetiopathogenesis of myxomatous mitral valve disease (MMVD) in the dog. Furthermore, there is evidence that valve endothelial cells (VECs) also contribute to disease development. In addition to examining native valve tissue to understand MMVD, another strategy is to separately examine VIC and VEC biology under in vitro culture conditions. The aim of this study was to isolate and characterise canine mitral VICs and VECs from normal dog valves using a combination of morphology, immunohistochemistry and reverse transcription PCR (RT-PCR).

Canine mitral VECs and VICs were isolated and cultured in vitro. The two cell populations exhibited different morphologies and growth patterns. VECs, but not VICs, expressed the endothelial markers, platelet endothelial cell adhesion molecule (PECAM-1 or CD31) and acetylated low density lipoprotein (Dil-Ac-LDL). Both VECs and VICs expressed vimentin and embryonic non-smooth muscle myosin heavy chain (SMemb), an activated mesenchymal cell marker. The myofibroblast marker, alpha smooth muscle actin (α -SMA), was detected at the mRNA level in both VEC and VIC cultures, but only at the protein level in VIC cultures. The morphological heterogeneity and expression of non-endothelial phenotypic markers in VEC cultures suggested that a mixture of cell types was present, which might be due to cell contamination and/or endothelial mesenchymal transition (EndoMT). The use of a specific endothelial culture medium for primary VEC cultures enhanced the endothelial properties of the cells and reduced α -SMA and SMemb expression.

1	Culture and characterisation of canine mitral valve interstitial and endothelial cells		
2			
3	M-M. Liu ^a , T.C. Flanagan ^b , C-C. Lu ^a , A. T. French ^c , D.J. Argyle ^a , B.M. Corcoran ^{a,*}		
4			
5	^a Royal (Dick) School of Veterinary Studies and The Roslin Institute, the University of		
6	Edinburgh, Easterbush, Roslin, Mid-Lothian, Scotland, UK EH25 9RG.		
7	^b School of Medicine & Medical Science, Health Sciences Centre, University College Dublin,		
8	Belfield, Dublin 4, Ireland.		
9	^c University of Glasgow School of Veterinary Medicine, Bearsden Road, Glasgow, G61 1QH,		
10	Scotland, UK		
11			
12	*Corresponding Author: Tel: +44 (0)131 651 9100		
13	E-mail address: Brendan.Corcoran@ed.ac.uk (Brendan Corcoran)		
14			
15			
16			

17 Abstract

18 Valve interstitial cells (VICs) have an important role in the aetiopathogenesis of 19 myxomatous mitral valve disease (MMVD) in the dog. Furthermore, there is evidence that 20 valve endothelial cells (VECs) also contribute to disease development. In addition to 21 examining native valve tissue to understand MMVD, another strategy is to separately 22 examine VIC and VEC biology under in vitro culture conditions. The aim of this study was to isolate and characterise canine mitral VICs and VECs from normal dog valves using a 23 24 combination of morphology, immunohistochemistry and reverse transcription PCR (RT-25 PCR).

26

Canine mitral VECs and VICs were isolated and cultured in vitro. The two cell 27 populations exhibited different morphologies and growth patterns. VECs, but not VICs, 28 29 expressed the endothelial markers, platelet endothelial cell adhesion molecule (PECAM-1 or 30 CD31) and acetylated low density lipoprotein (Dil-Ac-LDL). Both VECs and VICs expressed vimentin and embryonic non-smooth muscle myosin heavy chain (SMemb), an activated 31 32 mesenchymal cell marker. The myofibroblast marker, alpha smooth muscle actin (α -SMA), 33 was detected at the mRNA level in both VEC and VIC cultures, but only at the protein level 34 in VIC cultures. The morphological heterogeneity and expression of non-endothelial phenotypic markers in VEC cultures suggested that a mixture of cell types was present, 35 36 which might be due to cell contamination and/or endothelial mesenchymal transition 37 (EndoMT). The use of a specific endothelial culture medium for primary VEC cultures 38 enhanced the endothelial properties of the cells and reduced α -SMA and SMemb expression.

39

Keywords: Valve interstitial cell; Valve endothelial cell; Cell culture; Canine; Myxomatous
mitral valve disease.

43 Introduction

44 Myxomatous mitral valve disease (MMVD) is characterised by extensive changes to 45 the valve extra cellular matrix (ECM), resulting in valve distortion and mechanical instability, 46 and leading to valve insufficiency and mitral regurgitation (Black et al., 2005; Hadian et al., 47 2007; Aupperle et al., 2009, 2010; Han et al., 2010, 2013b). The mechanisms that lead to 48 valve degeneration are not fully understood, but are presumed to involve changes in valve interstitial cell (VIC) and endothelial cell (VEC) phenotype and function (Corcoran et al., 49 50 2004; Disatian et al., 2008; Han et al., 2008, 2013a). Endothelial changes are characterised by 51 detachment, denuding and apoptosis with splitting of the basement membrane and increased 52 expression of basement membrane proteins, including laminin (Aupperle et al., 2009; Han et 53 al., 2013a). VICs show an activated myofibroblast phenotype and their numbers decline in 54 the overly myxomatous areas and proliferate close to the valve surface, most noticeably 55 where there has been endothelial damage (Disatian et al., 2008; Han et al., 2008, 2013a).

56

57 VICs are the cells contributing to ECM production and remodelling. Their synthetic 58 activity is driven mainly by the transforming growth factor beta (TGF- β) cytokine 59 superfamily, but is also thought to be modified by signals from the endothelial cells (Olsen et 60 al., 2003; Disatian et al., 2010; Aupperle and Disatian, 2012; Orton et al., 2012; Han et al., 61 2013a). This VEC-VIC interaction has led to the hypothesis that VEC activation is 62 fundamental to valve matrix homeostasis and that MMVD may be driven by mechanisms 63 involving VEC damage and leading to aberrant matrix remodelling.

64

Future examination of MMVD will require a greater understanding of VEC-VIC interactions and one approach is to do so using in vitro models. Canine mitral VICs have been previously isolated and cultured using a primary explant technique which favours 68 culturing migratory cells only, but there was limited phenotypic characterisation, and there 69 are no reports of successful cultures of canine VECs (Heaney et al., 2009). The aims of this 70 study, therefore, were (1) to develop robust methods for the culture and phenotypic 71 characterisation of canine mitral VICs and VECs based on a reported protocol for porcine 72 aortic valve cell isolation (Gould and Butcher, 2010), and (2) to examine canine mitral valve 73 cell biology under the culture conditions. Furthermore, as it is recognised to be difficult to maintain endothelial cell phenotype in VEC culture, different culture protocols were 74 75 evaluated.

76

77 Materials and methods

78 *Cell isolation and culture*

Cells were isolated from 12 healthy canine mitral valves showing no MMVD evidence on gross morphological inspection. The dogs were of different breeds, aged 2-5 years and with equal sex distribution. All dogs were euthanased for reasons other than cardiac disease. Samples were collected with full owner consent and sampling conformed to institutional and national ethical guidelines.

84

85 Mitral valves were removed within minutes of euthanasia under sterile conditions, 86 rinsed in cold sterile phosphate buffered saline (PBS) and placed on ice until transferred to a 87 tissue culture laminar hood. Excess annular tissue and chordae tendineae were removed, and 88 samples were placed in 35 mm petri dishes and incubated with 5 mL pre-warmed collagenase solution (600 Units/ml) in 5% CO₂ for 10 min at 37 °C. VECs were removed by gently 89 90 rotating a dry sterile swab over the surface of the leaflet and occasionally dabbing the swab 91 into the collagenase solution. The collagenase solution was aspirated and the suspended cells 92 pelleted by centrifuging at 1000g for 5 min at room temperature. The supernatant was

93 aspirated and the cells were re-suspended in advanced DMEM/F-12 medium (Life 94 Technologies) with 10% foetal bovine serum (FBS), 1% penicillin/streptomycin and 1% L-95 glutamine (AF-12). Cells were centrifuged, re-suspended again and seeded on culture flasks 96 or plates pre-coated with 2% gelatin and incubated at 37 °C in 5% CO₂. After VEC 97 collection, tissue was placed in a 15 mL conical tube with 10 mL collagenase solution (600 98 U/mL) and incubated for 18 h at 37 °C in 5% CO₂. After digestion, the tissue was 99 homogenized in the collagenase solution. The released cells were then seeded onto flasks and 100 cultured as described for VECs in AF-12 medium.

101

Media changes occurred at 2 day intervals for both populations. At confluence, cells were passaged by either trypsinization (0.05% - 1% Trypsin /EDTA) or by using 1 x TrypLE Express (Life Technologies). In general, VECs were expanded up to passage three and VICs were cultured up to passage 8.

106

107 Comparison of two culture media for canine mitral VECs

108 To evaluate the effect of specific endothelial culture medium on primary VEC 109 properties, a canine endothelial basal medium kit (Cn 211k-500, Cell Application; EBM) was 110 compared with the AF-12 medium for VECs. VEC preparations (at passages 1-2) from 111 different dogs (n = 3) were used for the medium comparison experiment. Prior to the medium 112 switch, cells were stored in freezing medium (70% AF-12 medium/20% FBS/10% 113 dimethylsulphoxide (DMSO; Invitrogen)) and cryopreserved in -150 °C freezer or in liquid 114 nitrogen freezer. Cells were revived in a 37 °C water bath, and were washed with pre-warmed 115 AF-12 medium by centrifuging. After removal of the supernatant, the cells were re-suspended 116 in canine EBM and seeded onto culture plates pre-coated with 2% gelatin and incubated at 117 37 °C in 5% CO₂. Control cultures were maintained in AF-12 for the same VEC preparation.

118 VEC cultures in EBM and AF-12 were harvested at confluence and evaluated for endothelial119 and mesenchymal marker expression.

120

121 *Reverse transcription-polymerase chain reaction (RT-PCR)*

Total RNA was isolated from both cell types using the RNeasy Mini Kit (Qiagen), according to the manufacturer's protocol. Extracted RNA was treated with RNase-free DNase I and concentration of RNA was measured using NanoDrop (Thermal Scientific). RNA was diluted using nuclease-free water (Qiagen) and was denatured at 65 $^{\circ}$ C for 5 min, then placed immediately on ice. Complementary DNA was synthesized using an Omniscript Reverse Transcription Kit (Qiagen) at 40 $^{\circ}$ C for 1 h.

128

129 The following genes (Table 1) were selected to characterise the VECs and VICs: 130 CD31 and von Willebrand Factor (vWF) (endothelial markers); alpha-smooth muscle actin 131 (a-SMA), embryonic form of non-smooth muscle myosin (SMemb) and transgelin (SM22) 132 (activated VIC (aVIC) markers); *vimentin* (mesenchymal origin cells). GAPDH was used as a 133 housekeeping gene. The primers of vWF-1, α -SMA and GAPDH were kind gifts (Hannah 134 Hodgkiss-Geere, the University of Edinburgh), and vWF3 primer sequence was derived from 135 Fulton et al. (2000). All other primers were originally designed using Primer 3 Input v.4.0 136 and nucleotide sequences of the above markers were obtained from the NCBI Gene Bank or 137 Ensembl databases.

138

PCR amplification used the Gotaq PCR Core System (Promega). Initial denaturation
at 95 °C for 5 min was followed by 30 cycles of 95 °C for 1 min, 58-61 °C for 1 min, 72 °C
for 1 min and a final extension at 72 °C for 10 min. PCR products were analysed using a

standard 2% agarose gel electrophoresis. Molecular Imager Gel Doc system (Bio-Rad) was
used to visualize the results after electrophoresis.

144

145 *Immunocytochemistry*

146 The cell suspension (200 μ L) was added to duplicate BD Falcon Culture slides, precoated with 2% gelatin for VECs, at a density of 1 x 10⁴ cell/well and incubated in 5% CO₂ at 147 148 37 °C for 24-48 h. Slides were directly fixed with acetone for 10 min at -20 °C, washed twice 149 in PBS, incubated for 1 h in blocking buffer (10% goat serum/0.1% Tween 20/PBS) and then 150 overnight in a humidity chamber at 4 °C with primary antibodies against CD31, vimentin, 151 SMemb and α -SMA (Table 2). Slides were washed twice with PBS and incubated with 100 152 µL goat anti-rabbit IgG or goat anti-mouse IgG secondary antibody for 1 h at room 153 temperature in a dark humid chamber. The slides were then washed three times in PBS, 4',6-154 diamidino-2-phenylindole (DAPI) mounted (Vectashield Mounting Medium, Vector Inc), and 155 examined using a Leitz fluorescence microscope (Leica).

156

157 DiI- acetylated-low density lipoprotein (DiI-Ac-LDL) labelling

Acetylated low density lipoprotein conjugated with fluorescence dye 1, 1'dioctadecyl-3, 3, 3', 3'-tetramethyl-indocarbocyanine perchlorate (DiI-Ac-LDL) was used to detect endothelial cells. The cell culture protocol prior to labelling was the same as that described for immunocytochemistry. DiI-Ac-LDL (200 μ L; 5 μ g/mL; BT902, Biomedical Technologies) reagent was added to one cell chamber and incubated for 4 h in 5% CO₂ at 37 °C. Slides were washed three times with PBS and were examined by fluorescence and bright field microscopy.

165

166 *Image capture and processing*

167 Cell morphology was observed by light microscopy during the culture period and 168 assessed for cell morphology and confluence. Representative images were captured using a 169 camera connected Zeiss Axiover 40 microscope. Immunofluorescence and DiI-Ac-LDL 170 fluorescent channel images were captured using a Leica Firecam (Leica). For optimal 171 visualization of the fluorescence signals, the antibody and DAPI staining images were 172 converted into grayscale with Image J software. Original antibody and DAPI channel colour 173 images were merged using Adobe Photoshop CS6 software (Adobe System). For DiI-Ac-174 LDL labelling, bright field images were taken from the same sampled area demonstrating cell 175 distribution and were converted into grayscale images with ImageJ software.

176

177 **Results**

178 *Cell culture morphology and growth*

By using current collagenase digestion techniques, VEC and VIC were successfully isolated from the same mitral valve (n = 12) and grew well in all sub-cultures. The two culture types showed differences in cell morphology and growth pattern.

182

Endothelial colony-forming cell clusters were randomly distributed in VEC culture flasks 1-2 days post primary isolation, and clustering promoted outgrowth. Cells had cobblestone morphology typical of endothelial cells in culture, and most noticeably at confluence (Fig.1). Confluence of passage 0 VECs was usually reached within 6-8 days in one T80 flask when most cells exhibited growth contact inhibition and formed a monolayer. Some cell polymorphism was also noted with spindle shapes and multiple cytoplasmic extensions, which increased in numbers with increasing passages.

191 Cultured VICs typically had a spindle shape reminiscent of a fibroblast-like 192 morphology, although heterogeneous morphologies were also observed, and the cells 193 proliferated readily in culture (Fig.1). At confluence, the cells tended to form a whorl-like 194 pattern with multiple layers and did not exhibit growth contact inhibition (Fig. 1). Primary 195 passage 0 VICs usually reached at confluence in a T80 flask within 3 days.

196

197 *VEC phenotypic characterization*

Phenotypic characterization was performed on 3-4 VEC cell lines (at passages 2-3) derived from different dogs. By RT-PCR, VECs expressed CD31, vimentin, as well as α -SMA, SMemb and SM22 (Fig. 2). Expression of vWF was not apparent in canine mitral VECs, and this finding was consistently observed with three different vWF primer pairs. All vWF primer pairs were validated by probing cDNA of two endothelial nature cell lines (one human umbilical vascular endothelial cell line and one canine haemangiosarcoma cell line), in which the expression was clearly identified (data not show).

205

By immunofluorescence, CD31 was identified in VECs, particularly at cell-cell junctions (Fig. 3). VECs also extensively expressed vimentin and SMemb in the cytoplasm (Fig. 3), while α -SMA expression was limited and only detected in a few cells from one culture. The endothelial identity of the VEC culture was also confirmed by DiI-Ac-LDL uptake (Fig. 3).

211

212 Effects of culture media on VEC culture characteristics

Replacing standard culture medium (AF-12) with EBM helped to maintain a more endothelial phenotype as determined by morphological assessment. Cells in both media showed Dil-Ac-LDL up-take and CD31 expression, but staining was more intense in cells

216	cultured with EBM (Fig. 4). At the transcriptional level, higher CD31 expression was
217	observed in EBM cultures compared with AF-12 cultures, while α -SMA and SMemb
218	expression was reduced in the EBM cultures compared to AF-12 cultures (Fig. 5).

219 *VIC phenotypic characterization*

220 Phenotypic characterization was performed on 3-5 VIC cell lines (at passages 4-7) 221 derived from different dogs. By RT-PCR, a-SMA, vimentin, SM22 and SMemb were expressed in VIC cultures in contrast to CD31 (Fig. 2). By immunofluorescence, VIC 222 223 cultures were also consistently negative for CD31 expression (Fig. 6). Similar to VECs, VICs 224 showed extensive cytoplasmic expression of vimentin and SMemb. α -SMA expression was 225 observed in the cytoplasm, although the expression level varied between different cultures, 226 with an estimated average of 10% cell expression. No VIC cultures showed DiI-Ac-LDL 227 uptake (Fig. 6).

228

229 Discussion

230 VICs have been previously cultured from dog mitral valves from normal dogs and in one study from dogs with mild MMVD (Heaney et al., 2009; Waxman et al., 2012). In the 231 232 study by Heaney et al. (2009), cultures were initiated up to 36 h after death (Heaney et al., 233 2009). Furthermore, their protocol favoured migratory cells, resulting in suboptimal VIC 234 isolation and culture (Taylor et al., 2000). Waxman et al. (2012) isolated mitral VICs using 235 the same collagenase digestion technique as in our study, but from a limited number of 236 laboratory Beagles that were healthy or with mild MMVD. VIC characteristics in a standard 237 tissue culture system, however, were not evaluated. To date there are no reports of culture 238 and characterisation of VECs collected from canine valves immediately after death.

In the current study, canine mitral VECs and VICs were successfully isolated from the same subjects, cultured and characterized. Relatively pure cell populations were obtained and there was differential expression of a range of known mitral valve cell phenotypic markers. This provides fundamental information on canine mitral valve cell biology, which will benefit mitral valve disease research.

245

246 VECs are known to express CD31 and to show selective uptake of DiI-Ac-LDL 247 through endocytosis. These properties allow confirmation of an endothelial origin (Cuy et al., 2003; Gould and Butcher, 2010). The lack of expression of vWF was unexpected, but could 248 249 be due to a tissue-specific heterogeneous expression of vWF, as has been previously observed 250 in endothelium from different anatomic sites (Pusztaszeri et al., 2006). A previous report on 251 porcine mitral VECs also found vWF expression to be weak at the transcriptional level 252 (Flanagan et al., 2006). The majority of VECs exhibited typical cobblestone morphology, but 253 cellular pleomorphism was also present with spindle-shaped cells reminiscent of interstitial 254 cells and fibroblasts. This may be explained by valve-side heterogeneity (as found in aortic 255 valve VECs), VIC contamination or endothelial mesenchymal transition (EndoMT) to a VIC 256 phenotype (Simmons et al., 2005; Bischoff and Aikawa, 2011). In the current study, the 257 mRNA expression of aVIC markers (i.e. SMemb, SM22 and a-SMA) in VEC cultures 258 suggests the presence of VIC contamination and/or EndoMT in the culture system. VIC 259 contamination is a common issue in primary VEC cultures when using enzymatic digestion 260 (Butcher and Nerem, 2007; Cheung et al., 2008). However, cell contamination cannot fully 261 explain the aVIC characteristics in the VEC cultures, particularly the extensive protein 262 expression of the activated mesenchymal marker SMemb in VEC. To our knowledge, this is 263 the first study showing SMemb expression in adult VECs, which suggests that VECs in the current culture system are in an activated state and have embryonic cell potential reminiscentof EndoMT.

266

267 EndoMT does involve increased mesenchymal and decreased endothelial marker 268 expression (Hoerstrup et al., 1998; Cheung et al., 2008; Bischoff and Aikawa, 2011). Such 269 phenotypic transformation of VECs has been suggested to be similar to what occurs during 270 valvular development (Person et al., 2005; Butcher and Markwald, 2007; Butcher and Nerem, 271 2007; Bischoff and Aikawa, 2011). The complexity of the transition from cardiac 272 endothelium to interstitial cells involves multiple signalling pathways, but particularly 273 endothelial growth factors and member of the TGF- β superfamily (Armstrong and Bischoff, 274 2004). The possible EndoMT potential in adult valve cells has been examined previously 275 (Paruchuri et al., 2006; Paranya et al., 2001; Mahler et al., 2013). It has been hypothesized to 276 be one of the key remodelling mechanisms in heart valve diseases, where a subset of adult 277 VECs possess foetal cell characteristics similar to progenitor cells, and can replenish the VIC 278 population when needed, for example during disease or remodelling process (Paruchuri et al., 279 2006; Liu et al., 2007; Bischoff and Aikawa, 2011).

280

281 Due to cell contamination and/or in vitro differentiation, the primary VECs were less 282 stable particularly in long term cultures. Cell purification techniques and special culture 283 medium favouring endothelial growth are potential solutions to increase VEC purity 284 (Hoerstrup et al., 1998; Cheung et al., 2008; Gould and Butcher, 2010). In the current study, 285 these two methods were evaluated separately. Trial cell sorting experiments were performed 286 on the VECs from one canine mitral valve (data not shown). By using CD146 and flow 287 cytometry (Wills et al., 2009), over 75% of positive cells were detected in the passage 1 VEC 288 culture. The separated CD146-positive cells showed stronger endothelial and decreased 289 mesenchymal properties compared to the unsorted cells in short term cultures. However, 290 myofibroblast-like morphology re-appeared later when cells reached confluence. This 291 suggests that even though cell sorting can efficiently separate VECs from a mixed population 292 culture, a standard cell culture medium (i.e. AF-12) cannot maintain the endothelial 293 properties in long term cultures. To preserve endothelial features and prevent mesenchymal 294 differentiation, VECs typically require a specific culture medium containing growth factors 295 favouring endothelial growth (Paranya et al., 2001; Wylie-Sears et al., 2011; Ci et al., 2013). 296 The use of EBM instead of AF-12 resulted in a more consistent VEC phenotype, as 297 evidenced by enhanced endothelial marker expression and decreased α -SMA and SMemb 298 expression. However, EBM did not inhibit the proliferation of contaminating cell types. A 299 combination of cell sorting and of a specific endothelial medium culture could improve the 300 purity of VECs as well as maintain them for longer culture, however, further investigations 301 are required to demonstrate it. For unsorted VECs, early passage (less than passage 3) and 302 less confluent cells should be used for studies of VECs to guarantee a reasonable culture 303 purity (Gould and Butcher, 2010).

304

305 Primary VIC cultures exhibited a fibroblast-like morphology, which is consistent with 306 previous reports (Heaney et al., 2009; Waxman et al., 2012). However, some morphological 307 variation was observed and a different response to enzymatic dissociation was observed with 308 some cells showing stronger adhesion than others. This adherence difference has been 309 associated with differential expression of α -SMA, with near 100% expression in more 310 adhesive cells (Blevins et al., 2006). The heterogeneity in native VICs has been reviewed by 311 Liu et al. (2007), who suggested that there are at least five sub-phenotypes. These sub-312 populations respond to stimulation and environmental changes differently and have different 313 functions, but currently there is no definitive marker panel to identify each VIC subphenotype. In general, VICs do not express VEC markers and are vimentin positive (Liu et al., 2007; Heaney et al., 2009). VICs are further confirmed by the expression of SMemb, SM22 and α -SMA, which are thought to indicate VIC activation (aVICs) (Della Rocca et al., 2000; Rabkin et al., 2001; Wiester and Giachelli, 2003; Disatian et al., 2008; Han et al., 2008; Stephens et al., 2011).

319

320 Recently aVICs have attracted some attention in the valvular research field, as they 321 were found to play a key role in valve remodelling or stromal degeneration (Rabkin et al., 322 2001; Disatian et al., 2008; Gupta et al., 2009; Han et al., 2008). In diseased valves, there is 323 an increase of α -SMA positive cells close to the endothelium, and these aVICs are presumed 324 to contribute to valve damage and attempts at repair (Disatian et al., 2008; Han et al., 2008). 325 Moreover, in vitro studies suggest that the aVIC population is dynamic, mechanosensitive 326 and responsive to external stimulation. For example, culture conditions, such as cell density, 327 substrate types and movement, can affect the level of α -SMA expression (Engler et al., 2006; 328 Stephens et al., 2011; Xu et al., 2012). These findings may explain the discrepancy of α -SMA 329 expression observed between the current study and a previous report (Taylor et al., 2000). 330 Under culture conditions using three dimensional collagen substrates, the appearance of α -331 SMA positive cells suggests that a cell population analogous to the disease phenotype is 332 present, but this can be reversed by mechanical treatment of the cells, reverting them to a 333 more normal fibroblast phenotype (Waxman et al., 2012). Functional studies of mitral VICs 334 are required. Cultured mitral VICs are capable of ECM synthesis and are involved in wound 335 repair (Lester et al., 1993; Durbin et al., 2005; Flanagan et al., 2006; Fayet et al., 2007; Gupta 336 et al., 2008). The effects of various conditions on canine valve cell phenotype and function 337 requires further investigation if cultured VECs and VICs are to be used for modelling mitral 338 valve disease.

339 Conclusion

The characterisation of canine mitral VECs and VICs demonstrated phenotypic
heterogeneity under the static culture conditions used in the current study. These findings
should contribute to improving the understanding of canine mitral valve cellular biology, and
should benefit mitral valve disease research and tissue engineering studies.
Acknowledgements
Mengmeng Liu is the recipient of a China Scholarship Council PhD scholarship. Dr
Hannah Hodgkiss-Geere kindly provided the primers for vWF-1, α-SMA and GAPDH.

348

349 **Conflict of interest statement**

350 None of the authors of this paper has a financial or personal relationship with other

351 people or organizations that could inappropriately influence or bias the content of the paper.

352

353 **References**

354 355

Armstrong, E.J., Bischoff, J., 2004. Heart valve development: endothelial cell signaling and differentiation. Circulation Research 95, 459-470.

358

Aupperle, H., Disatian, S., 2012. Pathology, protein expression and signaling in myxomatous
mitral valve degeneration: comparison of dogs and humans. Journal of Veterinary Cardiology
14, 59-71.

Aupperle, H., Marz, I., Thielebein, J., Dinges, G., Schoon, H.A., 2010. Histomorphological
findings and expression of matrix metalloproteinases and their tissue specific inhibitors
(TIMPs) in normal tricuspid valves and in chronic tricuspid valvular disease in dogs. The
Veterinary Journal 183, 176-183.

Aupperle, H., Marz, I., Thielebein, J., Kiefer, B., Kappe, A., Schoon, H.A., 2009.
Immunohistochemical characterization of the extracellular matrix in normal mitral valves and
in chronic valve disease (endocardiosis) in dogs. Research in Veterinary Science 87, 277-283.

Bischoff, J., Aikawa, E., 2011. Progenitor cells confer plasticity to cardiac valve
endothelium. Journal of Cardiovascular Translational Research 4, 710-719.

- Black, A., French, A.T., Dukes-McEwan, J., Corcoran, B.M., 2005. Ultrastructural
 morphologic evaluation of the phenotype of valvular interstitial cells in dogs with
 myxomatous degeneration of the mitral valve. American Journal of Veterinary Research 66,
 1408-1414.
- 379
- Blevins, T.L., Carroll, J.L., Raza, A.M., Grande-Allen, K.J., 2006. Phenotypic
 characterization of isolated valvular interstitial cell subpopulations. The Journal of Heart
 Valve Disease 15, 815-822.
- 383
- Butcher, J.T., Markwald, R.R., 2007. Valvulogenesis: the moving target. Philosophical
 Transactions of the Royal Society B: Biological Sciences 362, 1489-1503.
- 386
- Butcher, J.T., Nerem, R.M., 2007. Valvular endothelial cells and the mechanoregulation of
 valvular pathology. Philosophical Transactions of the Royal Society B: Biological Sciences
 362, 1445-1457.
- 390
- Cheung, W.Y., Young, E.W.K., Simmons, C.A., 2008. Techniques for isolating and purifying
 porcine aortic valve endothelial cells. The Journal of Heart Valve Disease 17, 674-681.
- 393
- Ci, H.B., Ou, Z.J., Chang, F.J., Liu, D.H., He, G.W., Xu, Z., Yuan, H.Y., Wang, Z.P., Zhang,
 X., Ou, J.S., 2013. Endothelial microparticles increase in mitral valve disease and impair
 mitral valve endothelial function. American Journal of Physiology-Endocrinology And
 Metabolism 304, E695-E702.
- 398
- Corcoran, B.M., Black, A., Anderson, H., McEwan, J.D., French, A., Smith, P., Devine, C.,
 2004. Identification of surface morphologic changes in the mitral valve leaflets and chordae
 tendineae of dogs with myxomatous degeneration. American Journal of Veterinary Research
 65, 198-206.
- 403
- Cuy, J.L., Beckstead, B.L., Brown, C.D., Hoffman, A.S., Giachelli, C.M., 2003. Adhesive
 protein interactions with chitosan: Consequences for valve endothelial cell growth on
 tissue-engineering materials. Journal of Biomedical Materials Research Part A 67, 538-547.
- 407
- Della Rocca, F., Sartore, S., Guidolin, D., Bertiplaglia, B., Gerosa, G., Casarotto, D.,
 Pauletto, P., 2000. Cell composition of the human pulmonary valve: A comparative study
 with the aortic valve The VESALIO* project. Annals of Thoracic Surgery 70, 1594-1600.
- 411
- Disatian, S., Ehrhart, E.J., 3rd, Zimmerman, S., Orton, E.C., 2008. Interstitial cells from dogs
 with naturally occurring myxomatous mitral valve disease undergo phenotype transformation.
 Journal of Heart Valve Disease 17, 402-411; Discussion 412.
- 415
- 416 Disatian, S., Lacerda, C., Orton, E.C., 2010. Tryptophan hydroxylase 1 expression is
 417 increased in phenotype-altered canine and human degenerative myxomatous mitral valves.
 418 Journal of Heart Valve Disease 19, 71-78.
- 419
- 420 Durbin, A., Nadir, N.A., Rosenthal, A., Gotlieb, A.I., 2005. Nitric oxide promotes in vitro
- 421 interstitial cell heart valve repair. Cardiovascular Pathology 14, 12-18.
- 422
- 423 Engler, A.J., Sen, S., Sweeney, H.L., Discher, D.E., 2006. Matrix elasticity directs stem cell
- 424 lineage specification. Cell 126, 677-689.

- Fayet, C., Bendeck, M.P., Gotlieb, A.I., 2007. Cardiac valve interstitial cells secrete
 fibronectin and form fibrillar adhesions in response to injury. Cardiovascular Pathology 16,
 203-211.
- 429

Flanagan, T.C., Black, A., O'Brien, M., Smith, T.J., Pandit, A.S., 2006. Reference models for
mitral valve tissue engineering based on valve cell phenotype and extracellular matrix
analysis. Cells Tissues Organs 183, 12-23.

433

Fulton, D., Papapetropoulos, A., Zhang, X., Catravas, J.D., Hintze. T.H., Sessa, W.C., 2000.
Quantification of eNOS mRNA in the caninecardiac vasculature by competitive PCR.
American Journal of Physiology-Heart and Circulatory Physiology 278, H658-H665.

437

Gould, R.A., Butcher, J.T., 2010. Isolation of valvular endothelial cells. Journal of Visualized
Experiments 46, 2158-2162.

440

Gupta, V., Werdenberg, J.A., Mendez, J.S., Jane Grande-Allen, K., 2008. Influence of strain
on proteoglycan synthesis by valvular interstitial cells in three-dimensional culture. Acta
Biomaterialia 4, 88-96.

444

Gupta, V., Barzilla, J.E., Mendez, J.S., Stephens, E.H., Lee, E.L., Collard, C.D., Laucirica,
R., Weigel, P.H., Grande-Allen, K.J., 2009. Abundance and location of proteoglycans and
hyaluronan within normal and myxomatous mitral valves. Cardiovascular Pathology 18, 191197.

449

Hadian, M., Corcoran, B.M., Han, R.I., Grossmann, J.G., Bradshaw, J.P., 2007. Collagen
organization in canine myxomatous mitral valve disease: an x-ray diffraction study.
Biophysical Journal 93, 2472-2476.

453

Han, R.I., Black, A., Culshaw, G.J., French, A.T., Else, R.W., Corcoran, B.M., 2008.
Distribution of myofibroblasts, smooth muscle-like cells, macrophages, and mast cells in
mitral valve leaflets of dogs with myxomatous mitral valve disease. American Journal of
Veterinary Research 69, 763-769.

458

Han, R.I., Black, A., Culshaw, G., French, A.T., Corcoran, B.M., 2010. Structural and
cellular changes in canine myxomatous mitral valve disease: an image analysis study. Journal
of Heart Valve Disease 19, 60-70.

462

Han, R.I., Clark, C.H., Black, A., French, A., Culshaw, G.J., Kempson, S.A., Corcoran, B.M.,
2013a. Morphological changes to endothelial and interstitial cells and to the extra-cellular
matrix in canine myxomatous mitral valve disease (endocardiosis). The Veterinary Journal
197, 388-394.

467

Han, R.I., Impoco, G., Culshaw, G., French, A.T., Black, A., Corcoran, B.M., 2013b. Cell
maceration scanning electron microscopy and computer-derived porosity measurements in
assessment of connective tissue microstructure changes in the canine myxomatous mitral
valve. The Veterinary Journal 197, 502-505.

472

Heaney, A.M., Bulmer, B.J., Ross, C.R., Schermerhorn, T., 2009. A technique for in vitro
culture of canine valvular interstitial cells. Journal of Veterinary Cardiology 11, 1-7.

476 Hoerstrup, S.P., Zünd, G., Schoeberlein, A., Ye, Q., Vogt, P.R., Turina, M.I., 1998. 477 Fluorescence activated cell sorting: a reliable method in tissue engineering of a bioprosthetic 478 heart valve. The Annals of Thoracic Surgery 66, 1653-1657. 479 480 Lester, W.M., Damji, A.A., Gedeon, I., Tanaka, M., 1993. Interstitial cells from the atrial and 481 ventricular sides of the bovine mitral valve respond differently to denuding endocardial 482 injury. In Vitro Cellular & Developmental Biology 29A, 41-50. 483 484 Liu, A.C., Joag, V.R., Gotlieb, A.I., 2007. The emerging role of valve interstitial cell 485 phenotypes in regulating heart valve pathobiology. The American Journal of Pathology 171, 486 1407-1418. 487 488 Mahler, G.J., Farrar, E.J., Butcher, J.T., 2013. Inflammatory Cytokines Promote Mesenchymal Transformation in Embryonic and Adult Valve Endothelial Cells. 489 490 Arteriosclerosis Thrombosis and Vascular Biology 33, 121-130. 491 492 Olsen, L.H., Martinussen, T., Pedersen, H.D., 2003. Early echocardiographic predictors of 493 myxomatous mitral valve disease in dachshunds. Veterinary Research 152, 293-297. 494 495 Orton, E.C., Lacerda, C.M., MacLea, H.B., 2012. Signaling pathways in mitral valve 496 degeneration. Journal of Veterinary Cardiology 14, 7-17. 497 498 Paranya, G., Vineberg, S., Dvorin, E., Kaushal, S., Roth, S.J., Rabkin, E., Schoen, F.J., 499 Bischoff, J., 2001. Aortic valve endothelial cells undergo transforming growth factor- β mediated and non-transforming growth factor- β -mediated transdifferentiation in vitro. 500 501 American Journal of Pathology 159, 1335-1343. 502 503 Paruchuri, S., Yang, J.H., Aikawa, E., Melero-Martin, J.M., Khan, Z.A., Loukogeorgakis, S., 504 Schoen, F.J., Bischoff, J., 2006. Human pulmonary valve progenitor cells exhibit 505 endothelial/mesenchymal plasticity in response to vascular endothelial growth factor-A and 506 transforming growth factor-beta2. Circulation Research 99, 861-869. 507 508 Person, A.D., Klewer, S.E., Runyan, R.B., 2005. Cell biology of cardiac cushion 509 development. International Review of Cytology 243, 287-335. 510 511 Pusztaszeri, M.P., Seelentag, W., Bosman, F.T., 2006. Immunohistochemical expression of 512 endothelial markers CD31, CD34, von Willebrand factor, and Fli-1 in normal human tissues. 513 Journal of Histochemistry & Cytochemistry 54, 385-395. 514 515 Rabkin, E., Aikawa, M., Stone, J.R., Fukumoto, Y., Libby, P., Schoen, F.J., 2001. Activated 516 interstitial myofibroblasts express catabolic enzymes and mediate matrix remodeling in 517 myxomatous heart valves. Circulation 104, 2525-2532. 518 519 Simmons, C.A., Grant, G.R., Manduchi, E., Davies, P.F., 2005. Spatial heterogeneity of 520 endothelial phenotypes correlates with side-specific vulnerability to calcification in normal 521 porcine aortic valves. Circulation Research 96, 792-799. 522

Stephens, E.H., Durst, C.A., West, J.L., Grande-Allen, K.J., 2011. Mitral valvular interstitial
cell responses to substrate stiffness depend on age and anatomic region. Acta Biomaterialia 7,
75-82.

526

Taylor, P.M., Allen, S.P., Yacoub, M.H., 2000. Phenotypic and functional characterization of
interstitial cells from human heart valves, pericardium and skin. Journal of Heart Valve
Disease 9, 150-158.

530

Waxman, A.S., Kornreich, B.G., Gould, R.A., Sydney Moïse, N., Butcher, J.T., 2012.
Interactions between TGFβ1 and cyclic strain in modulation of myofibroblastic
differentiation of canine mitral valve interstitial cells in 3D culture. Journal of Veterinary
Cardiology 14, 211-221.

535

Wiester, L., Giachelli, C., 2003. Expression and function of the integrin alpha9beta1 in
bovine aortic valve interstitial cells. Journal of Heart Valve Disease 12, 605.

538

Wills, T.B., Heaney, A.M., Wardrop, K.J., Haldorson, G.J., 2009. Immunomagnetic isolation
of canine circulating endothelial and endothelial progenitor cells. Veterinary Clinical
Pathology 38(4), 437-442.

542

543 Wylie-Sears, J., Aikawa, E., Levine, R.A., Yang, J.H., Bischoff, J., 2011. Mitral valve 544 endothelial cells with osteogenic differentiation potential. Arteriosclerosis, Thrombosis, and 545 Vascular biology 31, 598-607.

546

Xu, S., Liu, A.C., Kim, H., Gotlieb, A.I., 2012. Cell density regulates in vitro activation of
 heart valve interstitial cells. Cardiovascular Pathology 21, 65-73.

Table 1. PCR sequences for PCR characterization of valve endothelial cells (VECs) and valve

552 interstitial cells (VICs).

Gene	Primer Sequence (5'-3')	Product size
	AATCCCAAATTCCACGTCAG	
CD31	GAATGGAGCACCACAGGTTT	346 bp
	CTGGGAGAAGAGAGTCACGG	
vWF1	GTGGATGGAGTACACGGCTT	235 bp
	GGCTGTACCTGGATGAGAGG	
vWF2	GACAGGACAGGCTCCTTTTG	228 bp
	AATATAGGGCCCCGGCTCACTCAA	
<i>vWF3</i> (Fulton et al., 2000)	ACATCCCCGGGCCTCTTCTCATTC	512 bp
	AAGAACGGCGTGATTCTGAG	
SM22	CGGTAGTGCCCATCATTCTT	269 bp
	GGGGATGGGACAAAAGGACA	
a-SMA	GCCACGTAGCAGAGCTTCTCCTTGA	525 bp
	AGAAGCGAGCTGGAAAACTG	
SMemb	TCTTGCTCTGTCCGATTCTG	252 bp
	GGAGCAGCAGAACAAGATCC	
Vimentin	AGACGTGCCAAAGAAGCATT	282 bp
	CATCAACGGGAAGTCCATCT	
GAPDH	GTGGAAGCAGGGATGATGTT	428 bp

553

554

- Table 2. Details of the antibodies used for canine mitral valve endothelial (VEC) and valve
- 557 interstitial cell (VIC) immunocytochemistry characterisation.

Primary antibody	Species raised	Cat Number & Supplier	Dilution
Anti-CD31	Rabbit polyclonal	Ab28364, Abcam, UK	1:200
Anti-SMemb	Rabbit polyclonal	Ab24761, Abcam, UK	1:400
Anti-vimentin	Mouse monoclonal	V6389, Sigma, USA	1:1600
Anti-a-SMA	Mouse monoclonal	A2547, Sigma, USA	1:400
Secondary antibody			
Anti-rabbit IgG (H+L)	Goat	A11011, Invitrogen, UK	1:500
Alexa Fluor 508			
Anti-mouse IgG (H+L) Alexa Fluor488	Goat	A10667, Invitrogen, UK	1:500

561 Figures Legends

- 562
- 563 **Figure 1.** Canine mitral valve cell morphologies.

564 Upper Panel; primary isolated canine mitral valve endothelial cell (VEC) morphology. A. At 565 the early stage, single colony clusters randomly appeared in the cultures. B. Once the cells 566 reached confluence, the majority exhibited a cobblestone morphology and were growth-567 contact inhibited. Scale bar = $100 \mu m$.

568 Lower Panel; primary cultured valve interstitial cell (VIC) morphology before confluence (C) 569 and at confluence (D). Cells in VIC cultures exhibited fibroblast-like morphology, and tended 570 to arrange in a 'hill and valley' pattern in confluent cultures. Scale bar = $200 \,\mu m$.

Figure 2. Canine mitral valve endothelial cells (VECs) and valve interstitial cells (VICs) characterised by RT-PCR. Upper panel: VECs; the endothelial cell marker (CD31) and the activated VIC markers (α -SMA, SMemb and SM22) were detected in VECs. Vimentin and GAPDH were used as internal controls. Lower panel: VICs; cells were negative for the VEC marker CD31, but positive for α -SMA, SMemb, SM22 and vimentin. cDNA template derived from a VEC culture was used to amplify CD31 transcripts in the same experiment as a positive control (data not shown). Vimentin and GAPDH were used as internal controls.

578 Figure 3. Canine mitral valve endothelial cells (VECs) characterised bv 579 immunohistochemistry and DiI-Ac-LDL. Upper panel; DiI-Ac-LDL uptake by VECs in 580 culture (A) and cell distribution illustrated by bright field microscopy (B). Scale bar = 100581 µm. Lower panel; characterization of canine mitral VEC culture by immunofluorescence. 582 Cell expressed CD31 (red in C), vimentin (green in G) and SMemb (red in K), but were 583 negative for α -SMA (O). DAPI (blue) stained cell nuclei. DAPI channel (A, E, I and M), target protein channel (B, F, J and N) and merged images (C, G, K and O) are shown 584 585 respectively for each marker. The same primary cultures cell lines served as negative controls 586 (primary antibodies were omitted; D, H and L). Cells from a VIC culture showing expression 587 of α -SMA served as a positive control (P). Scale bar = 50 μ m.

Figure 4. Endothelial properties of canine mitral valve endothelial cells (VECs) under
different culture conditions. EBM cultures (left panel) and AF-12 cultures of VECs (right
panel) showing uptake of DiI-Ac-LDL (upper panel) and expression of CD31 (lower panel).
There is a greater intensity of staining for EBM cultures. DAPI (blue) stained cell nuclei.
Scale bar = 100 µm

593

Figure 5. Canine mitral canine mitral valve endothelial cells (VECs) under different culture conditions, characterised by semi-quantitative RT-PCR (arbitrary units compared to the reference gene GAPDH). Cells in EBM demonstrated higher expression of CD31 and decreased expression of α -SMA and SMemb compared to VECs and VICs cultured in AF-12.

598

599 6. cells (VICs) Figure Canine mitral valve interstitial characterised bv 600 immunohistochemistry and DiI-Ac-LDL. Upper panel; lack of DiI-Ac-LDL uptake by VICs (A) and cell distribution illustrated by bright field microscopy (B). Scale bar = 200 μ m. 601 602 Lower panel; characterization of canine mitral VIC culture by immunofluorescence. Cells 603 expressed vimentin (green in G), SMemb (red in K) and α-SMA (green in O), but were 604 negative for CD31 (C). DAPI (blue) stained cell nuclei. DAPI channel (A, E, I and M), target 605 protein channel (B, F, J and N) and merged images (C, G, K and O) are shown respectively 606 for each marker. The same cell lines served as negative controls (primary antibodies were

- omitted; H, L and P). Cells from a VEC culture showing expression of α -SMA served as a positive control (D). Scale bar = 50 μ m.



Figure2 Click here to download high resolution image









DII-Ac-LDL



Figure5 Click here to download high resolution image















Highlights:

- Successful culturing and characterising of valve interstitial and endothelial cells from the same subject.
- First reported culturing of valve interstitial cells (VICs) that did not favour a migratory phenotype.
- First reported culturing of canine valve endothelial cells (VECs).
- Assessment of the value of multiple cell markers in cell phenotyping.
- Identification of CD31 expression and acetylated low density lipoprotein (Dil-Ac-LDL) uptake as the prime markers for differentiating VECs from VICs.
- Identification of phenotypic plasticity under culture conditions and optimal passage for attaining the purest VEC population.

Highlights:

1. Successful culture and characterisation of valve interstitial and endothelial cells

2. Identification of CD31 expression and Dil-Ac-LDL uptake as the prime markers for differentiating VECs from VICs.

3. A specific endothelial culture medium for primary VEC cultures enhanced the endothelial properties of the cells

We would like to thank the reviewers for their comments. There were instances where the line numbers quoted did not match those of the current version of the manuscript, but we think we have managed to respond to each comment in turn.

Reviewer 1

Point 1: "The authors are not clear on whether they were able to culture a pure VEC population that then subsequently differentiated (e.g. via EndMT) or whether the population was heterogeneous (VEC+VIC) to start with. Their aSMA data suggests the latter, but if they combined with IHC of normal canine MV, it might support that canine VEC possess aSMA expression in vivo."

Response: Indeed, sub-endothelial VIC contamination in primary VECs isolated by enzymatic digestion has been recognised as a common technical issue (Cheung et al., 2008; Hoerstrup et al., 1998). However, we cannot completely rule out the possibility of cell differentiation in culture. Firstly, the α -SMA expression in VECs was mainly identified at gene level and it was almost absent at protein level. It is difficult to tell whether the α -SMA gene expression was from differentiating VECs or contaminating cells or from both. Secondly, which is probably more robust evidence, is the extensive expression of activated mesenchymal marker SMemb expression in the VEC cultures. This marker has been found highly expressed in embryonic VECs during development or in activated VICs during valve disease processes, while in healthy adult heart valves, the expression has been described as absent/minimally expressed (Aikawa et al., 2006). Therefore, we propose that the canine mitral VECs we cultured in vitro may possess some embryonic differentiation potentials and we have addressed the above points in the discussion of the revised manuscript.

With regards to native value α -SMA expression, we did look at it in normal canine MV by immunofluorescence (data not shown); the expression was very limited and mainly restricted to atrialis, and this finding is consistent with our own and others previous reports (Han et al., 2008; Disatian et al., 2008).

Point 2: "Their method is largely similar to that of Gould et al (JoVE 2012), which raises the question of whether ECM coating is essential for VEC adhesion. It would be good to know, as that can be a real problem out of the gate for others."

Response: From our experience, the answer is no. We once accidentally cultured our VECs on polypropylene flasks without gelatin coating; the cells still adhered and proliferated, however, with a different morphology and the efficiency of adherence was not investigated (data not shown). Indeed, it has been reported that additional ECM coating to plastic culture flasks can enhance endothelial cell adherence, proliferation and affect endothelial cell morphologies (Relou et al., 1998; Schelling et al., 1988; Xia et al., 2011). Both gelatin and collagen are commonly used for valve endothelial culture in previous reports (Wylie-Sears et al., 2011; Gould and Butcher, 2010). For these reasons, we would recommend matrix coating for endothelial culture.

Point 3: "Also, it isn't clear from my reading of the manuscript whether the VEC cultured in this study ended up being suitable for follow-on experiments on VEC behavior. Are enough cells available at passage 3 without compromising homogeneity?

Response: The answer is yes. Primary VECs P0 yields to $0.66 \times 10^6 - 3.5 \times 10^6$ depending on dog size and valve leaflet collected(anterior/posterior or both), and on average 1.5×10^6 approximately. This can further yield to approximate 1.0×10^7 at P1, and around 1×10^8 at P2. This number should be sufficient to do routine experiments such as immunocytochemistry (usually 1×10^4 cells per sample)

and RT-PCR (usually $0.5-1 \times 10^6$ cells for RNA extraction). In terms of cell purity, according to our findings, VECs cultured in AF-12 medium (without additional purification or endothelial medium replacement) showed at least 50%-60% typical endothelial cobblestone morphology within 3 passages. We didn't do endothelial phenotypic quantification on immunocytochemistry, but we presumed the actual number would be higher than morphological observation, as CD31 or DiI-Ac-LDL positive cells were heterogeneous in morphology (not limited to cobblestone shape only).

Point 4: "The authors suggest EBM culture can improve EC phenotype homogeneity at a sacrifice of proliferation, but does this approach create enough cells for an experimental assay?"

Response: Yes. When used EBM for primary P0 VECs culture, there was cell apoptosis and endothelial colony disappear during the early stages, but the cells started to proliferate rapidly in later culture periods. For those VECs experiencing EBM switch at P1 or P2, they did not show dramatically decreased proliferation compared to AF-12 culture. To avoid confusion the P0 EBM finding has been removed from the manuscript.

Point 5: "Do the authors recommend using EBM during whatever treatment would be performed, or switching to AF-12 Media?"

Response: Once starting culture the VECs in EBM we would continue with EBM rather than switching to AF-12. For experiments specifically investigating VEC behaviour and function, we would recommend to use EBM rather than AF-12, as the former is more optimal in maintaining VEC phenotype.

Point 6: "As a minor but important comment, having standard deviation bars on Figure 6 would be helpful to assess the expected variation in isolation quality and/or subsequent culture performance of the phenotypes of these cells."

Authors' Response: We've added standard error bars on Figure 5 (original Figure 6) in the revised manuscript (n=3).

Reviewer 2

Point 1: "Not sure about numbering of subsections in M and M and in Results - what is editorial policy?"

Response: We've removed the numbering of subsections in the revised manuscript.

Point 2: "Line 88 - presumably valves showed no gross evidence of MMVD?"

Response: We've addressed this point in the revised manuscript; please see Line 84-85.

Point 3: "Line 102 - a bit more pre-amble on the media experiments would be useful, just to put them into context. Typo – passage (P1)."

Response: We've addressed this point in the revised manuscript; please see Line 109-121.

Point 4: "Line 110 - might be useful to add a third sub-heading on cell passage (see later comment)."

Response: We have addressed this point in Line 104-107 in the revised manuscript.

Point 5: "Lines 117, 135 and 149 - cells from which passage were used on these experiments - was this consistent?"

Response: We have addressed this point in Line 112, 200 and 222 in the revised manuscript.

Point 6: "Lines 137 and 149 - details of slides - chamber slides mentioned."

Response: We have addressed this point in Line 147-148 in the revised manuscript.

Point 7: "Line 164 - was detected by RT-PCR in VEC culture?"

Response: We have addressed this point in Line 201-202 in the revised manuscript.

Point 8: "Line 177 - was detected by RT-PCR for....?"

Response: We have addressed this point in Line 223-224 in the revised manuscript.

Point 9: "Move section on effect of culture medium (lines 182-196) to before section on VIC morphology and characterization to be consistent with Methods."

Response: We have moved this section as requested; see Line 214-220.

Point 10: "Line 194 - typo - EBM.Both..."

Response: We have addressed this point in the revised manuscript.

Point 11: "Line 201 - Waxmann reference not in references."

Response: We have addressed this point Line 534 in the revised manuscript.

Point 12: "Line 239 - ...although it did not preserve...?"

Response: We have addressed this point in the revised manuscript.

Point 13: "Line 256 - Liu et al (2007),..."

Response: We have addressed this point in the revised manuscript.

Point 14: "Line 265 - Han 2013 - a or b?"

Response: We cannot locate this comment in the original manuscript. Effectively has been resolved

Point 15: "Several references in reference list are incorrect or incomplete - e.g. lines 381 and 386, Paranya reference has a space in it. There is variability in how journals are cited."

Response: We have addressed these inconsistencies and errors in the reference section of the revised manuscript.

Point 16: "Images - quality of text on images is poor."

Response: We have improved the text quality in the revised manuscript.

Point 17: "Figure 3 - top panel association of positive staining with cells shown in the bright field image is not as convincing as the other images."

Response: We have checked the image and can confirm that the bright field image is correctly matched with the fluorescence image. However, the focusing of the bright field image was not optimal. We have tried to enhance that by adjusting brightness and contrast of the bright field image in the revised manuscript.

Point 18: "Figures 3 and 4 - for clarity legends should explain column titles - what do the target protein images show, what is the merged image."

Response: We have addressed this point in the revised manuscript. Please see legends of Figure 3 and 6 (Figure 4 in original manuscript).

Comments E Bloome

1. The current Figure 4 is a copy of Figure 6, I think Figure 4 should be what was Figure 5 in the previous submission.

Figure identities seem correct; will up-load fresh set of figure images.

2. Cannot find Bischoff (2001) reference in the text.

No Bischoff 2001 in the text or reference list.

Comments A. Philby

1. Please check author's initials, e.g. "D.A. Argyle".

Changed to D.J.

2. Some reformatting of the Corresponding author's details will be required.

Not sure what is required.

3. Most non-standard abbreviations should be defined when first used in the Abstract and again in the Main text. Abbreviations that do not need to be defined include PCR, ELISA, DNA and RNA. In the Abstract, please change "reverse transcription polymerase chain reaction (RT-PCR)" to "reverse transcription PCR".

MS checked and corrected

4. Groups of references cited in the text should be arranged chronologically according to the idiosyncratic journal style, e.g. "Aupperle et al., 2009; Aupperle et al., 2010; Black et al., 2005; Hadian et al., 2007; Han et al., 2010; Han et al., 2013b" should be cited as "Black et al., 2005; Hadian et al., 2007; Aupperle et al., 2009, 2010; Han et al., 2010, 2013b".

Corrected; EndNote seems to think TVJ format is different!

5. Most units should be separated from values by a space, although percentages should be written without a space. Similalry, when writing temperatures, the value should be separated from the symbol for degrees Celsius by a space, e.g. "37°C" should be written as "37 °C" using the Word symbol for degrees Celsius rather than a superscripted letter "o".

MS checked and corrected

6. Some reformatting of tables and figure legends will be required.

Not sure what is required?

7. Please check spacing and punctuation throughout the manuscript.

Done.

8. Please reformat the Highlights with bullet points and define abbreviations. Each Highlight should be no more than 125 characters including spaces.

Done.



ROYAL (DICK) SCHOOL OF VETERINARY STUDIES The University of Edinburgh Easter Bush Campus Roslin Midlothian EH25 9RG 0131 650 6241 http://www.vet.ed.ac.uk/

29/7/2014

Please find attached "Canine Mitral Valve Interstitial and Endothelial Cell Culture and Characterisation" for consideration for publication in TVJ.

The paper describes the culturing of valve interstitial and endothelial cells from canine mitral valves. This is important as there is increasing interest in using culture systems to investigate mechanisms of pathogenesis in canine mitral valve disease, not least because of the difficulties in sourcing native valves.

No previous study has reported on canine VECs, which are much harder to grow than VICs, and previous reports of VICs have used techniques that only cultured migratory phenotypes.

This study provides the basic information necessary for accurate VIC and VEC culture and the techniques required for accurate phenotyping.

Brendom Corcolan

Brendan Corcoran