THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Host Innate Immune Responses to Infection by Avian- and Bat-Borne Viruses

Citation for published version:

Giotis, ES, Matthews, DA & Smith, J 2021, 'Host Innate Immune Responses to Infection by Avian- and Bat-Borne Viruses', *Frontiers in Cellular and Infection Microbiology*. <https://doi.org/10.3389/fcimb.2021.651289>

Digital Object Identifier (DOI):

[10.3389/fcimb.2021.651289](https://doi.org/10.3389/fcimb.2021.651289)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Frontiers in Cellular and Infection Microbiology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Editorial: Host Innate Immune Responses to Infection by Avian- and Bat-Borne Viruses

Efstathios S. Giotis^{1,2*}, David A. Matthews³ and Jacqueline Smith⁴

¹ Section of Virology, School of Medicine, Imperial College London, London, United Kingdom, ² School of Life Sciences, University of Essex, Colchester, United Kingdom, ³ School of Cellular and Molecular Medicine, Faculty of Life Sciences, University of Bristol, Bristol, United Kingdom, ⁴ The Roslin Institute and Royal (Dick) School of Veterinary Studies R(D)SVS, University of Edinburgh, Edinburgh, United Kingdom

Keywords: avian, zoonotic, virus, innate immunity, bats (Chiroptera)

Editorial on the Research Topic

Host Innate Immune Responses to Infection by Avian- and Bat-Borne Viruses

OPEN ACCESS

Edited by:

Curtis Brandt,
University of Wisconsin–Madison,
United States

Reviewed by:

Teneema Kuriakose,
St. Jude Children's Research Hospital,
United States

*Correspondence:

Efstathios S. Giotis
e.giotis@imperial.ac.uk

Specialty section:

This article was submitted to
Virus and Host,
a section of the journal
Frontiers in Cellular
and Infection Microbiology

Received: 09 January 2021

Accepted: 01 February 2021

Published: 24 February 2021

Citation:

Giotis ES, Matthews DA and Smith J
(2021) Editorial: Host Innate Immune
Responses to Infection by Avian- and
Bat-Borne Viruses.
Front. Cell. Infect. Microbiol. 11:651289.
doi: 10.3389/fcimb.2021.651289

The COVID-19 pandemic has generated many urgent questions on the origin, trajectory, and host preference of its causative betacoronavirus SARS-CoV-2, as well as renewed focus on other potentially zoonotic viruses. Several species of birds and wild bats can serve as reservoirs and/or mechanical vectors for many infectious viruses including influenza-A, SARS-CoV, MERS, and Ebola. Although substantial progress has been made, there are still major gaps in understanding the emergence, transmission, and adaptation of zoonotic avian- and bat-borne viruses. A major challenge is the dearth of suitable infection and immunity models. Extrapolating data from infection studies in human cell lines or rodents is limiting, as evolutionarily optimized immune factors function differently in non-hosts.

Viral infection triggers anti-viral host defenses and an inflammatory response, broadly coordinated by type I interferon (IFN) and NF- κ B, respectively. IFN induces production of hundreds of “interferon-stimulated genes” (ISGs), limiting virus replication until the adaptive immune response can “clean-up.” Rapid initiation of innate immune responses depends on viral recognition by pattern recognition receptors (PRRs) which are expressed in tissues where the virus replicates. Despite the high degree of evolutionary conservation, and assumed similarity in function, there are significant differences between the immune gene repertoires of birds and mammals and bats and other mammals. Ducks utilize many of the mammalian PRRs (i.e. RIG-I, TLR7, and TLR3) and their downstream adaptors to detect and inhibit the replication of influenza viruses (Campbell and Magor). In chickens, key innate immune genes, such as *IRF3* and *RIG-I*, appear to have been lost (Chen et al., 2013). However, chicken cells are capable of initiating an interferon response after virus infection and inducing the expression of hundreds of ISGs (Giotis et al., 2016; Giotis et al., 2017).

Differences in IFN-inducing capacity between virus strains is an important parameter contributing to the host response to viral infection and virulence. Influenza-A virus circulates in multiple hosts causing economic burden in the poultry industry, human epidemics, and

occasionally pandemics. Influenza-A virus contains eight negative-strand RNA segments, which are thought to encode 10 viral proteins as well as seven newly identified proteins such as PB1-F2, PA-X whose role is not entirely clear (Ma et al.). The full-length PA-X and PB1-F2 proteins in a recombinant 2009 pH1N1 virus have been reported to enhance viral replication *in vitro*, enhancing viral virulence *in vivo* mainly through simultaneously mediated host innate immune response (Ma et al.).

IBDV (causing Infectious Bursal Disease) is another economically important (but not zoonotic) virus for the poultry industry. Very virulent strains of IBDV (vIBDV) emerged in the 1980s, causing up to 60% mortality in some commercial flocks for reasons that are poorly understood (Dulwich et al., 2017). Dulwich et al. extended these observations by demonstrating that a vIBDV strain is able to down-regulate type I IFN and pro-inflammatory cytokine responses compared to a classical field strain both *in vitro* and *in vivo*.

The successful control of virus infection requires the coordination of both the innate and adaptive immune systems. During chronic viral infections the increased level or duration of stimulation of virus-specific CD8 T-cells leads to non-functional state called T-cell exhaustion (Freeman et al., 2006). Recent studies have shown that targeting the immunoreceptor molecule PD1 and its ligand PD-L1 can reverse the exhausted T-cell response (Freeman et al., 2006). The chicken PD-1/PD-L1 might also be used in the treatment of oncogenic avian viruses such as Marek's disease virus (Reddy et al.).

Current knowledge on bat IFN responses is rudimentary, with most of this obtained in studies of the Australian flying fox (*Pteropus alecto*) (Clayton and Munir). Sequencing of bat genomes suggests that evolution of "metabolically costly" flight imposed adaptations on their innate immune-, DNA damage- and inflammatory-response systems (Zhang et al., 2013). Key antiviral immunity components are conserved in bats, e.g. IFNs and their receptors and ISGs. However, genes that activate the inflammasome and/or IFN pathways are either missing or have altered function (Clayton and Munir). Studies have inferred a pattern of constitutively expressed IFN- α in unstimulated *Pteropus alecto* cells, which may provide a "switched-on" defense mechanism that blunts virus replication and pathogenesis (Zhou et al., 2016), and can explain why bats are asymptomatic reservoirs of viruses. The remarkable ability of bats to coexist with a wide range of potentially zoonotic viruses is exemplified by the evolutionarily

distinct bat influenza-A-like viruses H17N10 and H18N11 (BatIVs). Unlike classical influenza-A viruses, the surface glycoproteins of BatIVs neither bind nor cleave sialic acid receptors, but rather use the trans-species conserved MHC-II proteins (Giotis et al., 2019) to gain cell entry. The scientific evidence so far indicate a limited spillover risk for BatIVs, but data is not conclusive enough to dismiss the possibility of zoonotic transmission (Giotis).

Full-length genome sequences obtained from COVID-19 patients demonstrated that SARS-CoV-2 shares 79.5% sequence identity to SARS-CoV BJ01 strain and a 96% sequence identity to the bat coronavirus RaTG13, suggestive that bats are the original source of the virus (Zhou et al., 2020). The receptor binding protein spike (S) viral gene is highly divergent to other CoVs but has a 93.1% nucleotide identity to RaTG13 (Zhou et al., 2020). Molecular modeling and functional studies revealed that the host surface protein ACE2 mediates cell entry of SARS-CoV-2 and the serine protease TMPRSS2 is essential for S priming (Minakshi et al.). Furthermore, the aftermath of SARS-CoV-2 internalisation is governed by a complex and largely unknown network of host-pathogen interactions. Transcriptomic analysis of human alveolar and bronchial epithelial cells confirmed that the *CSF1* gene, a known target of the microRNA miR-1207-5p, is over-expressed following SARS-CoV-2 infection (Bertolazzi et al.). *CSF1* enhances macrophage recruitment and activation and its overexpression may contribute to the acute inflammatory response observed in severe COVID-19.

In conclusion, recent virus outbreaks confirmed the inextricable nature of human and animal health and disease. Interdisciplinary approaches are urgently needed to assess and prevent the spillover of avian- and bat-borne viruses using an organismal biology approach and focusing on viral ecology, diversity, and interactions with the host.

AUTHOR CONTRIBUTIONS

EG wrote the first draft and JS and DM edited and commented on the draft. All authors contributed to the article and approved the submitted version.

REFERENCES

- Chen, S., Cheng, A., and Wang, M. (2013). Innate sensing of viruses by pattern recognition receptors in birds. *Vet. Res.* 44. doi: 10.1186/1297-9716-44-82
- Dulwich, K. L., Giotis, E. S., Gray, A., Nair, V., Skinner, M. A., and Broadbent, A. J. (2017). Differential gene expression in chicken primary B cells infected *ex vivo* with attenuated and very virulent strains of infectious bursal disease virus (IBDV). *J. Gen. Virol.* 98, 2918–2930. doi: 10.1099/jgv.0.000979
- Freeman, G. J., Wherry, E. J., Ahmed, R., and Sharpe, A. H. (2006). Reinvigorating exhausted HIV-specific T cells via PD-1-PD-L1 ligand blockade. *J. Exp. Med.* 203, 2223–2227. doi: 10.1084/jem.20061800
- Giotis, E. S., Robey, R. C., Skinner, N. G., Tomlinson, C. D., Goodbourn, S., and Skinner, M. A. (2016). Chicken interferome: avian interferon-stimulated genes identified by microarray and RNA-seq of primary chick embryo fibroblasts treated with a chicken type I interferon (IFN- α). *Vet. Res.* 47, 75. doi: 10.1186/s13567-016-0363-8
- Giotis, E. S., Ross, C. S., Robey, R. R., Nothdurft, A., Goodbourn, S., and Skinner, M. A. (2017). Constitutively elevated levels of SOCS1 suppress innate responses in DF-1 immortalised chicken fibroblast cells. *Sci. Rep.* 7, 17485. doi: 10.1038/s41598-017-17730-2
- Giotis, E. S., Carnell, G., Young, E. F., Ghannay, S., Soteropoulos, P., Wang, L. F., et al. (2019). Entry of the bat influenza H17N10 virus into mammalian cells is enabled by the MHC class II HLA-DR receptor. *Nat. Microbiol.* 4, 2035–2038. doi: 10.1038/s41564-019-0517-3
- Zhang, G., Cowled, C., Shi, Z., Huang, Z., Bishop-Lilly, K. A., Fang, X., et al. (2013). Comparative analysis of bat genomes provides insight into the evolution of flight and immunity. *Science* 339, 456–460. doi: 10.1126/science.1230835
- Zhou, P., Tachedjian, M., Wynne, J. W., Boyd, V., Cui, J., Smith, I., et al. (2016). Contraction of the type I IFN locus and unusual constitutive expression of IFN- α in bats. *Proc. Natl. Acad. Sci. U. S. A.* 113, 2696–2701. doi: 10.1073/pnas.1518240113

Zhou, P., Yang, X. L., Wang, X. G., Hu, B., Zhang, L., Zhang, W., et al. (2020). A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 579, 270–273. doi: 10.1038/s41586-020-2012-7

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Giotis, Matthews and Smith. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.