THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Baseline epidemiology and associated dog ecology study towards stepwise elimination of rabies in Kwara state, Nigeria

Citation for published version:

Al-Mustapha, A, Abubakar, AT, Oyewo, M, Bamidele, FO, Ibrahim, A, Shuaib, MO, Olugasa, B, Balogun, MS, Kia, G, Mazeri, S & Heikinheimo, A 2021, 'Baseline epidemiology and associated dog ecology study towards stepwise elimination of rabies in Kwara state, Nigeria', *Preventive Veterinary Medicine*, vol. 189, 105295. <https://doi.org/10.1016/j.prevetmed.2021.105295>

Digital Object Identifier (DOI):

[10.1016/j.prevetmed.2021.105295](https://doi.org/10.1016/j.prevetmed.2021.105295)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Preventive Veterinary Medicine

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Baseline epidemiology and associated dog ecology study towards stepwise elimination of rabies in Kwara state, Nigeria

Ahmad I. Al-Mustapha, ^{a,b,c} * Abubakar Ahmed Tijani, ^{c,d} Oyewo Muftau, ^{c,d}, Folashade O. Bamidele ^{c,d}, Ahmed Ibrahim ^c, Muhammad Shuaib Osu, ^e Babasola Olugasa, ^f Muhammad Shakir Balogun, ^g Grace Kia, ^{h,i} Stella Mazeri, ^j Annamari Heikinheimo, ^a

^a Department of Food Hygiene and Environmental Health, Faculty of Veterinary Medicine, University of Helsinki, Finland.

^b Department of Veterinary Public Health and Preventive Medicine, Faculty of Veterinary Medicine, University of Ibadan, Oyo State, Nigeria.

^c Department of Veterinary Services, Kwara State Ministry of Agriculture and Rural Development, Ilorin, Nigeria.

^d Nigerian Field Epidemiology and Laboratory Training Program, Asokoro, Abuja-Nigeria.

^e Federal Department of Veterinary and Pest Control Services, Federal Ministry of Agriculture and Rural Development, Abuja -Nigeria.

^f Department of Veterinary Public Health and Preventive Medicine, Centre for Control and Prevention of Zoonoses (CCPZ), Faculty of Veterinary medicine, University of Ibadan, Nigeria.

^g African Field Epidemiology Network, Abuja – Nigeria.

^h Department of Veterinary Public Health and Preventive Medicine, Faculty of Veterinary Medicine, Ahmadu Bello University, Zaria-Kaduna State, Nigeria.

ⁱ War Against Rabies Foundation, Abuja- Nigeria.

^j Royal (Dick) School of Veterinary Studies and The Roslin Institute, The University of Edinburgh, Edinburgh – UK.

*Correspondence:

Al-Mustapha, Ahmad Ibrahim

a.i.al-mustapha@sms.ed.ac.uk

Abstract

Understanding domestic dog population dynamics and ecology is crucial to any effective rabies control program. This study was conducted as part of the baseline epidemiological studies necessary for the establishment of the Kwara Rabies Rapid Alert System “KRRAS”. This study aimed to determine the dog population structure of Kwara State by assessing the dog ownership, vaccination status, and prevalence of dog bites.

A total of 1,460 questionnaires were administered to respondents in the three senatorial zones of the state using Open Data Kit (ODK) between June 2019 to January 2020.

Of the 1460 households surveyed, 293 (20.1%) owned at least one dog with an average of 2.25 dogs per household. The male to female ratio was 1.9:1 and 79.3% (n=523/659) of the owned dogs were local breeds. A total of 785 dogs was enumerated (659 dogs from 293 households and 126 free-roaming dogs) and 7811 persons which resulted in a dog-human ratio of 1:9.95. The estimated dog population is 376,789 (95% CI: 343,700 – 379,878). Only 31% (n=204/659) of households vaccinated their dogs against rabies. The prevalence of dog-bite was 13% (n=193/1460) of which only 27% of the victims (n=61/225) received post-exposure prophylaxis (PEP). Dog ownership was significantly impacted by the ethnicity of respondents. Hausa (OR: 3.76; 95% CI: 1.15 - 12.4; $p < 0.001$) and Nupe (OR: 4.48; 95% CI: 1.77-11.33; $p < 0.001$) respondents owned dogs than Yoruba respondents. The rabies vaccination status of owned dogs was significantly impacted by the level of education (OR: 5.03; 95% CI: 1.50 - 16.83; $p < 0.001$); history of previous dog bite incidents (OR: 1.74; 95% CI: 0.95 - 3.17; $p < 0.001$); the breed of the dog with exotic dogs being more vaccinated (OR: 2.79; 95% 0.64 - 12.05; $p < 0.001$). Similarly, Male dogs (OR: 1.49, 95% 1.03 – 2.86; $p < 0.001$) and partially confined dogs (OR: 1.09, 95% 0.45 – 2.11, $p < 0.001$) were found to be vaccinated against rabies.

The results of the study showed low dog vaccination coverage, and high number of free roaming dogs. Hence, a threat to public health. The low dog vaccination coverage is below the 70-80% target recommended for herd immunity by the World Health Organization.

Keywords: Rabies; KRRAS; Vaccination; Baseline dog ecology; Nigeria

Background

Rabies is one of the oldest and most terrifying diseases known to man (Horton et al., 2015). It is a fatal viral disease caused by one of the seven lineages of *Lyssavirus* with a distinct “bullet” shape belonging to the family *Rhabdoviridae* (Badrane et al., 2001). Domestic dogs account for over 99% of human death from rabies (WHO, 2013).

It causes approximately 59,000 (95% CI: 25 - 159,000) human deaths annually mainly in Asia and Africa with 40% of people bitten by suspect rabid animals are children under 15 years of age (OIE, 2017a). Annually, it is responsible for 3.7 million (95% CI: 1.6 - 10.4 million) disability-adjusted life years (DALYs) and 8.6 billion USD (95% CI: 2.9 - 23.5 billion) economic losses (Hampson et al., 2015).

Rabies is endemic in Nigeria and has remained one of the most important neglected diseases of public health concern in the country. Despite the under-reporting of dog-bites incidents, about 10,000 cases of dog-bites are reported annually (NCDC, 2018). The rabies viral antigen has been detected in the brain tissues of apparently healthy dogs slaughtered for human consumption (Suleiman et al., 2020; Mohammed et al., 2019; Kia et al., 2018; Hambolu et al., 2014; Garba et al., 2010).

Several factors such as poor vaccination strategy, lack of sufficient vaccines, presence of stray (community) dogs and illegal trade of dogs within and across countries are major contributory factors to the endemicity and transboundary movement of rabies in Nigeria (Kia et al., 2018; Ogo et al., 2011; Ogunkoya et al., 2008). The epidemiology of rabies has been described in

Nigeria with dogs as the main reservoir of the disease (Kia et al., 2018; Ameh et al., 2014; Garba et al., 2011). The presence of unvaccinated free-roaming dogs (FRD) amidst human settlements is a major contributor to the high incidence and maintenance of rabies in Nigeria.

Mass dog vaccination is the most cost-effective strategy for preventing human rabies (OIE, 2017b). Parenteral vaccinations using the inactivated vaccine accompanied by the use of baits for oral vaccination of dogs (Maki et al., 2017) are effective in controlling canine rabies. Dog vaccination reduces human deaths attributable to rabies and the need for post-exposure prophylaxis (PEP) as a part of dog bite patient care (Cliquet et al., 2018).

In Nigeria, there is no national or regional rabies control program. Hence, the importance of projects such as the Kwara Rabies Rapid Alert System (KRRAS) cannot be over-emphasized. KRRAS is a one health integrated surveillance system designed to improve the reporting of dog-bite cases and enhance the diagnosis of rabies. The project is divided into three phases: Baseline epidemiological assessment studies; Mass dog vaccinations; and the Surveillance of dog-bite cases using the existing disease surveillance and notification officers (DSNOs) in the state. This study, as well as three others, are the epidemiological basis for the KRRAS.

This study aimed to be the baseline epidemiological assessment study necessary for the mass dog vaccinations; and an improved rabies surveillance system using the one-health approach. This ecological study is critical to vaccine procurement, prioritization of intervention plans, and essential for the enactment of the dog leash law in Kwara State.

Methods

Study area

Kwara State is located in the southern guinea savannah zone of Nigeria between latitude 8.9669° N, and longitude 4.3874° E. The state has a population of 3,599,800 (NPC, 2020). It

is located in Northcentral Nigeria. The state has three agro-ecological and geographical zones (Northern, Central, and Southern) with vast land and with varying climatic conditions.

Period and course of the survey

A cross-sectional survey was carried out to generate a baseline dog population, dog to human ratio, vaccination coverage for the anti-rabies vaccine, and the pentavalent dog vaccines (DHLPP) across the state. The study also assessed the incidence of dog bites in Kwara state. We developed a structured questionnaire which comprised of: a) owner demographics b) dog ownership and vaccination status and c) incidence of dog bites and its management. The questionnaire was pre-tested in a pilot study and adjusted accordingly before being used in this survey.

The required sample size was computed using Epi-Info V.7.0 (CDC, Atlanta, USA). At a 95% confidence interval (95% CI), we hypothesized that 50% of households will have dogs in each senatorial zone (384 households per zone, 1152 for the state). Furthermore, we added a 25% non-response rate to the total required households for the state (n=288). Therefore, the minimum number of households included in this survey was 1,460.

The survey was conducted from June 2019 to January 2020 in Kwara state./ A multi-stage sampling (Kwara state → 3 senatorial zones → local government areas → communities → areas → streets → households) of the respondents was carried out from all the communities in the state. Furthermore, a systematic random sampling of households was conducted to select streets, households, and respondents. A sampling interval of five households was used for this study.

Organization of survey team and movement plan

The Kwara Rabies Research Team (Kw-RRT), supported by a team of volunteers, composed of field epidemiologists, Unilorin One Health students, and data collectors carried out the survey. They were trained on the methodology of the survey and were divided into five groups

of two members, plus a group leader. They were randomly assigned separate predetermined routes. The survey was conducted on each selected street. Due to the lack of organized road connections in semi-urban and rural areas, the polio-vaccination micro-plan was adopted, using the polio house markings as a guide. Starting from the first household on the right side, every 6th household was selected and an adult member (18 years and above) was interviewed. The survey team explained the purpose of the survey to each respondent and informed consent was sought before the administration of the questionnaire. The questionnaire was administered in English and in local language of the community when needed. The answers were recorded using the open data kit (ODK) application in mobile phone and uploaded the data to the African Field Epidemiology Network (AFENET) server.

Free Roaming Dog Enumeration Technique

We used Beck's definition of a free-roaming dog (FRD) as “Any dog observed without human supervision on public property or private property with immediate unrestrained access to public property” (Berman and Dunbar, 1983). Using the photographic recapture technique (Beck, 1973); we used photography to prevent counting a dog twice within the same area. Surveys were alternated between mornings and late evenings on two alternate days. On each day of the study, counting of dogs was carried out in the morning (between 6 and 9 a.m.) and in the evening (5.30 to 7.00 p.m.). Every dog sighted within a 5m radius of the road was photographed and recorded.

The free-roaming dog population in the three senatorial zone was estimated using the Beck (1973) formula as previously described by Mshelbwala et al (2017).

$$N = \frac{\sum(Mn)}{\sum m}$$

where N= the estimate of the dog population of Kwara state.

M= the number of dogs observed and photographed each time in the street and considered as "sighted and marked".

m= the number of dogs identified as being previously photographed, that is "re-sighted/re-captured"

n= Each days' observation (M) – previously observed dogs (m) (Table 2).

Data analysis

The household survey data were analyzed using Minitab 19.1.1 (Pennsylvania, USA). Qualitative data were presented as proportions whereas quantitative data were presented as mean and standard deviation. A univariable logistic regression analysis was conducted to assess the association between the socio-demographic variables and the outcome variables [dog ownership of the household (yes vs no), and the vaccination status of the dog against rabies (vaccinated vs not vaccinated)]. Variables with p-value ≤ 0.05 were selected for multivariable logistic regression analysis using the logit function and α at 0.25. A multivariable logistic regression analysis was performed and the variables with p-value ≤ 0.05 were retained in the final model. The odds ratio (OR) and its 95% confidence interval (CI) of the variables associated with the outcome variables were derived from the final multivariable logistic regression model.

RESULTS

Demography of the respondents

The questionnaire was administered in English and translated to the local language of the respondents (where necessary). Most of the respondents (72%, n=1045/1460) had either secondary or tertiary education. Tradesmen/Artisans accounted for over 47% of the respondents. The demographic characteristics of the respondents is shown in Table 1.

183 **Table 1.** The demographic structure of respondents in Kwara state (n=1460).

Variables	No of respondents (%)
Senatorial Zone	
Kwara South	364 (25)
Kwara North	562 (38.5)
Kwara Central	534 (36.5)
Gender	
Female	382 (26.1)
Male	1072 (73.5)
Prefer not to say	6 (0.4)
Religion	
Christianity	523 (35.8)
Islam	914 (62.6)
Others	3 (0.2)
Prefer not to say	21(1.4)
Level of education	
No western education	187 (13.5)
Primary	165 (11.9)
Secondary	581 (41.9)
Tertiary	424 (30.6)
Prefer not to say	73 (5)
Other	30 (2.1)
Occupation	
Artisan	286 (20.4)
Civil servant	128 (9.1)
Driver	20 (1.4)
Farmer	169 (12)
Fisherman	3 (0.2)
Laborer	12 (0.9)
Other professions	64 (4.6)
Security officer	17 (1.2)
Student	253 (18)
Trader/ businessman	376 (26.9)
Unemployed	72 (5.2)
Prefer not to say	60 (4.1)

184

185 ***Dog Ecology***

186 A street survey estimated a total of 126 free-roaming dogs (FRDs) across the three
 187 senatorial zones (Table 2). A household questionnaire survey found a total of 659 dogs
 188 owned by 293 respondents resulting in an average of 2.25 dogs per visited household.
 189 During the survey, 7811 people were counted and a total of 785 dogs (FRD and owned)

giving a dog to human ratio of 1:9.95. Therefore, with a total population of 3,599,800; we estimated a total dog population of 361,789 (95% CI: 343,700 – 379,878) in Kwara state. The descriptive statistics of owned dogs showed that the local breeds of dogs (Mongrel) accounted for 79.3% of owned dogs. Most respondents (52%, n=198/380) kept dogs for security whereas hunting dogs accounted for another 28% (n=105/380) of owned dogs. Based on the use of the dog, 69% (n=202/302) partially/ never confine their dogs. Most of the owned dogs (56%, n=369/659) were aged between 1-3 years (Table 3).

Table 2. Estimation of free-roaming dogs using photographic mark-recapture methods in Kwara state.

Senatorial zone	Mean number of dogs sighted and photographed per area (M)	No of dogs sighted that were previously photographed (m)	n = (M-m)	N
Kwara South	17	5	11	37.4
Kwara North	11	3	8	29
Kwara Central	30	10	20	60
Total	58	18	39	126.4

Each parameter is an average of 7 chosen areas from each senatorial zone.

Dog vaccination status

Of the 293 households surveyed, only 31% (n=204/659) were previously vaccinated against rabies. Among the dogs vaccinated against rabies, only 22% (n=45/204) had received DHLPP® (a recommended pentavalent vaccine for puppies at 8 weeks of age meant to provide immunity against Canine distemper, Hepatitis, Leptospirosis, Parainfluenza, and Parvovirus). With 52% (n=48/92) of all vaccinations administered at home (Table 3).

211 **Table 3.** Characteristics of owned dog population and vaccination status in Kwara
212 state.

Variables	Frequency (%)
Do you own dogs? (n=1460)	
Yes	293 (20.1)
No	1167 (79.9)
Gender of dog ^b	
Female	231 (35)
Male	428 (65)
Breed of dog ^b	
Cross-bred	61 (9.3)
Exotic	75 (11.4)
Local	523 (79.3)
Age of dogs ^b	
< 1 year	119 (18)
1-3 years	369 (56)
3-5 years	119 (18)
> 5 years	52 (8)
Use of dog (s) ^a	
Breeding	31 (8)
Herding	5 (1)
Hunting	105 (28)
Pet	41 (11)
Security	198 (52)
Confinement of dog ^a	
Always	80 (27)
Never	152 (52)
Partial	50 (17)
Vaccination against rabies ^b	
Yes	204 (31)
No	455 (69)
Vaccination against DHLPP [®] ^b	
Yes	145 (22)
No	514 (78)
Where vaccination against rabies was received ^a	
At home	152 (52)
Private Veterinary Clinic	88 (30)
Gov't Vet clinic	53 (18)
Pet green book ^a	
Yes	171 (26)
No	488 (74)
Age of first vaccination ^b	
3-6 months	99 (15)
6-12 months	86 (13)
1-2 years	172 (26)
2-3 years	92 (14)
3-4 years	46 (7)

213	4-5 years	86 (13)
214	5 years +	79 (12)
215	Dog population management methods	
216	Give away	36 (55)
217	I Keep my pups	10 (15)
218	Killed	4 (6)
219	Sold	15 (23)

221 a Parameters computed based on the number of households that own dogs (n=293)

222 b Parameters computed based on the number of owned dogs (n=659).

223

224 *Incidence and management of dog bites in Kwara state*

225 Of the 1460 respondents, 13% (n=193/1460) had a history of dog-bite with (63%, n=154/246)

226 being beaten once and (23%, n=56/246) were bitten twice (Table 4). About 13% (n=27/225)

227 and 32% (n=72/225) used antibiotics and traditional methods (herbs) to treat dog-bite wounds

228 (Table 4) However, only 27% of the dog bite victims received the post-exposure prophylaxis

229 (PEP) and rabies immunoglobulin (RIG).

230

231 **Table 4.** Evaluation of dog-bite incidence and it's management (n=193).

Variable	Frequency (%)
Ever bitten by a dog?	
No	1267 (87)
Yes	193 (13)
No of bites incidents	
1	154 (63)
2	56 (23)
3	21 (9)
5	15 (6)
Outcome of dog	
Alive	72 (36)
Died	11 (6)
I don't know	12 (6)
Killed	59 (30)
Ran away	46 (23)
Treatment given to the dog-bite victim	
Antibiotics	27 (13)
Non-specific	64 (28)
PEP and RIG	61 (27)
Traditional	73 (32)

232

233

234 *Analysis of factors affecting dog ownership and vaccination status in Kwara State.*

The multivariable regression analysis demonstrated that the dog ownership in Kwara state was found to be associated with the occupation and the ethnicity of the household respondents. The Hausa (OR: 3.76, 95% CI: 1.15-12.35) and Nupe (OR: 4.48, 95% CI: 1.77 - 11.33) ethnic groups are more likely to own dogs when compared with other tribes. Similarly, comparing to Artisan, other types of occupation were likely to be associated with owning the dogs (Table 5).

The rabies vaccination status of was found to be associated with the gender and breed of the dog, dog management, education level of the owner and the dog bite incident history of the owner. Male dogs (OR: 1.49, 95% 1.03 – 2.86), exotic breed (OR: 2.79: 95% 0.64 - 12.05) and partially confined dogs (OR: 1.09, 95% 0.45 – 2.11) are found to be vaccinated against rabies. Similarly, an owner that have a history of previous dog bite incidents (OR: 1.74, 95% 0.95 - 3.17) and those owners that have attended school education are more likely to have vaccinated their dogs against rabies (Table 5).

249 Table 5. Univariable and multivariable logistic regression analysis of the factors associated with the “ownership” and “vaccination status against
250 rabies” in dogs in Kwara State, Nigeria

Variable				Referent		Univariable analysis				Multivariable analysis			
Dog ownership				Odds Ratio (95% CI)	X^2	DF	p -value	Adjusted Ratio (95% CI)	X^2	DF	p -value		
	Religion	Islam	Christianity	1.05 (0.80 - 1.37)	0.13	1	0.937	-	-	-	-		
	Occupation	Artisan	Civil	1.49 (0.91 - 2.42)	61.17	10	0.001	1.49 (0.89 – 2.29)	60.91	10	0.0001		
			Servant										
			Driver		2.21 (0.84 - 5.80)				2.14 (0.71 – 4.26)				
			Farmer		2.63 (1.72 - 4.03)				3.41 (1.89 – 5.03)				
			Fisherman		0.00				0.00				
			Laborer		1.37 (0.36 - 5.22)				1.09 (0.29 – 4.62)				
			Security officer		1.49 (0.80 - 2.79)				1.1 (0.69 – 3.11)				
			Student		2.24 (0.79 - 6.32)				2.14 (0.78 – 5.99)				
			Trader/		0.91 (0.59 - 1.40)				1.1 (0.67 – 1.69)				
			Businessman										
			Unemployed	0.90 (0.46 - 1.76)				0.78 (0.30 – 1.51)					
	Ethnicity	Yoruba	Hausa	4.37 (1.34 - 14.14)	69.67	6	0.0001	3.76 (1.15-12.35)	69.30	6	0.0001		
			Nupe		2.26 (0.73 - 7.00)				4.48 (1.77 - 11.33)				
			Bokobaru		0.75 (0.53 - 1.04)				0.65 (0.44 - 0.96)				
			Fulani		0.22 (0.08 - 0.55)				0.04 (0.01 - 0.15)				
			Baruba		0.31 (0.21 - 0.47)				0.28 (0.17 - 0.44)				
			Other		4.99 (2.00 - 12.47)				2.37 (1.16 - 4.89)				
Vaccination status of dogs													
	Gender of dog	Female	Male	1.7 (0.97 - 2.98)	3.45	1	0.063	1.49 (1.03 – 2.86)	2.89	1	0.001		
	Confinement of dogs	Always	Never	0.48 (0.27 - 0.86)	7.94	3	0.047	0.46 (0.25 – 0.84)	6.48	3	0.002		
			Partial		1.00 (0.49- 2.06)				1.09 (0.45 – 2.11)				
			I don't know		0.56 (0.14 - 2.29)				0.50 (0.22 – 2.14)				

	Breed	Cross-bred	Exotic Local	2.79 (0.72 - 10.86) 0.33 (0.15 - 0.75)	18.42	2	0.0001	2.79 (0.64 - 12.05) 0.34 (0.13 - 0.85)	15.1	2	0.001
	Previous dog-bite incident	No	Yes	1.43 (0.83 - 2.46)	1.64	1	0.2	1.74 (0.95 - 3.17)	3.25	1	0.071
	Use of dog	Breeding	Herding Hunting Pet Security	4.33 (0.42 - 44.43) 1.96 (0.50 - 7.69) 2.29 (0.52 - 10.21) 2.01 (0.55 - 7.33)	1.88	4	0.757	- - - -	-	-	-
	Level of education	No formal education	Primary Secondary Tertiary Others	5.03(1.50 - 16.83) 3.24 (1.08 - 9.77) 7.81 (2.50 - 24.32) 8.5 (0.44 - 163.88)	16.55	4	0.002	5.94 (1.72 - 20.42) 3.04 (0.97 - 9.49) 6.99 (2.13 - 22.92) 6.15 (0.27 - 137.47)	14.4	3	0.006

251 %= Percentage; χ^2 = chi-square; DF- Degree of freedom; OR – Odds ratio; 95% CI – 95% confidence interval.

252

DISCUSSION

Dogs are responsible for 99% of all cases of human rabies (WHO, 2020). Therefore, understanding the domestic dog ecology and its population structure is the first step to an effective elimination program (Cleveland et al., 2006). Baseline epidemiological studies in rabies elimination cannot be overemphasized especially in vaccine procurement for mass vaccination campaigns. More so, they form the evidence-base for the Kwara Rabies Rapid Alert System -KRRAS-; a collaborative state-based project designed to achieve the global aim of eradicating dog-mediated human rabies by 2030 in Kwara state. KRRAS was designed using the five pillars of the global framework; Socio-cultural, Technical, Organization, Political, and Resources (STOP-R) in Kwara state. The total dog population is lower than reported for many other states in Nigeria. This makes Kwara State a good candidate for the first rabies elimination project in the country.

Dog ownership was not affected by religion ($p = 0.937$). This is in contrast to findings by Mauti et al (2017); Oboegbulem and Nwakonobi (1989) which showed significantly higher dog ownership amongst Christians. Most of the owned dogs were local breeds. This aligns with reports from Kwaghe et al., (2019) and Grace et al., (2018).

Male dogs were more abundant than female dogs in Kwara State (sex ratio of dogs 1.9:1). It is believed that male dogs are better guardians and hunters than female dogs (Kitala et al., 2001). This is as previously described by Otolorin et al (2014); Aiyedun and Olugasa, (2012). However, this does not agree with the findings of Kwaghe et al., (2019); Hambolu et al (2014) who reported higher female to male dog ratio. Dogs were mostly kept for security purposes. This is similar to reports from other parts of Nigeria and Africa (Kwaghe et al., 2019; Garba et al., 2017; Mauti et al., 2017). The median age of owned dogs were 1-3 years was similar to reports from all over the country that shows the average dog were older than 1 year (Otolorin et al., 2014; Hambolu et al., 2013). Also, Aiyedun and Olugasa (2012) reported 71% of dogs aged > 6 months old.

The dog to human ratio of approximately 1:10 is lower than 1: 7.8 reported in Abia state (Otolorin et al., 2014). Similarly, Kwaghe et al., (2019) reported a higher dog-human ratio of 1: 6.6. Other studies like those of Hambolu et al., (2014); Atuman et. al., (2014); and Garba et al., (2017) reported a dog-human ratio of 1:5.6; 1:4.1 and 1:5.4 respectively (Table S1). Aiyedun and Olugasa (2012) reported a dog-human ratio of 1:139 in Ilorin; which is lower than what this study has recorded in Kwara state. This might be because Aiyedun and Olugasa covered the Ilorin metropolis. The dog to human ratio reported in this study is within the range reported and modeled for most African countries (Knobel et al., 2005). Within Africa, several studies conducted reported a dog-human ratio of 1:14 in Tanzania (Gsell et al., 2012); 1:21.5 in Chad (Mindekem et al., 2005); 1:15 in Kenya (Kitala et al., 2001) and 1: 16 in Zimbabwe (Brooks R., 1990). Kwara State has a total landmass of 36,825 km², giving a total of 7.5 dogs/ km². This is within the range of 6 and 21 dogs/ km² that was reported by Kitala et al (2001) in Kenya (Table S1). Much higher dog densities (>1000 dogs/ km²) had been estimated for Lagos (Hambolu et al., 2014) and some South American countries (Davlin and Vonville, 2012).

The variations in the dog to human ratio in different areas of Nigeria might be attributable to the different socio-cultural, religious, and economic status of various states of the federation. The rabies vaccination rate is similar to those earlier reported in other parts of the country. This is much lower than the 69.6%; 64.1% and 49.5% reported in Niger (Garba et al., 2017), Lagos (Hambolu et al., 2014) and Abuja (Mshelbwala et al., 2017) respectively. It is, however, higher than the 26.4% and 21% reported in Bauchi (Atuman et al., 2014) and Nasarawa (Kwaghe et al., 2019) respectively. This vaccination coverage falls significantly below the 70-80% vaccination rate needed to boost herd immunity (WHO, 2003). With low vaccination coverage, the general public is at risk of rabies exposure and the need for improved public awareness on rabies, first aid for dog-bite victims, and availability of PEP especially in rural areas cannot be over-emphasized.

Only 26% of the vaccinated dogs had a vaccination certificate. This is because only licensed veterinarians are allowed to issue a signed vaccination certificate and most vaccinations were administered at home (possibly- by para veterinary technicians). This has impaired proper monitoring and surveillance of dog health and welfare.

The prevalence of dog-bite and its management was evaluated as an important component of rabies epidemiology. The dog-bite incidence was lower than the 31% and 26.4% reported in Niger and Bauchi states respectively (Garba et al., 2017; Atuman et al., 2014).

There is an urgent need for public education on rabies (mission of KRRAS) as only 15% of the respondents vaccinated their dogs at the appropriate age of 3-4 months. More so, only 27% of dog-bite victims received the PEP and RIG injections from a health facility. This is especially worrying as most of the respondents (73%) treated dog-bites with antibiotics (13%), non-specific treatment (28%), and traditional (32%) concoctions. Non-specific treatment included wound cleaning and the use of antibiotics. The high level of stray/free-roaming dogs seen in Kwara state is due to the lack of dog population control programs. Hence, the need to re-introduce effective stray dog population control described under the OIE terrestrial animal health code (OIE, 2019) during our mass vaccination campaigns. This un-controlled breeding coupled with their use for hunting has introduced and maintained several genera of lyssaviruses in the environment by introducing the sylvatic (wild) rabies cycle into the urban cycle. This could potentially increase the transmission intensity and spread of rabies.

Conclusion

The results of the study showed low canine vaccination coverage. This poses a serious public health threat. It is essential to control the population of free-roaming dogs. Furthermore, public awareness needs to be raised among certain subsets of the population with special emphasis on proper dog-bite treatment regimen and availability of pre-exposure and post-exposure prophylaxis for humans.

This baseline epidemiology and ecology study forms the evidence base that can be used for implementing an effective dog Mass Vaccination Campaign in the state. Information from this study is vital for vaccine procurement, proportionate distribution within the state, and serves as a basis for a valid comparison for our post-vaccination surveys. We recommend the establishment of a rabies desk office (RDO) and include dog-bite amongst the reportable diseases by the disease surveillance and notification officers (DSNOs) to the District Health Information System (DHIS2). We equally recommend further longitudinal studies to define the health and welfare challenges of dogs in Nigeria

Declarations

- Ethics approval and consent to participate

Ethical clearance was obtained from the various ethical review boards of the Kwara State: Ministry of Health (MoH), Agriculture and Rural Development (MoARD), and Education and Human Capital Development (MoEHCD), Ilorin - Nigeria (reference number: MOH/KS/EU/777/31). Informed consent was sought from the respondents and participants could opt-out at any time.

- Consent for publication

Not applicable

- Availability of data and materials

The survey instrument and datasets are available as supplementary data.

- Competing interests

The authors declare that they have no competing interests.

- Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

- Authors' contributions

AIA, AAT, FB, MO, MSO, MSB, GK, and AI were involved in planning the study and data collection. AIA drafted the manuscript. BO and AH did the overall review of the manuscript. All authors read and approved the final study.

- Acknowledgments

We are grateful to the African Field Epidemiology Network (AFENET) for the provision of field data collectors and their server for data upload. We acknowledge the dedication of Jimoh Mohammad, Abdulrahim Ibrahim, and Idris Olarewaju in data collection.

Supplementary data

Supplementary file 1- Survey instrument

Supplementary file 2

Table S1. Comparison of dog ecology across Nigeria and Africa

References

- Aiyedun, J., & Olugasa, B., 2012. Use of aerial photograph to enhance dog population census in Ilorin, Nigeria. *Sokoto Journal of Veterinary Sciences*, 10(1). <http://doi.10.4314/sokjvs.v10i1.5>
- Ameh, V., Dzikwi, A., & Umoh, J., 2014. Assessment of Knowledge, Attitude and Practice of Dog Owners to Canine Rabies in Wukari Metropolis, Taraba State Nigeria. *Global Journal of Health Science*, 6(5). <http://doi.10.5539/gjhs.v6n5p226>
- Atuman, Y., Ogunkoya, A., Adawa, D., Nok, A., & Biallah, M., 2014. Dog ecology, dog bites and rabies vaccination rates in Bauchi State, Nigeria. *International Journal of Veterinary Science and Medicine*, 2(1), 41-5. <http://doi.10.1016/j.ijvsm.2014.04.001>
- Badrane, H., Bahloul, C., Perrin, P., & Tordo, N., 2001. Evidence of two lyssavirus phylogroups with distinct pathogenicity and immunogenicity. *J Virol*, 75, 3268-76.
- Beck, A. M. (1973). *The ecology of stray dogs: a study of free-ranging urban animals*. Indiana, United States: Purdue University Press Books 3.
- Brooks, R., 1990. Survey of the dog population of Zimbabwe and its level of rabies vaccination. *Veterinary Records*, 127(24):592- 6.
- Cleveland, S., Kaare, M., Knobel, D., & Laurenson, M. K., 2006. Canine vaccination providing broader benefits for disease control. *Vet Microbiol*, 117, 43-50. <http://doi.10.1016/j.vetmic.2006.04.009>
- Cliquet, F., Guiot, A., Aubert, M., Robardet, E., Rupprecht, C., & Meslin, F., 2018. Oral vaccination of dogs: a well-studied and undervalued tool for achieving human and dog rabies elimination. *Veterinary Research*, 49(1). <http://doi.10.1186/s13567-018-0554-6>
- Coleman PG, Fevre EM, Cleaveland S., 2004. Estimating the public health impact of rabies. *Emerging infectious diseases*, 10(1):140–2.
- Davlin, S., & VonVille, H., 2012. Canine rabies vaccination and domestic dog population characteristics in the developing world: A systematic review. *Vaccine*, 30(24), 3492-3502. <http://doi.10.1016/j.vaccine.2012.03.069>.
- Garba, A., Oboegbulem, S. I., Junaidu, A. U., Magaji, A. A., Umoh, J. U., Ahmed, A., Masdooq A.A., 2010. Rabies virus antigen in the brains of apparently healthy dogs in Sokoto and Katsina States, Nigeria. *Nig J Parasitol*, 31, 123-5.
- Garba, A., Dzikwi, A., Kazeem, H., Makanju, O., Hambagba, F., & Abduazeez, N., 2017. Dog Ecology and Management in Niger State, Nigeria: A Basic Tool for Rabies Control. *Journal of Agriculture And Ecology Research International*, 12(1), 1-9. <http://doi.10.9734/jaeri/2017/32442>
- Grace S. N. Kia, Umoh, J.U. Kwaga, J.K.P., Zhen F. Fu et al., 2018. Molecular Characterization of a Rabies Virus Isolated from Trade Dogs in Plateau State, Nigeria. *Sokoto Journal of Veterinary Sciences* 16(2):54-62.
- Gsell AS, Knobel DL, Kazwala RR, Vounatsou P, Zinsstag J., 2012. Domestic dog demographic structure and dynamics relevant to rabies control planning in urban areas in Africa: the case of Iringa, Tanzania. *BMC Veterinary Research*. 8:236.
- Hambolu, S., Dzikwi, A., Kwaga, J., Kazeem, H., Umoh, J., & Hambolu, D., 2013. Rabies and Dog Bites Cases in Lagos State Nigeria: A Prevalence and Retrospective Studies (2006-2011). *Global Journal of Health Science*, 6(1).
- Hambolu, S., Dzikwi, A., Kwaga, J., Kazeem, H., Umoh, J., & Hambolu, D., 2014. Dog Ecology and Population Studies in Lagos State, Nigeria. *Global Journal of Health Science*, 6(2).

435 Hampson, K., Coudeville, L., Lembo, T., Sambo, M., Kieffer, A., & Attlan, M. et al., 2015.
 436 Estimating the Global Burden of Endemic Canine Rabies. *PLOS Neglected Tropical*
 437 *Diseases*, 9(4), e0003709. <http://doi.10.1371/journal.pntd.0003709>

438 Horton, D., McElhinney, L., Freuling, C., Marston, D., Banyard, A., & Goharriz, H. et al.
 439 2015. Complex Epidemiology of a Zoonotic Disease in a Culturally Diverse Region:
 440 Phylogeography of Rabies Virus in the Middle East. *PLOS Neglected Tropical*
 441 *Diseases*, 9(3), e0003569. <http://doi.10.1371/journal.pntd.0003569>
 442 <http://doi.10.1128/JVI.75.7.3268-3276.2001>

443 Kitala P, McDermott J, Kyule M, Gathuma J., 2001. Dog ecology and demography information
 444 to support the planning of rabies control in Machakos District, Kenya. *Acta Tropica*.
 445 78(3):217-230.

446 Knobel DL, Cleaveland S, Coleman PG, Fèvre EM, Meltzer MI, Miranda ME, Shaw A,
 447 Zinsstag J, Meslin FX., 2005. Re-evaluating the burden of rabies in Africa and Asia. *Bull*
 448 *World Health Organ* 83: 360 – 368.

449 Kwaghe, A., Okomah, D., Okoli, I., Kachalla, M., Aligana, M., Alabi, O., & Mshelbwala, G.,
 450 2019. Estimation of dog population in Nasarawa state Nigeria: a pilot study. *Pan African*
 451 *Medical Journal*, 34. <http://doi.10.11604/pamj.2019.34.25.16755>

452 Maki, J., Guiot, A., Aubert, M., Brochier, B., Cliquet, F., & Hanlon, C. et al., 2017. Oral
 453 vaccination of wildlife using a vaccinia–rabies-glycoprotein recombinant virus vaccine
 454 (RABORAL V-RG®): a global review. *Veterinary Research*, 48(1).
 455 <http://http://doi.org/10.1186/s13567-017-0459-9>

456 Mauti, S., Traoré, A., Sery, A., Bryssinckx, W., Hattendorf, J., & Zinsstag, J., 2017. First study
 457 on domestic dog ecology, demographic structure and dynamics in Bamako,
 458 Mali. *Preventive Veterinary Medicine*, 146, 44-51. <http://doi.10.1016/j.prevetmed.2017.07.009>

460 Mindekem R, Kayali U, Yemadji N, Ndoutamia AG, Zinsstag J., 2005. Impact of canine
 461 demography on rabies transmission in N'djamena, Chad. *Médecine Tropicale*. 65(1):53-
 462 58.

463 Mohammed, S., Umoh, J.U., Kia, G.S.N. 2019. Prevalence of rabies antigen in brain tissue of
 464 slaughtered dogs and public health factors associated with dog processing in Billiri,
 465 Gombe State, Nigeria. *Savannah Veterinary Journal*, 29-35. doi: 10.36759/svj.2019.050

466 Mshelbwala, P., Akinwolemiwa, D., Maikai, B., Otolurin, R., Maurice, N., & Weese, J., 2017.
 467 Dog ecology and its implications for rabies control in Gwagwalada, Federal Capital
 468 Territory, Abuja, Nigeria. *Zoonoses And Public Health*, 65(1), 168-176. <http://doi.10.1111/zph.12385>

470 National Population Commission. 2020. <http://population.city/nigeria/adm/kwara/> (accessed
 471 25 February 2020).

472 Nigerian Center for Disease Control (NCDC). 2018. <https://ncdc.gov.ng/diseases/factsheet/41>
 473 (accessed 13 February 2020)

474 Oboegbulem, S. I. and Nwakonobi, I. E., 1989. Population density and ecology of dogs in
 475 Nigeria: a pilot study. *Revue Scientifique et Technique de Office International des*
 476 *Epizooties*, vol. 8, pp. 733– 745.

477 Ogo, M.F.; Nel, L.; Sabeta, C.T., 2011. Phylogenetic Evidence of the Public and Veterinary
 478 Health Threat of Dog Rabies in Nigeria. *Niger. Vet. J.* 32, 40–44.

479 Ogunkoya, A.B., 2008. Review of rabies and problems of rabies in Nigeria. In *Proceedings of*
 480 *the National Conference/Work on Rabies*; Ahmadu Bello University: Zaria, Nigeria, pp.
 481 62–70.

- Otolorin, G., Umoh, J., & Dzikwi, A., 2014. Demographic and Ecological Survey of Dog Population in Aba, Abia State, Nigeria. *ISRN Veterinary Science*, 2014, 1-5. <http://doi.10.1155/2014/806849>
- Suleiman, M., Kwaga, J., Okubanjo, O., Abarshi, M., & Kia, G., 2020. Molecular study of rabies virus in slaughtered dogs in Billiri and Kaltungo local government areas of Gombe state, Nigeria. *Acta Tropica*, 207, 105461. doi: 10.1016/j.actatropica.2020.105461
- World Health Organization (WHO). 2013. Expert Consultation on Rabies. Second Report. Geneva.
- World Organization for Animal Health (OIE). 2017a. Rabies portal. <http://www.oie.int/animal-health-in-the-world/rabies-portal/> (accessed 13 February 2020)
- World Organization for Animal Health (OIE). 2017b. Report of the meeting of the OIE biologicals standards commission. World Organization for Animal Health, Paris
- World Organization for Animal Health (OIE). 2019. Stray Dog Population Control. <https://www.oie.int/doc/ged/D9926.PDF> (accessed 25 February 2020).

502 **Supplementary data**

503 **Table S1** Comparison of dog ecology across Nigeria and Africa

Location	Dog-human ratio	Dog density/km ²	Dog owning households (%)	Dogs/household	Average dog age	Male-Female ratio	Vaccination coverage (%)	References
Kwara State (NC)	1:10	7.5	20	2.25	1-3 years	1.9:1	31	This study
Bauchi (NE)	1:4.1	-	-	2.3	1-5 years	1:1.2	26.4	Atuman et al., 2014
Lagos (SW)	1:5.6	-	95	2.8	> 1 year	1:1.5	64.1	Hambolu et al., 2014
Abia (SE)	1:7.8	-	-	1.5	> 1 year	1.6:1	39.9	Otolorin et al., 2014
Bamako, Mali	1:121	56	9	0.13	3.2 years	2.8:1	-	Mauti et al., 2017

504

505 NC- North-central, NE- North-eastern; SW- South-western; SE-South-eastern Nigeria; % -percentage.

506

507