THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Developmental expression patterns of chemokines CXCL11, CXCL12 and their receptor CXCR7 in testes of common marmoset and human testes

Citation for published version:

Westernströer, B, Langenstroth, D, Kliesch, S, Troppmann, B, Redmann, K, Macdonald, J, Mitchell, R, Wistuba, J, Schlatt, S & Neuhaus, N 2015, 'Developmental expression patterns of chemokines CXCL11, CXCL12 and their receptor CXCR7 in testes of common marmoset and human testes', *Cell and tissue research*. <https://doi.org/10.1007/s00441-015-2164-1>

Digital Object Identifier (DOI):

[10.1007/s00441-015-2164-1](https://doi.org/10.1007/s00441-015-2164-1)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Cell and tissue research

Publisher Rights Statement:

This is the author's final and accepted manuscript.

The final publication is available at Springer via <http://link.springer.com/article/10.1007%2Fs00441-015-2164-1>

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Developmental expression patterns of the chemokines CXCL11, CXCL12 and its receptor CXCR7 in common marmoset and human testes

Journal:	<i>Cell and Tissue Research</i>
Manuscript ID:	CTR-14-0494.R2
Manuscript Type:	Regular Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Westernströer, Birgit; University of Münster, Centre of Reproductive Medicine and Andrology Langenstroth, Daniel; University of Münster, Centre of Reproductive Medicine and Andrology Kliesch, Sabine; University Clinic Muenster, Department of Clinical Andrology Troppmann, Britta; University of Münster, Centre of Reproductive Medicine and Andrology Redmann, Klaus; University of Münster, Centre of Reproductive Medicine and Andrology Macdonald, Joni; MRC/University of Edinburgh, Centre for Reproductive Health Mitchell, Rod; MRC/University of Edinburgh, Centre for Reproductive Health Wistuba, Joachim; University of Münster, Centre of Reproductive Medicine and Andrology Schlatt, Stefan; Univ Münster, CeRA Neuhaus, Nina; University of Münster, Centre of Reproductive Medicine and Andrology
Keywords:	CXCR7, CXCL12, CXCL11, primate testes, testicular development

University Hospital of Münster. Centre of Reproductive Medicine and Andrology. 48129 Münster

**To the Editor-in-Chief of
Cell and Tissue Research**

(via electronic submission)

Centre of Reproductive Medicine and Andrology

Univ.-Prof. Dr. rer. nat. Stefan Schlatt
Director

Mailing address:
Albert-Schweitzer-Campus 1, Gebäude D11
D48149 Münster
Street address:
Domagkstraße 11
D 48149 Münster
Direct dialing: +49 (0)2 51 - 83 - 56099
Fax: +49 (0)2 51 - 83 - 54800
Operator: +49 (0)2 51 - 83 - 0
stefan.schlatt@ukmuenster.de
www.ukmuenster.de

Münster, 21.02.2015

Resubmission of the revised manuscript (CTR-14-0494.R1) entitled 'Developmental expression patterns of the chemokines CXCL11, CXCL12 and its receptor CXCR7 in common marmoset and human testes'.

Dear Prof. Unsicker, dear Prof. Sutovsky,

thank you for the second review of our manuscript. We are delighted to see that the paper is close to acceptance. We revised the manuscript in accordance to the suggestions by referee 1. We herewith submit a slightly revised version with some changes to the discussion.

We are grateful that you consider our work for publication in Cell and Tissue Research.

Yours sincerely,

Prof. Dr. Stefan Schlatt

Response to Reviewer

Reviewer: 1

I re-examined the revised version of the manuscript. The authors have addressed all of my concerns and the manuscript has been modified accordingly.

Response: We thank the reviewer for this positive assessment.

1) However, I have noted that the same group has recently published a paper on the profiling of CXCR4 and CXCR7 during murine testis development (Westernstroer et al. 2014). The expression pattern of CXCR7 in the murine germ cells as well as the expression profile of CXCL12 (assessed by qRT-PCR) is not perfectly in line with data obtained on monkey and human testis. CXCR7 expression is retained in more differentiated murine germ cells, and murine CXCL12 is stable up to 21 postnatal day and increases after 37 postnatal days. In my opinion, the authors should discuss and comment on this in more details (page 21, lane 354).

Response (lines 322 – 333, 362 – 367): We expanded the discussion as suggested.

Title: Developmental expression patterns of the chemokines CXCL11, CXCL12 and its receptor CXCR7 in common marmoset and human testes

Authors: Birgit Westernströer¹, Daniel Langenstroth¹, Sabine Kliesch², Britta Troppmann¹, Klaus Redmann¹, Joni Macdonald³, Rod Mitchell³, Joachim Wistuba¹, Stefan Schlatt¹, Nina Neuhaus¹

¹ Centre of Reproductive Medicine and Andrology, CeRA, Albert-Schweizer-Campus 1, Building D11, 48149 Münster, Germany

² Department of Clinical Andrology, Centre of Reproductive Medicine and Andrology, CeRA, Albert-Schweitzer-Campus 1, Building D11, 48149 Münster, Germany

³ MRC/University of Edinburgh Centre for Reproductive Health, The Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh, EH164TJ, Scotland, UK

Corresponding author: Prof. Dr. Stefan Schlatt
Centre of Reproductive Medicine and Andrology
Albert-Schweizer-Campus 1 (D11)
48149 Muenster, Germany
Email: Stefan.Schlatt@ukmuenster.de

Key words: CXCR7, CXCL12, CXCL11, primate testes, testicular development

Abstract

The chemokine receptor CXCR7 interacts with the chemokines CXCL11 and CXCL12. During development this ligand receptor system (C-X-C) provokes cell type-specific responses in terms of migration, adhesion or ligand sequestration. It is active in zebrafish and rodents but no data are available for its presence or function in primate testes. Real-time quantitative PCR was performed in the monkeys to detect CXCL11, CXCL12 and CXCR7. At the protein level CXCL12 and CXCR7 were localized in marmoset (*Callitrix jacchus*) testes whereas CXCR7 patterns were determined for various stages of human testes. Morphometry and flow cytometry were applied to quantify CXCR7-positive cells in monkeys. Transcript levels and protein expression of CXCR7 were detectable throughout testicular development. In both species CXCR7 protein expression was restricted to premeiotic germ cells. In immature marmoset testes 69.9 % $\pm$ 9% of the total germ cell population were labelled for CXCR7 whereas in the adult 4.7% $\pm$ 2.7% were positive for CXCR7. CXCL12 mRNA was detectable in all developmental stages of marmosets. The CXCL12 protein was exclusively localized to Sertoli cells. This pattern of CXCL12/CXCR7 indicates its involvement in regulatory processes that may concert the interaction between undifferentiated germ cells and Sertoli cells.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

39 **Introduction:**

40 The (C-X-C-motif) chemokine receptor type 7 (CXCR7) is a orphanized G-protein-coupled
41 seven-span transmembrane receptor which binds the chemokines CXCL12 (also called
42 stromal-derived factor-1, SDF-1) and CXCL11 (also called interferon-inducible T cell α -
43 chemoattractant, I-TAC) (Balabanian et al. 2005; Burns et al. 2006). Whereas CXCL12 is
44 expressed under homeostatic conditions to modulate immune surveillance and development,
45 the expression of CXCL11 is upregulated upon stimuli and therefore belongs to the
46 inflammatory chemokines. In general, CXCL11 production is stimulated by interferons in a
47 variety of cells, including leukocytes, fibroblasts and endothelial cells (reviewed in: Sánchez-
48 Martin et al. 2013) and therefore supports the massive cell infiltration in several inflamed
49 tissues. The chemokines CXCL12 and CXCL11 bind also to the conventional chemokine
50 receptors CXCR4 and CXCR3, respectively (Balabanian et al. 2005). However, CXCR7
51 binds CXCL12 and CXCL11 with higher affinity and belongs to the group of atypical
52 chemokine receptors (ACKRs). Interestingly, ACKRs are unable to couple to G-proteins
53 following ligand binding and therefore fail to induce the full spectrum of classical G-protein-
54 coupled chemokine receptor signalling and cellular responses (Comerford et al. 2007;
55 Ulvmar et al. 2011; Cancellieri et al. 2013; Nibbs et al. 2013). Instead, recent studies have
56 demonstrated that CXCR7 can signal through versatile adapter molecules and acts as an
57 endogenous β -arrestin-biased signalling receptor (Rajagopal et al. 2010). In addition,
58 CXCR7 may serve as a co-receptor for CXCR4 and therefore enhances the classical
59 CXCL12-mediated G-protein signalling (Levoye et al. 2009; Luker et al. 2009).

60 In particular, the action of CXCR7 has been shown to be important for several steps in
61 developing organs (Sánchez-Martin et al. 2013). Interestingly, expression of CXCR7 was
62 detected in various cell types. For instance, expression was found on the cell surface of
63 mesenchymal stem cells (Liu et al. 2010) as well as on renal progenitor cells (Mazzeinghi et
64 al. 2008). However, in the primordium of the zebrafish, CXCR7 expression was restricted to

the somatic environment. More specifically, it has been demonstrated that a CXCL12 gradient in the somatic tissue facilitates the directed migration of CXCR4-positive primordial germ cells (PGCs) towards the future gonad (Boldajipour et al. 2008; Mahabaleshwar et al. 2008; Staton et al. 2011; Mahabaleshwar et al. 2012). In this regard, expression of CXCR7 is of particular importance as it acts as a scavenger for CXCL12 and is therefore responsible for the generation of a CXCL12 gradient (Boldajipour et al. 2008; Mahabaleshwar et al. 2008; Staton et al. 2011; Mahabaleshwar et al. 2012). In addition, the CXCL12/CXCR4 interaction is also of importance for PGC migration in mice, suggesting a conserved mechanism (Ara et al. 2003; Molyneaux et al. 2003).

Moreover, recent studies in juvenile as well as in adult mouse and human testes have revealed that CXCL12 is secreted by Sertoli cells and showed that this chemokine is required for the migration and maintenance of undifferentiated spermatogonia within the stem cell niche (Gilbert et al. 2009; Payne et al. 2010; Kanatsu-Shinohara et al. 2012; Yang et al. 2013). Transcripts of *Cxcr7* have been detected in developing mouse testes (Westernströer et al. 2014) as well in testes obtained from adult rats and humans (Eva et al. 1993; McIver et al. 2013). In addition, the cellular localization of *Cxcr7* in developing mouse testes has been shown in premeiotic germ cells suggesting that *Cxcr7* may be involved in the regulation of the spermatogonial population (Westernströer et al. 2014).

Regarding testicular development and testicular stem cell physiology, the common marmoset monkey is similar to the human (Li et al. 2005; Mitchell et al. 2008; Albert et al. 2010; Albert et al. 2012; Lin et al. 2012; McKinnell et al. 2013) and is therefore an appropriate animal model to study the CXCR7 axis in the developing testis further. Therefore, we used immature and adult human testicular tissues and employed the marmoset monkey (*Callitrix jacchus*) as a non-human primate animal model in order to elucidate the expression pattern of CXCR7 and its ligands at distinct stages of testicular development.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

90 The primary aim of the present study was to investigate mRNA expression levels of *CXCL12*,
91 *CXCL11* and *CXCR7* in marmoset monkeys during four different phases of testicular
92 development. Based on these results, the second aim was to identify and to characterize the
93 cellular localization of CXCR7 protein expression during testicular development in the
94 marmoset and human testis.

95

96 **Material and Methods**

97 **Animals, collection and processing of testicular tissue**

98 All experimental procedures were performed in compliance with the German Federal Law on
99 the Care and Use of Laboratory Animals (animal license No. 84-02.05.20.12.0.18, LANUV
100 NRW, Germany). Common marmoset monkeys, *Callitrix jacchus*, were maintained at the
101 institutional breeding facilities under standardized conditions. Marmosets were fed with
102 pellets from Altromin (Lage, Germany) with daily supplement of fresh vegetables and fruits.
103 In addition, they had unlimited access to tap water and were kept under 12h light / 12h
104 darkness cycles.

105 Monkeys (n = 4 / group) were sacrificed under anaesthesia (ketamine/xylazine solution in
106 saline; i.m.; 0.1 ml per 100 gram body weight) at the age of 16, 32, 52 and > 80 weeks *post-*
107 *partum* (*wpp*). Testes were dissected and one half was snap-frozen for subsequent RNA
108 expression analysis, whereas the remaining tissue was fixed in Bouin's solution for 24 hrs,
109 transferred into 70% EtOH and embedded in paraffin for immunohistochemical analyses.

110 For flow cytometric analysis, adult marmosets (n = 3) were sacrificed as outlined above.
111 Testes were dissected and transferred into pre-cooled minimum essential medium-alpha
112 (MEM-α, Life Technologies, Gibco, Darmstadt, Germany), followed by enzymatic digestion
113 for flow cytometry.

114 Human testicular tissues and preparation for histological analyses

115 Human adult testicular tissues were obtained from patients attending the department of
116 Clinical Andrology due to treatment of testicular tumours. Of such patients testicular tissue
117 which was assessed to be unaffected by the tumour and with complete spermatogenesis
118 was used in this study (Ethics Committee of the Medical Faculty of Muenster and the State
119 Medical Board no. 2006-588-f-S). Status of spermatogenesis was assessed according to
120 Bergmann and Kliesch (2010).

121 Tissue was fixed in 4% paraformaldehyde (PFA) for 24 hrs, before it was transferred into an
122 increasing ethanol series (30%, 50% and 70%) and embedded in paraffin. For histological
123 evaluation, 3 µm sections were prepared and were stained using periodic acid Schiff's
124 reagent (PAS staining) followed by haematoxylin counterstaining as previously described
125 (Brinkworth et al. 1995). Based on the histological analysis, testicular tissues (n = 2) with
126 normal spermatogenesis were selected for this study.

127 In addition, human testicular tissues (age: 2 weeks and 6 months) from the Centre for
128 Reproductive Health at the University of Edinburgh were used. Testes were obtained at
129 autopsy with consent of their legal guardian from boys who died from various causes
130 (excluding reproductive and endocrine abnormalities). Testes were fixed in 10% Neutral
131 Buffered Formalin (NBF) for at least 24 hrs and transferred into 70% EtOH for processing.

133 Immunofluorescence on testicular tissue

134 Paraffin was removed using Pro Taqs Clear (Pro Taqs Clear, Quartett Immunodiagnostika &
135 Biotechnologie, Berlin, Germany) and sections were rehydrated in a decreasing ethanol
136 series. After washing with phosphate-buffered saline (PBS), sections were incubated with 5%
137 (w/v) bovine serum albumin (BSA) in Tris-buffered saline (TBS) for 20 min at room

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

temperature (RT). Primary antibodies against CXCR7 (rabbit polyclonal anti-CXCR7, Abcam, dilution of 1:400, Cambridge, UK) and MAGE-A4 (mouse anti-MAGE-A4, provided by Prof. G. C. Spagnoli from the University Hospital of Basel Switzerland, dilution of 1:20) were applied to the sections and sections were incubated in a humid chamber at 4°C for 1 hr. Following two washing steps in PBS, the appropriate Alexa fluor 488-linked or 546-linked secondary antibodies, diluted in TBS/5% BSA, were applied for 45 min at RT in the dark. Incubation with non-specific immunoglobulin G (rabbit IgG and mouse IgG, dilution of 1:1000) was used as negative control. Cells were counterstained with Hoechst (Sigma-Aldrich, Steinheim, Germany).

Immunohistochemistry on testicular tissue

Sections were dewaxed and rehydrated as described above. After rinsing with distilled water and TBS, non-specific peroxidases were blocked with 3% (v/v) H₂O₂ for 15 min at RT. In order to block non-specific binding sites, sections were incubated with 5% (w/v) bovine serum albumin (BSA) for 20 min at RT. Subsequently, primary antibodies against CXCR7 (rabbit polyclonal anti-CXCR7, Abcam, dilution of 1:400, Cambridge, UK) and CXCL12 (mouse monoclonal anti-CXCL12, R&D system, dilution of 1:50, Wiesbaden, Germany) were applied and sections were incubated overnight in a humid chamber at 4°C. Incubation with corresponding non-specific immunoglobulin G (rabbit IgG or mouse IgG, respectively) antibodies served as negative control. After three washing times with TBS, sections were incubated with biotin conjugated secondary antibodies and subsequently labeled with a streptavidin-conjugated horseradish peroxidase (Sigma-Aldrich, Steinheim, Germany). Staining was visualized using 3,3'-diaminobenzidine as chromogen (Sigma-Aldrich, Steinheim, Germany), and haematoxylin as counterstain.

163 **Morphometric evaluation**

164 For morphometric evaluation of marmoset testicular tissues, 4- μ m sections were taken at 40-
165 μ m intervals. In tissue sections from infantile marmosets, a total of 100 seminiferous tubules
166 per animal (n = 3) were analyzed to determine labeling indices of CXCR7-positive germ cells.
167 In accordance with previous publications germ cell types were categorized into the following
168 three groups, cells with no observable contact with the basement membrane were classified
169 as gonocytes, those with some point of contact as pre-spermatogonia and those with full
170 contact with the basement membrane as spermatogonia (Sharpe et al. 2003b). Additionally,
171 the percentages of CXCR7-positive and negative gonocytes, pre-spermatogonia and
172 spermatogonia were determined.

174 **RNA isolation and relative gene expression analysis**

175 RNA extraction was performed using the miRNeasy Kit (Quiagen, Hilden, Germany) and
176 genomic DNA was removed by DNase treatment. cDNA was generated with the iScript
177 cDNA Synthesis Kit (Bio-Rad, Munich, Germany) using 200 ng of total RNA. Specific primers
178 were designed using Primer Express® 3.0 Software (Life Technologies, Darmstadt,
179 Germany). Optimal primer concentrations, primer specificity and PCR efficiency were
180 evaluated following the Power SYBR® Green PCR User Guide (Life Technologies,
181 Darmstadt, Germany). Primer sequences are summarized in Table 1. For quantitative real-
182 time PCR (qRT-PCR) analyses, cDNA was diluted 1:10 and 1.5 μ l were used for each 15 μ l
183 PCR reaction with Power SYBR® Green Mastermix (Life Technologies, Darmstadt,
184 Germany). The PCR programme consisted of initial steps of activation and denaturation
185 which were run once for 10 min at 95°C, followed by 40 cycles of denaturation (15 sec at
186 95°C), annealing and elongation (1 min at 60°C). qRT-PCRs were run on the StepOnePlus™
187 (Life Technologies, Darmstadt, Germany) and were subsequently analysed using the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

StepOne™ software 2.2 (Life Technologies, Darmstadt, Germany). To calculate relative expression levels of *CXCR7* (n = 4 / group), *CXCL12* (n = 4 / group) and *CXCL11* (n = 4 / group) in marmoset testes, the 2^(-ΔCT) method (Livak et al. 2001) was applied, using *TOP1* as reference gene. Each sample was measured in duplicate.

Enzymatic digestion and flow cytometry of testicular cells

Single-cell suspension of testicular cells was prepared by sequential enzymatic digestion. The seminiferous tubules were digested in MEM-α with 1 mg/mL collagenase I (Sigma Aldrich, Steinheim, Germany) in a water bath at 37°C for 40 min. The digestion was stopped by adding MEM-α containing 10% fetal bovine serum (FBS, Life Technologies, PAA, Darmstadt, Germany). Afterwards, the seminiferous tubular fragments were isolated from interstitial cells by sedimentation at unit gravity and the seminiferous tubular fragments were further digested with 1 mg/mL collagenase I, 2.2 mg/mL DNase I (Sigma Aldrich, Steinheim, Germany) and 0.5 mg/mL hyaluronidase (Sigma Aldrich, Steinheim, Germany) in MEM-α until a single-cell suspension was achieved. Finally, the single-cell suspension was centrifuged (1500 rpm, 4 min), the supernatant was removed and enzymatic digestion was stopped with FBS.

For haematoxylin staining, testicular single cells were fixed on poly L-lysine coverslips with 4% PFA for 15 min. After rinsing with PBS, cells were counterstained with haematoxylin at RT for 1 min and afterwards analyzed.

For flow cytometric analysis, single-cell suspensions were filtered through a 30 μm mesh and washed with 5% (w/v) goat serum. Subsequently, cells were collected by centrifugation (1500 rpm, 5 min), the supernatant was removed and cells were incubated with the primary antibody against CXCR7 (rabbit polyclonal anti-CXCR7, Abcam, dilution of 1:200, Cambridge, UK) on ice for 30 min. Incubation with the corresponding non-specific

immunoglobulin G (rabbit IgG, dilution of 1:2000) served as negative control. Following two washing steps in 5% (w/v) goat serum, cells were incubated on ice with an Alexa fluor 488-linked secondary antibody (goat anti-rabbit Alexa fluor 488, Life Technologies, Darmstadt, Germany, diluted in 5% goat serum for 30 min. Finally, stained cells were analysed using a BD FACS Aria II (Becton Dickinson).

Immunofluorescence staining of testicular single cells

Fixed testicular single cells on poly L-lysine coverslips were rinsed with PBS. Cells were then incubated with 5% (w/v) BSA at RT for 30 min. Subsequently, primary antibodies against CXCR7 (rabbit polyclonal anti-CXCR7, Abcam, dilution of 1:400, Cambridge, UK) were applied and cells were incubated in a humid chamber at 4 °C for 1 hr. Following two washing steps in PBS, the appropriate Alexa fluor 546-linked secondary antibody, diluted in TBS/5% BSA, was applied for 30 min at RT in the dark. Incubation with non-specific immunoglobulin G (rabbit IgG, dilution of 1:2000) was performed as negative control. Cells were counterstained with Hoechst (Sigma-Aldrich, Steinheim, Germany).

Histological sample evaluations and image acquisition

Samples were analyzed using an Olympus BX61 microscope (Melville, NY, USA) with an attached Retiga 400R camera (QImaging, Burnaby, BC, Canada). All images were acquired digitally using cellSens imaging software (Olympus, Münster, Germany) and were processed using Adobe Photoshop CS2 9.0 (Adobe Systems, CA, USA).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Statistical analysis

Statistical analyses were performed using Graph Pad Prism 5 (Graph Pad Software, USA). For analysis of qRT-PCR results the non-parametric Kruskal-Wallis test was performed followed by the Dunn’s multiple comparison test. For morphometric analyses the Student’s t-test was applied. For these two tests, significant differences between the groups are marked with *P<0.05.

Significant differences between the testis weights were determined using the “One way analysis of variance” followed by the “Tukey’s multiple comparison test”. Groups marked with different letters (a, b, c, d) are significantly different from each other (P<0.001).

Results

Testicular development, testis growth and expression pattern of *CXCL12*, *CXCL11* and *CXCR7* in postnatal marmoset testes.

Testicular tissues from four phases of testicular development (16 *wpp*, 32 *wpp*, 52 *wpp* and > 80 *wpp*) were selected to characterize the expression pattern of *CXCR7*, *CXCL12* and *CXCL11* in the marmoset monkey.

Histological examination of testicular tissues of marmoset monkeys at the age of 16 *wpp* (Fig. 1a) revealed the presence of gonocytes, pre-spermatogonia and spermatogonia. At the age of 32 *wpp*, differentiated germ cells up to spermatocytes were detected in seminiferous tubules of marmoset monkeys (Fig. 1b). At the age of 52 *wpp* (Fig. 1c), round as well as elongated spermatids were observed in the seminiferous tubules. Finally, spermatogenesis was completed in the seminiferous tubules of marmoset monkeys at the age of > 80 *wpp* (Fig. 1d). In these age groups, testis weights (Fig. 1e) increased significantly from 36.4 ± 6.3 mg (16 *wpp*) via 181.9 ± 35.7 mg (32 *wpp*) and 376.7 ± 72.0 mg (52 *wpp*) to 481.9 ± 36.5 mg (> 80 *wpp*).

Relative expression levels of *CXCR7* ($n = 4$ / group), *CXCL12* ($n = 4$ / group) and *CXCL11* ($n = 4$ / group) during postnatal testis development in the marmoset were determined using qRT-PCR analysis. Expression of *CXCL12* was high in marmoset testes at the age of 16 *wpp* (0.214445 ± 0.112887) and then decreased significantly (0.088170 ± 0.012248 ; 2.4-fold; 52 *wpp*) during testicular development (Fig. 1f). Compared to *CXCL12*, expression of *CXCL11* was 56-fold (16 *wpp*), 76-fold (32 *wpp*), 21-fold (52 *wpp*) and 23-fold lower (> 80 *wpp*) and did not change significantly during postnatal testis development (Fig. 1g). Expression of the receptor *CXCR7* (Fig. 1h) was high in marmoset testes at the age of 16 *wpp* (0.368448 ± 0.094921) and decreased significantly (0.092540 ± 0.013916 ; 4-fold; > 80 *wpp*) during postnatal development.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

CXCL12 and CXCR7 protein expression in postnatal marmoset testes

Immunohistochemistry was performed to determine the expression of the homeostatic chemokine CXCL12 and the corresponding receptor CXCR7 during postnatal testis development in marmoset monkeys.

In the testes of marmoset monkeys at the age of 16 *wpp* (Fig. 2a), 32 *wpp* (Fig. 2c, d), 52 *wpp* (Fig. 2e, f) and > 80 *wpp* (Fig. 2g, h), CXCL12 was exclusively localized in the cytoplasm of Sertoli cells. In contrast, expression of CXCR7 was restricted to the germ cell population. More specifically, in marmoset testes at the age of 16 *wpp* the expression of CXCR7 was found in a subset of gonocytes (Fig. 3c), pre-spermatogonia (Fig. 3e) and spermatogonia (Fig. 3g). Furthermore, morphometric analyses revealed that the CXCR7 labelling index of the entire germ cell population was $69.9 \pm 9.0\%$ (Fig. 3i), which comprised CXCR7-positive gonocytes (Fig. 3c), pre-spermatogonia (Fig. 3f) and spermatogonia (Fig. 3h) with the indices of $36.8 \pm 13.7\%$, $18.0 \pm 4.1\%$ and $15.1 \pm 9.8\%$, respectively (Fig. 3i). Double immunofluorescence stainings (Fig. 4a-l) revealed subpopulations of MAGE-A4⁺/CXCR7⁺ and MAGE-A4⁺/CXCR7⁻ gonocytes (Fig. 4d, e, f), pre-spermatogonia (Fig. 4g, h, i) and spermatogonia (Fig. 4j, k, l). However, subsets of MAGE-A4⁺/CXCR7⁻ pre-spermatogonia (Fig. 4g, h, i) were only rarely observed in tubules of marmoset testes at the age of 16 *wpp*.

During postnatal development CXCR7 expression persisted in premeiotic germ cells located at the basement membrane (Fig. 5a - f). CXCR7-positive A_{pale} (Fig. 5b, d) and A_{dark} (Fig. 5f) spermatogonia were identified by morphological evaluation in marmoset monkeys at the age of 32 *wpp*, 52 *wpp* and > 80 *wpp*.

Furthermore, adult marmoset testes (n = 3) were enzymatically digested and isolated single cells were subjected to haematoxylin staining which showed cell types with different sizes and morphology (Fig. 5g). Subsequent, flow cytometric assessment revealed expression of

CXCR7 on the cell surface of $4.7 \pm 2.7\%$ of cells in the single cell suspension obtained from adult marmoset testes (Table S2) and CXCR7 protein expression was confirmed by immunofluorescence staining of cells fixed on poly L-lysine slides (Fig. 5h). The qualitative observations of CXCR7-positive cells following immunofluorescence staining were consistent with the data obtained by flow cytometry.

CXCR7 protein profiles in postnatal human testes

A panel of rare postnatal human testis samples (2 weeks, 6 months and adult) was immunohistochemically stained and revealed that a subset of gonocytes, pre-spermatogonia as well as spermatogonia was positive for CXCR7 in immature human testes (Fig. 6a-c). Moreover, the expression of CXCR7 in adult human testes (Fig. 6d) was found in premeiotic germ cells at the basement membrane.

Discussion

We have illustrated a schematic drawing of the localization patterns of CXCL12 and CXCR7 which are pointing to a crucial role of this chemokine/receptor pair in the developing primate testis (Fig. 7). Interestingly, transcript levels of CXCL12 were constitutively expressed and immunohistochemical analyses further demonstrated protein expression of CXCL12 in the cytoplasm of Sertoli cells at all stages of postnatal testicular development which indicated a non-maturation-dependent expression of the homeostatic chemokine CXCL12 in the developing marmoset testis. After birth, the highest number of Sertoli cells per testis is established at around 4 month of age. Afterwards, the Sertoli cell population decreases and after 8 month of age Sertoli cells become morphologically mature (e.g. establishment of tight-junctions between adjacent Sertoli cells, formation of lumen in the seminiferous tubules) in

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

marmoset testes (Sharpe et al. 2003a; Li et al. 2005). We therefore assume this phenomenon to explain the significantly decreased transcript levels of *CXCL12* in the developing marmoset testis. The Sertoli cell-specific expression pattern of *CXCL12* described in this study is in line with previous mouse (Payne et al. 2010; Yang et al. 2013, Westernströer et al. 2014) and human data (Gilbert et al. 2009). Here we describe for marmosets a decline of *CXCL12* from week 16-52 (prepuberty to postpuberty). This mRNA pattern is different from that previously observed in mice. The mRNA levels of *Cxcl12* in developing mouse testes remained constant from birth until day 21 and then increased until adulthood (Westernströer et al. 2014). The two methods for quantification of mRNA are different (mouse: luciferase based normalization; marmoset: reference gene based normalization) and the reported levels can therefore not be easily compared. However in mice and marmosets transcripts for *CXCL12* are detectable and appear to be modified throughout pubertal development. Current results therefore suggest that the action of *CXCL12* is evolutionarily conserved. Functions of *CXCL12* have been described to be organ or tissue specific. This includes a role for cell proliferation or survival (Kortesidis et al. 2005; Burns et al. 2006; Mazzinghi et al. 2008), differentiation (Lazarini et al. 2003), adhesion (Burns et al. 2006; Mazzinghi et al. 2008) and the regulation of cell migration (Bhakta et al. 2006; Mazzinghi et al. 2008). Regarding germ cells, *CXCL12* has been shown to be secreted by the somatic environment thereby facilitating the colonization of the gonads by PGCs during mouse embryonic development (Doitsidou et al. 2002; Ara et al. 2003; Molyneaux et al. 2003; Dambly-Chaudiere et al. 2007; Boldajipour et al. 2008). Furthermore, recent studies in neonatal and adult mice revealed that *CXCL12* is required for maintenance of spermatogonial stem cells (SSCs) within their niches as well as for the homing and colonization process of SSCs into their niches following germ cell transplantation assays (Payne et al. 2010; Kanatsu-Shinohara et al. 2012; Yang et al. 2013). Moreover, we previously reported significantly increased transcript levels of *Cxcl12* following germ cell

depletion in mice and suggested that Cxcl12 may stimulate the spermatogenic repopulation of the seminiferous tubules during testicular recovery (Westernströer et al. 2014).

Whereas inflammatory chemokines (e.g. the C-X-C chemokine CXCL11) are mainly involved in the recruitment of leukocytes to inflamed tissue sites, homeostatic chemokines (e.g. the C-X-C chemokine CXCL12) are constitutively expressed in organs and have important functions in stem cell homing, tissue homeostasis and repair. Indications for testicular inflammation (e.g. massive cell infiltration by neutrophils, lymphocytes, macrophages and mast cells in the interstitium) (Guazzone et al. 2009) were not seen in the analyzed testes. Moreover, in contrast to CXCL12, only low transcript levels of CXCL11 were detected in developing marmoset testes which confirmed that inflammatory C-X-C chemokines, especially CXCL11 as a secondary ligand for CXCR7, is likely not of functional relevance for healthy marmoset testes.

Transcript levels of CXCR7 were also constitutively expressed and immunohistochemical analyses further demonstrated that the atypical chemokine receptor CXCR7 is exclusively expressed by premeiotic germ cells (gonocytes, pre-spermatogonia, spermatogonia including A_{pale} and A_{dark} spermatogonia) throughout various phases of postnatal testicular development in the marmoset and the human testis. In agreement with these data we found that Cxcr7 protein can be detected in gonocytes as well as in Sall4-positive undifferentiated spermatogonia of the developing mouse testis (Westernströer et al. 2014). Additionally, Cxcr7 was also expressed on differentiated germ cells, spermatids and interstitial cells. These results reveal an evolutionarily conserved expression pattern of CXCR7 in premeiotic germ cells in the mouse and both analysed primate species (Westernströer et al. 2014). The morphometric evaluation in immature marmoset testes has shown additionally that approximately 80% of migrating germ cells (gonocytes and pre-spermatogonia) express CXCR7. In general, recent studies in marmoset monkeys indicated that the migration of gonocytes most likely begins postnatally between birth and 4 months of age (Rune et al. 2016).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1991; Sharpe et al. 2003b). Interestingly, several MAGE-A4⁺/CXCR7⁻ gonocytes were observed, whereas subsets of MAGE-A4⁺/CXCR7⁻ pre-spermatogonia were only rarely seen in seminiferous tubules, suggesting that CXCR7 may be involved in migration and/or adhesion of germ cells in immature marmoset testes. In general, migration of centrally located germ cells into the basal part of the seminiferous epithelium is a cellular process critical for germ cell survival and establishment of A_{dark} and A_{pale} spermatogonia which represent the reserve and active SSC pool in adulthood in marmoset and human testis (McGuinness and Orth, 1992; Orth et al. 2000). Moreover, our morphometric evaluation in immature marmoset testes has shown that around 50% of the spermatogonia at the basement membrane express CXCR7. Furthermore, we demonstrated that A_{dark} and A_{pale} spermatogonia express CXCR7 in marmoset testes at the age of 32 *wpp*, 52 *wpp* and > 80 *wpp*. Therefore, we assume different roles for the atypical chemokine receptor CXCR7 in the developing marmoset testes. Tissue- and cell type-specific functions of CXCR7 have been described and depend on the cell type and tissue. This includes a role for cell survival or adhesion as described for example for in the breast cancer cell line MCF-7 (Burns et al. 2006) as well as in renal multipotent progenitor cells of the human kidney (Mazzeinghi et al. 2008).

In addition, using HEK-293 and MDA-MB-231 cells, it has been demonstrated that receptor heterodimerization between CXCR4 and CXCR7 may enhance CXCL12-induced cell migration (Levoye et al. 2009; Decaillot et al. 2011). Regarding the localization of CXCR4 in developing mammalian testes, different expression patterns have been shown for mouse and human (Gilbert et al. 2009; Yoon et al. 2009; Kanatsu-Shinohara et al. 2012; Yang et al. 2013, McIver et al. 2013). Whilst expression of CXCR4 was found in gonocytes, undifferentiated spermatogonia, Sertoli cells and interstitial cells of neonatal and adult mouse testes (Yoon et al. 2009; Kanatsu-Shinohara et al. 2012; Yang et al. 2013), CXCR4 protein was detected throughout the seminiferous tubules of adult human testes (Gilbert et al. 2009).

398 So far, the expression profile and the role of CXCR4 in developing marmoset testes remain
399 unknown. Nevertheless, receptor heterodimerization between CXCR4 and CXCR7
400 throughout testicular development in primate testes cannot be excluded but needs to be
401 investigated in further studies.

402 In conclusion, we show that CXCL12/CXCR7 network is present in primate testes. The
403 localization pattern suggests its involvement during migration, survival or adhesion in
404 premeiotic germ cells.

405

406

407 **Declaration of interest**

408 The authors declare that there is no conflict of interest that could be perceived as prejudicing
409 the impartiality of the research reported.

410 **Funding**

411 The study was supported by a grant from the Center of Interdisciplinary Clinical Research
412 Muenster (IZKF; Schl2/001/13), the Innovative Medical Research of the University of Münster
413 Medical School (KO 111014) and the DFG project (SCHL 394/11-2).

414

415 **Acknowledgments**

416 We gratefully acknowledge Martin Heuermann and Günter Stelke (Centre of Reproductive
417 Medicine and Andrology, Münster, Germany), for excellent animal caretaking as well as
418 Nicole Terwort, Reinhild Sandhowe, Jutta Salzig and Adelheid Kersebom for tremendous

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

technical assistance. Flow cytometric analysis was performed at the Max-Planck-Institute for
Molecular Biomedicine (MPI, Münster) with the help of Dr. Martin Stehling.

References

Albert S, Ehmcke J, Wistuba J, Eildermann K, Behr R, Schlatt S, Gromoll J (2010) Germ cell
dynamics in the testis of the postnatal common marmoset monkey (*Callithrix jacchus*).
Reproduction 140:733-742

Albert S, Wistuba J, Eildermann K, Ehmcke J, Schlatt S, Gromoll J, Kossack N (2012)
Comparative marker analysis after isolation and culture of testicular cells from the immature
marmoset. *Cells Tissues Organs* 196:543-554

Ara T, Nakamura Y, Egawa T, Sugiyama T, Abe K, Kishimoto T, Matsui Y, Nagasawa T
(2003) Impaired colonization of the gonads by primordial germ cells in mice lacking a
chemokine, stromal cell-derived factor-1 (SDF-1). *Proc Natl Acad Sci U S A* 100:5319-5323

Balabanian K, Lagane B, Infantino S, Chow KY, Harriague J, Moepps B, Arenzana-
Seisdedos F, Thelen M, Bachelier F (2005) The chemokine SDF-1/CXCL12 binds to and
signals through the orphan receptor RDC1 in T lymphocytes. *J Biol Chem* 280:35760-35766

Bergmann M, Kliesch S (2010) Testicular biopsy and histology. In: Nieschlag E (ed)
Andrology Male Reproductive Health and Dysfunction. Springer-Verlag Berlin Heidelberg, pp
155-167

Bhakta S, Hong P, Koc O (2006) The surface adhesion molecule CXCR4 stimulates
mesenchymal stem cell migration to stromal cell-derived factor-1 in vitro but does not
decrease apoptosis under serum deprivation. *Cardiovasc Revasc Med* 7:19-24

- 441 Boldajipour B, Mahabaleshwar H, Kardash E, Reichman-Fried M, Blaser H, Minina S, Wilson
442 D, Xu Q, Raz E (2008) Control of chemokine-guided cell migration by ligand sequestration.
443 Cell 132:463-473
- 444 Brinkworth MH, Weinbauer GF, Schlatt S, Nieschlag E (1995) Identification of male germ
445 cells undergoing apoptosis in adult rats. J Reprod Fertil 105:25-33
- 446 Burns JM, Summers BC, Wang Y, Melikian A, Berahovich R, Miao Z, Penfold ME, Sunshine
447 MJ, Littman DR, Kuo CJ, Wei K, McMaster BE, Wright K, Howard MC, Schall TJ (2006) A
448 novel chemokine receptor for SDF-1 and I-TAC involved in cell survival, cell adhesion, and
449 tumor development. J Exp Med 203:2201-2213
- 450 Burns JM, Summers BC, Wang Y, Melikian A, Berahovich R, Miao Z, Penfold ME, Sunshine
451 MJ, Littman DR, Kuo CJ, Wei K, McMaster BE, Wright K, Howard MC, Schall TJ (2006) A
452 novel chemokine receptor for SDF-1 and I-TAC involved in cell survival, cell adhesion, and
453 tumor development. J Exp Med 203:2201-2213
- 454 Cancellieri C, Vacchini A, Locati M, Bonecchi R, Borroni EM (2013) Atypical chemokine
455 receptors: from silence to sound. Biochem Soc Trans 41:231-236
- 456 Comerford I, Litchfield W, Harata-Lee Y, Nibbs RJ, McColl SR (2007) Regulation of
457 chemotactic networks by 'atypical' receptors. Bioessays 29:237-247
- 458 Dambly-Chaudiere C, Cubedo N, Ghysen A (2007) Control of cell migration in the
459 development of the posterior lateral line: antagonistic interactions between the chemokine
460 receptors CXCR4 and CXCR7/RDC1. BMC Dev Biol 7:23
- 461 Decaillot FM, Kazmi MA, Lin Y, Ray-Saha S, Sakmar TP, Sachdev P (2011) CXCR7/CXCR4
462 heterodimer constitutively recruits beta-arrestin to enhance cell migration. J Biol Chem
463 286:32188-32197

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Doitsidou M, Reichman-Fried M, Stebler J, Koprunner M, Dorries J, Meyer D, Esguerra CV, Leung T, Raz E (2002) Guidance of primordial germ cell migration by the chemokine SDF-1. *Cell* 111:647-659

Eva C, Sprengel R (1993) A novel putative G protein-coupled receptor highly expressed in lung and testis. *DNA Cell Biol* 12:393-399

Gilbert DC, Chandler I, McIntyre A, Goddard NC, Gabe R, Huddart RA, Shipley J (2009) Clinical and biological significance of CXCL12 and CXCR4 expression in adult testes and germ cell tumours of adults and adolescents. *J Pathol* 217:94-102

Guazzone VA, Jacobo P, Theas MS, Lustig L (2009) Cytokines and chemokines in testicular inflammation: A brief review. *Microsc Res Tech* 72:620-628

Kanatsu-Shinohara M, Inoue K, Takashima S, Takehashi M, Ogonuki N, Morimoto H, Nagasawa T, Ogura A, Shinohara T (2012) Reconstitution of mouse spermatogonial stem cell niches in culture. *Cell Stem Cell* 11:567-578

Kortesidis A, Zannettino A, Isenmann S, Shi S, Lapidot T, Gronthos S (2005) Stromal-derived factor-1 promotes the growth, survival, and development of human bone marrow stromal stem cells. *Blood* 105:3793-3801

Lazarini F, Tham TN, Casanova P, Arenzana-Seisdedos F, Dubois-Dalcq M (2003) Role of the alpha-chemokine stromal cell-derived factor (SDF-1) in the developing and mature central nervous system. *Glia* 42:139-148

Levoye A, Balabanian K, Baleux F, Bachelerie F, Lagane B (2009) CXCR7 heterodimerizes with CXCR4 and regulates CXCL12-mediated G protein signaling. *Blood* 113:6085-6093

Li LH, Donald JM, Golub MS (2005) Review on testicular development, structure, function, and regulation in common marmoset. *Birth Defects Res B Dev Reprod Toxicol* 74:450-469

- 487 Lin ZY, Imamura M, Sano C, Nakajima R, Suzuki T, Yamadera R, Takehara Y, Okano HJ,
488 Sasaki E, Okano H (2012) Molecular signatures to define spermatogenic cells in common
489 marmoset (*Callithrix jacchus*). *Reproduction* 143:597-609
- 490 Liu H, Xue W, Ge G, Luo X, Li Y, Xiang H, Ding X, Tian P, Tian X (2010) Hypoxic
491 preconditioning advances CXCR4 and CXCR7 expression by activating HIF-1alpha in MSCs.
492 *Biochem Biophys Res Commun* 401:509-515
- 493 Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time
494 quantitative PCR and the 2⁻(Delta Delta C(T)) Method. *Methods* 25:402-408
- 495 Luker K, Gupta M, Luker G (2009) Bioluminescent CXCL12 fusion protein for cellular studies
496 of CXCR4 and CXCR7. *BioTechniques* 47:625-632
- 497 Luker K, Gupta M, Luker G (2009) Bioluminescent CXCL12 fusion protein for cellular studies
498 of CXCR4 and CXCR7. *BioTechniques* 47:625-632
- 499 Mahabaleshwar H, Boldajipour B, Raz E (2008) Killing the messenger: The role of CXCR7 in
500 regulating primordial germ cell migration. *Cell Adh Migr* 2:69-70
- 501 Mahabaleshwar H, Tarbashevich K, Nowak M, Brand M, Raz E (2012) Beta-Arrestin Control
502 of Late Endosomal Sorting Facilitates Decoy Receptor Function and Chemokine Gradient
503 Formation. *Development* 139:2897-2902
- 504 Mazinghi B, Ronconi E, Lazzeri E, Sagrinati C, Ballerini L, Angelotti ML, Parente E,
505 Mancina R, Netti GS, Becherucci F, Gacci M, Carini M, Gesualdo L, Rotondi M, Maggi E,
506 Lasagni L, Serio M, Romagnani S, Romagnani P (2008) Essential but differential role for
507 CXCR4 and CXCR7 in the therapeutic homing of human renal progenitor cells. *J Exp Med*
508 205:479-490

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

509 McGuinness MP, Orth JM (1992) Gonocytes of male rats resume migratory activity
510 postnatally. *Eur J Cell Biol* 59:196-210

511 McIver SC, Loveland KL, Roman SD, Nixon B, Kitazawa R, McLaughlin EA (2013) The
512 chemokine CXCL12 and its receptor CXCR4 are implicated in human seminoma metastasis.
513 *Andrology* 1:517-529

514 McKinnell C, Mitchell RT, Morris K, Anderson RA, Kelnar CJ, Wallace WH, Sharpe RM
515 (2013) Perinatal germ cell development and differentiation in the male marmoset (*Callithrix*
516 *jacchus*): similarities with the human and differences from the rat. *Hum Reprod* 28:886-896

517 Mitchell RT, Cowan G, Morris KD, Anderson RA, Fraser HM, Mckenzie KJ, Wallace WH,
518 Kelnar CJ, Saunders PT, Sharpe RM (2008) Germ cell differentiation in the marmoset
519 (*Callithrix jacchus*) during fetal and neonatal life closely parallels that in the human. *Hum*
520 *Reprod* 23:2755-2765

521 Molyneaux KA, Zinszner H, Kunwar PS, Schaible K, Stebler J, Sunshine MJ, O'Brien W, Raz
522 E, Littman D, Wylie C, Lehmann R (2003) The chemokine SDF1/CXCL12 and its receptor
523 CXCR4 regulate mouse germ cell migration and survival. *Development* 130:4279-4286

524 Nibbs RJ, Graham GJ (2013) Immune regulation by atypical chemokine receptors. *Nat Rev*
525 *Immunol* 13:815-829

526 Orth JM, Jester WF, Li LH, Laslett AL (2000) Gonocyte-Sertoli cell interactions during
527 development of the neonatal rodent testis. *Curr Top Dev Biol* 50:103-124

528 Payne CJ, Gallagher SJ, Foreman O, Dannenberg JH, Depinho RA, Braun RE (2010) Sin3a
529 is required by sertoli cells to establish a niche for undifferentiated spermatogonia, germ cell
530 tumors, and spermatid elongation. *Stem Cells* 28:1424-1434

- 531 Rajagopal S, Kim J, Ahn S, Craig S, Lam CM, Gerard NP, Gerard C, Lefkowitz RJ (2010)
532 Beta-arrestin- but not G protein-mediated signaling by the "decoy" receptor CXCR7. *Proc*
533 *Natl Acad Sci U S A* 107:628-632
- 534 Rune GM, de Souza P, Merker HJ (1991) Ultrastructural and histochemical characterization
535 of marmoset (*Callithrix jacchus*) Leydig cells during postnatal development. *Anat Embryol*
536 (Berl) 183:179-191
- 537 Sanchez-Martin L, Sanchez-Mateos P, Cabanas C (2013) CXCR7 impact on CXCL12
538 biology and disease. *Trends in Molecular Medicine* 19:12-22
- 539 Sharpe RM, McKinnell C, Kivlin C, Fisher JS (2003a) Proliferation and functional maturation
540 of Sertoli cells, and their relevance to disorders of testis function in adulthood. *Reproduction*
541 125:769-784
- 542 Sharpe RM, Fraser HM, Brougham MF, McKinnell C, Morris KD, Kelnar CJ, Wallace WH,
543 Walker M (2003b) Role of the neonatal period of pituitary-testicular activity in germ cell
544 proliferation and differentiation in the primate testis. *Hum Reprod* 18:2110-2117
- 545 Staton AA, Knaut H, Giraldez AJ (2011) miRNA regulation of Sdf1 chemokine signaling
546 provides genetic robustness to germ cell migration. *Nat Genet* 43:204-211
- 547 Ulvmar MH, Hub E, Rot A (2011) Atypical chemokine receptors. *Exp Cell Res* 317:556-568
- 548 Westernströer B, Terwort N, Ehmcke J, Wistuba J, Schlatt S, Neuhaus N (2014) Profiling of
549 Cxcl12 receptors, Cxcr4 and Cxcr7 in murine testis development and a spermatogenic
550 depletion model indicates a role for Cxcr7 in controlling Cxcl12 activity. *PLoS ONE*
551 9(12):e112598
- 552 Yang QE, Kim D, Kaucher A, Oatley MJ, Oatley JM (2013) CXCL12-CXCR4 signaling is
553 required for the maintenance of mouse spermatogonial stem cells. *J Cell Sci* 126:1009-1020

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

554 Yoon KA, Chae YM, Cho JY (2009) FGF2 stimulates SDF-1 expression through the Erm
555 transcription factor in Sertoli cells. J Cell Physiol 220:245-256
556
557

For Peer Review

Figures:

Fig. 1 Marmoset monkey testes after birth - age-related changes in histology, weight and transcript levels. Micrographs showing PAS stained tissue sections of marmoset testes at the age of 16 *wpp* (*wpp* = weeks *post partum*; a), 32 *wpp* (b), 52 *wpp* (c) and > 80 *wpp* (d). Gonocytes (arrowhead), pre-spermatogonia (white arrow) and spermatogonia (black arrow) are the predominant germ cells in marmoset testes at the age of 16 *wpp* (a). Meiotic cells (primary spermatocytes) were present in the seminiferous tubules of marmosets at the age of 32 *wpp* (b), whilst germ cells up to the stage of elongated spermatids were detected in the seminiferous epithelium of marmoset monkeys at the age of 52 *wpp* (c). Finally, spermatogenesis was complete in all marmosets at the age of > 80 *wpp* (d). Scale bars represent 50 μ m. Changes in testis weight (e) are shown as mean $\pm$ SEM. Groups marked with different letters (a, b, c, d) are significantly different from each other ($p < 0.001$). Relative mRNA levels of *CXCL12* (f), *CXCL11* (g) and *CXCR7* (h) in marmoset testes at the age of 16 *wpp*, 32 *wpp*, 52 *wpp* and > 80 *wpp* were normalized using the housekeeping gene *TOP1* (2^{- Δ Ct}) method). Results are shown as mean $\pm$ SEM and significant differences between the groups are marked with * $P < 0.05$.

Fig. 2 Immunohistochemical detection of *CXCL12* in marmoset testes at the age of 16 *wpp* (*wpp* = weeks *post partum*; a), 32 *wpp* (c, d), 52 *wpp* (e, f) and > 80 *wpp* (g, h). Incubation with non-specific IgG antibody was performed as control and a representative image of a marmoset testis at the age of 16 *wpp* is shown in (b). During testicular development, the chemokine *CXCL12* is exclusively localized in the cytoplasm of Sertoli cells (white arrow). Neither relatively undifferentiated germ cells (gonocytes, arrow head; spermatogonia, black arrow) nor differentiating germ cells showed expression of *CXCL12*. Scale bars represent 20 μ m.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 3 Immunohistochemical detection of CXCR7 and labelling index of CXCR7-positive cells in marmoset testes at the age of 16 *wpp* (*wpp* = weeks *post partum*). A representative image is showing CXCR7 expression in marmoset testes at the age of 16 *wpp* (a). The staining revealed that subsets of germ cells express CXCR7 (black arrow), whilst the remainder are negative (arrow head) for CXCR7. Sertoli cells (white arrow) in the seminiferous tubules showed no expression of CXCR7. Corresponding IgG antibody was used as negative control and a representative image is shown in (b). High magnification of CXCR7 stained tissue sections revealed CXCR7-positive and negative gonocytes (c, d), pre-spermatogonia (e, f) and spermatogonia (g, h). Scale bars represent 10 μ m. Scores of CXCR7-positive and CXCR7-negative gonocytes (Gon+/Gon-), pre-spermatogonia (pre-SPG+/pre-SPG-) and spermatogonia (SPG+/SPG-) in marmoset testes at the age of 16 *wpp* are shown in (i). Results are shown as mean $\pm$ SEM and significant differences between the groups are marked with * $P < 0.05$.

Fig. 4 Representative images for expression of MAGE-A4 (green) and CXCR7 (red) in testis sections from marmosets at the age of 16 *wpp* (a - c). MAGE-A4 expression is seen in the majority of germ cells including gonocytes (d), pre-spermatogonia (g) and spermatogonia (j). Subpopulations of MAGE-A4⁺/CXCR7⁺ (arrow heads) and/or MAGE-A4⁺/CXCR7⁻ (arrows) gonocytes (d - f), pre-spermatogonia (g - i) as well as spermatogonia (j - l) can be identified. Note that MAGE-A4⁺/CXCR7⁻ pre-spermatogonia are occasionally seen in the seminiferous tubules of marmoset testes at the age of 16 *wpp*. All sections were counterstained with Hoechst (blue). Dotted lines indicate the basement membrane of the seminiferous tubules. Scale bars represent 10 μ m.

Fig. 5 Immunohistochemical detection of CXCR7 in marmoset testes at the age of 32 *wpp* (*wpp* = weeks *post partum*; a, b), 52 *wpp* (c, d) and > 80 *wpp* (e, f). In marmoset testes at the age of 32 *wpp*, 52 *wpp* and > 80 *wpp*, premeiotic germ cells (black arrow; including A_{pale} and A_{dark} spermatogonia) located at the basement membrane are labelled for CXCR7 whereas Sertoli cells (white arrow) showed no expression for CXCR7. Scale bars represent 20 μm (a – f). Following enzymatic digestion of adult marmoset testes, haematoxylin-stained single cells showed cell types with different morphology (g) as well as immunofluorescence staining revealed CXCR7-positive cells in this single cell suspension (h). As negative control, incubation with nonspecific IgG antibody was performed and a representative image is shown in (i). Scale bars represent 50 μm (g), 15 μm (h) and 60 μm (i).

Fig. 6 Immunohistochemical detection of CXCR7 in human testes at 2 weeks (a) and 6 months after birth (c) as well as in adult (d) human testes. Corresponding IgG antibody was used as negative control and a representative image of an immature testis (2 weeks) is shown in (b). In immature human testes (a; c), gonocytes (arrow head, no contact with the basement membrane), pre-spermatogonia (white arrows, point contact with the basement membrane) and spermatogonia (black arrows, full contact with the basement membrane) were labelled for CXCR7. In adult human testes (d), premeiotic germ cells (red arrow) located at the basement membrane were labelled for CXCR7. Scale bars represent 20 μm .

Fig. 7 Schematic drawing of CXCR7 (red) and CXCL12 (green) expression in seminiferous tubules of marmoset testes at the ages of 16 *wpp* (a), 32 *wpp* (b), 52 *wpp* (c) and > 80 *wpp* (*wpp* = weeks *post partum*, d). In marmoset testes at the age of 16 *wpp*, CXCL12 is highly expressed by Sertoli cells whereas the atypical chemokine receptor CXCR7 is expressed by few gonocytes, pre-spermatogonia and spermatogonia (a). In the seminiferous epithelium of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

632 marmoset testes at the ages of 32 wpp (b), 52 wpp (c) and > 80 wpp (d), CXCL12 is still
633 highly expressed by Sertoli cells whereas CXCR7 is expressed by premeitotic germ cells,
634 including A_{pale} and A_{dark} spermatogonia. The expression pattern of CXCL12 and CXCR7
635 indicates a CXCL12/CXCR7 network in primate testes.

636

637

For Peer Review

Title: Developmental expression patterns of the chemokines CXCL11, CXCL12 and its receptor CXCR7 in common marmoset and human testes

Authors: Birgit Westernströer¹, Daniel Langenstroth¹, Sabine Kliesch², Britta Troppmann¹, Klaus Redmann¹, Joni Macdonald³, Rod Mitchell³, Joachim Wistuba¹, Stefan Schlatt¹, Nina Neuhaus¹

¹ Centre of Reproductive Medicine and Andrology, CeRA, Albert-Schweizer-Campus 1, Building D11, 48149 Münster, Germany

² Department of Clinical Andrology, Centre of Reproductive Medicine and Andrology, CeRA, Albert-Schweitzer-Campus 1, Building D11, 48149 Münster, Germany

³ MRC/University of Edinburgh Centre for Reproductive Health, The Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh, EH164TJ, Scotland, UK

Corresponding author: Prof. Dr. Stefan Schlatt

Centre of Reproductive Medicine and Andrology

Albert-Schweizer-Campus 1 (D11)

48149 Muenster, Germany

Email: Stefan.Schlatt@ukmuenster.de

Key words: CXCR7, CXCL12, CXCL11, primate testes, testicular development

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

21 **Abstract**

22 The chemokine receptor CXCR7 interacts with the chemokines CXCL11 and CXCL12.
23 During development this ligand receptor system (C-X-C) provokes cell type-specific
24 responses in terms of migration, adhesion or ligand sequestration. It is active in zebrafish
25 and rodents but no data are available for its presence or function in primate testes. Real-time
26 quantitative PCR was performed in the monkeys to detect CXCL11, CXCL12 and CXCR7. At
27 the protein level CXCL12 and CXCR7 were localized in marmoset (*Callitrix jacchus*) testes
28 whereas CXCR7 patterns were determined for various stages of human testes. Morphometry
29 and flow cytometry were applied to quantify CXCR7-positive cells in monkeys. Transcript
30 levels and protein expression of CXCR7 were detectable throughout testicular development.
31 In both species CXCR7 protein expression was restricted to premeiotic germ cells. In
32 immature marmoset testes 69.9 % $\pm$ 9% of the total germ cell population were labelled for
33 CXCR7 whereas in the adult 4.7% $\pm$ 2.7% were positive for CXCR7. CXCL12 mRNA was
34 detectable in all developmental stages of marmosets. The CXCL12 protein was exclusively
35 localized to Sertoli cells. This pattern of CXCL12/CXCR7 indicates its involvement in
36 regulatory processes that may concert the interaction between undifferentiated germ cells
37 and Sertoli cells.

38

39 Introduction:

40 The (C-X-C-motif) chemokine receptor type 7 (CXCR7) is a orphanized G-protein-coupled
41 seven-span transmembrane receptor which binds the chemokines CXCL12 (also called
42 stromal-derived factor-1, SDF-1) and CXCL11 (also called interferon-inducible T cell α -
43 chemoattractant, I-TAC) (Balabanian et al. 2005; Burns et al. 2006). Whereas CXCL12 is
44 expressed under homeostatic conditions to modulate immune surveillance and development,
45 the expression of CXCL11 is upregulated upon stimuli and therefore belongs to the
46 inflammatory chemokines. In general, CXCL11 production is stimulated by interferons in a
47 variety of cells, including leukocytes, fibroblasts and endothelial cells (reviewed in: Sánchez-
48 Martín et al. 2013) and therefore supports the massive cell infiltration in several inflamed
49 tissues. The chemokines CXCL12 and CXCL11 bind also to the conventional chemokine
50 receptors CXCR4 and CXCR3, respectively (Balabanian et al. 2005). However, CXCR7
51 binds CXCL12 and CXCL11 with higher affinity and belongs to the group of atypical
52 chemokine receptors (ACKRs). Interestingly, ACKRs are unable to couple to G-proteins
53 following ligand binding and therefore fail to induce the full spectrum of classical G-protein-
54 coupled chemokine receptor signalling and cellular responses (Comerford et al. 2007;
55 Ulvmar et al. 2011; Cancellieri et al. 2013; Nibbs et al. 2013). Instead, recent studies have
56 demonstrated that CXCR7 can signal through versatile adapter molecules and acts as an
57 endogenous β -arrestin-biased signalling receptor (Rajagopal et al. 2010). In addition,
58 CXCR7 may serve as a co-receptor for CXCR4 and therefore enhances the classical
59 CXCL12-mediated G-protein signalling (Levoye et al. 2009; Luker et al. 2009).

60 In particular, the action of CXCR7 has been shown to be important for several steps in
61 developing organs (Sánchez-Martín et al. 2013). Interestingly, expression of CXCR7 was
62 detected in various cell types. For instance, expression was found on the cell surface of
63 mesenchymal stem cells (Liu et al. 2010) as well as on renal progenitor cells (Mazzeinghi et
64 al. 2008). However, in the primordium of the zebrafish, CXCR7 expression was restricted to

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

65 the somatic environment. More specifically, it has been demonstrated that a CXCL12
66 gradient in the somatic tissue facilitates the directed migration of CXCR4-positive primordial
67 germ cells (PGCs) towards the future gonad (Boldajipour et al. 2008; Mahabaleshwar et al.
68 2008; Staton et al. 2011; Mahabaleshwar et al. 2012). In this regard, expression of CXCR7 is
69 of particular importance as it acts as a scavenger for CXCL12 and is therefore responsible
70 for the generation of a CXCL12 gradient (Boldajipour et al. 2008; Mahabaleshwar et al. 2008;
71 Staton et al. 2011; Mahabaleshwar et al. 2012). In addition, the CXCL12/CXCR4 interaction
72 is also of importance for PGC migration in mice, suggesting a conserved mechanism (Ara et
73 al. 2003; Molyneaux et al. 2003).

74 Moreover, recent studies in juvenile as well as in adult mouse and human testes have
75 revealed that CXCL12 is secreted by Sertoli cells and showed that this chemokine is required
76 for the migration and maintenance of undifferentiated spermatogonia within the stem cell
77 niche (Gilbert et al. 2009; Payne et al. 2010; Kanatsu-Shinohara et al. 2012; Yang et al.
78 2013). Transcripts of *Cxcr7* have been detected in developing mouse testes (Westernströer
79 et al. 2014) as well in testes obtained from adult rats and humans (Eva et al. 1993; McIver et
80 al. 2013). In addition, the cellular localization of *Cxcr7* in developing mouse testes has been
81 shown in premeiotic germ cells suggesting that *Cxcr7* may be involved in the regulation of
82 the spermatogonial population (Westernströer et al. 2014).

83 Regarding testicular development and testicular stem cell physiology, the common marmoset
84 monkey is similar to the human (Li et al. 2005; Mitchell et al. 2008; Albert et al. 2010; Albert
85 et al. 2012; Lin et al. 2012; McKinnell et al. 2013) and is therefore an appropriate animal
86 model to study the CXCR7 axis in the developing testis further. Therefore, we used immature
87 and adult human testicular tissues and employed the marmoset monkey (*Callitrix jacchus*) as
88 a non-human primate animal model in order to elucidate the expression pattern of CXCR7
89 and its ligands at distinct stages of testicular development.

The primary aim of the present study was to investigate mRNA expression levels of *CXCL12*, *CXCL11* and *CXCR7* in marmoset monkeys during four different phases of testicular development. Based on these results, the second aim was to identify and to characterize the cellular localization of CXCR7 protein expression during testicular development in the marmoset and human testis.

Material and Methods

Animals, collection and processing of testicular tissue

All experimental procedures were performed in compliance with the German Federal Law on the Care and Use of Laboratory Animals (animal license No. 84-02.05.20.12.0.18, LANUV NRW, Germany). Common marmoset monkeys, *Callitrix jacchus*, were maintained at the institutional breeding facilities under standardized conditions. Marmosets were fed with pellets from Altromin (Lage, Germany) with daily supplement of fresh vegetables and fruits. In addition, they had unlimited access to tap water and were kept under 12h light / 12h darkness cycles.

Monkeys (n = 4 / group) were sacrificed under anaesthesia (ketamine/xylazine solution in saline; i.m.; 0.1 ml per 100 gram body weight) at the age of 16, 32, 52 and > 80 weeks *post-partum* (*wpp*). Testes were dissected and one half was snap-frozen for subsequent RNA expression analysis, whereas the remaining tissue was fixed in Bouin's solution for 24 hrs, transferred into 70% EtOH and embedded in paraffin for immunohistochemical analyses.

For flow cytometric analysis, adult marmosets (n = 3) were sacrificed as outlined above. Testes were dissected and transferred into pre-cooled minimum essential medium-alpha (MEM- α , Life Technologies, Gibco, Darmstadt, Germany), followed by enzymatic digestion for flow cytometry.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Human testicular tissues and preparation for histological analyses

Human adult testicular tissues were obtained from patients attending the department of Clinical Andrology due to treatment of testicular tumours. Of such patients testicular tissue which was assessed to be unaffected by the tumour and with complete spermatogenesis was used in this study (Ethics Committee of the Medical Faculty of Muenster and the State Medical Board no. 2006-588-f-S). Status of spermatogenesis was assessed according to Bergmann and Kliesch (2010).

Tissue was fixed in 4% paraformaldehyde (PFA) for 24 hrs, before it was transferred into an increasing ethanol series (30%, 50% and 70%) and embedded in paraffin. For histological evaluation, 3 µm sections were prepared and were stained using periodic acid Schiff's reagent (PAS staining) followed by haematoxylin counterstaining as previously described (Brinkworth et al. 1995). Based on the histological analysis, testicular tissues (n = 2) with normal spermatogenesis were selected for this study.

In addition, human testicular tissues (age: 2 weeks and 6 months) from the Centre for Reproductive Health at the University of Edinburgh were used. Testes were obtained at autopsy with consent of their legal guardian from boys who died from various causes (excluding reproductive and endocrine abnormalities). Testes were fixed in 10% Neutral Buffered Formalin (NBF) for at least 24 hrs and transferred into 70% EtOH for processing.

Immunofluorescence on testicular tissue

Paraffin was removed using Pro Taqs Clear (Pro Taqs Clear, Quartett Immunodiagnostika & Biotechnologie, Berlin, Germany) and sections were rehydrated in a decreasing ethanol series. After washing with phosphate-buffered saline (PBS), sections were incubated with 5% (w/v) bovine serum albumin (BSA) in Tris-buffered saline (TBS) for 20 min at room

1
2
3
4 138 temperature (RT). Primary antibodies against CXCR7 (rabbit polyclonal anti-CXCR7, Abcam,
5
6 139 dilution of 1:400, Cambridge, UK) and MAGE-A4 (mouse anti-MAGE-A4, provided by Prof.
7
8 140 G. C. Spagnoli from the University Hospital of Basel Switzerland, dilution of 1:20) were
9
10 141 applied to the sections and sections were incubated in a humid chamber at 4°C for 1 hr.
11
12 142 Following two washing steps in PBS, the appropriate Alexa fluor 488-linked or 546-linked
13
14 143 secondary antibodies, diluted in TBS/5% BSA, were applied for 45 min at RT in the dark.
15
16 144 Incubation with non-specific immunoglobulin G (rabbit IgG and mouse IgG, dilution of
17
18 145 1:1000) was used as negative control. Cells were counterstained with Hoechst (Sigma-
19
20 146 Aldrich, Steinheim, Germany).
21
22
23
24
25
26
27
28

29 148 **Immunohistochemistry on testicular tissue**

30 149 Sections were dewaxed and rehydrated as described above. After rinsing with distilled water
31 150 and TBS, non-specific peroxidases were blocked with 3% (v/v) H₂O₂ for 15 min at RT. In
32 151 order to block non-specific binding sites, sections were incubated with 5% (w/v) bovine
33 152 serum albumin (BSA) for 20 min at RT. Subsequently, primary antibodies against CXCR7
34 153 (rabbit polyclonal anti-CXCR7, Abcam, dilution of 1:400, Cambridge, UK) and CXCL12
35 154 (mouse monoclonal anti-CXCL12, R&D system, dilution of 1:50, Wiesbaden, Germany) were
36 155 applied and sections were incubated overnight in a humid chamber at 4°C. Incubation with
37 156 corresponding non-specific immunoglobulin G (rabbit IgG or mouse IgG, respectively)
38 157 antibodies served as negative control. After three washing times with TBS, sections were
39 158 incubated with biotin conjugated secondary antibodies and subsequently labeled with a
40 159 streptavidin-conjugated horseradish peroxidase (Sigma-Aldrich, Steinheim, Germany).
41 160 Staining was visualized using 3,3'-diaminobenzidine as chromogen (Sigma-Aldrich,
42 161 Steinheim, Germany), and haematoxylin as counterstain.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Morphometric evaluation

For morphometric evaluation of marmoset testicular tissues, 4- μ m sections were taken at 40- μ m intervals. In tissue sections from infantile marmosets, a total of 100 seminiferous tubules per animal (n = 3) were analyzed to determine labeling indices of CXCR7-positive germ cells. In accordance with previous publications germ cell types were categorized into the following three groups, cells with no observable contact with the basement membrane were classified as gonocytes, those with some point of contact as pre-spermatogonia and those with full contact with the basement membrane as spermatogonia (Sharpe et al. 2003b). Additionally, the percentages of CXCR7-positive and negative gonocytes, pre-spermatogonia and spermatogonia were determined.

RNA isolation and relative gene expression analysis

RNA extraction was performed using the miRNeasy Kit (Quiagen, Hilden, Germany) and genomic DNA was removed by DNase treatment. cDNA was generated with the iScript cDNA Synthesis Kit (Bio-Rad, Munich, Germany) using 200 ng of total RNA. Specific primers were designed using Primer Express® 3.0 Software (Life Technologies, Darmstadt, Germany). Optimal primer concentrations, primer specificity and PCR efficiency were evaluated following the Power SYBR® Green PCR User Guide (Life Technologies, Darmstadt, Germany). Primer sequences are summarized in Table 1. For quantitative real-time PCR (qRT-PCR) analyses, cDNA was diluted 1:10 and 1.5 μ l were used for each 15 μ l PCR reaction with Power SYBR® Green Mastermix (Life Technologies, Darmstadt, Germany). The PCR programme consisted of initial steps of activation and denaturation which were run once for 10 min at 95°C, followed by 40 cycles of denaturation (15 sec at 95°C), annealing and elongation (1 min at 60°C). qRT-PCRs were run on the StepOnePlus™ (Life Technologies, Darmstadt, Germany) and were subsequently analysed using the

188 StepOne™ software 2.2 (Life Technologies, Darmstadt, Germany). To calculate relative
189 expression levels of *CXCR7* (n = 4 / group), *CXCL12* (n = 4 / group) and *CXCL11* (n = 4 /
190 group) in marmoset testes, the $2^{-\Delta\text{CT}}$ method (Livak et al. 2001) was applied, using *TOP1* as
191 reference gene. Each sample was measured in duplicate.

193 Enzymatic digestion and flow cytometry of testicular cells

194 Single-cell suspension of testicular cells was prepared by sequential enzymatic digestion.
195 The seminiferous tubules were digested in MEM- α with 1 mg/mL collagenase I (Sigma
196 Aldrich, Steinheim, Germany) in a water bath at 37°C for 40 min. The digestion was stopped
197 by adding MEM- α containing 10% fetal bovine serum (FBS, Life Technologies, PAA,
198 Darmstadt, Germany). Afterwards, the seminiferous tubular fragments were isolated from
199 interstitial cells by sedimentation at unit gravity and the seminiferous tubular fragments were
200 further digested with 1 mg/mL collagenase I, 2.2 mg/mL DNase I (Sigma Aldrich, Steinheim,
201 Germany) and 0.5 mg/mL hyaluronidase (Sigma Aldrich, Steinheim, Germany) in MEM- α
202 until a single-cell suspension was achieved. Finally, the single-cell suspension was
203 centrifuged (1500 rpm, 4 min), the supernatant was removed and enzymatic digestion was
204 stopped with FBS.

205 For haematoxylin staining, testicular single cells were fixed on poly L-lysine coverslips with
206 4% PFA for 15 min. After rinsing with PBS, cells were counterstained with haematoxylin at
207 RT for 1 min and afterwards analyzed.

208 For flow cytometric analysis, single-cell suspensions were filtered through a 30 μm mesh and
209 washed with 5% (w/v) goat serum. Subsequently, cells were collected by centrifugation
210 (1500 rpm, 5 min), the supernatant was removed and cells were incubated with the primary
211 antibody against CXCR7 (rabbit polyclonal anti-CXCR7, Abcam, dilution of 1:200,
212 Cambridge, UK) on ice for 30 min. Incubation with the corresponding non-specific

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

immunoglobulin G (rabbit IgG, dilution of 1:2000) served as negative control. Following two washing steps in 5% (w/v) goat serum, cells were incubated on ice with an Alexa fluor 488-linked secondary antibody (goat anti-rabbit Alexa fluor 488, Life Technologies, Darmstadt, Germany, diluted in 5% goat serum for 30 min. Finally, stained cells were analysed using a BD FACSAria II (Becton Dickinson).

Immunofluorescence staining of testicular single cells

Fixed testicular single cells on poly L-lysine coverslips were rinsed with PBS. Cells were then incubated with 5% (w/v) BSA at RT for 30 min. Subsequently, primary antibodies against CXCR7 (rabbit polyclonal anti-CXCR7, Abcam, dilution of 1:400, Cambridge, UK) were applied and cells were incubated in a humid chamber at 4 °C for 1 hr. Following two washing steps in PBS, the appropriate Alexa fluor 546-linked secondary antibody, diluted in TBS/5% BSA, was applied for 30 min at RT in the dark. Incubation with non-specific immunoglobulin G (rabbit IgG, dilution of 1:2000) was performed as negative control. Cells were counterstained with Hoechst (Sigma-Aldrich, Steinheim, Germany).

Histological sample evaluations and image acquisition

Samples were analyzed using an Olympus BX61 microscope (Melville, NY, USA) with an attached Retiga 400R camera (QImaging, Burnaby, BC, Canada). All images were acquired digitally using cellSens imaging software (Olympus, Münster, Germany) and were processed using Adobe Photoshop CS2 9.0 (Adobe Systems, CA, USA).

236 Statistical analysis

237 Statistical analyses were performed using Graph Pad Prism 5 (Graph Pad Software, USA).

238 For analysis of qRT-PCR results the non-parametric Kruskal-Wallis test was performed
239 followed by the Dunn's multiple comparison test. For morphometric analyses the Student's t-
240 test was applied. For these two tests, significant differences between the groups are marked
241 with * $P < 0.05$.

242 Significant differences between the testis weights were determined using the "One way
243 analysis of variance" followed by the "Tukey's multiple comparison test". Groups marked with
244 different letters (a, b, c, d) are significantly different from each other ($P < 0.001$).

245

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Testicular development, testis growth and expression pattern of CXCL12, CXCL11 and CXCR7 in postnatal marmoset testes.

Testicular tissues from four phases of testicular development (16 wpp, 32 wpp, 52 wpp and > 80 wpp) were selected to characterize the expression pattern of CXCR7, CXCL12 and CXCL11 in the marmoset monkey.

Histological examination of testicular tissues of marmoset monkeys at the age of 16 wpp (Fig. 1a) revealed the presence of gonocytes, pre-spermatogonia and spermatogonia. At the age of 32 wpp, differentiated germ cells up to spermatocytes were detected in seminiferous tubules of marmoset monkeys (Fig. 1b). At the age of 52 wpp (Fig. 1c), round as well as elongated spermatids were observed in the seminiferous tubules. Finally, spermatogenesis was completed in the seminiferous tubules of marmoset monkeys at the age of > 80 wpp (Fig. 1d). In these age groups, testis weights (Fig. 1e) increased significantly from 36.4 ± 6.3 mg (16 wpp) via 181.9 ± 35.7 mg (32 wpp) and 376.7 ± 72.0 mg (52 wpp) to 481.9 ± 36.5 mg (> 80 wpp).

Relative expression levels of CXCR7 (n = 4 / group), CXCL12 (n = 4 / group) and CXCL11 (n = 4 / group) during postnatal testis development in the marmoset were determined using qRT-PCR analysis. Expression of CXCL12 was high in marmoset testes at the age of 16 wpp (0.214445 ± 0.112887) and then decreased significantly (0.088170 ± 0.012248 ; 2.4-fold; 52 wpp) during testicular development (Fig.1f). Compared to CXCL12, expression of CXCL11 was 56-fold (16 wpp), 76-fold (32 wpp), 21-fold (52 wpp) and 23-fold lower (> 80 wpp) and did not change significantly during postnatal testis development (Fig. 1g). Expression of the receptor CXCR7 (Fig. 1h) was high in marmoset testes at the age of 16 wpp (0.368448 ± 0.094921) and decreased significantly (0.092540 ± 0.013916 ; 4-fold; > 80 wpp) during postnatal development.

271 **CXCL12 and CXCR7 protein expression in postnatal marmoset testes**

272 Immunohistochemistry was performed to determine the expression of the homeostatic
273 chemokine CXCL12 and the corresponding receptor CXCR7 during postnatal testis
274 development in marmoset monkeys.

275 In the testes of marmoset monkeys at the age of 16 *wpp* (Fig. 2a), 32 *wpp* (Fig. 2c, d), 52
276 *wpp* (Fig. 2e, f) and > 80 *wpp* (Fig. 2g, h), CXCL12 was exclusively localized in the
277 cytoplasm of Sertoli cells. In contrast, expression of CXCR7 was restricted to the germ cell
278 population. More specifically, in marmoset testes at the age of 16 *wpp* the expression of
279 CXCR7 was found in a subset of gonocytes (Fig. 3c), pre-spermatogonia (Fig. 3e) and
280 spermatogonia (Fig. 3g). Furthermore, morphometric analyses revealed that the CXCR7
281 labelling index of the entire germ cell population was $69.9 \pm 9.0\%$ (Fig. 3i), which comprised
282 CXCR7-positive gonocytes (Fig. 3c), pre-spermatogonia (Fig. 3f) and spermatogonia (Fig.
283 3h) with the indices of $36.8 \pm 13.7\%$, $18.0 \pm 4.1\%$ and $15.1 \pm 9.8\%$, respectively (Fig. 3i).
284 Double immunofluorescence stainings (Fig. 4a-l) revealed subpopulations of MAGE-
285 A4⁺/CXCR7⁺ and MAGE-A4⁺/CXCR7⁻ gonocytes (Fig. 4d, e, f), pre-spermatogonia (Fig. 4g, h,
286 i) and spermatogonia (Fig. 4j, k, l). However, subsets of MAGE-A4⁺/CXCR7⁻ pre-
287 spermatogonia (Fig. 4g, h, i) were only rarely observed in tubules of marmoset testes at the
288 age of 16 *wpp*.

289 During postnatal development CXCR7 expression persisted in premeiotic germ cells located
290 at the basement membrane (Fig. 5a - f). CXCR7-positive A_{pale} (Fig. 5b, d) and A_{dark} (Fig. 5f)
291 spermatogonia were identified by morphological evaluation in marmoset monkeys at the age
292 of 32 *wpp*, 52 *wpp* and > 80 *wpp*.

293 Furthermore, adult marmoset testes (n = 3) were enzymatically digested and isolated single
294 cells were subjected to haematoxylin staining which showed cell types with different sizes
295 and morphology (Fig. 5g). Subsequent, flow cytometric assessment revealed expression of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

CXCR7 on the cell surface of $4.7 \pm 2.7\%$ of cells in the single cell suspension obtained from adult marmoset testes (Table S2) and CXCR7 protein expression was confirmed by immunofluorescence staining of cells fixed on poly L-lysine slides (Fig. 5h). The qualitative observations of CXCR7-positive cells following immunofluorescence staining were consistent with the data obtained by flow cytometry.

CXCR7 protein profiles in postnatal human testes

A panel of rare postnatal human testis samples (2 weeks, 6 months and adult) was immunohistochemically stained and revealed that a subset of gonocytes, pre-spermatogonia as well as spermatogonia was positive for CXCR7 in immature human testes (Fig. 6a-c). Moreover, the expression of CXCR7 in adult human testes (Fig. 6d) was found in premeiotic germ cells at the basement membrane.

Discussion

We have illustrated a schematic drawing of the localization patterns of CXCL12 and CXCR7 which are pointing to a crucial role of this chemokine/receptor pair in the developing primate testis (Fig. 7). Interestingly, transcript levels of CXCL12 were constitutively expressed and immunohistochemical analyses further demonstrated protein expression of CXCL12 in the cytoplasm of Sertoli cells at all stages of postnatal testicular development which indicated a non-maturation-dependent expression of the homeostatic chemokine CXCL12 in the developing marmoset testis. After birth, the highest number of Sertoli cells per testis is established at around 4 month of age. Afterwards, the Sertoli cell population decreases and after 8 month of age Sertoli cells become morphologically mature (e.g. establishment of tight-junctions between adjacent Sertoli cells, formation of lumen in the seminiferous tubules) in

1
2
3
4 320 marmoset testes (Sharpe et al. 2003a; Li et al. 2005). We therefore assume this
5
6 321 phenomenon to explain the significantly decreased transcript levels of *CXCL12* in the
7
8 322 developing marmoset testis. The Sertoli cell-specific expression pattern of *CXCL12*
9
10 323 described in this study is in line with previous mouse (Payne et al. 2010; Yang et al. 2013,
11
12 324 Westernströer et al. 2014) and human data (Gilbert et al. 2009). Here we describe for
13
14 325 marmosets a decline of *CXCL12* from week 16-52 (prepuberty to postpuberty). This mRNA
15
16 326 pattern is different from that previously observed in mice. The mRNA levels of *Cxcl12* in
17
18 327 developing mouse testes remained constant from birth until day 21 and then increased until
19
20 328 adulthood (Westernströer et al. 2014). The two methods for quantification of mRNA are
21
22 329 different (mouse: luciferase based normalization; marmoset: reference gene based
23
24 330 normalization) and the reported levels can therefore not be easily compared. However in
25
26 331 mice and marmosets transcripts for *CXCL12* are detectable and appear to be modified
27
28 332 throughout pubertal development. Current results therefore suggest that the action of
29
30 333 *CXCL12* is evolutionarily conserved. Functions of *CXCL12* have been described to be organ
31
32 334 or tissue specific. This includes a role for cell proliferation or survival (Kortesidis et al. 2005;
33
34 335 Burns et al. 2006; Mazzinghi et al. 2008), differentiation (Lazarini et al. 2003), adhesion
35
36 336 (Burns et al. 2006; Mazzinghi et al. 2008) and the regulation of cell migration (Bhakta et al.
37
38 337 2006; Mazzinghi et al. 2008). Regarding germ cells, *CXCL12* has been shown to be secreted
39
40 338 by the somatic environment thereby facilitating the colonization of the gonads by PGCs
41
42 339 during mouse embryonic development (Doitsidou et al. 2002; Ara et al. 2003; Molyneaux et
43
44 340 al. 2003; Dambly-Chaudiere et al. 2007; Boldajipour et al. 2008). Furthermore, recent studies
45
46 341 in neonatal and adult mice revealed that *CXCL12* is required for maintenance of
47
48 342 spermatogonial stem cells (SSCs) within their niches as well as for the homing and
49
50 343 colonization process of SSCs into their niches following germ cell transplantation assays
51
52 344 (Payne et al. 2010; Kanatsu-Shinohara et al. 2012; Yang et al. 2013). Moreover, we
53
54 345 previously reported significantly increased transcript levels of *Cxcl12* following germ cell
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

depletion in mice and suggested that Cxcl12 may stimulate the spermatogenic repopulation of the seminiferous tubules during testicular recovery (Westernströer et al. 2014).

Whereas inflammatory chemokines (e.g. the C-X-C chemokine CXCL11) are mainly involved in the recruitment of leukocytes to inflamed tissue sites, homeostatic chemokines (e.g. the C-X-C chemokine CXCL12) are constitutively expressed in organs and have important functions in stem cell homing, tissue homeostasis and repair. Indications for testicular inflammation (e.g. massive cell infiltration by neutrophils, lymphocytes, macrophages and mast cells in the interstitium) (Guazzone et al. 2009) were not seen in the analyzed testes. Moreover, in contrast to CXCL12, only low transcript levels of CXCL11 were detected in developing marmoset testes which confirmed that inflammatory C-X-C chemokines, especially CXCL11 as a secondary ligand for CXCR7, is likely not of functional relevance for healthy marmoset testes.

Transcript levels of CXCR7 were also constitutively expressed and immunohistochemical analyses further demonstrated that the atypical chemokine receptor CXCR7 is exclusively expressed by premeiotic germ cells (gonocytes, pre-spermatogonia, spermatogonia including A_{pale} and A_{dark} spermatogonia) throughout various phases of postnatal testicular development in the marmoset and the human testis. In agreement with these data we found that Cxcr7 protein can be detected in gonocytes as well as in Sall4-positive undifferentiated spermatogonia of the developing mouse testis (Westernströer et al. 2014). Additionally, Cxcr7 was also expressed on differentiated germ cells, spermatids and interstitial cells. These results reveal an evolutionarily conserved expression pattern of CXCR7 in premeiotic germ cells in the mouse and both analysed primate species (Westernströer et al. 2014). The morphometric evaluation in immature marmoset testes has shown additionally that approximately 80% of migrating germ cells (gonocytes and pre-spermatogonia) express CXCR7. In general, recent studies in marmoset monkeys indicated that the migration of gonocytes most likely begins postnatally between birth and 4 months of age (Rune et al.

16

1991; Sharpe et al. 2003b). Interestingly, several MAGE-A4⁺/CXCR7⁺ gonocytes were observed, whereas subsets of MAGE-A4⁺/CXCR7⁺ pre-spermatogonia were only rarely seen in seminiferous tubules, suggesting that CXCR7 may be involved in migration and/or adhesion of germ cells in immature marmoset testes. In general, migration of centrally located germ cells into the basal part of the seminiferous epithelium is a cellular process critical for germ cell survival and establishment of A_{dark} and A_{pale} spermatogonia which represent the reserve and active SSC pool in adulthood in marmoset and human testis (McGuinness and Orth, 1992; Orth et al. 2000). Moreover, our morphometric evaluation in immature marmoset testes has shown that around 50% of the spermatogonia at the basement membrane express CXCR7. Furthermore, we demonstrated that A_{dark} and A_{pale} spermatogonia express CXCR7 in marmoset testes at the age of 32 *wpp*, 52 *wpp* and > 80 *wpp*. Therefore, we assume different roles for the atypical chemokine receptor CXCR7 in the developing marmoset testes. Tissue- and cell type-specific functions of CXCR7 have been described and depend on the cell type and tissue. This includes a role for cell survival or adhesion as described for example for in the breast cancer cell line MCF-7 (Burns et al. 2006) as well as in renal multipotent progenitor cells of the human kidney (Mazzeinghi et al. 2008).

In addition, using HEK-293 and MDA-MB-231 cells, it has been demonstrated that receptor heterodimerization between CXCR4 and CXCR7 may enhance CXCL12-induced cell migration (Levoye et al. 2009; Decaillot et al. 2011). Regarding the localization of CXCR4 in developing mammalian testes, different expression patterns have been shown for mouse and human (Gilbert et al. 2009; Yoon et al. 2009; Kanatsu-Shinohara et al. 2012; Yang et al. 2013, McIver et al. 2013). Whilst expression of CXCR4 was found in gonocytes, undifferentiated spermatogonia, Sertoli cells and interstitial cells of neonatal and adult mouse testes (Yoon et al. 2009; Kanatsu-Shinohara et al. 2012; Yang et al. 2013), CXCR4 protein was detected throughout the seminiferous tubules of adult human testes (Gilbert et al. 2009).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

398 So far, the expression profile and the role of CXCR4 in developing marmoset testes remain
399 unknown. Nevertheless, receptor heterodimerization between CXCR4 and CXCR7
400 throughout testicular development in primate testes cannot be excluded but needs to be
401 investigated in further studies.

402 In conclusion, we show that CXCL12/CXCR7 network is present in primate testes. The
403 localization pattern suggests its involvement during migration, survival or adhesion in
404 premeiotic germ cells.

407 **Declaration of interest**

408 The authors declare that there is no conflict of interest that could be perceived as prejudicing
409 the impartiality of the research reported.

410 **Funding**

411 The study was supported by a grant from the Center of Interdisciplinary Clinical Research
412 Muenster (IZKF; Schl2/001/13), the Innovative Medical Research of the University of Münster
413 Medical School (KO 111014) and the DFG project (SCHL 394/11-2).

415 **Acknowledgments**

416 We gratefully acknowledge Martin Heuermann and Günter Stelke (Centre of Reproductive
417 Medicine and Andrology, Münster, Germany), for excellent animal caretaking as well as
418 Nicole Terwort, Reinhild Sandhowe, Jutta Salzig and Adelheid Kersebom for tremendous

technical assistance. Flow cytometric analysis was performed at the Max-Planck-Institute for Molecular Biomedicine (MPI, Münster) with the help of Dr. Martin Stehling.

421

422 References

423 Albert S, Ehmcke J, Wistuba J, Eildermann K, Behr R, Schlatt S, Gromoll J (2010) Germ cell
424 dynamics in the testis of the postnatal common marmoset monkey (*Callithrix jacchus*).
425 *Reproduction* 140:733-742

426 Albert S, Wistuba J, Eildermann K, Ehmcke J, Schlatt S, Gromoll J, Kossack N (2012)
427 Comparative marker analysis after isolation and culture of testicular cells from the immature
428 marmoset. *Cells Tissues Organs* 196:543-554

429 Ara T, Nakamura Y, Egawa T, Sugiyama T, Abe K, Kishimoto T, Matsui Y, Nagasawa T
430 (2003) Impaired colonization of the gonads by primordial germ cells in mice lacking a
431 chemokine, stromal cell-derived factor-1 (SDF-1). *Proc Natl Acad Sci U S A* 100:5319-5323

432 Balabanian K, Lagane B, Infantino S, Chow KY, Harriague J, Moepps B, Arenzana-
433 Seisdedos F, Thelen M, Bachelier F (2005) The chemokine SDF-1/CXCL12 binds to and
434 signals through the orphan receptor RDC1 in T lymphocytes. *J Biol Chem* 280:35760-35766

435 Bergmann M, Kliesch S (2010) Testicular biopsy and histology. In: Nieschlag E (ed)
436 *Andrology Male Reproductive Health and Dysfunction*. Springer-Verlag Berlin Heidelberg, pp
437 155-167

438 Bhakta S, Hong P, Koc O (2006) The surface adhesion molecule CXCR4 stimulates
439 mesenchymal stem cell migration to stromal cell-derived factor-1 in vitro but does not
440 decrease apoptosis under serum deprivation. *Cardiovasc Revasc Med* 7:19-24

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

441 Boldajipour B, Mahabaleshwar H, Kardash E, Reichman-Fried M, Blaser H, Minina S, Wilson
442 D, Xu Q, Raz E (2008) Control of chemokine-guided cell migration by ligand sequestration.
443 Cell 132:463-473

444 Brinkworth MH, Weinbauer GF, Schlatt S, Nieschlag E (1995) Identification of male germ
445 cells undergoing apoptosis in adult rats. J Reprod Fertil 105:25-33

446 Burns JM, Summers BC, Wang Y, Melikian A, Berahovich R, Miao Z, Penfold ME, Sunshine
447 MJ, Littman DR, Kuo CJ, Wei K, McMaster BE, Wright K, Howard MC, Schall TJ (2006) A
448 novel chemokine receptor for SDF-1 and I-TAC involved in cell survival, cell adhesion, and
449 tumor development. J Exp Med 203:2201-2213

450 Burns JM, Summers BC, Wang Y, Melikian A, Berahovich R, Miao Z, Penfold ME, Sunshine
451 MJ, Littman DR, Kuo CJ, Wei K, McMaster BE, Wright K, Howard MC, Schall TJ (2006) A
452 novel chemokine receptor for SDF-1 and I-TAC involved in cell survival, cell adhesion, and
453 tumor development. J Exp Med 203:2201-2213

454 Cancellieri C, Vacchini A, Locati M, Bonecchi R, Borroni EM (2013) Atypical chemokine
455 receptors: from silence to sound. Biochem Soc Trans 41:231-236

456 Comerford I, Litchfield W, Harata-Lee Y, Nibbs RJ, McColl SR (2007) Regulation of
457 chemotactic networks by 'atypical' receptors. Bioessays 29:237-247

458 Dambly-Chaudiere C, Cubedo N, Ghysen A (2007) Control of cell migration in the
459 development of the posterior lateral line: antagonistic interactions between the chemokine
460 receptors CXCR4 and CXCR7/RDC1. BMC Dev Biol 7:23

461 Decaillot FM, Kazmi MA, Lin Y, Ray-Saha S, Sakmar TP, Sachdev P (2011) CXCR7/CXCR4
462 heterodimer constitutively recruits beta-arrestin to enhance cell migration. J Biol Chem
463 286:32188-32197

- Doitsidou M, Reichman-Fried M, Stebler J, Koprunner M, Dorries J, Meyer D, Esguerra CV, Leung T, Raz E (2002) Guidance of primordial germ cell migration by the chemokine SDF-1. *Cell* 111:647-659
- Eva C, Sprengel R (1993) A novel putative G protein-coupled receptor highly expressed in lung and testis. *DNA Cell Biol* 12:393-399
- Gilbert DC, Chandler I, McIntyre A, Goddard NC, Gabe R, Huddart RA, Shipley J (2009) Clinical and biological significance of CXCL12 and CXCR4 expression in adult testes and germ cell tumours of adults and adolescents. *J Pathol* 217:94-102
- Guazzone VA, Jacobo P, Theas MS, Lustig L (2009) Cytokines and chemokines in testicular inflammation: A brief review. *Microsc Res Tech* 72:620-628
- Kanatsu-Shinohara M, Inoue K, Takashima S, Takehashi M, Ogonuki N, Morimoto H, Nagasawa T, Ogura A, Shinohara T (2012) Reconstitution of mouse spermatogonial stem cell niches in culture. *Cell Stem Cell* 11:567-578
- Kortesidis A, Zannettino A, Isenmann S, Shi S, Lapidot T, Gronthos S (2005) Stromal-derived factor-1 promotes the growth, survival, and development of human bone marrow stromal stem cells. *Blood* 105:3793-3801
- Lazarini F, Tham TN, Casanova P, Arenzana-Seisdedos F, Dubois-Dalcq M (2003) Role of the alpha-chemokine stromal cell-derived factor (SDF-1) in the developing and mature central nervous system. *Glia* 42:139-148
- Levoye A, Balabanian K, Baleux F, Bachelier F, Lagane B (2009) CXCR7 heterodimerizes with CXCR4 and regulates CXCL12-mediated G protein signaling. *Blood* 113:6085-6093
- Li LH, Donald JM, Golub MS (2005) Review on testicular development, structure, function, and regulation in common marmoset. *Birth Defects Res B Dev Reprod Toxicol* 74:450-469

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

487 Lin ZY, Imamura M, Sano C, Nakajima R, Suzuki T, Yamadera R, Takehara Y, Okano HJ,
488 Sasaki E, Okano H (2012) Molecular signatures to define spermatogenic cells in common
489 marmoset (*Callithrix jacchus*). *Reproduction* 143:597-609

490 Liu H, Xue W, Ge G, Luo X, Li Y, Xiang H, Ding X, Tian P, Tian X (2010) Hypoxic
491 preconditioning advances CXCR4 and CXCR7 expression by activating HIF-1alpha in MSCs.
492 *Biochem Biophys Res Commun* 401:509-515

493 Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time
494 quantitative PCR and the 2⁻(Delta Delta C(T)) Method. *Methods* 25:402-408

495 Luker K, Gupta M, Luker G (2009) Bioluminescent CXCL12 fusion protein for cellular studies
496 of CXCR4 and CXCR7. *BioTechniques* 47:625-632

497 Luker K, Gupta M, Luker G (2009) Bioluminescent CXCL12 fusion protein for cellular studies
498 of CXCR4 and CXCR7. *BioTechniques* 47:625-632

499 Mahabaleshwar H, Boldajipour B, Raz E (2008) Killing the messenger: The role of CXCR7 in
500 regulating primordial germ cell migration. *Cell Adh Migr* 2:69-70

501 Mahabaleshwar H, Tarbashevich K, Nowak M, Brand M, Raz E (2012) Beta-Arrestin Control
502 of Late Endosomal Sorting Facilitates Decoy Receptor Function and Chemokine Gradient
503 Formation. *Development* 139:2897-2902

504 Mazinghi B, Ronconi E, Lazzeri E, Sagrinati C, Ballerini L, Angelotti ML, Parente E,
505 Mancina R, Netti GS, Becherucci F, Gacci M, Carini M, Gesualdo L, Rotondi M, Maggi E,
506 Lasagni L, Serio M, Romagnani S, Romagnani P (2008) Essential but differential role for
507 CXCR4 and CXCR7 in the therapeutic homing of human renal progenitor cells. *J Exp Med*
508 205:479-490

- 509 McGuinness MP, Orth JM (1992) Gonocytes of male rats resume migratory activity
510 postnatally. *Eur J Cell Biol* 59:196-210
- 511 McIver SC, Loveland KL, Roman SD, Nixon B, Kitazawa R, McLaughlin EA (2013) The
512 chemokine CXCL12 and its receptor CXCR4 are implicated in human seminoma metastasis.
513 *Andrology* 1:517-529
- 514 McKinnell C, Mitchell RT, Morris K, Anderson RA, Kelnar CJ, Wallace WH, Sharpe RM
515 (2013) Perinatal germ cell development and differentiation in the male marmoset (*Callithrix*
516 *jacchus*): similarities with the human and differences from the rat. *Hum Reprod* 28:886-896
- 517 Mitchell RT, Cowan G, Morris KD, Anderson RA, Fraser HM, McKenzie KJ, Wallace WH,
518 Kelnar CJ, Saunders PT, Sharpe RM (2008) Germ cell differentiation in the marmoset
519 (*Callithrix jacchus*) during fetal and neonatal life closely parallels that in the human. *Hum*
520 *Reprod* 23:2755-2765
- 521 Molyneaux KA, Zinszner H, Kunwar PS, Schaible K, Stebler J, Sunshine MJ, O'Brien W, Raz
522 E, Littman D, Wylie C, Lehmann R (2003) The chemokine SDF1/CXCL12 and its receptor
523 CXCR4 regulate mouse germ cell migration and survival. *Development* 130:4279-4286
- 524 Nibbs RJ, Graham GJ (2013) Immune regulation by atypical chemokine receptors. *Nat Rev*
525 *Immunol* 13:815-829
- 526 Orth JM, Jester WF, Li LH, Laslett AL (2000) Gonocyte-Sertoli cell interactions during
527 development of the neonatal rodent testis. *Curr Top Dev Biol* 50:103-124
- 528 Payne CJ, Gallagher SJ, Foreman O, Dannenberg JH, Depinho RA, Braun RE (2010) Sin3a
529 is required by sertoli cells to establish a niche for undifferentiated spermatogonia, germ cell
530 tumors, and spermatid elongation. *Stem Cells* 28:1424-1434

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

531 Rajagopal S, Kim J, Ahn S, Craig S, Lam CM, Gerard NP, Gerard C, Lefkowitz RJ (2010)
532 Beta-arrestin- but not G protein-mediated signaling by the "decoy" receptor CXCR7. Proc
533 Natl Acad Sci U S A 107:628-632

534 Rune GM, de Souza P, Merker HJ (1991) Ultrastructural and histochemical characterization
535 of marmoset (*Callithrix jacchus*) Leydig cells during postnatal development. Anat Embryol
536 (Berl) 183:179-191

537 Sanchez-Martin L, Sanchez-Mateos P, Cabanas C (2013) CXCR7 impact on CXCL12
538 biology and disease. Trends in Molecular Medicine 19:12-22

539 Sharpe RM, McKinnell C, Kivlin C, Fisher JS (2003a) Proliferation and functional maturation
540 of Sertoli cells, and their relevance to disorders of testis function in adulthood. Reproduction
541 125:769-784

542 Sharpe RM, Fraser HM, Brougham MF, McKinnell C, Morris KD, Kelnar CJ, Wallace WH,
543 Walker M (2003b) Role of the neonatal period of pituitary-testicular activity in germ cell
544 proliferation and differentiation in the primate testis. Hum Reprod 18:2110-2117

545 Staton AA, Knaut H, Giraldez AJ (2011) miRNA regulation of Sdf1 chemokine signaling
546 provides genetic robustness to germ cell migration. Nat Genet 43:204-211

547 Ulvmar MH, Hub E, Rot A (2011) Atypical chemokine receptors. Exp Cell Res 317:556-568

548 Westernströer B, Terwort N, Ehmcke J, Wistuba J, Schlatt S, Neuhaus N (2014) Profiling of
549 Cxcl12 receptors, Cxcr4 and Cxcr7 in murine testis development and a spermatogenic
550 depletion model indicates a role for Cxcr7 in controlling Cxcl12 activity. PLoS ONE
551 9(12):e112598

552 Yang QE, Kim D, Kaucher A, Oatley MJ, Oatley JM (2013) CXCL12-CXCR4 signaling is
553 required for the maintenance of mouse spermatogonial stem cells. J Cell Sci 126:1009-1020

554 Yoon KA, Chae YM, Cho JY (2009) FGF2 stimulates SDF-1 expression through the Erm
555 transcription factor in Sertoli cells. J Cell Physiol 220:245-256

556

557

For Peer Review

Figures:

Fig. 1 Marmoset monkey testes after birth - age-related changes in histology, weight and transcript levels. Micrographs showing PAS stained tissue sections of marmoset testes at the age of 16 *wpp* (*wpp* = weeks *post partum*; a), 32 *wpp* (b), 52 *wpp* (c) and > 80 *wpp* (d). Gonocytes (arrowhead), pre-spermatogonia (white arrow) and spermatogonia (black arrow) are the predominant germ cells in marmoset testes at the age of 16 *wpp* (a). Meiotic cells (primary spermatocytes) were present in the seminiferous tubules of marmosets at the age of 32 *wpp* (b), whilst germ cells up to the stage of elongated spermatids were detected in the seminiferous epithelium of marmoset monkeys at the age of 52 *wpp* (c). Finally, spermatogenesis was complete in all marmosets at the age of > 80 *wpp* (d). Scale bars represent 50 μ m. Changes in testis weight (e) are shown as mean $\pm$ SEM. Groups marked with different letters (a, b, c, d) are significantly different from each other ($p < 0.001$). Relative mRNA levels of *CXCL12* (f), *CXCL11* (g) and *CXCR7* (h) in marmoset testes at the age of 16 *wpp*, 32 *wpp*, 52 *wpp* and > 80 *wpp* were normalized using the housekeeping gene *TOP1* (2^{- Δ Ct}) method). Results are shown as mean $\pm$ SEM and significant differences between the groups are marked with * $P < 0.05$.

Fig. 2 Immunohistochemical detection of *CXCL12* in marmoset testes at the age of 16 *wpp* (*wpp* = weeks *post partum*; a), 32 *wpp* (c, d), 52 *wpp* (e, f) and > 80 *wpp* (g, h). Incubation with non-specific IgG antibody was performed as control and a representative image of a marmoset testis at the age of 16 *wpp* is shown in (b). During testicular development, the chemokine *CXCL12* is exclusively localized in the cytoplasm of Sertoli cells (white arrow). Neither relatively undifferentiated germ cells (gonocytes, arrow head; spermatogonia, black arrow) nor differentiating germ cells showed expression of *CXCL12*. Scale bars represent 20 μ m.

Fig. 3 Immunohistochemical detection of CXCR7 and labelling index of CXCR7-positive cells in marmoset testes at the age of 16 *wpp* (*wpp* = weeks *post partum*). A representative image is showing CXCR7 expression in marmoset testes at the age of 16 *wpp* (a). The staining revealed that subsets of germ cells express CXCR7 (black arrow), whilst the remainder are negative (arrow head) for CXCR7. Sertoli cells (white arrow) in the seminiferous tubules showed no expression of CXCR7. Corresponding IgG antibody was used as negative control and a representative image is shown in (b). High magnification of CXCR7 stained tissue sections revealed CXCR7-positive and negative gonocytes (c, d), pre-spermatogonia (e, f) and spermatogonia (g, h). Scale bars represent 10 μ m. Scores of CXCR7-positive and CXCR7-negative gonocytes (Gon+/Gon-), pre-spermatogonia (pre-SPG+/pre-SPG-) and spermatogonia (SPG+/SPG-) in marmoset testes at the age of 16 *wpp* are shown in (i). Results are shown as mean $\pm$ SEM and significant differences between the groups are marked with * $P < 0.05$.

Fig. 4 Representative images for expression of MAGE-A4 (green) and CXCR7 (red) in testis sections from marmosets at the age of 16 *wpp* (a - c). MAGE-A4 expression is seen in the majority of germ cells including gonocytes (d), pre-spermatogonia (g) and spermatogonia (j). Subpopulations of MAGE-A4⁺/CXCR7⁺ (arrow heads) and/or MAGE-A4⁺/CXCR7⁻ (arrows) gonocytes (d - f), pre-spermatogonia (g - i) as well as spermatogonia (j - l) can be identified. Note that MAGE-A4⁺/CXCR7⁻ pre-spermatogonia are occasionally seen in the seminiferous tubules of marmoset testes at the age of 16 *wpp*. All sections were counterstained with Hoechst (blue). Dotted lines indicate the basement membrane of the seminiferous tubules. Scale bars represent 10 μ m.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 5 Immunohistochemical detection of CXCR7 in marmoset testes at the age of 32 *wpp* (*wpp* = weeks *post partum*; a, b), 52 *wpp* (c, d) and > 80 *wpp* (e, f). In marmoset testes at the age of 32 *wpp*, 52 *wpp* and > 80 *wpp*, premeiotic germ cells (black arrow; including A_{pale} and A_{dark} spermatogonia) located at the basement membrane are labelled for CXCR7 whereas Sertoli cells (white arrow) showed no expression for CXCR7. Scale bars represent 20 µm (a – f). Following enzymatic digestion of adult marmoset testes, haematoxylin-stained single cells showed cell types with different morphology (g) as well as immunofluorescence staining revealed CXCR7-positive cells in this single cell suspension (h). As negative control, incubation with nonspecific IgG antibody was performed and a representative image is shown in (i). Scale bars represent 50 µm (g), 15 µm (h) and 60 µm (i).

Fig. 6 Immunohistochemical detection of CXCR7 in human testes at 2 weeks (a) and 6 months after birth (c) as well as in adult (d) human testes. Corresponding IgG antibody was used as negative control and a representative image of an immature testis (2 weeks) is shown in (b). In immature human testes (a; c), gonocytes (arrow head, no contact with the basement membrane), pre-spermatogonia (white arrows, point contact with the basement membrane) and spermatogonia (black arrows, full contact with the basement membrane) were labelled for CXCR7. In adult human testes (d), premeiotic germ cells (red arrow) located at the basement membrane were labelled for CXCR7. Scale bars represent 20 µm.

Fig. 7 Schematic drawing of CXCR7 (red) and CXCL12 (green) expression in seminiferous tubules of marmoset testes at the ages of 16 *wpp* (a), 32 *wpp* (b), 52 *wpp* (c) and > 80 *wpp* (*wpp* = weeks *post partum*, d). In marmoset testes at the age of 16 *wpp*, CXCL12 is highly expressed by Sertoli cells whereas the atypical chemokine receptor CXCR7 is expressed by few gonocytes, pre-spermatogonia and spermatogonia (a). In the seminiferous epithelium of

632 marmoset testes at the ages of 32 wpp (b), 52 wpp (c) and > 80 wpp (d), CXCL12 is still
633 highly expressed by Sertoli cells whereas CXCR7 is expressed by premeitotic germ cells,
634 including A_{pale} and A_{dark} spermatogonia. The expression pattern of CXCL12 and CXCR7
635 indicates a CXCL12/CXCR7 network in primate testes.

636

637

219x372mm (300 x 300 DPI)

219x414mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

219x588mm (300 x 300 DPI)

219x588mm (300 x 300 DPI)

219x310mm (300 x 300 DPI)

219x220mm (300 x 300 DPI)

325x228mm (300 x 300 DPI)

Supplemental Table 1: Westernströer *et al.*

Primer sequences used for real-time quantitative PCR		
Target	Forward Primer (5' - 3')	Reverse Primer (5' - 3')
<i>calCXCR7</i>	CGTCATCGGCATGATTGC	TCTTGGCCTGGACGTTTAC
<i>calCXCR4</i>	GGAGGGCATCAGTATATACACTTCAG	AAGCAGGGTTCCTTTATGGAGTC
<i>calCXCL12</i>	CACTCCAAACTGTGCCCTTCA	TGCACACTTGTCTGTTGTTGTTCT
<i>calCXCL11</i>	GCTGTGATATTCTGCGCTATGG	AGACAGCGTCCTCTGTTGAACA
<i>calTOP1</i>	CCGAAGGGACCTGAAAAGTG	CAGTCTCTGCACAGCCTTCTTCT
<i>cal</i> = <i>Callitrix jacchus</i>		

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplemental Table 2: Westernströer *et al.*

Flow cytometric assessment of CXCR7-positive cells in adult marmoset testes (n = 3)		
Animal Nr.:	Alexa ⁺ [%]	
	CXCR7	Rabbit IgG
1	7.6	0.5
2	2.4	0.9
3	4.0	0.3
Average ± SD	4.7 ± 2.7	0.6 ± 0.3
SD = Standard derivation		