THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Curiosity and the CAT: Considering Cognitive Analytic Therapy as an intervention for psychosis

Citation for published version:

Taylor, PJ, Perry, A, Hutton, P, Seddon, C & Tan, R 2014, 'Curiosity and the CAT: Considering Cognitive Analytic Therapy as an intervention for psychosis', *Psychosis: Psychological, Social and Integrative Approaches*, vol. 7, no. 3, pp. 276-278. <https://doi.org/10.1080/17522439.2014.956785>

Digital Object Identifier (DOI):

[10.1080/17522439.2014.956785](https://doi.org/10.1080/17522439.2014.956785)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Psychosis: Psychological, Social and Integrative Approaches

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Abstract

In the interests of providing patient choice and developing more effective interventions for people struggling with psychosis it is important that alternative talking therapies are explored. Cognitive Analytic Therapy is an integrative therapy which has specific features that that may be particularly suited to psychosis. The current report briefly outlines this approach and its potential benefits.

Keyword: Cognitive Analytic Therapy; Psychosis; Intervention

Curiosity and the CAT: Considering Cognitive Analytic Therapy as an Intervention for Psychosis

There is a growing use of “talking therapies” in treating psychotic disorders. Cognitive Behavioural Therapy (CBT) is arguably the most widely studied talking therapy for treating psychosis, and is recommended for use in UK guidelines (National Institute for Health and Care Excellence, 2014). However, CBT does not work for everyone and effect sizes are typically small to moderate (Sarin, Wallin & Widerlov, 2011; Wykes, Steel, Everitt & Tarrier, 2008). Moreover, different psychosocial interventions for psychosis may impact upon different outcomes (Turner, van der Gaag, Karyotaki & Cuijpers, 2014). The importance of patient choice within the UK has also been emphasised (Department of Health, 2012). With these issues in mind it is vital that we remain curious about alternative therapies for those with experiences of psychosis.

Cognitive Analytic Therapy (CAT; Ryle & Kerr, 2002) represents an alternative talking therapy. CAT is an integrative therapy, which draws upon many of the principles of cognitive therapy, but combines these with ideas from personal construct, object relations and social developmental theory. CAT involves a process of mapping out historical interpersonal experiences and identifying enactments of similar relationship patterns in a person’s day-to-day life. A theoretical CAT model of psychosis has been developed and illustrated in individual case studies (Kerr et al., 2003; Kerr, Crowley & Beard, 2006; Perry, 2012). This model provides an account of how certain ways of relating to self and others (‘Reciprocal Roles’ [RRs]) develop over time and are enacted in day-to-day life. The formation of these RRs is largely socially mediated. Within this model, psychotic experiences can be understood as representing disordered or extreme expressions of RRs (Kerr, Birkett & Chanen, 2003). For example, paranoia may represent an underlying RR of ‘victimised and targeted’ in relation to an ‘abusive and threatening’ other (possibly a product of early abuse or hostile

parenting). A malicious voice may result from early experiences of a critical other that is first *internalised* as a RR and then *externalised* (or dissociated) as a voice perhaps because it was too painful (Perry, 2012). CAT is a *fundamentally* interpersonal model, adopting a dialogical perspective of the self (i.e., that the self emerges and develops through the ongoing interaction with others) that emphasises social, cultural and interpersonal influences, and so differs to interpersonal varieties of CBT (Ryle & Kerr, 2002).

CAT has a number of strengths for working with experiences of psychosis: 1) The emphasis in CAT upon interpersonal relationships is relevant as many of the difficulties that are considered indicative of psychosis are both inter and intrapersonal in nature (e.g., paranoia, voices; Berry, Wearden, Barrowclough & Liversidge, 2006). 2) The dialogical component (making links between ‘self-self’ and ‘self-other’ relationships) may be particularly useful in explaining some of the processes by which ‘normal’ everyday experiences, as well as more traumatic experiences, can lead to seemingly bizarre experiences. CAT-specific tools such as the reformulation letter provide a means of mapping out these links between early experiences and difficulties. 3) Viewing psychosis in terms of early interactions is consistent with the link between interpersonal trauma and psychosis (Read, van Os, Morrison & Ross, 2005). 4) CAT provides a coherent framework working with ruptures in therapy and difficulties around engagement. 5) The dialogical nature of CAT provides a means of representing stigma (fundamentally a social process), where present, as a central part of the phenomenology of the psychosis, rather than a secondary difficulty.

A Recent review suggests support for the efficacy of CAT in a variety of disorders, most often complex presentations (e.g., personality disorder; Calvert & Kellett, in press). Positive findings have emerged within a recent trial of CAT for bipolar disorder (Evans & Kellett, 2014). However, research into CAT for those with psychosis is still in its early phases. Initial evidence of the safety and acceptability of CAT for individuals with

experiences of psychosis exists from case-studies and small case-series (Graham, 1995; Kerr et al., 2001), and trials of multi-modal interventions (Gleeson et al., 2012). Based on the Medical Research Council's guidance concerning the evaluation of complex interventions, we believe the next step is to further confirm the safety and acceptability of CAT through a larger case-series of CAT for individuals with psychosis (Craig et al., 2012).

References

- Berry, K., Wearden, A., Barrowclough, C., & Liversidge, T. (2006). Attachment styles, interpersonal relationships and psychotic phenomena in a non-clinical student sample. *Personality and Individual Differences*, 41, 707-718. doi: <http://dx.doi.org/10.1016/j.paid.2006.03.009>
- Calvert, R., & Kellett, S. (in press). Cognitive analytic therapy: A review of the outcome evidence base for treatment. *Psychology & Psychotherapy*. doi: 10.1111/papt.12020
- Craig, P., Diepe, P., Macintyre, S., Michie, S., Nazareth, I., & Petticrew, M. (2008). Developing and evaluating complex interventions: New guidance. www.mrc.ac.uk/complexinterventionsguidance
- Deptment of Health. (2012). *Liberating the NHS: NO decision about me, without me*
Retrieved from https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/216980/Liberating-the-NHS-No-decision-about-me-without-me-Government-response.pdf
- Gleeson, J. F. M., Chanen, A., Cotton, S. M., Pearce, T., Newman, B., & McCutcheon, L. (2012). Treating co-occurring first-episode psychosis and borderline personality: A pilot randomized controlled trial. *Early Intervention in Psychiatry*, 6, 21-29. doi: 10.1111/j.1751-7893.2011.00306.x
- Graham, C. (1995). Dissociative psychosis: An atypical presentation and response to cognitive-analytic therapy. *Irish Journal of Psychological Medicine*, 12, 109-111.
- Evans, M. & Kellett, S. (2014). RCT in CAT and bipolar disorder. Paper presented at the ACAT annual national conference 2014, Liverpool.
- Kerr, I. B. (2001). Brief cognitive analytic therapy for post-acute manic psychosis on a psychiatric intensive care unit. *Clinical Psychology & Psychotherapy*, 8, 117-129. doi: 10.1002/cpp.251

- Kerr, I. B., Birkett, P. B., & Chanen, A. (2003). Clinical and service implications of a cognitive analytic therapy model of psychosis. *Australian & New Zealand Journal of Psychiatry*, 37, 515-523.
- Kerr, I. B., Crowley, V., & Beard, H. (2006). A cognitive analytic therapy-based approach to psychotic disorder. In J. O. Johannessen, B. V. Martindale & J. Cullberg (Eds.), *Evolving Psychosis* (pp. 172-184). London: ISPS.
- National Institute for health and Care Excellence. (2014). *Psychosis and schizophrenia in adults: Treatment and management* Retrieved from <http://publications.nice.org.uk/psychosis-and-schizophrenia-in-adults-treatment-and-management-cg178/recommendations#first-episode-psychosis-2>
- Perry, A. (2012). CAT with people who hear distressing voices. *Reformulation, summer*, 16-22.
- Read, J., van Os, J., Morrison, A. P., & Ross, C. A. (2005). Childhood trauma, psychosis and schizophrenia: A literature review with theoretical and clinical implications. *Acta Psychiatrica Scandinavica*, 112, 330-350. doi: 10.1111/j.1600-0447.2005.00634.x
- Ryle, A., & Kerr, I. B. (2002). *Introducing cognitive analytic therapy: Principles and practice*. Oxford: Wiley-Blackwell.
- Sarin, F., Wallin, L., & Widerlov, B. (2011). Cognitive behavior therapy for schizophrenia: A meta-analytical review of randomized controlled trials. *Nordic Journal of Psychiatry*, 65, 162-174. doi: 10.3109/08039488.2011.577188
- Turner, D. T., van der Gaag, M., Karyotaki, E., & Cuijpers, P. (2014). Psychological interventions for psychosis: A meta-analysis of comparative outcome studies. *American Journal of Psychiatry*, 171, 523-538. doi: 10.1176/appi.ajp.2013.13081159

Wykes, T., Steel, C., Everitt, B., & Tarrier, N. (2008). Cognitive behavior therapy for schizophrenia: Effect sizes, clinical models, and methodological rigor. *Schizophrenia Bulletin*, 34, 523-537. doi: 10.1093/schbul/sbm114