THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Preserved extra-foveal processing of object semantics in Alzheimer's disease

Citation for published version:

Cimminella, F, D'Innocenzo, G, Della Sala, S, Iavarone, A, Musella, C & Coco, MI 2022, 'Preserved extra-foveal processing of object semantics in Alzheimer's disease', *Journal of Geriatric Psychiatry and Neurology*, vol. 35, no. 3, pp. 418-433. <https://doi.org/10.1177/08919887211016056>

Digital Object Identifier (DOI):

[10.1177/08919887211016056](https://doi.org/10.1177/08919887211016056)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Geriatric Psychiatry and Neurology

Publisher Rights Statement:

The final version of this paper has been published in Journal of Geriatric Psychiatry and Neurology, Vol/Issue, Month/Year by SAGE Publications Ltd, All rights reserved. © Francesco Cimminella, Giorgia D'Innocenzo, Sergio Della Sala, Alessandro Iavarone, Caterina Musella and Moreno I. Coco, 2021. It is available at: <https://journals.sagepub.com/doi/10.1177/08919887211016056>

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Preserved Extra-Foveal Processing of Object Semantics in Alzheimer's Disease

Francesco Cimminella^{1,2*}, Giorgia D'Innocenzo^{3*}, Sergio Della Sala¹, Alessandro Iavarone⁴,
Caterina Musella⁵, Moreno I. Coco^{3,6†}

¹ Human Cognitive Neuroscience, Psychology, University of Edinburgh, Edinburgh, UK

² Laboratory of Experimental Psychology, Suor Orsola Benincasa University, Naples, Italy

³ Faculdade de Psicologia, Universidade de Lisboa, Lisbon, Portugal

⁴ Neurological Unit, CTO Hospital, AORN "Ospedali dei Colli", Naples, Italy

⁵ Associazione Italiana Malattia d'Alzheimer (AIMA sezione Campania), Naples, Italy

⁶ School of Psychology, The University of East London, London, UK

Author Note

Francesco Cimminella <https://orcid.org/0000-0001-5403-2719>

Giorgia D'Innocenzo <https://orcid.org/0000-0002-1388-2693>

Sergio Della Sala <https://orcid.org/0000-0003-4719-508X>

Alessandro Iavarone <https://orcid.org/0000-0002-9596-5811>

Caterina Musella <https://orcid.org/0000-0001-6264-7705>

Moreno I. Coco <https://orcid.org/0000-0002-2825-4200>

The data and R script to reproduce visualisations and models are available in the Open Science Framework (<https://osf.io/gyutd/>).

Correspondence should be sent to moreno.cocoi@gmail.com

* Indicates equal contribution.

Abstract

Alzheimer's disease (AD) patients underperform on a range of tasks requiring semantic processing, but it is unclear whether this impairment is due to a generalised loss of semantic knowledge or to issues in accessing and selecting such information from memory. The objective of this eye-tracking visual search study was to determine whether semantic expectancy mechanisms known to support object recognition in healthy adults are preserved in AD patients. Furthermore, as AD patients are often reported to be impaired in accessing information in extra-foveal vision, we investigated whether that was also the case in our study. Twenty AD patients and twenty age-matched controls searched for a target object among an array of distractors presented extra-foveally. The distractors were either semantically related or unrelated to the target (e.g., a car in an array with other vehicles or kitchen items). Results showed that semantically related objects were detected with more difficulty than semantically unrelated objects by both groups, but more markedly by the AD group. Participants looked earlier and for longer at the critical objects when these were semantically unrelated to the distractors. Our findings show that AD patients can process the semantics of objects and access it in extra-foveal vision. This suggests that their impairments in semantic processing may reflect difficulties in accessing semantic information rather than a generalised loss of semantic memory.

Keywords: Visual search; Alzheimer's disease; semantic processing; eye-tracking

Preserved Extra-Foveal Processing of Object Semantics in Alzheimer's Disease

A hallmark of Alzheimer's disease (AD) and degenerative cognitive deficits leading to AD (e.g., Mild Cognitive Impairment - MCI) is their detrimental impact on semantic memory (Chan et al., 1993; Daum, Riesch, Sartori, & Birbaumer, 1996; Hodges & Patterson, 1995; Martin & Fedio, 1983; Mulatti, Calia, De Caro, & Della Sala, 2014), defined as conceptually organised knowledge not tied to specific episodes (Gold & Budson, 2008; Nebes, 1989). From an early stage of the disease, AD patients present with disruptions in word finding abilities (Bandera, Della Sala, Laiacona, Luzzatti, & Spinnler, 1991; Spaan, Raaijmakers, & Jonker, 2003), name fewer pictures in object naming tasks than age-matched controls (Holmes, Fitch, & Ellis, 2006), underperform on word-to-picture matching tasks (Adlam, Bozeat, Arnold, Watson, & Hodges, 2006) and their verbal recall does not benefit from cues (Ivanou et al., 2005; Spaan, Raaijmakers, & Jonker, 2005). In addition, relative to healthy controls, AD patients produce more semantic-superordinate errors when asked to generate names of objects belonging to a specific semantic category (see Salmon et al., 1999, for a review) or they generate less words of a particular semantic category than words beginning with a particular letter (Phillips, Della Sala, & Trivelli, 1996), which suggests specific difficulties with semantic knowledge rather than generic disruptions linked to lexical retrieval (Monsch et al., 1994).

The findings reported above have at times been interpreted as reflecting a breakdown of semantic knowledge in AD (Adlam et al., 2006; Hodges, Salmon, & Butters, 1992; Salmon et al., 1999), but AD patients have shown normal semantic abilities when tested on tasks which can be performed more automatically and include a priming component. Semantic priming implies the automatic activation of information related to a processed stimulus, which results in faster responses when, for example, a stimulus is preceded by a semantically related cue compared to when the preceding cue is unrelated to the target stimulus (e.g., Gold & Budson, 2008). Studies that have used semantic priming in AD have reported levels of performance comparable to – or higher than –

80 those achieved by age-matched controls (Chertkow et al., 1994; Hernández et al., 2008 and
81 Nakamura et al., 2000).

82 The control of selective attention is also known to decline starting from the early stage of
83 AD, and this is mediated by task requirements (Baddeley, Baddeley, Bucks, & Wilcock, 2001).
84 Therefore, evidence of preserved semantic abilities in AD may be linked to tasks that can be
85 performed more automatically and require reduced attentional control (Ober, 2002; Perri et al.,
86 2003). Consistent with the distinction between automatic and controlled memory processes (Jacoby,
87 1991; Moscovitch, 1992), AD may, in fact, induce a selective impairment in the explicit access and
88 intentional retrieval of semantic knowledge (Nebes, 1992; Ober, 2002; Rogers & Friedman, 2008),
89 rather than its indiscriminate loss. Tippet and colleagues (2004), for example, tested AD patients
90 and healthy age-matched individuals on a lexical fluency task, a comparison task and a verb
91 generation task, and systematically varied the number of alternatives participants were faced with.
92 In the verb generation task, participants had to produce a verb related to a visually presented noun;
93 in the high selection condition the noun was associated with many appropriate responses (e.g., *boat*
94 *-> sail, dock, capsize*), whereas in the low selection condition the noun could only be matched with
95 one or few responses (e.g., *piano -> play*). The results of this study showed that AD patients were
96 disproportionately impaired in the high-selection conditions, but performed closely to healthy
97 controls when the demands on selective attention were reduced (Tippet et al., 2004). Similarly,
98 Rich and colleagues (2002) showed that semantic judgements of AD patients in a sorting task were
99 impaired when the retrieval context was unstructured (i.e., when the number of groups to sort items
100 into was not specified), but performed normally under more structured conditions (i.e., when
101 instead the number of groups was specified by the experimenter).

102 When investigating semantic mechanisms and their functioning in healthy and neuro-
103 pathological ageing, it is also important to consider the type of stimuli used (e.g., verbal vs. non-
104 verbal). As argued by Paivio's Dual Coding theory (Paivio, 1991; Paivio & Csapo, 1973), words
105 have a mnemonic disadvantage over pictures because they are encoded into the semantic system

106 through a verbal route only, whereas pictures rely on an additional visual route. Thus, words can be
107 mainly accessed through conceptual processing, whereas pictures can also benefit from perceptual
108 processing (O'Connor & Ally, 2010), which facilitates their subsequent retrieval¹ (Ally, McKeever,
109 et al., 2009). The mnemonic superiority of pictures over words is already present in childhood
110 (Whitehouse, Maybery, & Durkin, 2006), it becomes increasingly pronounced as we age (Ally et
111 al., 2008) and it is particularly evident in AD patients (Ally, Gold, et al., 2009b), probably due to
112 the fact that pictures allow AD patients to access semantic information that is less accessible with
113 lexical stimuli (Rich et al., 2002). In support of this, some AD patients show preserved information
114 about the use of objects that they cannot name (Bartolo, Della Sala, & Cubelli, 2016). Therefore, it
115 is possible that AD selectively impairs the top-down access to – and retrieval of – lexical
116 information, and that this impairment may account for previous findings of reduced performance on
117 semantic memory tasks that have used verbal stimuli (Cuetos, Arce, Martínez, & Ellis, 2017).

118 As such, the main objective of the present study was to determine whether AD patients
119 utilise semantic knowledge when asked to perform a visual search for pictorial stimuli. To this aim,
120 we also examined patients' eye movement behaviour, which can be taken as an implicit index of
121 semantic processing and help unveil its more automatic functioning (see Ramzaoui et al., 2018 for a
122 recent review on the topic). Eye movement responses can index memory processes even when
123 explicit conscious recollection seems to fail (Hannula et al., 2010). For instance, in visual search
124 tasks with complex scenes, target detection is affected by previous knowledge of the specific
125 contexts in which objects are more likely to occur (Henderson & Hayes, 2017; Loftus &
126 Mackworth, 1978), which indicates that semantic knowledge affects the way in which we extract
127 visual information. In particular, overt attention appears to be guided by a top-down analysis of the
128 conceptual features that are shared (or not) between the search target and the other objects in the

¹ A lengthy discussion of Dual Coding theory is beyond the scope of this study, hence we refer interested readers to previous literature on the topic (e.g., McBride and Doshier, 2002; Nelson, Reed and McEvoy 1977; or Paivio, 2014).

129 visual scene (e.g., Belke, Humphreys, Watson, Derrick, Meyer, & Telling, 2008; Nuthmann, de
130 Groot, Huettig, & Olivers, 2019; Ruotolo, Kalénine, & Bartolo, 2020). The guidance provided by
131 the object features can also be visual, as observers searching for a cued target (e.g., a red ball) can
132 preferentially look at visually similar (e.g., a red apple) than dissimilar (e.g., a yellow banana)
133 objects (Alexander & Zelinsky, 2011; Schmidt & Zelinsky, 2009). Moreover, eye movements on a
134 target object precede its explicit identification by as many as 25 fixations (Holm, Eriksson, &
135 Andersson, 2008), and longer fixations are observed when a target object embedded into a scene is
136 changed, even when the change is not consciously detected by the participant (Hollingworth,
137 Williams, & Henderson, 2001). Gaze patterns are also particularly revelatory about cognitive and
138 attentional processes in patient populations (Hannula et al., 2010; Molitor, Ko, & Ally, 2015), and
139 they can be used to discriminate between healthy older adults and individuals suffering from
140 neurodegenerative disease such as MCI and AD (Crutcher et al., 2009; Oyama et al., 2019) or to
141 identify those individuals at risk of developing such diseases (Ryan et al., 2019; Zola et al., 2013).

142 It is generally agreed that objects are recognised through the activation of their conceptual
143 representations and of related concepts sharing similar features (Chertkow et al., 1994; Kiefer &
144 Pulvermüller, 2012). It has been shown that AD patients underperform in detecting, recognising and
145 categorising objects (Laatu, Revonsuo, Jäykkä, Portin, & Rinne, 2003), which supports the
146 hypothesis of a generalised impairment in semantic memory. However, as already discussed above,
147 the semantic impairment observed in AD may more specifically relate to the explicit top-down
148 selection of a single object among a pool of co-activated semantic competitors. Evidence in support
149 of this proposition comes from a recent eye-tracking study conducted by Seckin and colleagues
150 (2016). In this study, healthy older adults, patients with a generalised object naming impairment
151 (non-semantic aphasia) and patients with a semantic variant of the object naming impairment
152 (semantic aphasia) were asked to search for a target object in an array with fifteen other distractor
153 objects placed in an imaginary circle at extra-foveal distance from the centre of the display. The
154 critical manipulation of this study was that half of the distractors belonged to the same semantic

category as the target, whereas the remaining distractors belonged to different semantic categories than that of the target. A key finding was that, although all participants tended to fixate more on the distractors that were semantically related to the target object, this behaviour was particularly pronounced in the patients with semantic aphasia. This result probably reflected the difficulty of semantic aphasia patients in isolating the target object from its competitors, as their explicit knowledge of its word denotation was blurred by the co-activated semantically related distractors.

As well as issues related to semantic processing, AD patients may display additional difficulties in visual search due to reductions in their field of focal attention (Rösler et al 2000), which may specifically disrupt processing of information in extra-foveal vision. Boucart and colleagues (2014) investigated this question in AD patients by using a categorization task (i.e., spot-the-animal) in which they presented participants with pairs of natural images displayed to the right and left side of a central fixation cross. Their results showed that AD patients were less accurate than healthy older adults, who were in turn less accurate than younger adults, in the categorization task. As argued by Boucart et al., this impairment may stem from deficits in the magnocellular pathway of AD patients (Gilmore & Whitehouse, 1995; Kergoat et al., 2002; Sartucci et al., 2010), which result in degraded extra-foveal processing of objects' features. In sum, even though mechanisms of semantic processing have often been shown to be impaired in AD patients, it is not yet clear whether the reported impairments are due to a generalised loss of semantic knowledge or whether they may specifically relate to the explicit control and access of this information. Moreover, despite evidence of reduced extra-foveal information intake in this group, it is yet to be established whether this also occurs when the semantics of the visual context is manipulated to generate a more automatic and implicit "pop-out" effect.

Therefore, the aims of the present study were twofold: (1) to establish whether processing of semantic information about objects is preserved in AD and, if so, (2) to determine whether it can occur through extra-foveal vision, as shown in our previous work on healthy younger adults (Cimminella, et al. 2020). We used the experimental paradigm we previously developed, reported in

181 Cimminella et al., which we adapted to make it suitable for use with participants with AD and
182 healthy older adults (see the Methods section for further details). In this paradigm, participants
183 search for a target object, which is either related (e.g., *a car*) or unrelated (e.g., *a banana*) to other
184 semantically homogenous distractors (e.g., *other vehicles*; see Figure 1 for an example of the
185 design). This manipulation critically differs from Seckin et al., (2016), where distractors belonging
186 to different semantic categories were presented together with semantically related distractors. We
187 previously showed that in young adults, our manipulation led to a “semantic pop-out” effect
188 whereby participants looked at a critical object earlier, and for longer, when it was semantically
189 unrelated than related to the other distractors (Cimminella et al, 2020). This finding indicated that,
190 in young adults, the rapid and complex extraction of object semantics occurs in extra-foveal vision,
191 and it influences the early allocation of overt attention (see also Nuthmann et al., 2019; and Borges
192 et al., 2020, for corroborating results in healthy older adults using naturalistic scenes).

193 In the context of a neuropathology associated with semantic impairment, such as AD, if
194 semantic processes are compromised, we should not observe any effect of semantic relatedness on
195 either manual responses or eye movements. In practice, the time taken to visually identify and
196 accurately recognize the search target should not vary depending on the semantic context in which
197 it is presented. In contrast, if semantic processes are, to a certain degree, preserved, we would
198 expect search responses to be mediated by the semantic relatedness of the objects in the array,
199 consistent with previous findings on younger adults. We reasoned that, if AD patients’ attentional
200 abilities are sufficiently preserved to enable them to process information extra-foveally, then we
201 would expect shorter latencies to first fixations onto semantically unrelated objects. This would
202 indicate that object semantics are accessed in extra-foveal vision, with parafoveal semantic
203 processing driving target selection in AD patients and healthy older adults alike. If processing also
204 continued during foveation, we would expect unrelated objects to be looked at for longer due to the
205 heightened complexity of the semantic context. Evidence of this integration cost in AD would
206 suggest that retrieving operations to identify the search object are also partly preserved. Finally, we

observed ceiling search performance on young adults in our previous study, but for older adults, a semantically related search object (e.g., a car) in target-absent trials may lead to greater memory demands, i.e., the object is not present in the context. This, in turn, may result in a higher rate of false detections due to co-activation of semantically homogenous distractors (e.g., other vehicles) that need to be suppressed.

Methods

Participants

A group of 20 patients with a diagnosis of mild to moderate AD (14 women) and a group of 20 healthy controls (9 women) took part in the study. Both patients and controls were native Italian speakers recruited from the Neurological and Stroke Unit at CTO Hospital, Naples (15 patients and 10 controls) and the Alzheimer's Disease Italian Association AIMA Campania (5 patients and 10 controls). The two groups were matched on age (AD: $M = 71.15$, $SD = 8.50$; Control: $M = 69.35$, $SD = 5.52$; $t(32.60) = .79$, $p = .43$) and years of education (AD: $M = 8.90$, $SD = 4.29$; Control: $M = 10.50$, $SD = 4.37$; $t(37.99) = -1.17$, $p = .25$). The diagnosis of AD was based on family and medical history interview and neuroimaging evidence (available for 16² out of 20 patients), according to the NINCDS-ADRDA criteria (McKhann et al., 1984). The diagnosis was also based on neuropsychological testing, including the Italian version of the mini-mental state examination (MMSE, Folstein, Folstein, & McHugh, 1975; Frisoni, Rozzini, Bianchetti, & Trabucchi, 1993), the frontal assessment battery (FAB, Dubois, Slachevsky, Litvan, & Pillon, 2000; Iavarone et al., 2004), the Rey's auditory verbal learning test (Ricci, Graef, Blundo, & Miller, 2012), word fluency (Costa et al., 2014), Raven's coloured progressive matrices (Ambra et al., 2016), freehand copying of drawings, and copying of drawings with landmarks (Gainotti, Miceli, & Caltagirone, 1977). We refer the reader to Table 1 for descriptive statistics of these neuropsychological. Inclusion criteria for both AD patients and healthy controls were: 1) MMSE raw score ≥ 16 ; 2) between 50 and 90

² MRI and PET: 3 patients; MRI and CAT: 1 patient; only MRI 8 patients, only PET 4 patients.

years of age; 3) no less than 3 years of schooling; 4) normal or corrected-to-normal vision with no history of eye surgery; 5) no history of neurological (other than AD) and/or psychiatric disorders; 6) no history of alcohol or substance abuse and/or use of medications likely to affect cognitive functioning; 7) able to understand the instructions and perform the task.

Eighteen AD patients³ and all healthy controls also completed a neuropsychological battery of tests at the end of the eye-tracking visual search task (refer to Table 2 for descriptive and inferential t-test statistics comparing the two groups). Specifically, they completed: (1) the Albert's line cancellation task (Fullerton, Mcsherry, & Stout, 1986), which tests for spatial neglect; (2) a word reading and an object naming task from the "Esame Neuropsicologico Per l'Afasia" (neuropsychological test battery for aphasia - ENPA, Capasso & Miceli, 2001), to evaluate potential impairments in reading and object recognition (i.e., agnosia) and (3) the forward digit span from the Wechsler adult intelligence scale (WAIS-IV, Wechsler, 2008), which tests short-term memory, to examine the capacity of our participants to retain the identity of the cued object across the search task. All participants were at ceiling on the Albert's task and displayed similar performances on word reading and object naming. Even though healthy controls had a significantly higher digit span than AD patients, we note that this latter group was able to correctly remember an average of 4.83 digits, which suggests that their short-term working memory was sufficiently preserved to cope with the visual search task. All participants were naive to the purpose of the study and unfamiliar with the stimulus material. The study was approved by the Psychology Research Ethics Committee at the School of Philosophy, Psychology and Language Sciences at the University of Edinburgh (Ref: 12-1617) prior to starting the data collection; written consent was collected at the beginning of each session.

Design

³ Two AD patients reported fatigue and did not want to continue with the additional neuropsychological assessments.

254 A 2x2x2x2 mixed factorial design was used with *Semantic Relatedness* (unrelated, related),
255 *Visual Saliency* (non-salient, salient), and *Target* (absent, present) as within-participant variables,
256 and *Group* (control, AD) as a between-participant variable.

257 **Stimuli**

258 We used the 128 five-object arrays from Cimminella et al. (2020) and created a further 32
259 arrays (i.e., a total of 160 items) by crossing the visual saliency (non-salient, salient) and the
260 semantic relatedness (unrelated, related) of 40 critical objects (i.e., 40 * 4); refer to Supplementary
261 Material A for miniatures of the experimental arrays. The position of the critical object in the array
262 was rotated across items to prevent location biases. A total of 200 objects were taken from the Bank
263 of Standardized Stimuli (BOSS) database (Brodeur, Dionne-Dostie, Montreuil, & Lepage, 2010;
264 Brodeur, Guérard, & Bouras, 2014), and used to create the experimental arrays (i.e., 40 as critical
265 objects and 160 as distractors). All objects were univocally classifiable into 20 semantic categories
266 (see Cimminella et al., 2020, for details on the semantic category norms). In addition, all objects
267 were named by a sample of Italian participants to obtain naming norms in Italian (the reader is
268 referred to Supplementary Material B for the norming of the materials). No object picture was
269 presented more than once to avoid any uncontrolled effect that may derive from repeated exposures
270 to the same stimulus.

271 The arrays were presented on a uniform white background, at a resolution of 1024 x 768
272 pixels and at a viewing distance of 75 cm. Object pictures measured 150 x 150 pixels (4.25° x 4.47°
273 of visual angle) and were arranged on an imaginary circle such that the midpoint of each object was
274 equidistant from the centre of the array (corresponding to the central fixation cross) and from the
275 two adjacent object midpoints. The circle had a fixed radius of 344 pixels (9.75°), while the
276 distance between objects was 404.40 pixels (11.46°).

277 We manipulated the visual saliency of the critical object by changing its brightness/contrast
278 and hue/saturation with GIMP (Version 2.8.2) to make it visually more conspicuous in the salient
279 than in the non-salient condition. We validated the effectiveness of our manipulation with the

280 Walther and Koch's Matlab Saliency Toolbox (2006): a Wilcoxon signed-rank test confirmed that
281 the critical object was ranked among the most visually conspicuous regions of the array in the
282 salient condition ($Mdn = 1$) and the least in the non-salient condition ($Mdn = 4$), $p < .001$, z -
283 coefficient = -7.98.

284 We implemented the semantic relatedness manipulation by creating object arrays with all
285 objects belonging to the same semantic category (related), or all distractors of the same semantic
286 category but the critical object of a different one (unrelated). To validate the semantic relatedness
287 manipulation, we computed the mean of the semantic similarity score of the critical object with
288 every other distractor in each experimental array using Latent Semantic Analysis (LSA, Landauer &
289 Dumais, 1997; Landauer, Foltz, & Laham, 1998), trained on co-occurrences of words (as
290 implemented by Hoffman, Lambon Ralph, and Rogers 2013) and on labels of objects (as normed by
291 Brodeur et al. 2010, 2014). A t-test confirmed that semantic similarity between the critical object
292 and all other distractors was significantly higher in the semantically related ($M = .44$, $SD = .24$) than
293 unrelated condition ($M = .01$, $SD = .08$), $t(39) = 10.52$, $p < .001$, Cohen's $d = 1.66$.

294 We also included 40 five-object filler arrays, 32 arrays from Cimminella et al. (2020), and 8
295 new arrays created using objects from the BOSS database (Brodeur et al., 2010, 2014). The filler
296 items ensured a balanced distribution of target detection response (i.e., 50% object present, 50%
297 object absent); and were not systematically manipulated in semantic relatedness or visual saliency
298 nor will they be analysed. Each participant saw the same 40 fillers and 40 unique experimental
299 arrays, which were counterbalanced across the conditions of visual saliency, semantic relatedness,
300 and target (absent, present) using a Latin square rotation. In the target-present condition, the cue
301 word of the search target, i.e., the target name, always referred to a critical object depicted in the
302 array. In the target-absent condition, the cue word did not refer to any object in the arrays, but it was
303 either semantically related to the critical object depicted in the display and unrelated to all other
304 semantically homogenous distractor objects ($M = .03$, $SD = .11$), or semantically related to all
305 objects ($M = .30$, $SD = .21$), $t(39) = 7.02$, $p < .001$. Cohen's $d = 1.11$ (the reader is referred to

306 Supplementary Material C for the full list of target-present and target-absent experimental arrays
307 used in this study). Finally, we used the Bank of Local Analyzer Responses (BOLAR) method
308 (Zelinsky, 2003) to examine whether the critical object was visually more similar to the distractors
309 when semantically related rather than unrelated to them (Cimminella et al., 2020). We found that
310 this was indeed the case (related: $M = .52$, $SD = .13$; unrelated: $M = .49$, $SD = .13$; $t(39) = 2.39$, $p =$
311 $.02$, Cohen's $d = 0.38$) and so controlled for the potential confounding effects of visual similarity on
312 search by including it as a quasi-experimental predictor in our models (see the Statistical Analysis
313 sub-section below for details).

314 **Apparatus**

315 Visual stimuli were displayed on a 19-inch Sony SDM E96D monitor, and eye movements
316 were recorded using a Gazepoint GP3 HD 150 eye-tracker (150 Hz sampling rate). A pilot study on
317 a pool of young participants ensured that the Gazepoint GP3 HD 150 eye-tracker was able to detect
318 semantic relatedness effects on eye movements in a similar way to the EyeLink 1000 (SR Research,
319 1,000 Hz), used in Cimminella et al. (2020), despite its lower sampling rate; see Supplementary
320 Material D for the pilot study. A forehead and chin rest were used to stabilise participants' viewing
321 position. The experiment was built using OpenSesame (Version 3.1.9, Mathôt, Schreij, &
322 Theeuwes, 2012) and the PyGaze Python plug-in (Dalmaijer, Mathôt, & Van der Stigchel, 2014)
323 was used to acquire the eye-movement data.

324 **Procedure**

325 At the beginning of each experimental session, the eye tracker was calibrated using a 9-point
326 grid appearing on the monitor. Each trial began with a fixation cross at the centre of the display; as
327 soon as participants looked at it, the experimenter, who was monitoring their gaze position,
328 manually pressed the F key on their keyboard to trigger the cue word indicating the search target.
329 The word was visible on the screen for 1,200 ms, followed by another central fixation cross for 150
330 ms, and then the object array was displayed. Participants received written and verbal instructions
331 and were asked to perform the task as quickly and accurately as possible by verbally responding *sì*

(Italian for *yes*) if the target was present in the array, or *no* if it was not. The experimenter recorded participants' responses by pressing the left or the right arrow key, respectively. If the participant answered *no*, the experiment progressed to the next trial. If the participant answered *yes*, the object array was replaced by a number array, and the participant was asked to verbally indicate the number matching the target location. This provided us with an additional verification of the search accuracy; responses were regarded as correct only if both the search response and the number response were accurate. If participants did not respond within 5,000 ms, a null response was logged (see Figure 2 for an example of a trial run). Each participant completed 4 practice trials and 80 randomized trials (40 experimental and 40 filler trials). If a patient could not remember the task, the experiment was stopped, and the instructions and the practice session repeated before continuing the session. The experimental session lasted approximately 45 minutes.

Data Analysis

Data processing

Fixations and saccade events were extracted from the raw eye movement samples with the I2MC algorithm, which is particularly suitable for noisy and low frequency data (Hessels, Niehorster, Kemner, & Hooze, 2017), using MATLAB (R2016b). Out of the 1,600 experimental trials considered in the analysis (i.e., 40 trials x 40 participants), we discarded 4 trials because of machine error (no eye movement was recorded) and 57 trials of the target-present condition where the number responses were inaccurate. On the remaining trials (1,539), we analysed the response accuracy. The eye-movement measures were computed only on accurate trials in which the critical object was fixated at least once (1,178 trials).

Dependent variables

We considered *response accuracy* to measure the target detection performance (a binary variable coded as 0 = Incorrect; 1 = Correct), and the following measures from the fixation data: *latency to first fixation*, which is the time (in ms) between the onset of the array and the first fixation on the critical object; and *first-pass dwell time*, that is the ratio between the sum of all

fixations during the first inspection of the critical object and all other fixations during the trial. The latencies to first fixation were z-scored to account for general slowing effect associated to AD (Faust, Balota, Spieler, & Ferraro, 1999).

Statistical analyses.

Linear and generalised linear mixed-effects models (G/LMM), as implemented by the lme4 package (Bates et al., 2015) in R (version 4.0.0), were used to analyse the data. We adopted the same forward best-path model selection procedure as in Cimminella et al. (2020), whereby we start by evaluating the random variables and then proceed by adding the predictors as main effects and in interaction, one-by-one, while testing whether the inclusion of each new parameter would significantly improve the log-likelihood of the model fit using chi-square tests. The only difference here with this approach is that the fixed effect of group was retained during model selection even when found non-significant to test for its potential interactions with the variables of interest of our study. The fixed effects considered, and centred to reduce co-linearity, were: *Semantic Relatedness* (Unrelated, Related), *Visual Saliency* (Non-salient, Salient), *Target* (Absent, Present), *Group* (Control, AD), and *Visual Similarity*, which was obtained by splitting the 160 items into two groups (Dissimilar, Similar) based on the median score obtained with the BOLAR (i.e., .52). The random variables included in the models, both as intercepts and slopes, were Participant (40) and Item (160). We report the coefficients⁴, standard errors, t-values (LMM), and z-values (GLMM) of those predictors that were retained in the final models. The p-values were based on Satterthwaite approximation for denominator degrees of freedom for LMMs models, on asymptotic Wald tests for GLMM models, and computed using the lmerTest R package (Kuznetsova, Brockhoff, & Christensen, 2017).

The data and R script to reproduce visualisations and analyses are available in the Open Science Framework (<https://osf.io/gyutd/>)

⁴ Bi-variate correlation matrices detailing the association of all predictors in each G/LME model are reported in Supplementary Material E.

Results

Model coefficients for response accuracy⁵, latency to first fixation and first-pass dwell time are reported in Table 3. On response accuracy (Figure 3), we found a significant main effect of group, whereby AD patients performed less accurately than healthy controls. There was also a significant main effect of semantic relatedness, and a significant interaction between semantic relatedness and target: accuracy was higher when the critical object was semantically unrelated than related to the distractors, with this difference being greater in target-absent than target-present trials.

When looking at latency to first fixation (Figure 4), we found that the critical object was looked at earlier when it was semantically unrelated to the other distractors than when it belonged to the same semantic category, and on target-present compared to target-absent trials. No other predictor (e.g., visual saliency or visual similarity) was found to be significant, either as main effect or interaction.

On first-pass dwell time (Figure 5), we found longer dwell times for the healthy age-matched controls than for the AD patients, when the critical object was semantically unrelated rather than related to the distractors and in target-present trials compared to target-absent trials. Moreover, when the critical object was semantically unrelated to the distractors, this resulted in longer dwell times in the healthy control group than in the AD group, and in target-present compared to target-absent trials. Importantly, we did not find any effect of visual saliency or visual similarity, either as main effect or as interaction, in any of the measures reported herein⁶.

Discussion

⁵ On this measure, we demonstrated using a binomial test that the performance of AD patients was significantly higher than chance across all experimental conditions ($p < .01$), indicating that they were able to complete the task.

⁶ We replicated the significant main effects of semantic relatedness and target when analysing overall dwell time, i.e., the ratio between the sum of all fixations on the critical object and the sum of all remaining fixations occurring during the trial (unrelated: $M = .47$, $SD = .26$; related: $M = .39$, $SD = .24$; $\beta = -.10$, $SE = .01$, $t = -7.79$, $p < .001$; target-absent: $M = .24$, $SD = .14$; target-present: $M = .61$, $SD = .21$; $\beta = .36$, $SE = .01$, $t = 27.32$, $p < .001$).

402 A frequent feature of AD patients' cognitive profile is an impairment in tasks requiring
403 explicit access to semantic knowledge, such as category fluency (Salmon et al., 1999), word-to-
404 picture matching (Adlam et al., 2006) and object naming (Holmes et al., 2006). There is evidence,
405 however, that some implicit semantic abilities related to automatic processing may be preserved
406 (e.g., Chertkow et al., 1994; Perri et al., 2003; Perry & Hodges, 1999), and eye-movement
407 responses can be revelatory of such implicit processes (Hannula et al., 2010).

408 In the present study, we investigated whether AD patients' ability to process semantic
409 information of visual objects is preserved, and whether such processing can occur in extra-foveal
410 vision. AD patients and healthy age-matched controls searched for a target (e.g., a car), which was
411 either semantically related or unrelated to a set of semantically homogenous distractors (e.g., an
412 array of vehicles or kitchen utensils, respectively). The target object was either visually available in
413 the display (target-present trials), or it was replaced by a semantically related competitor of the
414 target (target-absent trials), and it was either visually salient or non-salient to evaluate potential
415 bottom-up stimulus driven effects. So, if the semantics of objects were processed in extra-foveal
416 vision, we expected semantically unrelated targets to be prioritised in early overt attention and be
417 easier to recognize as deviant from the context of the array. On the contrary, we expected that a
418 semantically related critical object would be recognized with more difficulties in target-absent trials,
419 as the on-going competition of co-activated semantic distractors would increase the chance of false
420 detection, i.e., it requires the involvement of memory processes.

421 Results revealed a significant main effect of group on search accuracy, with AD patients
422 performing worse than age-matched controls, even if significantly above chance across all
423 experimental conditions, which is in line with previous evidence that AD can lead to impaired
424 object identification (Porter et al., 2010; Laatu, et, al., 2003; Tales et al., 2002). However, regardless
425 of the group, all participants detected the critical object more accurately when it was semantically
426 unrelated to the distractors compared to when it belonged to the same semantic category. This result
427 indicates that the semantic information of the critical object was processed enough to lead to a

428 significant difference in its identification, and crucially, this effect was also observed in the AD
429 group. In contrast to Seckin, et al., 2016, we did not observe greater difficulty to recognize
430 semantically related objects in our neuropathological population, i.e., AD, compared to healthy
431 older adults, which may reflect underlying differences between our patients and individuals with
432 more specific conditions such as semantic aphasia. Another important finding was the poorer
433 recognition performance for semantically related objects in target-absent trials, which indicates that
434 our participants found it harder to resolve the ongoing competition of semantically related
435 distractors, especially because they needed to rely more strongly on memory for the critical object,
436 which was no longer visually available in the context. Note that we did not find any difference
437 associated with the semantic relatedness of the critical object on the detection accuracy of young
438 participants in Cimminella et al. (2020), but this discrepancy can be accounted for by the fact that
439 performance was at ceiling in that study.

440 On eye-movement measures, we found that both AD patients and healthy controls looked at
441 the critical object significantly earlier, and for longer, when it was semantically unrelated compared
442 to when it was related to the other distractors. These findings align with our previous work on
443 healthy young participants which used the same experimental paradigm (Cimminella et al., 2020),
444 and complement previous work using naturalistic scenes that also demonstrated extra-foveal
445 processing in healthy older adults (Borges, Fernandes, & Coco, 2020) and unravelled the links to
446 the brain activity time-locked to it (Coco, Nuthmann, & Dimigen, 2020). These results strongly
447 suggest that extra-foveal processing of object semantics occurred in AD patients, and as for the
448 healthy older adults and the younger adults of our previous studies, it contributed to guiding overt
449 visual attention. The solid evidence of intact processing of object semantics observed on both the
450 allocation of visual attention and the target detection accuracy suggests that the semantic deficits
451 found in previous studies of AD patients may be associated with factors other than a complete loss
452 of semantic knowledge. This core finding has important methodological, theoretical, and practical
453 implications for AD research.

454 Methodologically, our results point to two key factors that call for attention when assessing
455 semantic abilities in AD patients: whether the task requires explicit or implicit access to semantic
456 knowledge; and the nature of the stimuli used, i.e., verbal or non-verbal. Semantic impairments in
457 AD have been typically reported in category fluency, object naming and word-to-picture matching
458 tasks (Holmes et al., 2006; Monsch et al., 1994; Salmon et al., 1999; Spaan et al., 2003), which
459 require participants to intentionally access their semantic knowledge and retrieve relevant
460 information. In contrast, in the present study we used a visual search task, which effectively entails
461 a cueing component, i.e., the target of the search is prompted to the participant, whereby explicit
462 retrieval of semantic information is not required. We argue that this resulted in the relatively
463 automatic activation of a specific target template that facilitated processing of the associated
464 semantic information, thus leading to the emergence of preserved semantic abilities in AD patients.
465 The other factor that may have contributed to our findings is that the stimuli in our task consisted of
466 non-verbal material (i.e., pictures of objects), which is known to have a cognitive advantage over
467 verbal material that demands more complex strategies of lexical matching and retrieval (Ally, Gold,
468 & Budson, 2009a; Ally et al., 2008). It is possible that studies that employ verbal material, which
469 requires additional processing effort and resources compared to pictures, may find AD-related
470 semantic impairments which would not be observed in studies that employ the same tasks but with
471 visual material (see O'Connor & Ally, 2010 for an example of a study about the effects of stimulus
472 format on memory tasks). In fact, a limitation of the present study is that we did not directly
473 compare semantics relatedness effects using both verbal and non-verbal material. One objective of
474 our current investigations on this topic is to utilise a similar experimental paradigm as in this study,
475 but to probe search using verbal stimuli only, i.e., a target word followed by an array of words. We
476 trust this will help elucidate the role played by the stimulus material on semantic processing in AD
477 patients.

478 Our study also highlights the methodological usefulness of eye-movement responses in
479 research on clinical populations, because they can index implicit mechanisms of semantic

processing (Hannula et al., 2010). In the present study, eye movements revealed preserved extra-foveal capture by object semantics in AD patients, which counters previous evidence of impaired extra-foveal processing in this population (Boucart, Bubbico, Szaffarczyk, & Pasquier, 2014; Rösler, Mapstone, Hays-Wicklund, Gitelman, & Weintraub, 2005). We attribute this discrepancy to the semantic pop-out effect induced by our experimental paradigm: when patients were faced with a single object that was semantically divergent from the surrounding context, this facilitated its covert processing in extra-foveal vision and elected it as a possible target to saccade to. This explanation would align with previous evidence of similar performance in AD patients and healthy controls in instances where the selective demands of the task are reduced (e.g., Tippet et al., 2004). An important question that stems from this finding is whether similar evidence of preserved semantic processing would manifest if all objects were clustered around central fixation and placed within foveal vision. This experimental manipulation would remove the need for overt eye-movements, whereby performance would depend solely on covert attention. Given the current results, we would expect target detection accuracy to show the same advantage for a semantically unrelated compared to a related critical object in AD patients, as such a measure is independent of the overt allocation of visual attention.

Analysis of the first-pass dwell times also indicated preserved semantic abilities in the AD group, as a semantically unrelated critical object was looked at more than a semantically related one, although this difference was less pronounced in the AD patients than it was in the healthy controls. The evidence of longer foveation mediated by object semantics hints at other mechanisms of target identification, closely related with linguistic processing, which may be spared by AD too. Psycholinguistics research on young adults has, in fact, clearly shown effects of semantic relatedness during spoken language situated in a visual context (e.g., arrays of visual objects; Huettig & Altmann, 2005; or Coco et al., 2016, for photo-realistic stimuli); which has led to integrative proposals about the relationship between attention, memory and language (see Huettig et al., 2011, for an example of an integrative theoretical framework). Future research on AD should

506 address questions about situated language processing, focusing on whether evidence of spared
507 semantics would also emerge when language and vision are synchronously active.

508 Theoretically, our findings argue against a generalised loss of semantic memory in AD
509 (Adlam et al., 2006; Hodges et al., 1992; Salmon et al., 1999), and suggests that semantic
510 processing may be spared, at least in part, when the task can be performed relatively automatically
511 (Ober, 2002; Perri et al., 2003). Moreover, our results seem to align with a two-stage model of
512 recollection (Moscovitch, 2008), even if we realise that such a model was mainly conceived to
513 describe episodic memory. In the first-stage, semantic information is implicitly retrieved from
514 memory, which is reflected in the responses indirectly associated to memory processes (e.g., eye
515 movement). Then, in a second slower stage, such retrieved information becomes explicitly
516 available, influencing downstream responses (e.g., the manual response of target detection). This
517 would be consistent with evidence of gradual semantic degradation in patients with semantic
518 aphasia (Seckin et al., 2016): as the pathology degrades semantic memory, concepts become
519 progressively more blurred rather than being entirely lost.

520 Our results also have implications for neuropsychological diagnostics criteria and theoretical
521 accounts of semantic memory. Additional correlation analyses of the AD patients'
522 neuropsychological profiles and semantic relatedness effects on their search responses, in fact,
523 failed to show any significant association between the two (see Appendix for a description of this
524 analysis and their results). A key implication of this dissociation is that visual guidance by object
525 semantics during a search task is an index of cognitive functioning which escapes what is assessed
526 using classic neuropsychological tests. Additionally, the lack of an association between verbal
527 semantic abilities of our AD patients (e.g., RAVLT) and visual guidance by object semantics may
528 argue against the notion of a unitary semantic memory system (Bright, Moss, & Tyler, 2004;
529 Lambon, Patterson, & Hodges, 1997) and suggests instead that the semantic system may be divided
530 into modality-specific subsystems, which interact with one another but separately deal with pictures
531 and words (Paivio, 2014; Saffran, Branch Coslett, Martin, & Boronat, 2003). If we assume this

532 division, it follows that AD may involve preserved semantic processing for visual material and a
533 specific impairment in the semantic processing of verbal material.

534 As a point of discussion, and differently from our previous study on young participants, the
535 analysis of the probability that the first fixation would fall on the critical object did not show any
536 effect of semantic relatedness⁷. This suggests that, regardless of pathology, older adults require
537 more fixations to orient their gaze in the visual scene, which is in line with the general slowing
538 associated with cognitive aging (Crawford, Smith, & Berry, 2017; Rösler et al., 2005). Moreover,
539 our study did not show any other key difference in fixation behaviour between the healthy control
540 participants and the AD group, beyond the significantly decreased first fixations time on the critical
541 object in patients (see Fernández et al., 2018 for a similar finding). That being said, differences on
542 eye-movement responses of AD patients have mostly been found on saccades (e.g., Fletcher &
543 Sharpe, 1986; Yang, Wang, Su, Xiao, & Kapoula, 2013). Our analysis comparing the mean saccade
544 amplitude of the two groups did not show any significant difference and hence it was not reported.
545 This result must be taken cautiously due to the relatively low sampling rate of our eye-tracker that
546 may not have been sensitive enough to detect finer differences, which calls for future research
547 specifically looking at saccade measures in visual search paradigms adapted to the purpose (e.g., by
548 varying systematically the distance of the target and distractor objects in the arrays).

549 In sum, our study suggests that AD patients retain semantic knowledge and are able to
550 access it when the task involves more implicit processes and utilises non-verbal stimuli, probably
551 indicating that they present deficits when this knowledge has, instead, to be intentionally and
552 explicitly retrieved. These findings also have some practical implications, as determining which
553 conditions can enable the emergence of preserved semantic processes in AD patients may allow the
554 development of strategies aimed at facilitating their access to and use of semantic knowledge more
555 effectively, which in turn could increase patients' quality of life.

⁷ We did not include the full description of this analysis, as it was not significant.

556

557

558

Acknowledgments

559

560

We thank Dr Jennifer Hall for her assistance with language editing and careful proofreading of the manuscript.

561

Conflict of Interest Statement

562

The authors declare that there is no conflict of interest.

563

Appendix

Pearson correlation was used to determine whether the neuropsychological performances of the AD patients were related to their manual and eye-movement responses to the visual search task. The aim of this analysis was to qualify the nature of semantic processing during visual search in relation to neuropsychological performance. Evidence of a dissociation between neuropsychological results and search responses would suggest that visual guidance by object semantics may be independent from what is classically assessed by off-the-shelf tests.

To get at a synthetic measure of semantic relatedness effects on search responses (i.e., accuracy, latency to first fixation and first-pass dwell time), we first computed by-participant averages across trials and separately for the two conditions of Semantic Relatedness and then calculated their difference (Unrelated - Related). This difference score reflects the amount of change that a semantically unrelated critical object implied on search responses compared to a semantically related object. For example, if a semantically unrelated object were expected to be looked at earlier than a semantically related object, we would expect the difference score for latency of first fixation to be negative. All measures were standardised and centred prior to this analysis.

Results revealed significant positive correlations between the mini mental state examination and two tests of short-term memory (RAVLT.IR and DSF), which is congruent with the type of functions assessed by the MMSE. Likewise, we observed significant positive correlations between RCPM, CD and CDL which are all tests tapping into visuo-spatial functions. Most importantly, however, we did not observe significant correlations between any of the neuropsychological tests and search responses. The only significant negative correlation observed on search responses was between the latency of first fixation and first-pass dwell time, which aligns with the main results reported on these measures: an unrelated object is looked at earlier but for longer compared to a semantically related object. Correlations which were significant at $p < .05$ are displayed in Figure 6, and the full results of these correlation analyses are reported in Table 4.

*Table 4: Pearson correlation analysis examining the neuropsychological profiles of AD participants together with their responses to the visual search task (accuracy, latency to first fixation, first-pass dwell time). On search responses, we first computed by-participant averages across trials for the two conditions of Semantic Relatedness then derived a difference score (Unrelated - Related). This measure reflects the differential that semantic relatedness had on search responses. All measures have been standardised and centred before correlating them. Significant correlations are marked using bold face, and the associated p-value indicated using asterisks (*p < 0.05; **p < 0.01; ***p < 0.001).*

	MMSE	FAB	RAVLT (IR)	RAVLT (DR)	WF	RCPM	CD	CDL	WR	ON	DSF	Accuracy	Latency to First Fixation
MMSE													
FAB	0.00												
RAVLT (IR)	0.51*	0.14											
RAVLT (DR)	0.28	-0.34	0.62**										
WF	0.27	0.40	0.18	-0.15									
RCPM	0.36	0.55*	0.56*	0.15	0.43								
CD	0.31	0.22	0.28	0.03	0.54*	0.69***							
CDL	0.13	0.07	0.12	-0.03	0.32	0.45	0.69**						
WR	0.07	0.29	0.15	-0.04	0.32	0.13	-0.05	-0.24					
ON	-0.09	0.16	0.23	0.20	-0.01	0.12	-0.07	-0.10	-0.02				
DSF	0.76***	0.46	0.33	-0.06	0.38	0.40	0.35	0.01	0.34	-0.03			
Accuracy	-0.28	-0.03	-0.06	-0.03	-0.09	-0.07	-0.08	-0.12	0.46	-0.09	-0.14		
Latency to First Fixation	-0.11	-0.29	-0.16	0.22	-0.02	-0.27	0.13	0.22	-0.23	-0.04	-0.27	-0.12	
First pass Dwell Time	0.07	0.06	-0.10	-0.23	-0.04	-0.14	-0.05	-0.11	0.33	0.25	0.38	0.08	-0.53*

Note. MMSE: Mini mental state examination; FAB: frontal assessment battery; RAVLT: Rey's auditory verbal learning test (IR, immediate recall; DR, delayed recall); WF: word fluency; RCPM: Raven's coloured progressive matrices; CD: freehand copying of drawings; CDL: copying drawings with landmarks; WR: word reading; ON: Object naming; DSF: Digit span forwards. The horizontal line in the table distinguishes between neuropsychological and search responses.

Figure 6: Pearson correlation plot examining the neuropsychological profiles of AD participants and their responses to the visual search task (accuracy, latency to first fixation, first-pass dwell time). On search responses, we first computed by-participant averages across trials for the two conditions of Semantic Relatedness then calculated a difference score (Unrelated - Related). This measure would reflect the differential that semantic relatedness had on search responses for each participant. All measures have been standardised and centred before correlating them. Only correlations significant at $p < 0.05$ are presented and empty cells indicate non-significant correlations. The colour indicates the strength of the correlation from red (-1) to blue (1).

Note. MMSE: Mini mental state examination; FAB: frontal assessment battery; RAVLT: Rey's auditory verbal learning test (IR, immediate recall; DR, delayed recall); WF: word fluency; RCPM: Raven's coloured progressive matrices; CD: freehand copying of drawings; CDL: copying drawings with landmarks; WR: word reading; ON: Object naming; DSF: Digit span forwards.

References

- Adlam, A. L. R., Bozeat, S., Arnold, R., Watson, P., & Hodges, J. R. (2006). Semantic knowledge in mild cognitive impairment and mild Alzheimer's disease. *Cortex*, 42(5), 675–684. [https://doi.org/10.1016/S0010-9452\(08\)70404-0](https://doi.org/10.1016/S0010-9452(08)70404-0)
- Alexander, R. G., & Zelinsky, G. J. (2011). Visual similarity effects in categorical search. *Journal of Vision*, 11(8), 1–15. <https://doi.org/10.1167/11.8.9>
- Ally, B. A., Gold, C. A., & Budson, A. E. (2009a). An evaluation of recollection and familiarity in Alzheimer's disease and mild cognitive impairment using receiver operating characteristics. *Brain and Cognition*, 69(3), 504–513. <https://doi.org/10.1016/j.bandc.2008.11.003>
- Ally, B. A., Gold, C. A., & Budson, A. E. (2009b). The picture superiority effect in patients with Alzheimer's disease and mild cognitive impairment. *Neuropsychologia*, 47(2), 595–598. <https://doi.org/10.1016/j.neuropsychologia.2008.10.010>
- Ally, B. A., McKeever, J. D., Waring, J. D., & Budson, A. E. (2009). Preserved frontal memorial processing for pictures in patients with mild cognitive impairment. *Neuropsychologia*, 47(10), 2044–2055. <https://doi.org/10.1016/j.neuropsychologia.2009.03.015>
- Ally, B. A., Waring, J. D., Beth, E. H., McKeever, J. D., Milberg, W. P., & Budson, A. E. (2008). Aging memory for pictures: Using high-density event-related potentials to understand the effect of aging on the picture superiority effect. *Neuropsychologia*, 46(2), 679–689. <https://doi.org/10.1016/j.neuropsychologia.2007.09.011>
- Ambra, F. I., Iavarone, A., Ronga, B., Chieffi, S., Carnevale, G., Iaccarino, L., ... Garofalo, E. (2016). Qualitative patterns at Raven's colored progressive matrices in mild cognitive impairment and Alzheimer's disease. *Aging Clinical and Experimental Research*, 28(3), 561–565. <https://doi.org/10.1007/s40520-015-0438-9>
- Baddeley, A. D., Baddeley, H. A., Bucks, R. S., & Wilcock, G. K. (2001). Attentional control in Alzheimer's disease. *Brain*, 124(8), 1492–1508. <https://doi.org/10.1093/brain/124.8.1492>
- Bandera, L., Della Sala, S., Laiacona, M., Luzzatti, C., & Spinnler, H. (1991). Generative

- 647 associative naming in dementia of Alzheimer's type. *Neuropsychologia*, 29(4), 291–304.
648 [https://doi.org/10.1016/0028-3932\(91\)90043-8](https://doi.org/10.1016/0028-3932(91)90043-8)
- 649 Bartolo, A., Della Sala, S., & Cubelli, R. (2016). The sign of “undue contact” in the Object Use
650 Test. *Cortex*, 75, 235–236. <https://doi.org/10.1016/j.cortex.2015.06.016>
- 651 Bates, D., Mächler, M., Bolker, B. M., Walker, S. C., Machler, M., Bolker, B. M., ... Walker, S. C.
652 (2015). Fitting Linear Mixed-Effects Models using lme4. *Journal of Statistical Software*,
653 67(1), 1–48. <https://doi.org/10.18637/jss.v067.i01>
- 654 Borges, M. T., Fernandes, E. G., & Coco, M. I. (2020). Age-related differences during visual
655 search: the role of contextual expectations and cognitive control mechanisms. *Aging*,
656 *Neuropsychology, and Cognition*, 27(4), 489–516.
657 <https://doi.org/10.1080/13825585.2019.1632256>
- 658 Boucart, M., Bubbico, G., Szaffarczyk, S., & Pasquier, F. (2014). Animal spotting in Alzheimer's
659 disease: An eye tracking study of object categorization. *Journal of Alzheimer's Disease*, 39(1),
660 181–189. <https://doi.org/10.3233/JAD-131331>
- 661 Bright, P., Moss, H., & Tyler, L. K. (2004). Unitary vs multiple semantics: PET studies of word and
662 picture processing. *Brain and Language*, 89(3), 417–432.
663 <https://doi.org/10.1016/j.bandl.2004.01.010>
- 664 Brodeur, M. B., Dionne-Dostie, E., Montreuil, T., & Lepage, M. (2010). The Bank of Standardized
665 Stimuli (BOSS), a New Set of 480 Normative Photos of Objects to Be Used as Visual Stimuli
666 in Cognitive Research. *PLoS ONE*, 5(5), 1–13. Retrieved from
667 <http://10.0.5.91/journal.pone.0010773>
- 668 Brodeur, M. B., Guérard, K., & Bouras, M. (2014). Bank of Standardized Stimuli (BOSS) phase ii:
669 930 new normative photos. *PLoS ONE*, 9(9), 1–10.
670 <https://doi.org/10.1371/journal.pone.0106953>
- 671 Capasso, R., & Miceli, G. (2001). *Esame Neuropsicologico per l' Afasia (E.N.P.A.)*. Milano:
672 Springer.

- 673 Chan, A. S., Butters, N., Paulsen, J. S., Salmon, D. P., Swenson, M. R., & Maloney, L. T. (1993).
674 An assessment of the semantic network in patients with Alzheimer's disease. *Journal of*
675 *Cognitive Neuroscience*, 5(2), 254–261. <https://doi.org/10.1162/jocn.1993.5.2.254>
- 676 Chertkow, H., Bub, D., Bergman, H., Bruemmer, A., Merling, A., & Rothfleisch, J. (1994).
677 Increased Semantic Priming in Patients with Dementia of the Alzheimer's Type. *Journal of*
678 *Clinical and Experimental Neuropsychology*, 16(4), 608–622.
679 <https://doi.org/10.1080/01688639408402672>
- 680 Cimminella, F., Della Sala, S., & Coco, M. I. (2020). Extra-foveal Processing of Object Semantics
681 Guides Early Overt Attention During Visual Search. *Attention, Perception, & Psychophysics*,
682 82, 655–670. <https://doi.org/0.3758/s13414-019-01906-1>
- 683 Coco, M. I., Keller, F., & Malcolm, G. L. (2016). Anticipation in Real-World Scenes: The Role of
684 Visual Context and Visual Memory. *Cognitive Science*, 40(8), 1995–2024.
685 <https://doi.org/10.1111/cogs.12313>
- 686 Coco, M. I., Nuthmann, A., & Dimigen, O. (2019). Fixation-related brain potentials during
687 semantic integration of object–scene information. *Journal of Cognitive Neuroscience*, 32(4),
688 571–589. https://doi.org/10.1162/jocn_a_01504
- 689 Costa, A., Bagoj, E., Monaco, M., Zabberoni, S., De Rosa, S., Papantonio, A. M., ... Carlesimo, G.
690 A. (2014). Standardization and normative data obtained in the Italian population for a new
691 verbal fluency instrument, the phonemic/semantic alternate fluency test. *Neurological*
692 *Sciences*, 35(3), 365–372. <https://doi.org/10.1007/s10072-013-1520-8>
- 693 Crawford, T. J., Smith, E. S., & Berry, D. M. (2017). Eye gaze and aging: Selective and combined
694 effects of working memory and inhibitory control. *Frontiers in Human Neuroscience*, 11, 563.
695 <https://doi.org/10.3389/fnhum.2017.00563>
- 696 Crutcher, M. D., Calhoun-Haney, R., Manzanares, C. M., Lah, J. J., Levey, A. I., & Zola, S. M.
697 (2009). Eye tracking during a visual paired comparison task as a predictor of early dementia.
698 *American Journal of Alzheimer's Disease and Other Dementias*, 24(3), 258–266.

- 699 <https://doi.org/10.1177/1533317509332093>
- 700 Cuetos, F., Arce, N., Martínez, C., & Ellis, A. W. (2017). Word recognition in Alzheimer's disease:
701 Effects of semantic degeneration. *Journal of Neuropsychology*, 11(1), 26–39.
702 <https://doi.org/10.1111/jnp.12077>
- 703 Dalmaijer, E. S., Mathôt, S., & Van der Stigchel, S. (2014). PyGaze: an open-source, cross-platform
704 toolbox for minimal-effort programming of eyetracking experiments. *Behavior Research*
705 *Methods*, 46(4), 913–921. <https://doi.org/10.3758/s13428-013-0422-2>
- 706 Daum, I., Riesch, G., Sartori, G., & Birbaumer, N. (1996). Semantic memory impairment in
707 Alzheimer's disease. *Journal of Clinical and Experimental Neuropsychology*, 18(5), 648–665.
- 708 Dubois, B., Slachevsky, A., Litvan, I., & Pillon, B. (2000). The FAB: A frontal assessment battery
709 at bedside. *Neurology*, 55(11), 1621–1626. <https://doi.org/10.1212/WNL.55.11.1621>
- 710 Faust, M. E., Balota, D. A., Spieler, D. H., & Ferraro, F. R. (1999). Individual differences in
711 information-processing rate and amount: Implications for group differences in response
712 latency. *Psychological Bulletin*, 125(6), 777–799. [https://doi.org/10.1037/0033-](https://doi.org/10.1037/0033-2909.125.6.777)
713 [2909.125.6.777](https://doi.org/10.1037/0033-2909.125.6.777)
- 714 Fernández, G., Orozco, D., Agamennoni, O., Schumacher, M., Sañudo, S., Biondi, J., & Parra, M.
715 A. (2018). Visual Processing during Short-Term Memory Binding in Mild Alzheimer's
716 Disease. *Journal of Alzheimer's Disease : JAD*, 63(1), 185–194. [https://doi.org/10.3233/JAD-](https://doi.org/10.3233/JAD-170728)
717 [170728](https://doi.org/10.3233/JAD-170728)
- 718 Fletcher, W. A., & Sharpe, J. A. (1986). Saccadic eye movement dysfunction in Alzheimer's
719 disease. *Annals of Neurology*, 20(4), 464–471. <https://doi.org/10.1002/ana.410200405>
- 720 Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). “Mini-mental state”: A practical method
721 for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*,
722 12(3), 189–198. [https://doi.org/10.1016/0022-3956\(75\)90026-6](https://doi.org/10.1016/0022-3956(75)90026-6)
- 723 Frisoni, G. B., Rozzini, R., Bianchetti, A., & Trabucchi, M. (1993). Principal Lifetime Occupation
724 and MMSE Score in Elderly Persons. *Journals of Gerontology*, 48(6), S310–S314.

- 725 <https://doi.org/10.1093/geronj/48.6.S310>
- 726 Fullerton, K. J., Mcsherry, D., & Stout, R. W. (1986). Albert's test: a neglected test of perceptual
727 neglect (Fullerton 1986). *The Lancet*, 327(8478), P430-432. <https://doi.org/10.1016/S0140->
728 6736(86)92381-0
- 729 Gainotti, G., Miceli, G., & Caltagirone, C. (1977). Constructional apraxia in left brain-damaged
730 patients: a planning disorder? *Cortex*, 13(2), 109–118. <https://doi.org/10.1016/s0010->
731 9452(77)80002-6
- 732 Gilmore, G. C., & Whitehouse, P. J. (1995). Contrast sensitivity in Alzheimer's Disease: A 1-year
733 longitudinal analysis. *Optomtry and Vision Science*, 72(2), 83–91.
- 734 Gold, C. A., & Budson, A. E. (2008). Memory loss in Alzheimer's disease: Implications for
735 development of therapeutics. *October*, 8(12), 1879–1891.
736 <https://doi.org/10.1586/14737175.8.12.1879>.
- 737 Hannula, D. E., Althoff, R. R., Warren, D. E., Riggs, L., Cohen, N. J., & Ryan, J. D. (2010). Worth
738 a glance: using eye movements to investigate the cognitive neuroscience of memory. *Frontiers*
739 *in Human Neuroscience*, 4(October), 166. <https://doi.org/10.3389/fnhum.2010.00166>
- 740 Henderson, J. M., & Hayes, T. R. (2017). Meaning-based guidance of attention in scenes as
741 revealed by meaning maps. *Nature Human Behaviour*, 1(10), 743–747.
742 <https://doi.org/10.1038/s41562-017-0208-0>
- 743 Hernández, M., Costa, A., Juncadella, M., Sebastián-Gallés, N., & Reñé, R. (2008). Category-
744 specific semantic deficits in Alzheimer's disease: A semantic priming study.
745 *Neuropsychologia*, 46(4), 935–946. <https://doi.org/10.1016/j.neuropsychologia.2007.11.018>
- 746 Hessels, R. S., Niehorster, D. C., Kemner, C., & Hooge, I. T. C. (2017). Noise-robust fixation
747 detection in eye movement data: Identification by two-means clustering (I2MC). *Behavior*
748 *Research Methods*, 49(5), 1802–1823. <https://doi.org/10.3758/s13428-016-0822-1>
- 749 Hodges, J. R., & Patterson, K. (1995). Is semantic memory consistently impaired early in the course
750 of Alzheimer's Disease? Neuroanatomical and diagnostic implications. *Neuropsychologia*,

- 751 33(4), 441–459.
- 752 Hodges, J. R., Salmon, D. P., & Butters, N. (1992). Semantic memory impairment in Alzheimer's
753 disease: Failure of access or degraded knowledge? *Neuropsychologia*, 30(4), 301–314.
- 754 Hoffman, P., Lambon, R. M., & Rogers, T. T. (2013). Semantic diversity: A measure of semantic
755 ambiguity based on variability in the contextual usage of words. *Behavior Research Methods*,
756 45(3), 718–730. <https://doi.org/10.3758/s13428-012-0278-x>
- 757 Hollingworth, A., Williams, C. C., & Henderson, J. M. (2001). To see and remember: Visually
758 specific information is retained in memory from previously attended objects in natural scenes.
759 *Psychonomic Bulletin and Review*, 8(4), 761–768. <https://doi.org/10.3758/BF03196215>
- 760 Holm, L., Eriksson, J., & Andersson, L. (2008). Looking as if you know: Systematic object
761 inspection precedes object recognition. *Journal of Vision*, 8(4), 1–7.
762 <https://doi.org/10.1167/8.4.14>
- 763 Holmes, S., Fitch, F. J., & Ellis, A. (2006). Age of acquisition affects object recognition and
764 naming in patients with Alzheimer's disease. *Journal of Clinical and Experimental*
765 *Neuropsychology*, 28(6), 1010–1022. <https://doi.org/10.1080/13803390591004392>
- 766 Huettig, F., & Altmann, G. T. M. (2005). *Word meaning and the control of eye fixation : semantic*
767 *competitor effects and the visual world paradigm*. 96(1), 23–32.
768 <https://doi.org/10.1016/j.cognition.2004.10.003>
- 769 Huettig, F., Olivers, C. N. L., & Hartsuiker, R. J. (2011). Looking, language, and memory: bridging
770 research from the visual world and visual search paradigms. *Acta Psychologica*, 137(2), 138–
771 150. <https://doi.org/10.1016/j.actpsy.2010.07.013>
- 772 Iavarone, A., Ronga, B., Pellegrino, L., Loré, E., Vitaliano, S., Galeone, F., & Carlomagno, S.
773 (2004). The Frontal Assessment Battery (FAB): normative data from an Italian sample and
774 performances of patients with Alzheimer's disease and frontotemporal dementia. *Functional*
775 *Neurology*, 19(3), 191–195.
- 776 Ivanoiu, A., Adam, S., Van Der Linden, M., Salmon, E., Juillerat, A. C., Mulligan, R., & Seron, X.

- 777 (2005). Memory evaluation with a new cued recall test in patients with mild cognitive
778 impairment and Alzheimer's disease. *Journal of Neurology*, 252(1), 47–55.
779 <https://doi.org/10.1007/s00415-005-0597-2>
- 780 Jacoby, L. L. (1991). A process dissociation framework: Separating automatic from intentional uses
781 of memory. *Journal of Memory and Language*, 30(5), 513–541. [https://doi.org/10.1016/0749-](https://doi.org/10.1016/0749-596X(91)90025-F)
782 596X(91)90025-F
- 783 Kergoat, H., Kergoat, M., Justino, L., Chertkow, H., Robillard, A., & Bergman, H. (2002). Visual
784 retinocortical function in dementia of the Alzheimer type. *Gerontology*, 48, 197–203.
785 <https://doi.org/10.1159/000058350>
- 786 Kiefer, M., & Pulvermüller, F. (2012). Conceptual representations in mind and brain: Theoretical
787 developments, current evidence and future directions. *Cortex*, 48(7), 805–825.
788 <https://doi.org/10.1016/j.cortex.2011.04.006>
- 789 Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in
790 Linear Mixed Effects Models. *Journal of Statistical Software*, 82(13), 1–26.
791 <https://doi.org/10.18637/jss.v082.i13>
- 792 Laatu, S., Revonsuo, A., Jäykkä, H., Portin, R., & Rinne, J. O. (2003). Visual object recognition in
793 early Alzheimer's disease: deficits in semantic processing. *Acta Neurologica Scandinavica*,
794 108(2), 82–89.
- 795 Lambon, R. M., Patterson, K., & Hodges, J. R. (1997). The relationship between naming and
796 semantic knowledge for different categories in dementia of Alzheimer's type.
797 *Neuropsychologia*, 35(9), 1251–1260. [https://doi.org/10.1016/S0028-3932\(97\)00052-3](https://doi.org/10.1016/S0028-3932(97)00052-3)
- 798 Landauer, T. K., & Dumais, S. T. (1997). A Solution to Plato's Problem: The Latent Semantic
799 Analysis Theory of Acquisition, Induction, and Representation of Knowledge. *Psychological*
800 *Review*, 104(2), 211–240. <https://doi.org/10.1037/0033-295X.104.2.211>
- 801 Landauer, T. K., Foltz, P. W., & Laham, D. (1998). An introduction to latent semantic analysis.
802 *Discourse Processes*, 25, 259–284. <https://doi.org/10.1080/01638539809545028>

- 803 Loftus, G. R., & Mackworth, N. H. (1978). Cognitive determinants of fixation location during
804 picture viewing. *Journal of Experimental Psychology. Human Perception and Performance*,
805 4(4), 565–572. <https://doi.org/10.1037/0096-1523.4.4.565>
- 806 Martin, A., & Fedio, P. (1983). Word Production and Comprehension in Alzheimer’s Disease : The
807 Breakdown of Semantic Knowledge. *Brain and Language*, 19, 124–141.
- 808 Mathôt, S., Schreij, D., & Theeuwes, J. (2012). OpenSesame: An open-source, graphical
809 experiment builder for the social sciences. *Behavior Research Methods*, 44(2), 314–324.
810 <https://doi.org/10.3758/s13428-011-0168-7>
- 811 McBride, D. M., & Doshier, B. A. (2002). A comparison of conscious and automatic memory
812 processes for picture and word stimuli: A process dissociation analysis. *Consciousness and*
813 *Cognition*, 11(3), 423–460. [https://doi.org/10.1016/S1053-8100\(02\)00007-7](https://doi.org/10.1016/S1053-8100(02)00007-7)
- 814 McKhann, G., Drachman, D., Folstein, M., Katzman, R., Price, D., & Stadlan, E. M. (1984).
815 Clinical diagnosis of Alzheimer’s disease: report of the NINCDS-ADRDA Work Group under
816 the auspices of Department of Health and Human Services Task Force on Alzheimer’s
817 Disease. *Neurology*, 34(7), 939–944. <https://doi.org/10.1212/wnl.34.7.939>
- 818 Molitor, R. J., Ko, P. C., & Ally, B. A. (2015). Eye movements in Alzheimer’s disease. *Journal of*
819 *Alzheimer’s Disease*, 44(1), 1–12.
- 820 Monsch, A. U., Bondi, M. W., Butters, N., Paulsen, J. S., Salmon, D. P., Brugger, P., & Swenson,
821 M. R. (1994). A Comparison of Category and Letter Fluency in Alzheimer’s Disease and
822 Huntington’s Disease. *Neuropsychology*, 8(1), 25–30. [https://doi.org/10.1037/0894-](https://doi.org/10.1037/0894-4105.8.1.25)
823 4105.8.1.25
- 824 Moscovitch, M. (1992). Memory and working-with-memory: A component process model based on
825 modules and central systems. *Journal of Cognitive Neuroscience*, 4(3), 257–267.
826 <https://doi.org/10.1162/jocn.1992.4.3.257>
- 827 Moscovitch, M. (2008). The hippocampus as a “stupid,” domain-specific module: Implications for
828 theories of recent and remote memory, and of imagination. *Canadian Journal of Experimental*

- 829 *Psychology*, 62(1), 62–79. <https://doi.org/10.1037/1196-1961.62.1.62>
- 830 Mulatti, C., Calia, C., De Caro, M. F., & Della Sala, S. (2014). The cumulative semantic
831 interference effect in normal and pathological ageing. *Neuropsychologia*, 65, 125–130.
832 <https://doi.org/10.1016/j.neuropsychologia.2014.10.007>
- 833 Nakamura, H., Nakanishi, M., Hamanaka, T., Nakaaki, S., & Yoshida, S. (2000). Semantic priming
834 in patients with Alzheimer and semantic dementia. *Cortex*, 36(2), 151–162.
835 [https://doi.org/10.1016/S0010-9452\(08\)70521-5](https://doi.org/10.1016/S0010-9452(08)70521-5)
- 836 Nebes, R. D. (1989). Semantic memory in Alzheimer’s disease. *Psychological Bulletin*, 106(3),
837 377–394.
- 838 Nebes, R. D. (1992). Semantic memory dysfunction in Alzheimer’s disease: Disruption of semantic
839 knowledge or information-processing limitation? In L. R. Squire & N. Butters (Eds.),
840 *Neuropsychology of memory*, 2nd ed. (pp. 233–240). Retrieved from
841 [https://search.ebscohost.com/login.aspx?direct=true&db=psyh&AN=1992-98460-](https://search.ebscohost.com/login.aspx?direct=true&db=psyh&AN=1992-98460-020&lang=pt-pt&site=eds-live&scope=site)
842 [020&lang=pt-pt&site=eds-live&scope=site](https://search.ebscohost.com/login.aspx?direct=true&db=psyh&AN=1992-98460-020&lang=pt-pt&site=eds-live&scope=site)
- 843 Nelson, D. L., Reed, V. S., & McEvoy, C. L. (1977). Learning to order pictures and words: A
844 model of sensory and semantic encoding. *Journal of Experimental Psychology: Human*
845 *Learning & Memory*, 3(5), 485–497. <https://doi.org/10.1037/0278-7393.3.5.485>
- 846 Nuthmann, A., De Groot, F., Huettig, F., & Olivers, C. N. L. (2019). Extra-foveal attentional
847 capture by object semantics. *PLoS ONE*, 14(5), 1–19.
848 <https://doi.org/10.1371/journal.pone.0217051>
- 849 O’Connor, M. K., & Ally, B. A. (2010). Using stimulus form change to understand memorial
850 familiarity for pictures and words in patients with mild cognitive impairment and Alzheimer’s
851 disease. *Neuropsychologia*, 48(7), 2068–2074.
852 <https://doi.org/10.1016/j.neuropsychologia.2010.03.027>
- 853 Ober, B. A. (2002). RT and non-RT methodology for semantic priming research with Alzheimer’s
854 disease patients: A critical review. *Journal of Clinical and Experimental Neuropsychology*,

- 855 24(7), 883–911. <https://doi.org/10.1076/jcen.24.7.883.8384>
- 856 Oyama, A., Takeda, S., Ito, Y., Nakajima, T., Takami, Y., Takeya, Y., ... Morishita, R. (2019).
857 Novel Method for Rapid Assessment of Cognitive Impairment Using High-Performance Eye-
858 Tracking Technology. *Scientific Reports*, 9(1), 1–9. [https://doi.org/10.1038/s41598-019-](https://doi.org/10.1038/s41598-019-49275-x)
859 49275-x
- 860 Paivio, A. (1991). Dual coding theory: Retrospect and current status. *Canadian Journal of*
861 *Psychology/Revue Canadienne de Psychologie*, 45(3), 255–287.
862 <https://doi.org/10.1037/h0084295>
- 863 Paivio, A. (2014). Intelligence, dual coding theory, and the brain. *Intelligence*, 47, 141–158.
864 <https://doi.org/10.1016/j.intell.2014.09.002>
- 865 Paivio, A., & Csapo, K. (1973). Picture superiority in free recall: Imagery or dual coding? *Cognitive*
866 *Psychology*, 5(2), 176–206. [https://doi.org/10.1016/0010-0285\(73\)90032-7](https://doi.org/10.1016/0010-0285(73)90032-7)
- 867 Perri, R., Carlesimo, G. A., Zannino, G. D., Mauri, M., Muolo, B., Pettenati, C., & Caltagirone, C.
868 (2003). Intentional and automatic measures of specific-category effect in the semantic
869 impairment of patients with Alzheimer's disease. *Neuropsychologia*, 41(11), 1509–1522.
870 [https://doi.org/10.1016/S0028-3932\(03\)00075-7](https://doi.org/10.1016/S0028-3932(03)00075-7)
- 871 Perry, R. J., & Hodges, J. R. (1999). Attention and executive deficits in Alzheimer's disease. A
872 critical review. *Brain*, 122(3), 383–404. <https://doi.org/10.1093/brain/122.3.383>
- 873 Phillips, L. H., Della Sala, S., & Trivelli, C. (1996). Fluency deficits in patients with Alzheimer's
874 disease and frontal lobe lesions. *European Journal of Neurology*, 3(2), 102–108.
875 <https://doi.org/10.1111/j.1468-1331.1996.tb00200.x>
- 876 Porter, G., Leonards, U., Wilcock, G., Haworth, J., Troscianko, T., & Tales, A. (2010). New
877 insights into feature and conjunction search: II. Evidence from Alzheimer's disease. *Cortex*,
878 46(5), 637–649. <https://doi.org/10.1016/j.cortex.2009.04.014>
- 879 Ramzaoui, H., Faure, S., & Spotorno, S. (2018). Alzheimer's Disease, Visual Search, and
880 Instrumental Activities of Daily Living: A Review and a New Perspective on Attention and

- 881 Eye Movements. *Journal of Alzheimer's Disease*, 66(3), 901–925.
882 <https://doi.org/10.3233/JAD-180043>
- 883 Ricci, M., Graef, S., Blundo, C., & Miller, L. A. (2012). Using the Rey Auditory Verbal Learning
884 Test (RAVLT) to differentiate Alzheimer's dementia and behavioural variant fronto-temporal
885 dementia. *The Clinical Neuropsychologist*, 26(6), 926–941.
886 <https://doi.org/10.1080/13854046.2012.704073>
- 887 Rich, J. B., Park, N. W., Dopkins, S., & Brandt, J. (2002). What do Alzheimer's disease patients
888 know about animals? It depends on task structure and presentation format. *Journal of the*
889 *International Neuropsychological Society*, 8(1), 83–94.
890 <https://doi.org/10.1017/S1355617702811080>
- 891 Rogers, S. L., & Friedman, R. B. (2008). The underlying mechanisms of semantic memory loss in
892 Alzheimer's disease and semantic dementia. *Neuropsychologia*, 46(1), 12–21.
893 <https://doi.org/10.1016/j.neuropsychologia.2007.08.010>
- 894 Rösler, A., Mapstone, M., Hays-Wicklund, A., Gitelman, D. R., & Weintraub, S. (2005). The
895 “zoom lens” of focal attention in visual search: Changes in aging and Alzheimer's disease.
896 *Cortex*, 41(4), 512–519. [https://doi.org/10.1016/S0010-9452\(08\)70191-6](https://doi.org/10.1016/S0010-9452(08)70191-6)
- 897 Ruotolo, F., Kalénine, S., & Bartolo, A. (2020). Activation of manipulation and function knowledge
898 during visual search for objects. *Journal of Experimental Psychology: Human Perception and*
899 *Performance*, 46(1), 66–90. <https://doi.org/10.1037/xhp0000696>
- 900 Ryan, J. D., Shen, K., & Liu, Z. (2019). The intersection between the oculomotor and hippocampal
901 memory systems: empirical developments and clinical implications. *Annals of the New York*
902 *Academy of Sciences*, 1–27. <https://doi.org/10.1111/nyas.14256>
- 903 Saffran, E. M., Branch Coslett, H., Martin, N., & Boronat, C. B. (2003). Access to knowledge from
904 pictures but not words in a patient with progressive fluent aphasia. *Language and Cognitive*
905 *Processes*, 18(5–6), 725–757. <https://doi.org/10.1080/01690960344000107>
- 906 Salmon, D. P., Butters, N., & Chan, A. S. (1999). The deterioration of semantic memory in

- 907 Alzheimer's disease. *Canadian Journal of Experimental Psychology/Revue Canadienne de*
 908 *Psychologie Expérimentale*, 53(1), 108–117. <https://doi.org/10.1037/h0087303>
- 909 Sartucci, F., Borghetti, D., Bocci, T., Murri, L., Orsini, P., Porciatti, V., ... Domenici, L. (2010).
 910 Dysfunction of the magnocellular stream in Alzheimer's disease evaluated by pattern
 911 electroretinograms and visual evoked potentials. *Brain Research Bulletin*, 82(3–4), 169–176.
 912 <https://doi.org/10.1016/j.brainresbull.2010.04.001>
- 913 Schmidt, J., & Zelinsky, G. J. (2009). Search guidance is proportional to the categorical specificity
 914 of a target cue. *Quarterly Journal of Experimental Psychology*, 62(10), 1904–1914.
 915 <https://doi.org/10.1080/17470210902853530>
- 916 Seckin, M., Mesulam, M. M., Voss, J. L., Huang, W., Rogalski, E. J., & Hurley, R. S. (2016). Am I
 917 looking at a cat or a dog? Gaze in the semantic variant of primary progressive aphasia is
 918 subject to excessive taxonomic capture. *Journal of Neurolinguistics*, 37, 68–81.
 919 <https://doi.org/10.1016/j.jneuroling.2015.09.003>
- 920 Spaan, P. E. J., Raaijmakers, J. G. W., & Jonker, C. (2003). Alzheimer's disease versus normal
 921 ageing: A review of the efficiency of clinical and experimental memory measures. *Journal of*
 922 *Clinical and Experimental Neuropsychology*, 25(2), 216–233.
 923 <https://doi.org/10.1076/jcen.25.2.216.13638>
- 924 Spaan, P. E. J., Raaijmakers, J. G. W., & Jonker, C. (2005). Early assessment of dementia: The
 925 contribution of different memory components. *Neuropsychology*, 19(5), 629–640.
 926 <https://doi.org/10.1037/0894-4105.19.5.629>
- 927 Tales, A., Butler, S. R., Fossey, J., Gilchrist, I. D., Jones, R. W., & Troscianko, T. (2002). Visual
 928 search in Alzheimer's disease: A deficiency in processing conjunctions of features.
 929 *Neuropsychologia*, 40(12), 1849–1857. [https://doi.org/10.1016/S0028-3932\(02\)00073-8](https://doi.org/10.1016/S0028-3932(02)00073-8)
- 930 Tippet, L. J., Gendall, A., Farah, M. J., & Thompson-Schill, S. L. (2004). Selection Ability in
 931 Alzheimer's Disease: Investigation of a Component of Semantic Processing. *Neuropsychology*,
 932 18(1), 163–173. <https://doi.org/10.1037/0894-4105.18.1.163>

- 933 Walther, D., & Koch, C. (2006). Modeling attention to salient proto-objects. *Neural Networks*,
934 19(9), 1395–1407. <https://doi.org/10.1016/j.neunet.2006.10.001>
- 935 Wechsler, D. (2008). *Wechsler Adult Intelligence Scale - Fourth Edition (WAIS-IV)* (Fourth).
936 <https://doi.org/10.1037/t15169-000>
- 937 Whitehouse, A. J. O., Maybery, M. T., & Durkin, K. (2006). The development of the picture-
938 superiority effect. *British Journal of Developmental Psychology*, 24(4), 767–773.
939 <https://doi.org/10.1348/026151005X74153>
- 940 Yang, Q., Wang, T., Su, N., Xiao, S., & Kapoula, Z. (2013). Specific saccade deficits in patients
941 with Alzheimer’s disease at mild to moderate stage and in patients with amnesic mild
942 cognitive impairment. *Age*, 35(4), 1287–1298. <https://doi.org/10.1007/s11357-012-9420-z>
- 943 Zelinsky, G. J. (2003). Detecting changes between real-world objects using spatiochromatic filters.
944 *Psychonomic Bulletin and Review*, 10(3), 533–555. <https://doi.org/10.3758/BF03196516>
- 945 Zola, S. M., Manzanares, C. M., Clopton, P., Lah, J. J., & Levey, A. I. (2013). A behavioral task
946 predicts conversion to mild cognitive impairment and Alzheimer’s disease. *American Journal*
947 *of Alzheimer’s Disease and Other Dementias*, 28(2), 179–184.
948 <https://doi.org/10.1177/1533317512470484>
949
950

Tables

Table 1: Minimum, Maximum, Mean, and Standard Deviation (SD) of the Neuropsychological Tests' Scores (Raw) of the AD group.

Neuropsychological Test		Minimum	Maximum	Mean	SD
RAVLT	IR	6	31	19.90	7.97
	DR	0	4	1.15	1.39
WF		5	35	21.40	9.08
FAB		7	15	10.37	2.09
RCPM		9	31	17.32	5.09
CD		0	11	6.32	2.85
CDL		21	70	58.53	11.35

Note. RAVLT: Rey's auditory verbal learning test (IR, immediate recall; DR, delayed recall); WF: word fluency; FAB: frontal assessment battery; RCPM: Raven's coloured progressive matrices; CD: freehand copying of drawings; CDL: copying drawings with landmarks.

Table 2: Means, Standard Deviations (in parenthesis), t-values, and p-values for the Scores on MMSE, Word Reading Test, Object Naming Test, and Digit Span Forwards.

Variables		AD	Control	t-value	p-value
MMSE	Raw	20.85 (2.68)	28.40 (1.35)	- 11.24	< 0.001
	Corrected	21.46 (2.37)	28.86 (2.08)	- 11.36	< 0.001
WR		9.61 (0.50)	9.85 (0.37)	- 1.66	0.11
ON		9.56 (0.78)	9.85 (0.37)	- 1.46	0.16
DSF		4.83 (0.99)	5.65 (0.93)	- 2.62	0.01

Note. The word reading (WR) test, the object naming (ON) test, and the digit span forwards (DSF) were administered to 18 (out of 20) AD patients, and all the healthy controls.

Table 3: Linear and Generalised Linear Mixed-effects Model Outputs.

Dependent Variable	Predictor	β	SE	z-value or t-value	p-value
Accuracy	Intercept	2.21	0.15	14.50	< 0.001
	Semantic Relatedness	- 0.93	0.21	- 4.56	< 0.001
	Group	- 1.32	0.23	- 5.70	< 0.001
	Target	0.22	0.24	0.95	0.34
	Visual Similarity	- 0.36	0.20	- 1.80	0.07
	Semantic Relatedness:Target	1.58	0.34	4.67	< 0.001
Latency to first fixation (z-scores)	Intercept	0.02	0.04	0.41	0.68
	Group	0.04	0.05	0.74	0.46
	Target	- 0.31	0.05	- 6.06	< 0.001
	Semantic Relatedness	0.43	0.09	4.91	< 0.001
	Visual Similarity	0.15	0.08	1.80	0.07
	Group:Target	- 0.01	0.10	- 0.14	0.89
	Group:Semantic Relatedness	0.02	0.13	0.20	0.84
	Group:Visual Similarity	0.09	0.10	0.90	0.37
First-pass dwell time	Intercept	0.35	0.01	36.01	< 0.001
	Target	0.31	0.02	18.46	< 0.001
	Semantic Relatedness	- 0.11	0.01	- 7.53	< 0.001
	Group	- 0.03	0.02	- 2.30	0.02
	Semantic Relatedness:Target	- 0.05	0.02	- 2.30	0.02
	Semantic Relatedness:Group	0.05	0.02	2.26	0.02

Note. Predictors are listed in the table in the same order as they were entered in the models. The predictors used in the different models are: Semantic Relatedness (Unrelated = -.49, Related = .51), Group (Control = -.49, AD = .51), Target (Absent = -.48, Present = .52), and Visual Similarity (Dissimilar = -.51, Similar = .49). The z-value is reported for the generalised linear mixed-effects model with binomial link of the response accuracy, and t-values instead are reported for all other linear mixed-effects models (i.e., latency to first fixation and first-pass dwell time).

Figures

Figure 1. Experimental design and example of an object array, which included a critical object (e.g., *car*, highlighted in red) plus 4 distractors. In target-present trials, the target name cued the critical object as the target. In target-absent trials, the target name cued an object that was not visually depicted in the array, but it was semantically related to the critical object and thus to the distractors in the semantically related but not the unrelated condition (e.g., airplane). The written cues for the targets were presented in Italian.

Figure 2. Example of a trial. The target name was cued (“automobile” is the Italian translation for “car”) after the presentation of a fixation cross. Then, the same fixation cross appeared for 150 ms, followed by the object array. If the participant indicated that a target was found, the object array was replaced with a number array, and she/he was asked to remember the location of the object by verbally indicating the corresponding number in the array. When the participant responded that the target was not found, the object array was immediately followed by the next trial.

1042 *Figure 3.* Mean response accuracy (proportions) for the Control (Left Panel) and AD group (Right Panel), in Target-
1043 Present and Absent trials (on the x-axis), in the Semantically Unrelated (dark grey) vs. Related (light grey) condition.
1044 Error bars represent 95% confidence intervals around the mean.

1057 *Figure 4.* Mean latency to first fixation (z-scores) on the critical object for the control (Left Panel) and AD group
 1058 (Right Panel), in Target-Present and -Absent trials (on the x-axis), with the Semantic Relatedness of the critical object
 1059 marked using line types and colours (Unrelated: dark grey, solid line; Related: light grey, dashed line). Error bars
 1060 represent 95% confidence intervals around the mean.

Figure 5. Mean first-pass dwell time (proportions) on the critical object for the control (Left Panel) and AD group (Right Panel), in Target-Present and -Absent trials (on the x-axis), with the Semantic Relatedness of the critical object marked using line types and colours (Unrelated: dark grey, solid line; Related: light grey, dashed line). Error bars represent 95% confidence intervals around the mean.