



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### The trans-ancestral genomic architecture of glycemc traits

**Citation for published version:**

The Meta-Analysis of Glucose and Insulin-related Traits Consortium (MAGIC) 2021, 'The trans-ancestral genomic architecture of glycemc traits', *Nature Genetics*, vol. 53. <https://doi.org/10.1038/s41588-021-00852-9>

**Digital Object Identifier (DOI):**

[10.1038/s41588-021-00852-9](https://doi.org/10.1038/s41588-021-00852-9)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Nature Genetics

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# The Trans-Ancestral Genomic Architecture of Glycemic Traits

Ji Chen<sup>1,2#</sup>, Cassandra N. Spracklen<sup>3,4#</sup>, Gaëlle Marenne<sup>2,5#</sup>, Arushi Varshney<sup>6#</sup>, Laura J Corbin<sup>7,8#</sup>, Jian'an Luan<sup>9</sup>, Sara M Willems<sup>9</sup>, Ying Wu<sup>3</sup>, Xiaoshuai Zhang<sup>9,10</sup>, Momoko Horikoshi<sup>11,12,13</sup>, Thibaud S Boutin<sup>14</sup>, Reedik Mägi<sup>15</sup>, Johannes Waage<sup>16,17</sup>, Achilleas Pitsilides<sup>18</sup>, Ruifang Li-Gao<sup>19</sup>, Kei Hang Katie Chan<sup>20,21,22</sup>, Jie Yao<sup>23</sup>, Mila D Anasanti<sup>24</sup>, Audrey Y Chu<sup>25</sup>, Annique Claringbould<sup>26</sup>, Jani Heikkinen<sup>24</sup>, Jaeyoung Hong<sup>18</sup>, Jouke-Jan Hottenga<sup>27,28</sup>, Shaofeng Huo<sup>29</sup>, Marika A. Kaakinen<sup>30,24</sup>, Tin Louie<sup>31</sup>, Winfried März<sup>32,33,34</sup>, Hortensia Moreno-Macias<sup>35</sup>, Anne Ndungu<sup>12</sup>, Sarah C. Nelson<sup>31</sup>, Ilja M. Nolte<sup>36</sup>, Kari E North<sup>37</sup>, Chelsea K. Raulerson<sup>3</sup>, Debashree Ray<sup>38</sup>, Rebecca Rohde<sup>37</sup>, Denis Rybin<sup>18</sup>, Claudia Schurmann<sup>39,40</sup>, Xueling Sim<sup>41,42,43</sup>, Loz Southam<sup>2</sup>, Isobel D Stewart<sup>9</sup>, Carol A. Wang<sup>44</sup>, Yujie Wang<sup>37</sup>, Peitao Wu<sup>18</sup>, Weihua Zhang<sup>45,46</sup>, Tarunveer S. Ahluwalia<sup>16,17,47</sup>, Emil VR Appel<sup>48</sup>, Lawrence F. Bielak<sup>49</sup>, Jennifer A. Brody<sup>50</sup>, Noël P Burt<sup>51</sup>, Claudia P Cabrera<sup>52,53</sup>, Brian E Cade<sup>54,55</sup>, Jin Fang Chai<sup>41</sup>, Xiaoran Chai<sup>56,57</sup>, Li-Ching Chang<sup>58</sup>, Chien-Hsiun Chen<sup>58</sup>, Brian H Chen<sup>59</sup>, Kumaraswamy Naidu Chitrala<sup>60</sup>, Yen-Feng Chiu<sup>61</sup>, Hugoline G. de Haan<sup>19</sup>, Graciela E Delgado<sup>34</sup>, Ayse Demirkan<sup>62,30</sup>, Qing Duan<sup>3,63</sup>, Jorgen Engmann<sup>64</sup>, Segun A Fatumo<sup>65,66,67</sup>, Javier Gayán<sup>68</sup>, Franco Giulianini<sup>69</sup>, Jung Ho Gong<sup>20</sup>, Stefan Gustafsson<sup>70</sup>, Yang Hai<sup>71</sup>, Fernando P Hartwig<sup>72,7</sup>, Jing He<sup>73</sup>, Yoriko Heianza<sup>74</sup>, Tao Huang<sup>75</sup>, Alicia Huerta-Chagoya<sup>76,77</sup>, Mi Yeong Hwang<sup>78</sup>, Richard A. Jensen<sup>50</sup>, Takahisa Kawaguchi<sup>79</sup>, Katherine A Kentistou<sup>80,81</sup>, Young Jin Kim<sup>78</sup>, Marcus E Kleber<sup>34</sup>, Ishminder K Kooner<sup>46</sup>, Shuiqing Lai<sup>20</sup>, Leslie A Lange<sup>82</sup>, Carl D Langefeld<sup>83</sup>, Marie Lauzon<sup>23</sup>, Man Li<sup>84</sup>, Symen Ligthart<sup>62</sup>, Jun Liu<sup>62,85</sup>, Marie Loh<sup>86,45</sup>, Jirong Long<sup>87</sup>, Valeriya Lyssenko<sup>88,89</sup>, Massimo Mangino<sup>90,91</sup>, Carola Marzi<sup>92,93</sup>, May E Montasser<sup>94</sup>, Abhishek Nag<sup>12</sup>, Masahiro Nakatochi<sup>95</sup>, Damia Noce<sup>96</sup>, Raymond Noordam<sup>97</sup>, Giorgio Pistis<sup>98</sup>, Michael Preuss<sup>39,99</sup>, Laura Raffield<sup>3</sup>, Laura J. Rasmussen-Torvik<sup>100</sup>, Stephen S Rich<sup>101,102</sup>, Neil R Robertson<sup>11,12</sup>, Rico Rueedi<sup>103,104</sup>, Kathleen Ryan<sup>94</sup>, Serena Sanna<sup>98,26</sup>, Richa Saxena<sup>105,106,107</sup>, Katharina E Schraut<sup>80,81</sup>, Bengt Sennblad<sup>108</sup>, Kazuya Setoh<sup>79</sup>, Albert V Smith<sup>109,110</sup>, Lorraine Southam<sup>111,112</sup>, Thomas Sparsø<sup>48</sup>, Rona J Strawbridge<sup>113,114</sup>, Fumihiko Takeuchi<sup>115</sup>, Jingyi Tan<sup>23</sup>, Stella Trompet<sup>97,116</sup>, Erik van den Akker<sup>117,118,119</sup>, Peter J van der Most<sup>36</sup>, Niek Verweij<sup>120,121</sup>, Mandy Vogel<sup>122</sup>, Heming Wang<sup>54,55</sup>, Chaolong Wang<sup>123,124</sup>, Nan Wang<sup>125,126</sup>, Helen R Warren<sup>52,53</sup>, Wanqing Wen<sup>87</sup>, Tom Wilsgaard<sup>127</sup>, Andrew Wong<sup>128</sup>, Andrew R Wood<sup>1</sup>, Tian Xie<sup>36</sup>, Mohammad Hadi Zafarmand<sup>129,130</sup>, Jing-Hua Zhao<sup>131</sup>, Wei Zhao<sup>49</sup>, Najaf Amin<sup>62,85</sup>, Zorayr Arzumanyan<sup>23</sup>, Arne Astrup<sup>132</sup>, Stephan JL Bakker<sup>133</sup>, Damiano Baldassarre<sup>134,135</sup>, Marian Beekman<sup>117</sup>, Richard N Bergman<sup>136</sup>, Alain Bertoni<sup>137</sup>, Matthias Blüher<sup>138</sup>, Lori L. Bonnycastle<sup>139</sup>, Stefan R Bornstein<sup>140</sup>, Donald W Bowden<sup>141</sup>, Qiuyin Cai<sup>73</sup>, Archie Campbell<sup>142,143</sup>, Harry Campbell<sup>80</sup>, Yi Cheng Chang<sup>144,145,146</sup>, Eco J.C. de Geus<sup>27,28</sup>, Abbas Dehghan<sup>62</sup>, Shufa Du<sup>147</sup>, Gudny Eiriksdottir<sup>110</sup>, Aliko Eleni Farmaki<sup>148,149</sup>, Mattias Frånberg<sup>150</sup>, Christian Fuchsberger<sup>96</sup>, Yutang Gao<sup>151</sup>, Anette P Gjesing<sup>48</sup>, Anuj Goel<sup>152,12</sup>, Sohee Han<sup>78</sup>, Catharina A Hartman<sup>153</sup>, Christian Herder<sup>154,155,156</sup>, Andrew A. Hicks<sup>96</sup>, Chang-Hsun Hsieh<sup>157,158</sup>, Willa A. Hsueh<sup>159</sup>, Sahoko Ichiara<sup>160</sup>, Michiya Igase<sup>161</sup>, M. Arfan Ikram<sup>62</sup>, W. Craig Johnson<sup>31</sup>, Marit E Jørgensen<sup>17,162</sup>, Peter K Joshi<sup>80</sup>, Rita R Kalyani<sup>163</sup>, Fouad R. Kandeel<sup>164</sup>, Tomohiro Katsuya<sup>165,166</sup>, Chiea Chuen Khor<sup>124</sup>, Wieland Kiess<sup>122</sup>, Ivana Kolcic<sup>167</sup>, Teemu Kuulasmaa<sup>168</sup>, Johanna Kuusisto<sup>169</sup>, Kristi Läll<sup>15</sup>, Kelvin Lam<sup>23</sup>, Deborah A Lawlor<sup>170,8</sup>, Nanette R. Lee<sup>171,172</sup>, Rozenn N. Lemaitre<sup>50</sup>, Honglan Li<sup>173</sup>, Lifelines Cohort Study<sup>174</sup>, Shih-Yi Lin<sup>175,176,177</sup>, Jaana Lindström<sup>178</sup>, Allan Linneberg<sup>179,180</sup>, Jianjun Liu<sup>124,181</sup>, Carlos Lorenzo<sup>182</sup>, Tatsuaki Matsubara<sup>183</sup>, Fumihiko Matsuda<sup>79</sup>, Geltrude Mingrone<sup>184</sup>, Simon Mooijaart<sup>97</sup>, Sanghoon Moon<sup>78</sup>, Toru Nabika<sup>185</sup>, Girish N. Nadkarni<sup>39</sup>, Jerry L. Nadler<sup>186</sup>, Mari Nelis<sup>15</sup>, Matt J Neville<sup>11,187</sup>, Jill M Norris<sup>188</sup>, Yasumasa Ohya<sup>189</sup>, Annette Peters<sup>190,93,191</sup>, Patricia A. Peyser<sup>49</sup>, Ozren Polasek<sup>167,192</sup>, Qibin Qi<sup>193</sup>, Dennis Raven<sup>153</sup>, Dermot F Reilly<sup>194</sup>, Alex Reiner<sup>195</sup>, Fernando Rivadeneira<sup>196</sup>, Kathryn Roll<sup>23</sup>, Igor Rudan<sup>197</sup>, Charumathi Sabanayagam<sup>56,198</sup>, Kevin Sandow<sup>23</sup>, Naveed Sattar<sup>199</sup>, Annette Schürmann<sup>200,201</sup>, Jinxiu Shi<sup>202</sup>, Heather M Stringham<sup>43,42</sup>, Kent D. Taylor<sup>23</sup>, Tanya M. Teslovich<sup>203</sup>, Betina Thuesen<sup>179</sup>, Paul RHJ Timmers<sup>80,204</sup>, Elena Tremoli<sup>135</sup>, Michael Y Tsai<sup>205</sup>, Andre Uitterlinden<sup>196</sup>, Rob M van Dam<sup>41,181,206</sup>, Diana van Heemst<sup>97</sup>, Astrid van Hylckama Vlieg<sup>19</sup>, Jana V Van Vliet-Ostaptchouk<sup>36</sup>, Jagadish Vangipurapu<sup>207</sup>, Henrik Vestergaard<sup>48,208</sup>, Tao Wang<sup>193</sup>, Ko Willems van Dijk<sup>209,210,211</sup>, Tatijana Zemunik<sup>212</sup>, Goncalo R Abecasis<sup>43</sup>, Linda S. Adair<sup>147,213</sup>, Carlos Alberto Aguilar-

Salinas<sup>214,215,216</sup>, Marta E Alarcón-Riquelme<sup>217,218</sup>, Ping An<sup>219</sup>, Larissa Aviles-Santa<sup>220</sup>, Diane M  
 Becker<sup>221</sup>, Lawrence J Beilin<sup>222</sup>, Sven Bergmann<sup>103,104,223</sup>, Hans Bisgaard<sup>16</sup>, Corri Black<sup>224</sup>, Michael  
 Boehnke<sup>43,42</sup>, Eric Boerwinkle<sup>225,226</sup>, Bernhard O Böhm<sup>227,228</sup>, Klaus Bønnelykke<sup>16</sup>, D I. Boomsma<sup>27,28</sup>,  
 Erwin P. Bottinger<sup>39,229,230</sup>, Thomas A Buchanan<sup>231,232,126</sup>, Mickaël Canouil<sup>233,234</sup>, Mark J Caulfield<sup>52,53</sup>,  
 John C. Chambers<sup>86,45,46,235,236</sup>, Daniel I. Chasman<sup>69,237</sup>, Yii-Der Ida Chen<sup>23</sup>, Ching-Yu Cheng<sup>56,198</sup>, Francis  
 S. Collins<sup>139</sup>, Adolfo Correa<sup>238</sup>, Francesco Cucca<sup>98</sup>, H. Janaka de Silva<sup>239</sup>, George Dedoussis<sup>240</sup>, Sölve  
 Elmståhl<sup>241</sup>, Michele K. Evans<sup>242</sup>, Ele Ferrannini<sup>243</sup>, Luigi Ferrucci<sup>244</sup>, Jose C Florez<sup>245,246,107</sup>, Paul W  
 Franks<sup>89,247</sup>, Timothy M Frayling<sup>1</sup>, Philippe Froguel<sup>233,234,248</sup>, Bruna Gigante<sup>249</sup>, Mark O. Goodarzi<sup>250</sup>,  
 Penny Gordon-Larsen<sup>147,213</sup>, Harald Grallert<sup>92,93</sup>, Niels Grarup<sup>48</sup>, Sameline Grimsgaard<sup>127</sup>, Leif  
 Groop<sup>251,252</sup>, Vilundur Gudnason<sup>110,253</sup>, Xiuqing Guo<sup>23</sup>, Anders Hamsten<sup>114</sup>, Torben Hansen<sup>48</sup>,  
 Caroline Hayward<sup>204</sup>, Susan R. Heckbert<sup>254</sup>, Bernardo L Horta<sup>72</sup>, Wei Huang<sup>202</sup>, Erik Ingelsson<sup>255</sup>,  
 Pankow S James<sup>256</sup>, Marjo-Ritta Jarvelin<sup>257,258,259,260</sup>, Jost B Jonas<sup>261,262,263</sup>, J. Wouter Jukema<sup>116,264</sup>,  
 Pontiano Kaleebu<sup>265</sup>, Robert Kaplan<sup>193,195</sup>, Sharon L.R. Kardia<sup>49</sup>, Norihiro Kato<sup>115</sup>, Sirkka M. Keinänen-  
 Kiukaanniemi<sup>266,267</sup>, Bong-Jo Kim<sup>78</sup>, Mika Kivimäki<sup>268</sup>, Heikki A. Koistinen<sup>269,270,271</sup>, Jaspal S.  
 Kooner<sup>46,235,236,272</sup>, Antje Körner<sup>122</sup>, Peter Kovacs<sup>138,273</sup>, Diana Kuh<sup>128</sup>, Meena Kumari<sup>274</sup>, Zoltan  
 Kutalik<sup>275,104</sup>, Markku Laakso<sup>169</sup>, Timo A. Lakka<sup>276,277,278</sup>, Lenore J Launer<sup>60</sup>, Karin Leander<sup>279</sup>, Huaixing  
 Li<sup>29</sup>, Xu Lin<sup>29</sup>, Lars Lind<sup>280</sup>, Cecilia Lindgren<sup>12,281,282</sup>, Simin Liu<sup>20</sup>, Ruth J.F. Loos<sup>39,99</sup>, Patrik KE  
 Magnusson<sup>283</sup>, Anubha Mahajan<sup>12</sup>, Andres Metspalu<sup>15</sup>, Dennis O Mook-Kanamori<sup>19,284</sup>, Trevor A  
 Mori<sup>222</sup>, Patricia B Munroe<sup>52,53</sup>, Inger Njølstad<sup>127</sup>, Jeffrey R O'Connell<sup>94</sup>, Albertine J Oldehinkel<sup>153</sup>, Ken  
 K Ong<sup>9</sup>, Sandosh Padmanabhan<sup>285</sup>, Colin N.A. Palmer<sup>286</sup>, Nicholette D Palmer<sup>141</sup>, Oluf Pedersen<sup>48</sup>,  
 Craig E Pennell<sup>44</sup>, David J Porteous<sup>142,287</sup>, Peter P. Pramstaller<sup>96</sup>, Michael A. Province<sup>219</sup>, Bruce M.  
 Psaty<sup>50,254,288</sup>, Lu Qi<sup>289</sup>, Leslie J. Raffel<sup>290</sup>, Rainer Rauramaa<sup>278</sup>, Susan Redline<sup>54,55</sup>, Paul M Ridker<sup>69,291</sup>,  
 Frits R. Rosendaal<sup>19</sup>, Timo E. Saaristo<sup>292,293</sup>, Manjinder Sandhu<sup>294</sup>, Jouko Saramies<sup>295</sup>, Neil  
 Schneiderman<sup>296</sup>, Peter Schwarz<sup>140,297,201</sup>, Laura J. Scott<sup>43,42</sup>, Elizabeth Selvin<sup>38</sup>, Peter Sever<sup>272</sup>, Xiao-ou  
 Shu<sup>87</sup>, P Eline Slagboom<sup>117</sup>, Kerrin S Small<sup>90</sup>, Blair H Smith<sup>298</sup>, Harold Snieder<sup>36</sup>, Tamar Sofer<sup>299,246</sup>,  
 Thorkild I.A. Sørensen<sup>48,300,7,8</sup>, Tim D Spector<sup>90</sup>, Alice Stanton<sup>301</sup>, Claire J Steves<sup>90,302</sup>, Michael  
 Stumvoll<sup>138</sup>, Liang Sun<sup>29</sup>, Yasuharu Tabara<sup>79</sup>, E Shyong Tai<sup>181,41,303</sup>, Nicholas J Timpson<sup>7,8</sup>, Anke  
 Tönjes<sup>138</sup>, Jaakko Tuomilehto<sup>304,305,306</sup>, Teresa Tusie<sup>77,307</sup>, Matti Uusitupa<sup>308</sup>, Pim van der Harst<sup>120,26</sup>,  
 Cornelia van Duijn<sup>85,62</sup>, Veronique Vitart<sup>204</sup>, Peter Vollenweider<sup>309</sup>, Tanja GM Vrijkotte<sup>129</sup>, Lynne E  
 Wagenknecht<sup>310</sup>, Mark Walker<sup>311</sup>, Ya X Wang<sup>262</sup>, Nick J Wareham<sup>9</sup>, Richard M Watanabe<sup>125,232,126</sup>,  
 Hugh Watkins<sup>152,12</sup>, Wen B Wei<sup>312</sup>, Ananda R Wickremasinghe<sup>313</sup>, Gonneke Willemsen<sup>27,28</sup>, James F  
 Wilson<sup>80,204</sup>, Tien-Yin Wong<sup>56,198</sup>, Jer-Yuarn Wu<sup>58</sup>, Anny H Xiang<sup>314</sup>, Lisa R Yanek<sup>221</sup>, Loïc Yengo<sup>315</sup>,  
 Mitsuhiro Yokota<sup>316</sup>, Eleftheria Zeggini<sup>111,317,318</sup>, Wei Zheng<sup>87</sup>, Alan B Zonderman<sup>60</sup>, Jerome I Rotter<sup>23</sup>,  
 Anna L Gloyn<sup>11,12,187,319</sup>, Mark I. McCarthy<sup>11,320,187,12@</sup>, Josée Dupuis<sup>18</sup>, James B Meigs<sup>321,246,107</sup>, Robert A  
 Scott<sup>9</sup>, Inga Prokopenko<sup>30,24</sup>, Aaron Leong<sup>322,323,237</sup>, Ching-Ti Liu<sup>18</sup>, Stephen CJ Parker<sup>6,324#</sup>, Karen L.  
 Mohlke<sup>3#</sup>, Claudia Langenberg<sup>9#</sup>, Eleanor Wheeler<sup>2,9#</sup>, Andrew P. Morris<sup>325,326,327,12#</sup>, Inês Barroso<sup>1,2,9#</sup>  
 on behalf of the Meta-Analysis of Glucose and Insulin-related Traits Consortium (MAGIC)

<sup>1</sup>Exeter Centre of Excellence for Diabetes Research (EXCEED), Genetics of Complex Traits, University  
 of Exeter Medical School, University of Exeter, Exeter, UK, <sup>2</sup>Department of Human Genetics,  
 Wellcome Sanger Institute, Hinxton, Cambridge, UK, <sup>3</sup>Department of Genetics, University of North  
 Carolina, Chapel Hill, NC, USA, <sup>4</sup>Department of Biostatistics and Epidemiology, University of  
 Massachusetts, Amherst, MA, USA, <sup>5</sup>Inserm, Univ Brest, EFS, UMR 1078, GGB, Brest, France,  
<sup>6</sup>Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI,  
 USA, <sup>7</sup>MRC Integrative Epidemiology Unit, University of Bristol, Bristol, UK, <sup>8</sup>Department of  
 Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK, <sup>9</sup>MRC  
 Epidemiology Unit, Institute of Metabolic Science, University of Cambridge, Cambridge, UK,  
<sup>10</sup>Department of Biostatistics, School of Public Health, Shandong University, Jinan, Shandong, China,  
<sup>11</sup>Oxford Centre for Diabetes, Endocrinology and Metabolism, Radcliffe Department of Medicine,  
 University of Oxford, Oxford, UK, <sup>12</sup>Wellcome Centre for Human Genetics, University of Oxford,  
 Oxford, UK, <sup>13</sup>Laboratory for Genomics of Diabetes and Metabolism, RIKEN Centre for Integrative

103 Medical Sciences, Yokohama, Japan, <sup>14</sup>Medical Research Council Human Genetics Unit, Institute for  
 104 Genetics and Molecular Medicine, Edinburgh, UK, <sup>15</sup>Estonian Genome Center, Institute of Genomics,  
 105 University of Tartu, Tartu, Estonia, <sup>16</sup>COPSAC, Copenhagen Prospective Studies on Asthma in  
 106 Childhood, Herlev and Gentofte Hospital, University of Copenhagen, Copenhagen, Denmark, <sup>17</sup>Steno  
 107 Diabetes Center Copenhagen, Gentofte, Denmark, <sup>18</sup>Department of Biostatistics, Boston University  
 108 School of Public Health, Boston, MA, USA, <sup>19</sup>Department of Clinical Epidemiology, Leiden University  
 109 Medical Center, Leiden, The Netherlands, <sup>20</sup>Department of Epidemiology, Brown University School of  
 110 Public Health, Brown University, Providence, RI, USA, <sup>21</sup>Department of Biomedical Sciences, City  
 111 University of Hong Kong, Hong Kong SAR, China, <sup>22</sup>Department of Electrical Engineering, City  
 112 University of Hong Kong, Hong Kong SAR, China, <sup>23</sup>The Institute for Translational Genomics and  
 113 Population Sciences, Department of Pediatrics, The Lundquist Institute for Biomedical Innovation at  
 114 Harbor-UCLA Medical Center, Torrance, CA, USA, <sup>24</sup>Department of Metabolism, Digestion, and  
 115 Reproduction, Imperial College London, London, UK, <sup>25</sup>Division of Preventive Medicine, Brigham and  
 116 Women's Hospital, Boston, MA, USA, <sup>26</sup>Department of Genetics, University of Groningen, University  
 117 Medical Center Groningen, Groningen, The Netherlands, <sup>27</sup>Department of Biological Psychology,  
 118 Faculty of Behaviour and Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam, The  
 119 Netherlands, <sup>28</sup>Amsterdam Public Health Research Institute, Amsterdam Universities Medical Center,  
 120 Amsterdam, The Netherlands, <sup>29</sup>CAS Key Laboratory of Nutrition, Metabolism and Food Safety,  
 121 Shanghai Institute of Nutrition and Health, University of Chinese Academy of Sciences, Chinese  
 122 Academy of Sciences, Shanghai, China, <sup>30</sup>Section of Statistical Multi-omics, Department of Clinical  
 123 and Experimental Research, University of Surrey, Guildford, Surrey, UK, <sup>31</sup>Department of  
 124 Biostatistics, University of Washington, Seattle, WA, USA, <sup>32</sup>SYNLAB Academy, SYNLAB Holding  
 125 Deutschland GmbH, Mannheim, Germany, <sup>33</sup>Clinical Institute of Medical and Chemical Laboratory  
 126 Diagnostics, Medical University Graz, Graz, Austria, <sup>34</sup>Vth Department of Medicine (Nephrology,  
 127 Hypertensiology, Rheumatology, Endocrinology, Diabetology), Medical Faculty Mannheim,  
 128 Heidelberg University, Mannheim, Baden-Württemberg, Germany, <sup>35</sup>Department of Economics,  
 129 Metropolitan Autonomous University, Mexico City, Mexico, <sup>36</sup>Department of Epidemiology,  
 130 University of Groningen, University Medical Center Groningen, Groningen, The Netherlands, <sup>37</sup>CVD  
 131 Genetic Epidemiology Computational Laboratory, Gillings School of Global Public Health, University  
 132 of North Carolina, Chapel Hill, NC, USA, <sup>38</sup>Department of Epidemiology, Johns Hopkins Bloomberg  
 133 School of Public Health, Baltimore, MD, USA, <sup>39</sup>The Charles Bronfman Institute for Personalized  
 134 Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA, <sup>40</sup>HPI Digital Health Center,  
 135 Digital Health and Personalized Medicine, Hasso Plattner Institute, Potsdam, Germany, <sup>41</sup>Saw Swee  
 136 Hock School of Public Health, National University of Singapore and National University Health  
 137 System, Singapore, Singapore, <sup>42</sup>Center for Statistical Genetics, University of Michigan, Ann Arbor,  
 138 MI, USA, <sup>43</sup>Department of Biostatistics, School of Public Health, University of Michigan, Ann Arbor,  
 139 MI, USA, <sup>44</sup>School of Medicine and Public Health, College of Health, Medicine and Wellbeing, The  
 140 University of Newcastle, Newcastle, NSW, Australia, <sup>45</sup>Department of Epidemiology and Biostatistics,  
 141 Imperial College London, London, UK, <sup>46</sup>Department of Cardiology, Ealing Hospital, London North  
 142 West Healthcare NHS Trust, Middlesex, UK, <sup>47</sup>The Bioinformatics Centre, Department of Biology,  
 143 University of Copenhagen, Copenhagen, Denmark, <sup>48</sup>Novo Nordisk Foundation Center for Basic  
 144 Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen,  
 145 Copenhagen, Denmark, <sup>49</sup>Department of Epidemiology, School of Public Health, University of  
 146 Michigan, Ann Arbor, MI, USA, <sup>50</sup>Department of Medicine, Cardiovascular Health Research Unit,  
 147 University of Washington, Seattle, WA, USA, <sup>51</sup>Metabolism Program, Program in Medical and  
 148 Population Genetics, Broad Institute, Cambridge, MA, USA, <sup>52</sup>Department of Clinical Pharmacology,  
 149 William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen  
 150 Mary University of London, London, UK, <sup>53</sup>NIHR Barts Cardiovascular Biomedical Research Centre,  
 151 Queen Mary University of London, London, UK, <sup>54</sup>Department of Medicine, Sleep and Circadian  
 152 Disorders, Brigham and Women's Hospital, Boston, MA, USA, <sup>55</sup>Department of Medicine, Sleep  
 153 Medicine, Harvard Medical School, Boston, MA, USA, <sup>56</sup>Ocular Epidemiology, Singapore Eye Research

154 Institute, Singapore National Eye Centre, Singapore, Singapore, <sup>57</sup>Department of Ophthalmology,  
 155 National University of Singapore and National University Health System, Singapore, Singapore,  
 156 <sup>58</sup>Institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan, Taiwan, <sup>59</sup>Department of  
 157 Epidemiology, The Herbert Wertheim School of Public Health and Human Longevity Science, UC San  
 158 Diego, La Jolla, CA, USA, <sup>60</sup>Laboratory of Epidemiology and Population Sciences, National Institute on  
 159 Aging, National Institutes of Health, Baltimore, MD, USA, <sup>61</sup>Institute of Population Health Sciences,  
 160 National Health Research Institutes, Miaoli, Taiwan, <sup>62</sup>Department of Epidemiology, Erasmus Medical  
 161 Center, Rotterdam, The Netherlands, <sup>63</sup>Department of Statistics, University of North Carolina at  
 162 Chapel Hill, Chapel Hill, NC, USA, <sup>64</sup>Institute of Cardiovascular Science, UCL, London, UK, <sup>65</sup>Uganda  
 163 Medical Informatics Centre (UMIC), MRC/UVR and London School of Hygiene & Tropical Medicine  
 164 (Uganda Research Unit), Entebbe, Uganda, <sup>66</sup>London School of Hygiene & Tropical Medicine, London,  
 165 UK, <sup>67</sup>H3Africa Bioinformatics Network (H3ABioNet) Node, Centre for Genomics Research and  
 166 Innovation, NABDA/FMST, Abuja, Nigeria, <sup>68</sup>Bioinfosol, Sevilla, Spain, <sup>69</sup>Division of Preventive  
 167 Medicine, Brigham and Women's Hospital, Boston, MA, USA, <sup>70</sup>Department of Medical Sciences,  
 168 Molecular Epidemiology and Science for Life Laboratory, Uppsala University, Uppsala, Sweden,  
 169 <sup>71</sup>Department of Statistics, The University of Auckland, Science Center, Auckland, New Zealand,  
 170 <sup>72</sup>Postgraduate Program in Epidemiology, Federal University of Pelotas, Pelotas, RS, Brazil,  
 171 <sup>73</sup>Department of Medicine, Epidemiology, Vanderbilt University Medical Center, Nashville, TN, USA,  
 172 <sup>74</sup>Department of Epidemiology, Tulane University Obesity Research Center,, Tulane University, New  
 173 Orleans, USA, <sup>75</sup>Department of Epidemiology and Biostatistics, School of Public Health, Peking  
 174 University, Beijing, China, <sup>76</sup>Molecular Biology and Genomic Medicine Unit, National Council for  
 175 Science and Technology, Mexico City, Mexico, <sup>77</sup>Molecular Biology and Genomic Medicine Unit,  
 176 National Institute of Medical Sciences and Nutrition, Mexico City, Mexico, <sup>78</sup>Division of Genome  
 177 Science, Department of Precision Medicine, National Institute of Health, Cheongju-si,  
 178 Chungcheongbuk-do, South Korea, <sup>79</sup>Center for Genomic Medicine, Kyoto University Graduate  
 179 School of Medicine, Kyoto, Japan, <sup>80</sup>Centre for Global Health Research, Usher Institute, University of  
 180 Edinburgh, Edinburgh, Scotland, <sup>81</sup>Centre for Cardiovascular Sciences, Queen's Medical Research  
 181 Institute, University of Edinburgh, Edinburgh, Scotland, <sup>82</sup>Department of Medicine, Division of  
 182 Biomedical Informatics and Personalized Medicine, University of Colorado Anschutz Medical  
 183 Campus, Denver, CO, USA, <sup>83</sup>Department of Biostatistics and Data Science, Wake Forest School of  
 184 Medicine, Winston-Salem, NC, USA, <sup>84</sup>Department of Medicine, Division of Nephrology and  
 185 Hypertension, University of Utah, Salt Lake City, UT, USA, <sup>85</sup>Nuffield Department of Population  
 186 Health, University of Oxford, Oxford, UK, <sup>86</sup>Lee Kong Chian School of Medicine, Nanyang  
 187 Technological University, Singapore, Singapore, <sup>87</sup>Division of Epidemiology, Department of Medicine,  
 188 Vanderbilt Epidemiology Center, Vanderbilt University Medical Center, Nashville, TN, USA,  
 189 <sup>88</sup>Department of Clinical Science, Center for Diabetes Research, University of Bergen, Bergen,  
 190 Norway, <sup>89</sup>Department of Clinical Sciences, Lund University Diabetes Centre, Lund University,  
 191 Malmo, Sweden, <sup>90</sup>Department of Twin Research and Genetic Epidemiology, School of Life Course  
 192 Sciences, King's College London, London, UK, <sup>91</sup>NIHR Biomedical Research Centre, Guy's and St  
 193 Thomas' Foundation Trust, London, UK, <sup>92</sup>Institute of Epidemiology, Research Unit of Molecular  
 194 Epidemiology, Helmholtz Zentrum München Research Center for Environmental Health, Neuherberg,  
 195 Bavaria, Germany, <sup>93</sup>German Center for Diabetes Research (DZD), Neuherberg, Bavaria, Germany,  
 196 <sup>94</sup>Department of Medicine, Division of Endocrinology, Diabetes, and Nutrition, University of  
 197 Maryland School of Medicine, Baltimore, MD, USA, <sup>95</sup>Public Health Informatics Unit, Department of  
 198 Integrated Sciences, Nagoya University Graduate School of Medicine, Nagoya, Japan, <sup>96</sup>Institute for  
 199 Biomedicine, Eurac Research, Bolzano, BZ, Italy, <sup>97</sup>Department of Internal Medicine, Section of  
 200 Gerontology and Geriatrics, Leiden University Medical Center, Leiden, The Netherlands, <sup>98</sup>Istituto di  
 201 Ricerca Genetica e Biomedica (IRGB), Consiglio Nazionale delle Ricerche (CNR), Monserrato, Italy,  
 202 <sup>99</sup>The Mindich Child Health and Development Institute for Personalized Medicine, Icahn School of  
 203 Medicine at Mount Sinai, New York, NY, USA, <sup>100</sup>Department of Preventive Medicine, Northwestern  
 204 University Feinberg School of Medicine, Chicago, IL, USA, <sup>101</sup>Center for Public Health Genomics,

205 University of Virginia, Charlottesville, VA, USA, <sup>102</sup>Department of Public Health Sciences, University of  
 206 Virginia, Charlottesville, VA, USA, <sup>103</sup>Department of Computational Biology, University of Lausanne,  
 207 Lausanne, Switzerland, <sup>104</sup>Swiss Institute of Bioinformatics, Lausanne, Switzerland, <sup>105</sup>Center for  
 208 Genomic Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA,  
 209 <sup>106</sup>Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital,  
 210 Boston, MA, USA, <sup>107</sup>Program in Medical and Population Genetics,, Broad Institute, Cambridge, MA,  
 211 USA, <sup>108</sup>Department of Cell and Molecular Biology,, National Bioinformatics Infrastructure Sweden,,  
 212 Science for Life Laboratory, Uppsala University, Uppsala, Sweden, <sup>109</sup>Department of Biostatistics,  
 213 University of Michigan, Ann Arbor, MI, USA, <sup>110</sup>Icelandic Heart Association, Kopavogur, Iceland,  
 214 <sup>111</sup>Institute of Translational Genomics, Helmholtz Zentrum München – German Research Center for  
 215 Environmental Health, Neuherberg, Germany, <sup>112</sup>Wellcome Sanger Institute, Hinxton, Cambridge,  
 216 UK, <sup>113</sup>Institute of Health and Wellbeing, University of Glasgow, Glasgow, Glasgow, UK,  
 217 <sup>114</sup>Department of Medicine Solna, Cardiovascular medicine, Karolinska Institutet, Stockholm,  
 218 Sweden, <sup>115</sup>National Center for Global Health and Medicine, Tokyo, Japan, <sup>116</sup>Department of  
 219 Cardiology, Leiden University Medical Center, Leiden, The Netherlands, <sup>117</sup>Department of Biomedical  
 220 Data Sciences, Molecular Epidemiology, Leiden University Medical Center, Leiden, The Netherlands,  
 221 <sup>118</sup>Department of Pattern Recognition & Bioinformatics, Delft University of Technology, Delft, The  
 222 Netherlands, <sup>119</sup>Department of Biomedical Data Sciences, Leiden Computational Biology Center,  
 223 Leiden University Medical Center, Leiden, The Netherlands, <sup>120</sup>Department of Cardiology, University  
 224 of Groningen, University Medical Center Groningen, Groningen, The Netherlands, <sup>121</sup>Genomics plc,  
 225 Oxford, UK, <sup>122</sup>Center of Pediatric Research, University Children’s Hospital Leipzig, University of  
 226 Leipzig Medical Center, Leipzig, Germany, <sup>123</sup>Department of Epidemiology and Biostatistics, School of  
 227 Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan,  
 228 China, <sup>124</sup>Genome Institute of Singapore, Agency for Science, Technology and Research, Singapore,  
 229 Singapore, <sup>125</sup>Department of Preventive Medicine, Keck School of Medicine of USC, Los Angeles, CA,  
 230 USA, <sup>126</sup>USC Diabetes and Obesity Research Institute, Keck School of Medicine of USC, Los Angeles,  
 231 CA, USA, <sup>127</sup>Department of Community Medicine, Faculty of Health Sciences, UIT the Arctic  
 232 University of Norway, Tromsø, Norway, <sup>128</sup>MRC Unit for Lifelong Health & Ageing at UCL, London,  
 233 UK, <sup>129</sup>Department of Public Health, Amsterdam Public Health Research Institute, Amsterdam  
 234 Universities Medical Center, Amsterdam, The Netherlands, <sup>130</sup>Department of Clinical Epidemiology,  
 235 Biostatistics, and Bioinformatics, Amsterdam Public Health Research Institute, Amsterdam  
 236 Universities Medical Center, Amsterdam, The Netherlands, <sup>131</sup>Department of Public Health and  
 237 Primary Care, School of Clinical Medicine, University of Cambridge, Cambridge, UK, <sup>132</sup>Department of  
 238 Nutrition, Exercise, and Sports, Faculty of Science, University of Copenhagen, Copenhagen,  
 239 Denmark, <sup>133</sup>Department of Internal Medicine, University of Groningen, University Medical Center  
 240 Groningen, Groningen, The Netherlands, <sup>134</sup>Department of Medical Biotechnology and Translational  
 241 Medicine, University of Milan, Milan, Italy, <sup>135</sup>Centro Cardiologico Monzino, IRCCS, Milan, Italy,  
 242 <sup>136</sup>Diabetes and Obesity Research Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA,  
 243 <sup>137</sup>Department of Epidemiology and Prevention, Division of Public Health Sciences, Wake Forest  
 244 School of Medicine, Winston-Salem, NC, USA, <sup>138</sup>Medical Department III – Endocrinology,  
 245 Nephrology, Rheumatology, University of Leipzig Medical Center, Leipzig, Germany, <sup>139</sup>Medical  
 246 Genomics and Metabolic Genetics Branch, National Human Genome Research Institute, National  
 247 Institutes of Health, Bethesda, MD, USA, <sup>140</sup>Department for Prevention and Care of Diabetes, Faculty  
 248 of Medicine Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany, <sup>141</sup>Department  
 249 of Biochemistry, Wake Forest School of Medicine, Winston-Salem, NC, USA, <sup>142</sup>Centre for Genomic  
 250 and Experimental Medicine, Institute of Genetics & Molecular Medicine, University of Edinburgh,  
 251 Western General Hospital, Edinburgh, UK, <sup>143</sup>Usher Institute, University of Edinburgh, Edinburgh, UK,  
 252 <sup>144</sup>Department of Internal Medicine, National Taiwan University Hospital, Taipei, Taiwan, <sup>145</sup>Graduate  
 253 Institute of Medical Genomics and Proteomics, National Taiwan University, Taipei, Taiwan,  
 254 <sup>146</sup>Institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan, <sup>147</sup>Department of Nutrition,  
 255 Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC, USA,

<sup>148</sup>Department of Population Science and Experimental Medicine, Institute of Cardiovascular Science,  
 University College London, London, UK, <sup>149</sup>Department of Nutrition and Dietetics, School of Health  
 Science and Education, Harokopio University of Athens, Athens, Greece, <sup>150</sup>Department of Medicine  
 Solna, Cardiovascular medicine, Stockholm, Sweden, <sup>151</sup>Department of Epidemiology, Shanghai  
 Cancer Institute, Shanghai, China, <sup>152</sup>Division of Cardiovascular Medicine, Radcliffe Department of  
 Medicine, University of Oxford, Oxford, UK, <sup>153</sup>Department of Psychiatry, Interdisciplinary Center  
 Psychopathy and Emotion Regulation, University of Groningen, University Medical Center  
 Groningen, Groningen, The Netherlands, <sup>154</sup>Institute for Clinical Diabetology, German Diabetes  
 Center, Leibniz Center for Diabetes Research at Heinrich Heine University Düsseldorf, Düsseldorf,  
 Germany, <sup>155</sup>Division of Endocrinology and Diabetology, Medical Faculty, Heinrich Heine University  
 Düsseldorf, Düsseldorf, Germany, <sup>156</sup>German Center for Diabetes Research (DZD), Düsseldorf,  
 Germany, <sup>157</sup>Internal Medicine, Endocrine & Metabolism, Tri-Service General Hospital, Taipei,  
 Taiwan, <sup>158</sup>School of Medicine, National Defense Medical Center, Taipei, Taiwan, <sup>159</sup>Internal  
 Medicine, Endocrinology, Diabetes & Metabolism, Diabetes and Metabolism Research Center, The  
 Ohio State University Wexner Medical Center, Columbus, OH, USA, <sup>160</sup>Department of Environmental  
 and Preventive Medicine, Jichi Medical University School of Medicine, Shimotsuke, Japan,  
<sup>161</sup>Department of Anti-aging Medicine, Ehime University Graduate School of Medicine, Toon, Japan,  
<sup>162</sup>National Institute of Public Health, University of Southern Denmark, Odense, Denmark,  
<sup>163</sup>Department of Medicine, Endocrinology, Diabetes & Metabolism, Johns Hopkins University School  
 of Medicine, Baltimore, MD, USA, <sup>164</sup>Clinical Diabetes, Endocrinology & Metabolism, Translational  
 Research & Cellular Therapeutics, Beckman Research Institute of the City of Hope, Duarte, CA, USA,  
<sup>165</sup>Department of Clinical Gene Therapy, Osaka University Graduate School of Medicine, Suita, Japan,  
<sup>166</sup>Department of Geriatric and General Medicine, Osaka University Graduate School of Medicine,  
 Suita, Japan, <sup>167</sup>Department of Public Health, University of Split School of Medicine, Split, Croatia,  
<sup>168</sup>Institute of Biomedicine, Bioinformatics Center, University of Eastern Finland, Kuopio, Finland,  
<sup>169</sup>Department of Medicine, University of Eastern Finland and Kuopio University Hospital, Kuopio,  
 Finland, <sup>170</sup>MRC Integrative Epidemiology Unit, University of Bristol, Bristol, Bristol, UK, <sup>171</sup>USC-Office  
 of Population Studies Foundation, University of San Carlos, Cebu City, Philippines, <sup>172</sup>Department of  
 Anthropology, Sociology and History, University of San Carlos, Cebu City, Philippines, <sup>173</sup>State Key  
 Laboratory of Oncogene and Related Genes & Department of Epidemiology, Shanghai Cancer  
 Institute, Renji Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China,  
<sup>174</sup>Groningen, The Netherlands, <sup>175</sup>Internal Medicine, Endocrine & Metabolism, Taichung Veterans  
 General Hospital, Taichung, Taiwan, <sup>176</sup>Center for Geriatrics and Gerontology,, Taichung Veterans  
 General Hospital, Taichung, Taiwan, <sup>177</sup>National Defense Medical Center, National Yang-Ming  
 University, Taipei, Taiwan, <sup>178</sup>Diabetes Prevention Unit, National Institute for Health and Welfare,  
 Helsinki, Finland, <sup>179</sup>Center for Clinical Research and Prevention, Bispebjerg and Frederiksberg  
 Hospital, Copenhagen, Denmark, <sup>180</sup>Department of Clinical Medicine, Faculty of Health and Medical  
 Sciences, University of Copenhagen, Copenhagen, Denmark, <sup>181</sup>Yong Loo Lin School of Medicine,  
 National University of Singapore and National University Health System, Singapore, Singapore,  
<sup>182</sup>Department of Medicine, University of Texas Health Sciences Center, San Antonio, TX, USA,  
<sup>183</sup>Department of Internal Medicine, Aichi Gakuin University School of Dentistry, Nagoya, Japan,  
<sup>184</sup>Department of Diabetes, Diabetes, & Nutritional Sciences, James Black Centre, King's College  
 London, London, UK, <sup>185</sup>Department of Functional Pathology, Shimane University School of  
 Medicine, Izumo, Japan, <sup>186</sup>Department of Medicine and Pharmacology, New York Medical College  
 School of Medicine, Valhalla, NY, USA, <sup>187</sup>Oxford NIHR Biomedical Research Centre, Oxford University  
 Hospitals NHS Foundation Trust, Oxford, UK, <sup>188</sup>Colorado School of Public Health, University of  
 Colorado Anschutz Medical Campus, Aurora, CO, USA, <sup>189</sup>Department of Geriatric Medicine and  
 Neurology, Ehime University Graduate School of Medicine, Toon, Japan, <sup>190</sup>Institute of Epidemiology,  
 Helmholtz Zentrum München Research Center for Environmental Health, Neuherberg, Bavaria,  
 Germany, <sup>191</sup>Institute for Medical Information Processing, Biometry, and Epidemiology, Ludwig-  
 Maximilians University Munich, Munich, Bavaria, Germany, <sup>192</sup>Gen-info Ltd, Zagreb, Croatia,

<sup>193</sup>Department of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, NY, USA, <sup>194</sup>Genetics and Pharmacogenomics, Merck Sharp & Dohme Corp., Kenilworth, NJ, USA, <sup>195</sup>Department of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA, USA, <sup>196</sup>Department of Internal Medicine, Erasmus Medical Center, Rotterdam, The Netherlands, <sup>197</sup>Centre for Global Health, The Usher Institute, University of Edinburgh, Edinburgh, UK, <sup>198</sup>Ophthalmology & Visual Sciences Academic Clinical Program (Eye ACP), Duke-NUS Medical School, Singapore, Singapore, <sup>199</sup>BHF Glasgow Cardiovascular Research Centre, Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, UK, <sup>200</sup>Department of Experimental Diabetology, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany, <sup>201</sup>German Center for Diabetes Research (DZD e.V.), Neuherberg, Germany, <sup>202</sup>Department of Genetics, Shanghai-MOST Key Laboratory of Health and Disease Genomics, Chinese National Human Genome Center at Shanghai (CHGC) and Shanghai Academy of Science & Technology (SAST), Shanghai, China, <sup>203</sup>Sarepta Therapeutics, Cambridge, Massachusetts, USA, <sup>204</sup>Medical Research Council Human Genetics Unit, Institute for Genetics and Cancer, University of Edinburgh, Edinburgh, UK, <sup>205</sup>Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, MN, USA, <sup>206</sup>Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA, USA, <sup>207</sup>Institute of Clinical Medicine, Internal Medicine, University of Eastern Finland, Kuopio, Finland, <sup>208</sup>Department of Medicine, Bornholms Hospital, Rønne, Denmark, <sup>209</sup>Department of Internal Medicine, Division of Endocrinology, Leiden University Medical Center, Leiden, The Netherlands, <sup>210</sup>Laboratory for Experimental Vascular Medicine, Leiden University Medical Center, Leiden, The Netherlands, <sup>211</sup>Department of Human Genetics, Leiden University Medical Center, Leiden, The Netherlands, <sup>212</sup>Department of Human Biology, University of Split School of Medicine, Split, Croatia, <sup>213</sup>Carolina Population Center, University of North Carolina, Chapel Hill, NC, USA, <sup>214</sup>Department of Endocrinology and Metabolism, Instituto Nacional de Ciencias Medicas y Nutricion, Mexico City, Mexico, <sup>215</sup>Unidad de Investigación de Enfermedades Metabólicas, Instituto Nacional de Ciencias Médicas y Nutrición and Tec Salud, Mexico City, Mexico, <sup>216</sup>Instituto Tecnológico y de Estudios Superiores de Monterrey Tec Salud, Mexico City, Mexico, <sup>217</sup>Department of Medical Genomics, Pfizer/University of Granada/Andalusian Government Center for Genomics and Oncological Research (GENYO), Granada, Spain, <sup>218</sup>Institute for Environmental Medicine, Chronic Inflammatory Diseases, Karolinska Institutet, Solna, Sweden, <sup>219</sup>Department of Genetics, Division of Statistical Genomics, Washington University School of Medicine, St. Louis, MO, USA, <sup>220</sup>Clinical and Health Services Research, National Institute on Minority Health and Health Disparities, Bethesda, MD, USA, <sup>221</sup>Department of Medicine, General Internal Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, USA, <sup>222</sup>Medical School, Royal Perth Hospital Unit, University of Western Australia, Perth, WA, Australia, <sup>223</sup>Department of Integrative Biomedical Sciences, University of Cape Town, Cape Town, South Africa, <sup>224</sup>Aberdeen Centre for Health Data Science, 1:042 Polwarth Building,, School of Medicine, Medical, Science and Nutrition, University of Aberdeen, Foresterhill, Aberdeen, UK, <sup>225</sup>Human Genetics Center, School of Public Health, The University of Texas Health Science Center at Houston, Houston, TX, USA, <sup>226</sup>Human Genome Sequencing Center, Baylor College of Medicine, Houston, TX, USA, <sup>227</sup>Division of Endocrinology and Diabetes, Graduate School of Molecular Endocrinology and Diabetes, University of Ulm, Ulm, Baden-Württemberg, Germany, <sup>228</sup>LKC School of Medicine, Nanyang Technological University, Singapore and Imperial College London, UK, Singapore, <sup>229</sup>Hasso Plattner Institute for Digital Health at Mount Sinai, Icahn School of Medicine at Mount Sinai, New York, NY, USA, <sup>230</sup>Digital Health Center, Hasso Plattner Institut, University Potsdam, Potsdam, Germany, <sup>231</sup>Department of Medicine, Keck School of Medicine of USC, Los Angeles, CA, USA, <sup>232</sup>Department of Physiology and Neuroscience, Keck School of Medicine of USC, Los Angeles, CA, USA, <sup>233</sup>INSERM UMR 1283 / CNRS UMR 8199, European Institute for Diabetes (EGID), Université de Lille, Lille, France, <sup>234</sup>INSERM UMR 1283 / CNRS UMR 8199, European Institute for Diabetes (EGID), Institut Pasteur de Lille, Lille, France, <sup>235</sup>Imperial College Healthcare NHS Trust, Imperial College London, London, UK, <sup>236</sup>MRC-PHE Centre for Environment and Health, Imperial College London, London, UK, <sup>237</sup>Harvard Medical School, Boston,



358 MA, USA, <sup>238</sup>Department of Medicine, Jackson Heart Study, University of Mississippi Medical Center,  
 359 Jackson, MS, USA, <sup>239</sup>Department of Medicine, Faculty of Medicine, University of Kelaniya, Ragama,  
 360 Sri Lanka, <sup>240</sup>Department of Nutrition and Dietetics, School of Health Science and Education,  
 361 Harokopio University of Athens, Kallithea, Greece, <sup>241</sup>Department of Clinical Sciences, Lund  
 362 University, Malmö, Sweden, <sup>242</sup>Laboratory of Epidemiology and Population Sciences, National  
 363 Institute on Aging Intramural Research Program, National Institutes of Health, Baltimore, MD, USA,  
 364 <sup>243</sup>CNR Institute of Clinical Physiology, Pisa, Italy, <sup>244</sup>Intramural Research Program, National Institute  
 365 of Aging, Baltimore, MD, USA, <sup>245</sup>Diabetes Unit and Center for Genomic Medicine, Massachusetts  
 366 General Hospital, Boston, MA, USA, <sup>246</sup>Department of Medicine, Harvard Medical School, Boston,  
 367 MA, USA, <sup>247</sup>Department of Public Health and Clinical Medicine, Umeå University, Umeå, Sweden,  
 368 <sup>248</sup>Department of Genomics of Common Disease, Imperial College London, London, UK,  
 369 <sup>249</sup>Department of Medicine, Cardiovascular medicine, Karolinska Institutet, Stockholm, Sweden,  
 370 <sup>250</sup>Department of Medicine, Division of Endocrinology, Diabetes & Metabolism, Cedars-Sinai Medical  
 371 Center, Los Angeles, CA, USA, <sup>251</sup>Diabetes Centre, Lund University, Sweden, <sup>252</sup>Finnish Institute of  
 372 Molecular Medicine, Helsinki University, Helsinki, Finland, <sup>253</sup>Faculty of Medicine, School of health  
 373 sciences, University of Iceland, Reykjavik, Iceland, <sup>254</sup>Department of Epidemiology, Cardiovascular  
 374 Health Research Unit, University of Washington, Seattle, WA, USA, <sup>255</sup>Department of Medicine,  
 375 Division of Cardiovascular Medicine, Stanford University School of Medicine, Stanford University,  
 376 Stanford, CA, USA, <sup>256</sup>Division of Epidemiology and Community Health, University of Minnesota,  
 377 Minneapolis, MN, USA, <sup>257</sup>Department of Epidemiology and Biostatistics, MRC-PHE Centre for  
 378 Environment and Health, School of Public Health, Imperial College London, London, UK, <sup>258</sup>Center for  
 379 Life Course Health Research, Faculty of Medicine, University of Oulu, Oulu, Finland, <sup>259</sup>Unit of  
 380 Primary Health Care, Oulu University Hospital, OYS, Oulu, Finland, <sup>260</sup>Department of Life Sciences,  
 381 College of Health and Life Sciences, Brunel University London, London, UK, <sup>261</sup>Department of  
 382 Ophthalmology, Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany, <sup>262</sup>Beijing  
 383 Institute of Ophthalmology, Beijing Ophthalmology and Visual Science Key Lab, Beijing Tongren Eye  
 384 Center, Beijing Tongren Hospital, Capital Medical University, Beijing, China, <sup>263</sup>Institute of Molecular  
 385 and Clinical Ophthalmology Basel IOB, Basel, Switzerland, <sup>264</sup>Netherlands Heart Institute, Utrecht,  
 386 The Netherlands, <sup>265</sup>MRC/UVRI and LSHTM (Uganda Research Unit), Entebbe, Uganda, <sup>266</sup>Faculty of  
 387 Medicine, Institute of Health Sciences, University of Oulu, Oulu, Finland, <sup>267</sup>Unit of General Practice,  
 388 Oulu University Hospital, Oulu, Finland, <sup>268</sup>Department of Epidemiology and Public Health, UCL,  
 389 London, UK, <sup>269</sup>Department of Public Health Solutions, Finnish Institute for Health and Welfare,  
 390 Helsinki, Finland, <sup>270</sup>Department of Medicine, University of Helsinki and Helsinki University Central  
 391 Hospital, Helsinki, Finland, <sup>271</sup>Minerva Foundation Institute for Medical Research, Helsinki, Finland,  
 392 <sup>272</sup>National Heart and Lung Institute, Imperial College London, London, UK, <sup>273</sup>IFB Adiposity Diseases,  
 393 University of Leipzig Medical Center, Leipzig, Germany, <sup>274</sup>Institute for Social and Economic Research,  
 394 University of Essex, Colchester, UK, <sup>275</sup>University Institute of Primary Care and Public Health, Division  
 395 of Biostatistics, University of Lausanne, Lausanne, Switzerland, <sup>276</sup>Institute of Biomedicine, School of  
 396 Medicine, University of Eastern Finland, Finland, <sup>277</sup>Department of Clinical Physiology and Nuclear  
 397 Medicine, Kuopio University Hospital, Kuopio, Finland, <sup>278</sup>Foundation for Research in Health Exercise  
 398 and Nutrition, Kuopio Research Institute of Exercise Medicine, Kuopio, Finland, <sup>279</sup>Institute of  
 399 Environmental Medicine, Cardiovascular and Nutritional Epidemiology, Karolinska Institutet,  
 400 Stockholm, Sweden, <sup>280</sup>Department of Medical Sciences, Uppsala, Sweden, <sup>281</sup>Big Data Institute,  
 401 Nuffield Department of Medicine, University of Oxford, Oxford, UK, <sup>282</sup>Nuffield Department of  
 402 Women's and Reproductive Health, University of Oxford, Oxford, UK, <sup>283</sup>Department of Medical  
 403 Epidemiology and Biostatistics and the Swedish Twin Registry, Karolinska Institutet, Stockholm,  
 404 Sweden, <sup>284</sup>Department of Public Health and Primary Care, Leiden University Medical Center, Leiden,  
 405 The Netherlands, <sup>285</sup>Institute of Cardiovascular and Medical Sciences, University of Glasgow,  
 406 Glasgow, UK, <sup>286</sup>Division of Population Health and Genomics, School of Medicine, University of  
 407 Dundee, Ninewells Hospital and Medical School, Dundee, UK, <sup>287</sup>Centre for Cognitive Ageing and  
 408 Cognitive Epidemiology, University of Edinburgh, Edinburgh, UK, <sup>288</sup>Department of Health Services,

Cardiovascular Health Research Unit, University of Washington, Seattle, WA, USA, <sup>289</sup>Department of Epidemiology, Tulane University School of Public Health and Tropical Medicine, New Orleans, LA, USA, <sup>290</sup>Department of Pediatrics, Genetic and Genomic medicine, University of California, Irvine, Irvine, CA, USA, <sup>291</sup>Havard Medical School, Boston, MA, USA, <sup>292</sup>Tampere, Finnish Diabetes Association, Tampere, Finland, <sup>293</sup>Pirkanmaa Hospital District, Tampere, Finland, <sup>294</sup>Department of Medicine, University of Cambridge, Cambridge, UK, <sup>295</sup>South Karelia Central Hospital, Lappeenranta, Finland, <sup>296</sup>Department of Psychology, University of Miami, Miami, FL, USA, <sup>297</sup>Paul Langerhans Institute Dresden of the Helmholtz Center Munich, University Hospital and Faculty of Medicine, Dresden, Germany, <sup>298</sup>Division of Population Health and Genomics, Ninewells Hospital and Medical School, University of Dundee, Dundee, UK, <sup>299</sup>Division of Sleep and Circadian Disorders, Brigham and Women's Hospital, Boston, MA, USA, <sup>300</sup>Department of Public Health, Section of Epidemiology, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark, <sup>301</sup>Department of Molecular and Cellular Therapeutics, Royal College of Surgeons in Ireland, Dublin, Ireland, <sup>302</sup>Department of Aging and Health, Guy's and St Thomas' Foundation Trust, London, UK, <sup>303</sup>Cardiovascular and Metabolic Disease Signature Research Program, Duke-NUS Medical School, Singapore, Singapore, <sup>304</sup>Department of Public Health Solutions, National Institute for Health and Welfare, Helsinki, Finland, <sup>305</sup>Department of Public Health, University of Helsinki, Helsinki, Finland, <sup>306</sup>Saudi Diabetes Research Group, King Abdulaziz University, Jeddah, Saudi Arabia, <sup>307</sup>Department of Genomic Medicine and Environmental Toxicology, Instituto de Investigaciones Biomedicas, Universidad Nacional Autonoma de Mexico, Mexico City, Mexico, <sup>308</sup>Department of Public Health and Clinical Nutrition, University of Eastern Finland, Finland, <sup>309</sup>Department of Medicine, Internal Medicine, Lausanne University Hospital (CHUV), Lausanne, Switzerland, <sup>310</sup>Department of Public Health Sciences, Wake Forest School of Medicine, Winston-Salem, NC, USA, <sup>311</sup>Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK, <sup>312</sup>Beijing Tongren Eye Center, Beijing Key Laboratory of Intraocular Tumor Diagnosis and Treatment, Beijing Ophthalmology & Visual Sciences Key Lab, Beijing Tongren Hospital, Capital Medical University, Beijing, China, China, <sup>313</sup>Department of Public Health, Faculty of Medicine, University of Kelaniya, Ragama, Sri Lanka, <sup>314</sup>Department of Research and Evaluation, Kaiser Permanente of Southern California, Pasadena, CA, USA, <sup>315</sup>Institute for Molecular Bioscience, The University of Queensland, Queensland, Australia, <sup>316</sup>Kurume University School of Medicine, Japan, <sup>317</sup>Wellcome Sanger Institute, Hinxton, UK, <sup>318</sup>TUM School of Medicine, Technical University of Munich and Klinikum Rechts der Isar, Munich, Germany, <sup>319</sup>Department of Pediatrics, Division of Endocrinology, Stanford School of Medicine, Stanford, CA, USA, <sup>320</sup>Wellcome Centre for Human Genetics, Nuffield Department of Medicine, University of Oxford, Oxford, UK, <sup>321</sup>Department of Medicine, Division of General Internal Medicine, Massachusetts General Hospital, Boston, MA, USA, <sup>322</sup>Department of Medicine, General Internal Medicine, Massachusetts General Hospital, Boston, MA, USA, <sup>323</sup>Department of Medicine, Diabetes Unit and Endocrine Unit, Massachusetts General Hospital, Boston, MA, USA, <sup>324</sup>Department of Human Genetics, University of Michigan, Ann Arbor, MI, USA, <sup>325</sup>Centre for Genetics and Genomics Versus Arthritis, Division of Musculoskeletal and Dermatological Sciences, The University of Manchester, Manchester, UK, <sup>326</sup>Centre for Musculoskeletal Research, Division of Musculoskeletal and Dermatological Sciences, The University of Manchester, Manchester, UK, <sup>327</sup>Department of Biostatistics, University of Liverpool, Liverpool, UK

# Denote shared authorship contributions

@ Current address: Genentech, South San Francisco, CA

**\*Corresponding author:** Inês Barroso, Exeter Centre of Excellence for Diabetes Research (EXCEED), Exeter Medical School, University of Exeter, Exeter, UK, +44 1392 408221, [ines.barroso@exeter.ac.uk](mailto:ines.barroso@exeter.ac.uk)

# **Abstract**

Glycemic traits are used to diagnose and monitor type 2 diabetes, and cardiometabolic health. To date, most genetic studies of glycemic traits have focused on individuals of European ancestry. Here, we aggregated genome-wide association studies in up to 281,416 individuals without diabetes (30% non-European ancestry) with fasting glucose, 2h-glucose post-challenge, glycated hemoglobin, and fasting insulin data. Trans-ancestry and single-ancestry meta-analyses identified 242 loci (99 novel;  $P < 5 \times 10^{-8}$ ), 80% with no significant evidence of between-ancestry heterogeneity. Analyses restricted to European ancestry individuals with equivalent sample size would have led to 24 fewer new loci. Compared to single-ancestry, equivalent sized trans-ancestry fine-mapping reduced the number of estimated variants in 99% credible sets by a median of 37.5%. Genomic feature, gene-expression and gene-set analyses revealed distinct biological signatures for each trait, highlighting different underlying biological pathways. Our results increase understanding of diabetes pathophysiology by use of trans-ancestry studies for improved power and resolution.

Fasting glucose (FG), 2h-glucose post-challenge (2hGlu), and glycated hemoglobin (HbA1c) are glycemic traits used to diagnose diabetes<sup>1</sup>. In addition, HbA1c is the most commonly used biomarker to monitor glucose control in patients with diabetes. Fasting insulin (FI) reflects a combination of insulin secretion and insulin resistance, both components of type 2 diabetes (T2D), and insulin clearance<sup>2</sup>. Collectively, all four glycemic traits are useful to better understand T2D pathophysiology<sup>3-5</sup> and cardiometabolic outcomes<sup>6</sup>.

To date, genome-wide association studies (GWAS) and analysis of Metabochip and exome arrays have identified >120 loci associated with glycemic traits in individuals without diabetes<sup>7-15</sup>. However, despite considerable differences in the prevalence of T2D risk factors across ancestries<sup>16-18</sup>, most glycemic trait GWAS have insufficient representation of individuals of non-European ancestry. Additionally, they have limited resolution for fine-mapping of causal variants and for effector transcript identification. Here, we present large-scale trans-ancestry meta-analyses of GWAS for four glycemic traits in individuals without diabetes. We aimed to identify additional glycemic trait-associated loci; investigate the portability of loci and genetic scores across ancestries; leverage differences in effect allele frequency (EAF), effect size, and linkage disequilibrium (LD) across diverse populations to conduct fine-mapping and aid causal variant/effector transcript identification; and compare the genetic architecture of glycemic traits to further identify the cell-types and target tissues most influenced by these traits which inform T2D pathophysiology.

## Results

### Study design and definitions

To identify loci associated with glycemic traits FG, 2hGlu, FI, and HbA1c, we aggregated GWAS in up to 281,416 individuals without diabetes, ~30% of whom were of non-European ancestry [13% East Asian, 7% Hispanic, 6% African-American, 3% South Asian, and 2% sub-Saharan African (Ugandan data only available for HbA1c)]. Each cohort imputed data to the 1000 Genomes Project reference panel<sup>19</sup> (phase 1 v3, March 2012, or later; **Methods, Supplementary Table 1, Extended Data Figure 1, Supplementary Note**). Up to ~49.3 million variants were directly genotyped or imputed, with between 38.6 million (2hGlu) and 43.5 million variants (HbA1c) available for analysis after exclusions based on minor allele count (MAC < 3) and imputation quality (imputation  $r^2$  or INFO score < 0.40) in each cohort. FG, 2hGlu and FI analyses were adjusted for BMI<sup>15</sup> but for simplicity they are abbreviated as FG, 2hGlu and FI (**Methods**).

We first performed trait-specific fixed-effect meta-analyses *within* each ancestry using METAL<sup>20</sup> (**Methods**). We defined “single-ancestry lead” variants as the strongest trait-associated variants ( $P < 5 \times 10^{-8}$ ) within a 1Mb region in an ancestry (**Table 1**). Within each ancestry and each autosome, we used approximate conditional analyses in GCTA<sup>21,22</sup>, to identify “single-ancestry index variants” ( $P < 5 \times 10^{-8}$ ) that exert conditionally distinct effects on the trait (**Table 1, Methods, Supplementary Note**). This approach identified 124 FG, 15 2hGlu, 48 FI and 139 HbA1c variants that were significant in at least one ancestry (**Supplementary Table 2**).

Next, we conducted trait-specific *trans-ancestry* meta-analyses using MANTRA (**Methods, Supplementary Table 1, Supplementary Note**) to identify genome-wide significant “trans-ancestry lead variants”, defined as the most significant trait-associated variant across all ancestries ( $\log_{10}$  Bayes Factor [BF] > 6, equivalent to  $P < 5 \times 10^{-8}$ )<sup>23</sup> (**Table 1, Methods**). Here, we present trans-ancestry results as our primary results (**Supplementary Table 2**).

Causal variants are expected to affect related glycemic traits and may be shared across ancestries. Therefore, we combined all single-ancestry lead variants, single-ancestry index variants, and/or trans-ancestry lead variants (for any trait) mapping within 500Kb of each other, into a single “trans-ancestry locus” bounded by 500Kb flanking sequences (**Table 1, Extended Data Figure 2**). As defined,

a trans-ancestry locus may contain multiple causal variants affecting one or more glycemic traits, exerting their effect in one or more ancestry.

### Glycemic trait locus discovery

Trans-ancestry meta-analyses identified 235 trans-ancestry loci, of which 59 contained lead variants for more than one trait. In addition, we identified seven “single-ancestry loci” that did not contain any trans-ancestry lead variants (**Table 1, Supplementary Table 2**). Of the 242 combined loci, 99 (including 6 of the 7 single-ancestry) had not been previously associated with any of the four glycemic traits or with T2D, at the time of analysis (**Figure 1, Supplementary Table 3, Supplementary note**). However, based on recent East Asian and trans-ancestry T2D GWAS meta-analyses<sup>23-27</sup>, the lead variants at 27/99 novel glycemic trait loci have strong evidence of association with T2D ( $P < 10^{-4}$ ; 13 loci with  $P < 5 \times 10^{-8}$ ), suggesting they are also important in T2D pathophysiology (**Supplementary Tables 2 and 4**).

Of the six single-ancestry novel loci, three were unique to non-European ancestry individuals (**Supplementary Table 3**). An African American association for FI (lead variant rs12056334) near *LOC100128993* (an uncharacterized RNA gene; **Supplementary Note**), an African American association for FG (lead variant rs61909476) near *ETS1* and a Hispanic association for FG (lead variant rs12315677) within *PIK3C2G* (**Supplementary Table 3**). Despite broadly similar EAF across ancestries, rs61909476 was significantly associated with FG only in African American individuals (EAF ~7%,  $b = 0.0812$  mmol/l,  $SE = 0.01$  mmol/l,  $P = 3.9 \times 10^{-8}$  vs EAF 10-17%,  $b = 0.002$  mmol/l,  $se = 0.003$ - $0.017$  mmol/l,  $P = 0.44$ - $0.95$  in all other ancestries, **Supplementary table 2, Supplementary note**). The nearest gene, *ETS1*, encodes a transcription factor that is expressed in mouse pancreatic  $\beta$ -cells, and its overexpression decreases glucose-stimulated insulin secretion in mouse islets<sup>28</sup>. Located within the *PIK3C2G* gene, rs12315677 has an 84% EAF in Hispanic (70-94% in other ancestries) and is significantly associated with FG in this ancestry alone ( $b = 0.0387$  mmol/l,  $SE = 0.0075$  mmol/l,  $P = 4.0 \times 10^{-8}$  vs  $b = -0.0128$ - $0.010$  mmol/l,  $SE = 0.003$ - $0.018$  mmol/l,  $P = 0.14$ - $0.76$  in all other ancestries, **Supplementary note**). In mice, deletion of *Pik3c2g* leads to a phenotype characterized by reduced glycogen storage in the liver, hyperlipidemia, adiposity, and insulin resistance with increasing age, or after a high fat diet<sup>29</sup>. Instances of similar EAFs but differing effect sizes between populations, could be due to genotype-by-environment or other epistatic effects. Alternatively, lower imputation accuracy in smaller sample sizes could deflate effect sizes, although imputation quality for these variants was good (average  $r^2 = 0.81$ ). Finally, the variants detected here may be in LD with ancestry-specific causal variants not interrogated here that differ in frequency across ancestries. However, we could not find evidence of rarer alleles in the cognate populations from the 1000G project (**Supplementary Table 5**). The final three single-ancestry loci were identified in individuals of European ancestry (**Supplementary note**).

Next, by rescaling the standard errors of allelic effect sizes to artificially boost the sample size of the European meta-analysis to match that of trans-ancestry meta-analysis, we determined that 21 of the novel trans-ancestry loci would not have been discovered with an equivalent sample size comprised exclusively of European ancestry individuals (**Supplementary note**). Their discovery was due to the higher EAF and/or larger effect size in non-European ancestry populations. In particular, two loci (near *LINC00885* and *MIR4278*) contain East Asian and African American single-ancestry lead variants, respectively, suggesting that these specific ancestries may be driving the trans-ancestry discovery (**Supplementary Tables 2-3**). Combined with the three single-ancestry non-European loci described above, our results show that 24% (24/99) of novel loci were discovered due to the contribution of non-European ancestry participants, strengthening the argument for expanding genetic studies in diverse populations.

### Allelic architecture of glycemic traits

Single-ancestry and trans-ancestry results combined increased the number of established loci for FG to 102 (182 signals, 53 novel loci), FI to 66 (95 signals, 49 novel loci), 2hGlu to 21 (28 signals, 11 novel loci), and HbA1c to 127 (218 signals, 62 novel loci) (**Supplementary Table 2**), with significant overlap across traits (**Extended Data Figure 3**). We also detected ( $P < 0.05$  or  $\log_{10}BF > 0$ ) the vast majority (~90%) of previously established glycemic signals, 70-88% of which attained genome-wide significance (**Supplementary Note, Supplementary Table 6**). Given that analyses for FG, FI, and 2hGlu were performed adjusted for BMI, we confirmed that collider bias did not influence >98% of signals discovered (**Supplementary note**)<sup>31</sup>. As expected, given the greater power due to increased sample sizes, new association signals tended to have smaller effect sizes and/or EAFs in European ancestry individuals compared to established signals (**Extended Data Figure 4**).

#### Characterization of lead variants across ancestries

To better understand the transferability of trans-ancestry lead variants across ancestries, we investigated the pairwise EAF correlation and the pairwise summarized heterogeneity of effect sizes between ancestries<sup>32</sup> (**Methods, Supplementary Note**). Consistent with population history and evolution, these results demonstrated considerable EAF correlation ( $\rho^2 > 0.70$ ) between European and Hispanic, European and South Asian, and Hispanic and South Asian populations, consistent across all four traits, and between African Americans and Ugandans for HbA1c (**Extended Data Figure 5**). Despite significant EAF correlations, some pairwise comparisons exhibited strong evidence for effect size heterogeneity between ancestries that was less consistent between traits (**Extended Data Figure 5**). However, sensitivity analyses demonstrated that, across all comparisons, the evidence for heterogeneity is driven by a small number of variants, with between 81.5% (for HbA1c) and 85.7% of trans-ancestry lead variants (for FG) showing no evidence for trans-ancestry heterogeneity ( $P > 0.05$ ) (**Supplementary Note**).

#### Trait variance explained by associated loci

The trait variance explained by genome-wide significant loci was assessed using the single-ancestry variants only or a combination of single-ancestry and trans-ancestry variants (**Supplementary Table 7**) with betas extracted from the relevant single-ancestry meta-analysis results (**Methods**). The variance explained was assessed by linear regression in a subset of the contributing cohorts (**Methods, Supplementary Tables 8-11**). In general, the approach that explained the most variance was to begin with the trans-ancestry lead variants that had  $P < 0.1$  in the relevant single-ancestry meta-analysis, then add in all single-ancestry variants that were not in LD with the trans-ancestry variants ( $LD\ r^2 < 0.1$ ) (List C, **Supplementary Tables 8-11, Figure 2**). Using this approach, the mean variance in the trait distribution explained was between 0.7% (2hGlu in EUR) and 6% (HbA1c in AA). The European-based estimates explained more variance relative to previous estimates of 2.8% for FG and 1.7% for HbA1c<sup>33</sup> (**Supplementary Note**).

#### Transferability of EUR ancestry-derived polygenic scores

To investigate the transferability of polygenic scores across ancestries we used the PRS-CSauto software<sup>34</sup> to first build polygenic scores for each glycemic trait based on European ancestry data. However, the training set for 2hGlu was too small so this trait was excluded. To build the polygenic scores (PGS), for each trait we first removed five of the largest European cohorts from the European ancestry meta-analysis. These five cohorts were meta-analyzed and used as our European ancestry test dataset, for each trait. The remaining European ancestry cohorts were also meta-analyzed and used as the training dataset, from which we derived a PGS for each trait (**Methods**). We used PRS-CSauto to revise the effect size estimates for the variants in the score (obtained from the training European datasets) based on the LD of the test population. PRS-CSauto does not have LD reference panels for South Asian or Hispanic ancestry and as such we were unable to test the transferability of the PGS into those populations. The “gtx” package<sup>35</sup> (**Methods**) was used to obtain the  $R^2$  for each test population (**Figure 3, Supplementary Table 12**). Consistent with other complex traits<sup>36</sup>, the

European ancestry-derived PGS had greater predictive power into test data of European ancestry than other ancestry groups.

### Fine-mapping

We fine-mapped, 231 trans-ancestry and six single-ancestry autosomal loci (**Supplementary Table 2, Supplementary note**). Using FINEMAP with ancestry-specific LD and an average LD matrix across ancestries, we conducted fine-mapping both within (161 loci with single-ancestry lead variants) and across ancestries (231 loci) for each trait (**Methods**). Because 59 of the 231 trans-ancestry loci were associated with more than one trait, we conducted trans-ancestry fine-mapping for a total of 305 locus-trait associations. Of these 305 locus-trait combinations, FINEMAP estimated the presence of a single causal variant at 186 loci (61%), while multiple distinct causal variants were implicated at 126 loci (39%), for a total of 464 causal variants (**Figure 4A**).

### Credible sets for causal variants

At each locus, we next constructed credible sets (CS) for each causal variant that account for  $\geq 99\%$  of the posterior probability of association (PPA). We identified 21 locus-trait associations (at 19 loci) for which the 99% CS included a single variant, and we highlight four examples (**Methods, Supplementary Note, Figure 4B, Supplementary Table 13**).

At *MTNR1B* and *SIX3* we identified, respectively, rs10830963 (PPA>0.999, for both HbA1c and FG) and rs12712928 (PPA=0.997, for FG) as the likely causal variants. At both loci previous studies confirm these variants affect transcriptional activity<sup>37,38,39</sup> (**Supplementary note**). At a locus near *PFKM* associated with HbA1c, trans-ancestry fine-mapping identified rs12819124 (PPA>0.999) as the likely causal variant. This variant has been previously associated with mean corpuscular hemoglobin<sup>40</sup>, suggesting an effect on HbA1c via the red blood cell (RBC, **Supplementary note**). At *HBB*, we identified rs334 (PPA>0.999; Glu7Val) as the likely causal variant associated with HbA1c. rs334 is a causal variant of sickle cell anemia<sup>41</sup>, previously associated with urinary albumin-to-creatinine ratio in Caribbean Hispanic individuals<sup>42</sup>, severe malaria in a Tanzanian study population<sup>43</sup>, hematocrit and mean corpuscular volume in Hispanic/Latino populations<sup>44</sup>, and RBC distribution in Ugandan individuals<sup>45</sup>, all pointing to a variant effect on HbA1c via non-glycemic pathways.

The remaining locus-trait associations with a single variant in the 99% CS (**Supplementary Table 13**) point to variants that could be prioritized for functional follow-up to elucidate impact on glycemic trait physiology.

At an additional 156 locus-trait associations trans-ancestry fine-mapping identified 99% CS with 50 or fewer variants (**Figure 4B, Supplementary Table 13**). Consistent with the potential for >1 causal variant in a locus, 74 locus-trait associations contained 88 variants with PPA>0.90 that are strong candidate causal variants (**Supplementary Table 14**). For example, 10 are coding variants including several missense such as the *HBB* Glu7Val mentioned above, *GCKR* Leu446Pro, *RREB1* Asp1771Asn, *G6PC2* Pro324Ser, *GLP1R* Ala316Thr, and *TMPRSS6* Val736Ala, each of which have been proposed or shown to affect gene function<sup>12,46-50</sup>. We additionally identified *AMPD3* Val311Leu (PPA=0.989) and *TMC6* Trp125Arg (PPA>0.999) variants associated with HbA1c which were previously detected in an exome array analysis but had not been fine-mapped with certainty due to the absence of backbone GWAS data<sup>30</sup>. Our fine-mapping now suggest these variants are likely causal and identify their cognate genes as effector transcripts.

Finally, we evaluated the resolution obtained in the trans-ancestry versus single-ancestry fine-mapping (**Methods, Supplementary Note**). We compared the number of variants in 99% CS across 98 locus-trait associations which, as suggested by FINEMAP, had a single causal variant in both trans-ancestry and single-ancestry analyses. Fine-mapping within and across ancestries was conducted

using the same set of variants. At 8 of 98 locus-trait associations single-ancestry fine-mapping identified a single variant in the CS. In addition, at 72 of the 98 locus-trait associations, the number of variants in the 99% CS was smaller in the trans-ancestry fine-mapping (**Figure 4C**), which likely reflects the larger sample size and differences in LD structure, EAFs, and effect sizes across diverse populations. To quantify the estimated improvement in fine-mapping resolution attributable to the multi-ancestry GWAS, we then compared 99% CS sizes from the trans-ancestry fine-mapping to single-ancestry-specific data emulating the same total sample size by rescaling the standard errors (**Methods**). Of the 72 locus-trait associations with estimated improved fine-mapping in trans-ancestry analysis, resolution at 38 (53%) was improved because of the larger sample size in the trans-ancestry fine-mapping analysis (**Figure 4C**), and this estimated improved resolution would likely have been obtained in a European-only fine-mapping effort with equivalent sample size. However, at 34 (47%) loci, the inclusion of samples from multiple diverse populations yielded the estimated improved resolution. On average, ancestry differences led to a reduction in the median number of variants in the 99% CS from 24 to 15 variants (37.5% median reduction; **Figure 4C**), demonstrating the value of conducting fine-mapping across ancestries.

### HbA1c Signal Classification

HbA1c-associated variants can exert their effects on HbA1c levels through both glycemic and non-glycemic pathways<sup>7,51</sup> and their correct classification can affect T2D diagnostic accuracy<sup>7,52</sup>. Using prior association results for other glycemic, RBC, and iron traits, and a fuzzy clustering approach we classified variants into their most likely mode of action (**Methods, Supplementary note**). Of the 218 HbA1c-associated variants, 27 (12%) could not be characterized due to missing data and 23 (11%) could not be classified into a “known” class (**Supplementary note**). The remaining signals were classified as principally: a) glycemic (n=53; 24%), b) affecting iron levels/metabolism (n=12; 6%), or c) RBC traits (n=103; 47%). A genetic risk score (GRS) composed of all HbA1c-associated signals was strongly associated with T2D risk (OR=2.4, 95% CI 2.3-2.5,  $P=2.7 \times 10^{-298}$ ). However, when using partitioned GRSs composed of these different classes of variants (**Methods**), we found the T2D association was mainly driven by variants influencing HbA1c through glycemic pathways (OR=2.6, 95% CI 2.5-2.8,  $P=2.3 \times 10^{-250}$ ), with weaker evidence of association (despite the larger number of variants in the GRS) and a more modest risk (OR=1.4, 95% CI 1.2-1.7,  $P=4.7 \times 10^{-4}$ ) imparted by signals in the mature RBC cluster that were not glycemic (i.e. where those specific variants had  $P>0.05$  for FI, 2hGlu and FG) (**Extended Data Figure 6, Supplementary note**). This contrasts our previous finding where we found no significant association between a risk score of non-glycemic variants and T2D<sup>7</sup>. Our current results could be partly driven by T2D cases being diagnosed based on HbA1c levels that may be influenced by the non-glycemic signals, or by glycemic effects not captured by FI, 2hGlu or FG measures.

### Biological signatures of glycemic trait associated loci

To better understand distinct and shared biological signatures underlying variant-trait associations, we conducted genomic feature enrichment, eQTL co-localization, and tissue and gene-set enrichment analyses across all four traits.

#### Epigenomic landscape of trait-associated variants

We explored the genomic context underlying glycemic trait loci by computing overlap enrichment for annotations such as coding, conserved regions, and super enhancers merged across multiple cell types<sup>53-55</sup> using the GREGOR tool<sup>56</sup>. We observed that FG, FI and HbA1c signals (**Supplementary Table 7**) were significantly ( $P<8.4 \times 10^{-4}$ , Bonferroni threshold for 59 annotations) enriched in evolutionarily conserved regions (**Fig 5A, Extended Data Figure 7, Supplementary Table 15**).



We then considered epigenomic landscapes defined in individual cell/tissue types. Previously, stretch enhancers (StrE, enhancer chromatin states  $\geq 3\text{kb}$  in length) in pancreatic islets were shown to be highly cell-specific and strongly enriched with T2D risk signals<sup>57</sup>. Considering StrEs across 31 cell-types<sup>39</sup>, FG and 2hGlu signals showed the highest enrichment in islets (FG: fold-enrichment=4.70,  $P=2.7 \times 10^{-24}$ ; 2hGlu: fold-enrichment=5.51,  $P=3.6 \times 10^{-4}$  **Figure 5A, Supplementary Table 16**), highlighting the importance of islets for these traits. FI signals were enriched in skeletal muscle (fold-enrichment=3.17,  $P=7.8 \times 10^{-6}$ ) and adipose StrEs (fold-enrichment=3.27,  $P=1.8 \times 10^{-7}$ ) consistent with these tissues as targets of insulin action (**Figure 5A**). StrEs in individual cell types showed higher enrichment than super enhancers merged across cell types, highlighting the importance of cell-specific analyses (**Figure 5A**). HbA1c signals were enriched in StrEs of multiple cell types and tissues, but have the strongest enrichment in K562 leukemia derived cells (fold-enrichment=3.24,  $P=1.2 \times 10^{-7}$ , **Figure 5A**). Among the “hard” glycemic and red blood cell (mature + reticulocyte) HbA1c signals, glycemic signals were enriched in islet StrEs (fold-enrichment=3.96,  $P=3.7 \times 10^{-16}$ ) while red blood cell signals were enriched in K562 StrEs (fold-enrichment=7.5,  $P=2.08 \times 10^{-14}$ , **Figure 5B, Supplementary Table 17**). These analyses suggest that these glycemic trait-associated variants influence the function of tissue-specific enhancers.

Independent analyses with fGWAS<sup>58</sup> and GARFIELD<sup>59</sup> yielded consistent results (**Extended Data Figures 8 and 9, Supplementary Tables 16 and 18**). Notably, FI signals at a lenient threshold of  $P < 10^{-5}$  were enriched in liver StrEs using GARFIELD (odds ratio=1.92,  $P=1.7 \times 10^{-4}$ ) (**Extended Data Figure 9A**). This suggests that liver regulatory annotations are relevant for FI GWAS signals, but that we lack power to detect significant enrichment using the genome-wide significant loci and the current set of reference annotations.

We next explored the 27 loci driving the FI enrichment in adipose and skeletal muscle, 11 of which overlapped StrEs in both tissues (**Figure 5C**). At the *COL4A2* locus, variants within an intronic region overlap StrEs in adipose tissue, skeletal muscle, and a human skeletal muscle myoblast (HSMM) cell line that are not shared across other cell/tissue types. Among these, rs9555695 (in the 99% CS) also overlaps accessible chromatin regions in adipose (**Figure 5D**). At a narrow signal with no proxy variants ( $\text{LD } r^2 > 0.7$  in Europeans), the lead trans-ancestry variant rs62271373 (PPA = 0.94) located in an intergenic region ~25kb from the *LINC01214* gene overlaps StrEs specific to adipose and HSMM and an active enhancer chromatin state in skeletal muscle (**Figure 5E**). Collectively, the tissue-specific epigenomic signatures at GWAS signals provide an opportunity to nominate tissues where these variants are likely to be active. This map may help future efforts to deconvolute GWAS signals into tissue-specific disease pathology.

### Co-localization of GWAS and eQTLs

Among the 99 novel glycemic trait loci, we identified co-localized eQTLs at 34 loci in blood, pancreatic islets, subcutaneous or visceral adipose, skeletal muscle, or liver, providing suggestive evidence of causal genes (**Supplementary Table 19**). The co-localized eQTLs include several genes previously reported at glycemic trait loci: *ADCY5*, *CAMK1D*, *IRS1*, *JAZF1*, and *KLF14*<sup>60-62</sup>. For some additional loci, the co-localized genes have prior evidence for a role in glycemic regulation. For example, the lead trans-ancestry variant and likely causal variant, rs1799815 (PPA=0.993), associated with FI is the strongest variant associated with expression of *INSR*, encoding the insulin receptor, in subcutaneous adipose from METSIM ( $P=2 \times 10^{-9}$ ) and GTEx ( $P=5 \times 10^{-6}$ ). The A allele at rs1799815 is associated with higher FI and lower expression of *INSR*, consistent with the relationship between insulin resistance and reduced *INSR* function<sup>63</sup>. In a second example, rs841572, the trans-ancestry lead variant associated with FG, has the highest PPA (PPA=0.535) among the 20 variants in the 99% CS and is in strong LD ( $r^2=0.87$ ) with the lead eQTL variant (rs841576, also in the 99% CS) associated with *SLC2A1* expression in blood (eQTLGen  $P=1 \times 10^{-8}$ ). *SLC2A1*, also known as *GLUT1*,

encodes the major glucose transporter in brain, placenta, and erythrocytes, and is responsible for glucose entry into the brain<sup>64</sup>. rs841572-A is associated with lower FG and lower *SLC2A1* expression. While rare missense variants in *SLC2A1* are an established cause of seizures and epilepsy<sup>65</sup>, our data suggest that *SLC2A1* variants also affect plasma glucose levels within a population. These co-localized signals provide possible regulatory mechanisms for variant effects on genes to influence glycemic traits.

The co-localized eQTLs also provide new insights into the mechanisms at glycemic trait loci. For example, rs9884482 (in the 99% CS) is associated with FI and *TET2* expression in subcutaneous adipose ( $P=2\times 10^{-20}$ ); rs9884482 is in high LD ( $r^2=0.96$  in Europeans) with the lead *TET2* eQTL variant (rs974801). *TET2* encodes a DNA-demethylase that can affect transcriptional repression<sup>66</sup>. Adipose *Tet2* expression is reduced in diet-induced insulin resistance in mice<sup>67</sup>, and knockdown of *Tet2* blocked adipogenesis<sup>67,68</sup>. Consistently, in human adipose tissue, rs9884482-C was associated with lower *TET2* expression and higher FI. In a second example, rs617948 is associated with HbA1c (in the 99% CS) and is the lead variant associated with *C2CD2L* expression in blood (eQTLGen  $P=3\times 10^{-96}$ ). *C2CD2L*, also known as *TMEM24*, regulates pulsatile insulin secretion and facilitates release of insulin pool reserves<sup>69,70</sup>. rs617948-G was associated with higher HbA1c and lower *C2CD2L*, providing evidence for a role of this insulin secretion protein in glucose homeostasis. Our HbA1c “soft” clustering assigned this signal to both the “unknown” (0.51 probability) and “reticulocyte” (0.42 probability) clusters. rs617948 is strongly associated with HbA1c ( $P<6.8\times 10^{-8}$ ), but not with FG, FI or 2hGlu ( $P>0.05$ , **Supplementary Table 20, Supplementary Note**). This suggests an effect of this variant on reticulocyte biology, and on insulin secretion, potentially influencing HbA1c levels through different tissues, and providing a plausible explanation for the classification as “unknown”.

### Tissue Expression

Consistent with effector transcript expression analysis using GTEx data<sup>30</sup>, we found significant differences in tissue expression across the glycemic trait signals. FG signals were enriched for genes expressed in the pancreas ( $FDR<0.05$ ), while there were an insufficient number of significant associations in 2hGlu to identify enrichment for any tissue or cell type at  $FDR<0.2$  threshold. FI signals were enriched for connective tissue and cells (which includes adipose tissue), endocrine glands, blood cells, and muscles ( $FDR<0.2$ ) and HbA1c signals were significantly enriched for genes expressed in the pancreas, hemic, and immune system ( $FDR<0.05$ ) (**Figure 6, Supplementary Table 21**). Consistent with previous analysis<sup>30</sup>, FI-enrichment for connective tissue was driven by adipose tissue (subcutaneous and visceral), while the newly described enrichment with endocrine glands was driven by the adrenal glands and cortex (**Supplementary Table 21**). Beyond enrichment for genes expressed in glycemic-related tissues, HbA1c signals were enriched with genes expressed in blood, consistent with the role of RBC in this trait and our previous results<sup>30</sup>.

The association between FI signals and genes expressed in adrenal glands is notable, suggesting a possible direct role for these genes in insulin resistance. These genes might influence cortisol levels, which could contribute to insulin resistance and FI levels through impaired insulin receptor signaling in peripheral tissues, as well as influencing body fat distribution, stimulate lipolysis, and other indirect mechanisms<sup>71,72</sup>.

### Gene-set Analyses

Next, we performed gene-set analysis using DEPICT (**Methods**). In keeping with previous results<sup>30</sup>, we found distinct gene-sets enriched ( $FDR<0.05$ ) for each glycemic trait except 2hGlu, which had insufficient associations to have power in this analysis. FG-associated variants highlighted gene-sets involved in metabolism and gene-sets involved in general cellular function such as “cytoplasmic vesicle membrane” and “circadian clock” (**Figure 7A**). In contrast, in addition to metabolism-related

gene-sets, FI-associated variants highlighted pathways related to growth, cancer and reproduction (**Figure 7B**). This is consistent with the role of insulin as a mitogenic hormone, and with epidemiological links between insulin and certain types of cancer<sup>73</sup> and reproductive disorders such as polycystic ovary syndrome<sup>74</sup>. HbA1c-associated variants highlighted many gene-sets (**Figure 7C**), including those linked to metabolism and hematopoiesis, again recapitulating our postulated effects of variants on glucose and RBC biology. Additional pathways from HbA1c-associated variants also highlighted previous “CREBP PPI” and lipid biology related to T2D<sup>75</sup> and HbA1c<sup>76</sup>, respectively, and potential new biology through which variants may influence HbA1c.

## Discussion

Here we describe a large glycemic trait meta-analysis of GWAS for which 30% of the population was composed of East Asian, Hispanic, African-American, South Asian and sub-Saharan African participants. This effort identified 242 loci (235 trans-ancestry and seven single-ancestry), which jointly explain between 0.7% (2hGlu in European ancestry individuals) and 6% (HbA1c in African American ancestry individuals) of the variance in glycemic traits in any given ancestry. While 114/242 loci are associated with T2D ( $P < 10^{-4}$ ; 83 loci with  $P < 5 \times 10^{-8}$ , **Supplementary Table 4**), absence of strong evidence of association at the remaining loci ( $P \geq 10^{-4}$ ) suggests that for alleles more frequent than 5% we can exclude T2D ORs  $\geq 1.07$  with 80% power ( $\alpha = 5 \times 10^{-8}$ ; and ORs  $\geq 1.05$  for  $\alpha = 10^{-4}$ ) given a current study of 228,499 T2D cases and 1,178,783 controls<sup>27</sup>. We identified 486 signals associated with glycemic traits, of which eight have MAF  $< 1\%$ , and 45 have  $1\% \leq \text{MAF} < 5\%$  in all ancestries, highlighting that 89% of signals identified are common in at least one ancestry studied.

A key aim of our study was to evaluate the added advantage of including population diversity in genetic discovery and fine-mapping efforts. Beyond the larger sample size included in the trans-ancestry meta-analysis, we were able to estimate the contribution of non-European ancestry data in locus discovery and fine-mapping resolution. We found that 24 of the 99 newly discovered loci owe their discovery to the inclusion of East Asian, Hispanic, African-American, South Asian and sub-Saharan African participant data, due to differences in EAF and effect sizes across ancestries.

Comparison of 295 trans-ancestry lead variants (315 locus-trait associations) across ancestries demonstrated that between 81.5% (HbA1c) and 85.7% (FG) of the trans-ancestry lead variants had no evidence of trans-ancestry heterogeneity in allelic effects ( $P > 0.05$ ).

Given sample size and power limitations, genome-wide significant trait-associated variants in a single-ancestry explain only a modest proportion of trait variance in that ancestry (**Figure 2**). We demonstrate that trans-ancestry lead variants explain more trait variance than the ancestry-specific variants (**Figure 2**). This shows that even though some trans-ancestry lead variants are not genome-wide significant in all ancestries, they contribute to the genetic architecture of the trait in most ancestries.

We evaluated for the first time the transferability of European ancestry-derived glycemic trait PGS into other ancestries. Consistent with other traits<sup>36,77,78</sup>, we confirm that European ancestry-derived PGS perform much worse when the test dataset is from a different ancestry. Each trait-specific PGS improves trait variance explained by between 3.5-fold (HbA1c) and 6-fold (FG) in the European dataset (**Figure 3, Supplementary Table 12**) compared to a score built only from trans-ancestry lead variants and European index variants (**Figure 2, Supplementary tables 9-12**).

Despite development of approaches to derive polygenic risk scores<sup>79</sup>, we note the difficulty in using summary level data to build a PGS in one ancestry and then apply it in test datasets of different ancestry. While PRS-CSauto<sup>34</sup> is able to use summary level data, revision of the effect size estimates

to account for LD required reference panels that matched the ancestry of the test dataset. However, the current software lacks appropriate reference panels for many ancestries, precluding its broad application. Future developments of trans-ancestry PGS are required for improved cross-ancestry performance.

We show that fine-mapping resolution is improved in trans-ancestry, compared to single-ancestry fine-mapping efforts. In ~50% of our loci, we showed that the improvement was due to differences in EAF, effect size, or LD structure between ancestries, and not just due to the overall increased sample size available for trans-ancestry fine-mapping. By performing trans-ancestry fine-mapping, and co-localizing GWAS signals with eQTL signals and coding variants, we identified new candidate causal genes. Altogether, these results motivate continued expansion of genetic and genomic efforts in diverse populations to improve understanding of these traits in groups disproportionately affected by T2D.

Given data on four different glycemic traits and their utility to diagnose and monitor T2D and metabolic health, we also sought to characterize biological features underlying these traits. We show that despite significant sharing of loci across the four traits, each trait is also characterized by unique features based on StrE, gene expression and gene-set signatures. Combining genetic data from these traits with T2D data will further elucidate pathways driving normal physiology and pathophysiology, and help further develop useful predictive scores for disease classification and management<sup>4,5</sup>.

## References

- 1 in *Use of Glycated Haemoglobin (HbA1c) in the Diagnosis of Diabetes Mellitus: Abbreviated Report of a WHO Consultation* (World Health Organization Copyright © World Health Organization 2011., 2011).
- 2 Goodarzi, M. O. *et al.* Fasting insulin reflects heterogeneous physiological processes: role of insulin clearance. *American journal of physiology. Endocrinology and metabolism* **301**, E402-408, doi:10.1152/ajpendo.00013.2011 (2011).
- 3 Dimas, A. S. *et al.* Impact of type 2 diabetes susceptibility variants on quantitative glycemic traits reveals mechanistic heterogeneity. *Diabetes* **63**, 2158-2171, doi:10.2337/db13-0949 (2014).
- 4 Udler, M. S. *et al.* Type 2 diabetes genetic loci informed by multi-trait associations point to disease mechanisms and subtypes: A soft clustering analysis. *PLoS medicine* **15**, e1002654, doi:10.1371/journal.pmed.1002654 (2018).
- 5 Udler, M. S., McCarthy, M. I., Florez, J. C. & Mahajan, A. Genetic Risk Scores for Diabetes Diagnosis and Precision Medicine. *Endocrine reviews* **40**, 1500-1520, doi:10.1210/er.2019-00088 (2019).
- 6 Sarwar, N. *et al.* Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: a collaborative meta-analysis of 102 prospective studies. *Lancet* **375**, 2215-2222, doi:10.1016/s0140-6736(10)60484-9 (2010).
- 7 Wheeler, E. *et al.* Impact of common genetic determinants of Hemoglobin A1c on type 2 diabetes risk and diagnosis in ancestrally diverse populations: A transethnic genome-wide meta-analysis. *PLoS medicine* **14**, e1002383, doi:10.1371/journal.pmed.1002383 (2017).

926 8 Dupuis, J. *et al.* New genetic loci implicated in fasting glucose homeostasis and their  
927 impact on type 2 diabetes risk. *Nature genetics* **42**, 105-116, doi:10.1038/ng.520  
928 (2010).

929 9 Manning, A. K. *et al.* A genome-wide approach accounting for body mass index  
930 identifies genetic variants influencing fasting glycemic traits and insulin resistance.  
931 *Nature genetics* **44**, 659-669, doi:10.1038/ng.2274 (2012).

932 10 Walford, G. A. *et al.* Genome-Wide Association Study of the Modified Stumvoll  
933 Insulin Sensitivity Index Identifies BCL2 and FAM19A2 as Novel Insulin Sensitivity  
934 Loci. *Diabetes* **65**, 3200-3211, doi:10.2337/db16-0199 (2016).

935 11 Horikoshi, M. *et al.* Discovery and Fine-Mapping of Glycaemic and Obesity-Related  
936 Trait Loci Using High-Density Imputation. *PLoS genetics* **11**, e1005230,  
937 doi:10.1371/journal.pgen.1005230 (2015).

938 12 Mahajan, A. *et al.* Identification and functional characterization of G6PC2 coding  
939 variants influencing glycemic traits define an effector transcript at the G6PC2-  
940 ABCB11 locus. *PLoS genetics* **11**, e1004876, doi:10.1371/journal.pgen.1004876  
941 (2015).

942 13 Hwang, J. Y. *et al.* Genome-wide association meta-analysis identifies novel variants  
943 associated with fasting plasma glucose in East Asians. *Diabetes* **64**, 291-298,  
944 doi:10.2337/db14-0563 (2015).

945 14 Chen, P. *et al.* Multiple nonglycemic genomic loci are newly associated with blood  
946 level of glycated hemoglobin in East Asians. *Diabetes* **63**, 2551-2562,  
947 doi:10.2337/db13-1815 (2014).

948 15 Scott, R. A. *et al.* Large-scale association analyses identify new loci influencing  
949 glycemic traits and provide insight into the underlying biological pathways. *Nature*  
950 *genetics* **44**, 991-1005, doi:10.1038/ng.2385 (2012).

951 16 Spanakis, E. K. & Golden, S. H. Race/ethnic difference in diabetes and diabetic  
952 complications. *Current diabetes reports* **13**, 814-823, doi:10.1007/s11892-013-0421-  
953 9 (2013).

954 17 Tillin, T. *et al.* Insulin resistance and truncal obesity as important determinants of the  
955 greater incidence of diabetes in Indian Asians and African Caribbeans compared with  
956 Europeans: the Southall And Brent REvisited (SABRE) cohort. *Diabetes care* **36**, 383-  
957 393, doi:10.2337/dc12-0544 (2013).

958 18 Whincup, P. H. *et al.* Early emergence of ethnic differences in type 2 diabetes  
959 precursors in the UK: the Child Heart and Health Study in England (CHASE Study).  
960 *PLoS medicine* **7**, e1000263, doi:10.1371/journal.pmed.1000263 (2010).

961 19 Auton, A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68-74,  
962 doi:10.1038/nature15393 (2015).

963 20 Willer, C. J., Li, Y. & Abecasis, G. R. METAL: fast and efficient meta-analysis of  
964 genomewide association scans. *Bioinformatics (Oxford, England)* **26**, 2190-2191,  
965 doi:10.1093/bioinformatics/btq340 (2010).

966 21 Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide  
967 complex trait analysis. *American journal of human genetics* **88**, 76-82,  
968 doi:10.1016/j.ajhg.2010.11.011 (2011).

969 22 Yang, J. *et al.* Conditional and joint multiple-SNP analysis of GWAS summary statistics  
970 identifies additional variants influencing complex traits. *Nature genetics* **44**, 369-375,  
971 s361-363, doi:10.1038/ng.2213 (2012).

972 23 Genome-wide association study of 14,000 cases of seven common diseases and  
973 3,000 shared controls. *Nature* **447**, 661-678, doi:10.1038/nature05911 (2007).

974 24 Mahajan, A. *et al.* Trans-ancestry genetic study of type 2 diabetes highlights the  
975 power of diverse populations for discovery and translation. *medRxiv*,  
976 2020.2009.2022.20198937, doi:10.1101/2020.09.22.20198937 (2020).

977 25 Mahajan, A. *et al.* Fine-mapping type 2 diabetes loci to single-variant resolution  
978 using high-density imputation and islet-specific epigenome maps. *Nature genetics*  
979 **50**, 1505-1513, doi:10.1038/s41588-018-0241-6 (2018).

980 26 Spracklen, C. N. *et al.* Identification of type 2 diabetes loci in 433,540 East Asian  
981 individuals. *Nature* **582**, 240-245, doi:10.1038/s41586-020-2263-3 (2020).

982 27 Vujkovic, M. *et al.* Discovery of 318 new risk loci for type 2 diabetes and related  
983 vascular outcomes among 1.4 million participants in a multi-ancestry meta-analysis.  
984 *Nature genetics* **52**, 680-691, doi:10.1038/s41588-020-0637-y (2020).

985 28 Luo, Y. *et al.* Transcription factor Ets1 regulates expression of thioredoxin-interacting  
986 protein and inhibits insulin secretion in pancreatic beta-cells. *PLoS one* **9**, e99049,  
987 doi:10.1371/journal.pone.0099049 (2014).

988 29 Braccini, L. *et al.* PI3K-C2gamma is a Rab5 effector selectively controlling endosomal  
989 Akt2 activation downstream of insulin signalling. *Nature communications* **6**, 7400,  
990 doi:10.1038/ncomms8400 (2015).

991 30 Ng, N. H. J. *et al.* Tissue-Specific Alteration of Metabolic Pathways Influences  
992 Glycemic Regulation. *bioRxiv*, 790618, doi:10.1101/790618 (2019).

993 31 Aschard, H., Vilhjalmsen, B. J., Joshi, A. D., Price, A. L. & Kraft, P. Adjusting for  
994 heritable covariates can bias effect estimates in genome-wide association studies.  
995 *American journal of human genetics* **96**, 329-339, doi:10.1016/j.ajhg.2014.12.021  
996 (2015).

997 32 Lee, J. J. *et al.* Gene discovery and polygenic prediction from a genome-wide  
998 association study of educational attainment in 1.1 million individuals. *Nature*  
999 *genetics* **50**, 1112-1121, doi:10.1038/s41588-018-0147-3 (2018).

1000 33 Nolte, I. M. *et al.* Missing heritability: is the gap closing? An analysis of 32 complex  
1001 traits in the Lifelines Cohort Study. *European journal of human genetics : EJHG* **25**,  
1002 877-885, doi:10.1038/ejhg.2017.50 (2017).

1003 34 Ge, T., Chen, C. Y., Ni, Y., Feng, Y. A. & Smoller, J. W. Polygenic prediction via  
1004 Bayesian regression and continuous shrinkage priors. *Nature communications* **10**,  
1005 1776, doi:10.1038/s41467-019-09718-5 (2019).

1006 35 Dastani, Z. *et al.* Novel loci for adiponectin levels and their influence on type 2  
1007 diabetes and metabolic traits: a multi-ethnic meta-analysis of 45,891 individuals.  
1008 *PLoS genetics* **8**, e1002607, doi:10.1371/journal.pgen.1002607 (2012).

1009 36 Martin, A. R. *et al.* Clinical use of current polygenic risk scores may exacerbate health  
1010 disparities. *Nature genetics* **51**, 584-591, doi:10.1038/s41588-019-0379-x (2019).

1011 37 Gaulton, K. J. *et al.* Genetic fine mapping and genomic annotation defines causal  
1012 mechanisms at type 2 diabetes susceptibility loci. *Nature genetics* **47**, 1415-1425,  
1013 doi:10.1038/ng.3437 (2015).

1014 38 Spracklen, C. N. *et al.* Identification and functional analysis of glycemic trait loci in  
1015 the China Health and Nutrition Survey. *PLoS genetics* **14**, e1007275,  
1016 doi:10.1371/journal.pgen.1007275 (2018).

1017 39 Varshney, A. *et al.* Genetic regulatory signatures underlying islet gene expression  
1018 and type 2 diabetes. *Proceedings of the National Academy of Sciences of the United*  
1019 *States of America* **114**, 2301-2306, doi:10.1073/pnas.1621192114 (2017).

1020 40 Kichaev, G. *et al.* Leveraging Polygenic Functional Enrichment to Improve GWAS  
1021 Power. *American journal of human genetics* **104**, 65-75,  
1022 doi:10.1016/j.ajhg.2018.11.008 (2019).

1023 41 Shriner, D. & Rotimi, C. N. Whole-Genome-Sequence-Based Haplotypes Reveal Single  
1024 Origin of the Sickle Allele during the Holocene Wet Phase. *American journal of*  
1025 *human genetics* **102**, 547-556, doi:10.1016/j.ajhg.2018.02.003 (2018).

1026 42 Kramer, H. J. *et al.* African Ancestry-Specific Alleles and Kidney Disease Risk in  
1027 Hispanics/Latinos. *Journal of the American Society of Nephrology : JASN* **28**, 915-922,  
1028 doi:10.1681/asn.2016030357 (2017).

1029 43 Ravenhall, M. *et al.* Novel genetic polymorphisms associated with severe malaria and  
1030 under selective pressure in North-eastern Tanzania. *PLoS genetics* **14**, e1007172,  
1031 doi:10.1371/journal.pgen.1007172 (2018).

1032 44 Hodonsky, C. J. *et al.* Genome-wide association study of red blood cell traits in  
1033 Hispanics/Latinos: The Hispanic Community Health Study/Study of Latinos. *PLoS*  
1034 *genetics* **13**, e1006760, doi:10.1371/journal.pgen.1006760 (2017).

1035 45 Gurdasani, D. *et al.* Uganda Genome Resource Enables Insights into Population  
1036 History and Genomic Discovery in Africa. *Cell* **179**, 984-1002.e1036,  
1037 doi:10.1016/j.cell.2019.10.004 (2019).

1038 46 Rees, M. G. *et al.* Cellular characterisation of the GCKR P446L variant associated with  
1039 type 2 diabetes risk. *Diabetologia* **55**, 114-122, doi:10.1007/s00125-011-2348-5  
1040 (2012).

1041 47 Bonomo, J. A. *et al.* The ras responsive transcription factor RREB1 is a novel  
1042 candidate gene for type 2 diabetes associated end-stage kidney disease. *Human*  
1043 *molecular genetics* **23**, 6441-6447, doi:10.1093/hmg/ddu362 (2014).

1044 48 Wessel, J. *et al.* Low-frequency and rare exome chip variants associate with fasting  
1045 glucose and type 2 diabetes susceptibility. *Nature communications* **6**, 5897,  
1046 doi:10.1038/ncomms6897 (2015).

1047 49 Scott, R. A. *et al.* A genomic approach to therapeutic target validation identifies a  
1048 glucose-lowering GLP1R variant protective for coronary heart disease. *Science*  
1049 *translational medicine* **8**, 341ra376, doi:10.1126/scitranslmed.aad3744 (2016).

1050 50 Nai, A. *et al.* TMPRSS6 rs855791 modulates hepcidin transcription in vitro and serum  
1051 hepcidin levels in normal individuals. *Blood* **118**, 4459-4462, doi:10.1182/blood-  
1052 2011-06-364034 (2011).

1053 51 Soranzo, N. *et al.* Common variants at 10 genomic loci influence hemoglobin A(1)(C)  
1054 levels via glycemic and nonglycemic pathways. *Diabetes* **59**, 3229-3239,  
1055 doi:10.2337/db10-0502 (2010).

1056 52 Sarnowski, C. *et al.* Impact of Rare and Common Genetic Variants on Diabetes  
1057 Diagnosis by Hemoglobin A1c in Multi-Ancestry Cohorts: The Trans-Omics for  
1058 Precision Medicine Program. *American journal of human genetics* **105**, 706-718,  
1059 doi:10.1016/j.ajhg.2019.08.010 (2019).

1060 53 Kundaje, A. *et al.* Integrative analysis of 111 reference human epigenomes. *Nature*  
1061 **518**, 317-330, doi:10.1038/nature14248 (2015).

1062 54 Nagel, M. *et al.* Meta-analysis of genome-wide association studies for neuroticism in  
1063 449,484 individuals identifies novel genetic loci and pathways. *Nature genetics* **50**,  
1064 920-927, doi:10.1038/s41588-018-0151-7 (2018).

1065 55 Savage, J. E. *et al.* Genome-wide association meta-analysis in 269,867 individuals  
1066 identifies new genetic and functional links to intelligence. *Nature genetics* **50**, 912-  
1067 919, doi:10.1038/s41588-018-0152-6 (2018).

1068 56 Schmidt, E. M. *et al.* GREGOR: evaluating global enrichment of trait-associated  
1069 variants in epigenomic features using a systematic, data-driven approach.  
1070 *Bioinformatics (Oxford, England)* **31**, 2601-2606, doi:10.1093/bioinformatics/btv201  
1071 (2015).

1072 57 Parker, S. C. *et al.* Chromatin stretch enhancer states drive cell-specific gene  
1073 regulation and harbor human disease risk variants. *Proceedings of the National*  
1074 *Academy of Sciences of the United States of America* **110**, 17921-17926,  
1075 doi:10.1073/pnas.1317023110 (2013).

1076 58 Pickrell, J. K. Joint analysis of functional genomic data and genome-wide association  
1077 studies of 18 human traits. *American journal of human genetics* **94**, 559-573,  
1078 doi:10.1016/j.ajhg.2014.03.004 (2014).

1079 59 Iotchkova, V. *et al.* GARFIELD classifies disease-relevant genomic features through  
1080 integration of functional annotations with association signals. *Nature genetics* **51**,  
1081 343-353, doi:10.1038/s41588-018-0322-6 (2019).

1082 60 van de Bunt, M. *et al.* Transcript Expression Data from Human Islets Links Regulatory  
1083 Signals from Genome-Wide Association Studies for Type 2 Diabetes and Glycemic  
1084 Traits to Their Downstream Effectors. *PLoS genetics* **11**, e1005694,  
1085 doi:10.1371/journal.pgen.1005694 (2015).

1086 61 Civelek, M. *et al.* Genetic Regulation of Adipose Gene Expression and Cardio-  
1087 Metabolic Traits. *American journal of human genetics* **100**, 428-443,  
1088 doi:10.1016/j.ajhg.2017.01.027 (2017).

1089 62 Scott, L. J. *et al.* The genetic regulatory signature of type 2 diabetes in human  
1090 skeletal muscle. *Nature communications* **7**, 11764, doi:10.1038/ncomms11764  
1091 (2016).

1092 63 Ben Harouch, S., Klar, A. & Falik Zaccai, T. C. in *GeneReviews((R))* (eds M. P. Adam *et*  
1093 *al.*) (University of Washington, Seattle  
1094 University of Washington, Seattle. GeneReviews is a registered trademark of the University  
1095 of Washington, Seattle. All rights reserved., 1993).

1096 64 Agus, D. B. *et al.* Vitamin C crosses the blood-brain barrier in the oxidized form  
1097 through the glucose transporters. *The Journal of clinical investigation* **100**, 2842-  
1098 2848, doi:10.1172/jci119832 (1997).

1099 65 Wolking, S. *et al.* Focal epilepsy in glucose transporter type 1 (Glut1) defects: case  
1100 reports and a review of literature. *Journal of neurology* **261**, 1881-1886,  
1101 doi:10.1007/s00415-014-7433-5 (2014).

1102 66 Guallar, D. *et al.* RNA-dependent chromatin targeting of TET2 for endogenous  
1103 retrovirus control in pluripotent stem cells. *Nature genetics* **50**, 443-451,  
1104 doi:10.1038/s41588-018-0060-9 (2018).

1105 67 Bian, F. *et al.* TET2 facilitates PPARGgamma agonist-mediated gene regulation and  
1106 insulin sensitization in adipocytes. *Metabolism: clinical and experimental* **89**, 39-47,  
1107 doi:10.1016/j.metabol.2018.08.006 (2018).



1108 68 Yoo, Y. *et al.* TET-mediated hydroxymethylcytosine at the Pparg locus is  
1109 required for initiation of adipogenic differentiation. *International journal of obesity*  
1110 (2005) **41**, 652-659, doi:10.1038/ijo.2017.8 (2017).

1111 69 Lees, J. A. *et al.* Lipid transport by TMEM24 at ER-plasma membrane contacts  
1112 regulates pulsatile insulin secretion. *Science (New York, N.Y.)* **355**,  
1113 doi:10.1126/science.aah6171 (2017).

1114 70 Pottekat, A. *et al.* Insulin biosynthetic interaction network component, TMEM24,  
1115 facilitates insulin reserve pool release. *Cell reports* **4**, 921-930,  
1116 doi:10.1016/j.celrep.2013.07.050 (2013).

1117 71 Androulakis, I *et al.* Patients with apparently nonfunctioning adrenal incidentalomas  
1118 may be at increased cardiovascular risk due to excessive cortisol secretion. *The*  
1119 *Journal of clinical endocrinology and metabolism* **99**, 2754-2762,  
1120 doi:10.1210/jc.2013-4064 (2014).

1121 72 Altieri, B. *et al.* Adrenocortical tumors and insulin resistance: What is the first step?  
1122 *International journal of cancer* **138**, 2785-2794, doi:10.1002/ijc.29950 (2016).

1123 73 Johansson, M. *et al.* The influence of obesity-related factors in the etiology of renal  
1124 cell carcinoma-A mendelian randomization study. *PLoS medicine* **16**, e1002724,  
1125 doi:10.1371/journal.pmed.1002724 (2019).

1126 74 Diamanti-Kandarakis, E. & Dunaif, A. Insulin resistance and the polycystic ovary  
1127 syndrome revisited: an update on mechanisms and implications. *Endocrine reviews*  
1128 **33**, 981-1030, doi:10.1210/er.2011-1034 (2012).

1129 75 Morris, A. P. *et al.* Large-scale association analysis provides insights into the genetic  
1130 architecture and pathophysiology of type 2 diabetes. *Nature genetics* **44**, 981-990,  
1131 doi:10.1038/ng.2383 (2012).

1132 76 Leong, A. *et al.* Mendelian Randomization Analysis of Hemoglobin A(1c) as a Risk  
1133 Factor for Coronary Artery Disease. *Diabetes care* **42**, 1202-1208, doi:10.2337/dc18-  
1134 1712 (2019).

1135 77 Duncan, L. *et al.* Analysis of polygenic risk score usage and performance in diverse  
1136 human populations. *Nature communications* **10**, 3328, doi:10.1038/s41467-019-  
1137 11112-0 (2019).

1138 78 Mostafavi, H. *et al.* Variable prediction accuracy of polygenic scores within an  
1139 ancestry group. *eLife* **9**, doi:10.7554/eLife.48376 (2020).

1140 79 Choi, S. W., Mak, T. S. & O'Reilly, P. F. Tutorial: a guide to performing polygenic risk  
1141 score analyses. *Nature protocols* **15**, 2759-2772, doi:10.1038/s41596-020-0353-1  
1142 (2020).

1143 80 D'Orazio, P. *et al.* Approved IFCC recommendation on reporting results for blood  
1144 glucose (abbreviated). *Clinical chemistry* **51**, 1573-1576,  
1145 doi:10.1373/clinchem.2005.051979 (2005).

1146 81 Voight, B. F. *et al.* The metabochip, a custom genotyping array for genetic studies of  
1147 metabolic, cardiovascular, and anthropometric traits. *PLoS genetics* **8**, e1002793,  
1148 doi:10.1371/journal.pgen.1002793 (2012).

1149 82 Abecasis, G. R. *et al.* An integrated map of genetic variation from 1,092 human  
1150 genomes. *Nature* **491**, 56-65, doi:10.1038/nature11632 (2012).

1151 83 Li, Y., Willer, C. J., Ding, J., Scheet, P. & Abecasis, G. R. MaCH: using sequence and  
1152 genotype data to estimate haplotypes and unobserved genotypes. *Genetic*  
1153 *epidemiology* **34**, 816-834, doi:10.1002/gepi.20533 (2010).

1154 84 Pei, Y. F., Zhang, L., Li, J. & Deng, H. W. Analyses and comparison of imputation-  
1155 based association methods. *PloS one* **5**, e10827, doi:10.1371/journal.pone.0010827  
1156 (2010).

1157 85 Winkler, T. W. *et al.* Quality control and conduct of genome-wide association meta-  
1158 analyses. *Nature protocols* **9**, 1192-1212, doi:10.1038/nprot.2014.071 (2014).

1159 86 Devlin, B. & Roeder, K. Genomic control for association studies. *Biometrics* **55**, 997-  
1160 1004 (1999).

1161 87 Morris, A. P. Transethnic meta-analysis of genomewide association studies. *Genetic*  
1162 *epidemiology* **35**, 809-822, doi:10.1002/gepi.20630 (2011).

1163 88 Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from  
1164 polygenicity in genome-wide association studies. *Nature genetics* **47**, 291-295,  
1165 doi:10.1038/ng.3211 (2015).

1166 89 Dastani, Z. *et al.* Novel loci for adiponectin levels and their influence on type 2  
1167 diabetes and metabolic traits: a multi-ethnic meta-analysis of 45,891 individuals.  
1168 *PLoS genetics* **8**, e1002607, doi:10.1371/journal.pgen.1002607 (2012).

1169 90 Benner, C. *et al.* FINEMAP: efficient variable selection using summary data from  
1170 genome-wide association studies. *Bioinformatics (Oxford, England)* **32**, 1493-1501,  
1171 doi:10.1093/bioinformatics/btw018 (2016).

1172 91 Astle, W. J. *et al.* The Allelic Landscape of Human Blood Cell Trait Variation and Links  
1173 to Common Complex Disease. *Cell* **167**, 1415-1429.e1419,  
1174 doi:10.1016/j.cell.2016.10.042 (2016).

1175 92 Canela-Xandri, O., Rawlik, K. & Tenesa, A. An atlas of genetic associations in UK  
1176 Biobank. *Nature genetics* **50**, 1593-1599, doi:10.1038/s41588-018-0248-z (2018).

1177 93 Benyamin, B. *et al.* Novel loci affecting iron homeostasis and their effects in  
1178 individuals at risk for hemochromatosis. *Nature communications* **5**, 4926,  
1179 doi:10.1038/ncomms5926 (2014).

1180 94 Binesh, N. & Rezghi, M. Fuzzy clustering in community detection based on  
1181 nonnegative matrix factorization with two novel evaluation criteria. *Applied Soft*  
1182 *Computing* **69**, 689-703 (2018).

1183 95 Scott, R. A. *et al.* An Expanded Genome-Wide Association Study of Type 2 Diabetes in  
1184 Europeans. *Diabetes* **66**, 2888-2902, doi:10.2337/db16-1253 (2017).

1185 96 Ernst, J. *et al.* Mapping and analysis of chromatin state dynamics in nine human cell  
1186 types. *Nature* **473**, 43-49, doi:10.1038/nature09906 (2011).

1187 97 Mikkelsen, T. S. *et al.* Comparative epigenomic analysis of murine and human  
1188 adipogenesis. *Cell* **143**, 156-169, doi:10.1016/j.cell.2010.09.006 (2010).

1189 98 Ernst, J. & Kellis, M. ChromHMM: automating chromatin-state discovery and  
1190 characterization. *Nature methods* **9**, 215-216, doi:10.1038/nmeth.1906 (2012).

1191 99 Battle, A., Brown, C. D., Engelhardt, B. E. & Montgomery, S. B. Genetic effects on  
1192 gene expression across human tissues. *Nature* **550**, 204-213,  
1193 doi:10.1038/nature24277 (2017).

1194 100 Zhernakova, D. V. *et al.* Identification of context-dependent expression quantitative  
1195 trait loci in whole blood. *Nature genetics* **49**, 139-145, doi:10.1038/ng.3737 (2017).

1196 101 Westra, H. J. *et al.* Systematic identification of trans eQTLs as putative drivers of  
1197 known disease associations. *Nature genetics* **45**, 1238-1243, doi:10.1038/ng.2756  
1198 (2013).

- 102 Joehanes, R. *et al.* Integrated genome-wide analysis of expression quantitative trait loci aids interpretation of genomic association studies. *Genome biology* **18**, 16, doi:10.1186/s13059-016-1142-6 (2017).
- 103 Pers, T. H. *et al.* Biological interpretation of genome-wide association studies using predicted gene functions. *Nature communications* **6**, 5890, doi:10.1038/ncomms6890 (2015).

## Acknowledgments

The authors thank all investigators, staff members, and study participants for their contribution to all participating studies. The funders had no role in study design, data collection, analysis, decision to publish, or preparation of the manuscript. The authors received no specific funding for this work. A full list of funding and individual and study acknowledgments appears in the **Supplementary Note**.

## Author contributions

Project coordination: I.B.

Writing group: J.C., C.N.S., G.M., A.V., L.J.C., S.C.J.P., K.L.M., C.L., E.W., A.P.M., I.B.

Central analysis group: J.C., C.N.S., G.M., A.V., L.J.C., J.L., S.W., Y.W., X.Z., M.H., T.S.B., R.M., J.W., A.P., R.L., K.H.K.C., J.Y., M.D.A., A.Y.C., A.C., J.H., S.H., M.A.K., T.L., W.M., H.M-M., A.N., S.C.N., K.N., C.K.R., D.R., R.R., D.R., C.S., X.S., L.S., I.D.S., C.A.W., Y.W., P.W., W.Z., J.I.R., A.L.G., M.I.M., J.D., J.B.M., R.A.S., I.P., A.L., C.T.L., S.C.J.P., K.L.M., C.L., E.W., A.P.M., I.B.

Cohort analysts: T.S.A., E.V.R.A., L.F.B., J.A.B., N.P.B., C.P.C., B.E.C., J.C., X.C., L.C., C.C., B.H.C., K.C., Y.C., H.G.d., G.E.D., A.D., Q.D., J.E., S.A.F., J.G., F.G., J.G., S.G., Y.H., F.P.H., J.H., Y.H., T.H., A.H., M.H., R.A.J., T.K., K.A.K., Y.K., M.E.K., I.K.K., S.L., L.A.L., C.D.L., M.L., M.L., S.L., J.L., M.L., J.L., V.L., M.M., C.M., M.E.M., A.N., M.N., D.N., R.N., G.P., M.P., L.R., L.J.R., S.S.R., N.R.R., R.R., K.R., S.S., R.S., K.E.S., B.S., K.S., A.V.S., L.S., T.S., R.J.S., F.T., J.T., S.T., E.v., P.J.v., N.V., M.V., H.W., C.W., N.W., H.R.W., W.W., T.W., A.W., A.R.W., T.X., M.Z., J.Z., W.Z.

Cohort genotyping and phenotyping:

N.A., Z.A., A.A., S.J.L.B., D.B., M.B., R.N.B., A.B., M.B., L.L.B., S.R.B., D.W.B., Q.C., A.C., H.C., Y.C., E.J.C.d., A.D., S.D., G.E., A.F., M.F., C.F., Y.G., A.P.G., A.G., S.H., C.A.H., C.H., A.A.H., C.H., W.A.H., S.I., M.I., M.Arfañl., W.CraigJ., M.E.J., P.K.J., R.R.K., F.R.K., T.K., C.K., W.K., I.K., T.K., J.K., K.L., K.L., D.A.L., N.R.L., R.N.L., H.L., S.L., J.L., A.L., J.L., C.L., T.M., F.M., G.M., S.M., S.M., T.N., G.N.N., J.L.N., M.N., M.J.N., J.M.N., Y.O., A.P., P.A.P., O.P., Q.Q., D.R., D.F.R., A.R., F.R., K.R., I.R., C.S., K.S., N.S., A.S., J.S., H.M.S., K.D.T., T.M.T., B.T., P.RHJ.T., E.T., M.Y.T., A.U., R.M.v., D.v., A.v., J.V.V., J.V., H.V., T.W., K.W., T.Z.

Cohort oversight and/or principal investigator:

G.R.A., L.S.A., C.AlbertoA., M.E.A., P.A., L.A., D.M.B., L.J.B., S.B., H.B., C.B., M.B., E.B., B.O.B., K.B., D.I.B., E.P.B., T.A.B., M.C., M.J.C., J.C.C., D.I.C., Y.C., C.C., F.S.C., A.C., F.C., H.d., G.D., S.E., M.K.E., E.F., L.F., J.C.F., P.W.F., T.M.F., P.F., B.G., M.O.G., P.G., H.G., N.G., S.G., L.G., V.G., X.G., A.H., T.H., C.H., S.R.H., B.L.H., W.H., E.I., P.S.J., M.J., J.B.J., J.WouterJ., P.K., R.K., S.L.R.K., N.K., S.M.K., B.K., M.K., H.A.K., J.S.K., A.K., P.K., D.K., M.K., Z.K., M.L., T.A.L., L.J.L., K.L., H.L., X.L., L.L., C.L., S.L., R.J.F.L., P.KE.M., A.M., A.M., D.O.M., T.A.M., P.B.M., I.N., J.R.O., A.J.O., K.K.O., S.P., C.N.A.P., N.D.P., O.P., C.E.P., D.J.P., P.P.P., M.A.P., B.M.P., L.Q., L.J.R., R.R., S.R., P.M.R., F.R.R., T.E.S., M.S., J.S.N.S., P.S., L.J.S., E.S., P.S., X.S., P.ElineS., K.S.S., B.H.S., H.S., T.S., T.I.A.S., T.D.S., A.S., C.J.S., M.S., L.S., Y.T., E.T., N.J.T., A.T., J.T., T.T., M.U., P.v., C.v., , P.V., T.GM.V., L.E.W., M.W., Y.X.W., N.J.W., R.M.W., H.W., W.B.W., A.R.W., G.W., J.F.W., T.W., J.W., A.H.X., L.R.Y., L.Y., M.Y., E.Z., W.Z., A.B.Z., J.I.R., A.L.G., M.I.M., J.D., J.B.M., R.A.S., I.P., A.L., C.L., S.C.J.P., K.L.M., C.L., E.W., A.P.M., I.B.

## Competing interests statement

A. Astrup is the recipient of honoraria as speaker for a wide range of Danish and international concerns and receives royalties from textbooks, and from popular diet and cookery books. A. Astrup is also co-inventor of a number of patents, including Methods of inducing weight loss, treating obesity and preventing weight gain (licensee Gelesis, USA) and Biomarkers for predicting degree of weight loss (licensee Nestec SA, CH), owned by the University of Copenhagen, in accordance with Danish law. I. Barroso and spouse own stock in GlaxoSmithKline and Incyte Corporation. B.H. Chen is now an employee of Life Epigenetics, Inc.; all work was completed prior to employment at Life Epigenetics. A.Y. Chu is now an employee of Merck & Co.; all work was completed prior to employment by Merck & Co. J.C. Florez has received consulting honoraria from Janssen. J. Gayan is now an employee of F. Hoffmann-La Roche Ltd, and owns stock of Roche and GlaxoSmithKline. A.L. Gloyn has received honoraria from Merck and Novo Nordisk. As of June 2019, ALG discloses that her spouse is an employee of Genentech and hold stock options in Roche. E. Ingelsson is now an employee of GSK; all work was completed prior to his employment by GSK. W. März has received grants and/or personal fees from the following companies/corporations: Siemens Healthineers, Aegerion Pharmaceuticals, AMGEN, Astrazeneca, Sanofi, Alexion Pharmaceuticals, BASF, Abbott Diagnostics Numares AG, Berlin-Chemie, Akzea Therapeutics, Bayer Vital GmbH, bestbion dx GmbH, Boehringer Ingelheim Pharma GmbH Co KG, Immundiagnostik GmbH, Merck Chemicals GmbH, MSD Sharp and Dohme GmbH, Novartis Pharma GmbH, Olink Proteomics, and Synlab Holding Deutschland GmbH. M.I. McCarthy has served on advisory panels for Pfizer, NovoNordisk, Zoe Global and received honoraria from Merck, Pfizer, NovoNordisk and Eli Lilly. He holds stock options in Zoe Global and has received research funding from Abbvie, Astra Zeneca, Boehringer Ingelheim, Eli Lilly, Janssen, Merck, NovoNordisk, Pfizer, Roche, Sanofi Aventis, Servier, Takeda. He is now an employee of Genentech and a holder of Roche stock. J.B. Meigs has consulted for Quest Diagnostics, Inc., who manufacturers of an HbA1c assay. M.E. Montasser has received grant funding from Regeneron Pharmaceuticals. M.E. Montasser is also an inventor on a patent that was published by the United States Patent and Trademark Office on December 6, 2018 under Publication Number US 2018-0346888, and international patent application that was published on December 13, 2018 under Publication Number WO-2018/226560; all work was completed before these COI arose, and are unrelated to this work. D. Mook-Kanamori is a part-time clinical research consultant for Metabolon. J.L. Nadler is a member of the Scientific Advisory Board for Veralox Therapeutics Inc. C.N.A. Palmer has received research support from GlaxoSmithKline and AstraZeneca unrelated to this project. B.M. Psaty serves on the Steering Committee of the Yale Open Data Access Project funded by Johnson & Johnson. N. Sattar has consulted for Astrazeneca, Boehringer Ingelheim, Eli Lilly, Novo Nordisk, Napp and Sanofi and received grant support from Boehringer Ingelheim. R.A. Scott is an employee and shareholder of GlaxoSmithKline. T. Spector is the founder of Zoe Global Ltd. J. Tuomilehto receives research support from Bayer, is a consultant for Eli Lilly, and holds stock in Orion Pharma and Aktivolabs Ltd.

## Figure Legends

**Figure 1 - Summary of all 242 loci identified in this study.** 235 trans-ancestry loci are shown in orange (novel) or black (established) along with seven single-ancestry loci (blue) represented by nearest gene. Each locus is mapped to corresponding chromosome (outer segment). Each set of rows shows the results from the trans-ancestry analysis (orange) and each of the ancestries: European (purple), African American (tan), East Asian (grey), South Asian (green), Hispanic (yellow), sub-Saharan African (Ugandan-pink). Loci with a corresponding type 2 diabetes signal are represented by red circles in the middle of the plot.

**Figure 2 – Trait variance explained by associated loci.** The boxplots show the maximum, first quartile, median, third quartile and minimum of trait variance explained when using a genetic score with single-ancestry lead and index variants (EUR, AA, EAS, HISP and SAS) or a combination of individual trait trans-ancestry lead variants and single-ancestry lead and index variants (TA+EUR,

TA+AA, TA+EAS, TA+HISP and TA+SAS). Variance explained for each trait (FG, FI and HbA1c) in each ancestry is shown on different panels and in different colors. Data points represent the variance explained in individual cohorts used in this analysis.  $R^2$  was estimated in 1 to 11 cohorts with sample sizes ranging from 489 to 9,758 (**Supplementary Tables 8-11**).

**Figure 3 – Transferability of PGS across ancestries.** For each trait, the barplots represent trait variance explained when using a European ancestry-derived PGS in European, East Asian and African American test datasets. Variance explained (the height of each bar) for each trait (FG, FI and HbA1c) in each ancestry is shown on different panels and in different colors.

**Figure 4 - Trans-ancestry fine-mapping.** A) Number of plausible causal variants at each locus-trait association derived from FINEMAP. B) Number of variants within each 99% credible set. Twenty-one locus-trait associations at 19 loci were mapped to a single variant in the 99% credible set. C) Fine-mapping resolution. For each of the 98 locus-trait associations with a predicted single causal variant in both trans-ancestry and single-ancestry analyses, the number of variants included in the 99% credible set in the single-ancestry fine-mapping (x axis; logarithmic scale) is plotted against those in the trans-ancestry fine-mapping (y axis; logarithmic scale). Trans-ancestry and single-ancestry fine-mapping were based on the same set of variants. After removing eight locus-trait associations with one variant in the 99% credible sets in both trans-ancestry and single-ancestry analyses, there were 18 locus-trait associations (in grey) where trans-ancestry fine-mapping did not improve the resolution of fine-mapping results (i.e. number of variants in the 99% credible set did not decrease). Of the 72 locus-trait associations with improved trans-ancestry fine-mapping resolution (blue and red) further analyses in European fine-mapping emulating the total sample size in trans-ancestry fine-mapping demonstrated that 34 locus-trait associations (in red) were improved because of both total sample size and differences across ancestries, while 38 locus-trait associations (in blue) were only improved due to increased sample size in the original trans-ancestry fine-mapping analysis.

**Figure 5 - Epigenomic landscape of trait-associated variants.** A: Enrichment of GWAS variants to overlap genomic regions including ‘Static Annotations’ which are common or ‘static’ across cell types and ‘Stretch Enhancers’ which are identified in each tissue/cell type. The numbers of signals for each trait are indicated in parentheses. Enrichment was calculated using GREGOR<sup>56</sup>. One-sided test for significance (red) is determined after Bonferroni correction to account for 59 total annotations tested for each trait; nominal significance ( $P < 0.05$ ) is indicated in yellow. B: Enrichment for HbA1c GWAS signals partitioned into “hard” Glycemic and Red Blood Cell cluster (signals from “hard” mature Red Blood Cell and reticulocyte clusters together) to overlap annotations including StrEs in Islets and the blood-derived leukemia cell line K562, respectively (additional partitioned results in **Supplementary Table 17**). C: Individual FI GWAS signals that drive enrichment in Adipose and Skeletal Muscle StrEs. D, E: Genome browser shots of FI GWAS signals – intronic region of the *COL4A2* gene (D) and an inter-genic region ~25kb from *LINC01214* gene (E) showing GWAS SNPs (lead and LD  $r^2 > 0.8$  proxies), ATAC-seq signal tracks and chromatin state annotations in different tissues/cell types.

**Figure 6 - Tissues and cell types significantly enriched for genes within glycemic-associated loci.** Top panel FG-associated loci, middle panel FI-associated loci, bottom panel HbA1c-associated loci. FDR thresholds are shown in red ( $q < 0.05$ ), orange ( $q < 0.2$ ), black ( $q \geq 0.2$ ).

**Figure 7 - Gene-set enrichment analyses.** Results from affinity-propagation clustering of significantly enriched gene-sets ( $FDR < 0.05$ ) identified by DEPICT for A) FG, B) FI, and C) HbA1c. Each node is a meta gene-set which is represented by an exemplar gene-set within the meta gene-set. For example, in B. “chronic myeloid leukemia” is an exemplar gene-set representing a much broader meta gene-set relating to cancer and represented in the zoomed in section on the right. Similarities between the meta gene-sets are represented by Pearson correlation coefficients ( $r > 0.3$ ). The nodes are

1348 colored according to the minimum gene-set enrichment p-value for gene-sets in that meta gene-set..  
1349 PPI=protein-protein interaction network.  
1350

## Tables

**Table 1 – Glossary of terms** - This study combined analyses of trait-associations across multiple correlated glycemic traits and across multiple ancestries, which has presented challenges in our ability to apply commonly used terms with clarity. For this reason, we define below terms often used in the field with variable meaning, as well as definitions of new terms used in this study.

Term	Definition
EA (Effect allele)	The effect allele was that defined by METAL based on trans-ancestry FG results and aligned such that the same allele was kept as the effect allele across all ancestries and traits, irrespective of its allele frequency or effect size for that particular ancestry and trait, in this way the effect allele is not necessarily the trait-increasing allele.
Single-ancestry lead variant	Variant with the smallest p-value amongst all with $P < 5 \times 10^{-8}$ , within a 1Mb region, based on analysis of a single trait in a single ancestry.
Single-ancestry index variants	Variants identified by GCTA analysis of each autosome, and that appear to exert conditionally distinct effects on a given trait in a given ancestry ( $P < 5 \times 10^{-8}$ ). As defined, these include the single-ancestry lead variants.
Trans-ancestry lead variant	Variant identified by trans-ethnic meta-analysis of a given trait that has the strongest association for that trait ( $\log_{10}BF > 6$ , which is broadly equivalent to $P < 5 \times 10^{-8}$ ) within a 1Mb region.
Single-ancestry locus	1Mb region centred on a single-ancestry lead variant which does not contain a lead variant identified in the trans-ancestry meta-analysis (i.e., does not contain a trans-ancestry lead variant).
Signal	Conditionally independent association between a trait and a set of variants in LD with each other and which is noted by the corresponding index variant.
Trans-ancestry locus	A genomic interval that contains trans-ancestry trait-specific lead variants, with/out additional single-ancestry index variants, for one or more traits. This region is defined by starting at the telomere of each chromosome and selecting the first single-ancestry index variant or trans-ancestry lead variant for any trait. If other trans-ancestry lead variants or single-ancestry index variants mapped within 500kb of the first signal, then they were merged into the same locus. This process was repeated until there were no more signals within 500kb of the previous variant. A 500kb interval was added to the beginning of the first signal, and the end of the last signal to establish the final boundary of the trans-ancestry locus ( <b>Extended Data Figure 2</b> ). As defined, a trans-ancestry locus may not have a single lead trans-ancestry variant, but may instead contain multiple trans-ancestry lead variants, one for each trait.

## Online Methods

### *Study design and participants*

This study included trait data from four glycemic traits: fasting glucose (FG), fasting insulin (FI), 2hr post-challenge glucose (2hGlu), and glycated hemoglobin (HbA1c). The total number of contributing cohorts ranged from 41 (2hGlu) to 131 (FG), and the maximum sample size for each trait ranged from 85,916 (2hGlu) to 281,416 (FG) (**Supplementary Table 1**). Ancestry was initially defined at the cohort level, but within each cohort ancestry was confirmed with genetic data with ancestry outliers removed (**Supplementary Table 1**). Overall, European ancestry (EUR) participants dominated the sample size for all traits, representing between 68.0% (HbA1c) to 73.8% (2hGlu) of the overall sample size. African Americans (AA) represented between 1.7% (2hGlu) to 5.9% (FG) of participants; individuals of Hispanic ancestry (HISP) represented between 6.8% (FG) to 14.6% (2hGlu) of participants; individuals of East-Asian ancestry (EAS) represented between 9.9% (2hGlu) to 15.4% (HbA1c) of participants; and South-Asian ancestry (SAS) individuals represented between 0% (no contribution to 2hGlu) to 4.4% (HbA1c) of participants. Data from Ugandan participants were only available for the HbA1c analysis and represented 2% of participants.

### *Phenotypes*

Analyses included data for FG and 2hGlu measured in mmol/l, FI measured in pmol/l, and HbA1c in % [where possible, studies reported HbA1c as a National Glycohemoglobin Standardization Program (NGSP) percent]. Similar to previous MAGIC efforts<sup>7</sup>, individuals were excluded if they had type 1 or type 2 diabetes (defined by physician diagnosis); reported use of diabetes-relevant medication(s); or had a FG  $\geq 7$  mmol/L, 2hGlu  $\geq 11.1$  mmol/L, or HbA1c  $\geq 6.5\%$ , as detailed in **Supplementary Table 1**. 2hGlu measures were obtained 120 minutes after a glucose challenge in an oral glucose tolerance test (OGTT). Measures for FG and FI taken from whole blood were corrected to plasma level using the correction factor 1.13<sup>80</sup>.

### *Genotyping, quality control, and imputation*

Each participating cohort performed study-level quality control, imputation, and association analyses following a shared analysis plan. Cohorts were genotyped using commercially available genome-wide arrays or the Illumina CardioMetabochip (Metabochip) array (**Supplementary Table 1**)<sup>81</sup>. Prior to imputation, each cohort performed stringent sample and variant quality control (QC) to ensure only high-quality variants were kept in the genotype scaffold for imputation. Sample quality control checks included removing samples with low call rate  $< 95\%$ , extreme heterozygosity, sex mismatch with X chromosome variants, duplicates, first- or second-degree relatives (unless by design), or ancestry outliers. Following sample QC, cohorts applied variant QC thresholds for call rate ( $< 95\%$ ), Hardy-Weinberg Equilibrium (HWE)  $P < 1 \times 10^{-6}$ , and minor allele frequency (MAF). Full details of QC thresholds and exclusions by participating cohort are available in **Supplementary Table 1**.

Imputation was performed up to the 1000 Genomes Project phase 1 (v3) cosmopolitan reference panel<sup>82</sup>, with a small number of cohorts imputing up to the 1000 Genomes phase 3 panel<sup>19</sup> or population-specific reference panels (**Supplementary Table 1**).

### *Study level association analyses*

Each of the glycemic traits (FG, natural log FI, and 2hGlu) were regressed on BMI (except HbA1c), study-specific covariates, and principal components (unless implementing a linear mixed model). Analyses for FG, FI, and 2hGlu were adjusted for BMI as we had previously shown this did not materially affect results for FG and 2hGlu but improved our ability to detect FI-associated loci<sup>15</sup>. For simplicity, we refer to the traits as FG, FI and 2hGlu. For a discussion on collider bias see **Supplementary Note section 2c**. Both the raw and rank-based inverse normal transformed residuals from the regression were tested for association with genetic variants using SNPTTEST<sup>23</sup> or



Mach2Qtl<sup>83,84</sup>. Poorly imputed variants, defined as imputation  $r^2 < 0.4$  or INFO score  $< 0.4$ , were excluded from downstream analyses (**Supplementary Table 1**). Following study level QC, approximately 12,229,036 variants (GWAS cohorts) and 1,999,204 variants (Metabochip cohorts) were available for analysis (**Supplementary Table 1**).

### **Centralized quality control**

Each contributing cohort shared their summary statistic results with the central analysis group who performed additional QC using EasyQC<sup>85</sup>. Allele frequency estimates were compared to estimates from 1000Gp1 reference panel<sup>82</sup>, and variants were excluded from downstream analyses if there was a minor allele frequency difference  $> 0.2$  for AA, EUR, HISP, and EAS populations against AFR, EUR, MXL, and ASN populations from 1000 Genomes Phase 1, respectively, or a minor allele frequency difference  $> 0.4$  for SAS against EUR populations. At this stage, additional variants were excluded from each cohort file if they met one of the following criteria: were tri-allelic; had a minor allele count (MAC)  $< 3$ ; demonstrated a standard error of the effect size  $\geq 10$ ; or were missing an effect estimate, standard error, or imputation quality. All data that survived QC (approximately 12,186,053 variants from GWAS cohorts and 1,998,657 variants from Metabochip cohorts) were available for downstream meta-analyses.

### **Single-ancestry meta-analyses**

Single-ancestry meta-analyses were performed within each ancestry group using the fixed-effects inverse variance meta-analysis implemented in METAL<sup>20</sup>. We applied a double-genomic control (GC) correction<sup>15,86</sup> to both the study-specific GWAS results and the single-ancestry meta-analysis results. Study-specific Metabochip results were GC-corrected using 4,973 SNPs included on the Metabochip array for replication of associations with QT-interval, a phenotype not correlated with our glycemic traits<sup>15</sup>.

### **Identification of single-ancestry index variants**

To identify distinct association index variants across each chromosome within each ancestry (**Table 1**), we performed approximate conditional analyses implemented in GCTA<sup>21</sup> using the --cojo-slc option (autosomes) and distance-based clumping (X chromosome). Linkage disequilibrium (LD) correlations for GCTA were estimated from a representative cohort from each ancestry: WGHS (EUR); CHNS (EAS); SINDI (SAS); BioMe (AA); SOL (HISP) and Uganda (for itself). The results from GCTA were comparable when using alternative cohorts for the LD reference. For any index variant with a QC flag which caused reason for concern, we performed manual inspection of forest plots to decide whether the signal was likely to be real (**Supplementary note**). Among 335 single-ancestry index variants across all traits, this manual inspection was done for 40 signals of which 32 passed and 8 failed after inspection. Thus, a total of 327 single-ancestry index variants passed and 8 failed.

### **Trans-ancestry meta-analyses**

To leverage power across all ancestries, we also conducted trait-specific trans-ancestry meta-analysis by combining the single-ancestry meta-analysis results using MANTRA (**Supplementary note**)<sup>87</sup>. We defined  $\log_{10}$ Bayes' Factor (BF)  $> 6$  as genome-wide significant, approximately comparable to  $P < 5 \times 10^{-8}$ .

### **Manual curation of trans-ancestry lead variants**

To ensure trans-ancestry lead variants were robust, we performed manual inspection of forest plots by at least two authors, for any variants with flags indicating possible QC issues (**Supplementary note**). Of 463 trans-ancestry lead variants across all traits, 184 passed without inspection, 131 passed after inspection, and 148 failed after inspection.

### **Comparison of TA lead variants across ancestries**

For each pair of ancestries, we calculated Pearson's correlation in EAFs for each trans-ancestry lead variant. The pairwise summarized heterogeneity of effect sizes between ancestries was then tested using the joint F-test of heterogeneity<sup>32</sup>. The test statistic is the sum of Cochran Q-statistics for heterogeneity across all trans-ancestry signals. Under the null hypothesis, the statistics follows the  $\chi^2$  distribution with n degrees of freedom, where n is the number of the trans-ancestry lead variants.

#### **LD-pruned variant lists**

Several downstream analyses (for example, genomic feature enrichment, genetic scores, and estimation of variance explained by associated variants) require independent LD-pruned variants ( $r^2 < 0.1$ ) to avoid double-counting variants which might otherwise be in LD with each other and that do not provide additional "independent" evidence. Therefore, for these analyses we generated different lists of either TA or single-ancestry LD pruned ( $r^2 < 0.1$ ) variants, keeping in each case the variant with the strongest evidence of association (**Supplementary Table 7**). Subsequently, we combined TA and single-ancestry variant lists and conducted further LD pruning. For some analyses, we took the TA pruned variant list and added single-ancestry signals if the LD  $r^2 < 0.1$ , while for others we started with the single-ancestry pruned lists and supplemented with TA lead variants if the LD  $r^2 < 0.1$ . One exception was the list used for eQTL co-localizations, which included all single-ancestry European signals (without LD pruning) and supplemented with any additional TA lead variants (starting from the variants with the most significant P-values) in EUR LD  $r^2 < 0.1$  with any of the variants already in list, and that reached at least  $P < 1 \times 10^{-5}$  in the European ancestry meta-analysis.

#### **Trait variance explained by associated loci**

To determine how much of the phenotypic variance of each trait could be explained by the corresponding trait-associated loci, variants were combined in a series of weighted genetic scores (GS). The analysis was performed in a subset of the cohorts included in the discovery GWAS (with representation from each ancestry) and in a smaller number of independent cohorts (European ancestry only). Up to three different GS were derived per trait (and for each ancestry) in order to evaluate the potential for the trans-ancestry meta-GWAS identified loci to provide additional information above and beyond that contributed by the ancestry-specific meta-analysis results. These GS comprised: List A - single-ancestry signals; List B - single-ancestry signals plus trans-ancestry signals; and List C - trans-ancestry signals plus single-ancestry signals (**Supplementary Table 7**). In the case of the European ancestry cohorts that contributed to the GWAS, we employed the method of Nolte *et al.*<sup>33</sup> to adjust the effect sizes (betas) from the GWAS for the contribution of that cohort, providing sets of cohort-specific effect sizes that were then used to generate the GS. The association between each GS and its corresponding trait was tested by linear regression and the adjusted  $R^2$  from the model extracted as an estimate of the variance explained.

#### **Transferability of polygenic scores (PGS) across ancestries**

We used the PRS-CSauto<sup>34</sup> software to first build European ancestry-derived PGS for each glycemic trait (FG, FI, 2hGlu, HbA1c) on the basis of summary statistics. However, PRS-CSauto does not perform well when the training dataset is relatively small and the genetic architecture is sparse<sup>34</sup>. Consequently, 2hGlu was excluded from this analysis. For each trait, to obtain European ancestry training and test datasets, we first removed all cohorts only genotyped on the Metabochip which were not included in this analysis. From the remaining cohorts we then removed five of the largest European cohorts contributing to the respective European ancestry meta-analysis. For each trait, these five cohorts were meta-analyzed and used as the European ancestry test dataset. Subsequently, the remaining European ancestry cohorts were also meta-analyzed and used as the European ancestry training dataset. For each of the other ancestries, cohorts only genotyped on the Metabochip were also removed, and the remaining cohorts were meta-analyzed, and used as the non-European ancestry test datasets. Variants with MAF < 0.05 or missing in over half of the

individuals in the training dataset were removed<sup>34,88</sup>. The PGS for each trait was built using PRS-CSauto with default settings<sup>34</sup> with the effect size estimates based on the European training dataset being revised based on an LD reference panel matching the test dataset. The proportion of the trait variance explained by the European ancestry-derived PGS ( $R^2$ ) was estimated using the R package “gtx”<sup>89</sup> based on the revised effect sizes and summary statistics from the test dataset for each ancestry.

### **Fine-mapping**

Of the 242 loci identified in this study, 237 were autosomal loci which we took forward for fine-mapping (**Supplementary Table 2**). We used the Bayesian fine-mapping method FINEMAP<sup>90</sup> (version 1.1) to refine association signals and attempt to identify likely causal variants at each locus. FINEMAP estimates the maximum number of causal variants at each locus, calculates the posterior probability of each variant being causal, and proposes the most likely configuration of causal variants. The posterior probabilities of the configurations in each locus were used to construct 99% credible sets.

We performed both single-ancestry and trans-ancestry fine-mapping. In both analyses, only data from cohorts genotyped on GWAS arrays were used, and analyses were limited to trans-ancestry lead variants and other single-ancestry lead variants present in at least 90% of the samples for each trait. For the single-ancestry fine-mapping, FINEMAP estimates the number of causal variants in a region up to a maximum number, which we set to be two plus the number of distinct signals identified from the GCTA signal selection. FINEMAP uses single-ancestry and trait-specific z-scores from the fixed-effect meta-analysis in METAL<sup>20</sup> and an ancestry-specific LD reference, which we created from a subset of cohorts (combined sample size > 30% of the sample size for that ancestry), weighting each cohort by sample size. In the trans-ancestry fine-mapping, FINEMAP was similarly used to estimate the number of causal variants starting with two, and trait-specific z-scores and LD maps were generated from the sample size weighted average of those used in the single-ancestry fine-mapping. The maximum number of causal variants was iteratively increased by one until it was larger than the number of causal variants supported by data (Bayes factor), which was the estimated maximum number of causal variants used in the final run of fine-mapping analysis.

To compare fine-mapping results obtained from the single-ancestry and trans-ancestry efforts, analyses were limited to fine-mapping regions with evidence for a single likely causal variant in both, enabling a straightforward comparison of credible sets (**Supplementary note**). To ensure any difference in the fine-mapping results was not driven by different sets of variants being present in the different analyses, we repeated the single-ancestry fine-mapping limited to the same set of variants used in the trans-ancestry fine-mapping. The fine-mapping resolution was assessed based on comparisons of the 99% credible sets in terms of number of variants included in the set, and length of the region. To assess whether the improvement in the trans-ancestry fine-mapping was due to differences in LD, increased sample size, or both, we repeated the trans-ancestry fine-mapping mimicking the sample size present in the single-ancestry fine-mapping by dividing the standard errors by the square root of the sample size ratio and compared the results with those from the single-ancestry fine-mapping.

### **Functional Annotation of trait-associated variants**

#### ***HbA1c signal classification***

There were 218 HbA1c-associated signals from either the single-ancestry (i.e. all GCTA-signals from any ancestry) or trans-ancestry meta-analyses. To classify these signals in terms of their likely mode of action (i.e., glycemic, erythrocytic, or other<sup>7</sup>), we examined association summary statistics for the

lead variants at the 218 signals in other large European datasets for 19 additional traits: three glycemic traits from this study (FG, 2hGlu and FI); seven mature red blood cell (RBC) traits<sup>91,92</sup> (red blood cell count, mean corpuscular volume, hematocrit, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration, hemoglobin concentration and red cell distribution width); five reticulocyte traits (reticulocyte count, reticulocyte fraction of red cells, immature fraction of reticulocytes, high light scatter reticulocyte count and high light scatter percentage of red cells)<sup>91,92</sup>, and four iron traits (serum iron, transferrin, transferrin saturation and ferritin)<sup>93</sup>. Of the 218 HbA1c signals, data were available for the lead (n=183) or proxy (European LD  $r^2 > 0.8$ , n = 8) variants at 191 signals.

The additional traits were clustered using hierarchical clustering to ensure biologically related traits would cluster together (**Supplementary note**). We then used a non-negative matrix factorization (NMF)<sup>94</sup> process to cluster the HbA1c signals. Each cluster was labelled as glycemic, reticulocyte, mature RBC, or iron related based on the strength of association of signals in the cluster to the glycemic, reticulocyte, mature RBC and iron traits (**Supplementary note**). To verify that our cluster naming was correct, we used HbA1c association results conditioned on either FG or iron traits, or type 2 diabetes association results (**Supplementary note**).

#### ***HbA1c genetic risk scores (GRSs) and T2D risk***

We constructed GRS for each cluster of HbA1c-associated signals (based on hard clustering) and tested the association of each cluster with T2D risk using samples from the UK Biobank. Pairs of HbA1c signals in LD ( $r^2 > 0.10$ ) were LD pruned by removing the signal with the less significant *P*-value of association with HbA1c. The GRS for each cluster was calculated based on the logarithm of odds ratios from the latest T2D study summary statistics<sup>95</sup> and UK Biobank genotypes imputed to the Haplotype Reference Consortium<sup>19</sup>. From 487,409 UK Biobank samples (age between 46 and 82 years, and 55% female), we excluded participants for the following reasons: 373 with mismatched sex; 9 not used in the kinship calculation; 78,365 non-European ancestry individuals; and 138,504 with missing T2D status, age, or sex information. We further removed 26,896 related participants (kinship  $> 0.088$ , preferentially removing individuals with the largest number of relatives and controls where a T2D case was related to a control). T2D cases were defined by: (i) a history of diabetes without metformin or insulin treatment, (ii) self-reported diagnosis of T2D, or (iii) diagnosis of T2D in a national registry (N = 17,022, age between 47 and 79 years, and 36% female). Controls were participants without a history of T2D (N = 226,240, age between 46 and 82 years, and 56% female). We tested for association between each GRS and T2D using logistic regression including covariates for age, sex, and the first five principal components. Significance of association was evaluated by a bootstrap approach to incorporate the variance of each HbA1c associated signal in the T2D summary data. To do this, we generated the GRS of each cluster 200 times by resampling the logarithm of odds ratio of each signal with T2D. For each non-glycemic class that had a GRS significantly associated with T2D, we performed sensitivity analyses to evaluate whether the association was driven from variants that also belonged to a glycemic cluster when using a soft clustering approach (the signals were classified as also glycemic in the soft clustering or had an association  $P \leq 0.05$  with any of the three glycemic traits).

#### ***Chromatin states***

To identify genetic variants within association signals that overlapped predicted chromatin states, we used a previously published, 13 chromatin state model that included 31 diverse tissues, including pancreatic islets, skeletal muscle, adipose, and liver<sup>39</sup>. Briefly, this model was generated from cell/tissue ChIP-seq data for H3K27ac, H3K27me3, H3K36me3, H3K4me1, and H3K4me3, and input control from a diverse set of publicly available data<sup>53,57,96,97</sup> using the ChromHMM program<sup>98</sup>. As reported previously<sup>39</sup>, StrEs were defined as contiguous enhancer chromatin state (Active Enhancer 1 and 2, Genic Enhancer and Weak Enhancer) segments longer than 3kb<sup>57</sup>.

### ***Enrichment of genetic variants in genomic features***

We used GREGOR (version 1.2.1) to calculate the enrichment of GWAS variants overlapping static and StrEs<sup>56</sup>. For calculating the enrichment of glycemic trait-associated variants in these annotations, we used the filtered list of trait-associated variants as described above (**Supplementary Table 7**) as input. For calculating the enrichment of sub-classified HbA1c variants, we included the list of loci characterized as Glycemic, another list of loci characterized as Reticulocyte or mature Red Blood Cell, collectively representing the red blood cell fraction, along with lists of iron related or unclassified loci (**Supplementary Table 17**). We used the following parameters in GREGOR enrichment analyses: European  $r^2$  threshold (for inclusion of variants in LD with the lead variant) = 0.8, LD window size = 1 Mb, and minimum neighbour number = 500.

We used fGWAS (version 0.3.6)<sup>58</sup> to calculate enrichment of glycemic trait-associated variants in static and StrE annotations using summary level GWAS results. We used the default fGWAS parameters for enrichment analyses for individual annotations for each trait. For each annotation, the model provided the natural log of maximum likelihood estimate of the enrichment parameter. Annotations were considered as significantly enriched if the log2 (parameter estimate) and respective 95% confidence intervals were above zero or significantly depleted if the log2 (parameter estimate) and respective 95% confidence intervals were below zero.

We tested enrichment of trait-associated variants in static and StrE annotations with GARFIELD (v2)<sup>59</sup>. We formatted annotation overlap files as required by the tool; prepared input data at two GWAS thresholds - of  $1 \times 10^{-5}$  and a more stringent  $1 \times 10^{-8}$  by pruning and clumping with default parameters (garfield-prep-chr script). We calculated enrichment in each individual annotation using garfield-test.R with  $-c$  option set to 0. We also calculated the effective number of annotations using the garfield-Meff-Padj.R script. We used the effective number of annotations for each trait to obtain Bonferroni corrected significance thresholds for enrichment for each trait.

### ***eQTL analyses***

To aid in the identification of candidate casual genes at the European-only and trans-ancestry association signals, we examined whether any of the lead variants associated with glycemic traits (**Supplementary Table 7**) were also associated with expression level (FDR < 5%) of nearby transcripts located within 1 Mb in existing eQTL data sets of blood, subcutaneous adipose, visceral adipose, skeletal muscle, and pancreatic islet samples<sup>60,61,99-102</sup>. LD was estimated from the collected cohort pairwise LD information, where available, else from the European samples in 1000G phase 3. GWAS and eQTL signals likely co-localize when the GWAS variant and the variant most strongly associated with the expression level of the corresponding transcript (eSNP) exhibit high pairwise LD ( $r^2 > 0.8$ ; 1000 Genomes Phase 3, EUR). At these signals, we conducted reciprocal conditional analyses to test association between the GWAS variant and transcript level when the eSNP was also included in the model, and vice versa. We report GWAS and eQTL signals as co-localized if the association for the eSNP was not significant (FDR  $\geq 5\%$ ) when conditioned on the GWAS variant; we also report signals from the eQTLGen whole blood meta-analysis data that meet only the LD threshold because conditional analysis was not possible.

### ***Tissue and gene-set analysis***

We performed enrichment analysis using DEPICT (Data-driven Expression-Prioritized Integration for Complex Traits) version 3, specifically developed for 1000 Genomes Project imputed meta-analysis data<sup>103</sup> to identify cell types and tissues in which genes at trait-associated variants were strongly expressed, and to detect enrichment of gene-sets or pathways. DEPICT data included human gene expression data for 19,987 genes in 10,968 reconstituted gene sets, and 209 tissues/cell types. Because gene expression data in DEPICT is based on European samples and LD, we selected trait-associated variants with  $P < 10^{-5}$  in the European meta-analysis and tested for enrichment of signals in

each reconstituted gene-set, and each tissue or cell type. Enrichment results with a false discovery rate (FDR)<0.05 were considered significant. We ran DEPICT based on association results for all traits among: (i) cohorts with genome-wide data, or (ii) all cohorts (genome-wide and Metabochip cohorts). Because results were broadly consistent between the two approaches, we present results from the analysis that contained all cohorts as it had greater statistical power.

## **Statistics and reproducibility**

### **Sample size**

No statistical method was used to predetermine sample size. We aimed to bring together the largest possible sample size with GWAS data from individuals of diverse ancestries (European, Hispanic, African American, East Asian, South Asian and sub-Saharan African) without diabetes and with data for one or more of the following traits: fasting glucose, fasting insulin, 2hr post-challenge glucose, and glycated hemoglobin. The sample sizes were 281,416 (FG), 213,650 (FI), 215,977 (HbA1c) and 85,916 (2hGlu) (**Supplementary Table 1**). Our sample size was sufficiently powered to detect common variant associations with each of the glycemic traits and was able to detect associations at 242 loci.

### **Randomization/ Blinding**

This is a study of continuous traits therefore there were no experiments to randomize and there was no “outcome” to which investigators needed to be blinded to.

### **Data exclusions**

Prior to conducting this study, we identified reasons for which data should be excluded from the analysis at either the cohort or summary level; these exclusions are as follows. Sample quality control checks included removing samples with low call rate < 95%, extreme heterozygosity, sex mismatch with X chromosome variants, duplicates, first- or second-degree relatives (unless by design), or ancestry outliers. Following sample QC, cohorts applied variant QC thresholds for call rate (< 95%), Hardy-Weinberg Equilibrium (HWE)  $P < 1 \times 10^{-6}$ , and minor allele frequency (MAF). Full details of QC thresholds and exclusions by participating cohort are available in **Supplementary Table 1**. Each contributing cohort shared their summary statistic results with the central analysis group who performed additional QC using EasyQC. Allele frequency estimates were compared to estimates from 1000Gp1 reference panel, and variants were excluded from downstream analyses if there was a minor allele frequency difference > 0.2 for AA, EUR, HISP, and EAS populations against AFR, EUR, MXL, and ASN populations from 1000 Genomes Phase 1, respectively, or a minor allele frequency difference > 0.4 for SAS against EUR populations. At this stage, additional variants were excluded from each cohort file if they met one of the following criteria: were tri-allelic; had a minor allele count (MAC) < 3; demonstrated a standard error of the effect size  $\geq 10$ ; imputation  $r^2 < 0.4$  or INFO score < 0.4; or were missing an effect estimate, standard error, or imputation quality.

### **Data Availability**

Ancestry-specific and overall meta-analysis summary level results are available through the MAGIC website (<https://www.magicinvestigators.org/>). Summary statistics are also available through the GWAS catalogue (<https://www.ebi.ac.uk/gwas/>) with the following accession codes: GCST90002225, GCST90002226, GCST90002227, GCST90002228, GCST90002229, GCST90002230, GCST90002231, GCST90002232, GCST90002233, GCST90002234, GCST90002235, GCST90002236, GCST90002237, GCST90002238, GCST90002239, GCST90002240, GCST90002241, GCST90002242, GCST90002243, GCST90002244, GCST90002245, GCST90002246, GCST90002247, and GCST90002248.

### Code availability

Source code implementing methods described in the paper are publicly available on <https://zenodo.org/badge/latestdoi/346687844>.

### References for Methods

- 80 D'Orazio, P. *et al.* Approved IFCC recommendation on reporting results for blood glucose (abbreviated). *Clinical chemistry* **51**, 1573-1576, doi:10.1373/clinchem.2005.051979 (2005).
- 81 Voight, B. F. *et al.* The metabochip, a custom genotyping array for genetic studies of metabolic, cardiovascular, and anthropometric traits. *PLoS genetics* **8**, e1002793, doi:10.1371/journal.pgen.1002793 (2012).
- 82 Abecasis, G. R. *et al.* An integrated map of genetic variation from 1,092 human genomes. *Nature* **491**, 56-65, doi:10.1038/nature11632 (2012).
- 83 Li, Y., Willer, C. J., Ding, J., Scheet, P. & Abecasis, G. R. MaCH: using sequence and genotype data to estimate haplotypes and unobserved genotypes. *Genetic epidemiology* **34**, 816-834, doi:10.1002/gepi.20533 (2010).
- 84 Pei, Y. F., Zhang, L., Li, J. & Deng, H. W. Analyses and comparison of imputation-based association methods. *PloS one* **5**, e10827, doi:10.1371/journal.pone.0010827 (2010).
- 85 Winkler, T. W. *et al.* Quality control and conduct of genome-wide association meta-analyses. *Nature protocols* **9**, 1192-1212, doi:10.1038/nprot.2014.071 (2014).
- 86 Devlin, B. & Roeder, K. Genomic control for association studies. *Biometrics* **55**, 997-1004 (1999).
- 87 Morris, A. P. Transethnic meta-analysis of genomewide association studies. *Genetic epidemiology* **35**, 809-822, doi:10.1002/gepi.20630 (2011).
- 88 Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nature genetics* **47**, 291-295, doi:10.1038/ng.3211 (2015).
- 89 Dastani, Z. *et al.* Novel loci for adiponectin levels and their influence on type 2 diabetes and metabolic traits: a multi-ethnic meta-analysis of 45,891 individuals. *PLoS genetics* **8**, e1002607, doi:10.1371/journal.pgen.1002607 (2012).
- 90 Benner, C. *et al.* FINEMAP: efficient variable selection using summary data from genome-wide association studies. *Bioinformatics (Oxford, England)* **32**, 1493-1501, doi:10.1093/bioinformatics/btw018 (2016).
- 91 Astle, W. J. *et al.* The Allelic Landscape of Human Blood Cell Trait Variation and Links to Common Complex Disease. *Cell* **167**, 1415-1429.e1419, doi:10.1016/j.cell.2016.10.042 (2016).
- 92 Canela-Xandri, O., Rawlik, K. & Tenesa, A. An atlas of genetic associations in UK Biobank. *Nature genetics* **50**, 1593-1599, doi:10.1038/s41588-018-0248-z (2018).
- 93 Benyamin, B. *et al.* Novel loci affecting iron homeostasis and their effects in individuals at risk for hemochromatosis. *Nature communications* **5**, 4926, doi:10.1038/ncomms5926 (2014).
- 94 Binesh, N. & Rezghi, M. Fuzzy clustering in community detection based on nonnegative matrix factorization with two novel evaluation criteria. *Applied Soft Computing* **69**, 689-703 (2018).
- 95 Scott, R. A. *et al.* An Expanded Genome-Wide Association Study of Type 2 Diabetes in Europeans. *Diabetes* **66**, 2888-2902, doi:10.2337/db16-1253 (2017).

1772 96 Ernst, J. *et al.* Mapping and analysis of chromatin state dynamics in nine human cell  
1773 types. *Nature* **473**, 43-49, doi:10.1038/nature09906 (2011).

1774 97 Mikkelsen, T. S. *et al.* Comparative epigenomic analysis of murine and human  
1775 adipogenesis. *Cell* **143**, 156-169, doi:10.1016/j.cell.2010.09.006 (2010).

1776 98 Ernst, J. & Kellis, M. ChromHMM: automating chromatin-state discovery and  
1777 characterization. *Nature methods* **9**, 215-216, doi:10.1038/nmeth.1906 (2012).

1778 99 Battle, A., Brown, C. D., Engelhardt, B. E. & Montgomery, S. B. Genetic effects on  
1779 gene expression across human tissues. *Nature* **550**, 204-213,  
1780 doi:10.1038/nature24277 (2017).

1781 100 Zhernakova, D. V. *et al.* Identification of context-dependent expression quantitative  
1782 trait loci in whole blood. *Nature genetics* **49**, 139-145, doi:10.1038/ng.3737 (2017).

1783 101 Westra, H. J. *et al.* Systematic identification of trans eQTLs as putative drivers of  
1784 known disease associations. *Nature genetics* **45**, 1238-1243, doi:10.1038/ng.2756  
1785 (2013).

1786 102 Joehanes, R. *et al.* Integrated genome-wide analysis of expression quantitative trait  
1787 loci aids interpretation of genomic association studies. *Genome biology* **18**, 16,  
1788 doi:10.1186/s13059-016-1142-6 (2017).

1789 103 Pers, T. H. *et al.* Biological interpretation of genome-wide association studies using  
1790 predicted gene functions. *Nature communications* **6**, 5890,  
1791 doi:10.1038/ncomms6890 (2015).

1792