

THE UNIVERSITY of EDINBURGH

Edinburgh Research Explorer

ArcGeomorphometry: A toolbox for geomorphometric characterization of DEMs in the ArcGIS environment

Citation for published version:

Stuart, N 2015, 'ArcGeomorphometry: A toolbox for geomorphometric characterization of DEMs in the ArcGIS environment', Computers and Geosciences. https://doi.org/10.1016/j.cageo.2015.09.020

Digital Object Identifier (DOI):

10.1016/j.cageo.2015.09.020

Link: Link to publication record in Edinburgh Research Explorer

Document Version: Peer reviewed version

Published In: Computers and Geosciences

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1	ArcGeomorphometry: A toolbox for geomorphometric characterization of DEMs	
2	in the ArcGIS environment	
3 4 5	Juan P. Rigol-Sanchez ^{a,b,*} , Neil Stuart ^c , Antonio Pulido-Bosch ^a	
5	"Water Personness and Environmental Coology Department of Pielogy and Coology	
0 7	Water Resources and Environmental Geology, Department of Biology and Geology,	
/ 0	^b Department of Coology University of Jacop E 22071 Jacop Spain	
0	⁶ Department of Geology, University of Jaen, E-250/1 Jaen, Spain	
9	⁵ Institute of Geography, School of GeoSciences, University of Edinburgh, Edinburgh	
10	EF18 9XP, Scotland, United Kingdom	
11	* Company and in a path on	
12	* Corresponding author.	
13	<i>E-mail addresses</i> : <u>jprigol@ual.es</u> (J. P. Rigol-Sanchez), <u>N.Stuart@ed.ac.uk</u> (N. Stuart),	
14	<u>apulido(@ual.es</u> (A. Pulido-Bosch).	
15		
16		
17	Abstract	
18	A software tool is described for the extraction of geomorphometric land surface	
19	variables and features from Digital Elevation Models (DEMs). The	
20	ArcGeomorphometry Toolbox consists of a series of Python/Numpy processing	
21	functions, presented through an easy-to-use graphical menu for the widely used ArcGIS	
22	package. Although many GIS provide some operations for analysing DEMs, the	
23	methods are often only partially implemented and can be difficult to find and used	
24	effectively. Since the results of automated characterisation of landscapes from DEMs	
25	are influenced by the extent being considered, the resolution of the source DEM and the	
26	size of the kernel (analysis window) used for processing, we have developed a tool to	
27	allow GIS users to flexibly apply several multi-scale analysis methods to parameterise	
28	and classify a DEM into discrete land surface units. Users can control the threshold	
29	values for land surface classifications. The size of the processing kernel can be used to	
30	identify land surface features across a range of landscape scales. The pattern of land	

31 surface units from each attempt at classification is displayed immediately and can then be processed in the GIS alongside additional data that can assist with a visual 32 33 assessment and comparison of a series of results. The functionality of the ArcGeomorphometry toolbox is described using an example DEM. 34

35

36 *Keywords:* Geomorphometry; DEM; GIS; Python; Numpy; Digital Terrain Analysis.

- 37
- 38 39

1. Introduction

40

41 The analysis and classification of the land surface at various landscape scales is 42 a prerequisite for many studies within the geosciences. In the last two decades 43 geomorphometry – the discipline of quantitative land-surface analysis – has undergone 44 rapid progress due to the flexibility and rapidity with which the required computations 45 can now be performed through the computerized analysis of digital elevation models 46 (DEMs) (Pike, 2000; Pike et al., 2009). DEM analysis is now used to characterise and to 47 extract relevant landscape features in fields as diverse as geomorphology, surface 48 hydrology, visual impact assessment, watershed management, land management, 49 cellular telecommunications, civil engineering, oceanography, ecology, soil science, 50 planetary science, wind energy planning. The almost global coverage of gridded DEMs 51 at resolutions between 30-90m, from sources such as the ASTER Global Digital 52 Elevation Model (GDEM) and the Shuttle Radar Topographic Mission (SRTM) has 53 renewed interest in semi-automatic methods for the characterisation of contrasting 54 landscapes and for consistently identifying what Lueder (1959) defines as second-order 55 of relief features such as mountain ranges and plains and third-order relief features such 56 as individual hills, mountains and valleys.

57 Although the basic DEM processing can be conducted almost automatically, 58 there is still a need for user interaction at various stages, for example to review the 59 effects of different analyses and parameterisations, to compare the results of alternative landscape segmentations and classifications and to interpret and to contextualize the 60 61 results, especially when performed at multiple scales. The ability to visually explore and 62 compare many results along with the availability of faster and friendlier GIS toolboxes 63 have been recognised as important new developments in geomorphometry software (Wood, 2009a; Gessler et al., 2009). Gessler et al. (2009) have identified a number of 64 65 topics needing research in the field of geomorphometry. They include, among others, algorithm development for true multi-scale characterisation, maintaining operational 66 67 ease of use despite increasing complexity of morphometric procedures, and tools for 68 static and dynamic visualisation of measures and surface objects. Consequently, there is

69 a need for multi-scale land surface analysis and visualization tools that facilitate 70 common tasks such as performing multi-scale analyses and exploring the results of 71 using different analysis window sizes and classification parameters and hence finding 72 appropriate settings for identifying landscape characteristics and specific 73 geomorphometric features.

74 Previously, the analysis of DEMs was usually conducted using specialist, stand-75 alone software programs. However, the widespread adoption of GIS in academic, 76 professional and commercial arenas, the increased processing power of these systems 77 for handling and visualising DEMs and the large volumes of spatial information now available in GIS formats are practical drivers for greater land surface analysis 78 79 functionality to be included within GIS. As one means of achieving this, we present 80 here the ArcGeomorphometry tools for geomorphometric characterisation of DEMs in 81 the ArcGIS environment. The tools are implemented in Python/Numpy and enable a 82 wide range of analyses to be conducted efficiently on DEMs. To understand the range 83 of methods presently supported, the more common digital methods for land surface 84 analysis are briefly reviewed. The functionality of the ArcGeomorphometry toolbox is 85 then presented and compared to other existing software to locate it between the more 86 comprehensive, specialist tools and the more limited functionality found in commercial 87 GIS. The key features and operations of ArcGeomorphometry are described and 88 illustrated using an example DEM. Conclusions are then drawn about the utility of the 89 ArcGeomorphometry tools and scope for its further enhancement indicated.

- 90
- 91

2. The analysis of the land surface using digital methods

92

93 Geomorphometry is the science of quantitative land-surface analysis (Pike, 94 1995). Information produced by geomorphometry supports the study of many earth 95 surface processes where landforms act as a controlling or boundary condition (Dehn et 96 al., 2001). Applicable at different scales, geomorphometric analysis can range from the 97 identification of localised landforms through to the characterisation of extensive 98 regional or continental landscapes (Pike, 2000). This leads to the important distinction 99 between specific and general geomorphometry (Evans, 1972). While specific 100 geomorphometry analyses the geometric and topological characteristics of 'landforms' 101 (i.e. bounded segments of a land surface that are discrete and may be discontinuous), 102 general geomorphometry analyses 'land surface form' (i.e. a continuous field that covers the whole globe) (Evans, 2012). Thus, the related variables are object-based and
field-based (see Evans and Minar, 2011, for a comprehensive classification of the
fundamental variables).

106 A variety of equations have been proposed to calculate the fundamental 107 geomorphometric variables. Well known examples include Evans (1972, 1979, 1980), 108 Band (1986), Jenson and Domingue (1988), Pennock et al. (1987), Zevenbergen and 109 Thorne (1987), Dikau (1989), Moore et al. (1993), Shary (1995), Wood (1996), 110 Florinsky (1998), Wilson and Gallant (2000), Shary et al. (2002) and Schmidt et al. 111 (2003). The present study is focused on the algorithms for the calculation of field local 112 variables, therefore methods for calculating object and regional variables (e.g stream 113 order, distance to stream, catchment area) are not discussed here. In this regard Evans' 114 approach is the most widely used method in relation to field local variables.

Evans' method is based on fitting a second-order polynomial function to elevation in a central point and its neighbours and then deriving local gradient and curvatures (mutually orthogonal — profile and plan curvatures, and minimum and maximum curvatures) from the function:

119
$$z = ax^2 + by^2 + cxy + dx + ey + f$$
 (1)

where *a* to *f* are quadratic coefficients, *x* and *y* are local spatial coordinates, and *z* is elevation. Gradient and curvatures ($[L^{-1}]$) can be derived as (Evans, 1972, 1979, 1980; Schmidt et al., 2003):

123
$$G = (d^2 + e^{2)^{1/2}}$$
 (2)

124
$$C_p = -\frac{ad^2 + 2cde + be^2}{(d^2 + e^2)(1 + d^2 + e^2)^{3/2}}$$
 (3)

125
$$C_c = -\frac{ae^2 - 2cde + bd^2}{(d^2 + e^2)^{3/2}}$$
 (4)

126
$$C_{p-\min} = -a - b - ((a-b)^2 + c^2)^{1/2}$$
 (5)

127
$$C_{p-\max} = -a - b + ((a-b)^2 + c^2)^{1/2}$$
 (6)

where *G* is gradient, C_p is profile curvature, C_c is contour curvature, C_{p-min} is minimum profile curvature, and C_{p-max} is maximum profile curvature.

Several extensions to Evans' method have been proposed (Zevenbergen and
Thorne, 1987; Shary, 1995; Wood, 1996; Shary et al., 2002). Zevenbergen and Thorne
(1987) extended Evans' method for estimating land surface slope gradient and curvature

133 by fitting a (partial) fourth-order polynomial surface to elevation values within a 134 processing 3×3 window centred on a particular cell of a DEM. Shary (1995) extended 135 Evans's method and proposed several new curvature measures, distinguishing those that 136 depend on gravity (i.e. slope) (e.g. rotor, difference curvature) from those that are independent of slope and are derived using only surface geometry (e.g. unsphericity,). 137 138 Shary (1995) used a quadratic polynomial function and a linear equation system as 139 Evans (1980) but forced the locally interpolated surface to match the elevation of the 140 central point of the 3×3 window centred at a particular cell (Schmidt et al., 2003). These 141 measures can be derived from Eq. (1) as (see Shary, 1995, Shary et al., 2002, and 142 Schmidt et al., 2003, for a complete set of formulae):

143
$$C_f = \frac{c(d^2 - e^2) - de(a - b)}{(d^2 + e^2)^{3/2}}$$
(7)

144
$$C_m = -\frac{a(1+e^2) - 2cde + b(1+d^2)}{2(1+d^2+e^2)^{3/2}}$$
 (8)

145
$$C_g = -\frac{ab-c^2}{(1+d^2+e^2)^2}$$
 (9)

146
$$C_{tr} = \frac{c(d^2 - e^2) - de(a - b)^2}{(d^2 + e^2)^2 (1 + d^2 + e^2)^2}$$
(10)

147
$$C_{tot} = a^2 + 2c^2 + b^2$$
 (11)

148
$$C_{t} = -\frac{ae^{2} - 2cde + bd^{2}}{(d^{2} + e^{2})(1 + d^{2} + e^{2})^{1/2}}$$
(12)

where C_f is flowpath curvature or *rotor*, C_m is mean curvature, C_g is total Gaussian curvature, C_{tr} is total ring curvature, C_{tot} is total curvature, and C_t is tangential curvature. Other proposed curvature measures can be derived combining curvatures (3) to (12) above. Shary et al. (2002) also proposed a pre-filtering for Evans algorithm for curvature calculation that does not emphasize grid directions, which they termed *modified Evans–Young* algorithm.

157
$$C_l = -\frac{ad^2 + 2cde + be^2}{(d^2 + e^2)}$$
 (13)

158
$$C_s = -\frac{ae^2 - 2cde + bd^2}{(d^2 + e^2)}$$
(14)

where C_l is longitudinal curvature, C_s is cross-sectional curvature, and a to f are 159 160 quadratic coefficients as above. Note that Eq. (13) and (14) are those rewritten by 161 Schmidt et al. (2003) for uniformity of equations (2) to (14) (cf. Wood, 1996; curvature 162 measures of dimension $[L^{-1}]$). Schmidt et al. (2003) reviewed and compared the 163 algorithms for land surface curvature calculation proposed by Evans (1980), 164 Zevenbergen and Thorne (1987) and Shary (1995). They concluded that a local surface 165 representation derived from quadratic models ('Evans' and 'Shary') is more useful to 166 consistently describe local surface curvature, and to model the land surface by basic 167 land elements.

168 Wood (1996) made an important contribution to multi-scale geomorphometric 169 analysis by implementing a generalisation of Evans' approach to broader operational 170 scales. Evans' original approach was limited to computing local slope gradient and 171 curvature of land surface by analysing only the cell values within a 3×3 window (or 172 kernel) of neighbouring cells. In high resolution (e.g. <5 m peg spacing) DEMs, this 173 may detect only micro-scale anomalies in the land surface. MacMillan and Shary (2009) 174 concluded that it is not possible to select any single fixed dimension for a moving 175 window that will perfectly capture the wavelength of all landform features of interest in 176 any given area. However, most geomorphometric variables are calculated by moving a 177 3×3 window across a DEM and calculating the values for the central cell in the window 178 (Pike et al., 2009; Dragut and Eisank, 2011; Wilson, 2012). For instance, curvature 179 values are typically computed within a 3×3 window, but clear advantages to computing 180 curvatures within a series of larger neighbourhood analysis windows have been 181 demonstrated by authors such as Dikau (1989), Wood (1996), and Smith et al. (2006) (MacMillan and Shary, 2009). 182

183 The fundamental geomorphometric variables constitute basic building blocks for 184 deriving combined indices such as the topographic wetness index (TWI) or the 185 topographic position index (TPI) and for performing further and more sophisticated land 186 surface analyses and classifications (Evans and Minar, 2011). The use of 187 geomorphometric field variables to identify landform classes and features dates back 188 over four decades (Wilson, 2012). Over the last twenty years, several methods have 189 been developed to automate the extraction of land surface features from DEMs (e.g. 190 Graff and Usery, 1993; Miliaresis and Argialas, 1999; Dymond et al., 1995; Wood, 191 1996; Schmidt and Hewitt, 2004; Dragut and Blaschke, 2006). Several widely applied 192 approaches to automated classification of land surface elements are based on

193 consideration of local surface shape as measured by slope gradient and signs or values 194 of curvatures (MacMillan and Shary, 2009). The capabilities of this approach are best 195 illustrated by Wood (1996) who used slope, cross-sectional and minimum and 196 maximum profile curvatures calculated within the analysis window to define six 197 categories of surface-specific elements: peaks, ridges, passes, channels, pits, and plains 198 (Hengl and Evans, 2009).

199 Blaszczynski (1997) proposed an alternative method for curvature calculation 200 and determining whether cells were on convex or concave parts of the land surface. His approach to curvature analysis was used for classifying a continuous landscape surface 201 202 represented by a DEM into a series of discrete areas representing geomorphometric 203 surface-specific elements or features such as crests, troughs, side slopes, open and 204 enclosed basins, inclined and horizontal flats. Blaszczynski (1997) showed how 205 convexity and concavity can be identified by modifying the calculation of the average 206 percent slope gradient for a centre cell within a kernel. The calculated value of this curvature measure or 'signed average local relief', $R^{s}_{0,0}$, assigned to the cell in the centre 207 208 of a $n \times n$ kernel (where *n* is odd and $n \ge 3$) is:

209
$$R_{0,0}^{s} = 100 \frac{1}{rN} \sum_{i} \sum_{j} \frac{(z_{0,0} - z_{i,j})}{\sqrt{(x_{0} - x_{j})^{2} - (y_{0} - y_{i})^{2}}}$$
(15)

where $z_{0,0}$ is the elevation of the cell at the kernel centre (x_0, y_0) , the $z_{i,j}$ are the elevation values in the surrounding cells within the kernel at positions i,j = -(n-1)/2, ..., (n-1)/2with respect to the kernel centre, *r* is DEM grid spatial resolution (i.e. cell size), *N* is the number of surrounding cells within the kernel, and x,y are the spatial coordinates of the cells.

Yokoyama et al. (2002) proposed a geomorphometric variable termed 'openness' which is related to local curvature. Openness is directly related to land surface line-of-sight and thus is derived taking the maximum angle of vision from a point on the land surface within a given maximum radial distance. The calculated value at each cell of a DEM is:

220
$$\phi_{0,0} = 90 - \beta_{0,0} = 90 - \frac{1}{N_D} \sum_{d=1,2,\dots,L}^{N_D} \left\{ \arctan\left(\frac{z_{0,0} - z_{i,j}^d}{r\sqrt{(x_0 - x_j^d)^2 - (y_0 - y_i^d)^2}}\right) \right\}$$
(16)

221

where $\phi_{0,0}$, $\beta_{0,0}$ and $z_{0,0}$ are the (positive) openness, the maximum elevation angle and the elevation of the cell at the kernel centre (x₀, y₀), respectively, the $z^{d}_{i,j}$ are the elevation values in the cells located on a profile along an azimuth $d \in D = \{0^\circ, 45^\circ, 90^\circ, 135^\circ, 180^\circ, 225^\circ, 270^\circ, 315^\circ\}$ and within the kernel at positions i,j = -(n-1)/2, ..., (n-1)/2with respect to the kernel centre, N_D is the number of azimuths or compass directions (8 in the original algorithm), L is kernel (half) size, r is DEM grid spatial resolution, and x,y are the spatial coordinates of the cells. Similarly, a negative openness was defined using the minimum elevation angle.

- 230
- 231

2.1 Software for digital land surface analysis

232

233 The analysis of DEMs was traditionally conducted using stand-alone programs 234 developed for scientific use such as MicroDEM (Guth et al., 1987), TAPES-G (Gallant 235 and Wilson, 1996), TARDEM (Tarboton, 1997), and TauDEM (Tarboton and Ames, 236 2001). Whilst some stand-alone programs made links with GIS to take advantage of 237 their superior facilities for viewing, panning and management of DEMs, others relied on 238 image processing software. LandSerf (Wood, 1998, 2009b) for example was a 239 comprehensive, multi-platform suite of programs for multi-scale land surface analysis 240 and visualisation, aimed at researchers and written in Java. It computed a variety of land 241 surface variables from a DEM (slope, aspect, profile, plan, longitudinal and cross-242 sectional curvature), enabling a variety of land surface features (channels, ridges, peaks, 243 passes, pits and plains) to be classified.

244 While specialised software such as Landserf will continue to be used by 245 researchers where comprehensiveness of functionality is paramount, we identify a 246 broader range of application areas in which users value the convenience of carrying out 247 preparatory data processing with the same software they will use for further analysis 248 and presentation of results. GIS software is now so widely adopted by many scientific 249 professionals, for whom land surface analysis is just one necessary step towards a final 250 result and the overhead of investing time to learn specialised software for 251 geomorphometry may not be justified. These users create a demand for more 252 comprehensive and accessible functionality for land surface analysis in mainstream GIS 253 software. Gessler et al. (2009) have recognised the scarcity of user-friendly and 254 computationally efficient GIS tools as the most serious bottleneck in semi-automated geomorphometric mapping. 255

256 Most GIS now include functions for computing the most common 257 geomorphometric operations on DEMs such as the maximum down-slope gradient,

8

258 slope aspect, convexity, and direction of down-gradient flow paths (Gallant and Wilson, 259 2000). However, in most GIS, only the simpler algorithms are often used and 260 implemented using only a 3×3 kernel. With many users typically working with only one 261 DEM product, land surface variables computed using windows of such limited 262 dimension will detect only variations in topography at one scale determined by the 263 DEM resolution (Gallant and Wilson, 2000). Dragut and Eisank (2011) have argued that 264 the capability for multi-scale extraction of landscape features is still lacking and may be 265 hindering studies of how landform elements are extracted and recognised from 266 continuous fields of elevation data.

267 There have been some previous attempts of providing ArcGIS toolboxes for 268 geomorphometric analysis. Currently, to the best knowledge of the authors, two 269 toolboxes are publicly available: the ArcGIS Geomorphometry Toolbox (Reuter, 2009) 270 and the ArcGIS Geomorphometry and Gradient Metrics Toolbox (Evans et al., 2014). 271 The ArcGIS Geomorphometry Toolbox is a comprehensive ArcGIS toolbox containing 272 a large number of geomorphometric algorithms. Current toolbox version 1.0.6 is only 273 compatible with ArcGIS version 10.0 (ArcGIS version to be retired in 2015; Esri, 274 2015). The toolbox is provided as a commercial software program (it is almost free for 275 pure research) (Reuter, 2009). The toolbox includes a large number of geomorphometric 276 functions. The geomorphometric functions provided are grouped under menus labelled: "Landforms", and "Terrain parameters". "Landforms" menu includes eleven algorithms 277 278 for land surface classification (Pennock et al., 1994; MacMillan and Pettapiece, 1997; 279 MacMillan et al., 2000; Meybeck et al., 2001; Park et al., 2001; Weiss, 2001; Reuter, 280 2004; Dobos et al., 2005; Iwahashi and Pike, 2007) and derivation of some combined 281 indices (Bolstad's et al. (1998) Landform Index, Weiss' (2001) TPI). "Terrain 282 parameters" menu includes several algorithms for the calculation of fundamental 283 geomorphometric variables such as slope, aspect, curvature (profile, plan, tangential), 284 stream order, and watershed area (MacMillan et al., 2000; Reuter, 2004; Esri, 2010), 285 and of alternative variables such as openness (Yokoyama et al., 2002), and some 286 combined indices such as TWI, TPI, mass balance index (Moller et al., 2008), and 287 elevation residuals (Wilson and Gallant, 2000). Fundamental field variables (e.g. slope 288 gradient, aspect, curvature) are calculated through a fixed neighbourhood operation by 289 moving a 3×3 window across a DEM utilising ArcGIS functions (Esri, 2010). Curvature 290 can also be calculated using two alternative formulae (not documented or referenced). 291 Some combined indices (e.g. TWI, TPI, elevation residuals) can be calculated using a

range of windows extents by utilising ArcGIS focal statistics functions (e.g. focal mean). The openness variable, requiring direct access to neighbour elevation values within the analysis window, can be obtained up to a window extent of 9×9 (Reuter, 2009). A basic description or reference (embedded in source code) of the algorithms provided is included. Separated documentation or toolbox help pages are not provided.

297 The ArcGIS Geomorphometry and Gradient Metrics Toolbox (Evans et al., 298 2014) is an ArcGIS toolbox containing various utilities and geomorphometric 299 algorithms. Current toolbox version 2.0 is compatible with ArcGIS versions 10.x and is 300 provided as open source (freeware). The toolbox is devised to support ecological 301 modelling and hence functions provided are grouped under menus labelled 302 "Directionality", "Statistics", "Texture and Configuration", and "Temperature and 303 Moisture". The first two menus include general purpose utilities and statistical functions 304 (e.g. correlation). "Texture and Configuration" menu includes functions for the 305 calculation of indices such as dissection (Evans, 1972), hierarchical slope position 306 (Murphy et al., 2010), surface curvature index (Bolstad and Lillesand, 1992), roughness 307 (i.e. local elevation variance), slope position (Gallant and Wilson, 2000), and surface 308 relief ratio (Pike, 1971). "Temperature and Moisture" menu include functions for the 309 calculation of indices such as compound topographic index (Moore et al., 1993), heat 310 load index (McCune and Keon, 2002), integrated moisture index (Iverson et al., 1997), 311 and site exposure index (Balice et al., 2000). Indices are calculated combining standard 312 ArcGIS functions (working through a fixed 3×3 window) such as slope gradient, aspect, 313 and curvature (Esri, 2014) with ArcGIS focal statistics functions operating at a range of 314 windows extents. A basic description (embedded in source code) of the tools is 315 included. A "Read Me.pdf" file including a description and references of the algorithms 316 is provided. Toolbox help pages are not provided. Both toolboxes above require the 317 ArcGIS Spatial Analyst extension to operate.

318 This brief review has considered a variety of software packages for conducting 319 geomorphometry and identified various user requirements that are not fully met by 320 existing software. A more comprehensive review of software for geomorphometry by 321 Wood (2009a) in which eight packages (ArcGIS Workstation, GRASS, ILWIS, 322 LandSerf, MicroDEM, RiverTools, SAGA and TAS) were assessed for their 323 geomorphometric capabilities concluded there is considerable scope for software that 324 fills the gap that still exists between comprehensive, specialist tools and the limited 325 functionality presently implemented by major GIS vendors.

Using Wood's (2009a) triangular diagram of the software landscape for geomorphometry, we are proposing a solution that fills the gap between the standalone tools and a standard install of a mainstream GIS (Fig. 1). The tool takes advantage of the power of the GIS to handle the large DEM sizes, whilst retaining ease of navigation through its custom user interface to a more sophisticated set of methods, including support for multi-scale analysis of DEMs.



Geomorphometry

Hydrology

Fig. 1. Positioning of the new tool within the existing software landscape for geomorphometry (modifiedfrom Wood, 2009a).

336

333

332

The next section describes the development environment and the functions implemented to create a more comprehensive and accessible tool set for conducting geomorphometry efficiently and productively in ArcGIS.

340

341 3. ArcGeomorphometry toolbox for ArcGIS

342

```
343 3.1 The ArcGIS development environment
```

344

According to recent reports, the Esri ArcGIS software is the most commonly used GIS worldwide (GITA, 2008; Daratech Inc., 2011). Esri's flagship product, ArcGIS for Desktop, is widely used in education, industry and several scientific research fields, especially in the geosciences. In many of these fields, there is a need to conduct geomorphometric analysis. ArcGIS for Desktop includes the Spatial Analyst extension that can be used for this purpose. While Spatial Analyst provides efficient methods for constructing DEMs from various source data formats, its explicit functions for geomorphometric analysis are limited and implemented using a fixed 3×3 kernel (e.g. slope gradient, aspect, and curvature based on the method described by Zevenbergen and Thorne, 1987).

355 ArcGIS supports several popular programming and scripting languages, 356 although Esri has officially embraced Python as the recommended programming 357 language for working with ArcGIS (Zandbergen, 2012; Esri, 2014). User created 358 Python scripts can be integrated into ArcGIS as script tools, which work just like 359 standard ArcGIS processing (geoprocessing) tools and can be accessed from the ArcGIS 360 user interface. Python Toolboxes are geoprocessing toolboxes created entirely in Python and the tools contained within, look, act, and work just like the Toolboxes and tools 361 362 created in any other way. This allows easy sharing of tools among users and researchers 363 and facilitates amendments and addition of new tools to the toolbox.

ArcGIS geoprocessing functionality is accessible through Python using ArcPy library. Of particular importance to this study, Numerical Python (NumPy) is a numerical library for scientific computing, including support for powerful Ndimensional array objects.

368 The ability to construct more complex functionality from the basic language 369 syntax, the widespread availability of the scripting language and the many types of 370 DEM data already available in ArcGIS raster data format led to the decision to develop 371 the extended functionality for geomorphometric analysis using the ArcGIS Python 372 environment. This new functionality was then made accessible to the user using a 373 Python Toolbox, which can be installed, shared and modified. By following the 374 conventions for Python Toolboxes design (Esri, 2014), the code for the GUI integrates 375 with the standard ArcToolbox with the result that, once loaded, ArcGeomorphometry 376 menus, dialogues, help pages, etc., appear seamlessly incorporated within ArcGIS.

377

378 *3.2* . *The ArcGeomorphometry tools*

379

The ArcGeomorphometry tools allow landscapes stored as DEMs in any ArcGIS
raster format to be analysed and classified and land surface features to be identified at

382 different spatial scales. Standard menus and dialogue boxes guide the user through a 383 series of steps required to produce a geomorphometric analysis or land surface 384 classification, without having to program these procedures (Rigol-Sanchez and Stuart, 385 2005). Users can conduct a series of classifications of a DEM into different land surface 386 features (i.e. surface-specific elements) and by quickly reviewing the results, can 387 progressively refine the classifications. ArcGeomorphometry focus on field local 388 variables and implements many of the commonly needed functions for 389 geomorphometric analysis of DEMs (Table 1). It currently provides functions for true 390 multi-scale land surface analysis and classification based on the methods proposed by 391 Evans (1972, 1979, 1980) and Wood (1996); Shary (1995) and Shary et al., (2002); 392 Blaszczynski (1997); and Yokoyama et al. (2002). These functions are grouped by 393 method under menus labelled "Evans-Wood Method"; "Shary Method"; "Average 394 Relief"; and "Openness" respectively. The algorithms provided under Evans-Wood and 395 Shary menus make use of Numpy functions to fit a bivariate quadratic polynomial (for 396 each DEM cell) to elevation values contained within the given window/kernel size 397 extent by least squares. Polynomial parameters are then used to obtain 398 geomorphometric variables. The algorithms under Average Relief and Openness make 399 use of Numpy array indexing capabilities.

- 400
- 401 **Table 1**
- 402 Functions of ArcGeomorphometry.

Function name	Description			
Average Relief functions				
average slope	Calculate average slope percent			
classified average relief	Reclassify signed average local relief grid using user defined slope and signed			
	average local relief cut-offs			
signed average relief	Calculate signed average local relief			
Openness functions				
negative openness	Calculate 8-direction average minimum elevation angle below surface			
positive openness	Calculate 8-direction average maximum elevation angle above surface			
Evans-Wood Method functions				
aspect	Compute slope orientation or aspect			
elevationSmoothed	Return elevation smoothed by quadratic function			
crossCurvature	Compute cross-sectional curvature			
feature	Classify DEM into surface-specific elements (pit, peak, ridge, channel, pass, plane)			
	using user-defined slope and curvature thresholds			
longCurvature	Compute longitudinal curvature			
maxProfCurvature	Compute maximum profile curvature			

minProfCurvature	Compute minimum profile curvature			
modified Evans-Young	Modified Evans-Young algorithm (pre-filtering)			
planCurvature	Compute plan curvature			
profileCurvature	Compute profile curvature			
Slope	Compute slope steepness			
Shary Method functions				
aspect	Compute slope orientation or aspect			
crossCurvature	Compute cross-profile curvature			
longCurvature	Compute longitudinal curvature			
maxProfCurvature	Compute maximum profile curvature			
meanCurvature	Compute mean curvature			
minProfCurvature	Compute minimum profile curvature			
planCurvature	Compute plan curvature			
profileCurvature	Compute profile curvature			
rotor	Compute rotor			
tangentialCurvature	Compute tangential curvature			
totalCurvature	Compute total curvature			
totalGaussianCurvature	Compute total Gaussian curvature			
totalRingCurvature	Compute total ring curvature			
slope	Compute slope steepness			
unsphericity	Compute unsphericity			

403

404 The toolbox runs on any computer running ArcGIS for Desktop 10.1 SP1 or 405 higher. It consists of a Python script that realise the analysis routines, user menu, 406 dialogue boxes and basic help. Additional help pages are stored as xml files. Installed 407 tools can also be run in a standalone mode by calling them from a Python window or 408 ModelBuilder, but are intended primarily to be operated through a graphical menu. Tools use linear map units, such as feet or meters, and consequently, it is assumed that 409 410 input DEM has a projected coordinate system. The maximum size of the input raster 411 DEM, i.e. maximum number of cells, is limited by available RAM on computer up to a 412 maximum RAM allocation per Python 32-bit process imposed by the operating system 413 (2GB). In practice, DEMs of 1.0E+08 cells can be processed in a standard personal 414 computer (4GB RAM, Core i3-2100 processor running at 3.10GHz) in periods from few 415 minutes to several hours depending on the function and kernel size selected (Fig. 2). As 416 indicated above, DEM analyses involving direct operations on neighbour cells values 417 such as cell sum, subtraction or multiplication can be efficiently performed in Numpy in 418 one step using array indexing. This is the case for functions under Average Relief and 419 Openness. DEM analyses based on more complex operations that require simultaneous 420 access to all neighbouring cell values within the kernel such as function fitting

421 procedures (e.g. Evans-Wood Method or Shary Method) have to be undertaken in two 422 steps (neighbour data load using array indexing; and kernel operation, e.g. function 423 fitting by least squares, solving a system of linear scalar equations for each DEM cell). 424 Typically, Numpy array views are used to access neighbouring cell values.

425



429

430 Fig. 2. Timings for some ArcGeomorphometry functions: (a) Computed using an input DEM of 6,000 431 columns by 6,000 rows (3.60E+07 cells) and increasing kernel sizes (3×3 , 5×5 , 11×11 , 21×21 , 41×41 , 432 91×91). (b) Computed using a kernel size of 21×21 and increasing input DEM sizes (276 columns × 173 433 rows, 1,702×903, 6,000×6,000, 10,880 x 10,880). Processing was performed using a standard personal 434 computer (4GB RAM, Core i3-2100 processor running at 3.10GHz).

435

436 The ArcGeomorphometry toolbox allows the user to perform multi-scale 437 geomorphometric analyses. Hence, in all cases once the input DEM is selected, the size of the processing kernel (or analysis window) for land surface analysis is selected by
typing in the desired square dimension (a circle diameter for openness). Any positive
odd kernel size is allowed, so that maximum size of analysis window is limited only by
the spatial dimensions of input DEM or available system resources.

The results of each land surface classification are graphically displayed if the tool is executed from within ArcGIS ArcMap or ArcScene applications. Thus the user can readily display further ArcGIS grids (such as gridded land cover data or a previous geomorphological mapping of an area) and overlay vector data sets such as contour lines on top of output grids (Fig. 3). Adding this contextual information facilitates an immediate visual assessment of results, which can highlight incongruities or give credence to elements of a landscape classification.

- 449
- 450
- 451 452

4. The operation of ArcGeomorphometry illustrated using an example DEM

453 Fig. 3(a) shows a sample DEM used to illustrate the operation of the 454 ArcGeomorphometry Toolbox. The data are included on the ArcGIS for Desktop 455 installation media. The DEM covers an area of 23.64km by 23.04km of the town of 456 Stowe, Vermont, USA, with a cell size of 30m by 30m (788 columns \times 768 rows). The 457 topography of the area corresponds to a moderately rugged mountainous terrain. The 458 maximum elevation value (1,319m) is located close to the upper-left corner of the area 459 (Green Mountains) and minimum value (134m) is located close to the lower border at 460 the bottom of the main valley (Little River).

461 Fig. 3 illustrates the processing of the DEM with ArcGeomorphometry Average 462 *Relief* tools using a range of kernel sizes. Once the input DEM is selected, the size of 463 the analysis window for land surface analysis is selected by typing in the desired square 464 dimension. Any positive odd kernel size is allowed, although 81×81 cells has been 465 found practically to be sufficient to extract many amplitudes of land surface features 466 from DEMs with ground resolutions in the range from 10-200m. 81 cells equate to a 467 2.4km \times 2.4km search window for a 30m DEM and for this terrain produce a very 468 smoothed output surface. Figs. 3(b) to (f) are graphical displays generated within 469 ArcGIS ArcMap. The spatial pattern of land surface features identified by the methods 470 is generally consistent with what would be interpreted from topographic mapping of the 471 area. When the classification is repeated using larger kernel sizes, the number and the

16

- 472 complexity of the land surface features identified is reduced and greater smoothing of
- the land surface occurs.
- 474



Fig. 3. Test DEM (a) and results of classification using ArcGeomorphometry Average Relief tool computed using a: (b) 3×3 kernel (90m×90m). (c) 11×11 kernel (330m×330m). (d) 21×21 kernel (630m×630m). (e) 41×41 kernel (1,230m×1,230m). (f) 81×81 kernel (2,430m×2,430m). Maps of classifications are overlain with a vector layer of contour lines at 50m interval. Note that the extent of the area that can be classified by the processing without edge effects is reduced as the kernel size increases.

The sequence of classified grids in Fig. 3(b) - (f) illustrate that, as expected, land surface features extracted by using large kernel sizes have comparably larger spatial dimensions than those identified by small kernel sizes. Land surface features classified by large kernels reflect the variations of topography at a broader scale, corresponding roughly to features whose dimensions are similar to the length of entire hillsides. It can also be seen that the classification using the "standard" 3×3 kernel produces an image with less coherence and a higher local variance from which it is more difficult to interpret land surface features. In this example using a 30m resolution DEM, the range of kernel sizes from 3×3 to 81×81 covers a range of landscape features from the microscale (0-30m) to the meso-scale (30-2,430m) (Dikau, 1989). Indicative timings for performing the above classifications are shown in Fig. 2. Fig. 4 shows four other results as perspective views of the processing of the sample DEM using different functions of the toolbox generated within ArcGIS ArcScene.

494



Fig. 4. 2.5D perspective views illustrating processing of test DEM using ArcGeomorphometry tools. (a) *Evans-Wood Method "feature"* classification calculated using a 11×11 kernel (330m×330m). (b) Shary *Method "unsphericity"* variable calculated using a 11×11 kernel. (c) Shary Method "plan curvature"
variable calculated using a 21×21 kernel (630m×630m). (d) Openness "positive openness" variable
calculated using a 31×31 kernel (930m×930m). Vertical scale is exaggerated by a factor of 1.5.

500 501

5. Conclusions and planned enhancements

502

503 The ArcGeomorphometry toolbox provides a means for conducting exploratory, 504 iterative and multi-scale land surface analysis with DEMs in the ArcGIS environment. 505 Operating through the graphical user interface, users can easily and flexibly select the 506 desired function from a comprehensive selection and vary the size of the kernel to 507 identify features from the land surface model at different scales. Parameter values can 508 be adjusted flexibly to enable analysis and classification of different land surface 509 elements on the basis of both curvature and degree of slope of the surface at various 510 scales. Because the results from each iteration are immediately available for detailed 511 inspection using the sophisticated visualisation techniques of GIS, users may browse, 512 zoom, query, reclassify and overlay additional data sets to determine when an 513 acceptable classification has been found. The results are produced in a format that can 514 be immediately interpreted, integrated with additional data, or analysed further using 515 any available ArcGIS functions. The toolbox are highly portable and functions can also 516 be used within ArcGIS ModelBuilder or other scripts, in both interactive and batch 517 processing modes.

518 If a reference data set exists, for example if a field survey has previously 519 produced geomorphological mapping for a given locality, the ArcGeomorphometry 520 tools can be used to determine kernel and threshold parameter values that classify a 521 DEM for this area into land surface units that conform with the mapping. Once these 522 parameters have been established, it may be possible to apply similar parameter settings 523 to recognise similar landscape features from a DEM of the same specification but 524 covering a more extensive area for which geomorphological mapping has not been 525 previously produced.

526 A few limitations apply to processing DEMs with the ArcGeomorphometry 527 Toolbox. While the time for the per-pixel algorithms to process a gridded DEM 528 increases quadratically as the DEM extent is increased, for neighbourhood algorithms 529 the time increases at faster rates as the size of the kernel is increased, since many more 530 cells have to be processed in the input layer to create a single value in the output grid. 531 The present tests of ArcGeomorphometry suggested that quite large kernel sizes (e.g. 532 81×81) may sometimes be required to extract some larger amplitude land surface 533 features. While there is no limitation in the software upon the size of kernel that can be 534 used, working with kernels much larger than those normally available in standard 535 systems leads to 'non-interactive' processing unless the DEM extent is quite small (Fig. 536 2). The availability of higher resolution DEM products, such as 10m products derived 537 from InSAR data or submetric LIDAR DEM data, while potentially allowing much 538 finer surface detail to be revealed, would lead to much longer processing times if such 539 high spatial resolution data sets were used for extracting features of the same 540 dimensionality and over similar extents as those in this illustration.

541 In the present version of ArcGeomorphometry, if any cell in the processing 542 kernel has a null value, then the output for the cell at the centre of that kernel will be 543 null. As a consequence, each edge of the classified DEM created by the processing will be reduced by one-half of the kernel size, leading to the overall dimensions of the outputgrid being reduced by the number of (rows=columns=k) in the kernel.

Future improvements envisaged for the toolbox include: (a) The storage of DEMs as binary files on disk to circumvent the limit of input DEM size imposed by the operating system. (b) To allow the user to constrain analyses to specific quadrants of the analysis kernel (e.g. where the resultant cell value is determined only by cells in the north-east or south-west quadrant of the kernel). This may be a simple way to explore directional dependence of some land surface features or the influence of particular orientations upon features on land surface geomorphometry.

553

554 Acknowledgments

555

556 Comments from two anonymous reviewers greatly helped in improving the 557 manuscript. This work benefited from a "British Council – la Caixa" fellowship and an 558 EU Marie Curie research grant to the Department of Geography, School of GeoSciences 559 at the University of Edinburgh, UK. This study was supported by Research Group 560 RNM-189 and the project CADESEM P11-RNM-8115 of the of the Junta de Andalucia 561 Regional Government, Spain.

562

563 **References**

- 564
- Balice, R.G., Miller, J.D., Oswald, B.P., Edminister, C., Yool, S.R., 2000. Forest
 Surveys and Wildfire Assessment in the Los Alamos; 1998–1999. Los Alamos, NM,
 USA: Los Alamos National Laboratory. LA-13714-MS. 12 pp.
- 568 Band, L.E., 1986. Topographic partition of watersheds with digital elevation models.
- 569 Water Resources Research 22 (1), 15-24.
- Blaszczynski, J.S., 1997. Landform characterization with geographic information
 systems. Photogrammetric Engineering & Remote Sensing 63 (2), 183-191.
- 572 Bolstad, P.V., Lillesand, T.M. 1992. Improved classification of forest vegetation in
- 573 northern Wisconsin through a rule-based combination of soils, terrain, and Landsat
 574 TM data. Forest Science 38(1), 5–20.
- Bolstad, P.V., Swank, W., Vose J., 1998. Predicting Southern Appalachian overstory
 vegetation with digital terrain data. Landscape Ecology 13, 271–283.

- 577 Daratech Inc., 2011. GIS/Geospatial Markets and Opportunities 2011. Daratech Inc.,
 578 URL: http://www.daratech.com (accessed 24 June 2014).
- 579 Dehn, M., Gartner, H., Dikau, R., 2001. Principles of semantic modelling of landform
 580 structures. Computers & Geosciences 27 (8), 1005–1010.
- 581 Dikau, R., 1989. The application of a digital relief model to landform analysis in
 582 geomorphology. In: J. Raper (Ed.), Three dimensional applications of GIS. Taylor
 583 and Francis, London, UK, pp. 55–77.
- Dobos, E., Daroussin, J., Montanarella, L., 2005. An SRTM-based procedure to
 delineate SOTER Terrain Units on 1:1 and 1:5 million scales, EUR 21571 EN.
 Office for Official Publications of the European Communities, Luxembourg, 55 pp.
- 587 Dragut, L., Blaschke, T., 2006. Automated classification of landform elements using
 588 object-based image analysis. Geomorphology 81, 330–344.
- 589 Dragut, L., Eisank, C., 2011. Object representations at multiple scales from digital
 590 elevation models. Geomorphology 129, 183–189.
- 591 Dymond, J.R., Derose, R.C., Harmsworth, G.R., 1995. Automated mapping of land
 592 components from digital elevation data. Earth Surface Processes and Landforms 20,
 593 131-137.
- Esri, 2010. ArcGIS 10.0 Help Library. Environmental Systems Research Institute Inc.,
 Esri Press, Redlands, California, USA. URL:
 http://help.arcgis.com/en/arcgisdesktop/10.0/help/index.html (accessed 12 April
 2015).
- 598 Esri, 2014. ArcGIS Help Library. Environmental Systems Research Institute Inc., Esri
 599 Press, Redlands, California, USA. URL: http://resources.arcgis.com/en/help
 600 (accessed 11 October 2014).
- 601 Esri, 2015. ArcGIS for Desktop Product Life Cycle Support Status. URL:
 602 http://downloads.esri.com/support/product%20life%20cycle/arcgis_desktop/ArcGIS
 603 PLC.pdf (accessed 24 April 2015).
- Evans, I.S., 1972. General geomorphometry, derivatives of altitude and descriptive
 statistics. In: Chorley, R.J. (Ed.), Spatial Analysis in Geomorphology. Methuen &
 Co. Ltd., London, UK, pp. 17–90.
- Evans, I.S., 1979. An integrated system of terrain analysis and slope mapping. Final
 report on grant DA-ERO-591-73-G0040, University of Durham, England.
- 609 Evans, I.S., 1980. An integrated system of terrain analysis for slope mapping.
- 610 Zeitschrift fur Geomorphologie 36, 274–295.

- 611 Evans, I.S., 2012. Geomorphometry and landform mapping: What is a landform?612 Geomorphology 137, 94–106.
- 613 Evans, I.S., Minar, J., 2011. A classification of geomorphometric variables. In: Hengl,
- T., Evans, I.S., Wilson, J.P., Gould, M. (Eds.), Proceedings of the Geomorphometry
- 615 2011 Conference, Redlands, California, USA. 105–108, URL:
 616 http://geomorphometry.org/EvansMinar2011 (accessed 11 May 2014).
- Evans, J.S., Oakleaf, J., Cushman, S.A., Theobald, D., 2014. An ArcGIS Toolbox for
 Surface Gradient and Geomorphometric Modeling, version 2.0-0. URL:
 http://evansmurphy.wix.com/evansspatial (accessed 2 December 2014).
- Florinsky, I.V., 1998. Accuracy of local topographic variables derived from digital
 elevation models. International Journal of Geographical Information Science 12, 47–
 622 62.
- Gallant, J.C., Wilson, J.P., 1996. TAPES-G: a grid-based terrain analysis program for
 the environmental sciences. Computers & Geosciences 22 (7), 713–722.
- Gallant, J.C., Wilson, J.P., 2000. Primary topographic attributes. In: Wilson, J.P.,
 Gallant, J.C., (Eds.), Terrain Analysis: Principles and Applications. John Wiley &
 Sons, New York, USA, pp. 51–85.
- Gessler, P., Pike, R., MacMillan, R.A., Hengl, T., Reuter, H.I., 2009. The future of
 geomorphometry. In: Hengl, T. and Reuter, H.I. (Eds.), Geomorphometry: Concepts,
 Software, Applications. Developments in Soil Science, Vol. 33, Elsevier,
 Amsterdam, Netherlands, pp. 637–652.
- GITA, 2008. GITA Geospatial Technology Report 2007/2008. Geospatial Information
 and Technology Association (GITA).). URL: http://www.gita.org (accessed 24 June
 2014).
- Graff, L.H., Usery, E.L., 1993. Automated Classification of Basic-Level Terrain
 Features in Digital Elevation Models. Photogrammetric Engineering and Remote
 Sensing 59 (9), 1409–1417.
- Guth, P.L., Ressler, E. K., Bacastow, T. S., 1987. Microcomputer program for
 manipulating large digital terrain models. Computers & Geosciences 13 (3), 209–
 213.
- Hengl, T., Evans, I.S., 2009. Mathematical and digital models of the land surface. In:
 Hengl, T. and Reuter, H.I. (Eds.), Geomorphometry: Concepts, Software,
 Applications. Developments in Soil Science, Vol. 33, Elsevier, Amsterdam,
 Netherlands, pp. 31–63.

22

- Iverson, L.R., Dale, M.E., Scott, C.T., Prasad, A., 1997. A GIS-derived integrated
 moisture index to predict forest composition and productivity of Ohio forests
 (U.S.A.). Landscape Ecology 12, 331–48.
- Iwahashi, J., Pike, R.J., 2007. Automated classifications of topography from DEMs by
 an unsupervised nested-means algorithm and a three-part geometric signature.
 Geomorphology 86, 409–440.
- Jenson, S.K., Domingue, J.O., 1988. Extracting topographic structure from DEM for
 GIS analysis. Photogrammetric Engineering and Remote Sensing 54 (11), 1593–
 1600.
- Lueder, D. R., 1959. Aerial Photographic Interpretation Principles and Application.
 McGraw-Hill, New York, USA, 452 pp.
- MacMillan, R.A., Pettapiece, W.W., 1997. Soil Landscape Models: automated
 landscape characterization and generation of soil–landscape models. Research
 Report No. 1. Agriculture and Agri-Food Canada, Research Branch, Lethbridge,
 Canada, 75 pp.
- MacMillan, R.A., Pettapiece, W.W., Nolan, S.C., Goddard, T.W., 2000. A generic
 procedure for automatically segmenting landforms into landform elements using
 DEMs, heuristic rules and fuzzy logic. Fuzzy Sets and Systems 113, 81–109.
- MacMillan, R.A., Shary, P.A., 2009. Landforms and landform elements in
 geomorphometry. In: Hengl, T. and Reuter, H.I. (Eds.), Geomorphometry: Concepts,
 Software, Applications. Developments in Soil Science, Vol. 33, Elsevier,
 Amsterdam, Netherlands, pp. 227–254.
- McCune, B., Keon, D., 2002. Equations for potential annual direct incident radiation
 and heat load index. Journal of Vegetation Science 13, 603–606.
- Meybeck, M., Green, P., Vorosmarty, C.J., 2001. A new typology for mountains and
 other relief classes: An application to global continental water resources and
 population distribution. Mountain Research and Development 21, 34–45.
- Miliaresis, G.Ch., Argialas, D.P., 1999. Segmentation of physiographic features from
 the global digital elevation model/G-GTOPO30. Computers & Geosciences 25 (7),
 715–728.
- Moller, M., Volk, M., Friedrich, K., Lymburne, L., 2008. Placing soil-genesis and
 transport processes into a landscape context: A multiscale terrain-analysis approach.
- 577 Journal of Plant Nutrition and Soil Science 171, 419–430.

- Moore, I.D., Lewis, A., Gallant, J.C., 1993. Terrain attributes: estimation methods and
 scale effects. In: Jakeman, A.J., Beck, M.B., McAleer, M.J. (Eds.), Modeling
 Change in Environmental Systems. John Wiley and Sons, New York, pp. 189–214.
- Murphy, M.A., Evans, J.S., Storfer, A., 2010. Quantifying Bufo boreas connectivity in
 Yellowstone National Park with landscape genetics. Ecology 91, 252–261.
- Park, S.J., McSweeney, K., Lowery, B., 2001. Identification of the spatial distribution
 of soils using a process-based terrain characterisation. Geoderma 103, 249–272.
- Pennock, D.J., Anderson, D.W., de Jong, E., 1994. Landscape-scale changes in indicators of soil quality due to cultivation in Saskatchewan, Canada. Geoderma 64, 1–19.
- Pennock, D.J, Zebarth, B.J., Jong, E., 1987. Landform classification and soil
 distribution in hummocky terrain, Sasketchewan, Canada. Geoderma 40, 297–315.
- 690 Pike, R.J., 1995. Geomorphometry process, practice and prospects. Zeitshcrift für
 691 Geomorphologie Suppl. 101, 221–238.
- 692 Pike, R.J., 2000. Geomorphometry diversity in quantitative surface analysis. Progress
 693 in Physical Geography 24 (1), 1–20.
- 694 Pike, R.J., Wilson, S.E., 1971. Elevation relief ratio, hypsometric integral, and
 695 geomorphic area altitude analysis. Bulletin of the Geological Society of America 82,
 696 1079–1084.
- Pike, R.J., Evans, I.S., Hengl, T., 2009. Geomorphometry: a brief guide. In: Hengl, T.
 and Reuter, H.I. (Eds.), Geomorphometry: Concepts, Software, Applications.
 Developments in Soil Science, Vol. 33, Elsevier, Amsterdam, Netherlands, pp. 3–
 30.
- Reuter, H.I., 2004. Spatial Crop and Soil Landscape Processes under Special
 Consideration of Relief Information in a Loess Landscape. Der Andere Verlag,
 Osnabruck, Germany, 251 pp.
- Reuter, H.I., 2009. ArcGIS Geomorphometry Toolbox. Gisxperts gbr, Germany. URL:
 http://www.ai-relief.org/ (accessed 22 April 2015).
- Rigol-Sanchez, J.P., Stuart, N., 2005. Multi-scale geomorphometric characterization of
 DEMs in the ArcGIS environment. Sixth International Conference on
 Geomorphology. Zaragoza, Spain.
- Schmidt, J., Evans, I. S., Brinkmann, J., 2003. Comparison of polynomial models for
 land surface curvature calculation. International Journal of Geographical
 Information Science 17 (8), 797–814.

- Schmidt, J., Hewitt, A., 2004. Fuzzy land element classification from DTMs based on
 geometry and terrain position. Geoderma 121, 243–256.
- Shary, P.A., 1995. Land surface in gravity points classification by complete system of
 curvatures. Mathematical Geology 27 (3), 373–390.
- Shary P.A., Sharaya L.S., Mitusov A.V., 2002. Fundamental quantitative methods of
 land surface analysis. Geoderma 107, 1-32.
- Smith, M.P., Zhu, A.X., Burt, J.E., Stiles, C., 2006. The effects of DEM resolution and
 neighborhood size on digital soil survey. Geoderma 137 (1–2), 58–68.
- Tarboton, D.G., 1997. A new method for the determination of flow directions and
 contributing areas in grid Digital Elevation Models. Water Resources Research 33
 (2), 309–319.
- Tarboton, D.G., Ames, D. P., 2001. Advances in the mapping of flow networks from
 digital elevation data. ASCE, World Water and Environmental Resources Congress,
 Orlando, Florida, USA, pp. 35–45.
- Weiss, A.D., 2001. Topographic position and landforms analysis. Poster Presentation,
 Esri Users Conference, San Diego, CA, USA.
- Wilson, J.P., 2012. Digital terrain modeling. Geomorphology 137, 107–121.
- Wilson, J.P., Gallant, J.C., 2000. Digital terrain analysis. In: Wilson, J.P., Gallant, J.C.
 (Eds.), Terrain Analysis: Principles and Applications. John Wiley and Sons, New
 York, pp. 1–27.
- Wood, J.W., 1996. The geomorphological characterisation of digital elevation models.
 (Unpublished) Ph.D. Dissertation, Department of Geography, University of
 Leicester, Leicester, UK, 456 pp. URL: http://www.soi.city.ac.uk/~jwo/phd/
 (accessed 11 June 2012).
- Wood, J., 1998. Modelling the continuity of surface form in DEMs. In: Poiker, T.,
 Chrisman, N. (Eds.), Proceedings of the 8th International Symposium on Spatial
 Data Handling, IGU, Vancouver, Canada, pp.725–736.
- Wood, J., 2009a. Overview of software packages used in geomorphometry. In: Hengl,
- T. and Reuter, H.I. (Eds.), Geomorphometry: Concepts, Software, Applications.
 Developments in Soil Science, Vol. 33, Elsevier, Amsterdam, Netherlands, pp. 257–
 267.
- 743 Wood, J., 2009b. Geomorphometry in LandSerf. In: Hengl, T. and Reuter, H.I. (Eds.),
- 744 Geomorphometry: Concepts, Software, Applications. Developments in Soil Science,
- 745 Vol. 33, Elsevier, Amsterdam, Netherlands, pp. 333–349.

- 746 Yokoyama, R., Shirasawa, M., Pike, R.J., 2002. Visualizing topography by openness: a
- new application of image processing to digital elevation models. Photogrammetric
 Engineering and Remote Sensing 68 (3), 257–265.
- Zandbergen, P.A., 2012, Python Scripting for ArcGIS. Esri Press, Redlands, California.
 368 pp.
- 751 Zevenbergen, L.W., Thorne, C.R., 1987. Quantitative analysis of land surface
- topography. Earth Surface Processes and Landforms 12, 47–56.
- 753