



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Supernumerary cheek tooth in a Byzantine horse from Theodosius Harbour, Istanbul, Turkey

Citation for published version:

Pasicka, E, Onar, V & Dixon, PM 2017, 'Supernumerary cheek tooth in a Byzantine horse from Theodosius Harbour, Istanbul, Turkey', *Equine Veterinary Education*, vol. 29, no. 5, pp. 266-269.
<https://doi.org/10.1111/eve.12462>

Digital Object Identifier (DOI):

[10.1111/eve.12462](https://doi.org/10.1111/eve.12462)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Equine Veterinary Education

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Supernumerary cheek tooth in a Byzantine horse from Theodosius Harbour, Istanbul, Turkey

E.Pasicka**, *V.Onar†* and *P.M. Dixon‡

**Department of Biostructure and Animal Physiology, Faculty of Veterinary Medicine, Wroclaw University of Environmental and Life Sciences, Poland;*

†Osteoarchaeology Practice and Research Centre, Department of Anatomy, Faculty of Veterinary Medicine, İstanbul University, Avcılar, Turkey;

‡The Royal (Dick) School of Veterinary Studies and The Roslin Institute, The University of Edinburgh, Easter Bush Veterinary Campus, Midlothian, Scotland EH25 9RG

**Corresponding author's email: edyta.pasicka@yahoo.com landline: 0048 71 3205746*

Keywords: Byzantium; cheek teeth; equine dentistry; horse; Istanbul; polyodontia

Summary

The subject was a mandible belonging to a morphologically mature horse of the late Byzantium period, discovered during excavations at Theodosius Harbor in Istanbul, Turkey that had a developmental molar tooth abnormality, i.e. a supernumerary molar tooth. This is an interesting case due to the rarity of supernumerary molars in archeozoological materials, and also because it is the only such case of equid polyodontia from the late Byzantium period from that archaeological site.

Introduction

Animal remains are among the materials most often acquired during exproation of archeological sites (Baker and Brothwell 1980; O'Connor 2000; Davis 2002; Bartosiewicz 2008; Lasota-Moskalewska 2008; Reitz and Wing 2008; Waldron 2009). Animal bone or dental remains from excavations are, mainly because of their stability over time, a source of invaluable information on the anatomy and morphology of the detected species (Bökönyi 1974; Baker and Brothwell 1980; O'Connor 2000; Davis 2002; Bartosiewicz 2008). Because of the presence of multiple skeletons in some sites, they are suitable for comparative, quantitative and qualitative analyses (von den Driesch 1976). These archeozoological findings also indicate the role which domesticated animals had in cultural development of communities at that time (Lasota-Moskalewska 2005), and how domestication affected the biological characteristics of those animals (Bökönyi 1974; Lasota-Moskalewska 2008). They are also a source of information on diseases of animals closely associated with humans (Bartosiewicz 2008; Waldron 2009). However, due to the fact that human consumption remnants prevail in archeozoological materials, animal remains with possible pathological abnormalities, including skull fragments with anomalies, are very rare (Hillson 2005; Lasota-Moskalewska 2008; Reitz and Wing 2008; Waldron 2009; Pasicka *et al* 2012, 2014).

Materials and methods

The analysed material consisted of a right-sided mandible (catalog no. MRY3467), belonging to a morphologically mature horse aged approximately 9-11 years old at the time of death. This age was estimated by examination of the very well preserved incisors, including assessment of the oval shape of their occlusal surface, and the presence of some of residual infundibula in all incisors (Fig 1). In this paper the Triadan system of equine dental

nomenclature (Fig 3) is used to identify individual teeth (Dixon and du Toit 2011). The well preserved undamaged right mandibular bone had loss of Triadan 406 and the presence of a caudally situated supernumerary molar tooth (Triadan 412). The attached rostral aspect of the left mandibular bone contained an incisor tooth and a portion of the left physiological diastema (Fig 2a-c). The presence of fully developed and erupted canine teeth confirms it was an adult male horse (Fig 1).

This specimen is a part of a collection owned by Osteoarchaeology Practice and Research Centre, Department of Anatomy, Faculty of Veterinary Medicine, İstanbul University. The mandible was mined during excavation at the site of Theodosius harbour at Yenikapi in Istanbul, Turkey. The age of this specimen was estimated by radiocarbon dating (^{14}C) as being from the period of Late Byzantium (15th century AD) (Onar *et al.* 2013). This jaw presents an anomaly in molar dentition uncommon for osteo-archaeological materials, as manifested by the presence of an additional cheek tooth (Lasota-Moskalewska 2008). It is also the only recorded occurrence of polyodontia in Equidae from the Byzantium period at the location in question (Onar *et al.* 2015).

Results and Discussion

Estimating the age, at which animals died on the basis of skeletal or dental remains, is hardly ever precise. Animals in prehistory were characterized by a slower ontogeny rate, compared to current species where there is a faster morphological puberty, manifested by more rapid dental development and closure of growth plates of long bones. Because of the geographical site of recovery of this skull, this horse was possibly an Arabian horse-type breed, whose incisor wear differs from other breeds (Muyllé 2011). Additionally, when determining the

age of an individual based on dental examinations one should consider that the age norms adopted in archaeozoological research have been established in modern species (Lasota-Moskalewska 2008). Visual examination of the incisor occlusal surfaces indicated the animal was 9-11 years old, but applying another method of ageing, namely radiographic examination of the reserve crowns and roots using the guidelines for modern horses of Dixon and Copeland (1993), this specimen could be aged as between 12 - 15 years of age when it died.

Anomalies in dentition occur in both man and animals (Hillson 2005; Reitz and Wing 2008; France 2009; Waldron 2009) and they can be divided into genetic, developmental, and acquired in origin (Baker and Brothwell 1980; Hillson 2005). Malocclusion is the most common equine dental disorder and is caused by uneven attrition of the cheek teeth occlusal surface, possibly due to dietary reasons (Lasota-Moskalewska 2008).

Among the common equine developmental dental abnormalities, one should list the atavistic polydontia (typical), associated with the occurrence of a rudimentary 105/205 tooth at the beginning of the row (*wolf tooth*, *dens lupinus*) (König and Liebich 2006).

Developmental dental abnormalities include anomalies of shape and position of teeth, reduced numbers (hypodontia) or even total absence of teeth (anodontia). Hypodontia must be differentiated from where a tooth has been lost due to disease or extracted during the animal's life (acquired anomaly), and the alveoli of such teeth shows signs of healing (Chaix *et al.* 1997).

Apparent supernumerary teeth may actually be due to retention of deciduous teeth. True polyodontia may be due to random divisions of dental primordia. Horses can also have

displaced polyodontia, exemplified by a dentigerous cyst, found on the dorsal aspect of the skull in horses (Jubb and Kennedy 1963).

The true prevalence of equid supernumerary teeth is unknown, but clinical surveys have shown it to occur more commonly in incisors than in cheek teeth (Bökönyi 1974; Dixon *et al.* 1999a, 1999b; Hillson 2005), and more commonly in younger than in adult horses (Bökönyi 1974; Dixon *et al.* 2005; Hillson 2005). However in donkeys, polyodontia was identified in 4-5% of cases aged 6 years and older (Rodrigues *et al.* 2013).

Examination of photographs and radiographs of this specimen showed loss of the Triadan 406 (– but no radiographic or gross anatomic evidence of alveolar disease was evident and so this loss is likely an artefactual post-mortem loss. On gross examination, there is a slight ventral swelling of the mandible, circa 3-4 cm in length beneath the Triadan 407 and 408, with a more focal 1-2 cm wide periosteal reaction beneath the cadual root of 407. Radiography does not show any abnormalities in the overlying 407 or 408 teeth, but confirmed the presence of a localised periostitis of the ventral mandible. In an equid of this age, this swelling is very likely to be due to a local mandibular trauma that occurred many months earlier. Younger (3-5 year old) equids commonly have mandibular swellings due to eruption cysts at this site (Dixon and du Toit 2011).

There exists an apparently artefactual, superficial, vertical fracture of the lateral aspect of the mandible between 407 and 408 – that is not apparent on radiography and so this fracture is also likely to be a post-mortem artefactual fracture (Fig 3).

The 411 that is normally the most caudal cheek tooth, has a normal occlusal surface, i.e. and contains the usual 6 pulp horns and the normal triangular occlusal shape of a mandibular Triadan 411 (Dixon and du Toit 2011). Lateral radiographs (Fig 3). of this tooth shows a wide reserve crown, and a poorly defined cadual root, as compared to all other cheek teeth roots in this specimen – but this wide reserve crown and delayed cadual root development is a common feature of the equid Triadan 411 mandibular tooth (Dixon and Copeland 1993).

As noted there is a supernumerary cheek tooth (412) present. Because of the absence of an antagonist tooth, this tooth has overgrown considerably (> 1cm) in height, particualry on its caudal aspect (Fig 3). If the animal had survived, this 412 overgrowth would have increased greatly and eventually caused a severe clinical problem by initially lacerating the tongue and soft tissues of the hard palate region during mastication and even later, possibly penetrating the hard palate (Dixon 2010). Food invariably becomes impacted into diastemata that commonly develop between the supernumeray and adjacent teeth leading to painful periodontal disease (Dixon *et al.* 1999b; 2005; Dixon 2010).

Radiographic examaintion of the apex of this supernumerary tooth indicates this to be a relatively recently (estimated to be less than 2- 3 years old) erupted tooth because there is no root (enamel free apical area) developed yet, even allowing that root development in a supernumerary teeth may not follow the usual pattern, and that this Byzantine period horse may not have grown and developed as quickly as modern horses. There is also a lucent area beneath the apex of the supernumerary tooth resembling an eruption cyst, with sclerosis of the adjacent alveolar lining. However, the height of the overgrowth on this tooth may indicate that the tooth has been erupted for possibly 4-6 years (Fig 3). This supernumerary tooth is also possibly dysplastic because does not taper (rostro-caudally) in an apical direction like a

normal cheek tooth (Dixon *et al.* 2012), but instead appears to be slightly wider more apically, even allowing that it is a young tooth. However, its structure is not that of a connated (more than one tooth joined together) supernumerary tooth (Dixon *et al.* 2010). Otherwise this tooth seems of normal morphology.

Based on the information gathered during analyses, we can conclude that the described mandible belonged to a morphologically adult horse, which survived with the described developmental polyodontia up to about 9 -11 years of age. There is no gross or radiographic evidence that the presence of this additional tooth was causing a clinical problem to this horse and it was very unlikely to have caused its death.

Authors declaration of interests

No conflicts of interests have been declared.

References

- Baker, J. R. and Brothwell, D. R. (1980) Animal Diseases in Archaeology. Academic Press, London. pp 235.
- Bartosiewicz, L. (2008) Taphonomy and palaeopathology in archaeozoology. *Geobios* **41**, 69-77.
- Bökönyi, S. (1974) History of domestic mammals in Central and Eastern Europe. Akadémiai Kiadó. Budapest.

171 Chaix, L., Bridault, A., Picavet, R. (1997) A Tamed Brown Bear (*Ursus arctos*L.) of the Late
 172 Mesolithic from La Grande-Rivoire (Isère, France)? *J Archaeol Sci* **24** (12), 1067-1074.
 173
 174 Davis, S.J.M. (2002) *The Archaeology of Animals*. Routledge, London.
 175
 176 Dixon, P.M. (2010) Abnormalities of development and eruption in: *Equine Dentistry*, 3rd
 177 edn., Eds: K.J. Easley, P.M. Dixon and J.S. Schumacher, Elsevier Saunders. pp 99-114.
 178
 179 Dixon, P.M. and Copeland, A.N. (1993) The radiological appearance of mandibular cheek
 180 teeth in ponies of different ages. *Equine Vet. Educ.* **5**, 317-323.
 181
 182 Dixon, P.M. and du Toit, N. (2011) Dental Anatomy. In: *Equine Dentistry*, 3rd edn., Eds: K.J
 183 Easley, P.M. Dixon and J.S. Schumacher. Elsevier Saunders. pp 51-76.
 184
 185 Dixon, P.M., Easley, K.J. and Ekmann A. (2005) Supernumerary teeth in the horse. *Clin.*
 186 *Tech. Equine Pract.* **4**, 155-161.
 187
 188 Dixon, P.M., Tremaine, W.H., Pickles, K., Kuhns, L., Hawe, C., McCann, J., McGorum,
 189 B.C., Railton, D.I. and Brammer, S. (1999a) Equine dental disease - a long term study of 400
 190 cases: Part I –Introduction and disorders of incisor, canine and first premolar teeth. *Equine*
 191 *Vet. J.* **31**, 369-377.
 192
 193 Dixon, P.M., Tremaine, W.H., Pickles, K., Kuhns, L., Hawe, C., McCann, J., McGorum B.C.,
 194 Railton, D.I. and Brammer, S. (1999b) Equine dental disease - a long term study of 400

195 cases: Part II - Disorders of development, eruption and variations in position of the cheek
 196 teeth. *Equine Vet. J.* **31**, 519-528.

197

198 France, D.L. (2009) Human and nonhuman bone identification. A color atlas. CRC Press
 199 Taylor and Francis Group.

200

201 Hillson, S. (2005) Teeth. 2ed edn. Cambridge University Press, Cambridge, UK.

202

203 Jubb, K.V.F. and Kennedy, P.C. (1963) Pathology of Domestic Animals. Vol 2. Academic
 204 Press, New York and London.

205

206 König H.E. and Liebich H-G. (2007). Veterinary Anatomy of Domestic Mammals. Textbook
 207 and Colour Atlas. 1st edn. Schattauer, Stuttgart-New York.

208

209 Lasota-Moskalewska A. (2005) Zwierzęta udomowione w dziejach ludzkości. Wydawnictwa
 210 Uniwersytetu Warszawskiego, Warszawa.

211

212 Lasota-Moskalewska A. (2008). Archeozoologia. Ssaki. Wydawnictwa Uniwersytetu
 213 Warszawskiego, Warszawa.

214 Muylle, S (2011) Ageing. In: Equine Dentistry, 3rd edn., Eds: K.J Easley, P.M. Dixon and J.S.
 215 Schumacher. Elsevier Saunders. pp 85-94.

216

217 O'Connor T. (2000) The archaeology of animal bones. Sutton Publishing, Stroud.

218

219 Onar V., Alpak H., Pazvant G., Armutak, A., Gezer İnce N. and Kiziltan Z. (2013) A Bridge
 220 from Byzantium to Modern Day İstanbul: An Overview of Animal Skeleton Remains Found
 221 During Metro and Marmaray Excavations. *J. Fac. Vet. Med. İstanbul Univ.* **39** (1), 1-8.
 222 Onar, V., Pazvant, G., Pasicka, E., Armutak, A. and Alpak, H. (2015) Byzantine Horse
 223 Skeletons of Theodosius Harbour: 2. Withers height estimation. *Revue Med. Vet.* **166** (1-2),
 224 30-42.
 225 Pasicka, E., Chrószcz, A., Janeczek, M. and Mucha, A. (2012) Craniometric analysis of Early
 226 Medieval horses *Equus przewalskii* f. *caballus* (Linnaeus, 1758) from chosen areas in Poland.
 227 *Turk. J. Vet. Anim. Sci.* **36**, 688-697.
 228
 229 Pasicka, E., Chrószcz, A., Tarnawski, K. and Janeczek, M. (2014) Characteristics of
 230 Maxillary Cheek Teeth in Horses *Equus przewalskii* f. *caballus* (LINNAEUS, 1758) from
 231 Early Medieval Excavations in Poland. *Global Veterinaria.* **12** (3), 336-344.
 232
 233 Reitz, E.J. and Wing, E.S. (2008) Zooarchaeology. 2ed Edn. Cambridge University Press,
 234 Cambridge, UK.
 235
 236 Rodrigues, J.B., Sanroman-Llorens, F., Bastos, E., San Roman, F. and Viegas, C.A. (2013)
 237 Polydontia in donkeys. *Equine Vet. Educ.* **25** (7), 363-367.
 238
 239 Von den Driesch, A. (1976) A guide to the measurement of animal bones from archaeological
 240 sites. Peabody Museum Bulletin 1, Harvard University, Massachusetts.
 241
 242 Waldron, T. (2009) Palaeopathology. Cambridge University Press, Cambridge, UK.
 243

244
245
246
247
248



249
250
251
252
253
254
255
256
257
258

Fig 1: Incisor teeth of a Byzantine period horse recovered from Theodosius Harbour. Infundibular remnants are still present in all incisors; including a well-defined, small “cup” in 302 and irregular shaped enamel “rings” (“marks”) in 303 and 403 (black arrows).

259
260
261
262
263



264
265
266
267
268
269
270
271

Fig 2a: Right mandible of a Byzantine period horse recovered from Theodosius Harbour: lateral view.

272 dyta Pasicka
273 <edyta.pasicka@yahoo.com>



274

275

276

277

278 Fig 2b: Close up view of a right mandible of a Byzantine horse recovered from Theodosius Harbour:
279 lateral view.

280 The 406 tooth – is missing (site indicated by star, 407, 408 , 409 , 410 , 411 (M₃), and an overgrown
281 supernumerary tooth (412 -arrow) are present

282

283

284

285

286

287




Fig 2c: Right mandible of a Byzantine perido horse recovered from Theodosius Harbour: dorsal view.
The 406 tooth is missing; and a supernumerary 412 (arrow) is present.




Fig 3: Lateral Radiograph of a mandible of a Byzantine period horse recovered from Theodosius Harbour with teeth labelled using the Triadan system:.

▲-indicates a swelling, circa 3-4 cm in length beneath the Triadan 407 and 408, with a more focal 1-2 cm wide periosteal reaction over the cadual root of 407; ↑- poorly defined, wide caudal root of 412.