THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

"Megalosaurus cf. superbus" from southeastern Romania: the oldest known Cretaceous carcharodontosaurid (Dinosauria: Theropoda) and its implications for earliest Cretaceous Europe-Gondwana connections

Citation for published version:

Csiki-Sava, Z, Brusatte, S & Vasile, S 2016, "'Megalosaurus cf. superbus' from southeastern Romania: the oldest known Cretaceous carcharodontosaurid (Dinosauria: Theropoda) and its implications for earliest Cretaceous Europe-Gondwana connections', *Cretaceous Research*, vol. 60, pp. 221-238.
<https://doi.org/10.1016/j.cretres.2015.12.004>

Digital Object Identifier (DOI):

[10.1016/j.cretres.2015.12.004](https://doi.org/10.1016/j.cretres.2015.12.004)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Cretaceous Research

Publisher Rights Statement:

Copyright © 2015 Elsevier Ltd. All rights reserved.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Manuscript Number: YCRES-D-15-00207R2

Title: "Megalosaurus cf. superbus" from southeastern Romania: the oldest known Cretaceous carcharodontosaurid (Dinosauria: Theropoda) and its implications for earliest Cretaceous Europe-Gondwana connections

Article Type: Full Length Article

Keywords: Southern Dobrogea; Valanginian; Carcharodontosauridae; cratonic Europe; palaeobiogeography

Corresponding Author: Dr. Zoltan Csiki-Sava, Ph.D.

Corresponding Author's Institution: University of Bucharest

First Author: Zoltan Csiki-Sava, Ph.D.

Order of Authors: Zoltan Csiki-Sava, Ph.D.; Stephen L Brusatte; Ștefan Vasile

Abstract: Some of the best records of continental vertebrates from the Cretaceous of Europe come from Romania, particularly two well-known occurrences of dwarfed and morphologically aberrant dinosaurs and other taxa that lived on islands (the Cornet and Hațeg Island faunas). Substantially less is known about those vertebrates living in the more stable, cratonic regions of Romania (and Eastern Europe as a whole), particularly during the earliest Cretaceous. We describe one of the few early Early Cretaceous fossils that have ever been found from these regions, the tooth of a large theropod dinosaur from Southern Dobrogea, which was discovered over a century ago but whose age and identification have been controversial. We identify the specimen as coming from the Valanginian stage of the Early Cretaceous, an incredibly poorly sampled interval in global dinosaur evolution, and as belonging to Carcharodontosauridae, a clade of derived, large-bodied apex predators whose earliest Cretaceous history is poorly known. Quantitative analyses demonstrate that the Romanian tooth shows affinities with a derived carcharodontosaurid subgroup, the Carcharodontosaurinae, which until now has been known solely from Gondwana. Our results suggest that this subgroup of colossal predators did not evolved vicariantly as Laurasia split from Gondwana, but originated earlier, perhaps in Europe. The carcharodontosaurine diversification may have been tied to a north-to-south trans-Tethyan dispersal that took place sometime between the Valanginian and Aptian, illustrating the importance of palaeogeographic ties between these two realms during the largely mysterious early-mid Early Cretaceous.

Dear Editors,

We are submitting here the latest, corrected version of our manuscript about an Early Cretaceous carcharodontosaurid dinosaur from southeastern Romania. We have implemented all the changes suggested by the Handling Associated Editor, as outlined below. In the case of the changes concerning the reference list, we have included two different versions, as explained below, due to uncertainties we have about the correct formatting style. We hope that this revised version is now suitable for publication in Cretaceous Research.

Sincerely,

Zoltan Csiki-Sava (also on behalf of the co-authors)

Editor comments:

Page 1 – email addresses removed from other authors than corresponding author, as instructed.

Page 2 – some of the former keywords were replaced by newly selected ones, as suggested.

Page 3, line 75 – ‘in prep.’, removed, this work is still ongoing and mentioning it is not entirely necessary.

Page 11, line 258 – ‘in part’ removed; this is a controversial detail of the local stratigraphy that is of no importance for the economy of this manuscript.

Page 18, line 441 – we have added 3 to ‘Figure’, thanks for spotting this omission!

Page 22, line 526 – the correct reference is Williamson and Brusatte, 2014, as it is correctly listed in the reference list. We have corrected this reference; again, we are grateful for noting this error.

Page 32, line 786 – ‘2102’ was replaced by ‘2012’; also, we have updated (here and in the reference list) the reference ‘Lü et al., 2014’, published in the meantime, to ‘Lü et al., 2016’.

Page 36, line 884 – ‘see below’ was removed from the text, as suggested.

Page 37, line 917 – ‘KcKenna’ corrected to ‘McKenna’

References:

We have checked the latest issue(s) of CR, and are somewhat confused as to the required formatting of the references. We have compared several recently published papers, and have found two different formatting styles, e.g., one that comes from our own recent paper (Csiki-Sava, Z. et al., 2016. The East Side Story - The Transylvanian latest Cretaceous continental

vertebrate record and its implications for understanding Cretaceous-Paleogene boundary events. *Cretaceous Research*, 57: 662-698. <http://dx.doi.org/10.1016/j.cretres.2015.09.003>) and one in the still more recent paper of Averianov and Sues (Averianov, A. and Sues, H.-D., 2016. Troodontidae (Dinosauria: Theropoda) from the Upper Cretaceous of Uzbekistan. *Cretaceous Research*, 59: 98-110. <http://dx.doi.org/10.1016/j.cretres.2015.11.005>). Accordingly, in our resubmission we have included two different versions of our revised MS, each one following one of the two styles.

In one of the versions, we have retained our original, chronological formatting, that coincides with that used by Averianov and Sues (2016), while the second version follows the formatting implemented in Csiki-Sava et al. (2016) and also suggested by the Handling Editor. In this second version, we have made all of the changes suggested by the Editor, except a few instances such as:

Page 49, lines 1196-1214: Csiki-Sava et al. (2013, 2015) were kept after Csiki and Grigorescu (1998), Csiki et al. (2010), according to the alphabetical order recommended.

Page 60, lines 1475-1484: Rauhut and Werner (1995) kept before Richter et al. (2013) and Royo Torres et al. (2009), according to the alphabetical ordering we followed.

Research highlights

- An isolated, large theropod dinosaur tooth from Romania is referred to *Carcharodontosauridae*
- The Romanian carcharodontosaurid is Valanginian in age, the oldest Cretaceous record of the clade
- This occurrence supports dispersal from Europe to west-Gondwana during the mid-Early Cretaceous

○ basal / indeterminate Carcharodontosauridae	Carcharodontosaurinae	
☆ Romanian carcharodontosaurine	△ Carcharodontosaurus	▲ Mapusaurus
	▽ Giganotosaurus	▼ Tyrannotitan

Megalosaurus
sp. superbus 204
15.C.M. Saw.

1 “*Megalosaurus cf. superbus*” from southeastern Romania: the oldest known Cretaceous
2 carcharodontosaurid (Dinosauria: Theropoda) and its implications for earliest Cretaceous
3 Europe-Gondwana connections
4
5

6 Zoltán Csiki-Sava^{1*}, Stephen L. Brusatte², Ștefan Vasile¹
7 ¹ Department of Geology, Faculty of Geology and Geophysics, University of Bucharest, 1
8 Nicolae Bălcescu Boulevard, 010041 Bucharest, Romania, ~~zoltan.csiki@g.unibuc.ro,~~
9 ~~yokozuna_uz@yahoo.com~~

10 ² School of GeoSciences, University of Edinburgh, Grant Institute, James Hutton Road,
11 Edinburgh, EH9 3FE, United Kingdom, ~~Stephen.Brusatte@ed.ac.uk~~

12 * Corresponding author
13 zoltan.csiki@g.unibuc.ro

15 ABSTRACT

16 Some of the best records of continental vertebrates from the Cretaceous of Europe come from
17 Romania, particularly two well-known occurrences of dwarfed and morphologically aberrant
18 dinosaurs and other taxa that lived on islands (the Cornet and Hățeg Island faunas).
19 Substantially less is known about those vertebrates living in the more stable, cratonic regions
20 of Romania (and Eastern Europe as a whole), particularly during the earliest Cretaceous. We
21 describe one of the few early Early Cretaceous fossils that have ever been found from these
22 regions, the tooth of a large theropod dinosaur from Southern Dobrogea, which was
23 discovered over a century ago but whose age and identification have been controversial. We
24 identify the specimen as coming from the Valanginian stage of the Early Cretaceous, an
25 incredibly poorly sampled interval in global dinosaur evolution, and as belonging to

Carcharodontosauridae, a clade of derived, large-bodied apex predators whose earliest Cretaceous history is poorly known. Quantitative analyses demonstrate that the Romanian tooth shows affinities with a derived carcharodontosaurid subgroup, the Carcharodontosaurinae, which until now has been known solely from Gondwana. Our results suggest that this subgroup of colossal predators did not evolved vicariantly as Laurasia split from Gondwana, but originated earlier, perhaps in Europe. The carcharodontosaurine diversification may have been tied to a north-to-south trans-Tethyan dispersal that took place sometime between the Valanginian and Aptian, illustrating the importance of palaeogeographic ties between these two realms during the largely mysterious early–mid Early Cretaceous.

Keywords

~~Romania~~Southern Dobrogea; ~~Lower Cretaceous~~Valanginian; ~~Theropoda~~;
Carcharodontosauridae; cratonic Europe; palaeobiogeography

1. Introduction

Romania boasts one of the best records of continental vertebrate fossils from the Cretaceous of Europe (e.g., Grigorescu, 1992, 2003; Csiki-Sava et al., 2015). The vast majority of fossils come from two well-known occurrences: the Early Cretaceous bauxite accumulations of Cornet, in the northern Apuseni Mountains (e.g., Jurcsák, 1982; Benton et al., 1997; Posmoşanu, 2003; Dyke et al., 2011), and the famous latest Cretaceous beds of the Haţeg, Rusca Montană and western Transylvanian basins of Transylvania, which have yielded the dinosaur-dominated ‘Haţeg Island fauna’ (e.g., Nopcsa, 1923; Weishampel et al., 1991; Benton et al. 2010; Codrea et al., 2010, 2012; Grigorescu, 2010; Vremir, 2010; Vasile and Csiki, 2011; Csiki-Sava et al., 2015). Both of these faunas inhabited islands that were part of

the vast Cretaceous European Archipelago of the Neo-Tethys Ocean. Based on their isolated geological settings and the many dwarfed and morphologically aberrant taxa that make up the faunas, both have been interpreted as insular assemblages that give a unique window into how island environments affected the evolution of long-extinct organisms (e.g., Benton et al., 1997, 2010; Csiki-Sava et al., 2015).

The great volume of research on these assemblages over the past century, particularly the ‘Hațeg Island fauna’, has concealed an inconvenient bias: the stable, non-island, cratonic regions of Romania have yielded only extremely rare Mesozoic continental vertebrate remains (i.e., the Moldavian, Moesian and Scythian platforms; Săndulescu, 1984; Mutihac and Mutihac, 2010; Fig. 1). This is mostly because Mesozoic deposits are located in the subsurface in these regions, with only limited subaerial exposures available in the structurally highest-lying parts of the Moesian Platform, in Central and Southern Dobrogea (Middle Jurassic–Upper Cretaceous), as well as in the northeastern-most corner of the Moldavian Platform, along the Prut Valley (lower Upper Cretaceous) (see, e.g., Mutihac and Mutihac, 2010). This bias is unfortunate because fossils from these settings could lead to a better understanding of how mainland and island faunas differed during the Cretaceous, and because the cratonic portion of Europe was an important biogeographic stepping stone between the north and south as the continents fragmented and sea levels fluctuated.

Although the cratonic regions of Romania have yielded few Cretaceous terrestrial fossils, these deposits are not totally barren. In fact, one of the first Mesozoic continental vertebrates ever recorded from Romania comes from one of these deposits, the Lower Cretaceous shallow marine limestones of Southern Dobrogea (Fig. 1). This specimen—the isolated but well-preserved tooth of a large theropod dinosaur—has often been overlooked. It was described a little over a century ago by Simionescu (1913; Fig. 2A), and until a few recent discoveries of very rare isolated specimens (Stoica and Csiki, 2002; Csiki-Sava et al.,

2013, ~~in prep.~~; Dragastan et al., 2014), it remained as the sole published record of Mesozoic terrestrial vertebrates from the cratonic areas of Romania. It has never been comprehensively described and its precise age and taxonomic affinities have yet to be clarified, despite its potential importance as a well-preserved fossil from a poorly sampled area that could have critical evolutionary and biogeographic implications.

We here present a comprehensive description of the Dobrogea tooth and discuss its relevance for understanding dinosaur evolution and biogeography. We review the peculiar history of how this specimen was collected and curated, thoroughly document its morphology and age, identify it based on comparison to a broad range of theropods, and outline its importance. It turns out that this specimen, although only a single tooth, has wide-ranging implications. We identify it as coming from the Valanginian stage of the Early Cretaceous, which is incredibly poorly sampled both in Europe and globally (Weishampel et al., 2004), and as belonging to a carcharodontosaurid, a group of derived, large-bodied apex predators whose earliest Cretaceous history is poorly known. Carcharodontosaurids were once thought to be a uniquely Gondwanan group, but recent discoveries show that the basal members of the group were more widespread during the late Early-middle Cretaceous (e.g., Sereno et al., 1996; Brusatte and Sereno, 2008). The Romanian tooth shows affinities with a derived carcharodontosaurid subgroup, the Carcharodontosaurinae, that until now has been known only from Gondwana. It suggests that this subgroup of enormous predators did not evolve vicariantly as Pangaea split, but originated earlier, and perhaps in Europe, suggesting faunal interchange between Europe and Gondwana during the ‘dark ages’ of the early Early Cretaceous.

Abbreviations: UAIC – University “Alexandru Ioan Cuza”, Iași, Romania.

2. History of collecting and curation

101 Only two dinosaurian fossils are currently known from the cratonic areas of Romania: an
102 isolated theropod tooth and an isolated caudal vertebral centrum. Both of these were reported
103 from the Lower Cretaceous deposits of Southern Dobrogea (southeastern Romania; Csiki-
104 Sava et al., 2013, see also below). Unfortunately, exact details of their discovery and places
105 of origin are lost, a fact that can hinder an assessment of their age and interpretation of their
106 phylogenetic and palaeobiogeographic significance. Our aim here is to gather and report all
107 available information concerning the collecting of specimen UAIC (SCM1) 615, that is, the
108 isolated theropod tooth reported by Simionescu (1913; Fig. 2A).

109 According to the existing information - unpublished museum labels and records, and
110 the preliminary publication of Simionescu (1913) - specimen UAIC (SCM1) 615 was
111 discovered in the surroundings of Cochirleni, a small village south of Cernavodă and close to
112 the right bank of the Danube, in Southern Dobrogea, southeastern Romania (Fig. 1), probably
113 shortly before 1913, the date of its publication by Simionescu (1913).

114 Although studied and preliminarily described by Simionescu, UAIC (SCM1) 615 was not
115 collected by Simionescu personally. Instead, it was donated by a certain “de Tomas” (also
116 mentioned as “de Thomas” in the registry of the Hârşova Museum) to V. Cotovu from
117 Hârşova (Central Dobrogea), a local teacher, archaeology and natural history aficionado, and
118 amateur fossil collector (see, e.g., Covacef, 1995). Cotovu, described by Simionescu himself
119 as the “zélé fondateur et directeur du muséum de Hârşova” (*enthusiastic founder and director*
120 *of the Hârşova Museum*; Simionescu, 1906: p. 2), had previously provided fossil specimens
121 from Southern Dobrogea for study to Simionescu, a nationally acknowledged popular science
122 writer and scientist, whom Cotovu knew personally (Brânzilă, 2010). These circumstances
123 are supported by the fact that in the original description, Simionescu figures the specimen as
124 being accessioned in the “Regional-Museum von Harschowa” (*Hârşova Regional Museum*;
125 Simionescu, 1913: p. 687, fig.1), a designation he also used to refer to other Dobrogean

specimens not collected by him first-hand (e.g., a specimen of *Nautilus pseudoelegans* from Cernavodă, or a fragmentary tooth-bearing palatal fragment referred to as *Coelodus* sp., also originating from Cochirleni; see Simionescu, 1906). Confirming this deduction, an isolated tooth appears accessioned in the old registry book of the Hârşova Museum (under specimen number 200) as “*Megalosaurus cf. superbus*”, with the mention that it was “described by Prof. Simionescu in the *Centralblatt f. min. etc.*”. This is also the case of the *Coelodus* sp. specimen from Cochirleni (specimen number 86), similarly clearly identified as being described by Simionescu in the registry book.

Both of these vertebrate remains from Dobrogea that were formerly part of the Hârşova Museum collections are currently accessioned in the palaeontology collections of the UAIC (Turculeţ and Brânzilă, 2012), suggesting that, at one moment, several specimens were transferred there from the Hârşova Museum. Although no details are known about this transfer, it is probable that it took place right before (or when) the Hârşova Museum, including a part of its collections, was burned and largely destroyed during WWI, in 1916, a time when Simionescu still held a position at the UAIC.

After its original description, specimen UAIC (SCM1) 615 underwent a minor amount of damage (see below, Description). Also, at some point between its description in 1913 and the early 1960s (when the specimen was found in its present state in the collections of the UAIC by academic staff members who are still alive today and recall the discovery; I. Turculeţ, personal communication, May 2013) it was glued into a limestone matrix holder, while it was obviously completely freed of the surrounding matrix when it was described and figured in 1913 (Fig. 2). The circumstances under which these alterations took place are unclear. It is a distinct possibility that they occurred sometimes during WWII, when, in the spring of 1944, the frontline between the German-Romanian and Soviet armies reached the Iaşi–Chişinău line. At this moment, the geological-palaeontological collections of the UAIC

151 were packed in crates, and moved together with its personnel and other possessions to Zlatna,
152 in the Apuseni Mountains (western Romania), to safeguard them from any potential damage.
153 Mounting the specimen into the limestone stand would have been a quick way to stabilize it,
154 as it appears that packaging and transport of the specimens was done in haste (M. Brânzilă,
155 personal communication, April 2103). If that was indeed the case, the mounting would have
156 taken place without the knowledge of Simionescu, who left Iași and the UAIC in 1929, being
157 invited to become a professor of Palaeontology at the University of Bucharest (Brânzilă,
158 2010). Then again, however, Simionescu himself or staff of the Hârșova Museum might have
159 re-mounted the tooth after its original description, or else the mounting might have taken
160 place after the return of the collections to Iași, after WWII.

161 Unfortunately, it is not documented whether the mounting was made using the
162 original matrix, or if a trough corresponding to the tooth outline was carved into a randomly
163 chosen limestone block. The apparently excellent fit between the tooth and the depression
164 housing it (Fig. 2B, 3) suggests that this operation was completed carefully, and accurate
165 carving of a fake holder is difficult to reconcile with the rush accompanying the evacuation of
166 the Iași University, in 1944. Alternatively, the presence of a hand-written old registration
167 number on the specimen holder would support its early re-mounting, while still at the
168 Hârșova Museum. As noted previously, the original Hârșova Museum registration number of
169 the specimen was 200, which does not correspond to that currently written both on the
170 limestone holder and on a paper sticker (204). However, according to the old collection
171 registry of the Hârșova Museum, specimen numbers 201 through 225 were given to a series
172 of “indeterminate (fossil) bone fragments” from the “Cochirleni quarries”. Thus, these
173 specimens (now apparently lost) came from the same locality as the tooth, and they were
174 collected and donated by the same person to the Museum who donated UAIC (SCM1) 615.
175 There is, thus, a (albeit admittedly remote) possibility that the registration numbers were

176 mixed up during the re-mounting of the specimen, which in this case took place at an early
177 date in the Hârşova Museum. If this is indeed the case, the limestone used as holder could
178 have been the same as the original matrix of the specimen.

179 To conclude, the history of recovery and curation of the historically important
180 dinosaurian specimen UAIC (SCM1) 615 is rather convoluted and clouded by many
181 uncertainties. The exact date of discovery remains conjectural, and the exact place of the
182 discovery (thus also the original geological context of the tooth) is even more ambiguous.
183 The current state of the specimen, and especially its mounted status, suggest a curatorial
184 history that produced a moderate amount of damage to, but also partially obscured the
185 detailed morphology of the specimen. The convergence of such unfortunate events makes
186 deciphering the age, identity and evolutionary significance of the specimen troublesome,
187 although many lines of evidence, carefully considered, allow us to draw reasonable
188 conclusions (see below).

190 3. Geological setting

191 According to the available collecting information, the isolated theropod tooth UAIC (SCM1)
192 615 was discovered at Cochirleni (sometimes noted more specifically as the “*Cochirleni*
193 *quarry*” or “*Cokerleni quarry*”). Cochirleni is a small village in southwestern Dobrogea
194 situated close to the right bank of the Danube, and about 9 km south of the main urban center
195 of the region, Cernavodă (Fig. 1). The geology of the area has been well studied, because of
196 the unique outcropping conditions and rich fossiliferous nature of the Lower Cretaceous
197 deposits (reviewed in Avram et al., 1996; Neagu et al., 1997; Dragastan et al., 1998).

198 Southern Dobrogea is a cratonic area corresponding to the southeastern corner of
199 Romania. Whether it is considered part of the larger Moesian Platform (Săndulescu, 1984;
200 Ionesi, 1994), or a distinct craton (the South-Dobrogean Platform; Mutihac and Mutihac,

2010), researchers agree that it became integrated into the main European Craton towards the end of the Jurassic, at the latest, with the consolidation of the Cimmerian (Early Alpine: Triassic–earliest Cretaceous) North Dobrogean fold-and-thrust belt (Seghedi, 2001; Hyppolite, 2002). The age of its basement is also controversial, with estimates ranging from Archaic–Early Proterozoic (Mutihac and Mutihac, 2010) to latest Proterozoic (Ionesi, 1994).

The Precambrian basement of Southern Dobrogea is overlain by a flat-lying sedimentary cover that begins with the lowermost Palaeozoic and ends with the uppermost Neogene. The sedimentary succession is interrupted by a few major, as well as several less important, sedimentary hiatuses that separate 5 main sedimentary sequences corresponding to the Cambrian–Upper Carboniferous, the Permian–Triassic, the Middle Jurassic–Cretaceous, the Eocene–?Oligocene, and the middle Badenian (middle Miocene)–Upper Pliocene. The Palaeozoic and lower Mesozoic are known only from the subsurface of Southern Dobrogea, but Cretaceous and Cenozoic deposits have limited exposures along the main water courses of the region (Ionesi, 1994; Mutihac and Mutihac, 2010).

The outcropping Cretaceous in Southern Dobrogea is represented mainly by shallow marine, carbonate platform deposits in the lower part of the system, replaced by more open-water, chalky facies towards the later part of the period (e.g., Avram et al., 1993, 1996; Dragastan et al., 1998; Dinu et al., 2007); these crop out only as isolated patches along the main watercourses of the region (Fig. 1).

The Lower Cretaceous Series consists of several lithostratigraphic units with complex, partially overlapping and interfingering relationships (Dragastan et al., 1998, 2014). The lowest (and only artificially) outcropping unit is the Purbeck-type, siliciclastic-evaporitic Upper Kimmeridgian–Lower Berriasian Amara Formation that represents lagoonal to continental environments. This unit is covered by the shallow-marine, richly fossiliferous and locally reefal limestone-dominated Cernavodă Formation (restricted-open lagoonal to

carbonate platform, Upper Berriasian–Lower Hauterivian). A time-correlative unit of the Cernavodă Formation, the limestone-dolomitic Dumbrăveni Formation (Upper Berriasian–Lower Hauterivian), is restricted to the southeastern part of Southern Dobrogea. The Cernavodă and Dumbrăveni formations are covered unconformably by dominantly calcareous deposits with hippuritoid (‘pachyodont’) coquinas, small reefs and lens-like orbitolinid accumulations, referred to the Barremian–Lower Aptian Ostrov Formation by Dragastan et al. (1998), but to the Ramadan Formation (in part) by Avram et al. (1993, 1996). These deposits, formed in littoral to lagoonal and open reef terrace environments, are in turn capped by the fluvial-lacustrine, siliciclastic deposits of the Gherghina Formation, with Middle–Upper Aptian kaolinitic clays and thin coal intercalations. The Lower Cretaceous succession ends with the transgressive, glauconite-bearing, coastal to sublittoral siliciclastic deposits of the Cochirleni Formation (uppermost Aptian–Albian).

The Upper Cretaceous has a significantly more patchy development, mainly restricted to the eastern part of Southern Dobrogea, excepting the weakly glauconitic, chalky-sandy Peștera Formation (Lower Cenomanian) and the marly Dobromiru Formation (Upper Cenomanian) that cover the western-central parts of the area. The younger Cuza Vodă (Turonian), Murfatlar (Santonian–Lower-Middle Campanian), and Satu Nou (Upper Campanian) formations are dominantly chalky, suggesting the instalment of a relatively deeper, offshore depositional environment; neither of these units is known from western Southern Dobrogea.

In total, the Lower Cretaceous of Southern Dobrogea was deposited in a shallow marine, near-shore setting, fluctuating between carbonate platform, lagoonal, coastal-tidal flat, and continental environments (see Avram et al., 1996; Dragastan et al., 1998). Its main characteristic features, such as the observed lithological variability, the areal distribution of the different units, and the presence of several unconformities within the series, are all linked

to eustatic sea-level changes that affected the Southern Dobrogean territory during the Early Cretaceous (Dragastan et al., 1998). The main emergent land in the area was represented by the Central Dobrogean Massif, lying north of the study area, almost completely subaerially exposed and actively eroding during the Cretaceous. Consequently, shallow-marine to continental deposits are restricted mainly to the northern part of Southern Dobrogea, close to its boundary with the Central Dobrogean Massif (marked by the Capidava-Ovidiu Fault), and are replaced by more open marine deposits southward. As summarized above, several littoral, and even continental, sequences occur in this succession, including deposits in the Amara, Cernavodă, Ramadan (~~in part;~~ Avram et al., 1996) and Cochirleni formations, whereas the Gherghina Formation is purely continental, with occasional minor marine interbeds produced during short-term ingressions of the sea.

In the Cernavodă-Cochirleni area the outcropping Mesozoic is restricted to the Lower Cretaceous, and includes deposits belonging to the Cernavodă, Ostrov (or Ramadan), Gherghina, and Cochirleni formations. While the lower–middle part of the Cernavodă Formation is well exposed and widely distributed in this area, its upper part (the lower Hauterivian Vederoasa Member) is unevenly developed. This member is missing in the classical succession from Cernavodă-Hinog, on the right bank of the Danube (Dragastan et al., 1998), but was recently identified in the more eastern Cernavodă-lock section (Dragastan et al., 2014). Similarly, the Ostrov Formation is represented in the area only by its upper subunit (the Lower Aptian Lipnița Member; Dragastan et al., 1998), covering unconformably and transgressively the Valanginian Alimanu Member of the Cernavodă Formation in the southern end of the Cernavodă-Hinog section (Dragastan et al., 1998), and the lower Hauterivian Vederoasa Member in the Cernavodă-lock section (Dragastan et al., 2014).

Northward of the Hinog area, Valanginian deposits of the Alimanu Member are overlain directly by the Middle–Upper Aptian continental deposits of the Gherghina

Formation. These continental deposits also cover the *Orbitolina*-bearing calcareous-clayey deposits of the Lipnița Member towards the south, marking the advancement of emerged areas towards the central parts of Southern Dobrogea, including the Cernavodă-Cochirleni area, during this time interval (Avram et al., 1996). Marine conditions returned in the study area again in the latest Aptian, with a transgression marked by widespread deposition of the glauconitic, siliciclastic coastal to innermost shelf deposits of the Cochirleni Formation. These uppermost Aptian to Albian sands and sandstones cover transgressively all the underlying deposits, belonging to the Cernavodă, Ostrov, or Gherghina formations. Siliciclastic shallow-marine sedimentation continued into the Early Cenomanian, with the chalky-glauconitic deposits of the Peștera Formation.

4. Palaeontology

The isolated theropod tooth UAIC (SCM1) 615 (formerly in the collections of the Hârșova Museum, registered with no. 200; Fig. 2A) was described in a short note by Simionescu (1913), who referred it to *Megalosaurus* cf. *superbus*, a taxon erected by Sauvage (1882) from the Gault ('mid'-Cretaceous: Albian) of the Paris Basin, France. The Gault material described by Sauvage (1882; see also Sauvage, 1876) includes several isolated teeth that were deemed by Simionescu (1913) to be more similar to the Cochirleni tooth than are the teeth of *Megalosaurus bucklandi* (Buckland, 1824). Subsequently, the French Gault material was referred to the new genus *Erectopus* by Huene (1923), who also noted differences between it and the type species *M. bucklandi*.

The convoluted taxonomic history of *Erectopus superbus* was recently reviewed by Allain (2005), who established that both the isolated teeth first mentioned by Sauvage (1876) and the skeletal elements described by Sauvage (1882) belong to the same taxon, for which the name *Erectopus superbus* was retained. Allain (2005) regarded *Erectopus* as a member of

Carnosauria (= basal Tetanurae), an opinion also shared by Molnar (1990) and Holtz et al. (2004a), whereas the latest review of the Tetanurae (Carrano et al., 2012, p. 254) considered *Erectopus superbus* “a non-carcharodontosaurian allosauroid, possibly a metriacanthosaurid.” Accordingly, if we are following the original assessment of Simionescu (1913) but updating with contemporary taxonomy, the Cochirleni theropod tooth should now be considered referable to the basal tetanuran *Erectopus superbus*. However, the referral of this tooth to *Erectopus superbus* (or a close relative) was considered to be unsupported by positive evidence by Molnar (1990) and Holtz et al. (2004a). In order to re-assess this referral and to understand the exact taxonomic and phylogenetic affinities of UAIC (SCM1) 615 (Fig. 2B, 3), we provide here a detailed description of its morphology followed by a thorough comparative study of this tooth based on large datasets of theropod dental measurements and discrete characters compiled by Hendrickx and Mateus (2014) and Hendrickx et al. (2015a).

We note that in his review of Romanian dinosaurs, Grigorescu (2003) erroneously considered UAIC (SCM1) 615 as being referred by Simionescu to the taxon *Megalosaurus dunkeri* Kohen (sic; actually, *Megalosaurus dunkeri* Dames, 1884). This is clearly a simple misreading of Simionescu’s identification. Additionally, such a referral is also contradicted by the absence of mesial serrations in the holotype tooth of *M. dunkeri*, considered by Carrano et al. (2012) to represent an indeterminate theropod. The Dobrogea tooth, on the other hand, has mesial serrations (see below).

4.1. Age of UAIC (SCM1) 615

The age of UAIC (SCM1) 615 has been contentious, due to the uncertainties concerning its place of origin. Although it is often mentioned as originating from Cochirleni village (e.g., Grigorescu, 2003; Turculeț and Brânzilă, 2012), this has not been definitively established. According to the original report of Simionescu (1913), the tooth came from the upper part of

the Lower Cretaceous limestone succession exposed in the cliffs extending from Cernavodă to Cochirleni along the right bank of the Danube. The corresponding entry from the Hârşova Museum registry states that it was found in the ‘Cochirleni quarry’, a location that presently cannot be identified precisely. The only rocks to be quarried in the area are the calcareous deposits of either the Cernavodă or Ostrov formations, particularly the ones that crop out in the Danube bank cliffs between Cernavodă-Hinog-Cochirleni. Finally, although the mention ‘Cochirleni’ is usually considered to refer to Cochirleni village, it should be mentioned that the cliff-forming hill that extends between Cernavodă and Cochirleni is also known by the same name (Fig. 1). Taking all of this evidence into consideration, it is thus reasonable to conclude that the tooth was most likely found in the Lower Cretaceous limestone succession exposed in the Danube cliffs between Cernavodă and Cochirleni.

Based on the location of the discovery, in the upper part of the local limestone succession, and the age of the deposits from Cernavodă-Cochirleni known to him, Simionescu (1913) considered the tooth to be of Barremian age. Subsequently, the age of the tooth was given as Valanginian–Barremian (Weishampel, 1990; Weishampel et al., 2004) or Valanginian (e.g., Grigorescu, 2003), but without any supporting information.

New attempts have been made to more precisely constrain the age of UAIC (SCM1) 615. Dragastan et al. (2014) recently sampled the limestone matrix holder of the tooth, and reported from these samples an assemblage of foraminiferans, ostracods and microproblematicae (= *incertae sedis* microorganisms) that characterize their ‘Biozone IX with *Meandrospira favrei*’, of latest Valanginian age in the local lithostratigraphic scheme. In parallel, we also sampled the same limestone holder – a yellowish white, friable lime mudstone – that yielded a poor and badly preserved calcareous nannoplankton assemblage with *Watznaueria barnesiae*, *W. ovata*, *Nannoconus steinmanni*, *N. kamptneri*, *N. globulus*, *Calcicalathina* sp., *Speetonia colligata* and *Cyclagelosphaera deflandrei* (M. C. Melinte-

Dobrinescu, personal communication, November 2013), an assemblage that suggests a Berriasian–Hauterivian age of the limestone holder.

Since it is not clear if the limestone holder came from the same site as the tooth itself, we managed to take a second sample from the limestone matrix still partly filling the pulp cavity of the tooth, which must definitively be identical with the rocks the tooth was found in. This second, much smaller sample yielded only very scarce specimens of *Watznaueria barnesiae*, *Cyclagelosphaera margerelii* and *Diazomatolithus lehmanni* (M. C. Melinte-Dobrinescu, personal communication, November 2013), the latter two taxa having a peak in abundance during the Berriasian and, especially, the Valanginian.

In the nannoplankton succession reported previously by Avram et al. (1993) and derived from a systematic sampling of the Southern Dobrogean Lower Cretaceous, the concurrent presence of *Speetonia colligata*, *Calcicalathina oblongata*, *Diazomatolithus lehmanni* and *Nannoconus steinmanni* was noted in samples derived from the Alimanu Member of the Cernavodă Formation. These assemblages were interpreted to represent the nannoplankton zone CC3 of Sissingh (1977), of late Valanginian age. A comparable age was assigned to a roughly similar nannoplankton assemblage reported from the Lower Cretaceous of the Mecsek Mountains, Hungary, by Császár et al. (2000).

Together, all the available evidence (Simionescu's original account, geographic and geologic records, foraminifera, ostracods, microproblematicae, and calcareous nannoplankton) thus suggests that UAIC (SCM1) 615 originates from the Alimanu Member of the Cernavodă Formation, and it is most probably of late Valanginian age.

4.2. Description and comparisons

Specimen UAIC (SCM1) 615 is a large (total length, as preserved, is about 100 mm; Figs. 2, 3) lateral tooth of a theropod dinosaur, with a crown base length (CBL) of 29 mm, crown

base width (CBW) of 16.25 mm, crown height (CH) of 85.5 mm, and apical length (AL) of 91 mm (terminology following Smith et al., 2005 and Hendrickx et al., 2015b). It is remarkably well preserved, with the enamel in pristine condition. It preserves most of the crown and a small basal part of the root, but the crown tip is broken off, with an estimated 5 mm missing in the apical region.

In its present state, the mesial edge and part of the mesial third of the tooth are embedded in the limestone holder (Fig. 2B), although the tooth was once removed (see above, History of collecting and curation; Fig. 2A). Accordingly, it is exposed so that all faces of the tooth are widely visible, including the root region, except for the mesial surface.

Only the basal-most part of the root is preserved, and it is more complete near the mesial margin (Fig. 3B, C). Here, broken areas around the crown-root contact area (cervix) reveal details of the pulp cavity development, as well as the pattern of the dentine thickness variation (Fig. 3B–D). The crown also exhibits a transverse break at about two-thirds of its length (not present so obviously in the original figure of the specimen in Simionescu, 1913), and adjacent to it, the distal carina is also slightly chipped distal to mid-length. The labial face is superficially split near this break (Fig. 3A), while a more prominent region of damage appears on the lingual face, where a large (13 x 5 mm), slightly triangular wedge is broken off, exposing the deeper parts of the dentine (Fig. 3C). The damage to the lingual side apparently occurred after the original description of the tooth (Fig. 2), an observation that is concordant with the complex curatorial history of the specimen.

The basal-most, exposed part of the mesial face lacks the enamel cover (Fig. 3C, D), suggesting that this area already belongs to the root region. The mesial edge of the preserved crown base appears to be wider than the distal one, and is largely rounded transversely. Accordingly, the basal cross-section is teardrop-shaped (lanceolate); it is rounded mesially, but narrows distally into a small carina (Fig. 3D). As mentioned above, the pulp cavity is

exposed basally, being partly filled with a whitish-light gray limestone that is reminiscent of the matrix holder lithology. The pulp cavity narrows rapidly towards the cervix, as it is about 7.1 mm wide (labiolingually) at the apical-most part of the preserved root, but only about 4.5 mm wide at the base of the crown. In parallel, the enamel-dentine wall of the tooth becomes thicker: it is 3.5 mm thick in the apical-most part, 4.4 mm at the base of the crown, but thickens to 5.0–5.8 mm near the apical-most part of the basal break of the crown (Fig. 3B). Mirroring the outside cross-section, the contour of the pulp cavity is also teardrop-shaped (Fig. 3D).

The tooth is ziphodont and only very slightly recurved distally. The distal edge is nearly straight across its length, being very mildly concave in its basal half and slightly convex near its apex (Fig. 2, 3A). Thus, the apex is placed roughly at the distal margin of the tooth crown base. The mesial edge, as shown in the original publication of Simionescu (1913), is strongly convex across its entire length (Fig. 2A). The tooth is labiolingually compressed (Fig. 3B), with a crown base ratio ($CBR = CBW/CBL$) of 0.56, within the normal range of variation of most theropods. This differs from the thinner teeth of some, but not all, carcharodontosaurids ($CBR < 0.50$), and the much thicker incrassate teeth of derived tyrannosauroids and conical teeth of spinosaurids ($CBR > 0.75$) (Serenó et al., 1996; Brusatte et al., 2010a; Hendrickx and Mateus, 2014; Hendrickx et al., 2015a).

The crown cross-section is slightly asymmetrical labiolingually when it is seen in distal view. In this view, when the carina is facing directly distally, one side of the crown has a more pronounced bulge than its counterpart (about 8.5 mm wide, measured from the carina, vs. 6 mm on the other side; Fig. 3B); based on comparisons with the teeth of *Mapusaurus* (Coria and Currie, 2006), the more bulging side can be interpreted as the lingual one. This asymmetry diminishes apically, where both sides become about equally convex. The distal carina itself twists slightly sideways (labially) in apical direction, such that it is located closer

to the labial face where it terminates at the crown apex, and the lingual face of the denticles is exposed distally (Fig. 3B, F). This twist of the distal carina is accompanied by a similar outline of the lingual side; in distal view, this is somewhat convex basally, but becomes flat to slightly concave in the apical two-thirds of the crown. A similar S-shaped curvature of the crown, albeit more pronounced and different in details, was also reported in *Mapusaurus* and *Giganotosaurus* (Coria and Currie, 2006), and in indeterminate carcharodontosaurid teeth from Morocco (Richter et al., 2013).

The distal carina extends along the entire tooth height (Fig. 3A–C). It is covered with minute serrations across its entire preserved length; the denticles are proximodistally subrectangular, with a mesiodistal long axis that is greater than the apicobasal long axis (Fig. 3E–H). They are either roughly perpendicular to the tooth margin, or their long axes are oriented obliquely, such that they point slightly apically. The tip of the apex is broken off, so it is not possible to determine whether the serrations continued over the apex of the tooth. There are approximately 12.5 serrations (denticles) per 5 millimetres at the midpoint of the carina. Serration shape and size remain relatively constant across the carina, although the serrations near the midpoint and closer to the base of the carina (12 denticles per 5 mm; Fig. 3G, H) are slightly smaller than those near the apex (9 denticles per 5 mm; Fig. 3E, F). Changes in serration size are gradual across the carina, not sudden or sporadic.

Although they are all more or less rectangular in shape, the apical denticles are relatively shorter proximodistally than the more basal ones. Most of the denticles have slightly rounded, asymmetrically convex triangular tips, instead of being simply squarred-off, and they do not hook as in troodontids and to a lesser extent abelisaurids (Hendrickx and Mateus, 2014). Other denticles near the apex, however, show a faint concavity along their tips, giving them a bilobate aspect, although this is both less conspicuous and far less regularly developed than reported in *Tyrannotitan* (Novas et al., 2005). The denticles are

separated by simple, linear grooves (interdenticular slits or sulcae) along their entire length.

The interdenticular space between adjacent denticles is broad, measuring more than a third of the apicobasal width of a denticle (Fig. 3E, G). This space continues onto the surface of the crown as a very short interdenticular sulcus (“blood groove” of Currie et al., 1990). These sulci are so short and indistinct that they are only visible under low angle light.

Little can be said about the mesial carina, as it is not visible in the current state of the specimen, buried in the limestone matrix. Based on the description of Simionescu (1913), however, it is covered across its length with minute serrations; these decrease in size towards the base of the crown. Simionescu (1913) reported approximately 15 serrations (denticles) per 5 millimetres at the midpoint of the carina, meaning that the mesial denticles are slightly smaller than those on the distal carina. The denticle size difference index (DSDI: Rauhut and Werner, 1995) is 1.2, within the range of variation of most theropods (Hendrickx and Mateus, 2014). As Simionescu (1913) already pointed out, the presence of a mesial carina that extends towards the base of the crown sets apart UAIC (SCM1) 615 from *Megalosaurus bucklandii* where this stops well above the cervix (Benson et al., 2008), and it is instead similar to ‘*M. superbis*’ (Sauvage, 1876, 1882) in this respect.

The external enamel surface exhibits two forms of ornamentation. First, the majority of the labial and lingual faces are covered by relatively smooth enamel that exhibits a subtle form of braided texture visible under low angle light (Fig. 3A, C, E). This texture is made up of a series of very faint, apico-basally running ridges; these are of unequal lengths, starting at different points of the crown height, but none extends the whole length of the crown. The two longest ridges are placed near the distal carina. The enamel is also finely granulated.

Second, near the carinae on both labial and lingual surfaces there are marginal undulations: wrinkles in the enamel that stand out in bas relief (Brusatte et al., 2007). These are much better preserved and visible near the distal carina, where they are so pronounced

that they are clearly observable in normal light (Fig. 3A–C, G, H). Here, about 17 unevenly developed wrinkles are present along the crown height; in the basal half of the crown, the wrinkles extend about 6.5 mm onto the crown. These are elongate, such that they are longer than twice the space separating each undulation. The wrinkles project obliquely (in the mesiobasal direction) relative to the carina. They are apically concave, with a near-horizontal segment on the crown, and curve apically as they approach the carina (at about 45°) with a tendency to become tangential to the distal edge. The wrinkles are especially well developed, prominent and closely spaced in the basal part of the crown (about 7 wrinkles/16 mm; Fig. 3C, G)), but become more widely spaced and indistinct apically (about 3 wrinkles/16 mm). Apically, however, the wrinkles are somewhat wider and longer, extending over about half of the crown fore-aft length. Again, a slight asymmetry is present between the two sides of the crown in wrinkle development as well, these being better expressed on the more rounded, convex lingual face, but less well expressed on the flatter labial face (Fig. 3A, C, H). On the presumed labial face, only some of the basal-most wrinkles, particularly the second and third one, appear well defined.

Towards the base of the crown a few of the wrinkles continue across the labial and lingual surfaces as very subtle transverse undulations. Most conspicuous of these is a 3.5 mm wide horizontal swelling that crosses the crown, at the level of wrinkles 2 and 3; this swelling is clearly visible on both sides of the crown (Fig. 3. A, C). There are no lateral flutes, apico-basal ridges, or longitudinal grooves on the labial or lingual faces, either in the centre of the tooth or paralleling the carinae. Instead, the labial and lingual faces are uniformly convex, giving the tooth its teardrop-shaped outline in cross section.

5. Discussion

5.1. *Identification of UAIC (SCM1) 615*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

501 The isolated tooth from Cochirleni can be referred to Theropoda based on its large size,
502 recurved and labiolingually compressed morphology, and presence of a continuous series of
503 well-defined serrations on the distal carina.

504 Besides theropods, certain derived crocodyliforms – the sebecosuchians of Colbert
505 (1946; see also Turner and Sertich, 2010; Pol and Powell, 2011; Rabi and Sebők, 2015) – are
506 also known to possess remarkably theropod-like, laterally compressed and serrated teeth, not
507 unlike the morphology shown by UAIC (SCM1) 615. However, most sebecosuchian teeth are
508 significantly smaller than the Southern Dobrogean specimen, especially in the case of the
509 Cretaceous members of the clade (e.g. *Baurusuchus*; Carvalho et al., 2005). Even the largest,
510 caniniform teeth of the largest representatives of Sebecosuchia, such as the Miocene
511 *Barinasuchus* (Paolillo and Linares, 2007), are somewhat smaller than UAIC (SCM1) 615;
512 moreover, these teeth are slightly conical and less laterally compressed than the Southern
513 Dobrogean tooth. Finally, it should be noted that the oldest known members of Sebecosuchia
514 appear beginning in the Late Cretaceous (e.g. Kellner et al., 2014), and are thus significantly
515 younger than UAIC (SCM1) 615. Similarly, ziphodont crocodyliform teeth (i.e. with true
516 denticles along their carinae) are reported in Europe only beginning in the Albian (Ösi et al.,
517 2015), and these are both significantly smaller and different in morphology from the
518 Dobrogean tooth. Taken together, these suggest that the hypothesis of sebecosuchian
519 affinities of UAIC (SCM1) 615 can be discarded with confidence, and it indeed represents a
520 theropod tooth.

521 We used four techniques to identify which type of theropod UAIC (SCM1) 615 likely
522 belongs to (see also Supplementary Material).

523 First, we conducted a Principal Components Analysis (PCA) based on a large
524 database that includes a broad and representative sample of theropod teeth. This dataset was
525 compiled by Hendrickx et al. (2015a), which built upon the earlier studies of Smith et al.

(2005) and Larson and Currie (2013), and it or a similar version has been used in recent studies to identify isolated theropod teeth (e.g., Williamson and Brusatte, ~~2013~~2014; Brusatte and Clark, 2015). It comprises nearly 1000 theropod teeth scored for six measurements (CBL, CBW, CH, AL, MC, and DC, the latter two measuring the density of serrations per 5 mm at the midpoint of the mesial and distal carina, respectively). UAIC (SCM1) 615 was added to this dataset, the data were log-transformed prior to analysis, missing values for measurements were estimated with a mean value for that measurement from across the sample, and then a PCA was run using a correlation matrix. The analysis was conducted in PAST v2.17 (Hammer et al., 2001).

In the resulting two dimensional morphospace (Fig. 4), UAIC (SCM1) 615 plots close to many teeth belonging to carcharodontosaurids, along with some teeth belonging to spinosaurids and tyrannosauroids. It falls within the convex hull (maximum morphospace occupation area) of carcharodontosaurids only, although it is closely outside of the edges of spinosaurid and tyrannosauroid space. It also falls within the 95% confidence interval ellipse for carcharodontosaurids, but not within the ellipse of any other group (Supplementary Information). This exercise indicates that UAIC (SCM1) 615 is most similar to carcharodontosaurids.

Secondly, we used the log-transformed dataset that we also used for the PCA to conduct a clustering analysis. We performed the analysis in PAST v2.17, using the paired group algorithm and the correlation similarity measure. In the resulting dendrogram, UAIC (SCM1) 615 groups with a handful of teeth belonging to carcharodontosaurids, tyrannosauroids, and *Allosaurus* (Supplementary Information).

Third, we used the tooth measurement database to conduct a discriminant analysis in PAST v3.0 (Hammer et al., 2001). This analysis uses pre-determined groups (in this case, taxonomic clusters) to create a morphospace in which these groups are maximally separated.

This allows teeth of unknown affinities, such as UAIC (SCM1) 615, to be classified according to which taxonomic group it is most similar to in this discriminant morphospace. In total, 67.79% of other teeth are classified correctly when they are treated as having uncertain affinities and their measurements are used to classify them in discriminant space, indicating that this exercise returns reasonable results. Our analysis classifies the Romanian tooth as a carcharodontosaurid. Furthermore, the analysis places UAIC (SCM1) 615 within the convex hulls for carcharodontosaurids and tyrannosauroids, and the 95% confidence ellipses for carcharodontosaurids, coelophysoids, and neovenatorids.

Fourth, we ran a phylogenetic analysis by including UAIC (SCM1) 615 in the discrete character dataset of theropod dental features published by Hendrickx and Mateus (2014). The Romanian specimen was scored as a lateral tooth in this analysis. The analysis was conducted in TNT (Goloboff et al., 2008), and resulted in 224 most parsimonious trees (686 steps, consistency index of 0.338, retention index of 0.566). The strict consensus topology is moderately well resolved and places the Romanian tooth as the sister taxon to *Carcharodontosaurus* (Supplementary Material). This sister taxon pair is recovered as the sister clade to a grouping of the derived carcharodontosaurids *Mapusaurus* and *Giganotosaurus*.

Several synapomorphies support the carcharodontosaurid affinities of UAIC (SCM1) 615. The sister group relationship with *Carcharodontosaurus* is supported by two features: a roughly straight distal margin of the crown (character 68) and pronounced marginal undulations in the enamel that are well visible in normal light (character 112). The broader clade of UAIC (SCM1) 615, *Carcharodontosaurus*, *Mapusaurus*, and *Giganotosaurus* (= Carcharodontosaurinae, as defined by Brusatte and Sereno, 2008, and Carrano et al., 2012) is linked by numerous characters, including: large teeth with a crown height greater than 6 cm (character 65), a bowed or sigmoid distal carina in distal view (character 82), marginal

undulations that are at least twice as long mesiodistally as the space separating each undulation (character 111), and marginal undulations present on both mesial and distal sides of the crown (character 113).

The Romanian specimen also lacks many keystone dental synapomorphies of other theropod clades, based on the clade diagnoses of Hendrickx and Mateus (2014) and other cladistic studies that include dental characters. UAIC (SCM1) 615 does not possess the hooked distal denticles of some Abelisauridae, the strongly labially deflected distal carina and pronounced transverse enamel undulations extending across the labial and lingual tooth faces of Ceratosauridae, the incrassate teeth with apicobasal enamel flutes and deeply veined enamel surface texture of Spinosauridae, and the large transverse undulations of some basal allosauroids (Hendrickx and Mateus, 2014). It also lacks the thickened incrassate teeth of derived tyrannosauroids (Brusatte et al., 2010a) and the large and strongly hooked (or pointed) denticles of troodontids and therizinosauroids (e.g., Turner et al., 2012; Brusatte et al., 2014; Hendrickx and Mateus, 2014). The large size, as well as recurved and ziphodont shape of UAIC (SCM1) 615 is strikingly different from the non-ziphodont therizinosauroids, ornithomimosaur, alvarezsauroids, and most troodontids, which have conical, leaf-shaped, or peg-like teeth (when teeth are present) (e.g., Holtz et al., 2004a; Turner et al., 2012; Brusatte et al., 2014). Finally, besides its remarkably large size, the presence of serrations indicates that UAIC (SCM1) 615 does not belong to groups such as alvarezsauroids, oviraptorosaurs, basal troodontids, or avialans, which have unserrated crowns (e.g., Turner et al., 2012; Hendrickx and Mateus, 2014).

In summary, the four analyses all support carcharodontosaurid affinities for UAIC (SCM1) 615. Both overall tooth proportions and discrete phylogenetic characters point to a carcharodontosaurid identification, and the discriminant function analysis and phylogenetic analysis both explicitly recover the tooth as a carcharodontosaurid. For this reason we refer

1 this tooth to Carcharodontosauridae. Moreover, it appears to belong to a clade that unites very
2 derived and large-sized carcharodontosaurids (*Carcharodontosaurus*, *Giganotosaurus*, and
3 *Mapusaurus*), separated as such and named Carcharodontosaurinae by Brusatte and Sereno
4 (2008) and Carrano et al. (2012). The well-resolved internal topology of this clade, as
5 recovered in our analysis, is congruent with results of previous analyses based on larger sets
6 of characters from across the skeleton (e.g., Coria and Currie, 2006; Brusatte and Sereno,
7 2008; Brusatte et al., 2009; Ortega et al., 2010; Eddy and Clarke, 2011; Canale et al., 2015),
8 and offers some support for considering the Romanian carcharodontosaurid from Southern
9 Dobrogea as more closely related to the African *Carcharodontosaurus* than to the clade of
10 the South American giant carcharodontosaurids *Giganotosaurus* or *Mapusaurus*.

11 Two final notes are worth adding. First, our analyses also incorporated
12 carcharodontosaurids that are usually found to be basal within the clade, such as
13 *Acrocanthosaurus* and *Eocarcharia* (e.g., Harris, 1998; Sereno and Brusatte, 2008; Carrano
14 et al., 2012) as well as a host of other allosauroids, including members of Neovenatoridae
15 (*Neovenator*, *Australovenator* and *Fukuiraptor*), a clade that is often recovered as sister-
16 taxon to carcharodontosaurids within Carcharodontosauria (e.g., Benson et al., 2010; Carrano
17 et al., 2012; but see Novas et al., 2013; Porfiri et al., 2014, for an alternate placement of
18 neovenatorids in general). Both PCA and phylogenetic analysis clearly identified UAIC
19 (SCM1) 615 as more closely comparable morphologically to derived carcharodontosaurids
20 than to either basal carcharodontosaurids or to any other allosauroid subclade.

21 Second, our datasets also included teeth of *Erectopus*, the genus erected for
22 ‘*Megalosaurus*’ *superbus* to which UAIC (SCM1) 615 was originally referred. Again, our
23 analyses clearly indicate that there are no close morphological and morphometric similarities
24 between the two, which is in accordance with the suggestion of Carrano et al. (2012) that
25 *Erectopus* represents a non-carcharodontosaurid taxon, while our analysis identifies UAIC

(SCM1) 615 as a carcharodontosaurid. Instead, *Erectopus* groups with abelisauroids in the phylogenetic analysis. This is somewhat surprisingly, as Allain (2005) and Carrano et al. (2012) both identified *Erectopus* as a tetanuran. It should be noted, however, that Albian-aged abelisauroids are known from the same general area (eastern France) as that yielding the material referred to *Erectopus* (Accarie et al., 1995; Carrano and Sampson, 2008), raising the intriguing possibility that this taxon may represent an abelisauroid instead of an allosauroid tetanuran as suggested by Allain (2005) and Carrano et al. (2012). However, it must be remembered that this phylogenetic analysis is based on dental characters only, so it is probably more likely that *Erectopus* is a tetanuran with a dentition convergent to some extent with those of certain abelisauroids.

5.2. Body size of UAIC (SCM1) 615

One of the most salient and remarkable features of UAIC (SCM1) 615 is its large size. In the large and comprehensive sample of theropod teeth from our dataset, tooth size (estimated based on crown height – CH, and used as a rough proxy of body size) ranges from 2.2 mm (in the dromaeosaurid *Saurornitholestes* and the coelurosaur of uncertain affinities *Richardoestesia*) to 117.1 mm in the gigantic tyrannosauroid *Tyrannosaurus*. The Romanian specimen UAIC (SCM1) 615, with a CH of 85.5 mm, is ranked in the 60-80% maximum size (~ CH) range of the sample, and has a CH that is 73% of the largest tyrannosauroid teeth. Most of the teeth in the dataset (over 61% of the 966 measured teeth) are very small to small (less than 25 mm CH), and less than 10% of these fall in the 60-100% CH size categories. Teeth larger than UAIC (SCM1) 615 make up less than 5% of the total sample, and they represent only five taxa: the megalosaurid *Torvosaurus*, the tyrannosauroid *Tyrannosaurus*, the basal carcharodontosaurid *Acrocanthosaurus*, and the derived carcharodontosaurines *Carcharodontosaurus* and *Giganotosaurus*. Compared to other carcharodontosaurids, UAIC

(SCM1) 615 is smaller than the largest teeth of *Acrocanthosaurus* (9% difference), *Carcharodontosaurus* (20%), and *Giganotosaurus* (12.5%) in the dataset, but is 13% bigger than the largest tooth of *Mapusaurus*.

It is thus reasonable to conclude that UAIC (SCM1) 615 belonged to a large-sized carcharodontosaurid, comparable to, even if somewhat smaller than, the truly gigantic carcharodontosaurines *Giganotosaurus* and *Carcharodontosaurus* (Serenio et al., 1996; Calvo and Coria, 1998; Therrien and Henderson, 2007), taxa that were recovered as possible close relatives of the Romanian carcharodontosaurid by our phylogenetic analysis. This, in turn, corroborates growing evidence that very large body size was acquired very early in carcharodontosaurid history, since the earliest potential members of the clade are already of relatively large size (Rauhut, 2011). The oldest potential carcharodontosaurid is *Veterupristisaurus*, represented by isolated vertebrae that indicate an animal between 8.5 and 10 meters in total body length (compared to 11.5+ meters in *Acrocanthosaurus* and more derived carcharodontosaurids) (Rauhut, 2011). These specimens are known from the uppermost Jurassic of Tanzania, eastern Africa (Rauhut, 2011; Carrano et al., 2012; see below), predating at most ~18 million years (Mya) the occurrence of likely even larger-sized carcharodontosaurids in the Valanginian of Southern Dobrogea, Romania.

The inferred large body size of the South Dobrogean theropod is also remarkable as virtually all other dinosaur remains reported previously from Romania (both from the Early Cretaceous Cornet assemblage and the much later, end Cretaceous Hăţeg Island fauna) are significantly smaller, and many have been interpreted as insular dwarfs (e.g., Weishampel et al., 1993, 2003; Benton et al., 2006, 2010; Stein et al., 2010; Ősi et al., 2014). Although other Romanian theropod dinosaurs were not particularly dwarfed (e.g. Brusatte et al., 2013), they were nonetheless small (Nopcsa, 1902; Csiki and Grigorescu, 1998; Csiki et al., 2010; Brusatte et al., 2013). This bias towards small bodied Romanian theropods was also

interpreted as a consequence of their insular habitat (Csiki and Grigorescu, 1998), as all previously reported theropod remains come from within the Carpathian Orogen, an area with an archipelago-type palaeogeography during the Cretaceous (Dercourt et al., 2000; Csontos and Vörös, 2004; Csiki-Sava et al., 2015). By contrast, UAIC (SCM1) 615 was found in shallow marine deposits bordering the emerged areas of Central Dobrogea, part of the stable cratonic areas of Europe and connected at least intermittently to the Ukrainian Shield since the Late Jurassic (Fig. 5A). Although cratonic Europe was also transformed into an archipelago of islands during much of the Cretaceous, these islands were often both larger in size and more stable in space and time than were the transient emerged areas of the Tethyan archipelagoes. As such, it is conceivable that the Southern Dobrogean carcharodontosaurid was less constrained by space or resource limitations than the Tethyan insular dinosaurs, allowing it to retain a large body size.

5.3. UAIC (SCM1) 615 and Valanginian dinosaur distribution

Besides documenting the presence of large-sized mainland carcharodontosaurids in the Lower Cretaceous of Romania, UAIC (SCM1) 615 is also important in that it fills a significant gap in our knowledge on the composition and distribution of the Early Cretaceous dinosaurs in Europe. In their review of dinosaur occurrences, Weishampel et al. (2004) listed 83 Early Cretaceous dinosaur localities spread throughout Europe, more than half of these being known from the later part (Barremian–Albian) of that epoch; only around a dozen localities were listed from each age of the early part of the Early Cretaceous (Berriasian, Valanginian, and Hauterivian). Even despite a significant increase in Early Cretaceous dinosaur discoveries in Europe in recent years (e.g., Royo-Torres et al., 2009; Cobos et al., 2010, 2014; Galton, 2009; Norman, 2010, 2013; Pereda-Suberbiola et al., 2011, 2012; Sachs and Hornung, 2013; Blows and Honeysett, 2014), these remain very strongly biased towards

western and southwestern Europe (especially the UK, France and Spain). Frustratingly, no occurrences are known from the entire central, eastern and southern Europe for the Berriasian–Hauterivian time interval except for two from Romania: the Berriasian–Valanginian locality of Cornet (e.g., Jurcsák and Popa, 1979, 1983; Jurcsák, 1982; Benton et al., 1997) in the northern Apuseni Mountains of northwestern Romania, and the carcharodontosaurid tooth (Simionescu, 1913) from the Valanginian of Cochirleni, in Southern Dobrogea, southeastern Romania we are describing here (Fig. 5A).

Our identification of the Romanian tooth as a carcharodontosaurid documents the presence of this clade in Europe in the very early Cretaceous. This is significant, as carcharodontosaurids were widely distributed tens of millions of years later, in the middle Cretaceous (Aptian to Cenomanian), in western Gondwana (Africa and South America, see below). Despite the recent discoveries documenting that the clade was also present in North America and Asia during the middle Cretaceous (e.g., Sereno et al., 1996; Currie and Carpenter, 2000; Brusatte et al., 2009, 2012), there has been only very few occurrences in Europe, most importantly the Barremian-aged *Concavenator* from Spain (Ortega et al., 2010; see below). The carcharodontosaurid tooth from Southern Dobrogea is substantially older than *Concavenator*, demonstrating that carcharodontosaurids appeared in Europe earlier than previously thought and were a long-term component of the European mainland Early Cretaceous faunas. It also suggests that habitat-related palaeobiological differentiation might have been already present between the cratonic, stable European mainland, with a dinosaur fauna made up of normal-sized (even very large) taxa, and the islands from the mobile Alpine areas of the Mediterranean Neo-Tethys, with by now dwarfed dinosaurs such as those described from the Berriasian–Valanginian Cornet assemblage in northwestern Romania (Benton et al., 2006).

This Valanginian carcharodontosaurid represents an important datapoint not only for the Romanian Lower Cretaceous, but also for that of wider Eurasia. The Valanginian is a poorly documented age in dinosaur evolution, with very few precisely dated fossil occurrences from anywhere in the world (e.g., Weishampel et al., 2004). The best record of Valanginian dinosaurs is from Europe, with fewer and less well dated occurrences known from Asia, some of which have debatable or controversial dates. These include sites in Japan (e.g., Manabe and Hasegawa, 1995; Matsukawa et al., 2006; but see Kusuhashi et al., 2009 and Evans and Matsumoto, 2015, supporting an alternative, younger age of these assemblages) and in Thailand (e.g., Buffetaut and Suteethorn, 1998, 2007, with age constraints according to Racey, 2009; Racey and Goodall, 2009). Occurrences of possible Valanginian age from China (e.g., Jerzykiewicz and Russell, 1991; Shen and Mateer, 1992; Lucas and Estep, 1998) are either poorly constrained as early Early Cretaceous, or were shown subsequently to be younger than Valanginian (Lucas, 2006; Tong et al., 2009). Rare dinosaur remains of possible Valanginian (or ‘Neocomian’) age were also reported from southern Africa (e.g., De Klerk et al., 2000) and, tentatively, from North America (e.g., Lucas, 1901; McDonald, 2011, with age assignments according to Sames et al., 2010; Cifelli et al., 2014).

As one of the two known reports of Valanginian dinosaurs in Europe east of France, the Southern Dobrogean dinosaur record fills a huge palaeogeographic gap between the western European and the eastern Asian dinosaur faunas. Moreover, none of these early Early Cretaceous dinosaur assemblages from outside Europe include carcharodontosaurids (see below), as theropods are represented by coelurosaurians interpreted either as compsognathids (Gishlick and Gauthier, 2007) or basal ornithomimosaurs (Choiniere et al., 2012) in southern Africa, metriacanthosaurid allosauroids (‘sinraptorids’) in Thailand (Buffetaut and Suteethorn, 2007), and indeterminate allosauroids (Pérez-Moreno et al., 1993), non-

carcharodontosaurid tetanurans (Carrano et al., 2012) or enantiornithine birds (Lacasa Ruiz, 1989), besides indeterminate taxa (Carrano et al., 2012), in western Europe. This may suggest that carcharodontosaurids had not achieved a wide geographic distribution by this point in time, and that their more cosmopolitan distribution came later, during the middle Cretaceous.

Finally, the presence of the Cochirleani carcharodontosaurid might hint at the presence of palaeobiogeographic provinciality between the western and the eastern parts of Europe, partly mirroring those reported from the later part of the Late Cretaceous (e.g., Le Loeuff and Buffetaut, 1995; Weishampel et al., 2010; Ősi et al., 2012; Csiki-Sava et al., 2015). In the reasonably well sampled, and significantly better known, western European dinosaur faunas, Valanginian large carnivorous dinosaurs include non-carcharodontosaurid tetanurans (*Becklespinax*), as well as indeterminate allosauroids or indeterminate theropods (often described as ‘*Megalosaurus*’ *dunkeri*, ‘*M.*’ *insignis* or ‘*M.*’ *oweni*), none of which can be referred positively to Carcharodontosauridae (Carrano et al., 2012). The apparently provincial geographic distribution of the large-bodied theropods suggests that some degree of faunal differentiation was occurring within the European mainland, most probably promoted by geographic distance. Notably, this intra-European differentiation in theropod assemblages appears to stand in contrast with the faunal homogeneity reported in the case of the ornithomimids from the UK and Romania (e.g., Galton, 2009). It is important, however, to re-emphasize at this point that the Valanginian dinosaur fossil record is both exceedingly poor and patchy, even in Europe. Accordingly, further discoveries are needed to verify and support (or contradict) the presence of such a distribution pattern pointing to palaeobiogeographic provinciality inside Europe, as the one suggested by our carcharodontosaurid identification for UAIC (SCM1) 615.

5.4. UAIC (SCM1) 615 and carcharodontosaurid evolution and palaeobiogeography

Carcharodontosauridae were long considered as an exclusively Gondwanan group of theropods (e.g., Allain, 2002; Novas et al., 2005) since their first discovery in northern Africa (e.g., Stromer, 1931), and subsequent description of a host of referred taxa from the Aptian–Cenomanian of Africa and South America (Coria and Salgado, 1995; Sereno et al., 1996; Novas et al., 2005; Coria and Currie, 2006; Brusatte and Sereno, 2007; Sereno and Brusatte, 2008; Cau et al., 2013). This view started to change with the identification of the Early Cretaceous (Aptian–Albian) *Acrocanthosaurus* from North America as a basal carcharodontosaurid (e.g., Sereno et al., 1996; Harris, 1998; Sereno 1999; Brusatte and Sereno, 2008), suggesting that the clade had a wider, Neopangean palaeobiogeographic distribution by the mid–late Early Cretaceous. Such a wide distribution, even a cosmopolitan one, was further supported by the discovery of definitive carcharodontosaurids in the Lower Cretaceous of Europe (Ortega et al., 2010), and in the upper Lower to lower Upper Cretaceous of China (Brusatte et al., 2009, 2010b, 2012; Mo et al., 2014; Lü et al., 2014, 2016).

Together, the available evidence pointed to an early, pre-mid Early Cretaceous origin of the carcharodontosaurids, followed by their dispersal across Laurasia and western Gondwana beginning at least by the Aptian (Fig. 5B), a scenario that is concordant with the tentatively suggested presence of early carcharodontosaurids in the Upper Jurassic of Tanzania, which are based on fragmentary specimens (Rauhut, 2011; Carrano et al., 2012). It is also concordant with the widespread appearance of carcharodontosaurids in the fossil record starting with the Aptian, when they are reported in Africa (*Eocarcharia*; Sereno and Brusatte, 2008), South America (Vickers-Rich et al., 1999), North America (*Acrocanthosaurus*; Stovall and Langston, 1950; Harris, 1998; Currie and Carpenter, 2000; Eddy and Clarke, 2011), Europe (Canudo and Ruiz-Omeñaca, 2003; Pereda-Suberbiola et al.,

2012), and eastern Asia (*Kelmaysaurus*; Brusatte et al., 2012; ~~Lü et al., 2014~~; Mo et al., 2014; Lü et al., 2016).

During the Albian–Turonian, carcharodontosaurids became especially abundant and diverse in Africa (*Carcharodontosaurus*, *Sauroniops*; Stromer, 1931; Sereno et al., 1996; Brusatte and Sereno, 2007; Le Loeuff et al., 2012; Cau et al., 2013; Richter et al., 2013) and South America (*Tyrannotitan*, *Giganotosaurus*, *Mapusaurus*, alongside with indeterminate carcharodontosaurids; Coria and Salgado, 1995; Calvo and Coria, 1998; Novas et al., 2005; Coria and Currie, 2006; Casal et al., 2009; Candeiro et al., 2011; Canale et al., 2015; Fig. 5B). They were still present during this time interval in other continents, as well: in North America with *Acrocanthosaurus* until the Albian (D’Emic et al., 2012), in Europe until the Cenomanian (Vullo et al., 2007; Csiki-Sava et al., 2015), and in Eastern Asia with *Shaochilong* until the Turonian (Brusatte et al., 2009, 2010b; see also Chure et al., 1999).

After dominating terrestrial ecosystems at least in Africa, South America and eastern Asia during the Albian–Turonian (Brusatte et al., 2009; Coria and Salgado, 2005; Novas et al., 2013), carcharodontosaurids were considered to disappear from the fossil record after the Turonian in both Asia (Brusatte et al., 2009) and South America (e.g., Coria and Salgado, 2005; Calvo et al., 2006; Novas et al., 2013), to be replaced by other groups of large theropods such as tyrannosaurids in parts of Laurasia and abelisaurids in parts of Gondwana. Canale et al. (2009) even cautioned against assigning isolated theropod teeth from post-Cenomanian deposits of South America to Carcharodontosauridae (e.g., Canudo et al., 2008; Casal et al., 2009; Salgado et al., 2009) due to their morphological similarity to those of the abelisaurid *Skorpiovenator*. Recently, however, more diagnostic cranial remains were reported to suggest the survival of carcharodontosaurids into the latest Cretaceous (Campanian–Maastrichtian) in Brazil (Azevedo et al., 2013).

Contrasting with this rich and relatively continuous fossil record of Carcharodontosauridae starting with the Aptian, the first half of its evolutionary history is very poorly documented (Fig. 5B). Prior to the identification of UAIC (SCM1) 615, only two occurrences of pre-Aptian Cretaceous carcharodontosaurids were reported, one from the Barremian of Spain (Ortega et al., 2010; Gasca et al., 2014) and the other from the Barremian of Thailand (Buffetaut and Suteethorn, 2012). The Early Cretaceous *Kelmaysaurus* from Xinjiang, western China, was recognized as a carcharodontosaurid of possibly ?Valanginian to Aptian in age by Brusatte et al. (2012), but the deposits yielding these remains (the Lianmugui, or Lianmuxin, Formation of the Tugulu Group) were dated as Aptian–Albian by Eberth et al. (2001; see also Tong et al., 2009). An important temporal gap – of about 20 to 28 millions of years, according to the dates in Gradstein et al. (2012) – thus stretched between the oldest, tentatively assigned carcharodontosaurids from the Oxfordian–Tithonian of Tanzania, including the formally erected *Veterupristisaurus* (Rauhut, 2011; see also Carrano et al., 2012), and those that started to appear in the fossil record in the Barremian and then spread widely during the Aptian. Referral of UAIC (SCM1) 615 to Carcharodontosauridae partially fills this frustrating gap, effectively halving this shadowy period in the evolutionary history of the group.

Furthermore, our analyses tentatively cluster the Dobrogean theropod with the derived members of the Carcharodontosaurinae to the exclusion of the more basal, but significantly younger non-carcharodontosaurine carcharodontosaurids *Eocarcharia* and *Acrocanthosaurus*. If this placement is correct, then the Romanian tooth indicates that Carcharodontosaurinae diverged from other carcharodontosaurids considerably earlier than hitherto recognized.

The previously known fossil record of the clade suggested that Carcharodontosaurinae originated sometime between the Aptian and Albian, as basal carcharodontosaurids (*Acrocanthosaurus*, *Concavenator*, *Eocarcharia*) were moderately diverse in the Barremian–

Aptian, followed by the appearance of many fossils of carcharodontosaurines beginning in the Albian (Fig. 5B). The proposed affinities of the oldest carcharodontosaurid material – including isolated teeth referred to as ‘*Megalosaurus*’ *ingens* – from the east African Upper Jurassic, considered to be reminiscent of the Aptian–Albian *Acrocanthosaurus* (Rauhut, 2011), was also consistent with this evolutionary scenario. Now, our identification of UAIC (SCM1) 615 as a carcharodontosaurid dinosaur sharing important dental apomorphies with the derived Carcharodontosaurinae advocates the emergence of this clade (or at least the very large size and dental morphology characterizing it) well before the Albian, during or even before the Valanginian, and relegates taxa such as *Eocarcharia*, *Acrocanthosaurus* and *Concavenator* (the dentition of *Shaochilong* is unknown) as late-surviving members of the basal carcharodontosaurid radiation, with a relatively plesiomorphic dentition.

Besides shifting the emergence of the carcharodontosaurines earlier in time, identification of UAIC (SCM1) 615 as a carcharodontosaurid also has interesting palaeobiogeographic implications. As already noted, recent discoveries show that Carcharodontosauridae is not an endemic Gondwanan clade as was once proposed (e.g., Novas et al., 2005), with the identification of its widespread, Pangaeian distribution during the late Early Cretaceous (Sereno et al., 1996; Harris, 1998; Chure et al., 1999; Sereno, 1999; Brusatte and Sereno, 2008; Ortega et al., 2010; Brusatte et al., 2009, 2012; Mo et al., 2014). However, within Carcharodontosauridae itself, some palaeogeographic patterns have been widely accepted. For example, it has been widely acknowledged that Carcharodontosaurinae is a endemic subclade of Gondwanan carcharodontosaurids (e.g., Sereno 1999; Holtz et al., 2004b; Brusatte and Sereno, 2007; Sereno and Brusatte, 2008; Novas et al., 2013), as previously all its recognized members were restricted strictly to either Africa (Stromer, 1931; Sereno et al., 1996; Brusatte and Sereno, 2007) or South America (Coria and Salgado, 1995; Novas et al., 2005; Coria and Currie, 2006). Moreover, intra-clade relationships of

Carcharodontosaurinae were still adhering to patterns of continental fragmentation and vicariant evolution, with a basal split between the Albian–Cenomanian African *Carcharodontosaurus* and the Giganotosaurini, uniting the similarly Albian–Cenomanian southern South American *Giganotosaurus* and *Mapusaurus* (together with *Tyrannotitan*, if this taxon is also recovered within Carcharodontosaurinae; e.g., Novas et al., 2005, 2013).

This scenario is now challenged by our finding that the Southern Dobrogean carcharodontosaurid UAIC (SCM1) 615 may nest inside Carcharodontosaurinae. If true, such an affinity would suggest that the origin of Carcharodontosaurinae was not a southern, vicariant by-product of the Gondwana-Laurasia separation, a major palaeogeographic event that is considered to have been well underway by the end of the Jurassic, and essentially completed by the mid-Early Cretaceous (see Weishampel et al., 2010). Indeed, during this time palaeogeographic connections and faunal interactions were virtually non-existent between the northern Tethyan (European) and southern Tethyan (western Gondwanan, but essentially African) areas of the Mediterranean (e.g., Canudo et al., 2009; ~~see below~~), which makes a vicariant hypothesis intuitive. However, if the Romanian tooth represents a carcharodontosaurine, then it implies a much more complicated palaeogeographic history of the clade, which is not so clearly linked to continental breakup.

The palaeogeographic position of the Southern Dobrogean carcharodontosaurine in cratonic Europe, north of the Neo-Tethys, together with its significantly older age compared to other carcharodontosaurines, could indicate that separation of the carcharodontosaurine lineage took part in Europe and not in western Gondwana as previously assumed. This would also mean that representatives of this lineage were subsequently – after the Barremian – introduced to Africa and South America via trans-Tethyan dispersal, most probably at a time when faunal interactions between the southern and northern margins of the Mediterranean Tethys were resumed, after the early Barremian (Canudo et al., 2009).

Alternatively, it can be hypothesized that appearance of carcharodontosaurines in Southern Dobrogea is a consequence of southern immigration originating in western Gondwana, often considered the place of origin for this clade. However, this scenario has several potential caveats. Although Europe has been considered as forming part of a larger Eurogondwanan palaeobioprovince during the early Early Cretaceous (Ezcurra and Agnolín, 2012), and occasional trans-Tethyan faunal connections have been recognized between Africa and Europe during Late Jurassic to Early Cretaceous times (e.g., Gheerbrant and Rage, 2006), these interchanges either pre-dated the Berriasian (e.g., Gardner et al., 2003; Knoll and Ruiz-Omeñaca, 2009), or post-dated the Barremian (Canudo et al., 2009; Torcida Fernández-Baldor et al., 2011), with no positive evidence for actual faunal exchanges taking place during the ‘Neocomian’ (Berriasian–Hauterivian) time interval.

More recently, some potential evidence has emerged for Gondwana-to-Europe interchange during the ‘Neocomian’. The presence of the basal rebbachisaurid *Histriasaurus* (Dalla Vecchia, 1998) in the upper Hauterivian–lower Barremian of Croatia has been cited as indicative of very early and very rapid northward dispersal of this clade from western Gondwana (southern South America; Carballido et al., 2012; Fanti et al., 2015). Timing of this particular dispersal event was even constrained to the Berriasian–Valanginian time interval (Fanti et al., 2015), which makes it roughly contemporaneous with the record of the Southern Dobrogean carcharodontosaurine. It was also suggested, however, that dispersal of the line leading to *Histriasaurus* was mediated by the northward drift of the Apulian Microplate (= Adria; see Bosselini, 2002), a continental sliver acting as a passive transportation mechanism (‘Noah’s Ark’; ~~KeKenna~~McKenna, 1973) for basal rebbachisaurids after its separation from mainland Africa (e.g., Torcida Fernández-Baldor et al., 2011). Furthermore, the palaeogeographical separation between Africa and Adria (and thus the effective movement of the presumed ark) is considered to be at most an incipient one

1 923 during the Early Cretaceous by Bossellini (2002) and Zarcone et al. (2010), with spatial
2 924 continuity still present between the two landmasses, while deep-water basins continued to
3
4 925 separate Adria from the European Craton. Accordingly, although the presence of
5
6
7 926 *Histriasaurus* can represent a case of northward range extension of rebbachisaurids during
8
9 927 the Berriasian–Valanginian, it took place not strictly speaking into Europe, but only reached
10
11 928 the northernmost extremity of Adria, a northerly peninsular extension of the African
12
13 929 mainland. It was only starting with the Barremian that rebbachisaurids dispersed as far north
14
15 930 as the European cratonic areas, including Iberia and the British Isles (Mannion, 2009;
16
17 931 Mannion et al., 2011; Torcida Fernández-Baldor et al., 2011), a time when faunal
18
19 932 interchanges between Europe and Africa are considered to have been well underway (e.g.,
20
21 933 Gheerbrant and Rage, 2006; Canudo et al., 2009).

22
23
24
25
26 934 Unlike *Histriasaurus*, the taxon represented by UAIC (SCM1) 615 was an inhabitant
27
28 935 of the European mainland. It is thus unclear to what extent the example of rebbachisaurid
29
30 936 range extension into (present-day) Europe during the early Early Cretaceous, as potentially
31
32 937 testified by the discovery of the Croatian taxon, would also be applicable for the Southern
33
34 938 Dobrogean carcharodontosaurine. The available evidence suggests that these two cases are
35
36 939 very different, and that faunal connections during this time interval are not documented
37
38 940 between the African and European cratons as already pointed out by Gheerbrant and Rage
39
40 941 (2006).

41
42
43
44
45 942 Absence of documented faunal interactions weakens support for a scenario of south-
46
47 943 to-north immigration of derived carcharodontosaurines in Europe at the very beginning of the
48
49 944 Cretaceous, and would argue instead for a local, European development to explain the
50
51 945 presence of a Valanginian carcharodontosaurine in Southern Dobrogea. The pre-Barremian
52
53 946 presence of carcharodontosaurids in Europe is also consistent with their appearance in the
54
55 947 Barremian–Aptian fossil record of Eastern Asia, with Europe acting as a stepping stone in the
56
57
58
59
60
61
62
63
64
65

eastward dispersal of the clade. Similarly, the presence of Aptian carcharodontosaurids in North America likely requires the presence of pre-Aptian members of the clade in Europe, since faunal exchanges between these two landmasses are known to have been halted before the Aptian (e.g., Kirkland et al., 1999). Interestingly, it appears that only basal carcharodontosaurids were able to spread into the northern Laurasian landmasses, while the derived carcharodontosaurines dispersed exclusively across the Neo-Tethys, into western Gondwana. The causes of these distribution patterns remain as yet unknown, and further support – in the form on new carcharodontosaurid discoveries from the early-middle part of the Early Cretaceous – is required to better uphold such a scenario.

We finally reiterate that if the Romanian tooth does not belong to a carcharodontosaurine, but instead is artefactually grouping with them in the phylogenetic analysis because of the very incomplete nature of the material, then the traditional story of Carcharodontosaurinae as a product of vicariant evolution driven by the breakup of Pangea will remain strongly supported. However, even in such case UAIC (SCM1) 615 would still record the presence of early-occurring large carcharodontosaurid theropods with a very characteristic carcharodontosaurine-type dentition in the eastern part of the European craton, adding to known early Early Cretaceous theropod (and dinosaur) diversity, and potentially documenting dinosaur faunal provinciality in Europe and worldwide.

6. Conclusions

We re-describe and interpret the affinities of one of the most significant historical dinosaurian specimens of Romania, an isolated but well-preserved theropod tooth from Southern Dobrogea. Our extensive analyses suggest carcharodontosaurid relationships for this tooth, while the available evidence – including novel calcareous nannoplankton sampling – supports its Valanginian age. The Southern Dobrogean theropod tooth represents the oldest record of

Carcharodontosauridae in the Cretaceous, and the second oldest globally, eclipsed only by a collection of isolated specimens from the Upper Jurassic of eastern Africa. As one of the only two known Valanginian dinosaurian occurrences from Central and Eastern Europe, this record advances our understanding of European dinosaur distribution during the early Early Cretaceous, and also fills an important palaeogeographic gap between Western European and Eastern Asian dinosaurian assemblages of the Valanginian.

Based on dental apomorphies, our analyses further identify UAIC (SCM1) 615 as a possible member of Carcharodontosaurinae, a subclade of derived and gigantic carcharodontosaurids formerly known to be restricted to the Albian–Cenomanian of western Gondwana (Africa and South America). If this finding is correct, the Southern Dobrogean specimen documents the emergence of Carcharodontosaurinae earlier than previously recognized, thus also indicating an earlier acquisition of their characteristically large size.

Based on currently known palaeogeographic and chronostratigraphic constraints on the evolution of Carcharodontosauridae, it appears that not only did this clade have a wide distribution, but that crucial events of its evolutionary history such as the emergence of the derived carcharodontosaurines took place north of the Tethys, in cratonic Europe, instead of western Gondwana and as the result of vicariant evolution driven by the Gondwana-Laurasia split, as was formerly suggested. In such a case, instead of endemic evolution the emergence of the western Gondwanan mid-Cretaceous carcharodontosaurines was the result of a north-to-south trans-Tethyan dispersal that took place somewhere between the Valanginian and the Aptian. Recognizing a potential carcharodontosaurine dispersal event from Europe into western Gondwana adds further support for the presence of important palaeogeographic ties between the two realms during the second half of the Early Cretaceous.

Acknowledgements

This research was supported by the National Research Council of Romania (CNCS) grant PN-IIID-PCE-2011-3-0381 and a Sepkoski grant of the Paleontological Society for Z.Cs.-S. S.L.B. is supported by a Marie Curie Career Integration Grant EC630652, the Division of Paleontology of the American Museum of Natural History, and the School of GeoSciences of the University of Edinburgh. He thanks Mátyás Vremir, Radu Totoianu, and Mark Norell for many hours of fun discussion on Romanian fossils, and for supporting his work and travel in Romania. We thank Mihai Brânzilă and Paul Țibuleac (UAIC) for access to the specimen, for allowing us to collect samples for the nannoplankton analyses, and for their help and collegiality during our visit to Iași, as well as Ilie Turculeț for sharing information about the history of the specimen. Mihaela C. Melinte-Dobrinescu has gracefully analyzed the nannoplankton samples derived from UAIC (SCM1) 615; her contribution was essential in assessing the age of the specimen. Finally, we thank the reviewers Eric W.A. Mulder (Denekamp, the Netherlands) and Xabier Pereda-Suberbiola (Bilbao, Spain), as well as Associated Editor Elena Jagt-Yazykova, for their useful comments and suggestions that helped improve ~~a~~previous versions of the manuscript.

References

- Accarie, H., Beaudoin, B., Dejax, J., Friès, G., Michard, J.G., Taquet, P., 1995. Découverte d'un dinosaure théropode nouveau (*Genusaurus sisteronis* n. g., n. sp.) dans l'Albien marin de Sisteron (Alpes-de-Haute-Provence, France) et extension au Crétacé inférieur de la lignée cératosaurienne. *Comptes Rendus de l'Académie des Sciences Paris, Ila* 320, 327–334.
- Allain, R., 2002. Discovery of a megalosaur (Dinosauria, Theropoda) in the Middle Bathonian of Normandy (France) and its implications for the phylogeny of basal Tetanurae. *Journal of Vertebrate Paleontology* 22, 548–563.

- 1022 Allain, R., 2005. The enigmatic theropod dinosaur *Erectopus superbus* (Sauvage, 1882) from
1
2 1023 the Lower Albian of Louppy-le-Chateau (Meuse, France). In: Carpenter, K. (Ed.), The
3
4 1024 Carnivorous Dinosaurs. Indiana University Press, Bloomington, pp. 72–86.
5
6
7 1025 Avram, E., Szasz, L., Antonescu, E., Baltreș, A., Iva, M., Melinte, M., Neagu, T., Rădan, S.,
8
9 1026 Tomescu, C., 1993. Cretaceous terrestrial and shallow marine deposits in northern South
10
11 1027 Dobrogea (SE Romania). Cretaceous Research 14, 265–305.
12
13
14 1028 Avram, E., Costea, I., Dragastan, O., Mușiu, R., Neagu, T., Șindilar, V., Vinogradov, C.,
15
16 1029 1996. Distribution of the Middle-Upper Jurassic and Cretaceous facies in the Romanian
17
18 1030 eastern part of the Moesian Platform. Revue Roumaine de Géologie 39-40, 3–33.
19
20
21 1031 Azevedo, R.P.F. de, Simbras, F.M., Furtado, M.R., Candeiro, C.R.A., Bergqvist, L.P., 2013.
22
23 1032 First Brazilian carcharodontosaurid and other new theropod dinosaur fossils from the
24
25 1033 Campanian-Maastrichtian Presidente Prudente Formation, São Paulo State, southeastern
26
27 1034 Brazil. Cretaceous Research 40, 131–142.
28
29
30 1035 Benson, R.B.J., Barrett, P.M., Powell, H.P., Norman, D.B., 2008. The taxonomic status of
31
32 1036 *Megalosaurus bucklandii* (Dinosauria, Theropoda) from the Middle Jurassic of Oxfordshire,
33
34 1037 UK. Palaeontology 51, 419–424.
35
36
37 1038 Benson, R.B.J., Carrano, M.T., Brusatte, S.L., 2010. A new clade of archaic large-bodied
38
39 1039 predatory dinosaurs (Theropoda: Allosauroidae) that survived to the latest Mesozoic.
40
41
42 1040 Naturwissenschaften 97, 71–78.
43
44
45 1041 Benton, M.J., Cook, E., Grigorescu, D., Popa, E., Tallódi, E., 1997. Dinosaurs and other
46
47 1042 tetrapods in an Early Cretaceous bauxite-filled fissure, northwestern Romania.
48
49
50 1043 Palaeogeography, Palaeoclimatology, Palaeoecology 130, 275–292.
51
52
53 1044 Benton, M.J., Minter, N.J., Posmoșanu, E., 2006. Dwarfing in ornithomimid dinosaurs from the
54
55 1045 Early Cretaceous of Romania. In: Csiki, Z. (Ed.), Mesozoic and Cenozoic Vertebrates and
56
57
58
59
60
61
62
63
64
65

Paleoenvironments; Tributes to the Career of Prof. Dan Grigorescu. Ars Docendi, Bucharest, pp. 79–87.

Benton, M.J., Csiki, Z., Grigorescu, D., Redelstorff, R., Sander, M., Stein, K., Weishampel, D.B., 2010. Dinosaurs and the island rule: the dwarfed dinosaurs from Hațeg Island. *Palaeogeography, Palaeoclimatology, Palaeoecology* 293, 438–454.

Blows, W.T., Honeysett, K., 2014. First Valanginian *Polacanthus foxii* (Dinosauria, Ankylosauria) from England, from the Lower Cretaceous of Bexhill, Sussex. *Proceedings of the Geologists' Association* 125, 233–251.

Bosselini, A., 2002. Dinosaurs “re-write” the geodynamics of the eastern Mediterranean and the paleogeography of the Apulia Platform. *Earth-Science Reviews* 59, 211–234.

Brânzilă, M. (ed.), 2010. *Academicianul Ion Simionescu - savant și dascăl al neamului*. Ed. Universității "Alexandru Ioan Cuza", Iași, 182 pp.

Brusatte, S.L., Sereno, P.C., 2007. A new species of *Carcharodontosaurus* (Dinosauria: Theropoda) from the Cenomanian of Niger and a revision of the genus. *Journal of Vertebrate Paleontology* 27, 902–916.

Brusatte, S.L., Sereno, P.C., 2008. Phylogeny of Allosauroidea (Dinosauria: Theropoda): comparative analysis and resolution. *Journal of Systematic Palaeontology* 6, 155–182.

Brusatte, S. L., Clark, N.D.L., 2015. Theropod dinosaurs from the Middle Jurassic (Bajocian-Bathonian) of Skye, Scotland. *Scottish Journal of Geology* 51, 157-164. doi: 10.1144/sjg2014-022

Brusatte, S., Benson, R.B.J., Carr, T.D., Williamson, T.E., Sereno, P.C., 2007. The systematic utility of theropod enamel wrinkles. *Journal of Vertebrate Paleontology* 27, 1052–1056.

Brusatte, S.L., Benson, R.B.J., Chure, D.J., Xu, X., Sullivan, C., Hone, D.W.E., 2009. The first definitive carcharodontosaurid (Dinosauria: Theropoda) from Asia and the delayed

ascent of tyrannosaurids. *Naturwissenschaften* 96, 1051–1058. Brusatte, S.L., Norell, M.A.,
 Carr, T.D., Erickson, G.M., Hutchinson, J.R., Balanoff, A.M., Bever, G.S., Choiniere, J.N.,
 Makovicky, P.J., Xu, X., 2010a. Tyrannosaur paleobiology: new research on ancient
 exemplar organisms. *Science* 329, 1481–1485.
 Brusatte, S., Chure, D.J., Benson, R.B.J., Xu, X., 2010b. The osteology of *Shaochilong*
maortuensis, a carcharodontosaurid (Dinosauria: Theropoda) from the Late Cretaceous of
 Asia. *Zootaxa* 2334, 1–46.
 Brusatte, S.L., Benson, R.B.J., Xu, X., 2012. A reassessment of *Kelmayisaurus petrolicus*, a
 large theropod dinosaur from the Early Cretaceous of China. *Acta Palaeontologica Polonica*
 57, 65–72.
 Brusatte, S.L., Vremir, M., Csiki-Sava, Z., Turner, A.H., Watanabe, A., Erickson, G.M.,
 Norell, M.A., 2013. The osteology of *Balaur bondoc*, an island-dwelling dromaeosaurid
 (Dinosauria: Theropoda) from the Late Cretaceous of Romania. *Bulletin of the American*
Museum of Natural History 374, 3–100. doi: 10.1206/798.1
 Brusatte, S.L., Lloyd, G.T., Wang, S.C., Norell, M.A., 2014. Gradual assembly of avian body
 plan culminated in rapid rates of evolution across the dinosaur-bird transition. *Current*
Biology 24, 2386–2392.
 Buckland, W., 1824. Notice on the *Megalosaurus* or great fossil lizard of Stonesfield.
Transactions of the Geological Society 21, 390–397.
 Buffetaut, E., Suteethorn, V., 1998. Early Cretaceous dinosaurs from Thailand and their
 bearing on the early evolution and biogeographical history of some groups of Cretaceous
 dinosaurs. In: Lucas, S.G., Kirkland, J.I., Estep, J.W. (Eds.), *Lower and Middle Cretaceous*
Terrestrial Ecosystems, New Mexico Museum of Natural History and Science Bulletin 14,
 pp. 205–210.

- 1095 Buffetaut, E., Suteethorn, V., 2007. A sinraptorid theropod (Dinosauria: Saurischia) from the
1
2 1096 Phu Kradung Formation of northeastern Thailand. Bulletin de la Société Géologique de
3
4
5 1097 France 178, 497–502.
6
7 1098 Buffetaut, E., Suteethorn, V., 2012. A carcharodontid theropod (Dinosauria, Saurischia) from
8
9
10 1099 the Sao Khua Formation (Early Cretaceous, Barremian) of Thailand. In: Royo-Torres, R.,
11
12 1100 Gascó, F., Alcalá, L. (Eds.), 10th Annual Meeting of the European Association of Vertebrate
13
14 1101 Palaeontologists. ¡Fundamental! 20, [Teruel](#), pp. 27–30.
15
16
17 1102 Calvo, J.O., Coria, A., 1998. New specimen of *Giganotosaurus carolinii* (Coria & Salgado,
18
19 1103 1995), supports it as the largest theropod ever found. Gaia 15, 117–122.
20
21
22 1104 Calvo, J.O., Gandossi, P., Porfiri, J.D., 2006. Dinosaur faunal replacement during
23
24 1105 Cenomanian times in Patagonia, Argentina. In: Evans, S.E., Barrett, P.M. (Eds.), 9th
25
26 1106 Mesozoic Terrestrial Ecosystems and Biota, [Manchester](#), UK, pp. 17–20.
27
28
29 1107 Canale, J.I., Scanferla, C.A., Agnolin, F.L., Novas, F.E., 2009. New carnivorous dinosaur
30
31 1108 from the Late Cretaceous of NW Patagonia and the evolution of abelisaurid theropods.
32
33
34 1109 Naturwissenschaften 96, 409–414.
35
36 1110 Canale, J.I., Novas, F.E., Pol, D., 2015. Osteology and phylogenetic relationships of
37
38
39 1111 *Tyrannotitan chubutensis* Novas, de Valais, Vickers-Rich and Rich, 2005 (Theropoda:
40
41 1112 Carcharodontosauridae) from the Lower Cretaceous of Patagonia, Argentina. Historical
42
43 1113 Biology 27, 1–32. doi: 10.1080/08912963.2013.861830.
44
45
46 1114 Candeiro, C.R.A., Fanti, F., Therrien, F., Lamanna, M.C., 2011. Continental fossil vertebrates
47
48
49 1115 from the mid-Cretaceous (Albian-Cenomanian) Alcântara Formation, Brazil, and their
50
51 1116 relationship with contemporaneous faunas from North Africa. Journal of African Earth
52
53 1117 Sciences 60, 79–92.
54
55
56 1118 Canudo, J.I., Ruiz-Omeñaca, J.I., 2003. Los restos directos de dinosaurios terópodos
57
58 1119 (excluyendo Aves) en España. Ciencias de la Tierra 26, 347–373.
59
60
61
62
63
64
65

- Canudo, J.I., Filippi, L.S., Salgado, L., Garrido, A.C., Cerda, I.A., Garcia, R., Otero, A.,
2008. Theropod teeth associated with a sauropod carcass in the Upper Cretaceous (Plottier
Formation) of Rincón de los Sauces. In: Colectivo Arqueológico y Paleontológico de Salas de
los Infantes (Ed.), Actas de las IV Jornadas Internacionales sobre Paleontología de
Dinosaurios y su Entorno, Salas de los Infantes, Burgos, pp. 321–330.
- Canudo, J.I. Barco, J.L., Pereda Suberbiola, X., Ruiz-Omeñaca, J.I., Salgado, L., Torcida
Fernández-Baldor, F., Gasulla, J.M., 2009. What Iberian dinosaurs reveal about the bridge
said to exist between Gondwana and Laurasia in the Early Cretaceous. Bulletin de la Société
Géologique de France, 180, 5–11.
- Carballido, J.L., Salgado, L., Pol, D., Canudo, J.I., Garrido, A., 2012. A new basal
rebbachisaurid (Sauropoda, Diplodocoidea) from the Early Cretaceous of the Neuquén Basin;
evolution and biogeography of the group. Historical Biology 24, 631–654.
- Carrano, M.T., Sampson, S.D., 2008. The phylogeny of Ceratosauria (Dinosauria:
Theropoda). Journal of Systematic Palaeontology 6, 183–236.
- Carrano, M.T., Benson, R.B.J., Sampson, S.D., 2012. The phylogeny of Tetanurae
(Dinosauria: Theropoda). Journal of Systematic Palaeontology 10, 211–300.
- Carvalho, I.S., Campos, A.C.A., Nobre, P.H., 2005. *Baurusuchus salgadoensis*, a new Crocodylomorpha
from the Bauru Basin (Cretaceous), Brazil. Gondwana Research 8, 11–30.
- Casal, G., Candeiro, C.R.A., Martinez, R., Ivany, E., Ibiricu, L., 2009. Dientes de Theropoda
(Dinosauria: Saurischia) de la Formación Bajo Barreal, Cretácico Superior, Provincia del
Chubut, Argentina. Géobios 42, 553–560.
- Cau, A., Dalla Vecchia, F.M., Fabbri, M., 2013. A thick-skulled theropod (Dinosauria,
Saurischia) from the Upper Cretaceous of Morocco with implications for carcharodontosaurid
cranial evolution. Cretaceous Research 40, 251–260.

- Choiniere, J.N., Forster, C.A., De Klerk, W.J., 2012. New information on *Nqwebasaurus*
thwazi, a coelurosaurian theropod from the Early Cretaceous Kirkwood Formation in South
Africa. *Journal of African Earth Sciences* 71–72, 1–17.
- Chure, D.J., Manabe, M., Tanimoto, M., Tomida, Y., 1999. An unusual theropod tooth from
the Mifune Group (Late Cenomanian to Early Turonian), Kumamoto, Japan. In: Tomida, Y.,
Rich, T.H., Vickers-Rich, P. (Eds.), *Proceedings of the Second Gondwanan Dinosaur*
Symposium. National Science Museum Monographs 15, Tokyo, pp. 291–296.
- Cifelli, R.L., Davis, B.M., Sames, B., 2014. Earliest Cretaceous mammals from the western
United States. *Acta Palaeontologica Polonica* 59, 31–52. doi:10.4202/app.2012.0089.
- Cobos, A., Royo-Torres, R., Luque, L., Alcalá, L., Mampel, L., 2010. An Iberian stegosaurs
paradise: The Villar del Arzobispo Formation (Tithonian–Berriasian) in Teruel (Spain).
Palaeogeography, Palaeoclimatology, Palaeoecology 293, 223–236.
- Cobos, A., Lockley, M.G., Gascó, F., Royo–Torres, R., Alcalá, L., 2014. Megatheropods as
apex predators in the typically Jurassic ecosystems of the Villar del Arzobispo Formation
(Iberian Range, Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology* 399, 31–41. doi:
10.1016/j.palaeo.2014.02.008.
- Codrea, V., Vremir, M., Jipa, C., Godefroit, P., Csiki, Z., Smith, T., Fărcaș, C., 2010. More
than just Nopcsa’s Transylvanian dinosaurs: A look outside the Hațeg Basin.
Palaeogeography, Palaeoclimatology, Palaeoecology 293, 391–405. doi:
10.1016/j.palaeo.2009.10.027.
- Codrea, V., Godefroit, P., Smith, T., 2012. First discovery of Maastrichtian (latest
Cretaceous) terrestrial vertebrates in Rusca Montană Basin (Romania). In: Godefroit, P. (Ed.)
Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems. Indiana University Press,
Bloomington, pp. 570–581.

- Colbert, E.H., 1946. *Sebecus*, representative of a peculiar suborder of fossil Crocodilia from Patagonia. Bulletin of the American Museum of Natural History 87(4), 217–270.
- Coria, R.A., Salgado, L., 1995. A new giant carnivorous dinosaur from the Cretaceous of Patagonia. Nature 377, 224–226.
- Coria, R.A., Salgado, L., 2005. Mid-Cretaceous turnover of saurischian dinosaur communities: evidence from the Neuquén Basin. In: Veiga, G.D., Spalletti, L.A., Howell, J.A., Schwartz, E. (Eds.), The Neuquén Basin, Argentina: a case study in sequence stratigraphy and basin dynamics. Geological Society, London, Special Publications 252, pp. 317–327.
- Coria, R.A., Currie, P.J., 2006. A new carcharodontosaurid (Dinosauria, Theropoda) from the Upper Cretaceous of Argentina. Geodiversitas 28, 71–118.
- Covacef, Z., 1995. Pionieri ai culturii românești în Dobrogea; Ioan Cotovu și Vasile Cotovu. Analele Dobrogei I(1), 127–134.
- Császár, G., Kollányi, K., Lantos, M., Lelkes, G. and Tardine Filác, E., 2000. A Hidasivölgyi Márga Formáció kora és képződési környezete. Földtani Közlöny 130(4), 695–723.
- Csiki, Z., Grigorescu, D., 1998. Small theropods of the Late Cretaceous of the Hațeg Basin (Western Romania) - an unexpected diversity at the top of the food chain. Oryctos 1, 87–104.
- Csiki, Z., Vremir, M., Brusatte, S.L., Norell, M.A., 2010. An aberrant island-dwelling theropod dinosaur from the Late Cretaceous of Romania. Proceedings of the National Academy of Sciences 107, 15357–15361.
- Csiki-Sava, Z., Codrea, V., Vasile, Ș., 2013. Early Cretaceous dinosaur remains from Dobrogea (southeastern Romania). In: Picot, L. (Ed.), Abstracts, 11th Annual Meeting of the European Association of Vertebrate Palaeontologists, Villers-sur-Mer, France, pp. 28.

- 1192 Csiki-Sava, Z., Buffetaut, E., Ósi, A., Pereda-Suberbiola, X., Brusatte, S.L., 2015. Island life
1193 in the Cretaceous - faunal composition, biogeography, evolution, and extinction of land-
1194 living vertebrates on the Late Cretaceous European archipelago. *Zookeys* 469, 1–161. doi:
1195 10.3897/zookeys.469.8439.
- 1196 Csontos, L., Vörös, A., 2004. Mesozoic plate tectonic reconstruction of the Carpathian
1197 region. *Palaeogeography, Palaeoclimatology, Palaeoecology* 210, 1–56. doi:
1198 10.1016/j.palaeo.2004.02.033.
- 1199 Currie, P.J., Carpenter, K., 2000. A new specimen of *Acrocanthosaurus atokensis*
1200 (Theropoda, Dinosauria) from the Lower Cretaceous Antlers Formation (Lower Cretaceous,
1201 Aptian) of Oklahoma, USA. *Geodiversitas* 22, 207–246.
- 1202 Currie, P.J., Rigby, J.K., Sloan, R.E., 1990. Theropod teeth from the Judith River Formation
1203 of southern Alberta, Canada. In: Carpenter, K., Currie, P.J. (Eds.), *Dinosaur Systematics:
1204 Approaches and Perspectives*. Cambridge University Press, [Cambridge, Boston](#), pp. 107–125.
- 1205 Dalla Vecchia, F.M., 1998. Remains of Sauropoda (Reptilia, Saurischia) in the Lower
1206 Cretaceous (Upper Hauterivian/Lower Barremian) limestones of SW Istria (Croatia).
1207 *Geologia Croatica* 5, 105–134.
- 1208 Dames, W., 1884. *Megalosaurus dunkeri*. *Sitzungsberichte Gesellschaft Naturforschender
1209 Freunde zu Berlin* 1884, 186–188.
- 1210 De Klerk, W.J., Forster, C.A., Sampson, S.D., Chinsamy, A., Ross, C.F., 2000. A new
1211 coelurosaurian dinosaur from the Early Cretaceous of South Africa. *Journal of Vertebrate
1212 Paleontology* 20, 324–332.
- 1213 D’Emic, M.D., Melstrom, K.M., Eddy, D.R., 2012. Paleobiology and geographic range of the
1214 large-bodied Cretaceous theropod dinosaur *Acrocanthosaurus atokensis*. *Palaeogeography,
1215 Palaeoclimatology, Palaeoecology* 333–334, 13–23.

- 1216 Dercourt, J., Gaetani, M., Vrielynck, B., Barrier, E., Biju-Duval, B., Brunet, M., Cadet, J.P.,
1
2 1217 Crasquin, S., Săndulescu, M., Eds. (2000) Atlas Peri-Tethys Palaeogeographical Maps.
3
4 1218 CCGM/CGMW, Paris, 269 pp.
5
6
7 1219 Dinu, C., Grădinaru, E., Stoica, M., Diaconescu, V., 2007. Dobrogea 2007 Field Trip
8
9
10 1220 Preparation and Assistance. University of Bucharest, 123 pp.
11
12 1221 Dragastan, O., Neagu, T., Bărbulescu, A., Pană, I., 1998. Jurasicul și Cretacicul din Dobrogea
13
14 1222 Centrală și de Sud. Bucharest, 249 pp.
15
16
17 1223 Dragastan, O.N., Antoniadu, C., Stoica, M., 2014. Biostratigraphy and zonation of the Lower
18
19 1224 Cretaceous succession from Cernavodă-lock section, South Dobrogea, eastern part of the
20
21 1225 Moesian Platform (Romania). Carpathian Journal of Earth and Environmental Sciences 9(1),
22
23 1226 231–260.
24
25
26 1227 Dyke, G.J., Benton, M.J., Posmoșanu, E., Naish, D., 2011. Early Cretaceous (Berriasian)
27
28 1228 birds and pterosaurs from the Cornet bauxite mine, Romania. Palaeontology 54, 79–95.
29
30
31 1229 Eberth, D.A., Brinkman, D.B., Chen, P.-J., Yuan, F.-T., Wu, X.-C., Li, G., Cheng, XianX.-
32
33 Shen, C., 2001. Sequence stratigraphy, paleoclimate patterns, and vertebrate fossil
34 1230 preservation in Jurassic–Cretaceous strata of the Juggar Basin, Xinjiang Autonomous Region,
35
36 1231 People’s Republic of China. Canadian Journal of Earth Sciences 38, 1627–1644.
37
38
39 1232 Eddy, D.R., Clarke, J.A., 2011. New information on the cranial anatomy of
40
41 1233 *Acrocanthosaurus atokensis* and its implications for the phylogeny of Allosauroidea
42
43 1234 (Dinosauria: Theropoda). PLoS ONE 6(3), e17932. doi:10.1371/journal.pone.0017932.
44
45 1235 Evans, S.E., Matsumoto, R., 2015. An assemblage of lizards from the Early Cretaceous of
46
47 1236 Japan. Palaeontologia Electronica 18.2.36A, 1–36.
48
49 1237 Ezcurra, M.D., Agnolín, F.L., 2012. A new global palaeobiogeographical model for the Late
50
51 1238 Mesozoic and Early Tertiary. Systematic Biology 61, 553–566. doi:10.1093/sysbio/syr115.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- Fanti, F., Cau, A., Cantelli, L., Hassine, M., Auditore, M., 2015. New information on *Tataouinea hannibalis* from the Early Cretaceous of Tunisia and implications for the tempo and mode of rebbachisaurid sauropod evolution. PLoS ONE 10(4), e0123475. doi:10.1371/journal.pone.0123475.
- Galton, P.M., 2009. Notes on Neocomian (Lower Cretaceous) ornithomimid dinosaurs from England - *Hypsilophodon*, *Valdosaurus*, "*Camptosaurus*", "*Iguanodon*" - and referred specimens from Romania and elsewhere. Revue de Paléobiologie 28, 211–273.
- Gardner, J.D., Evans, S.E., Sigogneau-Russell, D., 2003. New albanerpetontid amphibians from the Early Cretaceous of Morocco and Middle Jurassic of England. Acta Palaeontologica Polonica 48, 301–319.
- Gasca, J.M., Canudo, J.I., Moreno-Azanza, M., 2014. A large-bodied theropod (Tetanurae: Carcharodontosauria) from the Mirambel Formation (Barremian) of Spain. Neues Jahrbuch für Geologie und Paläontologie Abhandlungen 273, 13–23. doi: 10.1127/0077-7749/2014/0413.
- Gheerbrant, E., Rage, J.-C., 2006. Paleobiogeography of Africa: How distinct from Gondwana and Laurasia? Palaeogeography, Palaeoclimatology, Palaeoecology 241, 224–246. doi:10.1016/j.palaeo.2006.03.016.
- Gishlick, A.D., Gauthier, J.A., 2007. On the manual morphology of *Compsognathus longipes* and its bearing on the diagnosis of Compsognathidae. Zoological Journal of the Linnean Society 149, 569–581.
- Goloboff, P. A., Farris, J. S., Nixon, K. C., 2008. TNT, a free program for phylogenetic analysis. Cladistics 24, 774–786.
- Gradstein, F.M., Ogg, J.G., Schmitz, M.D., Ogg, G.M. (Eds.), 2012. The Geologic Time Scale 2012. Elsevier, [Amsterdam](#), 1144 pp.

- Grigorescu, D., 1992. Nonmarine Cretaceous Formations of Romania. In: Matter, N.J., Chen, P.-J. (Eds.), Aspects of Nonmarine Cretaceous Geology. China Ocean Press, Beijing, pp. 142–164.
- Grigorescu, D., 2003. Dinosaurs of Romania. *Comptes rendus Palevol* 2, 97–101.
- Grigorescu, D., 2010. The Latest Cretaceous fauna with dinosaurs and mammals from the Hațeg Basin — A historical overview. *Palaeogeography, Palaeoclimatology, Palaeoecology* 293, 271–282.
- Hammer, O., Harper, D.A.T., Ryan, P.D., 2001. Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4, 1–9.
- Harris, J.D., 1998. A reanalysis of *Acrocanthosaurus atokensis*, its phylogenetic status, and paleobiogeographic implications, based on a new specimen from Texas. *New Mexico Museum of Natural History and Science Bulletin* 13, 1–75.
- Hendrickx, C., Mateus, O., 2014. Abelisauridae (Dinosauria: Theropoda) from the Late Jurassic of Portugal and dentition-based phylogeny as a contribution for the identification of isolated theropod teeth. *Zootaxa* 3751(1), 1–74.
- Hendrickx, C., Mateus, O., Araújo, R., 2015a. The dentition of megalosaurid theropods. *Acta Palaeontologica Polonica* 60, 627–642. doi:10.4202/app.00056.2013.
- Hendrickx, C., Mateus, O., Araújo, R., 2015b. A proposed terminology of theropod teeth (Dinosauria, Saurischia). *Journal of Vertebrate Paleontology* 35(5): e982797. doi: 10.1080/02724634.2015.982797.
- Holtz, T.R., Jr., Molnar, R.E., Currie, P.J., 2004a. Basal Tetanurae. In: Weishampel, D.B., Dodson, P., Osmólska, H. (Eds.), *The Dinosauria*. Second Edition. University of California Press, ~~Berkeley, Los Angeles, London~~~~Berkeley~~, pp. 71–110.
- Holtz, T.R., Chapman, R.E., Lamanna, M.C., 2004b. Mesozoic biogeography of Dinosauria. In: Weishampel, D.B., Dodson, P., Osmólska, H. (Eds.), *The Dinosauria*. Second Edition.

- University of California Press, ~~Berkeley, Los Angeles, London~~~~Berkeley, Los Angeles~~, pp. 627–642.
- Huene, F. von, 1923. Carnivorous Saurischia in Europe since the Triassic. Bulletin of the Geological Society of America 34, 449–458.
- Hippolyte, J.-C., 2002. Geodynamics of Dobrogea (Romania): new constraints on the evolution of the Tornquist–Teisseyre Line, the Black Sea and the Carpathians. Tectonophysics 357, 33–53.
- Ionesi, L., 1994. Geologia unităților de platformă și a orogenului Nord-Dobrogean. Ed. Tehnică, Bucharest, 280 pp.
- Jerzykiewicz, T., Russell, D.A., 1991. Late Mesozoic stratigraphy and vertebrates of the Gobi Basin. Cretaceous Research 12, 345–377.
- Jurcsák, T., 1982. Occurences nouvelles des Sauriens mésozoïques de Roumanie. Vertebrata Hungarica 21, 175–184.
- Jurcsák, T., Popa, E., 1979. Dinozaurieni ornithopozi din bauxitele de la Cornet (Munții Pădurea Craiului). Nymphaea 7, 37–75.
- Jurcsák, T., Popa, E., 1983. La faune de dinosauriens du Bihor (Roumanie). In: Buffetaut, E., Mazin, J.M., Salmon, E. (Eds.), Actes du Symposium Paléontologique Georges Cuvier. Le Serpenteaire, Montbéliard, pp. 325–335.
- Kellner, A.W.A., Pinheiro, A.E.P., Campos, D.A., 2014. A new sebecid from the Paleogene of Brazil and the crocodyliform radiation after the K–Pg boundary. PLoS ONE 9(1), e81386. doi:10.1371/journal.pone.0081386.
- Kirkland, J.I., Cifelli, R.L., Britt, B.B., Burge, D.L., DeCourten, F.L., Eaton, J.G., Parrish, J.M., 1999. Distribution of vertebrate faunas in the Cedar Mountain Formation, east-central Utah. Utah Geological Survey Miscellaneous Publication 99-1, 201–217.

- Knoll, F., Ruiz-Omeñaca, J.I., 2009. Theropod teeth from the basalmost Cretaceous of Anoual (Morocco) and their palaeobiogeographical significance. *Geological Magazine* 146, 602–616.
- Kusuhashi, N., Matsumoto, A., Murakami, M., Tagami, T., Hirata, T., Iizuka, T., Handa, T., Matsuoka, H., 2006. Zircon U–Pb ages from tuff beds of the upper Mesozoic Tetori Group in the Shokawa district, Gifu Prefecture, central Japan. *The Island Arc* 15, 378–390.
- Lacasa Ruiz, A., 1989. Nuevo genero de ave fosil del yacimiento Neocomiense del Montsec (Provincia de Lerida, España). *Estudios geológicos* 45(5-6), 417–425.
- Larson, D.W., Currie, P.J., 2013. Multivariate analyses of small theropod dinosaur teeth and implications for paleoecological turnover through time. *PLoS ONE* 8(1), e54329. doi:10.1371/journal.pone.0054329.
- Le Loeuff, J., Buffetaut, E., 1995. The evolution of Late Cretaceous non-marine vertebrate faunas in Europe. In: Sun, A.-L., Wang, Y.-Q. (Eds.), *Sixth Symposium on Mesozoic Terrestrial Ecosystems and Biota, Short Papers*. China Ocean Press, Beijing, pp. 181–184.
- Le Loeuff, J., Lang, E., Cavin, L., Buffetaut, E., 2012. Between Tendaguru and Bahariya: on the age of the Early Cretaceous dinosaur sites from the Continental Intercalaire and other African formations. *Journal of Stratigraphy* 36, 486–502.
- Lucas, F.A., 1901. A new dinosaur, *Stegosaurus marshi*, from the Lower Cretaceous of South Dakota. *Proceedings of the United States National Museum* 23(1224), 591–592.
- Lucas, S.G., 2006. The *Psittacosaurus* biochron, Early Cretaceous of Asia. *Cretaceous Research* 27, 189–198.
- Lucas, S.G., Estep, J.W., 1998. Vertebrate biostratigraphy and biochronology of the Cretaceous of China. In: Lucas, S.G., Kirkland, J.I., Estep, J.W. (Eds.), *Lower and Middle Cretaceous Terrestrial Ecosystems*. New Mexico Museum of Natural History and Science Bulletin 14, pp. 1–20.

- Lü, J.-C., Xu, L., Pu, H.-Y., Jia, S.-H., Azuma, Y., Chang, H.-L., Zhang, J.-M., ~~2014~~2016.
Paleogeographical significance of carcharodontosaurid teeth from the late Early Cretaceous
of Ruyang, Henan Province of central China. *Historical Biology*, 28, 8–13. doi:
10.1080/08912963.2014.947287.
- Manabe, M., Hasegawa, Y., 1995. Diapsid fauna and its paleobiogeographical implication,
the Neocomian section of the Tetori Group. In: Sun, A., Wang, Y. (Eds.), *Sixth Symposium
on Mesozoic Terrestrial Ecosystems and Biota, Short Papers*, China Ocean Press, Beijing,
pp. 179.
- Mannion, P.D., 2009. A rebbachisaurid sauropod from the Lower Cretaceous of the Isle of
Wight, England. *Cretaceous Research* 30, 521–526.
- Mannion, P.D., Upchurch, P., Hutt, S., 2011. New rebbachisaurid (Dinosauria: Sauropoda)
material from the Wessex Formation (Barremian, Early Cretaceous), Isle of Wight, United
Kingdom. *Cretaceous Research* 32, 774–780.
- Matsukawa, M., Ito, M., Nishida, N., Koarai, K., Lockley, M.G., Nichols, D.J., 2006. The
Cretaceous Tetori biota in Japan and its evolutionary significance for terrestrial ecosystems in
Asia. *Cretaceous Research* 27, 199–225.
- McDonald, A.T., 2011. The taxonomy of species assigned to *Camptosaurus* (Dinosauria:
Ornithopoda). *Zootaxa* 2783, 52–68.
- McKenna, M.C., 1973. Sweepstakes, filters, corridors, Noah's Arks, and Beached Viking
Funeral Ships in palaeogeography. In: Tarling, D.H., Runcorn, S.K. (Eds.), *Implications of
Continental Drift to the Earth Sciences*. Academic Press, New York, pp. 295–308.
- Mo, J.-Y., Huang, C.-L., Xie, S.-W., Buffetaut, E., 2014. A megatheropod tooth from the
Early Cretaceous of Fusui, Guangxi, Southern China. *Acta Geologica Sinica (English
Edition)* 88, 6–12.

- 1362 Molnar, R.E., 1990. Problematic Theropoda: "Carnosaurs". In: Weishampel, D.B., Dodson,
1
2 1363 P., Osmólska, H. (Eds.), The Dinosauria. University of California Press, Berkeley, Los
3
4 Angeles, Oxford, pp. 306–317.
5 1364
6
7 1365 Mutihac, V., Mutihac, G., 2010. The geology of Romania, within the Central East European
8
9 geosstructural context. Ed. Didactică și Pedagogică, Bucharest, 690 pp.
10 1366
11
12 1367 Neagu, T., Dragastan, O., Csiki, Z., 1997. Early Cretaceous shelf paleocommunities of
13
14 1368 Cernavodă (South Dobrogea, SE Romania). Acta Palaeontologica Romaniae 1, 28–36.
15
16
17 1369 Nopcsa, F., 1902. Notizen über cretacische Dinosaurier. Sitzungsberichte der Kaiserlichen
18
19 1370 Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe 111, 93–114.
20
21
22 1371 Nopcsa, F., 1923. On the geological importance of the primitive reptilian fauna of the
23
24 1372 uppermost Cretaceous of Hungary; with a description of a new tortoise (*Kallokibotium*).
25
26
27 1373 Quarterly Journal of the Geological Society of London 79, 100–116.
28
29 1374 Norman, D.B., 2010. A taxonomy of iguanodontians (Dinosauria: Ornithopoda) from the
30
31 lower Wealden Group (Cretaceous: Valanginian) of southern England. Zootaxa 2489, 47–66.
32 1375
33
34 1376 Norman, D.B., 2013. On the taxonomy and diversity of Wealden iguanodontian dinosaurs
35
36 1377 (Ornithischia: Ornithopoda). Revue de Paléobiologie 32(2), 385–404.
37
38
39 1378 Novas, F.E., de Valais, S., Vickers-Rich, P.A., Rich, T.H., 2005. A large Cretaceous theropod
40
41 1379 from Patagonia, Argentina, and the evolution of carcharodontosaurids. Naturwissenschaften
42
43 1380 92, 226–230.
44
45
46 1381 Novas, F.E., Agnolín, F.L., Ezcurra, M.D., Porfiri, J., Canale, J.I., 2013. Evolution of the
47
48 carnivorous dinosaurs during the Cretaceous: The evidence from Patagonia. Cretaceous
49 1382
50
51 1383 Research 45, 174–215.
52
53
54 1384 Ortega, F., Escaso, F., Sanz, J.L., 2010. A bizarre, humped Carcharodontosauria (Theropoda)
55
56 1385 from the Lower Cretaceous of Spain. Nature 467, 203–206.
57
58
59
60
61
62
63
64
65

- 1386 Ósi, A., Rabi, M., Makádi, L., Szentesi, Z., Botfalvai, G., Gulyás, P., 2012. The Late
1387 Cretaceous continental vertebrate fauna from Iharkút (western Hungary): a review. In:
1388 Godefroit, P. (Ed.), *Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems*.
1389 Indiana University Press, Bloomington, pp. 533–569.
- 1390 Ósi, A., Codrea, V., Prondvai, E., Csiki-Sava, Z., 2014. New ankylosaurian material from the
1391 Upper Cretaceous of Transylvania. *Annales de Paléontologie* 100, 257–271. doi:
1392 10.1016/j.annpal.2014.02.001.
- 1393 Ósi, A., Rabi, M., Makádi, L., 2015. An enigmatic crocodyliform tooth from the bauxites of
1394 western Hungary suggests hidden mesoeucrocodylian diversity in the Early Cretaceous
1395 European archipelago. *PeerJ* 3, e1160. doi:10.7717/peerj.1160.
- 1396 Paolillo, A., Linares, O.J., 2007. Nuevos cocodrilos Sebecosuchia del Cenozoico
1397 Suramericano (Mesosuchia: Crocodylia). *Paleobiologia Neotropical* 3, 1–25.
- 1398 Pereda-Suberbiola, X., Ruiz-Omeñaca, J.I., Fernandez-Baldor, F.T., Maisch, M.W., Huerta,
1399 P., Contreras, R., Izquierdo, L.A., Huerta, D.M., Montero, V.U., Welle, J., 2011. A tall-
1400 spined ornithomimid dinosaur from the Early Cretaceous of Salas de los Infantes (Burgos,
1401 Spain). *Comptes Rendus Palevol* 10, 551–558.
- 1402 Pereda-Suberbiola, X., Ruiz-Omeñaca, J.I., Canudo, J.I., Torcida, F., Sanz, J.L., 2012.
1403 Dinosaur faunas from the Early Cretaceous (Valanginian–Albian) of Spain. In: Godefroit, P.
1404 (Ed.), *Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems*. ~~Life of the Past~~.
1405 Indiana University Press, Bloomington, pp. 378–407.
- 1406 Pérez-Moreno, B.P., Sanz, J.L., Sudre, J., Sigé, B., 1993. A theropod dinosaur from the
1407 Lower Cretaceous of southern France. *Revue de Paléobiologie*, Volume spéciale 7, 173–188.
- 1408 Pol, D., Powell, J.E., 2011. A new sebecid mesoeucrocodylian from the Rio Loro Formation
1409 (Palaeocene) of north-western Argentina. *Zoological Journal of the Linnean Society* 163, S7–
1410 S36.

- 1411 [Porfiri, J.D., Novas, F.E., Calvo, J.O., Agnolín, F.L., Ezcurra, M.D., Cerda, I.A., 2014.](#)
1412 [Juvenile specimen of *Megaraptor* \(Dinosauria, Theropoda\) sheds light about tyrannosauroid](#)
1413 [radiation. *Cretaceous Research* 51, 35–55. doi:10.1016/j.cretres.2014.04.007.](#)
- 1414 Posmoşanu, E., 2003. Iguanodontian dinosaurs from the Lower Cretaceous bauxite site from
1415 Romania. *Acta Palaeontologica Romaniae* 4, 431–439.
- ~~1416 Porfiri, J.D., Novas, F.E., Calvo, J.O., Agnolín, F.L., Ezcurra, M.D., Cerda, I.A., 2014.~~
~~1417 Juvenile specimen of *Megaraptor* (Dinosauria, Theropoda) sheds light about tyrannosauroid~~
~~1418 radiation. *Cretaceous Research*, 51, 35–55. doi:10.1016/j.cretres.2014.04.007.~~
- 1419 Rabi, M., Sebök, N., 2015. A revised Eurogondwana model: Late Cretaceous notosuchian
1420 crocodyliforms and other vertebrate taxa suggest the retention of episodic faunal links
1421 between Europe and Gondwana during most of the Cretaceous. *Gondwana Research* 28,
1422 1197–1211. doi:10.1016/j.gr.2014.09.015.
- 1423 Racey, A., 2009. Mesozoic red bed sequences from SE Asia and the significance of the
1424 Khorat Group of NE Thailand. In: Buffetaut, E., Cuny, G., Le Loeuff, J., Suteethorn, V.
1425 (Eds.), Late Palaeozoic and Mesozoic Ecosystems in SE Asia. ~~The~~ Geological Society,
1426 London, Special Publications s 315, pp. 41–67. doi: 10.1144/SP315.5.
- 1427 Racey, A., Goodall, J.G.S., 2009. Palynology and stratigraphy of the Mesozoic Khorat Group
1428 red bed sequences from Thailand. In: Buffetaut, E., Cuny, G., Le Loeuff, J., Suteethorn, V.
1429 (Eds.), Late Palaeozoic and Mesozoic Ecosystems in SE Asia. ~~The~~ Geological Society ~~of,~~
1430 London, Special Publications s 315, pp. 69–83. doi: 10.1144/SP315.6.
- 1431 Rauhut, O.W.M., 2011. Theropod dinosaurs from the Late Jurassic of Tendaguru (Tanzania).
1432 *Special Papers in Palaeontology* 86, 195–239.
- 1433 Rauhut, O.W.M., Werner, C., 1995. First record of the family Dromaeosauridae (Dinosauria:
1434 Theropoda) in the Cretaceous of Gondwana (Wadi Milk Formation, northern Sudan).
1435 *Paläontologische Zeitschrift* 69(3/4), 475–489.

- 1436 Richter, U., Mudroch, A., Buckley, L.G., 2013. Isolated theropod teeth from the Kem Kem
1
2 1437 Beds (Early Cenomanian) near Taouz, Morocco. *Paläontologische Zeitschrift* 87, 291–309.
3
4 1438 Royo-Torres, R., Cobos, A., Luque, L., Aberasturi, A., Espilez, E., Fierro, I., Gonzales, A.,
5
6 1439 Mampel, L., Alcalá, L., 2009. High European sauropod dinosaur diversity during Jurassic–
7
8 1440 Cretaceous transition in Riodeva (Teruel, Spain). *Palaeontology* 52, 1009–1027.
9
10 1441 Sachs, S., Hornung, J.J., 2013. Ankylosaur remains from the Early Cretaceous (Valanginian)
11
12 1442 of Northwestern Germany. *PLoS ONE* 8(4), e60571. doi:10.1371/journal.pone.0060571.
13
14 1443 Salgado, L., Canudo, J.I., Garrido, A.C., Ruiz-Omeñaca, J.I., Garcia, R.A., de la Fuente,
15
16 1444 M.S., Barco, J.L., Bollati, R., 2009. Upper Cretaceous vertebrates from El Anfiteatro area,
17
18 1445 Río Negro, Patagonia, Argentina. *Cretaceous Research* 30, 767–784.
19
20 1446 Sames, B., Cifelli, R.L., Schudack, M.E., 2010. The nonmarine Lower Cretaceous of the
21
22 1447 North American Western Interior foreland basin: New biostratigraphic results from ostracod
23
24 1448 correlations and early mammals, and their implications for paleontology and geology of the
25
26 1449 basin—An overview. *Earth-Science Reviews* 101, 207–224.
27
28 1450 Sauvage, H.E., 1876. Notes sur les reptiles fossiles. *Bulletin de la Société Géologique de*
29
30 1451 France 4, 435–442.
31
32 1452 Sauvage, H.E., 1882. Recherches sur les reptiles trouves dans le Gault de l'est du bassin de
33
34 1453 Paris. *Mémoires de la Société Géologique de France* 2, 1–42.
35
36 1454 Săndulescu, M., 1984. *Geotectonica României*. Ed. Tehnică, Bucharest, 329 pp.
37
38 1455 Seghedi, A., 2001. The North Dobrogea orogenic belt (Romania): a review. In: Ziegler, P.A.,
39
40 1456 Cavazza, W., Robertson, A.H.F., Crasquin-Soleau, S. (Eds.), *Peri-Tethys Memoir 6: Peri-*
41
42 1457 *Tethyan Rift/Wrench Basins and Passive Margins*. *Mémoires de la Musée National d'Histoire*
43
44 1458 *Naturelle*, Paris, pp. 237–257.
45
46 1459 Sereno, P.C., 1999. Dinosaurian biogeography: vicariance, dispersal and regional extinction.
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- In: Tomida, Y., Rich, T.H., Vickers-Rich, P. (Eds.), Proceedings of the Second Gondwanan
Dinosaur Symposium. National Science Museum Monographs 15, Tokyo, pp. 249–257.
- Sereno, P.C., Brusatte, S.L., 2008. Basal abelisaurid and carcharodontosaurid theropods from
the Lower Cretaceous Elrhaz Formation of Niger. *Acta Palaeontologica Polonica* 53, 15–46.
- Sereno, P.C., Dutheil, D.B., Iarochene, M., Larsson, H.C.E., Lyon, G.H., Magwene, P.M.,
Sidor, C.A., Varricchio, D.J., Wilson, J.A., 1996. Predatory dinosaurs from the Sahara and
Late Cretaceous faunal differentiation. *Science* 272, 986–991.
- Shen, Y.B., Mateer, N.J., 1992. An outline of the Cretaceous system in northern Xinjiang,
western China. In: Mateer, N.J., Chen, P.J. (Eds.), *Aspects of Nonmarine Cretaceous
Geology*. China Ocean Press, Beijing, pp. 49–77.
- Simionescu, I., 1906. Note sur l'age des calcaires de Cernavoda (Dobrogea). *Annales
Scientifiques de l'Université de Jassy* 4(1), 1–3.
- Simionescu, I., 1913. *Megalosaurus* aus der Unterkreide der Dobrogea. *Centralblatt für
Mineralogie, Geologie und Paläontologie* 1913(20), 686–687.
- Sissingh, W., 1977. Biostratigraphy of Cretaceous calcareous nannoplankton. *Geologie en
Mijnbouw* 56, 37–65.
- Smith, J.B., Vann, D.R., Dodson, P., 2005. Dental morphology and variation in theropod
dinosaurs: implications for the taxonomic identification of isolated teeth. *The Anatomical
Record A* 285A, 699–736.
- Stein, K., Csiki, Z., Curry Rogers, K., Weishampel, D.B., Redelstorff, R., Carballido, J.L.,
Sander, P.M., 2010. Small body size and extreme cortical bone remodeling indicate phyletic
dwarfism in *Magyarosaurus dacus* (Sauropoda: Titanosauria). *Proceedings of the National
Academy of Sciences* 107, 9258–9263.

- Stoica, M., Csiki, Z., 2002. An earliest Cretaceous (Purbeckian) vertebrate fauna from Southern Dobrogea (southeastern Romania). In: Grigorescu, D., Csiki, Z. (Eds.), 7th European Workshop on Vertebrate Palaeontology, Sibiu, Romania, [Ars Docendi, Bucharest](#), pp. 34.
- Stovall, J.W., Langston, W., Jr., 1950. *Acrocanthosaurus atokensis*, a new genus and species of Lower Cretaceous Theropoda from Oklahoma. *American Midland Naturalist* 43, 686–728.
- Stromer, E., 1931. Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wüsten Ägyptens. II. Wirbeltier-Reste der Baharijstufe (unterstes Cenoman). 10. Ein Skelett-Rest von *Carcharodontosaurus* nov. gen. *Abhandlungen der Bayerischen Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Abteilung*, Neue Folge 9, 1–23.
- Therrien, F., Henderson, D.M., 2007. My theropod is bigger than yours ... or not: estimating body size from skull length in theropods. *Journal of Vertebrate Paleontology* 27, 108–115.
- Tong, H., Claude, J., Suteethorn, V., Naksri, W., Buffetaut, E., 2009. Turtle assemblages of the Khorat Group (Late Jurassic–Early Cretaceous) of NE Thailand and their palaeobiogeographical significance. In: Buffetaut, E., Cuny, G., Le Loeuff, J., Suteethorn, V. (Eds.), *Late Palaeozoic and Mesozoic Ecosystems in SE Asia*. The Geological Society, London, Special Publications 315, pp. 141–151.
- Torcida Fernández-Baldor, F., Canudo, J.I., Huerta, P., Montero, D., Pereda Suberbiola, X., Salgado, L., 2011. *Demandasaurus darwini*, a new rebbachisaurid sauropod from the Early Cretaceous of the Iberian Peninsula. *Acta Palaeontologica Polonica* 56, 535–552.
- Turculeț, I., Brânzică, M., 2012. Muzeul colecțiilor paleontologice originale de la Universitatea "Alexandru Ioan Cuza" Iași. Editura Universității "Alexandru Ioan Cuza" Iași, Iași, 173 pp.
- Turner, A.H., Sertich, J.J.W., 2010. Phylogenetic history of *Simosuchus clarki* (Crocodyliformes: Notosuchia) from the Late Cretaceous of Madagascar. *Journal of Vertebrate Paleontology* 30(Supplement 1), 177–236.

- Turner, A.H., Makovicky, P.J., Norell, M.A., 2012. A review of dromaeosaurid systematics and paravian phylogeny. *Bulletin of the American Museum of Natural History* 371, 1–206.
- Vasile, Ș., Csiki, Z., 2011. New Maastrichtian microvertebrates from the Rusca Montană Basin (Romania). *Oltenia. Studii și comunicări. Științele Naturii* 27(1), 221–230.
- Vickers-Rich, P., Rich, T.H., Lanus, D.R., Rich, L.S.V., Vacca, R., 1999. "Big Tooth" from the Early Cretaceous of Chubut Province, Patagonia: a possible carcharodontosaurid. In: Tomida, Y., Rich, T.H., Vickers-Rich, P. (Eds.), *Proceedings of the Second Gondwanan Dinosaur Symposium*. National Science Museum Monographs 15, Tokyo, pp. 85–88.
- Vremir, M., 2010. New faunal elements from the Late Cretaceous (Maastrichtian) continental deposits of Sebeș area (Transylvania). *Terra Sebus. Acta Musei Sabesiensis* 2, 635–684.
- Vullo, R., Néraudeau, D., Lenglet, T., 2007. Dinosaur teeth from the Cenomanian of Charentes, western France: evidence for a mixed Laurasian-Gondwanan assemblage. *Journal of Vertebrate Paleontology* 27, 931–943.
- Weishampel, D.B., 1990. Dinosaurian distribution. In: Weishampel, D.B., Dodson, P., Osmólska, H. (Eds.), *The Dinosauria*. California University Press, [Berkeley, Los Angeles, Oxford](#), pp. 63–140.
- Weishampel, D.B., Grigorescu, D., Norman, D.B., 1991. The Dinosaurs of Transylvania. *National Geographic Research & Exploration* 7(2), 196–215.
- Weishampel, D.B., Norman, D.B., Grigorescu, D., 1993. *Telmatosaurus transsylvanicus* from the Late Cretaceous of Romania: the most basal hadrosaurid dinosaur. *Palaeontology* 36, 361–385.
- Weishampel, D.B., Jianu, C.M., Csiki, Z., Norman, D.B., 2003. Osteology and phylogeny of *Zalmoxes* (n. g.), an unusual euornithopod dinosaur from the latest Cretaceous of Romania. *Journal of Systematic Palaeontology* 1, 65–123.

- Weishampel, D.B., Barrett, P.M., Coria, R.A., Le Loeuff, J., Xu, X., Zhao, X-J., Sahni, A.,
Gomani, E.M., Noto, C.R., 2004. Dinosaur distribution. In: Weishampel, D.B., Dodson, P.,
Osmólska, H. (Eds.), *The Dinosauria*. Second Edition. University of California Press,
Berkeley, Los Angeles, London, pp. 517–606.
- Weishampel, D.B., Csiki, Z., Benton, M.J., Grigorescu, D., Codrea, V., 2010.
Palaeobiogeographic relationships of the Hațeg biota — Between isolation and innovation.
Palaeogeography, Palaeoclimatology, Palaeoecology 293, 419–437.
- Williamson, T.E., Brusatte, S.L., 2014. Small theropod teeth from the Late Cretaceous of the
San Juan Basin, northwestern New Mexico and their implications for understanding latest
Cretaceous dinosaur evolution. PLoS ONE 9(4), e93190. doi:10.1371/journal.pone.0093190.
- Zarcone, G., Cillari, F.M.P., Stefano, P.D., Guzzetta, D., Nicosia, U., 2010. A possible bridge
between Adria and Africa: New palaeobiogeographic and stratigraphic constraints on the
Mesozoic palaeogeography of the Central Mediterranean area. *Earth-Science Reviews* 103,
154–162.
- Figure captions
- Figure 1. Simplified geological map of the Cernavodă-Cochirleni area; inset shows the
position of the study area within Romania. Legend: 1. Quaternary: a. Holocene alluvia, b.
Pleistocene–Holocene loessoid deposits; 2. Pre-Quaternary Cenozoic (Middle Eocene and
Miocene) deposits; Cretaceous: 3. Peștera Formation, Lower Cenomanian; 4. Cochirleni
Formation; uppermost Aptian–Lower Albian; 5. Gherghina Formation, Middle–Upper
Aptian; 6. Ostrov (= Ramadan) Formation; Barremian–Lower Aptian; 7. Cernavodă
Formation, Alimanu Member, Berriasian–Valanginian; 8. Water courses. (Redrawn after
Dragastan et al., 1998, 2014).

Figure 2. Specimen UAIC (SCM1) 615, indeterminate carcharodontosaurid lateral tooth from Cochirleni, Southern Dobrogea. A. UAIC (SCM1) 615, as figured by Simionescu (1913); B. Current state of UAIC (SCM1) 615, mounted in a limestone holder.

Figure 3. Detailed morphology of UAIC (SCM1) 615, an indeterminate carcharodontosaurid lateral tooth from Cochirleni, Southern Dobrogea. UAIC (SCM1) 615 in A. labial? side; B., distal; C., lingual? side, and D., basal (mesial to the right) views. Details of the distal carina (marked with boxes in A, respectively C): apical part in E., labial? and F. distal views; basal part in G., lingual? and H., distal views. Scale bar: 1 cm (A–D), 5 mm (E–H).

Figure 4. Dental morphospace of the different theropod clades according to the results of the PCA analysis; UAIC (SCM1) 615 (red star) plots within the morphospace occupied by Carcharodontosauridae. See further details of this analysis, as well as other quantitative analyses used to identify the tooth that deliver similar results (cluster analysis, discriminant function analysis, phylogenetic analysis), in the Supplementary Material.

Figure 5. A. Palaeogeographic setting of the two early Early Cretaceous Romanian dinosaur occurrences: the Berriasian–Valanginian Cornet locality (orange star), located on a Neo-Tethyan archipelago island, and the Valanginian Cochirleni locality (red star), situated on the marginal areas of the Eastern European cratonic mainland. B. Global chronostratigraphic and palaeobiogeographic distribution of the Carcharodontosauridae, plotted on Middle Aptian (approx. 120 Mya) palaeogeographic map; red star marks the position of UAIC (SCM1) 615 from Southern Dobrogea. Legend: 1 – *Veterupristisaurus*, ‘*Megalosaurus*’ *ingens*, Carcharodontosauridae indet., Tanzania, Late Jurassic; 2 – *Concavenator*, Spain, Barremian;

1582 3 – Carcharodontosauridae indet., Thailand, Barremian; 4 – *Acrocanthosaurus*, southeastern
 1583 United States, Aptian–Albian; 5 – Carcharodontosauridae indet., Spain, Aptian; 6 –
 1584 *Eocarcharia*, Niger, Aptian–Albian; 7 – Carcharodontosauridae indet., Guangxi, China,
 1585 Aptian; 8 – Carcharodontosauridae indet., Henan, China, Aptian; 9 – *Kelmayisaurus*,
 1586 Xinjiang, China, Aptian–Albian; 10 – Carcharodontosauridae indet., France, Cenomanian; 11
 1587 – *Sauroniops*, Morocco, Cenomanian; 12 – Carcharodontosauridae indet., Japan,
 1588 Cenomanian–early Turonian; 13 – *Shaochilong*, Inner Mongolia, China, Turonian; 14 –
 1589 Carcharodontosauridae indet., São Paulo, Brazil, Campanian–Maastrichtian (for relevant
 1590 references, see text, 5.4.). Palaeogeographic maps, courtesy of Ron Blakey
 1591 (<http://cpgeosystems.com/>).

“*Megalosaurus cf. superbus*” from southeastern Romania: the oldest known Cretaceous
carcharodontosaurid (Dinosauria: Theropoda) and its implications for earliest Cretaceous
Europe-Gondwana connections

Zoltán Csiki-Sava^{1*}, Stephen L. Brusatte², Ștefan Vasile¹

¹ Department of Geology, Faculty of Geology and Geophysics, University of Bucharest, 1
Nicolae Bălcescu Boulevard, 010041 Bucharest, Romania

² School of GeoSciences, University of Edinburgh, Grant Institute, James Hutton Road,
Edinburgh, EH9 3FE, United Kingdom

* Corresponding author
zoltan.csiki@g.unibuc.ro

ABSTRACT

Some of the best records of continental vertebrates from the Cretaceous of Europe come from
Romania, particularly two well-known occurrences of dwarfed and morphologically aberrant
dinosaurs and other taxa that lived on islands (the Cornet and Hațeg Island faunas).
Substantially less is known about those vertebrates living in the more stable, cratonic regions
of Romania (and Eastern Europe as a whole), particularly during the earliest Cretaceous. We
describe one of the few early Early Cretaceous fossils that have ever been found from these
regions, the tooth of a large theropod dinosaur from Southern Dobrogea, which was
discovered over a century ago but whose age and identification have been controversial. We
identify the specimen as coming from the Valanginian stage of the Early Cretaceous, an
incredibly poorly sampled interval in global dinosaur evolution, and as belonging to
Carcharodontosauridae, a clade of derived, large-bodied apex predators whose earliest

Cretaceous history is poorly known. Quantitative analyses demonstrate that the Romanian tooth shows affinities with a derived carcharodontosaurid subgroup, the Carcharodontosaurinae, which until now has been known solely from Gondwana. Our results suggest that this subgroup of colossal predators did not evolved vicariantly as Laurasia split from Gondwana, but originated earlier, perhaps in Europe. The carcharodontosaurine diversification may have been tied to a north-to-south trans-Tethyan dispersal that took place sometime between the Valanginian and Aptian, illustrating the importance of palaeogeographic ties between these two realms during the largely mysterious early–mid Early Cretaceous.

Keywords

Southern Dobrogea; Valanginian; Carcharodontosauridae; cratonic Europe; palaeobiogeography

1. Introduction

Romania boasts one of the best records of continental vertebrate fossils from the Cretaceous of Europe (e.g., Grigorescu, 1992, 2003; Csiki-Sava et al., 2015). The vast majority of fossils come from two well-known occurrences: the Early Cretaceous bauxite accumulations of Cornet, in the northern Apuseni Mountains (e.g., Jurcsák, 1982; Benton et al., 1997; Posmoşanu, 2003; Dyke et al., 2011), and the famous latest Cretaceous beds of the Haţeg, Rusca Montană and western Transylvanian basins of Transylvania, which have yielded the dinosaur-dominated ‘Haţeg Island fauna’ (e.g, Nopcsa, 1923; Weishampel et al., 1991; Benton et al. 2010; Codrea et al., 2010, 2012; Grigorescu, 2010; Vremir, 2010; Vasile and Csiki, 2011; Csiki-Sava et al., 2015). Both of these faunas inhabited islands that were part of the vast Cretaceous European Archipelago of the Neo-Tethys Ocean. Based on their isolated

geological settings and the many dwarfed and morphologically aberrant taxa that make up the faunas, both have been interpreted as insular assemblages that give a unique window into how island environments affected the evolution of long-extinct organisms (e.g., Benton et al., 1997, 2010; Csiki-Sava et al., 2015).

The great volume of research on these assemblages over the past century, particularly the ‘Hațeg Island fauna’, has concealed an inconvenient bias: the stable, non-island, cratonic regions of Romania have yielded only extremely rare Mesozoic continental vertebrate remains (i.e., the Moldavian, Moesian and Scythian platforms; Săndulescu, 1984; Mutihac and Mutihac, 2010; Fig. 1). This is mostly because Mesozoic deposits are located in the subsurface in these regions, with only limited subaerial exposures available in the structurally highest-lying parts of the Moesian Platform, in Central and Southern Dobrogea (Middle Jurassic–Upper Cretaceous), as well as in the northeastern-most corner of the Moldavian Platform, along the Prut Valley (lower Upper Cretaceous) (see, e.g., Mutihac and Mutihac, 2010). This bias is unfortunate because fossils from these settings could lead to a better understanding of how mainland and island faunas differed during the Cretaceous, and because the cratonic portion of Europe was an important biogeographic stepping stone between the north and south as the continents fragmented and sea levels fluctuated.

Although the cratonic regions of Romania have yielded few Cretaceous terrestrial fossils, these deposits are not totally barren. In fact, one of the first Mesozoic continental vertebrates ever recorded from Romania comes from one of these deposits, the Lower Cretaceous shallow marine limestones of Southern Dobrogea (Fig. 1). This specimen—the isolated but well-preserved tooth of a large theropod dinosaur—has often been overlooked. It was described a little over a century ago by Simionescu (1913; Fig. 2A), and until a few recent discoveries of very rare isolated specimens (Stoica and Csiki, 2002; Csiki-Sava et al., 2013; Dragastan et al., 2014), it remained as the sole published record of Mesozoic terrestrial

vertebrates from the cratonic areas of Romania. It has never been comprehensively described and its precise age and taxonomic affinities have yet to be clarified, despite its potential importance as a well-preserved fossil from a poorly sampled area that could have critical evolutionary and biogeographic implications.

We here present a comprehensive description of the Dobrogea tooth and discuss its relevance for understanding dinosaur evolution and biogeography. We review the peculiar history of how this specimen was collected and curated, thoroughly document its morphology and age, identify it based on comparison to a broad range of theropods, and outline its importance. It turns out that this specimen, although only a single tooth, has wide-ranging implications. We identify it as coming from the Valanginian stage of the Early Cretaceous, which is incredibly poorly sampled both in Europe and globally (Weishampel et al., 2004), and as belonging to a carcharodontosaurid, a group of derived, large-bodied apex predators whose earliest Cretaceous history is poorly known. Carcharodontosaurids were once thought to be a uniquely Gondwanan group, but recent discoveries show that the basal members of the group were more widespread during the late Early-middle Cretaceous (e.g., Sereno et al., 1996; Brusatte and Sereno, 2008). The Romanian tooth shows affinities with a derived carcharodontosaurid subgroup, the Carcharodontosaurinae, that until now has been known only from Gondwana. It suggests that this subgroup of enormous predators did not evolve vicariantly as Pangaea split, but originated earlier, and perhaps in Europe, suggesting faunal interchange between Europe and Gondwana during the ‘dark ages’ of the early Early Cretaceous.

Abbreviations: UAIC – University “Alexandru Ioan Cuza”, Iași, Romania.

2. History of collecting and curation

Only two dinosaurian fossils are currently known from the cratonic areas of Romania: an isolated theropod tooth and an isolated caudal vertebral centrum. Both of these were reported from the Lower Cretaceous deposits of Southern Dobrogea (southeastern Romania; Csiki-Sava et al., 2013, see also below). Unfortunately, exact details of their discovery and places of origin are lost, a fact that can hinder an assessment of their age and interpretation of their phylogenetic and palaeobiogeographic significance. Our aim here is to gather and report all available information concerning the collecting of specimen UAIC (SCM1) 615, that is, the isolated theropod tooth reported by Simionescu (1913; Fig. 2A).

According to the existing information - unpublished museum labels and records, and the preliminary publication of Simionescu (1913) - specimen UAIC (SCM1) 615 was discovered in the surroundings of Cochirleni, a small village south of Cernavodă and close to the right bank of the Danube, in Southern Dobrogea, southeastern Romania (Fig. 1), probably shortly before 1913, the date of its publication by Simionescu (1913).

Although studied and preliminarily described by Simionescu, UAIC (SCM1) 615 was not collected by Simionescu personally. Instead, it was donated by a certain “de Tomas” (also mentioned as “de Thomas” in the registry of the Hârşova Museum) to V. Cotovu from Hârşova (Central Dobrogea), a local teacher, archaeology and natural history aficionado, and amateur fossil collector (see, e.g., Covacef, 1995). Cotovu, described by Simionescu himself as the “zélé fondateur et directeur du muséum de Hârşova” (*enthusiastic founder and director of the Hârşova Museum*; Simionescu, 1906: p. 2), had previously provided fossil specimens from Southern Dobrogea for study to Simionescu, a nationally acknowledged popular science writer and scientist, whom Cotovu knew personally (Brânzilă, 2010). These circumstances are supported by the fact that in the original description, Simionescu figures the specimen as being accessioned in the “Regional-Museum von Harschowa” (*Hârşova Regional Museum*; Simionescu, 1913: p. 687, fig.1), a designation he also used to refer to other Dobrogean

specimens not collected by him first-hand (e.g., a specimen of '*Nautilus' pseudoelegans* from Cernavodă, or a fragmentary tooth-bearing palatal fragment referred to as '*Coelodus*' sp., also originating from Cochirleni; see Simionescu, 1906). Confirming this deduction, an isolated tooth appears accessioned in the old registry book of the Hârşova Museum (under specimen number 200) as "*Megalosaurus cf. superbus*", with the mention that it was "described by Prof. Simionescu in the *Centralblatt f. min. etc.*". This is also the case of the '*Coelodus*' sp. specimen from Cochirleni (specimen number 86), similarly clearly identified as being described by Simionescu in the registry book.

Both of these vertebrate remains from Dobrogea that were formerly part of the Hârşova Museum collections are currently accessioned in the palaeontology collections of the UAIC (Turculeţ and Brânzilă, 2012), suggesting that, at one moment, several specimens were transferred there from the Hârşova Museum. Although no details are known about this transfer, it is probable that it took place right before (or when) the Hârşova Museum, including a part of its collections, was burned and largely destroyed during WWI, in 1916, a time when Simionescu still held a position at the UAIC.

After its original description, specimen UAIC (SCM1) 615 underwent a minor amount of damage (see below, Description). Also, at some point between its description in 1913 and the early 1960s (when the specimen was found in its present state in the collections of the UAIC by academic staff members who are still alive today and recall the discovery; I. Turculeţ, personal communication, May 2013) it was glued into a limestone matrix holder, while it was obviously completely freed of the surrounding matrix when it was described and figured in 1913 (Fig. 2). The circumstances under which these alterations took place are unclear. It is a distinct possibility that they occurred sometimes during WWII, when, in the spring of 1944, the frontline between the German-Romanian and Soviet armies reached the Iaşi–Chişinău line. At this moment, the geological-palaeontological collections of the UAIC

150 were packed in crates, and moved together with its personnel and other possessions to Zlatna,
151 in the Apuseni Mountains (western Romania), to safeguard them from any potential damage.
152 Mounting the specimen into the limestone stand would have been a quick way to stabilize it,
153 as it appears that packaging and transport of the specimens was done in haste (M. Brânzilă,
154 personal communication, April 2103). If that was indeed the case, the mounting would have
155 taken place without the knowledge of Simionescu, who left Iași and the UAIC in 1929, being
156 invited to become a professor of Palaeontology at the University of Bucharest (Brânzilă,
157 2010). Then again, however, Simionescu himself or staff of the Hârșova Museum might have
158 re-mounted the tooth after its original description, or else the mounting might have taken
159 place after the return of the collections to Iași, after WWII.

160 Unfortunately, it is not documented whether the mounting was made using the
161 original matrix, or if a trough corresponding to the tooth outline was carved into a randomly
162 chosen limestone block. The apparently excellent fit between the tooth and the depression
163 housing it (Fig. 2B, 3) suggests that this operation was completed carefully, and accurate
164 carving of a fake holder is difficult to reconcile with the rush accompanying the evacuation of
165 the Iași University, in 1944. Alternatively, the presence of a hand-written old registration
166 number on the specimen holder would support its early re-mounting, while still at the
167 Hârșova Museum. As noted previously, the original Hârșova Museum registration number of
168 the specimen was 200, which does not correspond to that currently written both on the
169 limestone holder and on a paper sticker (204). However, according to the old collection
170 registry of the Hârșova Museum, specimen numbers 201 through 225 were given to a series
171 of “indeterminate (fossil) bone fragments” from the “Cochirleni quarries”. Thus, these
172 specimens (now apparently lost) came from the same locality as the tooth, and they were
173 collected and donated by the same person to the Museum who donated UAIC (SCM1) 615.
174 There is, thus, a (albeit admittedly remote) possibility that the registration numbers were

175 mixed up during the re-mounting of the specimen, which in this case took place at an early
176 date in the Hârşova Museum. If this is indeed the case, the limestone used as holder could
177 have been the same as the original matrix of the specimen.

178 To conclude, the history of recovery and curation of the historically important
179 dinosaurian specimen UAIC (SCM1) 615 is rather convoluted and clouded by many
180 uncertainties. The exact date of discovery remains conjectural, and the exact place of the
181 discovery (thus also the original geological context of the tooth) is even more ambiguous.
182 The current state of the specimen, and especially its mounted status, suggest a curatorial
183 history that produced a moderate amount of damage to, but also partially obscured the
184 detailed morphology of the specimen. The convergence of such unfortunate events makes
185 deciphering the age, identity and evolutionary significance of the specimen troublesome,
186 although many lines of evidence, carefully considered, allow us to draw reasonable
187 conclusions (see below).

189 3. Geological setting

190 According to the available collecting information, the isolated theropod tooth UAIC (SCM1)
191 615 was discovered at Cochirleni (sometimes noted more specifically as the “*Cochirleni*
192 *quarry*” or “*Cokerleni quarry*”). Cochirleni is a small village in southwestern Dobrogea
193 situated close to the right bank of the Danube, and about 9 km south of the main urban center
194 of the region, Cernavodă (Fig. 1). The geology of the area has been well studied, because of
195 the unique outcropping conditions and rich fossiliferous nature of the Lower Cretaceous
196 deposits (reviewed in Avram et al., 1996; Neagu et al., 1997; Dragastan et al., 1998).

197 Southern Dobrogea is a cratonic area corresponding to the southeastern corner of
198 Romania. Whether it is considered part of the larger Moesian Platform (Săndulescu, 1984;
199 Ionesi, 1994), or a distinct craton (the South-Dobrogean Platform; Mutihac and Mutihac,

2010), researchers agree that it became integrated into the main European Craton towards the end of the Jurassic, at the latest, with the consolidation of the Cimmerian (Early Alpine: Triassic–earliest Cretaceous) North Dobrogean fold-and-thrust belt (Seghedi, 2001; Hyppolite, 2002). The age of its basement is also controversial, with estimates ranging from Archaic–Early Proterozoic (Mutihac and Mutihac, 2010) to latest Proterozoic (Ionesi, 1994).

The Precambrian basement of Southern Dobrogea is overlain by a flat-lying sedimentary cover that begins with the lowermost Palaeozoic and ends with the uppermost Neogene. The sedimentary succession is interrupted by a few major, as well as several less important, sedimentary hiatuses that separate 5 main sedimentary sequences corresponding to the Cambrian–Upper Carboniferous, the Permian–Triassic, the Middle Jurassic–Cretaceous, the Eocene–?Oligocene, and the middle Badenian (middle Miocene)–Upper Pliocene. The Palaeozoic and lower Mesozoic are known only from the subsurface of Southern Dobrogea, but Cretaceous and Cenozoic deposits have limited exposures along the main water courses of the region (Ionesi, 1994; Mutihac and Mutihac, 2010).

The outcropping Cretaceous in Southern Dobrogea is represented mainly by shallow marine, carbonate platform deposits in the lower part of the system, replaced by more open-water, chalky facies towards the later part of the period (e.g., Avram et al., 1993, 1996; Dragastan et al., 1998; Dinu et al., 2007); these crop out only as isolated patches along the main watercourses of the region (Fig. 1).

The Lower Cretaceous Series consists of several lithostratigraphic units with complex, partially overlapping and interfingering relationships (Dragastan et al., 1998, 2014). The lowest (and only artificially) outcropping unit is the Purbeck-type, siliciclastic-evaporitic Upper Kimmeridgian–Lower Berriasian Amara Formation that represents lagoonal to continental environments. This unit is covered by the shallow-marine, richly fossiliferous and locally reefal limestone-dominated Cernavodă Formation (restricted-open lagoonal to

carbonate platform, Upper Berriasian–Lower Hauterivian). A time-correlative unit of the Cernavodă Formation, the limestone-dolomitic Dumbrăveni Formation (Upper Berriasian–Lower Hauterivian), is restricted to the southeastern part of Southern Dobrogea. The Cernavodă and Dumbrăveni formations are covered unconformably by dominantly calcareous deposits with hippuritoid (‘pachyodont’) coquinas, small reefs and lens-like orbitolinid accumulations, referred to the Barremian–Lower Aptian Ostrov Formation by Dragastan et al. (1998), but to the Ramadan Formation (in part) by Avram et al. (1993, 1996). These deposits, formed in littoral to lagoonal and open reef terrace environments, are in turn capped by the fluvial-lacustrine, siliciclastic deposits of the Gherghina Formation, with Middle–Upper Aptian kaolinitic clays and thin coal intercalations. The Lower Cretaceous succession ends with the transgressive, glauconite-bearing, coastal to sublittoral siliciclastic deposits of the Cochirleani Formation (uppermost Aptian–Albian).

The Upper Cretaceous has a significantly more patchy development, mainly restricted to the eastern part of Southern Dobrogea, excepting the weakly glauconitic, chalky-sandy Peștera Formation (Lower Cenomanian) and the marly Dobromiru Formation (Upper Cenomanian) that cover the western-central parts of the area. The younger Cuza Vodă (Turonian), Murfatlar (Santonian–Lower-Middle Campanian), and Satu Nou (Upper Campanian) formations are dominantly chalky, suggesting the instalment of a relatively deeper, offshore depositional environment; neither of these units is known from western Southern Dobrogea.

In total, the Lower Cretaceous of Southern Dobrogea was deposited in a shallow marine, near-shore setting, fluctuating between carbonate platform, lagoonal, coastal-tidal flat, and continental environments (see Avram et al., 1996; Dragastan et al., 1998). Its main characteristic features, such as the observed lithological variability, the areal distribution of the different units, and the presence of several unconformities within the series, are all linked

to eustatic sea-level changes that affected the Southern Dobrogean territory during the Early Cretaceous (Dragastan et al., 1998). The main emergent land in the area was represented by the Central Dobrogean Massif, lying north of the study area, almost completely subaerially exposed and actively eroding during the Cretaceous. Consequently, shallow-marine to continental deposits are restricted mainly to the northern part of Southern Dobrogea, close to its boundary with the Central Dobrogean Massif (marked by the Capidava-Ovidiu Fault), and are replaced by more open marine deposits southward. As summarized above, several littoral, and even continental, sequences occur in this succession, including deposits in the Amara, Cernavodă, Ramadan (Avram et al., 1996) and Cochirleni formations, whereas the Gherghina Formation is purely continental, with occasional minor marine interbeds produced during short-term ingressions of the sea.

In the Cernavodă-Cochirleni area the outcropping Mesozoic is restricted to the Lower Cretaceous, and includes deposits belonging to the Cernavodă, Ostrov (or Ramadan), Gherghina, and Cochirleni formations. While the lower–middle part of the Cernavodă Formation is well exposed and widely distributed in this area, its upper part (the lower Hauterivian Vederoasa Member) is unevenly developed. This member is missing in the classical succession from Cernavodă-Hinog, on the right bank of the Danube (Dragastan et al., 1998), but was recently identified in the more eastern Cernavodă-lock section (Dragastan et al., 2014). Similarly, the Ostrov Formation is represented in the area only by its upper subunit (the Lower Aptian Lipnița Member; Dragastan et al., 1998), covering unconformably and transgressively the Valanginian Alimanu Member of the Cernavodă Formation in the southern end of the Cernavodă-Hinog section (Dragastan et al., 1998), and the lower Hauterivian Vederoasa Member in the Cernavodă-lock section (Dragastan et al., 2014).

Northward of the Hinog area, Valanginian deposits of the Alimanu Member are overlain directly by the Middle–Upper Aptian continental deposits of the Gherghina

Formation. These continental deposits also cover the *Orbitolina*-bearing calcareous-clayey deposits of the Lipnița Member towards the south, marking the advancement of emerged areas towards the central parts of Southern Dobrogea, including the Cernavodă-Cochirleni area, during this time interval (Avram et al., 1996). Marine conditions returned in the study area again in the latest Aptian, with a transgression marked by widespread deposition of the glauconitic, siliciclastic coastal to innermost shelf deposits of the Cochirleni Formation. These uppermost Aptian to Albian sands and sandstones cover transgressively all the underlying deposits, belonging to the Cernavodă, Ostrov, or Gherghina formations. Siliciclastic shallow-marine sedimentation continued into the Early Cenomanian, with the chalky-glauconitic deposits of the Peștera Formation.

4. Palaeontology

The isolated theropod tooth UAIC (SCM1) 615 (formerly in the collections of the Hârșova Museum, registered with no. 200; Fig. 2A) was described in a short note by Simionescu (1913), who referred it to *Megalosaurus* cf. *superbus*, a taxon erected by Sauvage (1882) from the Gault ('mid'-Cretaceous: Albian) of the Paris Basin, France. The Gault material described by Sauvage (1882; see also Sauvage, 1876) includes several isolated teeth that were deemed by Simionescu (1913) to be more similar to the Cochirleni tooth than are the teeth of *Megalosaurus bucklandi* (Buckland, 1824). Subsequently, the French Gault material was referred to the new genus *Erectopus* by Huene (1923), who also noted differences between it and the type species *M. bucklandi*.

The convoluted taxonomic history of *Erectopus superbus* was recently reviewed by Allain (2005), who established that both the isolated teeth first mentioned by Sauvage (1876) and the skeletal elements described by Sauvage (1882) belong to the same taxon, for which the name *Erectopus superbus* was retained. Allain (2005) regarded *Erectopus* as a member of

Carnosauria (= basal Tetanurae), an opinion also shared by Molnar (1990) and Holtz et al. (2004a), whereas the latest review of the Tetanurae (Carrano et al., 2012, p. 254) considered *Erectopus superbus* “a non-carcharodontosaurian allosauroid, possibly a metriacanthosaurid.” Accordingly, if we are following the original assessment of Simionescu (1913) but updating with contemporary taxonomy, the Cochirleni theropod tooth should now be considered referable to the basal tetanuran *Erectopus superbus*. However, the referral of this tooth to *Erectopus superbus* (or a close relative) was considered to be unsupported by positive evidence by Molnar (1990) and Holtz et al. (2004a). In order to re-assess this referral and to understand the exact taxonomic and phylogenetic affinities of UAIC (SCM1) 615 (Fig. 2B, 3), we provide here a detailed description of its morphology followed by a thorough comparative study of this tooth based on large datasets of theropod dental measurements and discrete characters compiled by Hendrickx and Mateus (2014) and Hendrickx et al. (2015a).

We note that in his review of Romanian dinosaurs, Grigorescu (2003) erroneously considered UAIC (SCM1) 615 as being referred by Simionescu to the taxon *Megalosaurus dunkeri* Kohen (sic; actually, *Megalosaurus dunkeri* Dames, 1884). This is clearly a simple misreading of Simionescu’s identification. Additionally, such a referral is also contradicted by the absence of mesial serrations in the holotype tooth of *M. dunkeri*, considered by Carrano et al. (2012) to represent an indeterminate theropod. The Dobrogea tooth, on the other hand, has mesial serrations (see below).

4.1. Age of UAIC (SCM1) 615

The age of UAIC (SCM1) 615 has been contentious, due to the uncertainties concerning its place of origin. Although it is often mentioned as originating from Cochirleni village (e.g., Grigorescu, 2003; Turculeț and Brânzilă, 2012), this has not been definitively established. According to the original report of Simionescu (1913), the tooth came from the upper part of

the Lower Cretaceous limestone succession exposed in the cliffs extending from Cernavodă to Cochirleni along the right bank of the Danube. The corresponding entry from the Hârşova Museum registry states that it was found in the ‘Cochirleni quarry’, a location that presently cannot be identified precisely. The only rocks to be quarried in the area are the calcareous deposits of either the Cernavodă or Ostrov formations, particularly the ones that crop out in the Danube bank cliffs between Cernavodă-Hinog-Cochirleni. Finally, although the mention ‘Cochirleni’ is usually considered to refer to Cochirleni village, it should be mentioned that the cliff-forming hill that extends between Cernavodă and Cochirleni is also known by the same name (Fig. 1). Taking all of this evidence into consideration, it is thus reasonable to conclude that the tooth was most likely found in the Lower Cretaceous limestone succession exposed in the Danube cliffs between Cernavodă and Cochirleni.

Based on the location of the discovery, in the upper part of the local limestone succession, and the age of the deposits from Cernavodă-Cochirleni known to him, Simionescu (1913) considered the tooth to be of Barremian age. Subsequently, the age of the tooth was given as Valanginian–Barremian (Weishampel, 1990; Weishampel et al., 2004) or Valanginian (e.g., Grigorescu, 2003), but without any supporting information.

New attempts have been made to more precisely constrain the age of UAIC (SCM1) 615. Dragastan et al. (2014) recently sampled the limestone matrix holder of the tooth, and reported from these samples an assemblage of foraminiferans, ostracods and microproblematicae (= *incertae sedis* microorganisms) that characterize their ‘Biozone IX with *Meandrospira favrei*’, of latest Valanginian age in the local lithostratigraphic scheme. In parallel, we also sampled the same limestone holder – a yellowish white, friable lime mudstone – that yielded a poor and badly preserved calcareous nannoplankton assemblage with *Watznaueria barnesiae*, *W. ovata*, *Nannoconus steinmanni*, *N. kamptneri*, *N. globulus*, *Calcicalathina* sp., *Speetonia colligata* and *Cyclagelosphaera deflandrei* (M. C. Melinte-

Dobrinescu, personal communication, November 2013), an assemblage that suggests a Berriasian–Hauterivian age of the limestone holder.

Since it is not clear if the limestone holder came from the same site as the tooth itself, we managed to take a second sample from the limestone matrix still partly filling the pulp cavity of the tooth, which must definitively be identical with the rocks the tooth was found in. This second, much smaller sample yielded only very scarce specimens of *Watznaueria barnesiae*, *Cyclagelosphaera margerelii* and *Diazomatolithus lehmanni* (M. C. Melinte-Dobrinescu, personal communication, November 2013), the latter two taxa having a peak in abundance during the Berriasian and, especially, the Valanginian.

In the nannoplankton succession reported previously by Avram et al. (1993) and derived from a systematic sampling of the Southern Dobrogean Lower Cretaceous, the concurrent presence of *Speetonia colligata*, *Calcicalathina oblongata*, *Diazomatolithus lehmanni* and *Nannoconus steinmanni* was noted in samples derived from the Alimanu Member of the Cernavodă Formation. These assemblages were interpreted to represent the nannoplankton zone CC3 of Sissingh (1977), of late Valanginian age. A comparable age was assigned to a roughly similar nannoplankton assemblage reported from the Lower Cretaceous of the Mecsek Mountains, Hungary, by Császár et al. (2000).

Together, all the available evidence (Simionescu's original account, geographic and geologic records, foraminifera, ostracods, microproblematicae, and calcareous nannoplankton) thus suggests that UAIC (SCM1) 615 originates from the Alimanu Member of the Cernavodă Formation, and it is most probably of late Valanginian age.

4.2. Description and comparisons

Specimen UAIC (SCM1) 615 is a large (total length, as preserved, is about 100 mm; Figs. 2, 3) lateral tooth of a theropod dinosaur, with a crown base length (CBL) of 29 mm, crown

base width (CBW) of 16.25 mm, crown height (CH) of 85.5 mm, and apical length (AL) of 91 mm (terminology following Smith et al., 2005 and Hendrickx et al., 2015b). It is remarkably well preserved, with the enamel in pristine condition. It preserves most of the crown and a small basal part of the root, but the crown tip is broken off, with an estimated 5 mm missing in the apical region.

In its present state, the mesial edge and part of the mesial third of the tooth are embedded in the limestone holder (Fig. 2B), although the tooth was once removed (see above, History of collecting and curation; Fig. 2A). Accordingly, it is exposed so that all faces of the tooth are widely visible, including the root region, except for the mesial surface.

Only the basal-most part of the root is preserved, and it is more complete near the mesial margin (Fig. 3B, C). Here, broken areas around the crown-root contact area (cervix) reveal details of the pulp cavity development, as well as the pattern of the dentine thickness variation (Fig. 3B–D). The crown also exhibits a transverse break at about two-thirds of its length (not present so obviously in the original figure of the specimen in Simionescu, 1913), and adjacent to it, the distal carina is also slightly chipped distal to mid-length. The labial face is superficially split near this break (Fig. 3A), while a more prominent region of damage appears on the lingual face, where a large (13 x 5 mm), slightly triangular wedge is broken off, exposing the deeper parts of the dentine (Fig. 3C). The damage to the lingual side apparently occurred after the original description of the tooth (Fig. 2), an observation that is concordant with the complex curatorial history of the specimen.

The basal-most, exposed part of the mesial face lacks the enamel cover (Fig. 3C, D), suggesting that this area already belongs to the root region. The mesial edge of the preserved crown base appears to be wider than the distal one, and is largely rounded transversely. Accordingly, the basal cross-section is teardrop-shaped (lanceolate); it is rounded mesially, but narrows distally into a small carina (Fig. 3D). As mentioned above, the pulp cavity is

exposed basally, being partly filled with a whitish-light gray limestone that is reminiscent of the matrix holder lithology. The pulp cavity narrows rapidly towards the cervix, as it is about 7.1 mm wide (labiolingually) at the apical-most part of the preserved root, but only about 4.5 mm wide at the base of the crown. In parallel, the enamel-dentine wall of the tooth becomes thicker: it is 3.5 mm thick in the apical-most part, 4.4 mm at the base of the crown, but thickens to 5.0–5.8 mm near the apical-most part of the basal break of the crown (Fig. 3B). Mirroring the outside cross-section, the contour of the pulp cavity is also teardrop-shaped (Fig. 3D).

The tooth is ziphodont and only very slightly recurved distally. The distal edge is nearly straight across its length, being very mildly concave in its basal half and slightly convex near its apex (Fig. 2, 3A). Thus, the apex is placed roughly at the distal margin of the tooth crown base. The mesial edge, as shown in the original publication of Simionescu (1913), is strongly convex across its entire length (Fig. 2A). The tooth is labiolingually compressed (Fig. 3B), with a crown base ratio ($CBR = CBW/CBL$) of 0.56, within the normal range of variation of most theropods. This differs from the thinner teeth of some, but not all, carcharodontosaurids ($CBR < 0.50$), and the much thicker incrassate teeth of derived tyrannosauroids and conical teeth of spinosaurids ($CBR > 0.75$) (Serenó et al., 1996; Brusatte et al., 2010a; Hendrickx and Mateus, 2014; Hendrickx et al., 2015a).

The crown cross-section is slightly asymmetrical labiolingually when it is seen in distal view. In this view, when the carina is facing directly distally, one side of the crown has a more pronounced bulge than its counterpart (about 8.5 mm wide, measured from the carina, vs. 6 mm on the other side; Fig. 3B); based on comparisons with the teeth of *Mapusaurus* (Coria and Currie, 2006), the more bulging side can be interpreted as the lingual one. This asymmetry diminishes apically, where both sides become about equally convex. The distal carina itself twists slightly sideways (labially) in apical direction, such that it is located closer

to the labial face where it terminates at the crown apex, and the lingual face of the denticles is exposed distally (Fig. 3B, F). This twist of the distal carina is accompanied by a similar outline of the lingual side; in distal view, this is somewhat convex basally, but becomes flat to slightly concave in the apical two-thirds of the crown. A similar S-shaped curvature of the crown, albeit more pronounced and different in details, was also reported in *Mapusaurus* and *Giganotosaurus* (Coria and Currie, 2006), and in indeterminate carcharodontosaurid teeth from Morocco (Richter et al., 2013).

The distal carina extends along the entire tooth height (Fig. 3A–C). It is covered with minute serrations across its entire preserved length; the denticles are proximodistally subrectangular, with a mesiodistal long axis that is greater than the apicobasal long axis (Fig. 3E–H). They are either roughly perpendicular to the tooth margin, or their long axes are oriented obliquely, such that they point slightly apically. The tip of the apex is broken off, so it is not possible to determine whether the serrations continued over the apex of the tooth. There are approximately 12.5 serrations (denticles) per 5 millimetres at the midpoint of the carina. Serration shape and size remain relatively constant across the carina, although the serrations near the midpoint and closer to the base of the carina (12 denticles per 5 mm; Fig. 3G, H) are slightly smaller than those near the apex (9 denticles per 5 mm; Fig. 3E, F). Changes in serration size are gradual across the carina, not sudden or sporadic.

Although they are all more or less rectangular in shape, the apical denticles are relatively shorter proximodistally than the more basal ones. Most of the denticles have slightly rounded, asymmetrically convex triangular tips, instead of being simply squarred-off, and they do not hook as in troodontids and to a lesser extent abelisaurids (Hendrickx and Mateus, 2014). Other denticles near the apex, however, show a faint concavity along their tips, giving them a bilobate aspect, although this is both less conspicuous and far less regularly developed than reported in *Tyrannotitan* (Novas et al., 2005). The denticles are

separated by simple, linear grooves (interdenticular slits or sulcae) along their entire length.

The interdenticular space between adjacent denticles is broad, measuring more than a third of the apicobasal width of a denticle (Fig. 3E, G). This space continues onto the surface of the crown as a very short interdenticular sulcus (“blood groove” of Currie et al., 1990). These sulci are so short and indistinct that they are only visible under low angle light.

Little can be said about the mesial carina, as it is not visible in the current state of the specimen, buried in the limestone matrix. Based on the description of Simionescu (1913), however, it is covered across its length with minute serrations; these decrease in size towards the base of the crown. Simionescu (1913) reported approximately 15 serrations (denticles) per 5 millimetres at the midpoint of the carina, meaning that the mesial denticles are slightly smaller than those on the distal carina. The denticle size difference index (DSDI: Rauhut and Werner, 1995) is 1.2, within the range of variation of most theropods (Hendrickx and Mateus, 2014). As Simionescu (1913) already pointed out, the presence of a mesial carina that extends towards the base of the crown sets apart UAIC (SCM1) 615 from *Megalosaurus bucklandii* where this stops well above the cervix (Benson et al., 2008), and it is instead similar to ‘*M. superbis*’ (Sauvage, 1876, 1882) in this respect.

The external enamel surface exhibits two forms of ornamentation. First, the majority of the labial and lingual faces are covered by relatively smooth enamel that exhibits a subtle form of braided texture visible under low angle light (Fig. 3A, C, E). This texture is made up of a series of very faint, apico-basally running ridges; these are of unequal lengths, starting at different points of the crown height, but none extends the whole length of the crown. The two longest ridges are placed near the distal carina. The enamel is also finely granulated.

Second, near the carinae on both labial and lingual surfaces there are marginal undulations: wrinkles in the enamel that stand out in bas relief (Brusatte et al., 2007). These are much better preserved and visible near the distal carina, where they are so pronounced

that they are clearly observable in normal light (Fig. 3A–C, G, H). Here, about 17 unevenly developed wrinkles are present along the crown height; in the basal half of the crown, the wrinkles extend about 6.5 mm onto the crown. These are elongate, such that they are longer than twice the space separating each undulation. The wrinkles project obliquely (in the mesiobasal direction) relative to the carina. They are apically concave, with a near-horizontal segment on the crown, and curve apically as they approach the carina (at about 45°) with a tendency to become tangential to the distal edge. The wrinkles are especially well developed, prominent and closely spaced in the basal part of the crown (about 7 wrinkles/16 mm; Fig. 3C, G)), but become more widely spaced and indistinct apically (about 3 wrinkles/16 mm). Apically, however, the wrinkles are somewhat wider and longer, extending over about half of the crown fore-aft length. Again, a slight asymmetry is present between the two sides of the crown in wrinkle development as well, these being better expressed on the more rounded, convex lingual face, but less well expressed on the flatter labial face (Fig. 3A, C, H). On the presumed labial face, only some of the basal-most wrinkles, particularly the second and third one, appear well defined.

Towards the base of the crown a few of the wrinkles continue across the labial and lingual surfaces as very subtle transverse undulations. Most conspicuous of these is a 3.5 mm wide horizontal swelling that crosses the crown, at the level of wrinkles 2 and 3; this swelling is clearly visible on both sides of the crown (Fig. 3. A, C). There are no lateral flutes, apico-basal ridges, or longitudinal grooves on the labial or lingual faces, either in the centre of the tooth or paralleling the carinae. Instead, the labial and lingual faces are uniformly convex, giving the tooth its teardrop-shaped outline in cross section.

5. Discussion

5.1. *Identification of UAIC (SCM1) 615*

The isolated tooth from Cochirleni can be referred to Theropoda based on its large size, recurved and labiolingually compressed morphology, and presence of a continuous series of well-defined serrations on the distal carina.

Besides theropods, certain derived crocodyliforms – the sebecosuchians of Colbert (1946; see also Turner and Sertich, 2010; Pol and Powell, 2011; Rabi and Sebők, 2015) – are also known to possess remarkably theropod-like, laterally compressed and serrated teeth, not unlike the morphology shown by UAIC (SCM1) 615. However, most sebecosuchian teeth are significantly smaller than the Southern Dobrogean specimen, especially in the case of the Cretaceous members of the clade (e.g. *Baurusuchus*; Carvalho et al., 2005). Even the largest, caniniform teeth of the largest representatives of Sebecosuchia, such as the Miocene *Barinasuchus* (Paolillo and Linares, 2007), are somewhat smaller than UAIC (SCM1) 615; moreover, these teeth are slightly conical and less laterally compressed than the Southern Dobrogean tooth. Finally, it should be noted that the oldest known members of Sebecosuchia appear beginning in the Late Cretaceous (e.g. Kellner et al., 2014), and are thus significantly younger than UAIC (SCM1) 615. Similarly, ziphodont crocodyliform teeth (i.e. with true denticles along their carinae) are reported in Europe only beginning in the Albian (Ösi et al., 2015), and these are both significantly smaller and different in morphology from the Dobrogean tooth. Taken together, these suggest that the hypothesis of sebecosuchian affinities of UAIC (SCM1) 615 can be discarded with confidence, and it indeed represents a theropod tooth.

We used four techniques to identify which type of theropod UAIC (SCM1) 615 likely belongs to (see also Supplementary Material).

First, we conducted a Principal Components Analysis (PCA) based on a large database that includes a broad and representative sample of theropod teeth. This dataset was compiled by Hendrickx et al. (2015a), which built upon the earlier studies of Smith et al.

(2005) and Larson and Currie (2013), and it or a similar version has been used in recent studies to identify isolated theropod teeth (e.g., Williamson and Brusatte, 2014; Brusatte and Clark, 2015). It comprises nearly 1000 theropod teeth scored for six measurements (CBL, CBW, CH, AL, MC, and DC, the latter two measuring the density of serrations per 5 mm at the midpoint of the mesial and distal carina, respectively). UAIC (SCM1) 615 was added to this dataset, the data were log-transformed prior to analysis, missing values for measurements were estimated with a mean value for that measurement from across the sample, and then a PCA was run using a correlation matrix. The analysis was conducted in PAST v2.17 (Hammer et al., 2001).

In the resulting two dimensional morphospace (Fig. 4), UAIC (SCM1) 615 plots close to many teeth belonging to carcharodontosaurids, along with some teeth belonging to spinosaurids and tyrannosauroids. It falls within the convex hull (maximum morphospace occupation area) of carcharodontosaurids only, although it is closely outside of the edges of spinosaurid and tyrannosauroid space. It also falls within the 95% confidence interval ellipse for carcharodontosaurids, but not within the ellipse of any other group (Supplementary Information). This exercise indicates that UAIC (SCM1) 615 is most similar to carcharodontosaurids.

Secondly, we used the log-transformed dataset that we also used for the PCA to conduct a clustering analysis. We performed the analysis in PAST v2.17, using the paired group algorithm and the correlation similarity measure. In the resulting dendrogram, UAIC (SCM1) 615 groups with a handful of teeth belonging to carcharodontosaurids, tyrannosauroids, and *Allosaurus* (Supplementary Information).

Third, we used the tooth measurement database to conduct a discriminant analysis in PAST v3.0 (Hammer et al., 2001). This analysis uses pre-determined groups (in this case, taxonomic clusters) to create a morphospace in which these groups are maximally separated.

This allows teeth of unknown affinities, such as UAIC (SCM1) 615, to be classified according to which taxonomic group it is most similar to in this discriminant morphospace. In total, 67.79% of other teeth are classified correctly when they are treated as having uncertain affinities and their measurements are used to classify them in discriminant space, indicating that this exercise returns reasonable results. Our analysis classifies the Romanian tooth as a carcharodontosaurid. Furthermore, the analysis places UAIC (SCM1) 615 within the convex hulls for carcharodontosaurids and tyrannosauroids, and the 95% confidence ellipses for carcharodontosaurids, coelophysoids, and neovenatorids.

Fourth, we ran a phylogenetic analysis by including UAIC (SCM1) 615 in the discrete character dataset of theropod dental features published by Hendrickx and Mateus (2014). The Romanian specimen was scored as a lateral tooth in this analysis. The analysis was conducted in TNT (Goloboff et al., 2008), and resulted in 224 most parsimonious trees (686 steps, consistency index of 0.338, retention index of 0.566). The strict consensus topology is moderately well resolved and places the Romanian tooth as the sister taxon to *Carcharodontosaurus* (Supplementary Material). This sister taxon pair is recovered as the sister clade to a grouping of the derived carcharodontosaurids *Mapusaurus* and *Giganotosaurus*.

Several synapomorphies support the carcharodontosaurid affinities of UAIC (SCM1) 615. The sister group relationship with *Carcharodontosaurus* is supported by two features: a roughly straight distal margin of the crown (character 68) and pronounced marginal undulations in the enamel that are well visible in normal light (character 112). The broader clade of UAIC (SCM1) 615, *Carcharodontosaurus*, *Mapusaurus*, and *Giganotosaurus* (= Carcharodontosaurinae, as defined by Brusatte and Sereno, 2008, and Carrano et al., 2012) is linked by numerous characters, including: large teeth with a crown height greater than 6 cm (character 65), a bowed or sigmoid distal carina in distal view (character 82), marginal

undulations that are at least twice as long mesiodistally as the space separating each undulation (character 111), and marginal undulations present on both mesial and distal sides of the crown (character 113).

The Romanian specimen also lacks many keystone dental synapomorphies of other theropod clades, based on the clade diagnoses of Hendrickx and Mateus (2014) and other cladistic studies that include dental characters. UAIC (SCM1) 615 does not possess the hooked distal denticles of some Abelisauridae, the strongly labially deflected distal carina and pronounced transverse enamel undulations extending across the labial and lingual tooth faces of Ceratosauridae, the incrassate teeth with apicobasal enamel flutes and deeply veined enamel surface texture of Spinosauridae, and the large transverse undulations of some basal allosauroids (Hendrickx and Mateus, 2014). It also lacks the thickened incrassate teeth of derived tyrannosauroids (Brusatte et al., 2010a) and the large and strongly hooked (or pointed) denticles of troodontids and therizinosauroids (e.g., Turner et al., 2012; Brusatte et al., 2014; Hendrickx and Mateus, 2014). The large size, as well as recurved and ziphodont shape of UAIC (SCM1) 615 is strikingly different from the non-ziphodont therizinosauroids, ornithomimosaur, alvarezsauroids, and most troodontids, which have conical, leaf-shaped, or peg-like teeth (when teeth are present) (e.g., Holtz et al., 2004a; Turner et al., 2012; Brusatte et al., 2014). Finally, besides its remarkably large size, the presence of serrations indicates that UAIC (SCM1) 615 does not belong to groups such as alvarezsauroids, oviraptorosaurs, basal troodontids, or avialans, which have unserrated crowns (e.g., Turner et al., 2012; Hendrickx and Mateus, 2014).

In summary, the four analyses all support carcharodontosaurid affinities for UAIC (SCM1) 615. Both overall tooth proportions and discrete phylogenetic characters point to a carcharodontosaurid identification, and the discriminant function analysis and phylogenetic analysis both explicitly recover the tooth as a carcharodontosaurid. For this reason we refer

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
600 this tooth to Carcharodontosauridae. Moreover, it appears to belong to a clade that unites very
601 derived and large-sized carcharodontosaurids (*Carcharodontosaurus*, *Giganotosaurus*, and
602 *Mapusaurus*), separated as such and named Carcharodontosaurinae by Brusatte and Sereno
603 (2008) and Carrano et al. (2012). The well-resolved internal topology of this clade, as
604 recovered in our analysis, is congruent with results of previous analyses based on larger sets
605 of characters from across the skeleton (e.g., Coria and Currie, 2006; Brusatte and Sereno,
606 2008; Brusatte et al., 2009; Ortega et al., 2010; Eddy and Clarke, 2011; Canale et al., 2015),
607 and offers some support for considering the Romanian carcharodontosaurid from Southern
608 Dobrogea as more closely related to the African *Carcharodontosaurus* than to the clade of
609 the South American giant carcharodontosaurids *Giganotosaurus* or *Mapusaurus*.

610 Two final notes are worth adding. First, our analyses also incorporated
611 carcharodontosaurids that are usually found to be basal within the clade, such as
612 *Acrocanthosaurus* and *Eocarcharia* (e.g., Harris, 1998; Sereno and Brusatte, 2008; Carrano
613 et al., 2012) as well as a host of other allosauroids, including members of Neovenatoridae
614 (*Neovenator*, *Australovenator* and *Fukuiraptor*), a clade that is often recovered as sister-
615 taxon to carcharodontosaurids within Carcharodontosauria (e.g., Benson et al., 2010; Carrano
616 et al., 2012; but see Novas et al., 2013; Porfiri et al., 2014, for an alternate placement of
617 neovenatorids in general). Both PCA and phylogenetic analysis clearly identified UAIC
618 (SCM1) 615 as more closely comparable morphologically to derived carcharodontosaurids
619 than to either basal carcharodontosaurids or to any other allosauroid subclade.

620 Second, our datasets also included teeth of *Erectopus*, the genus erected for
621 ‘*Megalosaurus*’ *superbus* to which UAIC (SCM1) 615 was originally referred. Again, our
622 analyses clearly indicate that there are no close morphological and morphometric similarities
623 between the two, which is in accordance with the suggestion of Carrano et al. (2012) that
624 *Erectopus* represents a non-carcharodontosaurid taxon, while our analysis identifies UAIC

(SCM1) 615 as a carcharodontosaurid. Instead, *Erectopus* groups with abelisauroids in the phylogenetic analysis. This is somewhat surprisingly, as Allain (2005) and Carrano et al. (2012) both identified *Erectopus* as a tetanuran. It should be noted, however, that Albian-aged abelisauroids are known from the same general area (eastern France) as that yielding the material referred to *Erectopus* (Accarie et al., 1995; Carrano and Sampson, 2008), raising the intriguing possibility that this taxon may represent an abelisauroid instead of an allosauroid tetanuran as suggested by Allain (2005) and Carrano et al. (2012). However, it must be remembered that this phylogenetic analysis is based on dental characters only, so it is probably more likely that *Erectopus* is a tetanuran with a dentition convergent to some extent with those of certain abelisauroids.

5.2. Body size of UAIC (SCM1) 615

One of the most salient and remarkable features of UAIC (SCM1) 615 is its large size. In the large and comprehensive sample of theropod teeth from our dataset, tooth size (estimated based on crown height – CH, and used as a rough proxy of body size) ranges from 2.2 mm (in the dromaeosaurid *Saurornitholestes* and the coelurosaur of uncertain affinities *Richardoestesia*) to 117.1 mm in the gigantic tyrannosauroid *Tyrannosaurus*. The Romanian specimen UAIC (SCM1) 615, with a CH of 85.5 mm, is ranked in the 60-80% maximum size (~ CH) range of the sample, and has a CH that is 73% of the largest tyrannosauroid teeth. Most of the teeth in the dataset (over 61% of the 966 measured teeth) are very small to small (less than 25 mm CH), and less than 10% of these fall in the 60-100% CH size categories. Teeth larger than UAIC (SCM1) 615 make up less than 5% of the total sample, and they represent only five taxa: the megalosaurid *Torvosaurus*, the tyrannosauroid *Tyrannosaurus*, the basal carcharodontosaurid *Acrocanthosaurus*, and the derived carcharodontosaurines *Carcharodontosaurus* and *Giganotosaurus*. Compared to other carcharodontosaurids, UAIC

(SCM1) 615 is smaller than the largest teeth of *Acrocanthosaurus* (9% difference), *Carcharodontosaurus* (20%), and *Giganotosaurus* (12.5%) in the dataset, but is 13% bigger than the largest tooth of *Mapusaurus*.

It is thus reasonable to conclude that UAIC (SCM1) 615 belonged to a large-sized carcharodontosaurid, comparable to, even if somewhat smaller than, the truly gigantic carcharodontosaurines *Giganotosaurus* and *Carcharodontosaurus* (Serenio et al., 1996; Calvo and Coria, 1998; Therrien and Henderson, 2007), taxa that were recovered as possible close relatives of the Romanian carcharodontosaurid by our phylogenetic analysis. This, in turn, corroborates growing evidence that very large body size was acquired very early in carcharodontosaurid history, since the earliest potential members of the clade are already of relatively large size (Rauhut, 2011). The oldest potential carcharodontosaurid is *Veterupristisaurus*, represented by isolated vertebrae that indicate an animal between 8.5 and 10 meters in total body length (compared to 11.5+ meters in *Acrocanthosaurus* and more derived carcharodontosaurids) (Rauhut, 2011). These specimens are known from the uppermost Jurassic of Tanzania, eastern Africa (Rauhut, 2011; Carrano et al., 2012; see below), predating at most ~18 million years (Mya) the occurrence of likely even larger-sized carcharodontosaurids in the Valanginian of Southern Dobrogea, Romania.

The inferred large body size of the South Dobrogean theropod is also remarkable as virtually all other dinosaur remains reported previously from Romania (both from the Early Cretaceous Cornet assemblage and the much later, end Cretaceous Hățeg Island fauna) are significantly smaller, and many have been interpreted as insular dwarfs (e.g., Weishampel et al., 1993, 2003; Benton et al., 2006, 2010; Stein et al., 2010; Ősi et al., 2014). Although other Romanian theropod dinosaurs were not particularly dwarfed (e.g. Brusatte et al., 2013), they were nonetheless small (Nopcsa, 1902; Csiki and Grigorescu, 1998; Csiki et al., 2010; Brusatte et al., 2013). This bias towards small bodied Romanian theropods was also

interpreted as a consequence of their insular habitat (Csiki and Grigorescu, 1998), as all previously reported theropod remains come from within the Carpathian Orogen, an area with an archipelago-type palaeogeography during the Cretaceous (Dercourt et al., 2000; Csontos and Vörös, 2004; Csiki-Sava et al., 2015). By contrast, UAIC (SCM1) 615 was found in shallow marine deposits bordering the emerged areas of Central Dobrogea, part of the stable cratonic areas of Europe and connected at least intermittently to the Ukrainian Shield since the Late Jurassic (Fig. 5A). Although cratonic Europe was also transformed into an archipelago of islands during much of the Cretaceous, these islands were often both larger in size and more stable in space and time than were the transient emerged areas of the Tethyan archipelagoes. As such, it is conceivable that the Southern Dobrogean carcharodontosaurid was less constrained by space or resource limitations than the Tethyan insular dinosaurs, allowing it to retain a large body size.

5.3. UAIC (SCM1) 615 and Valanginian dinosaur distribution

Besides documenting the presence of large-sized mainland carcharodontosaurids in the Lower Cretaceous of Romania, UAIC (SCM1) 615 is also important in that it fills a significant gap in our knowledge on the composition and distribution of the Early Cretaceous dinosaurs in Europe. In their review of dinosaur occurrences, Weishampel et al. (2004) listed 83 Early Cretaceous dinosaur localities spread throughout Europe, more than half of these being known from the later part (Barremian–Albian) of that epoch; only around a dozen localities were listed from each age of the early part of the Early Cretaceous (Berriasian, Valanginian, and Hauterivian). Even despite a significant increase in Early Cretaceous dinosaur discoveries in Europe in recent years (e.g., Royo-Torres et al., 2009; Cobos et al., 2010, 2014; Galton, 2009; Norman, 2010, 2013; Pereda-Suberbiola et al., 2011, 2012; Sachs and Hornung, 2013; Blows and Honeysett, 2014), these remain very strongly biased towards

western and southwestern Europe (especially the UK, France and Spain). Frustratingly, no occurrences are known from the entire central, eastern and southern Europe for the Berriasian–Hauterivian time interval except for two from Romania: the Berriasian–Valanginian locality of Cornet (e.g., Jurcsák and Popa, 1979, 1983; Jurcsák, 1982; Benton et al., 1997) in the northern Apuseni Mountains of northwestern Romania, and the carcharodontosaurid tooth (Simionescu, 1913) from the Valanginian of Cochirleni, in Southern Dobrogea, southeastern Romania we are describing here (Fig. 5A).

Our identification of the Romanian tooth as a carcharodontosaurid documents the presence of this clade in Europe in the very early Cretaceous. This is significant, as carcharodontosaurids were widely distributed tens of millions of years later, in the middle Cretaceous (Aptian to Cenomanian), in western Gondwana (Africa and South America, see below). Despite the recent discoveries documenting that the clade was also present in North America and Asia during the middle Cretaceous (e.g., Sereno et al., 1996; Currie and Carpenter, 2000; Brusatte et al., 2009, 2012), there has been only very few occurrences in Europe, most importantly the Barremian-aged *Concavenator* from Spain (Ortega et al., 2010; see below). The carcharodontosaurid tooth from Southern Dobrogea is substantially older than *Concavenator*, demonstrating that carcharodontosaurids appeared in Europe earlier than previously thought and were a long-term component of the European mainland Early Cretaceous faunas. It also suggests that habitat-related palaeobiological differentiation might have been already present between the cratonic, stable European mainland, with a dinosaur fauna made up of normal-sized (even very large) taxa, and the islands from the mobile Alpine areas of the Mediterranean Neo-Tethys, with by now dwarfed dinosaurs such as those described from the Berriasian–Valanginian Cornet assemblage in northwestern Romania (Benton et al., 2006).

This Valanginian carcharodontosaurid represents an important datapoint not only for the Romanian Lower Cretaceous, but also for that of wider Eurasia. The Valanginian is a poorly documented age in dinosaur evolution, with very few precisely dated fossil occurrences from anywhere in the world (e.g., Weishampel et al., 2004). The best record of Valanginian dinosaurs is from Europe, with fewer and less well dated occurrences known from Asia, some of which have debatable or controversial dates. These include sites in Japan (e.g., Manabe and Hasegawa, 1995; Matsukawa et al., 2006; but see Kusuhashi et al., 2009 and Evans and Matsumoto, 2015, supporting an alternative, younger age of these assemblages) and in Thailand (e.g., Buffetaut and Suteethorn, 1998, 2007, with age constraints according to Racey, 2009; Racey and Goodall, 2009). Occurrences of possible Valanginian age from China (e.g., Jerzykiewicz and Russell, 1991; Shen and Mateer, 1992; Lucas and Estep, 1998) are either poorly constrained as early Early Cretaceous, or were shown subsequently to be younger than Valanginian (Lucas, 2006; Tong et al., 2009). Rare dinosaur remains of possible Valanginian (or ‘Neocomian’) age were also reported from southern Africa (e.g., De Klerk et al., 2000) and, tentatively, from North America (e.g., Lucas, 1901; McDonald, 2011, with age assignments according to Sames et al., 2010; Cifelli et al., 2014).

As one of the two known reports of Valanginian dinosaurs in Europe east of France, the Southern Dobrogean dinosaur record fills a huge palaeogeographic gap between the western European and the eastern Asian dinosaur faunas. Moreover, none of these early Early Cretaceous dinosaur assemblages from outside Europe include carcharodontosaurids (see below), as theropods are represented by coelurosaurians interpreted either as compsognathids (Gishlick and Gauthier, 2007) or basal ornithomimosaurs (Choiniere et al., 2012) in southern Africa, metriacanthosaurid allosauroids (‘sinraptorids’) in Thailand (Buffetaut and Suteethorn, 2007), and indeterminate allosauroids (Pérez-Moreno et al., 1993), non-

carcharodontosaurid tetanurans (Carrano et al., 2012) or enantiornithine birds (Lacasa Ruiz, 1989), besides indeterminate taxa (Carrano et al., 2012), in western Europe. This may suggest that carcharodontosaurids had not achieved a wide geographic distribution by this point in time, and that their more cosmopolitan distribution came later, during the middle Cretaceous.

Finally, the presence of the Cochirleni carcharodontosaurid might hint at the presence of palaeobiogeographic provinciality between the western and the eastern parts of Europe, partly mirroring those reported from the later part of the Late Cretaceous (e.g., Le Loeuff and Buffetaut, 1995; Weishampel et al., 2010; Ősi et al., 2012; Csiki-Sava et al., 2015). In the reasonably well sampled, and significantly better known, western European dinosaur faunas, Valanginian large carnivorous dinosaurs include non-carcharodontosaurid tetanurans (*Becklespinax*), as well as indeterminate allosauroids or indeterminate theropods (often described as ‘*Megalosaurus*’ *dunkeri*, ‘*M.*’ *insignis* or ‘*M.*’ *oweni*), none of which can be referred positively to Carcharodontosauridae (Carrano et al., 2012). The apparently provincial geographic distribution of the large-bodied theropods suggests that some degree of faunal differentiation was occurring within the European mainland, most probably promoted by geographic distance. Notably, this intra-European differentiation in theropod assemblages appears to stand in contrast with the faunal homogeneity reported in the case of the ornithomimids from the UK and Romania (e.g., Galton, 2009). It is important, however, to re-emphasize at this point that the Valanginian dinosaur fossil record is both exceedingly poor and patchy, even in Europe. Accordingly, further discoveries are needed to verify and support (or contradict) the presence of such a distribution pattern pointing to palaeobiogeographic provinciality inside Europe, as the one suggested by our carcharodontosaurid identification for UAIC (SCM1) 615.

5.4. UAIC (SCM1) 615 and carcharodontosaurid evolution and palaeobiogeography

Carcharodontosauridae were long considered as an exclusively Gondwanan group of theropods (e.g., Allain, 2002; Novas et al., 2005) since their first discovery in northern Africa (e.g., Stromer, 1931), and subsequent description of a host of referred taxa from the Aptian–Cenomanian of Africa and South America (Coria and Salgado, 1995; Sereno et al., 1996; Novas et al., 2005; Coria and Currie, 2006; Brusatte and Sereno, 2007; Sereno and Brusatte, 2008; Cau et al., 2013). This view started to change with the identification of the Early Cretaceous (Aptian–Albian) *Acrocanthosaurus* from North America as a basal carcharodontosaurid (e.g., Sereno et al., 1996; Harris, 1998; Sereno 1999; Brusatte and Sereno, 2008), suggesting that the clade had a wider, Neopangean palaeobiogeographic distribution by the mid–late Early Cretaceous. Such a wide distribution, even a cosmopolitan one, was further supported by the discovery of definitive carcharodontosaurids in the Lower Cretaceous of Europe (Ortega et al., 2010), and in the upper Lower to lower Upper Cretaceous of China (Brusatte et al., 2009, 2010b, 2012; Mo et al., 2014; Lü et al., 2016).

Together, the available evidence pointed to an early, pre-mid Early Cretaceous origin of the carcharodontosaurids, followed by their dispersal across Laurasia and western Gondwana beginning at least by the Aptian (Fig. 5B), a scenario that is concordant with the tentatively suggested presence of early carcharodontosaurids in the Upper Jurassic of Tanzania, which are based on fragmentary specimens (Rauhut, 2011; Carrano et al., 2012). It is also concordant with the widespread appearance of carcharodontosaurids in the fossil record starting with the Aptian, when they are reported in Africa (*Eocarcharia*; Sereno and Brusatte, 2008), South America (Vickers-Rich et al., 1999), North America (*Acrocanthosaurus*; Stovall and Langston, 1950; Harris, 1998; Currie and Carpenter, 2000; Eddy and Clarke, 2011), Europe (Canudo and Ruiz-Omeñaca, 2003; Pereda-Suberbiola et al., 2012), and eastern Asia (*Kelmayisaurus*; Brusatte et al., 2012; Mo et al., 2014; Lü et al., 2016).

During the Albian–Turonian, carcharodontosaurids became especially abundant and diverse in Africa (*Carcharodontosaurus*, *Sauroniops*; Stromer, 1931; Sereno et al., 1996; Brusatte and Sereno, 2007; Le Loeuff et al., 2012; Cau et al., 2013; Richter et al., 2013) and South America (*Tyrannotitan*, *Giganotosaurus*, *Mapusaurus*, alongside with indeterminate carcharodontosaurids; Coria and Salgado, 1995; Calvo and Coria, 1998; Novas et al., 2005; Coria and Currie, 2006; Casal et al., 2009; Candeiro et al., 2011; Canale et al., 2015; Fig. 5B). They were still present during this time interval in other continents, as well: in North America with *Acrocanthosaurus* until the Albian (D’Emic et al., 2012), in Europe until the Cenomanian (Vullo et al., 2007; Csiki-Sava et al., 2015), and in Eastern Asia with *Shaochilong* until the Turonian (Brusatte et al., 2009, 2010b; see also Chure et al., 1999).

After dominating terrestrial ecosystems at least in Africa, South America and eastern Asia during the Albian–Turonian (Brusatte et al., 2009; Coria and Salgado, 2005; Novas et al., 2013), carcharodontosaurids were considered to disappear from the fossil record after the Turonian in both Asia (Brusatte et al., 2009) and South America (e.g., Coria and Salgado, 2005; Calvo et al., 2006; Novas et al., 2013), to be replaced by other groups of large theropods such as tyrannosaurids in parts of Laurasia and abelisaurids in parts of Gondwana. Canale et al. (2009) even cautioned against assigning isolated theropod teeth from post-Cenomanian deposits of South America to Carcharodontosauridae (e.g., Canudo et al., 2008; Casal et al., 2009; Salgado et al., 2009) due to their morphological similarity to those of the abelisaurid *Skorpiovenator*. Recently, however, more diagnostic cranial remains were reported to suggest the survival of carcharodontosaurids into the latest Cretaceous (Campanian–Maastrichtian) in Brazil (Azevedo et al., 2013).

Contrasting with this rich and relatively continuous fossil record of Carcharodontosauridae starting with the Aptian, the first half of its evolutionary history is very poorly documented (Fig. 5B). Prior to the identification of UAIC (SCM1) 615, only two

occurrences of pre-Aptian Cretaceous carcharodontosaurids were reported, one from the Barremian of Spain (Ortega et al., 2010; Gasca et al., 2014) and the other from the Barremian of Thailand (Buffetaut and Suteethorn, 2012). The Early Cretaceous *Kelmayisaurus* from Xinjiang, western China, was recognized as a carcharodontosaurid of possibly ?Valanginian to Aptian in age by Brusatte et al. (2012), but the deposits yielding these remains (the Lianmugui, or Lianmuxin, Formation of the Tugulu Group) were dated as Aptian–Albian by Eberth et al. (2001; see also Tong et al., 2009). An important temporal gap – of about 20 to 28 millions of years, according to the dates in Gradstein et al. (2012) – thus stretched between the oldest, tentatively assigned carcharodontosaurids from the Oxfordian–Tithonian of Tanzania, including the formally erected *Veterupristisaurus* (Rauhut, 2011; see also Carrano et al., 2012), and those that started to appear in the fossil record in the Barremian and then spread widely during the Aptian. Referral of UAIC (SCM1) 615 to Carcharodontosauridae partially fills this frustrating gap, effectively halving this shadowy period in the evolutionary history of the group.

Furthermore, our analyses tentatively cluster the Dobrogean theropod with the derived members of the Carcharodontosaurinae to the exclusion of the more basal, but significantly younger non-carcharodontosaurine carcharodontosaurids *Eocarcharia* and *Acrocanthosaurus*. If this placement is correct, then the Romanian tooth indicates that Carcharodontosaurinae diverged from other carcharodontosaurids considerably earlier than hitherto recognized.

The previously known fossil record of the clade suggested that Carcharodontosaurinae originated sometime between the Aptian and Albian, as basal carcharodontosaurids (*Acrocanthosaurus*, *Concavenator*, *Eocarcharia*) were moderately diverse in the Barremian–Aptian, followed by the appearance of many fossils of carcharodontosaurines beginning in the Albian (Fig. 5B). The proposed affinities of the oldest carcharodontosaurid material – including isolated teeth referred to as ‘*Megalosaurus*’ *ingens* – from the east African Upper

Jurassic, considered to be reminiscent of the Aptian–Albian *Acrocanthosaurus* (Rauhut, 2011), was also consistent with this evolutionary scenario. Now, our identification of UAIC (SCM1) 615 as a carcharodontosaurid dinosaur sharing important dental apomorphies with the derived Carcharodontosaurinae advocates the emergence of this clade (or at least the very large size and dental morphology characterizing it) well before the Albian, during or even before the Valanginian, and relegates taxa such as *Eocarcharia*, *Acrocanthosaurus* and *Concavenator* (the dentition of *Shaochilong* is unknown) as late-surviving members of the basal carcharodontosaurid radiation, with a relatively plesiomorphic dentition.

Besides shifting the emergence of the carcharodontosaurines earlier in time, identification of UAIC (SCM1) 615 as a carcharodontosaurid also has interesting palaeobiogeographic implications. As already noted, recent discoveries show that Carcharodontosauridae is not an endemic Gondwanan clade as was once proposed (e.g., Novas et al., 2005), with the identification of its widespread, Pangaeian distribution during the late Early Cretaceous (Sereno et al., 1996; Harris, 1998; Chure et al., 1999; Sereno, 1999; Brusatte and Sereno, 2008; Ortega et al., 2010; Brusatte et al., 2009, 2012; Mo et al., 2014). However, within Carcharodontosauridae itself, some palaeogeographic patterns have been widely accepted. For example, it has been widely acknowledged that Carcharodontosaurinae is a endemic subclade of Gondwanan carcharodontosaurids (e.g., Sereno 1999; Holtz et al., 2004b; Brusatte and Sereno, 2007; Sereno and Brusatte, 2008; Novas et al., 2013), as previously all its recognized members were restricted strictly to either Africa (Stromer, 1931; Sereno et al., 1996; Brusatte and Sereno, 2007) or South America (Coria and Salgado, 1995; Novas et al., 2005; Coria and Currie, 2006). Moreover, intra-clade relationships of Carcharodontosaurinae were still adhering to patterns of continental fragmentation and vicariant evolution, with a basal split between the Albian–Cenomanian African *Carcharodontosaurus* and the Giganotosaurini, uniting the similarly Albian–Cenomanian

southern South American *Giganotosaurus* and *Mapusaurus* (together with *Tyrannotitan*, if this taxon is also recovered within Carcharodontosaurinae; e.g., Novas et al., 2005, 2013).

This scenario is now challenged by our finding that the Southern Dobrogean carcharodontosaurid UAIC (SCM1) 615 may nest inside Carcharodontosaurinae. If true, such an affinity would suggest that the origin of Carcharodontosaurinae was not a southern, vicariant by-product of the Gondwana-Laurasia separation, a major palaeogeographic event that is considered to have been well underway by the end of the Jurassic, and essentially completed by the mid-Early Cretaceous (see Weishampel et al., 2010). Indeed, during this time palaeogeographic connections and faunal interactions were virtually non-existent between the northern Tethyan (European) and southern Tethyan (western Gondwanan, but essentially African) areas of the Mediterranean (e.g., Canudo et al., 2009), which makes a vicariant hypothesis intuitive. However, if the Romanian tooth represents a carcharodontosaurine, then it implies a much more complicated palaeogeographic history of the clade, which is not so clearly linked to continental breakup.

The palaeogeographic position of the Southern Dobrogean carcharodontosaurine in cratonic Europe, north of the Neo-Tethys, together with its significantly older age compared to other carcharodontosaurines, could indicate that separation of the carcharodontosaurine lineage took part in Europe and not in western Gondwana as previously assumed. This would also mean that representatives of this lineage were subsequently – after the Barremian – introduced to Africa and South America via trans-Tethyan dispersal, most probably at a time when faunal interactions between the southern and northern margins of the Mediterranean Tethys were resumed, after the early Barremian (Canudo et al., 2009).

Alternatively, it can be hypothesized that appearance of carcharodontosaurines in Southern Dobrogea is a consequence of southern immigration originating in western Gondwana, often considered the place of origin for this clade. However, this scenario has

several potential caveats. Although Europe has been considered as forming part of a larger Eurogondwanan palaeobioprovince during the early Early Cretaceous (Ezcurra and Agnolín, 2012), and occasional trans-Tethyan faunal connections have been recognized between Africa and Europe during Late Jurassic to Early Cretaceous times (e.g., Gheerbrant and Rage, 2006), these interchanges either pre-dated the Berriasian (e.g., Gardner et al., 2003; Knoll and Ruiz-Omeñaca, 2009), or post-dated the Barremian (Canudo et al., 2009; Torcida Fernández-Baldor et al., 2011), with no positive evidence for actual faunal exchanges taking place during the ‘Neocomian’ (Berriasian–Hauterivian) time interval.

More recently, some potential evidence has emerged for Gondwana-to-Europe interchange during the ‘Neocomian’. The presence of the basal rebbachisaurid *Histriasaurus* (Dalla Vecchia, 1998) in the upper Hauterivian–lower Barremian of Croatia has been cited as indicative of very early and very rapid northward dispersal of this clade from western Gondwana (southern South America; Carballido et al., 2012; Fanti et al., 2015). Timing of this particular dispersal event was even constrained to the Berriasian–Valanginian time interval (Fanti et al., 2015), which makes it roughly contemporaneous with the record of the Southern Dobrogean carcharodontosaurine. It was also suggested, however, that dispersal of the line leading to *Histriasaurus* was mediated by the northward drift of the Apulian Microplate (= Adria; see Bosselini, 2002), a continental sliver acting as a passive transportation mechanism (‘Noah’s Ark’; McKenna, 1973) for basal rebbachisaurids after its separation from mainland Africa (e.g., Torcida Fernández-Baldor et al., 2011). Furthermore, the palaeogeographical separation between Africa and Adria (and thus the effective movement of the presumed ark) is considered to be at most an incipient one during the Early Cretaceous by Bossellini (2002) and Zarcone et al. (2010), with spatial continuity still present between the two landmasses, while deep-water basins continued to separate Adria from the European Craton. Accordingly, although the presence of *Histriasaurus* can represent a case

of northward range extension of rebbachisaurids during the Berriasian–Valanginian, it took place not strictly speaking into Europe, but only reached the northernmost extremity of Adria, a northerly peninsular extension of the African mainland. It was only starting with the Barremian that rebbachisaurids dispersed as far north as the European cratonic areas, including Iberia and the British Isles (Mannion, 2009; Mannion et al., 2011; Torcida Fernández-Baldor et al., 2011), a time when faunal interchanges between Europe and Africa are considered to have been well underway (e.g., Gheerbrant and Rage, 2006; Canudo et al., 2009).

Unlike *Histriasaurus*, the taxon represented by UAIC (SCM1) 615 was an inhabitant of the European mainland. It is thus unclear to what extent the example of rebbachisaurid range extension into (present-day) Europe during the early Early Cretaceous, as potentially testified by the discovery of the Croatian taxon, would also be applicable for the Southern Dobrogean carcharodontosaurine. The available evidence suggests that these two cases are very different, and that faunal connections during this time interval are not documented between the African and European cratons as already pointed out by Gheerbrant and Rage (2006).

Absence of documented faunal interactions weakens support for a scenario of south-to-north immigration of derived carcharodontosaurines in Europe at the very beginning of the Cretaceous, and would argue instead for a local, European development to explain the presence of a Valanginian carcharodontosaurine in Southern Dobrogea. The pre-Barremian presence of carcharodontosaurids in Europe is also consistent with their appearance in the Barremian–Aptian fossil record of Eastern Asia, with Europe acting as a stepping stone in the eastward dispersal of the clade. Similarly, the presence of Aptian carcharodontosaurids in North America likely requires the presence of pre-Aptian members of the clade in Europe, since faunal exchanges between these two landmasses are known to have been halted before

the Aptian (e.g., Kirkland et al., 1999). Interestingly, it appears that only basal carcharodontosaurids were able to spread into the northern Laurasian landmasses, while the derived carcharodontosaurines dispersed exclusively across the Neo-Tethys, into western Gondwana. The causes of these distribution patterns remain as yet unknown, and further support – in the form on new carcharodontosaurid discoveries from the early-middle part of the Early Cretaceous – is required to better uphold such a scenario.

We finally reiterate that if the Romanian tooth does not belong to a carcharodontosaurine, but instead is artefactually grouping with them in the phylogenetic analysis because of the very incomplete nature of the material, then the traditional story of Carcharodontosaurinae as a product of vicariant evolution driven by the breakup of Pangea will remain strongly supported. However, even in such case UAIC (SCM1) 615 would still record the presence of early-occurring large carcharodontosaurid theropods with a very characteristic carcharodontosaurine-type dentition in the eastern part of the European craton, adding to known early Early Cretaceous theropod (and dinosaur) diversity, and potentially documenting dinosaur faunal provinciality in Europe and worldwide.

6. Conclusions

We re-describe and interpret the affinities of one of the most significant historical dinosaurian specimens of Romania, an isolated but well-preserved theropod tooth from Southern Dobrogea. Our extensive analyses suggest carcharodontosaurid relationships for this tooth, while the available evidence – including novel calcareous nannoplankton sampling – supports its Valanginian age. The Southern Dobrogean theropod tooth represents the oldest record of Carcharodontosauridae in the Cretaceous, and the second oldest globally, eclipsed only by a collection of isolated specimens from the Upper Jurassic of eastern Africa. As one of the only two known Valanginian dinosaurian occurrences from Central and Eastern Europe, this

record advances our understanding of European dinosaur distribution during the early Early Cretaceous, and also fills an important palaeogeographic gap between Western European and Eastern Asian dinosaurian assemblages of the Valanginian.

Based on dental apomorphies, our analyses further identify UAIC (SCM1) 615 as a possible member of Carcharodontosaurinae, a subclade of derived and gigantic carcharodontosaurids formerly known to be restricted to the Albian–Cenomanian of western Gondwana (Africa and South America). If this finding is correct, the Southern Dobrogean specimen documents the emergence of Carcharodontosaurinae earlier than previously recognized, thus also indicating an earlier acquisition of their characteristically large size.

Based on currently known palaeogeographic and chronostratigraphic constraints on the evolution of Carcharodontosauridae, it appears that not only did this clade have a wide distribution, but that crucial events of its evolutionary history such as the emergence of the derived carcharodontosaurines took place north of the Tethys, in cratonic Europe, instead of western Gondwana and as the result of vicariant evolution driven by the Gondwana-Laurasia split, as was formerly suggested. In such a case, instead of endemic evolution the emergence of the western Gondwanan mid-Cretaceous carcharodontosaurines was the result of a north-to-south trans-Tethyan dispersal that took place somewhere between the Valanginian and the Aptian. Recognizing a potential carcharodontosaurine dispersal event from Europe into western Gondwana adds further support for the presence of important palaeogeographic ties between the two realms during the second half of the Early Cretaceous.

Acknowledgements

This research was supported by the National Research Council of Romania (CNCS) grant PN-IIID-PCE-2011-3-0381 and a Sepkoski grant of the Paleontological Society for Z.Cs.-S.

S.L.B. is supported by a Marie Curie Career Integration Grant EC630652, the Division of Paleontology of the American Museum of Natural History, and the School of GeoSciences of the University of Edinburgh. He thanks Mátyás Vremir, Radu Totoianu, and Mark Norell for many hours of fun discussion on Romanian fossils, and for supporting his work and travel in Romania. We thank Mihai Brânzilă and Paul Țibuleac (UAIC) for access to the specimen, for allowing us to collect samples for the nannoplankton analyses, and for their help and collegiality during our visit to Iași, as well as Ilie Turculeț for sharing information about the history of the specimen. Mihaela C. Melinte-Dobrinescu has gracefully analyzed the nannoplankton samples derived from UAIC (SCM1) 615; her contribution was essential in assessing the age of the specimen. Finally, we thank the reviewers Eric W.A. Mulder (Denekamp, the Netherlands) and Xabier Pereda-Suberbiola (Bilbao, Spain), as well as Associated Editor Elena Jagt-Yazykova, for their useful comments and suggestions that helped improve previous versions of the manuscript.

References

- Accarie, H., Beaudoin, B., Dejax, J., Friès, G., Michard, J.G., Taquet, P., 1995. Découverte d'un dinosaure théropode nouveau (*Genusaurus sisteronis* n. g., n. sp.) dans l'Albien marin de Sisteron (Alpes-de-Haute-Provence, France) et extension au Crétacé inférieur de la lignée cératosaurienne. *Comptes Rendus de l'Académie des Sciences Paris, Ila* 320, 327–334.
- Allain, R., 2002. Discovery of a megalosaur (Dinosauria, Theropoda) in the Middle Bathonian of Normandy (France) and its implications for the phylogeny of basal Tetanurae. *Journal of Vertebrate Paleontology* 22, 548–563.
- Allain, R., 2005. The enigmatic theropod dinosaur *Erectopus superbus* (Sauvage, 1882) from the Lower Albian of Louppy-le-Chateau (Meuse, France). In: Carpenter, K. (Ed.), *The Carnivorous Dinosaurs*. Indiana University Press, Bloomington, pp. 72–86.

- Avram, E., Szasz, L., Antonescu, E., Baltreş, A., Iva, M., Melinte, M., Neagu, T., Rădan, S.,
Tomescu, C., 1993. Cretaceous terrestrial and shallow marine deposits in northern South
Dobrogea (SE Romania). *Cretaceous Research* 14, 265–305.
- Avram, E., Costea, I., Dragastan, O., Muşiu, R., Neagu, T., Şindilar, V., Vinogradov, C.,
1996. Distribution of the Middle-Upper Jurassic and Cretaceous facies in the Romanian
eastern part of the Moesian Platform. *Revue Roumaine de Géologie* 39-40, 3–33.
- Azevedo, R.P.F. de, Simbras, F.M., Furtado, M.R., Candeiro, C.R.A., Bergqvist, L.P., 2013.
First Brazilian carcharodontosaurid and other new theropod dinosaur fossils from the
Campanian-Maastrichtian Presidente Prudente Formation, São Paulo State, southeastern
Brazil. *Cretaceous Research* 40, 131–142.
- Benson, R.B.J., Barrett, P.M., Powell, H.P., Norman, D.B., 2008. The taxonomic status of
Megalosaurus bucklandii (Dinosauria, Theropoda) from the Middle Jurassic of Oxfordshire,
UK. *Palaeontology* 51, 419–424.
- Benson, R.B.J., Carrano, M.T., Brusatte, S.L., 2010. A new clade of archaic large-bodied
predatory dinosaurs (Theropoda: Allosauroidae) that survived to the latest Mesozoic.
Naturwissenschaften 97, 71–78.
- Benton, M.J., Cook, E., Grigorescu, D., Popa, E., Tallódi, E., 1997. Dinosaurs and other
tetrapods in an Early Cretaceous bauxite-filled fissure, northwestern Romania.
Palaeogeography, Palaeoclimatology, Palaeoecology 130, 275–292.
- Benton, M.J., Minter, N.J., Posmoşanu, E., 2006. Dwarfing in ornithomimid dinosaurs from the
Early Cretaceous of Romania. In: Csiki, Z. (Ed.), *Mesozoic and Cenozoic Vertebrates and
Paleoenvironments; Tributes to the Career of Prof. Dan Grigorescu*. Ars Docendi, Bucharest,
pp. 79–87.

- Benton, M.J., Csiki, Z., Grigorescu, D., Redelstorff, R., Sander, M., Stein, K., Weishampel, D.B., 2010. Dinosaurs and the island rule: the dwarfed dinosaurs from Hațeg Island. *Palaeogeography, Palaeoclimatology, Palaeoecology* 293, 438–454.
- Blows, W.T., Honeysett, K., 2014. First Valanginian *Polacanthus foxii* (Dinosauria, Ankylosauria) from England, from the Lower Cretaceous of Bexhill, Sussex. *Proceedings of the Geologists' Association* 125, 233–251.
- Bosselini, A., 2002. Dinosaurs “re-write” the geodynamics of the eastern Mediterranean and the paleogeography of the Apulia Platform. *Earth-Science Reviews* 59, 211–234.
- Brânzilă, M. (ed.), 2010. *Academicianul Ion Simionescu - savant și dascăl al neamului*. Ed. Universității "Alexandru Ioan Cuza", Iași, 182 pp.
- Brusatte, S.L., Sereno, P.C., 2007. A new species of *Carcharodontosaurus* (Dinosauria: Theropoda) from the Cenomanian of Niger and a revision of the genus. *Journal of Vertebrate Paleontology* 27, 902–916.
- Brusatte, S.L., Sereno, P.C., 2008. Phylogeny of Allosauroidea (Dinosauria: Theropoda): comparative analysis and resolution. *Journal of Systematic Palaeontology* 6, 155–182.
- Brusatte, S. L., Clark, N.D.L., 2015. Theropod dinosaurs from the Middle Jurassic (Bajocian-Bathonian) of Skye, Scotland. *Scottish Journal of Geology* 51, 157-164. doi: 10.1144/sjg2014-022
- Brusatte, S., Benson, R.B.J., Carr, T.D., Williamson, T.E., Sereno, P.C., 2007. The systematic utility of theropod enamel wrinkles. *Journal of Vertebrate Paleontology* 27, 1052–1056.
- Brusatte, S.L., Benson, R.B.J., Chure, D.J., Xu, X., Sullivan, C., Hone, D.W.E., 2009. The first definitive carcharodontosaurid (Dinosauria: Theropoda) from Asia and the delayed ascent of tyrannosaurids. *Naturwissenschaften* 96, 1051–1058.
- Brusatte, S.L., Norell, M.A., Carr, T.D., Erickson, G.M., Hutchinson, J.R., Balanoff, A.M., Bever, G.S., Choiniere, J.N.,

- 1071 Makovicky, P.J., Xu, X. , 2010a. Tyrannosaur paleobiology: new research on ancient
1072 exemplar organisms. *Science* 329, 1481–1485.
- 1073 Brusatte, S., Chure, D.J., Benson, R.B.J., Xu, X., 2010b. The osteology of *Shaochilong*
1074 *maortuensis*, a carcharodontosaurid (Dinosauria: Theropoda) from the Late Cretaceous of
1075 Asia. *Zootaxa* 2334, 1–46.
- 1076 Brusatte, S.L., Benson, R.B.J., Xu, X., 2012. A reassessment of *Kelmayisaurus petrolicus*, a
1077 large theropod dinosaur from the Early Cretaceous of China. *Acta Palaeontologica Polonica*
1078 57, 65–72.
- 1079 Brusatte, S.L., Vremir, M., Csiki-Sava, Z., Turner, A.H., Watanabe, A., Erickson, G.M.,
1080 Norell, M.A., 2013. The osteology of *Balaur bondoc*, an island-dwelling dromaeosaurid
1081 (Dinosauria: Theropoda) from the Late Cretaceous of Romania. *Bulletin of the American*
1082 *Museum of Natural History* 374, 3–100. doi: 10.1206/798.1
- 1083 Brusatte, S.L., Lloyd, G.T., Wang, S.C., Norell, M.A., 2014. Gradual assembly of avian body
1084 plan culminated in rapid rates of evolution across the dinosaur-bird transition. *Current*
1085 *Biology* 24, 2386–2392.
- 1086 Buckland, W., 1824. Notice on the *Megalosaurus* or great fossil lizard of Stonesfield.
1087 *Transactions of the Geological Society* 21, 390–397.
- 1088 Buffetaut, E., Suteethorn, V., 1998. Early Cretaceous dinosaurs from Thailand and their
1089 bearing on the early evolution and biogeographical history of some groups of Cretaceous
1090 dinosaurs. In: Lucas, S.G., Kirkland, J.I., Estep, J.W. (Eds.), *Lower and Middle Cretaceous*
1091 *Terrestrial Ecosystems*, New Mexico Museum of Natural History and Science Bulletin 14,
1092 pp. 205–210.
- 1093 Buffetaut, E., Suteethorn, V., 2007. A sinraptorid theropod (Dinosauria: Saurischia) from the
1094 Phu Kradung Formation of northeastern Thailand. *Bulletin de la Société Géologique de*
1095 *France* 178, 497–502.

- 1096 Buffetaut, E., Suteethorn, V., 2012. A carcharodontid theropod (Dinosauria, Saurischia) from
1
2 1097 the Sao Khua Formation (Early Cretaceous, Barremian) of Thailand. In: Royo-Torres, R.,
3
4 1098 Gascó, F., Alcalá, L. (Eds.), 10th Annual Meeting of the European Association of Vertebrate
5
6
7 1099 Palaeontologists. ¡Fundamental! 20, Teruel, pp. 27–30.
8
9
10 1100 Calvo, J.O., Coria, A., 1998. New specimen of *Giganotosaurus carolinii* (Coria & Salgado,
11
12 1101 1995), supports it as the largest theropod ever found. *Gaia* 15, 117–122.
13
14 1102 Calvo, J.O., Gandossi, P., Porfiri, J.D., 2006. Dinosaur faunal replacement during
15
16
17 1103 Cenomanian times in Patagonia, Argentina. In: Evans, S.E., Barrett, P.M. (Eds.), 9th
18
19 1104 Mesozoic Terrestrial Ecosystems and Biota. Manchester, UK, pp. 17–20.
20
21
22 1105 Canale, J.I., Scanferla, C.A., Agnolin, F.L., Novas, F.E., 2009. New carnivorous dinosaur
23
24 1106 from the Late Cretaceous of NW Patagonia and the evolution of abelisaurid theropods.
25
26
27 1107 *Naturwissenschaften* 96, 409–414.
28
29 1108 Canale, J.I., Novas, F.E., Pol, D., 2015. Osteology and phylogenetic relationships of
30
31 1109 *Tyrannotitan chubutensis* Novas, de Valais, Vickers-Rich and Rich, 2005 (Theropoda:
32
33
34 1110 Carcharodontosauridae) from the Lower Cretaceous of Patagonia, Argentina. *Historical*
35
36 1111 *Biology* 27, 1–32. doi: 10.1080/08912963.2013.861830.
37
38
39 1112 Candeiro, C.R.A., Fanti, F., Therrien, F., Lamanna, M.C., 2011. Continental fossil vertebrates
40
41 1113 from the mid-Cretaceous (Albian-Cenomanian) Alcântara Formation, Brazil, and their
42
43
44 1114 relationship with contemporaneous faunas from North Africa. *Journal of African Earth*
45
46 1115 *Sciences* 60, 79–92.
47
48
49 1116 Canudo, J.I., Ruiz-Omeñaca, J.I., 2003. Los restos directos de dinosaurios terópodos
50
51 1117 (excluyendo Aves) en España. *Ciencias de la Tierra* 26, 347–373.
52
53
54 1118 Canudo, J.I., Filippi, L.S., Salgado, L., Garrido, A.C., Cerda, I.A., Garcia, R., Otero, A.,
55
56 1119 2008. Theropod teeth associated with a sauropod carcass in the Upper Cretaceous (Plottier
57
58 1120 Formation) of Rincón de los Sauces. In: *Colectivo Arqueológico y Paleontológico de Salas de*
59
60

- los Infantes (Ed.), Actas de las IV Jornadas Internacionales sobre Paleontología de Dinosaurios y su Entorno, Salas de los Infantes, Burgos, pp. 321–330.
- Canudo, J.I. Barco, J.L., Pereda Suberbiola, X., Ruiz-Omeñaca, J.I., Salgado, L., Torcida Fernández-Baldor, F., Gasulla, J.M., 2009. What Iberian dinosaurs reveal about the bridge said to exist between Gondwana and Laurasia in the Early Cretaceous. *Bulletin de la Société Géologique de France*, 180, 5–11.
- Carballido, J.L., Salgado, L., Pol, D., Canudo, J.I., Garrido, A., 2012. A new basal rebbachisaurid (Sauropoda, Diplodocoidea) from the Early Cretaceous of the Neuquén Basin; evolution and biogeography of the group. *Historical Biology* 24, 631–654.
- Carrano, M.T., Sampson, S.D., 2008. The phylogeny of Ceratosauria (Dinosauria: Theropoda). *Journal of Systematic Palaeontology* 6, 183–236.
- Carrano, M.T., Benson, R.B.J., Sampson, S.D., 2012. The phylogeny of Tetanurae (Dinosauria: Theropoda). *Journal of Systematic Palaeontology* 10, 211–300.
- Carvalho, I.S., Campos, A.C.A., Nobre, P.H., 2005. *Baurusuchus salgadoensis*, a new Crocodylomorpha from the Bauru Basin (Cretaceous), Brazil. *Gondwana Research* 8, 11–30.
- Casal, G., Candeiro, C.R.A., Martinez, R., Ivany, E., Ibiricu, L., 2009. Dientes de Theropoda (Dinosauria: Saurischia) de la Formación Bajo Barreal, Cretácico Superior, Provincia del Chubut, Argentina. *Géobios* 42, 553–560.
- Cau, A., Dalla Vecchia, F.M., Fabbri, M., 2013. A thick-skulled theropod (Dinosauria, Saurischia) from the Upper Cretaceous of Morocco with implications for carcharodontosaurid cranial evolution. *Cretaceous Research* 40, 251–260.
- Choiniere, J.N., Forster, C.A., De Klerk, W.J., 2012. New information on *Nqwebasaurus thwazi*, a coelurosaurian theropod from the Early Cretaceous Kirkwood Formation in South Africa. *Journal of African Earth Sciences* 71–72, 1–17.

- Chure, D.J., Manabe, M., Tanimoto, M., Tomida, Y., 1999. An unusual theropod tooth from the Mifune Group (Late Cenomanian to Early Turonian), Kumamoto, Japan. In: Tomida, Y., Rich, T.H., Vickers-Rich, P. (Eds.), Proceedings of the Second Gondwanan Dinosaur Symposium. National Science Museum Monographs 15, Tokyo, pp. 291–296.
- Cifelli, R.L., Davis, B.M., Sames, B., 2014. Earliest Cretaceous mammals from the western United States. *Acta Palaeontologica Polonica* 59, 31–52. doi:10.4202/app.2012.0089.
- Cobos, A., Royo-Torres, R., Luque, L., Alcalá, L., Mampel, L., 2010. An Iberian stegosaurs paradise: The Villar del Arzobispo Formation (Tithonian–Berriasian) in Teruel (Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology* 293, 223–236.
- Cobos, A., Lockley, M.G., Gascó, F., Royo–Torres, R., Alcalá, L., 2014. Megatheropods as apex predators in the typically Jurassic ecosystems of the Villar del Arzobispo Formation (Iberian Range, Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology* 399, 31–41. doi: 10.1016/j.palaeo.2014.02.008.
- Codrea, V., Vremir, M., Jipa, C., Godefroit, P., Csiki, Z., Smith, T., Fărcaș, C., 2010. More than just Nopcsa’s Transylvanian dinosaurs: A look outside the Hațeg Basin. *Palaeogeography, Palaeoclimatology, Palaeoecology* 293, 391–405. doi: 10.1016/j.palaeo.2009.10.027.
- Codrea, V., Godefroit, P., Smith, T., 2012. First discovery of Maastrichtian (latest Cretaceous) terrestrial vertebrates in Rusca Montană Basin (Romania). In: Godefroit, P. (Ed.) *Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems*. Indiana University Press, Bloomington, pp. 570–581.
- Colbert, E.H., 1946. *Sebecus*, representative of a peculiar suborder of fossil Crocodilia from Patagonia. *Bulletin of the American Museum of Natural History* 87(4), 217–270.
- Coria, R.A., Salgado, L., 1995. A new giant carnivorous dinosaur from the Cretaceous of Patagonia. *Nature* 377, 224–226.

- Coria, R.A., Salgado, L., 2005. Mid-Cretaceous turnover of saurischian dinosaur communities: evidence from the Neuquén Basin. In: Veiga, G.D., Spalletti, L.A., Howell, J.A., Schwartz, E. (Eds.), *The Neuquén Basin, Argentina: a case study in sequence stratigraphy and basin dynamics*. Geological Society, London, Special Publications 252, pp. 317–327.
- Coria, R.A., Currie, P.J., 2006. A new carcharodontosaurid (Dinosauria, Theropoda) from the Upper Cretaceous of Argentina. *Geodiversitas* 28, 71–118.
- Covacef, Z., 1995. Pionieri ai culturii românești în Dobrogea; Ioan Cotovu și Vasile Cotovu. *Analele Dobrogei* I(1), 127–134.
- Császár, G., Kollányi, K., Lantos, M., Lelkes, G. and Tardine Filácz, E., 2000. A Hidasivölgyi Márga Formáció kora és képződési környezete. *Földtani Közlöny* 130(4), 695–723.
- Csiki, Z., Grigorescu, D., 1998. Small theropods of the Late Cretaceous of the Hațeg Basin (Western Romania) - an unexpected diversity at the top of the food chain. *Oryctos* 1, 87–104.
- Csiki, Z., Vremir, M., Brusatte, S.L., Norell, M.A., 2010. An aberrant island-dwelling theropod dinosaur from the Late Cretaceous of Romania. *Proceedings of the National Academy of Sciences* 107, 15357–15361.
- Csiki-Sava, Z., Codrea, V., Vasile, Ș., 2013. Early Cretaceous dinosaur remains from Dobrogea (southeastern Romania). In: Picot, L. (Ed.), *Abstracts, 11th Annual Meeting of the European Association of Vertebrate Palaeontologists, Villers-sur-Mer, France*, pp. 28.
- Csiki-Sava, Z., Buffetaut, E., Ősi, A., Pereda-Suberbiola, X., Brusatte, S.L., 2015. Island life in the Cretaceous - faunal composition, biogeography, evolution, and extinction of land-living vertebrates on the Late Cretaceous European archipelago. *Zookeys* 469, 1–161. doi: 10.3897/zookeys.469.8439.

- 1194 Csontos, L., Vörös, A., 2004. Mesozoic plate tectonic reconstruction of the Carpathian
1195 region. *Palaeogeography, Palaeoclimatology, Palaeoecology* 210, 1–56. doi:
1196 10.1016/j.palaeo.2004.02.033.
- 1197 Currie, P.J., Carpenter, K., 2000. A new specimen of *Acrocanthosaurus atokensis*
1198 (Theropoda, Dinosauria) from the Lower Cretaceous Antlers Formation (Lower Cretaceous,
1199 Aptian) of Oklahoma, USA. *Geodiversitas* 22, 207–246.
- 1200 Currie, P.J., Rigby, J.K., Sloan, R.E., 1990. Theropod teeth from the Judith River Formation
1201 of southern Alberta, Canada. In: Carpenter, K., Currie, P.J. (Eds.), *Dinosaur Systematics:
1202 Approaches and Perspectives*. Cambridge University Press, Cambridge, Boston, pp. 107–125.
- 1203 Dalla Vecchia, F.M., 1998. Remains of Sauropoda (Reptilia, Saurischia) in the Lower
1204 Cretaceous (Upper Hauterivian/Lower Barremian) limestones of SW Istria (Croatia).
1205 *Geologia Croatica* 5, 105–134.
- 1206 Dames, W., 1884. *Megalosaurus dunkeri*. *Sitzungsberichte Gesellschaft Naturforschender
1207 Freunde zu Berlin* 1884, 186–188.
- 1208 De Klerk, W.J., Forster, C.A., Sampson, S.D., Chinsamy, A., Ross, C.F., 2000. A new
1209 coelurosaurian dinosaur from the Early Cretaceous of South Africa. *Journal of Vertebrate
1210 Paleontology* 20, 324–332.
- 1211 D’Emic, M.D., Melstrom, K.M., Eddy, D.R., 2012. Paleobiology and geographic range of the
1212 large-bodied Cretaceous theropod dinosaur *Acrocanthosaurus atokensis*. *Palaeogeography,
1213 Palaeoclimatology, Palaeoecology* 333–334, 13–23.
- 1214 Dercourt, J., Gaetani, M., Vrielynck, B., Barrier, E., Biju-Duval, B., Brunet, M., Cadet, J.P.,
1215 Crasquin, S., Săndulescu, M., Eds. (2000) *Atlas Peri-Tethys Palaeogeographical Maps.
1216 CCGM/CGMW, Paris*, 269 pp.
- 1217 Dinu, C., Grădinaru, E., Stoica, M., Diaconescu, V., 2007. *Dobrogea 2007 Field Trip
1218 Preparation and Assistance*. University of Bucharest, 123 pp.

- 1219 Dragastan, O., Neagu, T., Bărbulescu, A., Pană, I., 1998. Jurassicul și Cretacicul din Dobrogea
Centrală și de Sud. Bucharest, 249 pp.
- 1220
- 1221 Dragastan, O.N., Antoniadă, C., Stoica, M., 2014. Biostratigraphy and zonation of the Lower
Cretaceous succession from Cernavodă-lock section, South Dobrogea, eastern part of the
Moesian Platform (Romania). Carpathian Journal of Earth and Environmental Sciences 9(1),
231–260.
- 1222
- 1223
- 1224
- 1225 Dyke, G.J., Benton, M.J., Posmoșanu, E., Naish, D., 2011. Early Cretaceous (Berriasian)
birds and pterosaurs from the Cornet bauxite mine, Romania. Palaeontology 54, 79–95.
- 1226
- 1227 Eberth, D.A., Brinkman, D.B., Chen, P.-J., Yuan, F.-T., Wu, X.-C., Li, G., Cheng, X.-S.,
2001. Sequence stratigraphy, paleoclimate patterns, and vertebrate fossil preservation in
Jurassic–Cretaceous strata of the Juggar Basin, Xinjiang Autonomous Region, People's
Republic of China. Canadian Journal of Earth Sciences 38, 1627–1644.
- 1228
- 1229
- 1230
- 1231 Eddy, D.R., Clarke, J.A., 2011. New information on the cranial anatomy of
Acrocanthosaurus atokensis and its implications for the phylogeny of Allosauroidea
(Dinosauria: Theropoda). PLoS ONE 6(3), e17932. doi:10.1371/journal.pone.0017932.
- 1232
- 1233
- 1234 Evans, S.E., Matsumoto, R., 2015. An assemblage of lizards from the Early Cretaceous of
Japan. Palaeontologia Electronica 18.2.36A, 1–36.
- 1235
- 1236 Ezcurra, M.D., Agnolín, F.L., 2012. A new global palaeobiogeographical model for the Late
Mesozoic and Early Tertiary. Systematic Biology 61, 553–566. doi:10.1093/sysbio/syr115.
- 1237
- 1238 Fanti, F., Cau, A., Cantelli, L., Hassine, M., Auditore, M., 2015. New information on
Tataouinea hannibalis from the Early Cretaceous of Tunisia and implications for the tempo
and mode of rebbachisaurid sauropod evolution. PLoS ONE 10(4), e0123475.
doi:10.1371/journal.pone.0123475.
- 1239
- 1240
- 1241

- Galton, P.M., 2009. Notes on Neocomian (Lower Cretaceous) ornithopod dinosaurs from England - *Hypsilophodon*, *Valdosaurus*, “*Camptosaurus*”, “*Iguanodon*” - and referred specimens from Romania and elsewhere. *Revue de Paléobiologie* 28, 211–273.
- Gardner, J.D., Evans, S.E., Sigogneau-Russell, D., 2003. New albanerpetontid amphibians from the Early Cretaceous of Morocco and Middle Jurassic of England. *Acta Palaeontologica Polonica* 48, 301–319.
- Gasca, J.M., Canudo, J.I., Moreno-Azanza, M., 2014. A large-bodied theropod (Tetanurae: Carcharodontosauria) from the Mirambel Formation (Barremian) of Spain. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen* 273, 13–23. doi: 10.1127/0077-7749/2014/0413.
- Gheerbrant, E., Rage, J.-C., 2006. Paleobiogeography of Africa: How distinct from Gondwana and Laurasia? *Palaeogeography, Palaeoclimatology, Palaeoecology* 241, 224–246. doi:10.1016/j.palaeo.2006.03.016.
- Gishlick, A.D., Gauthier, J.A., 2007. On the manual morphology of *Compsognathus longipes* and its bearing on the diagnosis of Compsognathidae. *Zoological Journal of the Linnean Society* 149, 569–581.
- Goloboff, P. A., Farris, J. S., Nixon, K. C., 2008. TNT, a free program for phylogenetic analysis. *Cladistics* 24, 774–786.
- Gradstein, F.M., Ogg, J.G., Schmitz, M.D., Ogg, G.M. (Eds.), 2012. The Geologic Time Scale 2012. Elsevier, Amsterdam, 1144 pp.
- Grigorescu, D., 1992. Nonmarine Cretaceous Formations of Romania. In: Matter, N.J., Chen, P.-J. (Eds.), *Aspects of Nonmarine Cretaceous Geology*. China Ocean Press, Beijing, pp. 142–164.
- Grigorescu, D., 2003. Dinosaurs of Romania. *Comptes rendus Palevol* 2, 97–101.

- Grigorescu, D., 2010. The Latest Cretaceous fauna with dinosaurs and mammals from the
Hațeg Basin — A historical overview. *Palaeogeography, Palaeoclimatology, Palaeoecology*
293, 271–282.
- Hammer, O., Harper, D.A.T., Ryan, P.D., 2001. Paleontological statistics software package
for education and data analysis. *Palaeontologia Electronica* 4, 1–9.
- Harris, J.D., 1998. A reanalysis of *Acrocanthosaurus atokensis*, its phylogenetic status, and
paleobiogeographic implications, based on a new specimen from Texas. *New Mexico*
Museum of Natural History and Science Bulletin 13, 1–75.
- Hendrickx, C., Mateus, O., 2014. Abelisauridae (Dinosauria: Theropoda) from the Late
Jurassic of Portugal and dentition-based phylogeny as a contribution for the identification of
isolated theropod teeth. *Zootaxa* 3751(1), 1–74.
- Hendrickx, C., Mateus, O., Araújo, R., 2015a. The dentition of megalosaurid theropods. *Acta*
Palaeontologica Polonica 60, 627–642. doi:10.4202/app.00056.2013.
- Hendrickx, C., Mateus, O., Araújo, R., 2015b. A proposed terminology of theropod teeth
(Dinosauria, Saurischia). *Journal of Vertebrate Paleontology* 35(5), e982797. doi:
10.1080/02724634.2015.982797.
- Holtz, T.R., Jr., Molnar, R.E., Currie, P.J., 2004a. Basal Tetanurae. In: Weishampel, D.B.,
Dodson, P., Osmólska, H. (Eds.), *The Dinosauria*. Second Edition. University of California
Press, Berkeley, Los Angeles, London, pp. 71–110.
- Holtz, T.R., Chapman, R.E., Lamanna, M.C., 2004b. Mesozoic biogeography of Dinosauria.
In: Weishampel, D.B., Dodson, P., Osmólska, H. (Eds.), *The Dinosauria*. Second Edition.
University of California Press, Berkeley, Los Angeles, London, pp. 627–642.
- Huene, F. von, 1923. Carnivorous Saurischia in Europe since the Triassic. *Bulletin of the*
Geological Society of America 34, 449–458.
- Hippolyte, J.-C., 2002. Geodynamics of Dobrogea (Romania): new constraints on the

- 1291 evolution of the Tornquist–Teisseyre Line, the Black Sea and the Carpathians.
1
- 2 1292 Tectonophysics 357, 33–53.
3
- 4 1293 Ionesi, L., 1994. Geologia unităților de platformă și a orogenului Nord-Dobrogean. Ed.
5
6
- 7 1294 Tehnică, Bucharest, 280 pp.
8
- 9 1295 Jerzykiewicz, T., Russell, D.A., 1991. Late Mesozoic stratigraphy and vertebrates of the Gobi
10
11
- 12 1296 Basin. Cretaceous Research 12, 345–377.
13
- 14 1297 Jurcsák, T., 1982. Occurences nouvelles des Sauriens mésozoïques de Roumanie. Vertebrata
15
16
- 17 1298 Hungarica 21, 175–184.
18
- 19 1299 Jurcsák, T., Popa, E., 1979. Dinozaurieni ornithopozi din bauxitele de la Cornet (Munții
20
21
- 22 1300 Pădurea Craiului). Nymphaea 7, 37–75.
23
- 24 1301 Jurcsák, T., Popa, E., 1983. La faune de dinosauriens du Bihor (Roumanie). In: Buffetaut,
25
26
- 27 1302 E., Mazin, J.M., Salmon, E. (Eds.), Actes du Symposium Paléontologique Georges Cuvier.
28
- 29 1303 Le Serpentaire, Montbéliard, pp. 325–335.
30
- 31 1304 Kellner, A.W.A., Pinheiro, A.E.P., Campos, D.A., 2014. A new sebecid from the Paleogene
32
33
- 34 1305 of Brazil and the crocodyliform radiation after the K–Pg boundary. PLoS ONE 9(1), e81386.
35
- 36 1306 doi:10.1371/journal.pone.0081386.
37
- 38 1307 Kirkland, J.I., Cifelli, R.L., Britt, B.B., Burge, D.L., DeCourten, F.L., Eaton, J.G., Parrish,
39
40
- 41 1308 J.M., 1999. Distribution of vertebrate faunas in the Cedar Mountain Formation, east-central
42
43
- 44 1309 Utah. Utah Geological Survey Miscellaneous Publication 99-1, 201–217.
45
- 46 1310 Knoll, F., Ruiz-Omeñaca, J.I., 2009. Theropod teeth from the basalmost Cretaceous of
47
48
- 49 1311 Anoual (Morocco) and their palaeobiogeographical significance. Geological Magazine 146,
50
51
- 52 1312 602–616.
53
- 54 1313 Kusuhashi, N., Matsumoto, A., Murakami, M., Tagami, T., Hirata, T., Iizuka, T., Handa, T.,
55
56
- 57 1314 Matsuoka, H., 2006. Zircon U–Pb ages from tuff beds of the upper Mesozoic Tetori Group in
58
59
- 1315 the Shokawa district, Gifu Prefecture, central Japan. The Island Arc 15, 378–390.
60
61
62
63
64
65

- Lacasa Ruiz, A., 1989. Nuevo genero de ave fosil del yacimiento Neocomiense del Montsec (Provincia de Lerida, España). Estudios geológicos 45(5-6), 417–425.
- Larson, D.W., Currie, P.J., 2013. Multivariate analyses of small theropod dinosaur teeth and implications for paleoecological turnover through time. PLoS ONE 8(1), e54329. doi:10.1371/journal.pone.0054329.
- Le Loeuff, J., Buffetaut, E., 1995. The evolution of Late Cretaceous non-marine vertebrate faunas in Europe. In: Sun, A.-L., Wang, Y.-Q. (Eds.), Sixth Symposium on Mesozoic Terrestrial Ecosystems and Biota, Short Papers. China Ocean Press, Beijing, pp. 181–184.
- Le Loeuff, J., Lang, E., Cavin, L., Buffetaut, E., 2012. Between Tendaguru and Bahariya: on the age of the Early Cretaceous dinosaur sites from the Continental Intercalaire and other African formations. Journal of Stratigraphy 36, 486–502.
- Lucas, F.A., 1901. A new dinosaur, *Stegosaurus marshi*, from the Lower Cretaceous of South Dakota. Proceedings of the United States National Museum 23(1224), 591–592.
- Lucas, S.G., 2006. The *Psittacosaurus* biochron, Early Cretaceous of Asia. Cretaceous Research 27, 189–198.
- Lucas, S.G., Estep, J.W., 1998. Vertebrate biostratigraphy and biochronology of the Cretaceous of China. In: Lucas, S.G., Kirkland, J.I., Estep, J.W. (Eds.), Lower and Middle Cretaceous Terrestrial Ecosystems. New Mexico Museum of Natural History and Science Bulletin 14, pp. 1–20.
- Lü, J.-C., Xu, L., Pu, H.-Y., Jia, S.-H., Azuma, Y., Chang, H.-L., Zhang, J.-M., 2016. Paleogeographical significance of carcharodontosaurid teeth from the late Early Cretaceous of Ruyang, Henan Province of central China. Historical Biology, 28, 8–13. doi: 10.1080/08912963.2014.947287.
- Manabe, M., Hasegawa, Y., 1995. Diapsid fauna and its paleobiogeographical implication, the Neocomian section of the Tetori Group. In: Sun, A., Wang, Y. (Eds.), Sixth Symposium

- on Mesozoic Terrestrial Ecosystems and Biota, Short Papers. China Ocean Press, Beijing, pp. 179.
- Mannion, P.D., 2009. A rebbachisaurid sauropod from the Lower Cretaceous of the Isle of Wight, England. *Cretaceous Research* 30, 521–526.
- Mannion, P.D., Upchurch, P., Hutt, S., 2011. New rebbachisaurid (Dinosauria: Sauropoda) material from the Wessex Formation (Barremian, Early Cretaceous), Isle of Wight, United Kingdom. *Cretaceous Research* 32, 774–780.
- Matsukawa, M., Ito, M., Nishida, N., Koarai, K., Lockley, M.G., Nichols, D.J., 2006. The Cretaceous Tetori biota in Japan and its evolutionary significance for terrestrial ecosystems in Asia. *Cretaceous Research* 27, 199–225.
- McDonald, A.T., 2011. The taxonomy of species assigned to *Camptosaurus* (Dinosauria: Ornithopoda). *Zootaxa* 2783, 52–68.
- McKenna, M.C., 1973. Sweepstakes, filters, corridors, Noah's Arks, and Beached Viking Funeral Ships in palaeogeography. In: Tarling, D.H., Runcorn, S.K. (Eds.), *Implications of Continental Drift to the Earth Sciences*. Academic Press, New York, pp. 295–308.
- Mo, J.-Y., Huang, C.-L., Xie, S.-W., Buffetaut, E., 2014. A megatheropod tooth from the Early Cretaceous of Fusui, Guangxi, Southern China. *Acta Geologica Sinica* (English Edition) 88, 6–12.
- Molnar, R.E., 1990. Problematic Theropoda: "Carnosaurs". In: Weishampel, D.B., Dodson, P., Osmólska, H. (Eds.), *The Dinosauria*. University of California Press, Berkeley, Los Angeles, Oxford, pp. 306–317.
- Mutihac, V., Mutihac, G., 2010. The geology of Romania, within the Central East European geostructural context. Ed. Didactică și Pedagogică, Bucharest, 690 pp.
- Neagu, T., Dragastan, O., Csiki, Z., 1997. Early Cretaceous shelf paleocommunities of Cernavodă (South Dobrogea, SE Romania). *Acta Palaeontologica Romaniae* 1, 28–36.

- 1366 Nopcsa, F., 1902. Notizen über cretacische Dinosaurier. Sitzungsberichte der Kaiserlichen
 1 Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe 111, 93–114.
 2 1367
 3
 4 1368 Nopcsa, F., 1923. On the geological importance of the primitive reptilian fauna of the
 5
 6
 7 1369 uppermost Cretaceous of Hungary; with a description of a new tortoise (*Kallokibotium*).
 8
 9
 10 1370 Quarterly Journal of the Geological Society of London 79, 100–116.
 11
 12 1371 Norman, D.B., 2010. A taxonomy of iguanodontians (Dinosauria: Ornithopoda) from the
 13
 14 1372 lower Wealden Group (Cretaceous: Valanginian) of southern England. Zootaxa 2489, 47–66.
 15
 16
 17 1373 Norman, D.B., 2013. On the taxonomy and diversity of Wealden iguanodontian dinosaurs
 18
 19 1374 (Ornithischia: Ornithopoda). Revue de Paléobiologie 32, 385–404.
 20
 21
 22 1375 Novas, F.E., de Valais, S., Vickers-Rich, P.A., Rich, T.H., 2005. A large Cretaceous theropod
 23
 24 1376 from Patagonia, Argentina, and the evolution of carcharodontosaurids. Naturwissenschaften
 25
 26 1377 92, 226–230.
 27
 28
 29 1378 Novas, F.E., Agnolín, F.L., Ezcurra, M.D., Porfiri, J., Canale, J.I., 2013. Evolution of the
 30
 31 1379 carnivorous dinosaurs during the Cretaceous: The evidence from Patagonia. Cretaceous
 32
 33
 34 1380 Research 45, 174–215.
 35
 36 1381 Ortega, F., Escaso, F., Sanz, J.L., 2010. A bizarre, humped Carcharodontosauria (Theropoda)
 37
 38
 39 1382 from the Lower Cretaceous of Spain. Nature 467, 203–206.
 40
 41 1383 Ósi, A., Rabi, M., Makádi, L., Szentesi, Z., Botfalvai, G., Gulyás, P., 2012. The Late
 42
 43 1384 Cretaceous continental vertebrate fauna from Iharkút (western Hungary): a review. In:
 44
 45
 46 1385 Godefroit, P. (Ed.), Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems.
 47
 48
 49 1386 Indiana University Press, Bloomington, pp. 533–569.
 50
 51 1387 Ósi, A., Codrea, V., Prondvai, E., Csiki-Sava, Z., 2014. New ankylosaurian material from the
 52
 53 1388 Upper Cretaceous of Transylvania. Annales de Paléontologie 100, 257–271. doi:
 54
 55
 56 1389 10.1016/j.annpal.2014.02.001.
 57
 58
 59
 60
 61
 62
 63
 64
 65

- 1390 Ósi, A., Rabi, M., Makádi, L., 2015. An enigmatic crocodyliform tooth from the bauxites of
1
2 1391 western Hungary suggests hidden mesoeucrocodylian diversity in the Early Cretaceous
3
4
5 1392 European archipelago. *PeerJ* 3, e1160. doi:10.7717/peerj.1160.
6
7 1393 Paolillo, A., Linares, O.J., 2007. Nuevos cocodrilos Sebecosuchia del Cenozoico
8
9
10 1394 Suramericano (Mesosuchia: Crocodylia). *Paleobiologia Neotropical* 3, 1–25.
11
12 1395 Pereda-Suberbiola, X., Ruiz-Omeñaca, J.I., Fernandez-Baldor, F.T., Maisch, M.W., Huerta,
13
14 1396 P., Contreras, R., Izquierdo, L.A., Huerta, D.M., Montero, V.U., Welle, J., 2011. A tall-
15
16
17 1397 spined ornithopod dinosaur from the Early Cretaceous of Salas de los Infantes (Burgos,
18
19 1398 Spain). *Comptes Rendus Palevol* 10, 551–558.
20
21
22 1399 Pereda-Suberbiola, X., Ruiz-Omeñaca, J.I., Canudo, J.I., Torcida, F., Sanz, J.L., 2012.
23
24 1400 Dinosaur faunas from the Early Cretaceous (Valanginian–Albian) of Spain. In: Godefroit, P.
25
26 1401 (Ed.), *Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems*. Indiana University
27
28
29 1402 Press, Bloomington, pp. 378–407.
30
31
32 1403 Pérez-Moreno, B.P., Sanz, J.L., Sudre, J., Sigé, B., 1993. A theropod dinosaur from the
33
34 1404 Lower Cretaceous of southern France. *Revue de Paléobiologie Volume spéciale* 7, 173–188.
35
36 1405 Pol, D., Powell, J.E., 2011. A new sebecid mesoeucrocodylian from the Rio Loro Formation
37
38
39 1406 (Palaeocene) of north-western Argentina. *Zoological Journal of the Linnean Society* 163, S7–
40
41 1407 S36.
42
43
44 1408 Porfiri, J.D., Novas, F.E., Calvo, J.O., Agnolín, F.L., Ezcurra, M.D., Cerda, I.A., 2014.
45
46 1409 Juvenile specimen of *Megaraptor* (Dinosauria, Theropoda) sheds light about tyrannosauroid
47
48
49 1410 radiation. *Cretaceous Research* 51, 35–55. doi:10.1016/j.cretres.2014.04.007.
50
51 1411 Posmoşanu, E., 2003. Iguanodontian dinosaurs from the Lower Cretaceous bauxite site from
52
53 1412 Romania. *Acta Palaeontologica Romaniae* 4, 431–439.
54
55
56 1413 Rabi, M., Sebők, N., 2015. A revised Eurogondwana model: Late Cretaceous notosuchian
57
58 1414 crocodyliforms and other vertebrate taxa suggest the retention of episodic faunal links
59
60
61
62
63
64
65

between Europe and Gondwana during most of the Cretaceous. *Gondwana Research* 28,
1197–1211. doi:10.1016/j.gr.2014.09.015.

Racey, A., 2009. Mesozoic red bed sequences from SE Asia and the significance of the
Khorat Group of NE Thailand. In: Buffetaut, E., Cuny, G., Le Loeuff, J., Suteethorn, V.
(Eds.), *Late Palaeozoic and Mesozoic Ecosystems in SE Asia*. Geological Society, London,
Special Publications 315, pp. 41–67. doi: 10.1144/SP315.5.

Racey, A., Goodall, J.G.S., 2009. Palynology and stratigraphy of the Mesozoic Khorat Group
red bed sequences from Thailand. In: Buffetaut, E., Cuny, G., Le Loeuff, J., Suteethorn, V.
(Eds.), *Late Palaeozoic and Mesozoic Ecosystems in SE Asia*. Geological Society, London,
Special Publications 315, pp. 69–83. doi: 10.1144/SP315.6.

Rauhut, O.W.M., 2011. Theropod dinosaurs from the Late Jurassic of Tendaguru (Tanzania).
Special Papers in Palaeontology 86, 195–239.

Rauhut, O.W.M., Werner, C., 1995. First record of the family Dromaeosauridae (Dinosauria:
Theropoda) in the Cretaceous of Gondwana (Wadi Milk Formation, northern Sudan.
Paläontologische Zeitschrift 69, 475–489.

Richter, U., Mudroch, A., Buckley, L.G., 2013. Isolated theropod teeth from the Kem Kem
Beds (Early Cenomanian) near Taouz, Morocco. *Paläontologische Zeitschrift* 87, 291–309.

Royo-Torres, R., Cobos, A., Luque, L., Aberasturi, A., Espilez, E., Fierro, I., Gonzales, A.,
Mampel, L., Alcalá, L., 2009. High European sauropod dinosaur diversity during Jurassic–
Cretaceous transition in Riodeva (Teruel, Spain). *Palaeontology* 52, 1009–1027.

Sachs, S., Hornung, J.J., 2013. Ankylosaur remains from the Early Cretaceous (Valanginian)
of Northwestern Germany. *PLoS ONE* 8(4), e60571. doi:10.1371/journal.pone.0060571.

Salgado, L., Canudo, J.I., Garrido, A.C., Ruiz-Omeñaca, J.I., Garcia, R.A., de la Fuente,
M.S., Barco, J.L., Bollati, R., 2009. Upper Cretaceous vertebrates from El Anfiteatro area,
Río Negro, Patagonia, Argentina. *Cretaceous Research* 30, 767–784.

- 1440 Sames, B., Cifelli, R.L., Schudack, M.E., 2010. The nonmarine Lower Cretaceous of the
1
2 1441 North American Western Interior foreland basin: New biostratigraphic results from ostracod
3
4 1442 correlations and early mammals, and their implications for paleontology and geology of the
5
6
7 1443 basin—An overview. *Earth-Science Reviews* 101, 207–224.
8
9
10 1444 Sauvage, H.E., 1876. Notes sur les reptiles fossiles. *Bulletin de la Société Géologique de*
11
12 1445 *France* 4, 435–442.
13
14 1446 Sauvage, H.E., 1882. Recherches sur les reptiles trouves dans le Gault de l'est du bassin de
15
16
17 1447 Paris. *Mémoires de la Société Géologique de France* 2, 1–42.
18
19 1448 Săndulescu, M., 1984. *Geotectonica României*. Ed. Tehnică, Bucharest, 329 pp.
20
21
22 1449 Seghedi, A., 2001. The North Dobrogea orogenic belt (Romania): a review. In: Ziegler, P.A.,
23
24 1450 Cavazza, W., Robertson, A.H.F., Crasquin-Soleau, S. (Eds.), *Peri-Tethys Memoir 6: Peri-*
25
26 1451 *Tethyan Rift/Wrench Basins and Passive Margins. Mémoires de la Musée National d'Histoire*
27
28
29 1452 *Naturelle, Paris*, pp. 237–257.
30
31
32 1453 Sereno, P.C., 1999. Dinosaurian biogeography: vicariance, dispersal and regional extinction.
33
34 1454 In: Tomida, Y., Rich, T.H., Vickers-Rich, P. (Eds.), *Proceedings of the Second Gondwanan*
35
36 1455 *Dinosaur Symposium. National Science Museum Monographs* 15, Tokyo, pp. 249–257.
37
38
39 1456 Sereno, P.C., Brusatte, S.L., 2008. Basal abelisaurid and carcharodontosaurid theropods from
40
41 1457 the Lower Cretaceous Elrhaz Formation of Niger. *Acta Palaeontologica Polonica* 53, 15–46.
42
43
44 1458 Sereno, P.C., Dutheil, D.B., Iarochene, M., Larsson, H.C.E., Lyon, G.H., Magwene, P.M.,
45
46 1459 Sidor, C.A., Varricchio, D.J., Wilson, J.A., 1996. Predatory dinosaurs from the Sahara and
47
48
49 1460 Late Cretaceous faunal differentiation. *Science* 272, 986–991.
50
51 1461 Shen, Y.B., Mateer, N.J., 1992. An outline of the Cretaceous system in northern Xinjiang,
52
53 1462 western China. In: Mateer, N.J., Chen, P.J. (Eds.), *Aspects of Nonmarine Cretaceous*
54
55
56 1463 *Geology. China Ocean Press, Beijing*, pp. 49–77.
57
58
59
60
61
62
63
64
65

- 1464 Simionescu, I., 1906. Note sur l'age des calcaires de Cernavoda (Dobrogea). Annales
1
2 1465 Scientifiques de l'Université de Jassy 4(1), 1–3.
3
4
5 1466 Simionescu, I., 1913. *Megalosaurus* aus der Unterkreide der Dobrogea. Centralblatt für
6
7 1467 Mineralogie, Geologie und Paläontologie 1913(20), 686–687.
8
9
10 1468 Sissingh, W., 1977. Biostratigraphy of Cretaceous calcareous nannoplankton. Geologie en
11
12 1469 Mijnbouw 56, 37–65.
13
14
15 1470 Smith, J.B., Vann, D.R., Dodson, P., 2005. Dental morphology and variation in theropod
16
17 1471 dinosaurs: implications for the taxonomic identification of isolated teeth. The Anatomical
18
19 1472 Record A 285A, 699–736.
20
21
22 1473 Stein, K., Csiki, Z., Curry Rogers, K., Weishampel, D.B., Redelstorff, R., Carballido, J.L.,
23
24 1474 Sander, P.M., 2010. Small body size and extreme cortical bone remodeling indicate phyletic
25
26 1475 dwarfism in *Magyarosaurus dacus* (Sauropoda: Titanosauria). Proceedings of the National
27
28 1476 Academy of Sciences 107, 9258–9263.
29
30
31 1477 Stoica, M., Csiki, Z., 2002. An earliest Cretaceous (Purbeckian) vertebrate fauna from
32
33 1478 Southern Dobrogea (southeastern Romania). In: Grigorescu, D., Csiki, Z. (Eds.), 7th European
34
35 1479 Workshop on Vertebrate Palaeontology, Sibiu, Romania. Ars Docendi, Bucharest, pp. 34.
36
37
38 1480 Stovall, J.W., Langston, W., Jr., 1950. *Acrocanthosaurus atokensis*, a new genus and species
39
40 1481 of Lower Cretaceous Theropoda from Oklahoma. American Midland Naturalist 43, 686–728.
41
42
43 1482 Stromer, E., 1931. Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wüsten
44
45 1483 Ägyptens. II. Wirbeltier-Reste der Baharijstufe (unterstes Cenoman). 10. Ein Skelett-Rest
46
47 1484 von *Carcharodontosaurus* nov. gen. Abhandlungen der Bayerischen Akademie der
48
49 1485 Wissenschaften, Mathematisch-naturwissenschaftliche Abteilung Neue Folge 9, 1–23.
50
51
52 1486 Therrien, F., Henderson, D.M., 2007. My theropod is bigger than yours ... or not: estimating
53
54 1487 body size from skull length in theropods. Journal of Vertebrate Paleontology 27, 108–115.
55
56
57
58
59
60
61
62
63
64
65

- 1488 Tong, H., Claude, J., Suteethorn, V., Naksri, W., Buffetaut, E., 2009. Turtle assemblages of
1
2 1489 the Khorat Group (Late Jurassic–Early Cretaceous) of NE Thailand and their
3
4
5 1490 palaeobiogeographical significance. In: Buffetaut, E., Cuny, G., Le Loeuff, J., Suteethorn, V.
6
7 1491 (Eds.), Late Palaeozoic and Mesozoic Ecosystems in SE Asia. The Geological Society,
8
9
10 1492 London, Special Publications 315, pp. 141–151.
- 11
12 1493 Torcida Fernández-Baldor, F., Canudo, J.I., Huerta, P., Montero, D., Pereda Suberbiola, X.,
13
14
15 1494 Salgado, L., 2011. *Demandasaurus darwini*, a new rebbachisaurid sauropod from the Early
16
17 1495 Cretaceous of the Iberian Peninsula. Acta Palaeontologica Polonica 56, 535–552.
- 18
19 1496 Turculeț, I., Brânzilă, M., 2012. Muzeul colecțiilor paleontologice originale de la
20
21
22 1497 Universitatea "Alexandru Ioan Cuza" Iași. Editura Universității "Alexandru Ioan Cuza" Iași,
23
24 1498 Iași, 173 pp.
- 25
26
27 1499 Turner, A.H., Sertich, J.J.W., 2010. Phylogenetic history of *Simosuchus clarki*
28
29 1500 (Crocodyliformes: Notosuchia) from the Late Cretaceous of Madagascar. Journal of
30
31
32 1501 Vertebrate Paleontology 30(Supplement 1), 177–236.
- 33
34 1502 Turner, A.H., Makovicky, P.J., Norell, M.A., 2012. A review of dromaeosaurid systematics
35
36 1503 and paravian phylogeny. Bulletin of the American Museum of Natural History 371, 1–206.
- 37
38
39 1504 Vasile, Ș., Csiki, Z., 2011. New Maastrichtian microvertebrates from the Rusca Montană
40
41 1505 Basin (Romania). Oltenia. Studii și comunicări. Științele Naturii 27(1), 221–230.
- 42
43
44 1506 Vickers-Rich, P., Rich, T.H., Lanus, D.R., Rich, L.S.V., Vacca, R., 1999. "Big Tooth" from
45
46 1507 the Early Cretaceous of Chubut Province, Patagonia: a possible carcharodontosaurid. In:
47
48
49 1508 Tomida, Y., Rich, T.H., Vickers-Rich, P. (Eds.), Proceedings of the Second Gondwanan
50
51 1509 Dinosaur Symposium. National Science Museum Monographs 15, Tokyo, pp. 85–88.
- 52
53
54 1510 Vremir, M., 2010. New faunal elements from the Late Cretaceous (Maastrichtian) continental
55
56 1511 deposits of Sebeș area (Transylvania). Terra Sebus. Acta Musei Sabesiensis 2, 635–684.

- Vullo, R., Néraudeau, D., Lenglet, T., 2007. Dinosaur teeth from the Cenomanian of Charentes, western France: evidence for a mixed Laurasian-Gondwanan assemblage. *Journal of Vertebrate Paleontology* 27, 931–943.
- Weishampel, D.B., 1990. Dinosaurian distribution. In: Weishampel, D.B., Dodson, P., Osmólska, H. (Eds.), *The Dinosauria*. California University Press, Berkeley, Los Angeles, Oxford, pp. 63–140.
- Weishampel, D.B., Grigorescu, D., Norman, D.B., 1991. The Dinosaurs of Transylvania. *National Geographic Research & Exploration* 7(2), 196–215.
- Weishampel, D.B., Norman, D.B., Grigorescu, D., 1993. *Telmatosaurus transsylvanicus* from the Late Cretaceous of Romania: the most basal hadrosaurid dinosaur. *Palaeontology* 36, 361–385.
- Weishampel, D.B., Jianu, C.M., Csiki, Z., Norman, D.B., 2003. Osteology and phylogeny of *Zalmoxes* (n. g.), an unusual euornithopod dinosaur from the latest Cretaceous of Romania. *Journal of Systematic Palaeontology* 1, 65–123.
- Weishampel, D.B., Barrett, P.M., Coria, R.A., Le Loeuff, J., Xu, X., Zhao, X-J., Sahni, A., Gomani, E.M., Noto, C.R., 2004. Dinosaur distribution. In: Weishampel, D.B., Dodson, P., Osmólska, H. (Eds.), *The Dinosauria*. Second Edition. University of California Press, Berkeley, Los Angeles, London, pp. 517–606.
- Weishampel, D.B., Csiki, Z., Benton, M.J., Grigorescu, D., Codrea, V., 2010. Palaeobiogeographic relationships of the Hațeg biota — Between isolation and innovation. *Palaeogeography, Palaeoclimatology, Palaeoecology* 293, 419–437.
- Williamson, T.E., Brusatte, S.L., 2014. Small theropod teeth from the Late Cretaceous of the San Juan Basin, northwestern New Mexico and their implications for understanding latest Cretaceous dinosaur evolution. *PLoS ONE* 9(4), e93190. doi:10.1371/journal.pone.0093190.

Zarcone, G., Cillari, F.M.P., Stefano, P.D., Guzzetta, D., Nicosia, U., 2010. A possible bridge between Adria and Africa: New palaeobiogeographic and stratigraphic constraints on the Mesozoic palaeogeography of the Central Mediterranean area. *Earth-Science Reviews* 103, 154–162.

Figure captions

Figure 1. Simplified geological map of the Cernavodă-Cochirleni area; inset shows the position of the study area within Romania. Legend: 1. Quaternary: a. Holocene alluvia, b. Pleistocene–Holocene loessoid deposits; 2. Pre-Quaternary Cenozoic (Middle Eocene and Miocene) deposits; Cretaceous: 3. Peștera Formation, Lower Cenomanian; 4. Cochirleni Formation; uppermost Aptian–Lower Albian; 5. Gherghina Formation, Middle–Upper Aptian; 6. Ostrov (= Ramadan) Formation; Barremian–Lower Aptian; 7. Cernavodă Formation, Alimanu Member, Berriasian–Valanginian; 8. Water courses. (Redrawn after Dragastan et al., 1998, 2014).

Figure 2. Specimen UAIC (SCM1) 615, indeterminate carcharodontosaurid lateral tooth from Cochirleni, Southern Dobrogea. A. UAIC (SCM1) 615, as figured by Simionescu (1913); B. Current state of UAIC (SCM1) 615, mounted in a limestone holder.

Figure 3. Detailed morphology of UAIC (SCM1) 615, an indeterminate carcharodontosaurid lateral tooth from Cochirleni, Southern Dobrogea. UAIC (SCM1) 615 in A. labial? side; B., distal; C., lingual? side, and D., basal (mesial to the right) views. Details of the distal carina (marked with boxes in A, respectively C): apical part in E., labial? and F. distal views; basal part in G., lingual? and H., distal views. Scale bar: 1 cm (A–D), 5 mm (E–H).

Figure 4. Dental morphospace of the different theropod clades according to the results of the PCA analysis; UAIC (SCM1) 615 (red star) plots within the morphospace occupied by Carcharodontosauridae. See further details of this analysis, as well as other quantitative analyses used to identify the tooth that deliver similar results (cluster analysis, discriminant function analysis, phylogenetic analysis), in the Supplementary Material.

Figure 5. A. Palaeogeographic setting of the two early Early Cretaceous Romanian dinosaur occurrences: the Berriasian–Valanginian Cornet locality (orange star), located on a Neo-Tethyan archipelago island, and the Valanginian Cochirleni locality (red star), situated on the marginal areas of the Eastern European cratonic mainland. B. Global chronostratigraphic and palaeobiogeographic distribution of the Carcharodontosauridae, plotted on Middle Aptian (approx. 120 Mya) palaeogeographic map; red star marks the position of UAIC (SCM1) 615 from Southern Dobrogea. Legend: 1 – *Veterupristisaurus*, ‘*Megalosaurus*’ *ingens*, Carcharodontosauridae indet., Tanzania, Late Jurassic; 2 – *Concavenator*, Spain, Barremian; 3 – Carcharodontosauridae indet., Thailand, Barremian; 4 – *Acrocanthosaurus*, southeastern United States, Aptian–Albian; 5 – Carcharodontosauridae indet., Spain, Aptian; 6 – *Eocarcharia*, Niger, Aptian–Albian; 7 – Carcharodontosauridae indet., Guangxi, China, Aptian; 8 – Carcharodontosauridae indet., Henan, China, Aptian; 9 – *Kelmayisaurus*, Xinjiang, China, Aptian–Albian; 10 – Carcharodontosauridae indet., France, Cenomanian; 11 – *Sauroniops*, Morocco, Cenomanian; 12 – Carcharodontosauridae indet., Japan, Cenomanian–early Turonian; 13 – *Shaochilong*, Inner Mongolia, China, Turonian; 14 – Carcharodontosauridae indet., São Paulo, Brazil, Campanian–Maastrichtian (for relevant references, see text, 5.4.). Palaeogeographic maps, courtesy of Ron Blakey (<http://cpgeosystems.com/>).

1
2 1 “*Megalosaurus cf. superbus*” from southeastern Romania: the oldest known Cretaceous
3
4 2 carcharodontosaurid (Dinosauria: Theropoda) and its implications for earliest Cretaceous
5
6 3 Europe-Gondwana connections
7
8 4
9
10 5
11
12 6 Zoltán Csiki-Sava^{1*}, Stephen L. Brusatte², Ștefan Vasile¹
13
14 7 ¹ Department of Geology, Faculty of Geology and Geophysics, University of Bucharest, 1
15
16 8 Nicolae Bălcescu Boulevard, 010041 Bucharest, Romania, ~~zoltan.csiki@g.unibuc.ro,~~
17
18 9 ~~yokozuna_uz@yahoo.com~~
19
20
21 10 ² School of GeoSciences, University of Edinburgh, Grant Institute, James Hutton Road,
22
23 11 Edinburgh, EH9 3FE, United Kingdom, ~~Stephen.Brusatte@ed.ac.uk~~
24
25 12 * Corresponding author
26
27 13 zoltan.csiki@g.unibuc.ro
28
29 14
30
31 15 ABSTRACT
32
33 16 Some of the best records of continental vertebrates from the Cretaceous of Europe come from
34
35 17 Romania, particularly two well-known occurrences of dwarfed and morphologically aberrant
36
37 18 dinosaurs and other taxa that lived on islands (the Cornet and Hațeg Island faunas).
38
39 19 Substantially less is known about those vertebrates living in the more stable, cratonic regions
40
41 20 of Romania (and Eastern Europe as a whole), particularly during the earliest Cretaceous. We
42
43 21 describe one of the few early Early Cretaceous fossils that have ever been found from these
44
45 22 regions, the tooth of a large theropod dinosaur from Southern Dobrogea, which was
46
47 23 discovered over a century ago but whose age and identification have been controversial. We
48
49 24 identify the specimen as coming from the Valanginian stage of the Early Cretaceous, an
50
51 25 incredibly poorly sampled interval in global dinosaur evolution, and as belonging to
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Carcharodontosauridae, a clade of derived, large-bodied apex predators whose earliest Cretaceous history is poorly known. Quantitative analyses demonstrate that the Romanian tooth shows affinities with a derived carcharodontosaurid subgroup, the Carcharodontosaurinae, which until now has been known solely from Gondwana. Our results suggest that this subgroup of colossal predators did not evolved vicariantly as Laurasia split from Gondwana, but originated earlier, perhaps in Europe. The carcharodontosaurine diversification may have been tied to a north-to-south trans-Tethyan dispersal that took place sometime between the Valanginian and Aptian, illustrating the importance of palaeogeographic ties between these two realms during the largely mysterious early–mid Early Cretaceous.

Keywords

~~Romania~~ Southern Dobrogea; ~~Lower Cretaceous~~ Valanginian; ~~Theropoda~~; Carcharodontosauridae; ~~cratonic Europe~~; palaeobiogeography

1. Introduction

Romania boasts one of the best records of continental vertebrate fossils from the Cretaceous of Europe (e.g., Grigorescu, 1992, 2003; Csiki-Sava et al., 2015). The vast majority of fossils come from two well-known occurrences: the Early Cretaceous bauxite accumulations of Cornet, in the northern Apuseni Mountains (e.g., Jurcsák, 1982; Benton et al., 1997; Posmoşanu, 2003; Dyke et al., 2011), and the famous latest Cretaceous beds of the Haţeg, Rusca Montană and western Transylvanian basins of Transylvania, which have yielded the dinosaur-dominated ‘Haţeg Island fauna’ (e.g., Nopcsa, 1923; Weishampel et al., 1991; Benton et al. 2010; Codrea et al., 2010, 2012; Grigorescu, 2010; Vremir, 2010; Vasile and Csiki, 2011; Csiki-Sava et al., 2015). Both of these faunas inhabited islands that were part of

the vast Cretaceous European Archipelago of the Neo-Tethys Ocean. Based on their isolated geological settings and the many dwarfed and morphologically aberrant taxa that make up the faunas, both have been interpreted as insular assemblages that give a unique window into how island environments affected the evolution of long-extinct organisms (e.g., Benton et al., 1997, 2010; Csiki-Sava et al., 2015).

The great volume of research on these assemblages over the past century, particularly the ‘Hațeg Island fauna’, has concealed an inconvenient bias: the stable, non-island, cratonic regions of Romania have yielded only extremely rare Mesozoic continental vertebrate remains (i.e., the Moldavian, Moesian and Scythian platforms; Săndulescu, 1984; Mutihac and Mutihac, 2010; Fig. 1). This is mostly because Mesozoic deposits are located in the subsurface in these regions, with only limited subaerial exposures available in the structurally highest-lying parts of the Moesian Platform, in Central and Southern Dobrogea (Middle Jurassic–Upper Cretaceous), as well as in the northeastern-most corner of the Moldavian Platform, along the Prut Valley (lower Upper Cretaceous) (see, e.g., Mutihac and Mutihac, 2010). This bias is unfortunate because fossils from these settings could lead to a better understanding of how mainland and island faunas differed during the Cretaceous, and because the cratonic portion of Europe was an important biogeographic stepping stone between the north and south as the continents fragmented and sea levels fluctuated.

Although the cratonic regions of Romania have yielded few Cretaceous terrestrial fossils, these deposits are not totally barren. In fact, one of the first Mesozoic continental vertebrates ever recorded from Romania comes from one of these deposits, the Lower Cretaceous shallow marine limestones of Southern Dobrogea (Fig. 1). This specimen—the isolated but well-preserved tooth of a large theropod dinosaur—has often been overlooked. It was described a little over a century ago by Simionescu (1913; Fig. 2A), and until a few recent discoveries of very rare isolated specimens (Stoica and Csiki, 2002; Csiki-Sava et al.,

2013,~~in prep.~~; Dragastan et al., 2014), it remained as the sole published record of Mesozoic terrestrial vertebrates from the cratonic areas of Romania. It has never been comprehensively described and its precise age and taxonomic affinities have yet to be clarified, despite its potential importance as a well-preserved fossil from a poorly sampled area that could have critical evolutionary and biogeographic implications.

We here present a comprehensive description of the Dobrogea tooth and discuss its relevance for understanding dinosaur evolution and biogeography. We review the peculiar history of how this specimen was collected and curated, thoroughly document its morphology and age, identify it based on comparison to a broad range of theropods, and outline its importance. It turns out that this specimen, although only a single tooth, has wide-ranging implications. We identify it as coming from the Valanginian stage of the Early Cretaceous, which is incredibly poorly sampled both in Europe and globally (Weishampel et al., 2004), and as belonging to a carcharodontosaurid, a group of derived, large-bodied apex predators whose earliest Cretaceous history is poorly known. Carcharodontosaurids were once thought to be a uniquely Gondwanan group, but recent discoveries show that the basal members of the group were more widespread during the late Early-middle Cretaceous (e.g., Sereno et al., 1996; Brusatte and Sereno, 2008). The Romanian tooth shows affinities with a derived carcharodontosaurid subgroup, the Carcharodontosaurinae, that until now has been known only from Gondwana. It suggests that this subgroup of enormous predators did not evolve vicariantly as Pangaea split, but originated earlier, and perhaps in Europe, suggesting faunal interchange between Europe and Gondwana during the ‘dark ages’ of the early Early Cretaceous.

Abbreviations: UAIC – University “Alexandru Ioan Cuza”, Iași, Romania.

2. History of collecting and curation

1
2 101 Only two dinosaurian fossils are currently known from the cratonic areas of Romania: an
3
4 102 isolated theropod tooth and an isolated caudal vertebral centrum. Both of these were reported
5
6 103 from the Lower Cretaceous deposits of Southern Dobrogea (southeastern Romania; Csiki-
7
8 104 Sava et al., 2013, see also below). Unfortunately, exact details of their discovery and places
9
10 105 of origin are lost, a fact that can hinder an assessment of their age and interpretation of their
11
12 106 phylogenetic and palaeobiogeographic significance. Our aim here is to gather and report all
13
14 107 available information concerning the collecting of specimen UAIC (SCM1) 615, that is, the
15
16 108 isolated theropod tooth reported by Simionescu (1913; Fig. 2A).

18
19 109 According to the existing information - unpublished museum labels and records, and
20
21 110 the preliminary publication of Simionescu (1913) - specimen UAIC (SCM1) 615 was
22
23 111 discovered in the surroundings of Cochirleni, a small village south of Cernavodă and close to
24
25 112 the right bank of the Danube, in Southern Dobrogea, southeastern Romania (Fig. 1), probably
26
27 113 shortly before 1913, the date of its publication by Simionescu (1913).

28
29 114 Although studied and preliminarily described by Simionescu, UAIC (SCM1) 615 was not
30
31 115 collected by Simionescu personally. Instead, it was donated by a certain “de Tomas” (also
32
33 116 mentioned as “de Thomas” in the registry of the Hârşova Museum) to V. Cotovu from
34
35 117 Hârşova (Central Dobrogea), a local teacher, archaeology and natural history aficionado, and
36
37 118 amateur fossil collector (see, e.g., Covacef, 1995). Cotovu, described by Simionescu himself
38
39 119 as the “zélé fondateur et directeur du muséum de Hârşova” (*enthusiastic founder and director*
40
41 120 *of the Hârşova Museum*; Simionescu, 1906: p. 2), had previously provided fossil specimens
42
43 121 from Southern Dobrogea for study to Simionescu, a nationally acknowledged popular science
44
45 122 writer and scientist, whom Cotovu knew personally (Brânzilă, 2010). These circumstances
46
47 123 are supported by the fact that in the original description, Simionescu figures the specimen as
48
49 124 being accessioned in the “Regional-Museum von Harschowa” (*Hârşova Regional Museum*;
50
51 125 Simionescu, 1913: p. 687, fig.1), a designation he also used to refer to other Dobrogean

specimens not collected by him first-hand (e.g., a specimen of '*Nautilus*' *pseudoelegans* from Cernavodă, or a fragmentary tooth-bearing palatal fragment referred to as '*Coelodus*' sp., also originating from Cochirleni; see Simionescu, 1906). Confirming this deduction, an isolated tooth appears accessioned in the old registry book of the Hârşova Museum (under specimen number 200) as "*Megalosaurus* cf. *superbus*", with the mention that it was "described by Prof. Simionescu in the *Centralblatt f. min. etc.*". This is also the case of the '*Coelodus*' sp. specimen from Cochirleni (specimen number 86), similarly clearly identified as being described by Simionescu in the registry book.

Both of these vertebrate remains from Dobrogea that were formerly part of the Hârşova Museum collections are currently accessioned in the palaeontology collections of the UAIC (Turculeţ and Brânzilă, 2012), suggesting that, at one moment, several specimens were transferred there from the Hârşova Museum. Although no details are known about this transfer, it is probable that it took place right before (or when) the Hârşova Museum, including a part of its collections, was burned and largely destroyed during WWI, in 1916, a time when Simionescu still held a position at the UAIC.

After its original description, specimen UAIC (SCM1) 615 underwent a minor amount of damage (see below, Description). Also, at some point between its description in 1913 and the early 1960s (when the specimen was found in its present state in the collections of the UAIC by academic staff members who are still alive today and recall the discovery; I. Turculeţ, personal communication, May 2013) it was glued into a limestone matrix holder, while it was obviously completely freed of the surrounding matrix when it was described and figured in 1913 (Fig. 2). The circumstances under which these alterations took place are unclear. It is a distinct possibility that they occurred sometimes during WWII, when, in the spring of 1944, the frontline between the German-Romanian and Soviet armies reached the Iaşi–Chişinău line. At this moment, the geological-palaeontological collections of the UAIC

1
2 151 were packed in crates, and moved together with its personnel and other possessions to Zlatna,
3
4 152 in the Apuseni Mountains (western Romania), to safeguard them from any potential damage.
5
6 153 Mounting the specimen into the limestone stand would have been a quick way to stabilize it,
7
8 154 as it appears that packaging and transport of the specimens was done in haste (M. Brânzilă,
9
10 155 personal communication, April 2103). If that was indeed the case, the mounting would have
11
12 156 taken place without the knowledge of Simionescu, who left Iași and the UAIC in 1929, being
13
14 157 invited to become a professor of Palaeontology at the University of Bucharest (Brânzilă,
15
16 158 2010). Then again, however, Simionescu himself or staff of the Hârșova Museum might have
17
18 159 re-mounted the tooth after its original description, or else the mounting might have taken
19
20
21 160 place after the return of the collections to Iași, after WWII.

22
23 161 Unfortunately, it is not documented whether the mounting was made using the
24
25 162 original matrix, or if a trough corresponding to the tooth outline was carved into a randomly
26
27 163 chosen limestone block. The apparently excellent fit between the tooth and the depression
28
29 164 housing it (Fig. 2B, 3) suggests that this operation was completed carefully, and accurate
30
31 165 carving of a fake holder is difficult to reconcile with the rush accompanying the evacuation of
32
33 166 the Iași University, in 1944. Alternatively, the presence of a hand-written old registration
34
35 167 number on the specimen holder would support its early re-mounting, while still at the
36
37 168 Hârșova Museum. As noted previously, the original Hârșova Museum registration number of
38
39 169 the specimen was 200, which does not correspond to that currently written both on the
40
41 170 limestone holder and on a paper sticker (204). However, according to the old collection
42
43 171 registry of the Hârșova Museum, specimen numbers 201 through 225 were given to a series
44
45 172 of “indeterminate (fossil) bone fragments” from the “Cochirleni quarries”. Thus, these
46
47 173 specimens (now apparently lost) came from the same locality as the tooth, and they were
48
49
50 174 collected and donated by the same person to the Museum who donated UAIC (SCM1) 615.
51
52 175 There is, thus, a (albeit admittedly remote) possibility that the registration numbers were
53

1
2 176 mixed up during the re-mounting of the specimen, which in this case took place at an early
3
4 177 date in the Hârşova Museum. If this is indeed the case, the limestone used as holder could
5
6 178 have been the same as the original matrix of the specimen.
7

8 179 To conclude, the history of recovery and curation of the historically important
9
10 180 dinosaurian specimen UAIC (SCM1) 615 is rather convoluted and clouded by many
11
12 181 uncertainties. The exact date of discovery remains conjectural, and the exact place of the
13
14 182 discovery (thus also the original geological context of the tooth) is even more ambiguous.
15
16 183 The current state of the specimen, and especially its mounted status, suggest a curatorial
17
18 184 history that produced a moderate amount of damage to, but also partially obscured the
19
20
21 185 detailed morphology of the specimen. The convergence of such unfortunate events makes
22
23 186 deciphering the age, identity and evolutionary significance of the specimen troublesome,
24
25 187 although many lines of evidence, carefully considered, allow us to draw reasonable
26
27 188 conclusions (see below).
28
29 189
30

31 190 3. Geological setting 32

33 191 According to the available collecting information, the isolated theropod tooth UAIC (SCM1)
34
35 192 615 was discovered at Cochirleni (sometimes noted more specifically as the “*Cochirleni*
36
37 193 *quarry*” or “*Cokerleni quarry*”). Cochirleni is a small village in southwestern Dobrogea
38
39 194 situated close to the right bank of the Danube, and about 9 km south of the main urban center
40
41 195 of the region, Cernavodă (Fig. 1). The geology of the area has been well studied, because of
42
43 196 the unique outcropping conditions and rich fossiliferous nature of the Lower Cretaceous
44
45 197 deposits (reviewed in Avram et al., 1996; Neagu et al., 1997; Dragastan et al., 1998).
46
47

48 198 Southern Dobrogea is a cratonic area corresponding to the southeastern corner of
49
50 199 Romania. Whether it is considered part of the larger Moesian Platform (Săndulescu, 1984;
51
52 200 Ionesi, 1994), or a distinct craton (the South-Dobrogean Platform; Mutihac and Mutihac,
53

201 2010), researchers agree that it became integrated into the main European Craton towards the
end of the Jurassic, at the latest, with the consolidation of the Cimmerian (Early Alpine:
Triassic–earliest Cretaceous) North Dobrogean fold-and-thrust belt (Seghedi, 2001;
Hyppolite, 2002). The age of its basement is also controversial, with estimates ranging from
Archaic–Early Proterozoic (Mutihac and Mutihac, 2010) to latest Proterozoic (Ionesi, 1994).

The Precambrian basement of Southern Dobrogea is overlain by a flat-lying
sedimentary cover that begins with the lowermost Palaeozoic and ends with the uppermost
Neogene. The sedimentary succession is interrupted by a few major, as well as several less
important, sedimentary hiatuses that separate 5 main sedimentary sequences corresponding to
the Cambrian–Upper Carboniferous, the Permian–Triassic, the Middle Jurassic–Cretaceous,
the Eocene–?Oligocene, and the middle Badenian (middle Miocene)–Upper Pliocene. The
Palaeozoic and lower Mesozoic are known only from the subsurface of Southern Dobrogea,
but Cretaceous and Cenozoic deposits have limited exposures along the main water courses
of the region (Ionesi, 1994; Mutihac and Mutihac, 2010).

The outcropping Cretaceous in Southern Dobrogea is represented mainly by shallow
marine, carbonate platform deposits in the lower part of the system, replaced by more open-
water, chalky facies towards the later part of the period (e.g., Avram et al., 1993, 1996;
Dragastan et al., 1998; Dinu et al., 2007); these crop out only as isolated patches along the
main watercourses of the region (Fig. 1).

The Lower Cretaceous Series consists of several lithostratigraphic units with
complex, partially overlapping and interfingering relationships (Dragastan et al., 1998, 2014).
The lowest (and only artificially) outcropping unit is the Purbeck-type, siliciclastic-evaporitic
Upper Kimmeridgian–Lower Berriasian Amara Formation that represents lagoonal to
continental environments. This unit is covered by the shallow-marine, richly fossiliferous and
locally reefal limestone-dominated Cernavodă Formation (restricted-open lagoonal to

carbonate platform, Upper Berriasian–Lower Hauterivian). A time-correlative unit of the Cernavodă Formation, the limestone-dolomitic Dumbrăveni Formation (Upper Berriasian–Lower Hauterivian), is restricted to the southeastern part of Southern Dobrogea. The Cernavodă and Dumbrăveni formations are covered unconformably by dominantly calcareous deposits with hippuritoid (‘pachyodont’) coquinas, small reefs and lens-like orbitolinid accumulations, referred to the Barremian–Lower Aptian Ostrov Formation by Dragastan et al. (1998), but to the Ramadan Formation (in part) by Avram et al. (1993, 1996). These deposits, formed in littoral to lagoonal and open reef terrace environments, are in turn capped by the fluvial-lacustrine, siliciclastic deposits of the Gherghina Formation, with Middle–Upper Aptian kaolinitic clays and thin coal intercalations. The Lower Cretaceous succession ends with the transgressive, glauconite-bearing, coastal to sublittoral siliciclastic deposits of the Cochirleani Formation (uppermost Aptian–Albian).

The Upper Cretaceous has a significantly more patchy development, mainly restricted to the eastern part of Southern Dobrogea, excepting the weakly glauconitic, chalky-sandy Peștera Formation (Lower Cenomanian) and the marly Dobromiru Formation (Upper Cenomanian) that cover the western-central parts of the area. The younger Cuza Vodă (Turonian), Murfatlar (Santonian–Lower-Middle Campanian), and Satu Nou (Upper Campanian) formations are dominantly chalky, suggesting the instalment of a relatively deeper, offshore depositional environment; neither of these units is known from western Southern Dobrogea.

In total, the Lower Cretaceous of Southern Dobrogea was deposited in a shallow marine, near-shore setting, fluctuating between carbonate platform, lagoonal, coastal-tidal flat, and continental environments (see Avram et al., 1996; Dragastan et al., 1998). Its main characteristic features, such as the observed lithological variability, the areal distribution of the different units, and the presence of several unconformities within the series, are all linked

to eustatic sea-level changes that affected the Southern Dobrogean territory during the Early Cretaceous (Dragastan et al., 1998). The main emergent land in the area was represented by the Central Dobrogean Massif, lying north of the study area, almost completely subaerially exposed and actively eroding during the Cretaceous. Consequently, shallow-marine to continental deposits are restricted mainly to the northern part of Southern Dobrogea, close to its boundary with the Central Dobrogean Massif (marked by the Capidava-Ovidiu Fault), and are replaced by more open marine deposits southward. As summarized above, several littoral, and even continental, sequences occur in this succession, including deposits in the Amara, Cernavodă, Ramadan (~~in part;~~ Avram et al., 1996) and Cochirleni formations, whereas the Gherghina Formation is purely continental, with occasional minor marine interbeds produced during short-term ingressions of the sea.

In the Cernavodă-Cochirleni area the outcropping Mesozoic is restricted to the Lower Cretaceous, and includes deposits belonging to the Cernavodă, Ostrov (or Ramadan), Gherghina, and Cochirleni formations. While the lower–middle part of the Cernavodă Formation is well exposed and widely distributed in this area, its upper part (the lower Hauterivian Vederoasa Member) is unevenly developed. This member is missing in the classical succession from Cernavodă-Hinog, on the right bank of the Danube (Dragastan et al., 1998), but was recently identified in the more eastern Cernavodă-lock section (Dragastan et al., 2014). Similarly, the Ostrov Formation is represented in the area only by its upper subunit (the Lower Aptian Lipnița Member; Dragastan et al., 1998), covering unconformably and transgressively the Valanginian Alimanu Member of the Cernavodă Formation in the southern end of the Cernavodă-Hinog section (Dragastan et al., 1998), and the lower Hauterivian Vederoasa Member in the Cernavodă-lock section (Dragastan et al., 2014).

Northward of the Hinog area, Valanginian deposits of the Alimanu Member are overlain directly by the Middle–Upper Aptian continental deposits of the Gherghina

Formation. These continental deposits also cover the *Orbitolina*-bearing calcareous-clayey deposits of the Lipnița Member towards the south, marking the advancement of emerged areas towards the central parts of Southern Dobrogea, including the Cernavodă-Cochirleni area, during this time interval (Avram et al., 1996). Marine conditions returned in the study area again in the latest Aptian, with a transgression marked by widespread deposition of the glauconitic, siliciclastic coastal to innermost shelf deposits of the Cochirleni Formation. These uppermost Aptian to Albian sands and sandstones cover transgressively all the underlying deposits, belonging to the Cernavodă, Ostrov, or Gherghina formations. Siliciclastic shallow-marine sedimentation continued into the Early Cenomanian, with the chalky-glauconitic deposits of the Peștera Formation.

4. Palaeontology

The isolated theropod tooth UAIC (SCM1) 615 (formerly in the collections of the Hârșova Museum, registered with no. 200; Fig. 2A) was described in a short note by Simionescu (1913), who referred it to *Megalosaurus* cf. *superbus*, a taxon erected by Sauvage (1882) from the Gault ('mid'-Cretaceous: Albian) of the Paris Basin, France. The Gault material described by Sauvage (1882; see also Sauvage, 1876) includes several isolated teeth that were deemed by Simionescu (1913) to be more similar to the Cochirleni tooth than are the teeth of *Megalosaurus bucklandi* (Buckland, 1824). Subsequently, the French Gault material was referred to the new genus *Erectopus* by Huene (1923), who also noted differences between it and the type species *M. bucklandi*.

The convoluted taxonomic history of *Erectopus superbus* was recently reviewed by Allain (2005), who established that both the isolated teeth first mentioned by Sauvage (1876) and the skeletal elements described by Sauvage (1882) belong to the same taxon, for which the name *Erectopus superbus* was retained. Allain (2005) regarded *Erectopus* as a member of

Carnosauria (= basal Tetanurae), an opinion also shared by Molnar (1990) and Holtz et al. (2004a), whereas the latest review of the Tetanurae (Carrano et al., 2012, p. 254) considered *Erectopus superbis* “a non-carcharodontosaurian allosauroid, possibly a metriacanthosaurid.” Accordingly, if we are following the original assessment of Simionescu (1913) but updating with contemporary taxonomy, the Cochirleni theropod tooth should now be considered referable to the basal tetanuran *Erectopus superbis*. However, the referral of this tooth to *Erectopus superbis* (or a close relative) was considered to be unsupported by positive evidence by Molnar (1990) and Holtz et al. (2004a). In order to re-assess this referral and to understand the exact taxonomic and phylogenetic affinities of UAIC (SCM1) 615 (Fig. 2B, 3), we provide here a detailed description of its morphology followed by a thorough comparative study of this tooth based on large datasets of theropod dental measurements and discrete characters compiled by Hendrickx and Mateus (2014) and Hendrickx et al. (2015a).

We note that in his review of Romanian dinosaurs, Grigorescu (2003) erroneously considered UAIC (SCM1) 615 as being referred by Simionescu to the taxon *Megalosaurus dunkeri* Kohen (sic; actually, *Megalosaurus dunkeri* Dames, 1884). This is clearly a simple misreading of Simionescu’s identification. Additionally, such a referral is also contradicted by the absence of mesial serrations in the holotype tooth of *M. dunkeri*, considered by Carrano et al. (2012) to represent an indeterminate theropod. The Dobrogea tooth, on the other hand, has mesial serrations (see below).

4.1. Age of UAIC (SCM1) 615

The age of UAIC (SCM1) 615 has been contentious, due to the uncertainties concerning its place of origin. Although it is often mentioned as originating from Cochirleni village (e.g., Grigorescu, 2003; Turculeț and Brânzică, 2012), this has not been definitively established. According to the original report of Simionescu (1913), the tooth came from the upper part of

the Lower Cretaceous limestone succession exposed in the cliffs extending from Cernavodă to Cochirleni along the right bank of the Danube. The corresponding entry from the Hârşova Museum registry states that it was found in the ‘Cochirleni quarry’, a location that presently cannot be identified precisely. The only rocks to be quarried in the area are the calcareous deposits of either the Cernavodă or Ostrov formations, particularly the ones that crop out in the Danube bank cliffs between Cernavodă-Hinog-Cochirleni. Finally, although the mention ‘Cochirleni’ is usually considered to refer to Cochirleni village, it should be mentioned that the cliff-forming hill that extends between Cernavodă and Cochirleni is also known by the same name (Fig. 1). Taking all of this evidence into consideration, it is thus reasonable to conclude that the tooth was most likely found in the Lower Cretaceous limestone succession exposed in the Danube cliffs between Cernavodă and Cochirleni.

Based on the location of the discovery, in the upper part of the local limestone succession, and the age of the deposits from Cernavodă-Cochirleni known to him, Simionescu (1913) considered the tooth to be of Barremian age. Subsequently, the age of the tooth was given as Valanginian–Barremian (Weishampel, 1990; Weishampel et al., 2004) or Valanginian (e.g., Grigorescu, 2003), but without any supporting information.

New attempts have been made to more precisely constrain the age of UAIC (SCM1) 615. Dragastan et al. (2014) recently sampled the limestone matrix holder of the tooth, and reported from these samples an assemblage of foraminiferans, ostracods and microproblematicae (= *incertae sedis* microorganisms) that characterize their ‘Biozone IX with *Meandrospira favrei*’, of latest Valanginian age in the local lithostratigraphic scheme. In parallel, we also sampled the same limestone holder – a yellowish white, friable lime mudstone – that yielded a poor and badly preserved calcareous nannoplankton assemblage with *Watznaueria barnesiae*, *W. ovata*, *Nannoconus steinmanni*, *N. kamptneri*, *N. globulus*, *Calcicalathina* sp., *Speetonia colligata* and *Cyclagelosphaera deflandrei* (M. C. Melinte-

Dobrinescu, personal communication, November 2013), an assemblage that suggests a Berriasian–Hauterivian age of the limestone holder.

Since it is not clear if the limestone holder came from the same site as the tooth itself, we managed to take a second sample from the limestone matrix still partly filling the pulp cavity of the tooth, which must definitively be identical with the rocks the tooth was found in. This second, much smaller sample yielded only very scarce specimens of *Watznaueria barnesia*, *Cyclagelosphaera margerelii* and *Diazomatolithus lehmanni* (M. C. Melinte-Dobrinescu, personal communication, November 2013), the latter two taxa having a peak in abundance during the Berriasian and, especially, the Valanginian.

In the nannoplankton succession reported previously by Avram et al. (1993) and derived from a systematic sampling of the Southern Dobrogean Lower Cretaceous, the concurrent presence of *Speetonia colligata*, *Calcicalathina oblongata*, *Diazomatolithus lehmanni* and *Nannoconus steinmanni* was noted in samples derived from the Alimanu Member of the Cernavodă Formation. These assemblages were interpreted to represent the nannoplankton zone CC3 of Sissingh (1977), of late Valanginian age. A comparable age was assigned to a roughly similar nannoplankton assemblage reported from the Lower Cretaceous of the Mecsek Mountains, Hungary, by Császár et al. (2000).

Together, all the available evidence (Simionescu's original account, geographic and geologic records, foraminifera, ostracods, microproblematicae, and calcareous nannoplankton) thus suggests that UAIC (SCM1) 615 originates from the Alimanu Member of the Cernavodă Formation, and it is most probably of late Valanginian age.

4.2. Description and comparisons

Specimen UAIC (SCM1) 615 is a large (total length, as preserved, is about 100 mm; Figs. 2, 3) lateral tooth of a theropod dinosaur, with a crown base length (CBL) of 29 mm, crown

base width (CBW) of 16.25 mm, crown height (CH) of 85.5 mm, and apical length (AL) of 91 mm (terminology following Smith et al., 2005 and Hendrickx et al., 2015b). It is remarkably well preserved, with the enamel in pristine condition. It preserves most of the crown and a small basal part of the root, but the crown tip is broken off, with an estimated 5 mm missing in the apical region.

In its present state, the mesial edge and part of the mesial third of the tooth are embedded in the limestone holder (Fig. 2B), although the tooth was once removed (see above, History of collecting and curation; Fig. 2A). Accordingly, it is exposed so that all faces of the tooth are widely visible, including the root region, except for the mesial surface.

Only the basal-most part of the root is preserved, and it is more complete near the mesial margin (Fig. 3B, C). Here, broken areas around the crown-root contact area (cervix) reveal details of the pulp cavity development, as well as the pattern of the dentine thickness variation (Fig. 3B–D). The crown also exhibits a transverse break at about two-thirds of its length (not present so obviously in the original figure of the specimen in Simionescu, 1913), and adjacent to it, the distal carina is also slightly chipped distal to mid-length. The labial face is superficially split near this break (Fig. 3A), while a more prominent region of damage appears on the lingual face, where a large (13 x 5 mm), slightly triangular wedge is broken off, exposing the deeper parts of the dentine (Fig. 3C). The damage to the lingual side apparently occurred after the original description of the tooth (Fig. 2), an observation that is concordant with the complex curatorial history of the specimen.

The basal-most, exposed part of the mesial face lacks the enamel cover (Fig. 3C, D), suggesting that this area already belongs to the root region. The mesial edge of the preserved crown base appears to be wider than the distal one, and is largely rounded transversely. Accordingly, the basal cross-section is teardrop-shaped (lanceolate); it is rounded mesially, but narrows distally into a small carina (Fig. 3D). As mentioned above, the pulp cavity is

exposed basally, being partly filled with a whitish-light gray limestone that is reminiscent of the matrix holder lithology. The pulp cavity narrows rapidly towards the cervix, as it is about 7.1 mm wide (labiolingually) at the apical-most part of the preserved root, but only about 4.5 mm wide at the base of the crown. In parallel, the enamel-dentine wall of the tooth becomes thicker: it is 3.5 mm thick in the apical-most part, 4.4 mm at the base of the crown, but thickens to 5.0–5.8 mm near the apical-most part of the basal break of the crown (Fig. 3B). Mirroring the outside cross-section, the contour of the pulp cavity is also teardrop-shaped (Fig. 3D).

The tooth is ziphodont and only very slightly recurved distally. The distal edge is nearly straight across its length, being very mildly concave in its basal half and slightly convex near its apex (Fig. 2, 3A). Thus, the apex is placed roughly at the distal margin of the tooth crown base. The mesial edge, as shown in the original publication of Simionescu (1913), is strongly convex across its entire length (Fig. 2A). The tooth is labiolingually compressed (Fig. 3B), with a crown base ratio ($CBR = CBW/CBL$) of 0.56, within the normal range of variation of most theropods. This differs from the thinner teeth of some, but not all, carcharodontosaurids ($CBR < 0.50$), and the much thicker incrassate teeth of derived tyrannosauroids and conical teeth of spinosaurids ($CBR > 0.75$) (Sereno et al., 1996; Brusatte et al., 2010a; Hendrickx and Mateus, 2014; Hendrickx et al., 2015a).

The crown cross-section is slightly asymmetrical labiolingually when it is seen in distal view. In this view, when the carina is facing directly distally, one side of the crown has a more pronounced bulge than its counterpart (about 8.5 mm wide, measured from the carina, vs. 6 mm on the other side; Fig. 3B); based on comparisons with the teeth of *Mapusaurus* (Coria and Currie, 2006), the more bulging side can be interpreted as the lingual one. This asymmetry diminishes apically, where both sides become about equally convex. The distal carina itself twists slightly sideways (labially) in apical direction, such that it is located closer

to the labial face where it terminates at the crown apex, and the lingual face of the denticles is exposed distally (Fig. 3B, F). This twist of the distal carina is accompanied by a similar outline of the lingual side; in distal view, this is somewhat convex basally, but becomes flat to slightly concave in the apical two-thirds of the crown. A similar S-shaped curvature of the crown, albeit more pronounced and different in details, was also reported in *Mapusaurus* and *Giganotosaurus* (Coria and Currie, 2006), and in indeterminate carcharodontosaurid teeth from Morocco (Richter et al., 2013).

The distal carina extends along the entire tooth height (Fig. 3A–C). It is covered with minute serrations across its entire preserved length; the denticles are proximodistally subrectangular, with a mesiodistal long axis that is greater than the apicobasal long axis (Fig. 3E–H). They are either roughly perpendicular to the tooth margin, or their long axes are oriented obliquely, such that they point slightly apically. The tip of the apex is broken off, so it is not possible to determine whether the serrations continued over the apex of the tooth. There are approximately 12.5 serrations (denticles) per 5 millimetres at the midpoint of the carina. Serration shape and size remain relatively constant across the carina, although the serrations near the midpoint and closer to the base of the carina (12 denticles per 5 mm; Fig. 3G, H) are slightly smaller than those near the apex (9 denticles per 5 mm; Fig. 3E, F). Changes in serration size are gradual across the carina, not sudden or sporadic.

Although they are all more or less rectangular in shape, the apical denticles are relatively shorter proximodistally than the more basal ones. Most of the denticles have slightly rounded, asymmetrically convex triangular tips, instead of being simply squared-off, and they do not hook as in troodontids and to a lesser extent abelisaurids (Hendrickx and Mateus, 2014). Other denticles near the apex, however, show a faint concavity along their tips, giving them a bilobate aspect, although this is both less conspicuous and far less regularly developed than reported in *Tyrannotitan* (Novas et al., 2005). The denticles are

separated by simple, linear grooves (interdenticular slits or sulcae) along their entire length. The interdenticular space between adjacent denticles is broad, measuring more than a third of the apicobasal width of a denticle (Fig. 3E, G). This space continues onto the surface of the crown as a very short interdenticular sulcus (“blood groove” of Currie et al., 1990). These sulci are so short and indistinct that they are only visible under low angle light.

Little can be said about the mesial carina, as it is not visible in the current state of the specimen, buried in the limestone matrix. Based on the description of Simionescu (1913), however, it is covered across its length with minute serrations; these decrease in size towards the base of the crown. Simionescu (1913) reported approximately 15 serrations (denticles) per 5 millimetres at the midpoint of the carina, meaning that the mesial denticles are slightly smaller than those on the distal carina. The denticle size difference index (DSDI: Rauhut and Werner, 1995) is 1.2, within the range of variation of most theropods (Hendrickx and Mateus, 2014). As Simionescu (1913) already pointed out, the presence of a mesial carina that extends towards the base of the crown sets apart UAIC (SCM1) 615 from *Megalosaurus bucklandii* where this stops well above the cervix (Benson et al., 2008), and it is instead similar to ‘*M. superbus*’ (Sauvage, 1876, 1882) in this respect.

The external enamel surface exhibits two forms of ornamentation. First, the majority of the labial and lingual faces are covered by relatively smooth enamel that exhibits a subtle form of braided texture visible under low angle light (Fig. 3A, C, E). This texture is made up of a series of very faint, apico-basally running ridges; these are of unequal lengths, starting at different points of the crown height, but none extends the whole length of the crown. The two longest ridges are placed near the distal carina. The enamel is also finely granulated.

Second, near the carinae on both labial and lingual surfaces there are marginal undulations: wrinkles in the enamel that stand out in bas relief (Brusatte et al., 2007). These are much better preserved and visible near the distal carina, where they are so pronounced

that they are clearly observable in normal light (Fig. 3A–C, G, H). Here, about 17 unevenly developed wrinkles are present along the crown height; in the basal half of the crown, the wrinkles extend about 6.5 mm onto the crown. These are elongate, such that they are longer than twice the space separating each undulation. The wrinkles project obliquely (in the mesiobasal direction) relative to the carina. They are apically concave, with a near-horizontal segment on the crown, and curve apically as they approach the carina (at about 45°) with a tendency to become tangential to the distal edge. The wrinkles are especially well developed, prominent and closely spaced in the basal part of the crown (about 7 wrinkles/16 mm; Fig. 3C, G)), but become more widely spaced and indistinct apically (about 3 wrinkles/16 mm). Apically, however, the wrinkles are somewhat wider and longer, extending over about half of the crown fore-aft length. Again, a slight asymmetry is present between the two sides of the crown in wrinkle development as well, these being better expressed on the more rounded, convex lingual face, but less well expressed on the flatter labial face (Fig. 3A, C, H). On the presumed labial face, only some of the basal-most wrinkles, particularly the second and third one, appear well defined.

Towards the base of the crown a few of the wrinkles continue across the labial and lingual surfaces as very subtle transverse undulations. Most conspicuous of these is a 3.5 mm wide horizontal swelling that crosses the crown, at the level of wrinkles 2 and 3; this swelling is clearly visible on both sides of the crown (Fig. 3. A, C). There are no lateral flutes, apico-basal ridges, or longitudinal grooves on the labial or lingual faces, either in the centre of the tooth or paralleling the carinae. Instead, the labial and lingual faces are uniformly convex, giving the tooth its teardrop-shaped outline in cross section.

5. Discussion

5.1. Identification of UAIC (SCM1) 615

1
2 501 The isolated tooth from Cochirleni can be referred to Theropoda based on its large size,
3
4 502 recurved and labiolingually compressed morphology, and presence of a continuous series of
5
6 503 well-defined serrations on the distal carina.
7

8 504 Besides theropods, certain derived crocodyliforms – the sebecosuchians of Colbert
9
10 505 (1946; see also Turner and Sertich, 2010; Pol and Powell, 2011; Rabi and Sebők, 2015) – are
11
12 506 also known to possess remarkably theropod-like, laterally compressed and serrated teeth, not
13
14 507 unlike the morphology shown by UAIC (SCM1) 615. However, most sebecosuchian teeth are
15
16 508 significantly smaller than the Southern Dobrogean specimen, especially in the case of the
17
18 509 Cretaceous members of the clade (e.g. *Baurusuchus*; Carvalho et al., 2005). Even the largest,
19
20 510 caniniform teeth of the largest representatives of Sebecosuchia, such as the Miocene
21
22 511 *Barinasuchus* (Paolillo and Linares, 2007), are somewhat smaller than UAIC (SCM1) 615;
23
24 512 moreover, these teeth are slightly conical and less laterally compressed than the Southern
25
26 513 Dobrogean tooth. Finally, it should be noted that the oldest known members of Sebecosuchia
27
28 514 appear beginning in the Late Cretaceous (e.g. Kellner et al., 2014), and are thus significantly
29
30 515 younger than UAIC (SCM1) 615. Similarly, ziphodont crocodyliform teeth (i.e. with true
31
32 516 denticles along their carinae) are reported in Europe only beginning in the Albian (Ősi et al.,
33
34 517 2015), and these are both significantly smaller and different in morphology from the
35
36 518 Dobrogean tooth. Taken together, these suggest that the hypothesis of sebecosuchian
37
38 519 affinities of UAIC (SCM1) 615 can be discarded with confidence, and it indeed represents a
39
40 520 theropod tooth.
41
42

43 521 We used four techniques to identify which type of theropod UAIC (SCM1) 615 likely
44
45 522 belongs to (see also Supplementary Material).
46

47 523 First, we conducted a Principal Components Analysis (PCA) based on a large
48
49 524 database that includes a broad and representative sample of theropod teeth. This dataset was
50
51 525 compiled by Hendrickx et al. (2015a), which built upon the earlier studies of Smith et al.
52
53

(2005) and Larson and Currie (2013), and it or a similar version has been used in recent studies to identify isolated theropod teeth (e.g., Williamson and Brusatte, ~~2013~~2014; Brusatte and Clark, 2015). It comprises nearly 1000 theropod teeth scored for six measurements (CBL, CBW, CH, AL, MC, and DC, the latter two measuring the density of serrations per 5 mm at the midpoint of the mesial and distal carina, respectively). UAIC (SCM1) 615 was added to this dataset, the data were log-transformed prior to analysis, missing values for measurements were estimated with a mean value for that measurement from across the sample, and then a PCA was run using a correlation matrix. The analysis was conducted in PAST v2.17 (Hammer et al., 2001).

In the resulting two dimensional morphospace (Fig. 4), UAIC (SCM1) 615 plots close to many teeth belonging to carcharodontosaurids, along with some teeth belonging to spinosaurids and tyrannosauroids. It falls within the convex hull (maximum morphospace occupation area) of carcharodontosaurids only, although it is closely outside of the edges of spinosaurid and tyrannosauroid space. It also falls within the 95% confidence interval ellipse for carcharodontosaurids, but not within the ellipse of any other group (Supplementary Information). This exercise indicates that UAIC (SCM1) 615 is most similar to carcharodontosaurids.

Secondly, we used the log-transformed dataset that we also used for the PCA to conduct a clustering analysis. We performed the analysis in PAST v2.17, using the paired group algorithm and the correlation similarity measure. In the resulting dendrogram, UAIC (SCM1) 615 groups with a handful of teeth belonging to carcharodontosaurids, tyrannosauroids, and *Allosaurus* (Supplementary Information).

Third, we used the tooth measurement database to conduct a discriminant analysis in PAST v3.0 (Hammer et al., 2001). This analysis uses pre-determined groups (in this case, taxonomic clusters) to create a morphospace in which these groups are maximally separated.

This allows teeth of unknown affinities, such as UAIC (SCM1) 615, to be classified according to which taxonomic group it is most similar to in this discriminant morphospace. In total, 67.79% of other teeth are classified correctly when they are treated as having uncertain affinities and their measurements are used to classify them in discriminant space, indicating that this exercise returns reasonable results. Our analysis classifies the Romanian tooth as a carcharodontosaurid. Furthermore, the analysis places UAIC (SCM1) 615 within the convex hulls for carcharodontosaurids and tyrannosauroids, and the 95% confidence ellipses for carcharodontosaurids, coelophysoids, and neovenatorids.

Fourth, we ran a phylogenetic analysis by including UAIC (SCM1) 615 in the discrete character dataset of theropod dental features published by Hendrickx and Mateus (2014). The Romanian specimen was scored as a lateral tooth in this analysis. The analysis was conducted in TNT (Goloboff et al., 2008), and resulted in 224 most parsimonious trees (686 steps, consistency index of 0.338, retention index of 0.566). The strict consensus topology is moderately well resolved and places the Romanian tooth as the sister taxon to *Carcharodontosaurus* (Supplementary Material). This sister taxon pair is recovered as the sister clade to a grouping of the derived carcharodontosaurids *Mapusaurus* and *Giganotosaurus*.

Several synapomorphies support the carcharodontosaurid affinities of UAIC (SCM1) 615. The sister group relationship with *Carcharodontosaurus* is supported by two features: a roughly straight distal margin of the crown (character 68) and pronounced marginal undulations in the enamel that are well visible in normal light (character 112). The broader clade of UAIC (SCM1) 615, *Carcharodontosaurus*, *Mapusaurus*, and *Giganotosaurus* (= Carcharodontosaurinae, as defined by Brusatte and Sereno, 2008, and Carrano et al., 2012) is linked by numerous characters, including: large teeth with a crown height greater than 6 cm (character 65), a bowed or sigmoid distal carina in distal view (character 82), marginal

undulations that are at least twice as long mesiodistally as the space separating each undulation (character 111), and marginal undulations present on both mesial and distal sides of the crown (character 113).

The Romanian specimen also lacks many keystone dental synapomorphies of other theropod clades, based on the clade diagnoses of Hendrickx and Mateus (2014) and other cladistic studies that include dental characters. UAIC (SCM1) 615 does not possess the hooked distal denticles of some Abelisauridae, the strongly labially deflected distal carina and pronounced transverse enamel undulations extending across the labial and lingual tooth faces of Ceratosauridae, the incrassate teeth with apicobasal enamel flutes and deeply veined enamel surface texture of Spinosauridae, and the large transverse undulations of some basal allosauroids (Hendrickx and Mateus, 2014). It also lacks the thickened incrassate teeth of derived tyrannosauroids (Brusatte et al., 2010a) and the large and strongly hooked (or pointed) denticles of troodontids and therizinosauroids (e.g., Turner et al., 2012; Brusatte et al., 2014; Hendrickx and Mateus, 2014). The large size, as well as recurved and ziphodont shape of UAIC (SCM1) 615 is strikingly different from the non-ziphodont therizinosauroids, ornithomimosaurs, alvarezsauroids, and most troodontids, which have conical, leaf-shaped, or peg-like teeth (when teeth are present) (e.g., Holtz et al., 2004a; Turner et al., 2012; Brusatte et al., 2014). Finally, besides its remarkably large size, the presence of serrations indicates that UAIC (SCM1) 615 does not belong to groups such as alvarezsauroids, oviraptorosaurs, basal troodontids, or avialans, which have unserrated crowns (e.g., Turner et al., 2012; Hendrickx and Mateus, 2014).

In summary, the four analyses all support carcharodontosaurid affinities for UAIC (SCM1) 615. Both overall tooth proportions and discrete phylogenetic characters point to a carcharodontosaurid identification, and the discriminant function analysis and phylogenetic analysis both explicitly recover the tooth as a carcharodontosaurid. For this reason we refer

this tooth to Carcharodontosauridae. Moreover, it appears to belong to a clade that unites very derived and large-sized carcharodontosaurids (*Carcharodontosaurus*, *Giganotosaurus*, and *Mapusaurus*), separated as such and named Carcharodontosaurinae by Brusatte and Sereno (2008) and Carrano et al. (2012). The well-resolved internal topology of this clade, as recovered in our analysis, is congruent with results of previous analyses based on larger sets of characters from across the skeleton (e.g., Coria and Currie, 2006; Brusatte and Sereno, 2008; Brusatte et al., 2009; Ortega et al., 2010; Eddy and Clarke, 2011; Canale et al., 2015), and offers some support for considering the Romanian carcharodontosaurid from Southern Dobrogea as more closely related to the African *Carcharodontosaurus* than to the clade of the South American giant carcharodontosaurids *Giganotosaurus* or *Mapusaurus*.

Two final notes are worth adding. First, our analyses also incorporated carcharodontosaurids that are usually found to be basal within the clade, such as *Acrocanthosaurus* and *Eocarcharia* (e.g., Harris, 1998; Sereno and Brusatte, 2008; Carrano et al., 2012) as well as a host of other allosauroids, including members of Neovenatoridae (*Neovenator*, *Australovenator* and *Fukuiraptor*), a clade that is often recovered as sister-taxon to carcharodontosaurids within Carcharodontosauria (e.g., Benson et al., 2010; Carrano et al., 2012; but see Novas et al., 2013; Porfiri et al., 2014, for an alternate placement of neovenatorids in general). Both PCA and phylogenetic analysis clearly identified UAIC (SCM1) 615 as more closely comparable morphologically to derived carcharodontosaurids than to either basal carcharodontosaurids or to any other allosauroid subclade.

Second, our datasets also included teeth of *Erectopus*, the genus erected for ‘*Megalosaurus*’ *superbus* to which UAIC (SCM1) 615 was originally referred. Again, our analyses clearly indicate that there are no close morphological and morphometric similarities between the two, which is in accordance with the suggestion of Carrano et al. (2012) that *Erectopus* represents a non-carcharodontosaurid taxon, while our analysis identifies UAIC

(SCM1) 615 as a carcharodontosaurid. Instead, *Erectopus* groups with abelisauroids in the phylogenetic analysis. This is somewhat surprisingly, as Allain (2005) and Carrano et al. (2012) both identified *Erectopus* as a tetanuran. It should be noted, however, that Albian-aged abelisauroids are known from the same general area (eastern France) as that yielding the material referred to *Erectopus* (Accarie et al., 1995; Carrano and Sampson, 2008), raising the intriguing possibility that this taxon may represent an abelisauroid instead of an allosauroid tetanuran as suggested by Allain (2005) and Carrano et al. (2012). However, it must be remembered that this phylogenetic analysis is based on dental characters only, so it is probably more likely that *Erectopus* is a tetanuran with a dentition convergent to some extent with those of certain abelisauroids.

5.2. Body size of UAIC (SCM1) 615

One of the most salient and remarkable features of UAIC (SCM1) 615 is its large size. In the large and comprehensive sample of theropod teeth from our dataset, tooth size (estimated based on crown height – CH, and used as a rough proxy of body size) ranges from 2.2 mm (in the dromaeosaurid *Saurornitholestes* and the coelurosaur of uncertain affinities *Richardoestesia*) to 117.1 mm in the gigantic tyrannosauroid *Tyrannosaurus*. The Romanian specimen UAIC (SCM1) 615, with a CH of 85.5 mm, is ranked in the 60-80% maximum size (~ CH) range of the sample, and has a CH that is 73% of the largest tyrannosauroid teeth.

Most of the teeth in the dataset (over 61% of the 966 measured teeth) are very small to small (less than 25 mm CH), and less than 10% of these fall in the 60-100% CH size categories.

Teeth larger than UAIC (SCM1) 615 make up less than 5% of the total sample, and they represent only five taxa: the megalosaurid *Torvosaurus*, the tyrannosauroid *Tyrannosaurus*, the basal carcharodontosaurid *Acrocanthosaurus*, and the derived carcharodontosaurines

Carcharodontosaurus and *Giganotosaurus*. Compared to other carcharodontosaurids, UAIC

(SCM1) 615 is smaller than the largest teeth of *Acrocanthosaurus* (9% difference), *Carcharodontosaurus* (20%), and *Giganotosaurus* (12.5%) in the dataset, but is 13% bigger than the largest tooth of *Mapusaurus*.

It is thus reasonable to conclude that UAIC (SCM1) 615 belonged to a large-sized carcharodontosaurid, comparable to, even if somewhat smaller than, the truly gigantic carcharodontosaurines *Giganotosaurus* and *Carcharodontosaurus* (Serenó et al., 1996; Calvo and Coria, 1998; Therrien and Henderson, 2007), taxa that were recovered as possible close relatives of the Romanian carcharodontosaurid by our phylogenetic analysis. This, in turn, corroborates growing evidence that very large body size was acquired very early in carcharodontosaurid history, since the earliest potential members of the clade are already of relatively large size (Rauhut, 2011). The oldest potential carcharodontosaurid is *Veterupristisaurus*, represented by isolated vertebrae that indicate an animal between 8.5 and 10 meters in total body length (compared to 11.5+ meters in *Acrocanthosaurus* and more derived carcharodontosaurids) (Rauhut, 2011). These specimens are known from the uppermost Jurassic of Tanzania, eastern Africa (Rauhut, 2011; Carrano et al., 2012; see below), predating at most ~18 million years (Mya) the occurrence of likely even larger-sized carcharodontosaurids in the Valanginian of Southern Dobrogea, Romania.

The inferred large body size of the South Dobrogean theropod is also remarkable as virtually all other dinosaur remains reported previously from Romania (both from the Early Cretaceous Cornet assemblage and the much later, end Cretaceous Hațeg Island fauna) are significantly smaller, and many have been interpreted as insular dwarfs (e.g., Weishampel et al., 1993, 2003; Benton et al., 2006, 2010; Stein et al., 2010; Ősi et al., 2014). Although other Romanian theropod dinosaurs were not particularly dwarfed (e.g. Brusatte et al., 2013), they were nonetheless small (Nopcsa, 1902; Csiki and Grigorescu, 1998; Csiki et al., 2010; Brusatte et al., 2013). This bias towards small bodied Romanian theropods was also

interpreted as a consequence of their insular habitat (Csiki and Grigorescu, 1998), as all previously reported theropod remains come from within the Carpathian Orogen, an area with an archipelago-type palaeogeography during the Cretaceous (Dercourt et al., 2000; Csontos and Vörös, 2004; Csiki-Sava et al., 2015). By contrast, UAIC (SCM1) 615 was found in shallow marine deposits bordering the emerged areas of Central Dobrogea, part of the stable cratonic areas of Europe and connected at least intermittently to the Ukrainian Shield since the Late Jurassic (Fig. 5A). Although cratonic Europe was also transformed into an archipelago of islands during much of the Cretaceous, these islands were often both larger in size and more stable in space and time than were the transient emerged areas of the Tethyan archipelagoes. As such, it is conceivable that the Southern Dobrogean carcharodontosaurid was less constrained by space or resource limitations than the Tethyan insular dinosaurs, allowing it to retain a large body size.

5.3. UAIC (SCM1) 615 and Valanginian dinosaur distribution

Besides documenting the presence of large-sized mainland carcharodontosaurids in the Lower Cretaceous of Romania, UAIC (SCM1) 615 is also important in that it fills a significant gap in our knowledge on the composition and distribution of the Early Cretaceous dinosaurs in Europe. In their review of dinosaur occurrences, Weishampel et al. (2004) listed 83 Early Cretaceous dinosaur localities spread throughout Europe, more than half of these being known from the later part (Barremian–Albian) of that epoch; only around a dozen localities were listed from each age of the early part of the Early Cretaceous (Berriasian, Valanginian, and Hauterivian). Even despite a significant increase in Early Cretaceous dinosaur discoveries in Europe in recent years (e.g., Royo-Torres et al., 2009; Cobos et al., 2010, 2014; Galton, 2009; Norman, 2010, 2013; Pereda-Suberbiola et al., 2011, 2012; Sachs and Hornung, 2013; Blows and Honeysett, 2014), these remain very strongly biased towards

western and southwestern Europe (especially the UK, France and Spain). Frustratingly, no occurrences are known from the entire central, eastern and southern Europe for the Berriasian–Hauterivian time interval except for two from Romania: the Berriasian–Valanginian locality of Cornet (e.g., Jurcsák and Popa, 1979, 1983; Jurcsák, 1982; Benton et al., 1997) in the northern Apuseni Mountains of northwestern Romania, and the carcharodontosaurid tooth (Simionescu, 1913) from the Valanginian of Cochirleni, in Southern Dobrogea, southeastern Romania we are describing here (Fig. 5A).

Our identification of the Romanian tooth as a carcharodontosaurid documents the presence of this clade in Europe in the very early Cretaceous. This is significant, as carcharodontosaurids were widely distributed tens of millions of years later, in the middle Cretaceous (Aptian to Cenomanian), in western Gondwana (Africa and South America, see below). Despite the recent discoveries documenting that the clade was also present in North America and Asia during the middle Cretaceous (e.g., Sereno et al., 1996; Currie and Carpenter, 2000; Brusatte et al., 2009, 2012), there has been only very few occurrences in Europe, most importantly the Barremian-aged *Concavenator* from Spain (Ortega et al., 2010; see below). The carcharodontosaurid tooth from Southern Dobrogea is substantially older than *Concavenator*, demonstrating that carcharodontosaurids appeared in Europe earlier than previously thought and were a long-term component of the European mainland Early Cretaceous faunas. It also suggests that habitat-related palaeobiological differentiation might have been already present between the cratonic, stable European mainland, with a dinosaur fauna made up of normal-sized (even very large) taxa, and the islands from the mobile Alpine areas of the Mediterranean Neo-Tethys, with by now dwarfed dinosaurs such as those described from the Berriasian–Valanginian Cornet assemblage in northwestern Romania (Benton et al., 2006).

This Valanginian carcharodontosaurid represents an important datapoint not only for the Romanian Lower Cretaceous, but also for that of wider Eurasia. The Valanginian is a poorly documented age in dinosaur evolution, with very few precisely dated fossil occurrences from anywhere in the world (e.g., Weishampel et al., 2004). The best record of Valanginian dinosaurs is from Europe, with fewer and less well dated occurrences known from Asia, some of which have debatable or controversial dates. These include sites in Japan (e.g., Manabe and Hasegawa, 1995; Matsukawa et al., 2006; but see Kusuhashi et al., 2009 and Evans and Matsumoto, 2015, supporting an alternative, younger age of these assemblages) and in Thailand (e.g., Buffetaut and Suteethorn, 1998, 2007, with age constraints according to Racey, 2009; Racey and Goodall, 2009). Occurrences of possible Valanginian age from China (e.g., Jerzykiewicz and Russell, 1991; Shen and Mateer, 1992; Lucas and Estep, 1998) are either poorly constrained as early Early Cretaceous, or were shown subsequently to be younger than Valanginian (Lucas, 2006; Tong et al., 2009). Rare dinosaur remains of possible Valanginian (or ‘Neocomian’) age were also reported from southern Africa (e.g., De Klerk et al., 2000) and, tentatively, from North America (e.g., Lucas, 1901; McDonald, 2011, with age assignments according to Sames et al., 2010; Cifelli et al., 2014).

As one of the two known reports of Valanginian dinosaurs in Europe east of France, the Southern Dobrogean dinosaur record fills a huge palaeogeographic gap between the western European and the eastern Asian dinosaur faunas. Moreover, none of these early Early Cretaceous dinosaur assemblages from outside Europe include carcharodontosaurids (see below), as theropods are represented by coelurosauians interpreted either as compsognathids (Gishlick and Gauthier, 2007) or basal ornithomimosaurs (Choiniere et al., 2012) in southern Africa, metriacanthosaurid allosauroids (‘sinraptorids’) in Thailand (Buffetaut and Suteethorn, 2007), and indeterminate allosauroids (Pérez-Moreno et al., 1993), non-

carcharodontosaurid tetanurans (Carrano et al., 2012) or enantiornithine birds (Lacasa Ruiz, 1989), besides indeterminate taxa (Carrano et al., 2012), in western Europe. This may suggest that carcharodontosaurids had not achieved a wide geographic distribution by this point in time, and that their more cosmopolitan distribution came later, during the middle Cretaceous.

Finally, the presence of the Cochirleani carcharodontosaurid might hint at the presence of palaeobiogeographic provinciality between the western and the eastern parts of Europe, partly mirroring those reported from the later part of the Late Cretaceous (e.g., Le Loeuff and Buffetaut, 1995; Weishampel et al., 2010; Ősi et al., 2012; Csiki-Sava et al., 2015). In the reasonably well sampled, and significantly better known, western European dinosaur faunas, Valanginian large carnivorous dinosaurs include non-carcharodontosaurid tetanurans (*Becklespinax*), as well as indeterminate allosauroids or indeterminate theropods (often described as '*Megalosaurus*' *dunkeri*, '*M.*' *insignis* or '*M.*' *oweni*), none of which can be referred positively to Carcharodontosauridae (Carrano et al., 2012). The apparently provincial geographic distribution of the large-bodied theropods suggests that some degree of faunal differentiation was occurring within the European mainland, most probably promoted by geographic distance. Notably, this intra-European differentiation in theropod assemblages appears to stand in contrast with the faunal homogeneity reported in the case of the ornithopods from the UK and Romania (e.g., Galton, 2009). It is important, however, to re-emphasize at this point that the Valanginian dinosaur fossil record is both exceedingly poor and patchy, even in Europe. Accordingly, further discoveries are needed to verify and support (or contradict) the presence of such a distribution pattern pointing to palaeobiogeographic provinciality inside Europe, as the one suggested by our carcharodontosaurid identification for UAIC (SCM1) 615.

5.4. UAIC (SCM1) 615 and carcharodontosaurid evolution and palaeobiogeography

Carcharodontosauridae were long considered as an exclusively Gondwanan group of theropods (e.g., Allain, 2002; Novas et al., 2005) since their first discovery in northern Africa (e.g., Stromer, 1931), and subsequent description of a host of referred taxa from the Aptian–Cenomanian of Africa and South America (Coria and Salgado, 1995; Sereno et al., 1996; Novas et al., 2005; Coria and Currie, 2006; Brusatte and Sereno, 2007; Sereno and Brusatte, 2008; Cau et al., 2013). This view started to change with the identification of the Early Cretaceous (Aptian–Albian) *Acrocanthosaurus* from North America as a basal carcharodontosaurid (e.g., Sereno et al., 1996; Harris, 1998; Sereno 1999; Brusatte and Sereno, 2008), suggesting that the clade had a wider, Neopangean palaeobiogeographic distribution by the mid–late Early Cretaceous. Such a wide distribution, even a cosmopolitan one, was further supported by the discovery of definitive carcharodontosaurids in the Lower Cretaceous of Europe (Ortega et al., 2010), and in the upper Lower to lower Upper Cretaceous of China (Brusatte et al., 2009, 2010b, 2012; Mo et al., 2014; Lü et al., 2014, 2016).

Together, the available evidence pointed to an early, pre-mid Early Cretaceous origin of the carcharodontosaurids, followed by their dispersal across Laurasia and western Gondwana beginning at least by the Aptian (Fig. 5B), a scenario that is concordant with the tentatively suggested presence of early carcharodontosaurids in the Upper Jurassic of Tanzania, which are based on fragmentary specimens (Rauhut, 2011; Carrano et al., 2012). It is also concordant with the widespread appearance of carcharodontosaurids in the fossil record starting with the Aptian, when they are reported in Africa (*Eocarcharia*; Sereno and Brusatte, 2008), South America (Vickers-Rich et al., 1999), North America (*Acrocanthosaurus*; Stovall and Langston, 1950; Harris, 1998; Currie and Carpenter, 2000; Eddy and Clarke, 2011), Europe (Canudo and Ruiz-Omeñaca, 2003; Pereda-Suberbiola et al.,

2012), and eastern Asia (*Kelmayisaurus*; Brusatte et al., 2012; ~~Lü et al., 2014~~; Mo et al., 2014; [Lü et al., 2016](#)).

During the Albian–Turonian, carcharodontosaurids became especially abundant and diverse in Africa (*Carcharodontosaurus*, *Saurodon*; Stromer, 1931; Sereno et al., 1996; Brusatte and Sereno, 2007; Le Loeuff et al., 2012; Cau et al., 2013; Richter et al., 2013) and South America (*Tyrannotitan*, *Giganotosaurus*, *Mapusaurus*, alongside with indeterminate carcharodontosaurids; Coria and Salgado, 1995; Calvo and Coria, 1998; Novas et al., 2005; Coria and Currie, 2006; Casal et al., 2009; Candeiro et al., 2011; Canale et al., 2015; Fig. 5B). They were still present during this time interval in other continents, as well: in North America with *Acrocanthosaurus* until the Albian (D’Emic et al., 2012), in Europe until the Cenomanian (Vullo et al., 2007; Csiki-Sava et al., 2015), and in Eastern Asia with *Shaochilong* until the Turonian (Brusatte et al., 2009, 2010b; see also Chure et al., 1999).

After dominating terrestrial ecosystems at least in Africa, South America and eastern Asia during the Albian–Turonian (Brusatte et al., 2009; Coria and Salgado, 2005; Novas et al., 2013), carcharodontosaurids were considered to disappear from the fossil record after the Turonian in both Asia (Brusatte et al., 2009) and South America (e.g., Coria and Salgado, 2005; Calvo et al., 2006; Novas et al., 2013), to be replaced by other groups of large theropods such as tyrannosaurids in parts of Laurasia and abelisaurids in parts of Gondwana. Canale et al. (2009) even cautioned against assigning isolated theropod teeth from post-Cenomanian deposits of South America to Carcharodontosauridae (e.g., Canudo et al., 2008; Casal et al., 2009; Salgado et al., 2009) due to their morphological similarity to those of the abelisaurid *Skorpiovenator*. Recently, however, more diagnostic cranial remains were reported to suggest the survival of carcharodontosaurids into the latest Cretaceous (Campanian–Maastrichtian) in Brazil (Azevedo et al., 2013).

Contrasting with this rich and relatively continuous fossil record of Carcharodontosauridae starting with the Aptian, the first half of its evolutionary history is very poorly documented (Fig. 5B). Prior to the identification of UAIC (SCM1) 615, only two occurrences of pre-Aptian Cretaceous carcharodontosaurids were reported, one from the Barremian of Spain (Ortega et al., 2010; Gasca et al., 2014) and the other from the Barremian of Thailand (Buffetaut and Suteethorn, 2012). The Early Cretaceous *Kelmayisaurus* from Xinjiang, western China, was recognized as a carcharodontosaurid of possibly ?Valanginian to Aptian in age by Brusatte et al. (2012), but the deposits yielding these remains (the Lianmugui, or Lianmuxin, Formation of the Tugulu Group) were dated as Aptian–Albian by Eberth et al. (2001; see also Tong et al., 2009). An important temporal gap – of about 20 to 28 millions of years, according to the dates in Gradstein et al. (2012) – thus stretched between the oldest, tentatively assigned carcharodontosaurids from the Oxfordian–Tithonian of Tanzania, including the formally erected *Veterupristisaurus* (Rauhut, 2011; see also Carrano et al., 2012), and those that started to appear in the fossil record in the Barremian and then spread widely during the Aptian. Referral of UAIC (SCM1) 615 to Carcharodontosauridae partially fills this frustrating gap, effectively halving this shadowy period in the evolutionary history of the group.

Furthermore, our analyses tentatively cluster the Dobrogean theropod with the derived members of the Carcharodontosaurinae to the exclusion of the more basal, but significantly younger non-carcharodontosaurine carcharodontosaurids *Eocarcharia* and *Acrocanthosaurus*. If this placement is correct, then the Romanian tooth indicates that Carcharodontosaurinae diverged from other carcharodontosaurids considerably earlier than hitherto recognized.

The previously known fossil record of the clade suggested that Carcharodontosaurinae originated sometime between the Aptian and Albian, as basal carcharodontosaurids (*Acrocanthosaurus*, *Concavenator*, *Eocarcharia*) were moderately diverse in the Barremian–

1
2 848 Aptian, followed by the appearance of many fossils of carcharodontosaurines beginning in
3
4 849 the Albian (Fig. 5B). The proposed affinities of the oldest carcharodontosaurid material –
5
6 850 including isolated teeth referred to as ‘*Megalosaurus*’ *ingens* – from the east African Upper
7
8 851 Jurassic, considered to be reminiscent of the Aptian–Albian *Acrocanthosaurus* (Rauhut,
9
10 852 2011), was also consistent with this evolutionary scenario. Now, our identification of UAIC
11
12 853 (SCM1) 615 as a carcharodontosaurid dinosaur sharing important dental apomorphies with
13
14 854 the derived Carcharodontosaurinae advocates the emergence of this clade (or at least the very
15
16 855 large size and dental morphology characterizing it) well before the Albian, during or even
17
18 856 before the Valanginian, and relegates taxa such as *Eocarcharia*, *Acrocanthosaurus* and
19
20
21 857 *Concavenator* (the dentition of *Shaochilong* is unknown) as late-surviving members of the
22
23 858 basal carcharodontosaurid radiation, with a relatively plesiomorphic dentition.
24

25 859 Besides shifting the emergence of the carcharodontosaurines earlier in time,
26
27 860 identification of UAIC (SCM1) 615 as a carcharodontosaurid also has interesting
28
29 861 palaeobiogeographic implications. As already noted, recent discoveries show that
30
31 862 Carcharodontosauridae is not an endemic Gondwanan clade as was once proposed (e.g.,
32
33 863 Novas et al., 2005), with the identification of its widespread, Pangaeian distribution during the
34
35 864 late Early Cretaceous (Sereno et al., 1996; Harris, 1998; Chure et al., 1999; Sereno, 1999;
36
37 865 Brusatte and Sereno, 2008; Ortega et al., 2010; Brusatte et al., 2009, 2012; Mo et al., 2014).
38
39 866 However, within Carcharodontosauridae itself, some palaeogeographic patterns have been
40
41 867 widely accepted. For example, it has been widely acknowledged that Carcharodontosaurinae
42
43 868 is a endemic subclade of Gondwanan carcharodontosaurids (e.g., Sereno 1999; Holtz et al.,
44
45 869 2004b; Brusatte and Sereno, 2007; Sereno and Brusatte, 2008; Novas et al., 2013), as
46
47
48 870 previously all its recognized members were restricted strictly to either Africa (Stromer, 1931;
49
50 871 Sereno et al., 1996; Brusatte and Sereno, 2007) or South America (Coria and Salgado, 1995;
51
52 872 Novas et al., 2005; Coria and Currie, 2006). Moreover, intra-clade relationships of
53

Carcharodontosaurinae were still adhering to patterns of continental fragmentation and vicariant evolution, with a basal split between the Albian–Cenomanian African *Carcharodontosaurus* and the Giganotosaurini, uniting the similarly Albian–Cenomanian southern South American *Giganotosaurus* and *Mapusaurus* (together with *Tyrannotitan*, if this taxon is also recovered within Carcharodontosaurinae; e.g., Novas et al., 2005, 2013).

This scenario is now challenged by our finding that the Southern Dobrogean carcharodontosaurid UAIC (SCM1) 615 may nest inside Carcharodontosaurinae. If true, such an affinity would suggest that the origin of Carcharodontosaurinae was not a southern, vicariant by-product of the Gondwana-Laurasia separation, a major palaeogeographic event that is considered to have been well underway by the end of the Jurassic, and essentially completed by the mid-Early Cretaceous (see Weishampel et al., 2010). Indeed, during this time palaeogeographic connections and faunal interactions were virtually non-existent between the northern Tethyan (European) and southern Tethyan (western Gondwanan, but essentially African) areas of the Mediterranean (e.g., Canudo et al., 2009; ~~see below~~), which makes a vicariant hypothesis intuitive. However, if the Romanian tooth represents a carcharodontosaurine, then it implies a much more complicated palaeogeographic history of the clade, which is not so clearly linked to continental breakup.

The palaeogeographic position of the Southern Dobrogean carcharodontosaurine in cratonic Europe, north of the Neo-Tethys, together with its significantly older age compared to other carcharodontosaurines, could indicate that separation of the carcharodontosaurine lineage took part in Europe and not in western Gondwana as previously assumed. This would also mean that representatives of this lineage were subsequently – after the Barremian – introduced to Africa and South America via trans-Tethyan dispersal, most probably at a time when faunal interactions between the southern and northern margins of the Mediterranean Tethys were resumed, after the early Barremian (Canudo et al., 2009).

Alternatively, it can be hypothesized that appearance of carcharodontosaurines in Southern Dobrogea is a consequence of southern immigration originating in western Gondwana, often considered the place of origin for this clade. However, this scenario has several potential caveats. Although Europe has been considered as forming part of a larger Eurogondwanan palaeobioprovince during the early Early Cretaceous (Ezcurra and Agnolín, 2012), and occasional trans-Tethyan faunal connections have been recognized between Africa and Europe during Late Jurassic to Early Cretaceous times (e.g., Gheerbrant and Rage, 2006), these interchanges either pre-dated the Berriasian (e.g., Gardner et al., 2003; Knoll and Ruiz-Omeñaca, 2009), or post-dated the Barremian (Canudo et al., 2009; Torcida Fernández-Baldor et al., 2011), with no positive evidence for actual faunal exchanges taking place during the ‘Neocomian’ (Berriasian–Hauterivian) time interval.

More recently, some potential evidence has emerged for Gondwana-to-Europe interchange during the ‘Neocomian’. The presence of the basal rebbachisaurid *Histriasaurus* (Dalla Vecchia, 1998) in the upper Hauterivian–lower Barremian of Croatia has been cited as indicative of very early and very rapid northward dispersal of this clade from western Gondwana (southern South America; Carballido et al., 2012; Fanti et al., 2015). Timing of this particular dispersal event was even constrained to the Berriasian–Valanginian time interval (Fanti et al., 2015), which makes it roughly contemporaneous with the record of the Southern Dobrogean carcharodontosaurine. It was also suggested, however, that dispersal of the line leading to *Histriasaurus* was mediated by the northward drift of the Apulian Microplate (= Adria; see Bosselini, 2002), a continental sliver acting as a passive transportation mechanism (‘Noah’s Ark’; ~~KeKenna~~McKenna, 1973) for basal rebbachisaurids after its separation from mainland Africa (e.g., Torcida Fernández-Baldor et al., 2011). Furthermore, the palaeogeographical separation between Africa and Adria (and thus the effective movement of the presumed ark) is considered to be at most an incipient one

1
2 923 during the Early Cretaceous by Bossellini (2002) and Zarcone et al. (2010), with spatial
3
4 924 continuity still present between the two landmasses, while deep-water basins continued to
5
6 925 separate Adria from the European Craton. Accordingly, although the presence of
7
8 926 *Histriasaurus* can represent a case of northward range extension of rebbachisaurids during
9
10 927 the Berriasian–Valanginian, it took place not strictly speaking into Europe, but only reached
11
12 928 the northernmost extremity of Adria, a northerly peninsular extension of the African
13
14 929 mainland. It was only starting with the Barremian that rebbachisaurids dispersed as far north
15
16 930 as the European cratonic areas, including Iberia and the British Isles (Mannion, 2009;
17
18 931 Mannion et al., 2011; Torcida Fernández-Baldor et al., 2011), a time when faunal
19
20 932 interchanges between Europe and Africa are considered to have been well underway (e.g.,
21
22 933 Gheerbrant and Rage, 2006; Canudo et al., 2009).

23
24
25 934 Unlike *Histriasaurus*, the taxon represented by UAIC (SCM1) 615 was an inhabitant
26
27 935 of the European mainland. It is thus unclear to what extent the example of rebbachisaurid
28
29 936 range extension into (present-day) Europe during the early Early Cretaceous, as potentially
30
31 937 testified by the discovery of the Croatian taxon, would also be applicable for the Southern
32
33 938 Dobrogean carcharodontosaurine. The available evidence suggests that these two cases are
34
35 939 very different, and that faunal connections during this time interval are not documented
36
37 940 between the African and European cratons as already pointed out by Gheerbrant and Rage
38
39 941 (2006).

40
41 942 Absence of documented faunal interactions weakens support for a scenario of south-
42
43 943 to-north immigration of derived carcharodontosaurines in Europe at the very beginning of the
44
45 944 Cretaceous, and would argue instead for a local, European development to explain the
46
47 945 presence of a Valanginian carcharodontosaurine in Southern Dobrogea. The pre-Barremian
48
49 946 presence of carcharodontosaurids in Europe is also consistent with their appearance in the
50
51 947 Barremian–Aptian fossil record of Eastern Asia, with Europe acting as a stepping stone in the
52

eastward dispersal of the clade. Similarly, the presence of Aptian carcharodontosaurids in North America likely requires the presence of pre-Aptian members of the clade in Europe, since faunal exchanges between these two landmasses are known to have been halted before the Aptian (e.g., Kirkland et al., 1999). Interestingly, it appears that only basal carcharodontosaurids were able to spread into the northern Laurasian landmasses, while the derived carcharodontosaurines dispersed exclusively across the Neo-Tethys, into western Gondwana. The causes of these distribution patterns remain as yet unknown, and further support – in the form of new carcharodontosaurid discoveries from the early-middle part of the Early Cretaceous – is required to better uphold such a scenario.

We finally reiterate that if the Romanian tooth does not belong to a carcharodontosaurine, but instead is artefactually grouping with them in the phylogenetic analysis because of the very incomplete nature of the material, then the traditional story of Carcharodontosaurinae as a product of vicariant evolution driven by the breakup of Pangea will remain strongly supported. However, even in such case UAIC (SCM1) 615 would still record the presence of early-occurring large carcharodontosaurid theropods with a very characteristic carcharodontosaurine-type dentition in the eastern part of the European craton, adding to known early Early Cretaceous theropod (and dinosaur) diversity, and potentially documenting dinosaur faunal provinciality in Europe and worldwide.

6. Conclusions

We re-describe and interpret the affinities of one of the most significant historical dinosaurian specimens of Romania, an isolated but well-preserved theropod tooth from Southern Dobrogea. Our extensive analyses suggest carcharodontosaurid relationships for this tooth, while the available evidence – including novel calcareous nannoplankton sampling – supports its Valanginian age. The Southern Dobrogean theropod tooth represents the oldest record of

Carcharodontosauridae in the Cretaceous, and the second oldest globally, eclipsed only by a collection of isolated specimens from the Upper Jurassic of eastern Africa. As one of the only two known Valanginian dinosaurian occurrences from Central and Eastern Europe, this record advances our understanding of European dinosaur distribution during the early Early Cretaceous, and also fills an important palaeogeographic gap between Western European and Eastern Asian dinosaurian assemblages of the Valanginian.

Based on dental apomorphies, our analyses further identify UAIC (SCM1) 615 as a possible member of Carcharodontosaurinae, a subclade of derived and gigantic carcharodontosaurids formerly known to be restricted to the Albian–Cenomanian of western Gondwana (Africa and South America). If this finding is correct, the Southern Dobrogean specimen documents the emergence of Carcharodontosaurinae earlier than previously recognized, thus also indicating an earlier acquisition of their characteristically large size. Based on currently known palaeogeographic and chronostratigraphic constraints on the evolution of Carcharodontosauridae, it appears that not only did this clade have a wide distribution, but that crucial events of its evolutionary history such as the emergence of the derived carcharodontosaurines took place north of the Tethys, in cratonic Europe, instead of western Gondwana and as the result of vicariant evolution driven by the Gondwana-Laurasia split, as was formerly suggested. In such a case, instead of endemic evolution the emergence of the western Gondwanan mid-Cretaceous carcharodontosaurines was the result of a north-to-south trans-Tethyan dispersal that took place somewhere between the Valanginian and the Aptian. Recognizing a potential carcharodontosaurine dispersal event from Europe into western Gondwana adds further support for the presence of important palaeogeographic ties between the two realms during the second half of the Early Cretaceous.

Acknowledgements

This research was supported by the National Research Council of Romania (CNCS) grant PN-III-D-PCE-2011-3-0381 and a Sepkoski grant of the Paleontological Society for Z.Cs.-S. S.L.B. is supported by a Marie Curie Career Integration Grant EC630652, the Division of Paleontology of the American Museum of Natural History, and the School of GeoSciences of the University of Edinburgh. He thanks Mátyás Vremir, Radu Totoianu, and Mark Norell for many hours of fun discussion on Romanian fossils, and for supporting his work and travel in Romania. We thank Mihai Brânzilă and Paul Țibuleac (UAIC) for access to the specimen, for allowing us to collect samples for the nannoplankton analyses, and for their help and collegiality during our visit to Iași, as well as Ilie Turculeț for sharing information about the history of the specimen. Mihaela C. Melinte-Dobrinescu has gracefully analyzed the nannoplankton samples derived from UAIC (SCM1) 615; her contribution was essential in assessing the age of the specimen. Finally, we thank the reviewers Eric W.A. Mulder (Denekamp, the Netherlands) and Xabier Pereda-Suberbiola (Bilbao, Spain), as well as Associated Editor Elena Jagt-Yazykova, for their useful comments and suggestions that helped improve ~~a~~previous versions of the manuscript.

References

- Accarie, H., Beaudoin, B., Dejax, J., Friès, G., Michard, J. G., & Taquet, P. (1995). Découverte d'un dinosaure théropode nouveau (*Genusaurus sisteronis* n. g., n. sp.) dans l'Albien marin de Sisteron (Alpes-de-Haute-Provence, France) et extension au Crétacé inférieur de la lignée cératosaurienne. *Comptes Rendus de l'Académie des Sciences Paris, IIa* 320, 327–334.
- Allain, R. (2002). Discovery of a megalosaur (Dinosauria, Theropoda) in the Middle Bathonian of Normandy (France) and its implications for the phylogeny of basal Tetanurae. *Journal of Vertebrate Paleontology*, 22, 548–563.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

- 1
21023 Allain, R., (2005). The enigmatic theropod dinosaur *Erectopus superbus* (Sauvage, 1882)
3
41024 from the Lower Albian of Louppy-le-Chateau (Meuse, France). In K. Carpenter, K. (Ed.),
5
61025 *The Carnivorous Dinosaurs* (pp. 72–86). Indiana University Press, Bloomington; Indiana
7
81026 University Press, pp. 72–86.
9
101027 Avram, E., Costea, I., Dragastan, O., Muțiu, R., Neagu, T., Șindilar, V., & Vinogradov, C.
11
121028 (1996). Distribution of the Middle-Upper Jurassic and Cretaceous facies in the Romanian
13
141029 eastern part of the Moesian Platform. *Revue Roumaine de Géologie*, 39-40, 3–33.
15
161030 Avram, E., Szasz, L., Antonescu, E., Baltreș, A., Iva, M., Melinte, M., Neagu, T., Rădan, S.,
17
181031 & Tomescu, C., (1993). Cretaceous terrestrial and shallow marine deposits in northern
19
201032 South Dobrogea (SE Romania). *Cretaceous Research*, 14, 265–305.
21
22
231033 Avram, E., Costea, I., Dragastan, O., Muțiu, R., Neagu, T., Șindilar, V., Vinogradov, C.,
24
251034 1996. Distribution of the Middle-Upper Jurassic and Cretaceous facies in the Romanian
26
271035 eastern part of the Moesian Platform. *Revue Roumaine de Géologie* 39-40, 3–33.
28
291036 Azevedo, R. P. F. de, Simbras, F. M., Furtado, M. R., Candeiro, C. R. A., & Bergqvist, L. P.,
30
311037 (2013). First Brazilian carcharodontosaurid and other new theropod dinosaur fossils from the
32
331038 Campanian-Maastrichtian Presidente Prudente Formation, São Paulo State, southeastern
34
351039 Brazil. *Cretaceous Research*, 40, 131–142.
36
371040 Benson, R. B. J., Barrett, P. M., Powell, H. P., & Norman, D. B., (2008). The taxonomic
38
391041 status of *Megalosaurus bucklandii* (Dinosauria, Theropoda) from the Middle Jurassic of
40
411042 Oxfordshire, UK. *Palaeontology*, 51, 419–424.
42
431043 Benson, R. B. J., Carrano, M. T., & Brusatte, S. L., (2010). A new clade of archaic large-
44
451044 bodied predatory dinosaurs (Theropoda: Allosauroidae) that survived to the latest Mesozoic.
46
471045 *Naturwissenschaften*, 97, 71–78.
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Benton, M. J., Cook, E., Grigorescu, D., Popa, E., & Tallódi, E. (1997). Dinosaurs and other tetrapods in an Early Cretaceous bauxite-filled fissure, northwestern Romania. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 130, 275–292.

Benton, M. J., Csiki, Z., Grigorescu, D., Redelstorff, R., Sander, M., Stein, K., & Weishampel, D. B. (2010). Dinosaurs and the island rule: the dwarfed dinosaurs from Hateg Island. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 293, 438–454.

Benton, M. J., Minter, N. J., & Posmoşanu, E. (2006). Dwarfing in ornithopod dinosaurs from the Early Cretaceous of Romania. In: Z. Csiki, Z. (Ed.), *Mesozoic and Cenozoic Vertebrates and Paleoenvironments; Tributes to the Career of Prof. Dan Grigorescu* (pp. 79–87). *Ars Docendi*, Bucharest: *Ars Docendi*, pp. 79–87.

Benton, M. J., Csiki, Z., Grigorescu, D., Redelstorff, R., Sander, M., Stein, K., Weishampel, D. B., 2010. Dinosaurs and the island rule: the dwarfed dinosaurs from Hateg Island. *Palaeogeography, Palaeoclimatology, Palaeoecology* 293, 438–454.

Blows, W. T., & Honeysett, K. (2014). First Valanginian *Polacanthus foxii* (Dinosauria, Ankylosauria) from England, from the Lower Cretaceous of Bexhill, Sussex. *Proceedings of the Geologists' Association*, 125, 233–251.

Bosselini, A. (2002). Dinosaurs “re-write” the geodynamics of the eastern Mediterranean and the paleogeography of the Apulia Platform. *Earth-Science Reviews*, 59, 211–234.

Brânzilă, M. (ed.); (2010). *Academicianul Ion Simionescu - savant şi dascăl al neamului* (p. 182). Iaşi: Ed. Universităţii "Alexandru Ioan Cuza", Iaşi, 182 pp.

Brusatte, S. L., Sereno, P. C., 2007. A new species of *Carcharodontosaurus* (Dinosauria: Theropoda) from the Cenomanian of Niger and a revision of the genus. *Journal of Vertebrate Paleontology* 27, 902–916.

Brusatte, S. L., Sereno, P. C., 2008. Phylogeny of Allosauroida (Dinosauria: Theropoda): comparative analysis and resolution. *Journal of Systematic Palaeontology* 6, 155–182.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

1
2 1071 Brusatte, S. L., Clark, N. D. L., 2015. Theropod dinosaurs from the Middle Jurassic (Bajocian-
3
4 1072 Bathonian) of Skye, Scotland. *Scottish Journal of Geology* 51, 157–164. doi: 10.1144/sjg2014-
5
6 1073 022
7
8 1074 Brusatte, S., Benson, R. B. J., Carr, T. D., Williamson, T. E., & Sereno, P. C. (2007). The
9
10 1075 systematic utility of theropod enamel wrinkles. *Journal of Vertebrate Paleontology*, 27,
11
12 1076 1052–1056.
13
14 1077 Brusatte, S. L., Benson, R. B. J., Chure, D. J., Xu, X., Sullivan, C., & Hone, D. W. E. (2009).
15
16 1078 The first definitive carcharodontosaurid (Dinosauria: Theropoda) from Asia and the
17
18 1079 delayed ascent of tyrannosaurids. *Naturwissenschaften*, 96, 1051–1058.
19
20 1080 Brusatte, S. L., Benson, R. B. J., & Xu, X. (2012). A reassessment of *Kelmayisaurus*
21
22 *petrolicus*, a large theropod dinosaur from the Early Cretaceous of China. *Acta*
23
24 *Palaeontologica Polonica*, 57, 65–72.
25
26 1083 Brusatte, S. L., & Clark, N. D. L. (2015). Theropod dinosaurs from the Middle Jurassic
27
28 (Bajocian-Bathonian) of Skye, Scotland. *Scottish Journal of Geology*, 51, 157–164. doi:
29
30 1084 10.1144/sjg2014-022
31
32 1085 Brusatte, S., Chure, D. J., Benson, R. B. J., & Xu, X. (2010b). The osteology of *Shaochilong*
33
34 *maortuensis*, a carcharodontosaurid (Dinosauria: Theropoda) from the Late Cretaceous of
35
36 1087 Asia. *Zootaxa*, 2334, 1–46.
37
38 1088 Brusatte, S. L., Lloyd, G. T., Wang, S. C., & Norell, M. A. (2014). Gradual assembly of
39
40 1089 avian body plan culminated in rapid rates of evolution across the dinosaur-bird transition.
41
42 1090 *Current Biology*, 24, 2386–2392.
43
44 1091 Brusatte, S. L., Norell, M. A., Carr, T. D., Erickson, G. M., Hutchinson, J. R., Balanoff, A.
45
46 1092 M., Bever, G. S., Choiniere, J. N., Makovicky, P. J., & Xu, X. (2010a). Tyrannosaur
47
48 1093 paleobiology: new research on ancient exemplar organisms. *Science*, 329, 1481–1485.
49
50 1094

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

1
2 1095 Brusatte, S., Chure, D.J., Benson, R.B.J., Xu, X., 2010b. The osteology of *Shaochilong*
3
4 1096 *maortuensis*, a carcharodontosaurid (Dinosauria: Theropoda) from the Late Cretaceous of
5
6 1097 Asia. *Zootaxa* 2334, 1–46.
7
8 1098 Brusatte, S.L., Benson, R.B.J., Xu, X., 2012. A reassessment of *Kelmayisaurus petrolicus*, a
9
10 1099 large theropod dinosaur from the Early Cretaceous of China. *Acta Palaeontologica Polonica*
11
12 1100 57, 65–72.
13
14 1101 Brusatte, S.L., Vremir, M., Csiki-Sava, Z., Turner, A.H., Watanabe, A., Erickson, G.M.,
15
16 1102 Norell, M.A., 2013. The osteology of *Balaur bondoc*, an island-dwelling dromaeosaurid
17
18 1103 (Dinosauria: Theropoda) from the Late Cretaceous of Romania. *Bulletin of the American*
19
20 1104 *Museum of Natural History* 374, 3–100. doi: 10.1206/798.1
21
22 1105 Brusatte, S.L., Lloyd, G.T., Wang, S.C., Norell, M.A., 2014. Gradual assembly of avian body
23
24 1106 plan culminated in rapid rates of evolution across the dinosaur–bird transition. *Current*
25
26 1107 *Biology* 24, 2386–2392.
27
28 1108 Brusatte, S. L., & Sereno, P. C. (2007). A new species of *Carcharodontosaurus* (Dinosauria:
29
30 1109 *Theropoda*) from the Cenomanian of Niger and a revision of the genus. *Journal of Vertebrate*
31
32 1110 *Paleontology*, 27, 902–916.
33
34 1111 Brusatte, S. L., & Sereno, P. C. (2008). Phylogeny of Allosauroida (Dinosauria: Theropoda):
35
36 1112 comparative analysis and resolution. *Journal of Systematic Palaeontology*, 6, 155–182.
37
38 1113 Brusatte, S. L., Vremir, M., Csiki-Sava, Z., Turner, A. H., Watanabe, A., Erickson, G. M., &
39
40 1114 Norell, M. A. (2013). The osteology of *Balaur bondoc*, an island-dwelling dromaeosaurid
41
42 1115 (Dinosauria: Theropoda) from the Late Cretaceous of Romania. *Bulletin of the American*
43
44 1116 *Museum of Natural History*, 374, 3–100. doi: 10.1206/798.1
45
46 1117 Buckland, W.; (1824). Notice on the *Megalosaurus* or great fossil lizard of Stonesfield.
47
48 1118 *Transactions of the Geological Society*, 21, 390–397.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

1
2 1119 Buffetaut, E., & Suteethorn, V., (1998). Early Cretaceous dinosaurs from Thailand and their
3
4 1120 bearing on the early evolution and biogeographical history of some groups of Cretaceous
5
6 1121 dinosaurs. In: ~~Lucas, S. G.~~ Lucas, ~~Kirkland, J. I.~~ Kirkland, ~~Estep, J. W.~~ Estep (Eds.), *Lower*
7
8 1122 *and Middle Cretaceous Terrestrial Ecosystems* (pp. 205–210), New Mexico Museum of
9
10 1123 Natural History and Science Bulletin, 14, pp. 205–210.
11
12 1124 Buffetaut, E., & Suteethorn, V., (2007). A sinraptorid theropod (Dinosauria: Saurischia)
13
14 1125 from the Phu Kradung Formation of northeastern Thailand. *Bulletin de la Société Géologique*
15
16 1126 *de France*, 178, 497–502.
17
18 1127 Buffetaut, E., & Suteethorn, V., (2012). A carcharodontid theropod (Dinosauria, Saurischia)
19
20 1128 from the Sao Khua Formation (Early Cretaceous, Barremian) of Thailand. In: ~~Royo-Torres,~~
21
22 ~~R. Royo-Torres,~~ ~~Gascó, F.~~ Gascó, & ~~Alcalá, L.~~ Alcalá (Eds.), *10th Annual Meeting of the*
23
24 *European Association of Vertebrate Palaeontologists* (pp. 27–30). Teruel; Fundamental!, 20,
25
26 pp. 27–30.
27
28 1132 Calvo, J. O., & Coria, A., (1998). New specimen of *Giganotosaurus carolinii* (Coria &
29
30 Salgado, 1995), supports it as the largest theropod ever found. *Gaia*, 15, 117–122.
31
32 1134 Calvo, J. O., Gandossi, P., & Porfiri, J. D., (2006). Dinosaur faunal replacement during
33
34 Cenomanian times in Patagonia, Argentina. In: ~~Evans, S. E.~~ Evans, & ~~Barrett, P. M.~~ Barrett
35
36 (Eds.), *9th Mesozoic Terrestrial Ecosystems and Biota*, (pp. 17–20). Manchester, UK, pp.
37
38 17–20.
39
40 1138 Canale, J. I., Scanferla, C. A., Agnolin, F. L., & Novas, F. E., (2009). New carnivorous
41
42 dinosaur from the Late Cretaceous of NW Patagonia and the evolution of abelisaurid
43
44 1139 theropods. *Naturwissenschaften*, 96, 409–414.
45
46 1140 Canale, J. I., Novas, F. E., & Pol, D., (2015). Osteology and phylogenetic relationships of
47
48 *Tyrannotitan chubutensis* Novas, de Valais, Vickers-Rich and Rich, 2005 (Theropoda:
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: Font: Italic

Formatted: Font: Not Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1143 Carcharodontosauridae) from the Lower Cretaceous of Patagonia, Argentina. *Historical*
1144 *Biology*, 27, 1–32. doi: 10.1080/08912963.2013.861830.
1145 Candeiro, C. R. A., Fanti, F., Therrien, F., & Lamanna, M. C. (2011). Continental fossil
1146 vertebrates from the mid-Cretaceous (Albian-Cenomanian) Alcântara Formation, Brazil, and
1147 their relationship with contemporaneous faunas from North Africa. *Journal of African Earth*
1148 *Sciences*, 60, 79–92.
1149 Canudo, J. I. Barco, J. L., Pereda Suberbiola, X., Ruiz-Omeñaca, J. I., Salgado, L., Torcida
1150 Fernández-Baldor, F., & Gasulla, J. M. (2009). What Iberian dinosaurs reveal about the
1151 bridge said to exist between Gondwana and Laurasia in the Early Cretaceous. *Bulletin de la*
1152 *Société Géologique de France*, 180, 5–11.
1153 Canudo, J. I., Filippi, L. S., Salgado, L., Garrido, A. C., Cerda, I. A., Garcia, R., & Otero, A.
1154 (2008). Theropod teeth associated with a sauropod carcass in the Upper Cretaceous (Plottier
1155 Formation) of Rincón de los Sauces. In Colectivo Arqueológico y Paleontológico de Salas de
1156 los Infantes (Ed.), *Actas de las IV Jornadas Internacionales sobre Paleontología de*
1157 *Dinosaurios y su Entorno* (pp. 321–330). Salas de los Infantes, Burgos.
1158 Canudo, J. I., & Ruiz-Omeñaca, J. I. (2003). Los restos directos de dinosaurios terópodos
1159 (excluyendo Aves) en España. *Ciencias de la Tierra* 26, 347–373.
1160 ~~Canudo, J.I., Filippi, L.S., Salgado, L., Garrido, A.C., Cerda, I.A., Garcia, R., Otero, A.,~~
1161 ~~2008. Theropod teeth associated with a sauropod carcass in the Upper Cretaceous (Plottier~~
1162 ~~Formation) of Rincón de los Sauces. In: Colectivo Arqueológico y Paleontológico de Salas de~~
1163 ~~los Infantes (Ed.), Actas de las IV Jornadas Internacionales sobre Paleontología de~~
1164 ~~Dinosaurios y su Entorno, Salas de los Infantes, Burgos, pp. 321–330.~~
1165 ~~Canudo, J.I. Barco, J.L., Pereda Suberbiola, X., Ruiz Omenaca, J.I., Salgado, L., Torcida~~
1166 ~~Fernández Baldor, F., Gasulla, J.M., 2009. What Iberian dinosaurs reveal about the bridge~~

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: (Default) Times
New Roman

Formatted: Font: Italic

~~said to exist between Gondwana and Laurasia in the Early Cretaceous. Bulletin de la Société
Géologique de France, 180, 5–11.~~

Carballido, J. L., Salgado, L., Pol, D., Canudo, J. I., & Garrido, A. (2012). A new basal
rebbachisaurid (Sauropoda, Diplodocoidea) from the Early Cretaceous of the Neuquén Basin;
evolution and biogeography of the group. *Historical Biology*, 24, 631–654.

~~Carrano, M. T., Benson, R. B. J., & Sampson, S. D. (2012). The phylogeny of Tetanurae
(Dinosauria: Theropoda). *Journal of Systematic Palaeontology*, 10, 211–300.~~

Carrano, M. T., & Sampson, S. D. (2008). The phylogeny of Ceratosauria (Dinosauria:
Theropoda). *Journal of Systematic Palaeontology*, 6, 183–236.

~~Carrano, M. T., Benson, R. B. J., Sampson, S. D., 2012. The phylogeny of Tetanurae
(Dinosauria: Theropoda). *Journal of Systematic Palaeontology* 10, 211–300.~~

Carvalho, I. S., Campos, A. C. A., & Nobre, P. H. (2005). *Baurusuchus salgadoensis*, a
new Crocodylomorpha from the Bauru Basin (Cretaceous), Brazil. *Gondwana Research*, 8,
11–30.

Casal, G., Candeiro, C. R. A., Martinez, R., Ivany, E., & Ibiricu, L. (2009). Dientes de
Theropoda (Dinosauria: Saurischia) de la Formación Bajo Barreal, Cretácico Superior,
Provincia del Chubut, Argentina. *Géobios*, 42, 553–560.

Cau, A., Dalla Vecchia, F. M., & Fabbri, M. (2013). A thick-skulled theropod (Dinosauria,
Saurischia) from the Upper Cretaceous of Morocco with implications for carcharodontosaurid
cranial evolution. *Cretaceous Research*, 40, 251–260.

Choiniere, J. N., Forster, C. A., & De Klerk, W. J. (2012). New information on
Nqwebasaurus thwazi, a coelurosaurian theropod from the Early Cretaceous Kirkwood

Formation in South Africa. *Journal of African Earth Sciences*, 71–72, 1–17.

Chure, D. J., Manabe, M., Tanimoto, M., & Tomida, Y. (1999). An unusual theropod tooth
from the Mifune Group (Late Cenomanian to Early Turonian), Kumamoto, Japan. In:

- 1
2 1192 ~~Tomida, Y. Tomida, Rich, T. H. Rich, & Vickers-Rich, P. Vickers-Rich~~ (Eds.), *Proceedings*
3
4 1193 *of the Second Gondwanan Dinosaur Symposium* (pp. 291–296). *Tokyo*: National Science
5
6 1194 Museum Monographs, 15, ~~Tokyo, pp. 291–296.~~
7
8 1195 Cifelli, R. L., Davis, B. M., & Sames, B. (2014). Earliest Cretaceous mammals from the
9
10 1196 western United States. *Acta Palaeontologica Polonica*, 59, 31–52.
11
12 1197 doi:10.4202/app.2012.0089.
13
14 1198 ~~Cobos, A., Lockley, M. G., Gascó, F., Royo-Torres, R., & Alcalá, L. (2014). Megatheropods~~
15
16 1199 ~~as apex predators in the typically Jurassic ecosystems of the Villar del Arzobispo Formation~~
17
18 1200 ~~(Iberian Range, Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 399, 31–41.~~
19
20
21 1201 ~~doi: 10.1016/j.palaeo.2014.02.008.~~
22
23 1202 Cobos, A., Royo-Torres, R., Luque, L., Alcalá, L., & Mampel, L. (2010). An Iberian
24
25 1203 stegosaurs paradise: The Villar del Arzobispo Formation (Tithonian–Berriasian) in Teruel
26
27 1204 (Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 293, 223–236.
28
29 1205 ~~Cobos, A., Lockley, M. G., Gascó, F., Royo-Torres, R., Alcalá, L., 2014. Megatheropods as~~
30
31 1206 ~~apex predators in the typically Jurassic ecosystems of the Villar del Arzobispo Formation~~
32
33 1207 ~~(Iberian Range, Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology* 399, 31–41. doi:~~
34
35 1208 ~~10.1016/j.palaeo.2014.02.008.~~
36
37 1209 ~~Codrea, V., Godefroit, P., & Smith, T. (2012). First discovery of Maastrichtian (latest~~
38
39 1210 ~~Cretaceous) terrestrial vertebrates in Rusca Montană Basin (Romania). In P. Godefroit (Ed.)~~
40
41 1211 ~~*Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems* (pp. 570–581).~~
42
43 1212 ~~Bloomington: Indiana University Press.~~
44
45 1213 Codrea, V., Vremir, M., Jipa, C., Godefroit, P., Csiki, Z., Smith, T., & Fărcaș, C. (2010).
46
47 1214 More than just Nopcsa’s Transylvanian dinosaurs: A look outside the Hațeg Basin.
48
49 1215 *Palaeogeography, Palaeoclimatology, Palaeoecology* 293, 391–405. doi:
50
51 1216 10.1016/j.palaeo.2009.10.027.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

- 1
2 1217 Codrea, V., Godefroit, P., Smith, T., 2012. First discovery of Maastrichtian (latest
3
4 1218 Cretaceous) terrestrial vertebrates in Rusca Montană Basin (Romania). In: Godefroit, P. (Ed.)
5
6 1219 Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems. Indiana University Press,
7
8 1220 Bloomington, pp. 570–581.
9
10 1221 Colbert, E. H. (1946). *Sebecus*, representative of a peculiar suborder of fossil Crocodilia
11
12 1222 from Patagonia. *Bulletin of the American Museum of Natural History*, 87(4), 217–270.
13
14 1223 Coria, R. A., & Currie, P. J. (2006). A new carcharodontosaurid (Dinosauria, Theropoda)
15
16 1224 from the Upper Cretaceous of Argentina. *Geodiversitas*, 28, 71–118.
17
18 1225 Coria, R. A., & Salgado, L. (1995). A new giant carnivorous dinosaur from the Cretaceous
19
20 1226 of Patagonia. *Nature*, 377, 224–226.
21
22 1227 Coria, R. A., & Salgado, L. (2005). Mid-Cretaceous turnover of saurischian dinosaur
23
24 1228 communities: evidence from the Neuquén Basin. In: Veiga, G. D., Veiga, Spalletti, L. A.
25
26 1229 Spalletti, Howell, J. A., Howell, & Schwartz, E. Schwartz (Eds.), *The Neuquén Basin,*
27
28 1230 *Argentina: a case study in sequence stratigraphy and basin dynamics* (pp. 317–327).
29
30 1231 Geological Society, London, Special Publications, 252, pp. 317–327.
31
32 1232 Coria, R. A., Currie, P. J., 2006. A new carcharodontosaurid (Dinosauria, Theropoda) from the
33
34 1233 Upper Cretaceous of Argentina. *Geodiversitas* 28, 71–118.
35
36 1234 Covacef, Z. (1995). Pionieri ai culturii românești în Dobrogea; Ioan Cotovu și Vasile
37
38 1235 Cotovu. *Analele Dobrogei*, I(1), 127–134.
39
40 1236 Császár, G., Kollányi, K., Lantos, M., Lelkes, G., and Tardine Filacz, E. (2000). A
41
42 1237 Hidasivölgyi Márga Formáció kora és képződési környezete. *Földtani Közlöny*, 130(4), 695–
43
44 1238 723.
45
46 1239 Csiki, Z., & Grigorescu, D. (1998). Small theropods of the Late Cretaceous of the Hateg
47
48 1240 Basin (Western Romania) - an unexpected diversity at the top of the food chain. *Oryctos*, 1,
49
50 1241 87–104.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

- Csiki, Z., Vremir, M., Brusatte, S. L., & Norell, M. A. (2010). An aberrant island-dwelling theropod dinosaur from the Late Cretaceous of Romania. *Proceedings of the National Academy of Sciences*, 107, 15357–15361.
- ~~Csiki-Sava, Z., Codrea, V., Vasile, Ș., 2013. Early Cretaceous dinosaur remains from Dobrogea (southeastern Romania). In: Picot, L. (Ed.), Abstracts, 11th Annual Meeting of the European Association of Vertebrate Palaeontologists, Villers-sur-Mer, France, pp. 28.~~
- Csiki-Sava, Z., Buffetaut, E., Ősi, A., Pereda-Suberbiola, X., & Brusatte, S. L. (2015). Island life in the Cretaceous - faunal composition, biogeography, evolution, and extinction of land-living vertebrates on the Late Cretaceous European archipelago. *Zookeys*, 469, 1–161. doi: 10.3897/zookeys.469.8439.
- ~~Csiki-Sava, Z., Codrea, V., & Vasile, Ș. (2013). Early Cretaceous dinosaur remains from Dobrogea (southeastern Romania). In L. Picot (Ed.), Abstracts, 11th Annual Meeting of the European Association of Vertebrate Palaeontologists (pp. 28). Villers-sur-Mer, France.~~
- Csontos, L., & Vörös, A. (2004). Mesozoic plate tectonic reconstruction of the Carpathian region. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 210, 1–56. doi: 10.1016/j.palaeo.2004.02.033.
- Currie, P. J., & Carpenter, K. (2000). A new specimen of *Acrocanthosaurus atokensis* (Theropoda, Dinosauria) from the Lower Cretaceous Antlers Formation (Lower Cretaceous, Aptian) of Oklahoma, USA. *Geodiversitas*, 22, 207–246.
- Currie, P. J., Rigby, J. K., & Sloan, R. E. (1990). Theropod teeth from the Judith River Formation of southern Alberta, Canada. In: ~~Carpenter, K.~~ Carpenter, & Currie, P. J. Currie (Eds.), *Dinosaur Systematics: Approaches and Perspectives* (pp. 107–125). Cambridge, Boston: Cambridge University Press, ~~pp. 107–125.~~

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

- Dalla Vecchia, F. M., (1998). Remains of Sauropoda (Reptilia, Saurischia) in the Lower Cretaceous (Upper Hauterivian/Lower Barremian) limestones of SW Istria (Croatia). *Geologia Croatica*, 5, 105–134.
- Dames, W., (1884). *Megalosaurus dunkeri*. *Sitzungsberichte Gesellschaft Naturforschender Freunde zu Berlin*, 1884, 186–188.
- De Klerk, W. J., Forster, C. A., Sampson, S. D., Chinsamy, A., & Ross, C. F. (2000). A new coelurosaurian dinosaur from the Early Cretaceous of South Africa. *Journal of Vertebrate Paleontology*, 20, 324–332.
- D’Emic, M. D., Melstrom, K. M., & Eddy, D. R., (2012). Paleobiology and geographic range of the large-bodied Cretaceous theropod dinosaur *Acrocanthosaurus atokensis*. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 333–334, 13–23.
- Dercourt, J., Gaetani, M., Vrielynck, B., Barrier, E., Biju-Duval, B., Brunet, M., Cadet, J. P., Crasquin, S., & Săndulescu, M., (Eds.) (2000). *Atlas Peri-Tethys Palaeogeographical Maps (269 p.)*. Paris: CCGM/CGMW, Paris, 269 pp.
- Dinu, C., Grădinaru, E., Stoica, M., & Diaconescu, V., (2007). *Dobrogea 2007 Field Trip Preparation and Assistance (123 p.)*. University of Bucharest, 123 pp.
- Dragastan, O. N., Antoniadu, C., & Stoica, M. (2014). Biostratigraphy and zonation of the Lower Cretaceous succession from Cernavodă-lock section, South Dobrogea, eastern part of the Moesian Platform (Romania). *Carpathian Journal of Earth and Environmental Sciences*, 9, 231–260.
- Dragastan, O., Neagu, T., Bărbulescu, A., & Pană, I., (1998). *Jurasicul și Cretacicul din Dobrogea Centrală și de Sud (249 p.)*. Bucharest, 249 pp.
- Dragastan, O. N., Antoniadu, C., Stoica, M., 2014. Biostratigraphy and zonation of the Lower Cretaceous succession from Cernavodă-lock section, South Dobrogea, eastern part of the

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

1
21289 | ~~Moesian Platform (Romania). *Carpathian Journal of Earth and Environmental Sciences* 9(1),~~
3
41290 | ~~231–260.~~
5
61291 | Dyke, G. J., Benton, M. J., Posmoşanu, E., & Naish, D. (2011). Early Cretaceous
7
81292 | (Berriasian) birds and pterosaurs from the Cornet bauxite mine, Romania. *Palaeontology*, 54,
9
101293 | 79–95.
11
121294 | Eberth, D. A., Brinkman, D. B., Chen, P.-J., Yuan, F.-T., Wu, X.-C., Li, G., & Cheng,
13
141295 | ~~Xian X.~~-Shen, C. (2001). Sequence stratigraphy, paleoclimate patterns, and vertebrate fossil
15
161296 | preservation in Jurassic–Cretaceous strata of the Juggar Basin, Xinjiang Autonomous Region,
17
181297 | People’s Republic of China. *Canadian Journal of Earth Sciences*, 38, 1627–1644.
19
201298 | Eddy, D. R., & Clarke, J. A. (2011). New information on the cranial anatomy of
21
221299 | *Acrocanthosaurus atokensis* and its implications for the phylogeny of Allosauroidae
23
241300 | (Dinosauria: Theropoda). *PLoS ONE*, 6(3), e17932. doi:10.1371/journal.pone.0017932.
25
261301 | Evans, S. E., & Matsumoto, R. (2015). An assemblage of lizards from the Early Cretaceous
27
281302 | of Japan. *Palaeontologia Electronica*, 18.2.36A, 1–36.
29
301303 | Ezcurra, M. D., & Agnolín, F. I. (2012). A new global palaeobiogeographical model for
31
321304 | the Late Mesozoic and Early Tertiary. *Systematic Biology*, 61, 553–566.
33
341305 | doi:10.1093/sysbio/syr115.
35
361306 | Fanti, F., Cau, A., Cantelli, L., Hassine, M., & Audatore, M. (2015). New information on
37
381307 | *Tataouinea hannibalis* from the Early Cretaceous of Tunisia and implications for the tempo
39
401308 | and mode of rebbachisaurid sauropod evolution. *PLoS ONE*, 10(4), e0123475.
41
421309 | doi:10.1371/journal.pone.0123475.
43
441310 | Galton, P. M. (2009). Notes on Neocomian (Lower Cretaceous) ornithomimid dinosaurs from
45
461311 | England - *Hypsilophodon*, *Valdosaurus*, “*Camptosaurus*”, “*Iguanodon*” - and referred
47
481312 | specimens from Romania and elsewhere. *Revue de Paléobiologie*, 28, 211–273.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

- Gardner, J. D., Evans, S. E., & Sigogneau-Russell, D. (2003). New albanerpetontid amphibians from the Early Cretaceous of Morocco and Middle Jurassic of England. *Acta Palaeontologica Polonica*, 48, 301–319.
- Gasca, J. M., Canudo, J. I., & Moreno-Azanza, M. (2014). A large-bodied theropod (Tetanurae: Carcharodontosauria) from the Mirambel Formation (Barremian) of Spain. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 273, 13–23. doi: 10.1127/0077-7749/2014/0413.
- Gheerbrant, E., & Rage, J.-C. (2006). Paleobiogeography of Africa: How distinct from Gondwana and Laurasia? *Palaeogeography, Palaeoclimatology, Palaeoecology*, 241, 224–246. doi:10.1016/j.palaeo.2006.03.016.
- Gishlick, A. D., & Gauthier, J. A. (2007). On the manual morphology of *Compsognathus longipes* and its bearing on the diagnosis of Compsognathidae. *Zoological Journal of the Linnean Society*, 149, 569–581.
- Goloboff, P. A., Farris, J. S., & Nixon, K. C. (2008). TNT, a free program for phylogenetic analysis. *Cladistics*, 24, 774–786.
- Gradstein, F. M., Ogg, J. G., Schmitz, M. D., & Ogg, G. M. (Eds.) (2012). *The Geologic Time Scale 2012 (1144 p.)*. Amsterdam: Elsevier, 1144 pp.
- Grigorescu, D. (1992). Nonmarine Cretaceous Formations of Romania. In: Matter, N. J., Matter, & Chen, P.-J., Chen (Eds.), *Aspects of Nonmarine Cretaceous Geology (pp. 142–164)*. Beijing: China Ocean Press, Beijing, pp. 142–164.
- Grigorescu, D. (2003). Dinosaurs of Romania. *Comptes rendus-Rendus Palevol*, 2, 97–101.
- Grigorescu, D. (2010). The Latest Cretaceous fauna with dinosaurs and mammals from the Hațeg Basin — A historical overview. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 293, 271–282.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

1
2 1337 Hammer, O., Harper, D. A. T., & Ryan, P. D. (2001). Paleontological statistics software
3
4 1338 package for education and data analysis. *Palaeontologia Electronica*, 4, 1–9.
5
6 1339 Harris, J. D. (1998). A reanalysis of *Acrocanthosaurus atokensis*, its phylogenetic status,
7
8 1340 and paleobiogeographic implications, based on a new specimen from Texas. *New Mexico*
9
10 1341 *Museum of Natural History and Science Bulletin*, 13, 1–75.
11
12 1342 Hendrickx, C., & Mateus, O. (2014). Abelisauridae (Dinosauria: Theropoda) from the Late
13
14 1343 Jurassic of Portugal and dentition-based phylogeny as a contribution for the identification of
15
16 1344 isolated theropod teeth. *Zootaxa*, 3751(1), 1–74.
17
18 1345 Hendrickx, C., Mateus, O., & Araújo, R. (2015a). The dentition of megalosaurid theropods.
19
20 1346 *Acta Palaeontologica Polonica*, 60, 627–642. doi:10.4202/app.00056.2013.
21
22 1347 Hendrickx, C., Mateus, O., & Araújo, R. (2015b). A proposed terminology of theropod
23
24 1348 teeth (Dinosauria, Saurischia). *Journal of Vertebrate Paleontology*, 35(5), e982797. doi:
25
26 1349 10.1080/02724634.2015.982797.
27
28
29 1350 Holtz, T. R., Chapman, R. E., & Lamanna, M. C. (2004b). Mesozoic biogeography of
30
31 1351 Dinosauria. In D. B. Weishampel, P. Dodson, & H. Osmólska (Eds.), *The Dinosauria*.
32
33 1352 *Second Edition* (pp. 627–642). Berkeley, Los Angeles, London: University of California
34
35 1353 Press.
36
37 1354 Holtz, T. R., Jr., Molnar, R. E., & Currie, P. J. (2004a). Basal Tetanurae. In: D. B.
38
39 1355 Weishampel, P. Dodson, & H. Osmólska ~~Weishampel, D. B., Dodson, P., Osmólska, H.~~
40
41 1356 (Eds.), *The Dinosauria. Second Edition* (pp. 71–110). University of California Press,
42
43 1357 Berkeley, Los Angeles, London: University of California Press ~~pp. 71–110.~~
44
45 1358 Holtz, T. R., Chapman, R. E., Lamanna, M. C., 2004b. Mesozoic biogeography of Dinosauria.
46
47 1359 In: Weishampel, D. B., Dodson, P., Osmólska, H. (Eds.), *The Dinosauria. Second Edition*.
48
49 1360 University of California Press, Berkeley, Los Angeles, pp. 627–642.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

- Huene, F. von- (1923). Carnivorous Saurischia in Europe since the Triassic. *Bulletin of the Geological Society of America*, 34, 449–458.
- Hippolyte, J.-C. (2002). Geodynamics of Dobrogea (Romania): new constraints on the evolution of the Tornquist–Teisseyre Line, the Black Sea and the Carpathians. *Tectonophysics*, 357, 33–53.
- Ionesi, L. (1994). *Geologia unităților de platformă și a orogenului Nord-Dobrogean* (p. 280). Ed. Tehnică, Bucharest: Ed. Tehnică, 280 pp.
- Jerzykiewicz, T., & Russell, D. A. (1991). Late Mesozoic stratigraphy and vertebrates of the Gobi Basin. *Cretaceous Research*, 12, 345–377.
- Jurcsák, T. (1982). Occurrences nouvelles des Sauriens mésozoïques de Roumanie. *Vertebrata Hungarica*, 21, 175–184.
- Jurcsák, T., & Popa, E. (1979). Dinozaurieni ornithopozi din bauxitele de la Cornet (Munții Pădurea Craiului). *Nymphaea*, 7, 37–75.
- Jurcsák, T., & Popa, E. (1983). La faune de dinosauriens du Bihor (Roumanie). In: ~~Buffetaut, E.~~ *Buffetaut*, ~~Mazin, J. M.~~ *Mazin*, ~~Salmon, E.~~ *Salmon* (Eds.), *Actes du Symposium Paléontologique Georges Cuvier* (pp. 325–335). ~~Le Serpentaire~~, Montbéliard: *Le Serpentaire*, pp. 325–335.
- Kellner, A. W. A., Pinheiro, A. E. P., & Campos, D. A. (2014). A new sebecid from the Paleogene of Brazil and the crocodyliform radiation after the K–Pg boundary. *PLoS ONE*, 9(1), e81386. doi:10.1371/journal.pone.0081386.
- Kirkland, J. I., Cifelli, R. L., Britt, B. B., Burge, D. L., DeCourten, F. L., Eaton, J. G., & Parrish, J. M. (1999). Distribution of vertebrate faunas in the Cedar Mountain Formation, east-central Utah. *Utah Geological Survey Miscellaneous Publication*, 99-1, 201–217.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

- Knoll, F., & Ruiz-Omeñaca, J. I. (2009). Theropod teeth from the basalmost Cretaceous of Anoual (Morocco) and their palaeobiogeographical significance. *Geological Magazine*, 146, 602–616.
- Kusuhashi, N., Matsumoto, A., Murakami, M., Tagami, T., Hirata, T., Iizuka, T., Handa, T., & Matsuoka, H. (2006). Zircon U–Pb ages from tuff beds of the upper Mesozoic Tetori Group in the Shokawa district, Gifu Prefecture, central Japan. *The Island Arc*, 15, 378–390.
- Lacasa Ruiz, A. (1989). Nuevo genero de ave fosil del yacimiento Neocomiense del Montsec (Provincia de Lerida, España). *Estudios geológicos*, 45(5-6), 417–425.
- Larson, D. W., & Currie, P. J. (2013). Multivariate analyses of small theropod dinosaur teeth and implications for paleoecological turnover through time. *PLoS ONE*, 8(1), e54329. doi:10.1371/journal.pone.0054329.
- Le Loeuff, J., & Buffetaut, E. (1995). The evolution of Late Cretaceous non-marine vertebrate faunas in Europe. In: Sun, A.-L. Sun, & Wang, Y.-Q. Wang (Eds.), *Sixth Symposium on Mesozoic Terrestrial Ecosystems and Biota, Short Papers* (pp. 181–184). China Ocean Press, Beijing: China Ocean Press pp. 181–184.
- Le Loeuff, J., Lang, E., Cavin, L., & Buffetaut, E. (2012). Between Tendaguru and Bahariya: on the age of the Early Cretaceous dinosaur sites from the Continental Intercalaire and other African formations. *Journal of Stratigraphy*, 36, 486–502.
- Lucas, F. A. (1901). A new dinosaur, *Stegosaurus marshi*, from the Lower Cretaceous of South Dakota. *Proceedings of the United States National Museum*, 23(1224), 591–592.
- Lucas, S. G. (2006). The *Psittacosaurus* biochron, Early Cretaceous of Asia. *Cretaceous Research*, 27, 189–198.
- Lucas, S. G., & Estep, J. W. (1998). Vertebrate biostratigraphy and biochronology of the Cretaceous of China. In Lucas, S. G. Lucas, Kirkland, J. I. Kirkland, Estep, J. W. Estep

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

- 1
2 1408 (Eds.), *Lower and Middle Cretaceous Terrestrial Ecosystems* (pp. 1–20). New Mexico
3
4 1409 Museum of Natural History and Science Bulletin, 14, pp. 1–20.
5
6 1410 Lü, J.-C., Xu, L., Pu, H.-Y., Jia, S.-H., Azuma, Y., Chang, H.-L., & Zhang, J.-M.,
7
8 1411 (2014, 2016). Paleogeographical significance of carcharodontosaurid teeth from the late Early
9
10 1412 Cretaceous of Ruyang, Henan Province of central China. *Historical Biology*, 28, 8–13. doi:
11
12 1413 10.1080/08912963.2014.947287.
13
14 1414 Manabe, M., & Hasegawa, Y., (1995). Diapsid fauna and its paleobiogeographical
15
16 1415 implication, the Neocomian section of the Tetori Group. In: Sun, A., Sun, & Wang, Y., Wang
17
18 1416 (Eds.), *Sixth Symposium on Mesozoic Terrestrial Ecosystems and Biota, Short Papers*, (pp.
19
20
21 1417 179). Beijing: China Ocean Press, pp. 179.
22
23 1418 Mannion, P. D., (2009). A rebbachisaurid sauropod from the Lower Cretaceous of the Isle of
24
25 1419 Wight, England. *Cretaceous Research*, 30, 521–526.
26
27 1420 Mannion, P. D., Upchurch, P., & Hutt, S., (2011). New rebbachisaurid (Dinosauria:
28
29 1421 Sauropoda) material from the Wessex Formation (Barremian, Early Cretaceous), Isle of
30
31 1422 Wight, United Kingdom. *Cretaceous Research*, 32, 774–780.
32
33 1423 Matsukawa, M., Ito, M., Nishida, N., Koarai, K., Lockley, M. G., Nichols, D. J., (2006). The
34
35 1424 Cretaceous Tetori biota in Japan and its evolutionary significance for terrestrial ecosystems in
36
37 1425 Asia. *Cretaceous Research*, 27, 199–225.
38
39 1426 McDonald, A. T., (2011). The taxonomy of species assigned to *Camptosaurus* (Dinosauria:
40
41 1427 Ornithopoda). *Zootaxa*, 2783, 52–68.
42
43 1428 McKenna, M. C., (1973). Sweepstakes, filters, corridors, Noah's Arks, and Beached Viking
44
45 1429 Funeral Ships in palaeogeography. In: Tarling, D. H., Tarling, Runcorn, S. K., Runcorn (Eds.),
46
47 1430 *Implications of Continental Drift to the Earth Sciences* (pp. 295–308). Academic Press, New
48
49 1431 York, Academic Press, pp. 295–308.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Mo, J.-Y., Huang, C.-L., Xie, S.-W., & Buffetaut, E. (2014). A megatheropod tooth from the Early Cretaceous of Fusui, Guangxi, Southern China. *Acta Geologica Sinica (English Edition)*, 88, 6–12.

Formatted: Font: Italic

Formatted: Font: Italic

Molnar, R. E. (1990). Problematic Theropoda: "Carnosaurs". In: Weishampel, D. B., Weishampel, D. B., Dodson, P., Dodson, P., & Osmólska, H. Osmólska (Eds.), *The Dinosauria* (pp. 306–317). University of California Press, Berkeley, Los Angeles, Oxford; University of California Press, 306–317.

Formatted: Font: Italic

Mutihac, V., & Mutihac, G. (2010). *The geology of Romania, within the Central East European geostructural context* (p. 690). Ed. Didactică și Pedagogică, Bucharest; Ed. Didactică și Pedagogică, 690 pp.

Formatted: Font: Italic

Neagu, T., Dragastan, O., & Csiki, Z. (1997). Early Cretaceous shelf paleocommunities of Cernavodă (South Dobrogea, SE Romania). *Acta Palaeontologica Romaniae*, 1, 28–36.

Formatted: Font: Italic

Formatted: Font: Italic

Nopcsa, F. (1902). Notizen über cretacische Dinosaurier. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe*, 111, 93–114.

Formatted: Font: Italic

Nopcsa, F. (1923). On the geological importance of the primitive reptilian fauna of the uppermost Cretaceous of Hungary; with a description of a new tortoise (*Kallokibotium*).

Quarterly Journal of the Geological Society of London, 79, 100–116.

Formatted: Font: Italic

Formatted: Font: Italic

Norman, D. B. (2010). A taxonomy of iguanodontians (Dinosauria: Ornithopoda) from the lower Wealden Group (Cretaceous: Valanginian) of southern England. *Zootaxa*, 2489, 47–66.

Formatted: Font: Italic

Formatted: Font: Italic

Norman, D. B. (2013). On the taxonomy and diversity of Wealden iguanodontian dinosaurs (Ornithischia: Ornithopoda). *Revue de Paléobiologie*, 32(2), 385–404.

Formatted: Font: Italic

Formatted: Font: Italic

Novas, F. E., Agnolín, F. L., Ezcurra, M. D., Porfiri, J., & Canale, J. I. (2013). Evolution of the carnivorous dinosaurs during the Cretaceous: The evidence from Patagonia. *Cretaceous Research*, 45, 174–215.

Formatted: Font: Italic

Formatted: Font: Italic

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Novas, F. E., de Valais, S., Vickers-Rich, P. A., & Rich, T. H. (2005). A large Cretaceous theropod from Patagonia, Argentina, and the evolution of carcharodontosaurids. *Naturwissenschaften*, 92, 226–230.
- ~~Novas, F. E., Agnolín, F. L., Ezcurra, M. D., Porfiri, J., Canale, J. I., 2013. Evolution of the~~
~~carnivorous dinosaurs during the Cretaceous: The evidence from Patagonia. Cretaceous~~
~~Research 45, 174–215.~~
- Ortega, F., Escaso, F., & Sanz, J. L. (2010). A bizarre, humped Carcharodontosauria (Theropoda) from the Lower Cretaceous of Spain. *Nature*, 467, 203–206.
- ~~Ősi, A., Codrea, V., Prondvai, E., & Csiki-Sava, Z. (2014). New ankylosaurian material from~~
~~the Upper Cretaceous of Transylvania. Annales de Paléontologie, 100, 257–271. doi:~~
~~10.1016/j.annpal.2014.02.001.~~
- Ősi, A., Rabi, M., Makádi, L., Szentesi, Z., Botfalvai, G., & Gulyás, P. (2012). The Late Cretaceous continental vertebrate fauna from Iharkút (western Hungary): a review. In: ~~Godefroit, P. Godefroit~~ (Ed.), *Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems* (pp. 533–569). ~~Indiana University Press~~, Bloomington, ~~Indiana University Press~~ pp. 533–569.
- ~~Ősi, A., Codrea, V., Prondvai, E., Csiki-Sava, Z., 2014. New ankylosaurian material from the~~
~~Upper Cretaceous of Transylvania. Annales de Paléontologie 100, 257–271. doi:~~
~~10.1016/j.annpal.2014.02.001.~~
- Ősi, A., Rabi, M., & Makádi, L. (2015). An enigmatic crocodyliform tooth from the bauxites of western Hungary suggests hidden mesoeucrocodylian diversity in the Early Cretaceous European archipelago. *PeerJ*, 3, e1160. doi:10.7717/peerj.1160.
- Paolillo, A., & Linares, O. J. (2007). Nuevos cocodrilos Sebecosuchia del Cenoicoico Suramericano (Mesosuchia: Crocodylia). *Paleobiologia Neotropical*, 3, 1–25.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Pereda-Suberbiola, X., Ruiz-Omeñaca, J. I., Canudo, J. I., Torcida, F., & Sanz, J. L. (2012).

Dinosaur faunas from the Early Cretaceous (Valanginian–Albian) of Spain. In P. Godefroit

(Ed.), *Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems* (pp. 378–407).

Formatted: Font: Italic

Bloomington: Indiana University Press.

Pereda-Suberbiola, X., Ruiz-Omeñaca, J. I., Fernandez-Baldor, F. T., Maisch, M. W.,

Huerta, P., Contreras, R., Izquierdo, L. A., Huerta, D. M., Montero, V. U., & Welle, J.

(2011). A tall-spined ornithopod dinosaur from the Early Cretaceous of Salas de los Infantes

(Burgos, Spain). *Comptes Rendus Palevol*, 10, 551–558.

Formatted: Font: Italic

Formatted: Font: Italic

~~Pereda-Suberbiola, X., Ruiz-Omeñaca, J. I., Canudo, J. I., Torcida, F., Sanz, J. L., 2012.~~

~~Dinosaur faunas from the Early Cretaceous (Valanginian–Albian) of Spain. In: Godefroit, P.~~

~~(Ed.), Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems. Life of the Past.~~

~~Indiana University Press, Bloomington, pp. 378–407.~~

Pérez-Moreno, B. P., Sanz, J. L., Sudre, J., & Sigé, B. (1993). A theropod dinosaur from

the Lower Cretaceous of southern France. *Revue de Paléobiologie, Volume spéciale* 7, 173–

188.

Formatted: Font: Italic

Formatted: Font: Italic

Pol, D., & Powell, J. E. (2011). A new sebecid mesoeucrocodylian from the Rio Loro

Formation (Palaeocene) of north-western Argentina. *Zoological Journal of the Linnean*

Society, 163, S7–S36.

Formatted: Font: Italic

Formatted: Font: Italic

Porfiri, J. D., Novas, F. E., Calvo, J. O., Agnolín, F. L., Ezcurra, M. D., & Cerda, I. A.

(2014). Juvenile specimen of *Megaraptor* (Dinosauria, Theropoda) sheds light about

tyrannosauroid radiation. *Cretaceous Research*, 51, 35–55. doi:10.1016/j.cretres.2014.04.007.

Formatted: Font: Italic

Formatted: Font: Italic

Posmoşanu, E. (2003). Iguanodontian dinosaurs from the Lower Cretaceous bauxite site

from Romania. *Acta Palaeontologica Romaniae*, 4, 431–439.

Formatted: Font: Italic

Formatted: Font: Italic

1
2 1503 Porfiri, J.D., Novas, F.E., Calvo, J.O., Agnolín, F.L., Ezcurra, M.D., Cerda, I.A., 2014.
3
4 1504 Juvenile specimen of *Megaraptor* (Dinosauria, Theropoda) sheds light about tyrannosauroid
5
6 1505 radiation. *Cretaceous Research*, 51, 35–55. doi:10.1016/j.cretres.2014.04.007.
7
8 1506 Rabi, M., & Sebök, N., (2015). A revised Eurogondwana model: Late Cretaceous
9
10 1507 notosuchian crocodyliforms and other vertebrate taxa suggest the retention of episodic faunal
11
12 1508 links between Europe and Gondwana during most of the Cretaceous. *Gondwana Research*,
13
14 1509 28, 1197–1211. doi:10.1016/j.gr.2014.09.015.
15
16 1510 Racey, A., (2009). Mesozoic red bed sequences from SE Asia and the significance of the
17
18 1511 Khorat Group of NE Thailand. In: Buffetaut, E. Buffetaut, Cuny, G. Cuny, Le Loeuff, J. Le
19
20 1512 Loeuff, & Suteethorn, V. Suteethorn (Eds.), *Late Palaeozoic and Mesozoic Ecosystems in SE*
21
22 1513 *Asia* (pp. 41–67). The Geological Society, London, Special Publications, 315, pp. 41–67. doi:
23
24 1514 10.1144/SP315.5.
25
26 1515 Racey, A., & Goodall, J. G. S., (2009). Palynology and stratigraphy of the Mesozoic Khorat
27
28 1516 Group red bed sequences from Thailand. In: Buffetaut, E. Buffetaut, Cuny, G. Cuny, Le
29
30 1517 Loeuff, J. Le Loeuff, & Suteethorn, V. Suteethorn (Eds.), *Late Palaeozoic and Mesozoic*
31
32 1518 *Ecosystems in SE Asia* (pp. 69–83). The Geological Society of, London, Special Publications,
33
34 1519 315, pp. 69–83. doi: 10.1144/SP315.6.
35
36 1520 Rauhut, O. W. M., (2011). Theropod dinosaurs from the Late Jurassic of Tendaguru
37
38 1521 (Tanzania). *Special Papers in Palaeontology*, 86, 195–239.
39
40 1522 Rauhut, O. W. M., & Werner, C., (1995). First record of the family Dromaeosauridae
41
42 1523 (Dinosauria: Theropoda) in the Cretaceous of Gondwana (Wadi Milk Formation, northern
43
44 1524 Sudan. *Paläontologische Zeitschrift*, 69(3/4), 475–489.
45
46 1525 Richter, U., Mudroch, A., & Buckley, L. G., (2013). Isolated theropod teeth from the Kem
47
48 1526 Kem Beds (Early Cenomanian) near Taouz, Morocco. *Paläontologische Zeitschrift*, 87, 291–
49
50 1527 309.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: Left, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

1
2 1528 Royo-Torres, R., Cobos, A., Luque, L., Aberasturi, A., Espilez, E., Fierro, I., Gonzales, A.,
3
4 1529 Mampel, L., & Alcalá, L. (2009). High European sauropod dinosaur diversity during
5
6 1530 Jurassic–Cretaceous transition in Riodeva (Teruel, Spain). *Palaeontology*, 52, 1009–1027.
7
8 1531 Sachs, S., & Hornung, J. J. (2013). Ankylosaur remains from the Early Cretaceous
9
10 1532 (Valanginian) of Northwestern Germany. *PLoS ONE*, 8(4), e60571.
11
12 1533 doi:10.1371/journal.pone.0060571.
13
14 1534 Salgado, L., Canudo, J. I., Garrido, A. C., Ruiz-Omeñaca, J. I., Garcia, R. A., de la Fuente,
15
16 1535 M. S., Barco, J. L., & Bollati, R. (2009). Upper Cretaceous vertebrates from El Anfiteatro
17
18 1536 area, Río Negro, Patagonia, Argentina. *Cretaceous Research*, 30, 767–784.
19
20 1537 Sames, B., Cifelli, R. L., & Schudack, M. E. (2010). The nonmarine Lower Cretaceous of
21
22 1538 the North American Western Interior foreland basin: New biostratigraphic results from
23
24 1539 ostracod correlations and early mammals, and their implications for paleontology and
25
26 1540 geology of the basin—An overview. *Earth-Science Reviews*, 101, 207–224.
27
28 1541 Sauvage, H. E. (1876). Notes sur les reptiles fossiles. *Bulletin de la Société Géologique de*
29
30 1542 *France*, 4, 435–442.
31
32 1543 Sauvage, H. E. (1882). Recherches sur les reptiles trouves dans le Gault de l'est du bassin
33
34 1544 de Paris. *Mémoires de la Société Géologique de France*, 2, 1–42.
35
36 1545 Săndulescu, M. (1984). *Geotectonica României* (p. 329). Ed. Tehnică, Bucharest, 329 pp.
37
38 1546 *Ed. Tehnică*.
39
40 1547 Seghedi, A. (2001). The North Dobrogea orogenic belt (Romania): a review. In: Ziegler, P.
41
42 1548 A. Ziegler, Cavazza, W. Cavazza, Robertson, A. H. F. Robertson, & Crasquin-Soleau, S.
43
44 1549 Crasquin-Soleau (Eds.), *Peri-Tethys Memoir 6: Peri-Tethyan Rift/Wrench Basins and Passive*
45
46 1550 *Margins* (pp. 237–257). Paris: Mémoires de la Musée National d'Histoire Naturelle, Paris, pp.
47
48 1551 237–257.
49
50 1552 Sereno, P. C. (1999). Dinosaurian biogeography: vicariance, dispersal and regional
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

1
2 extinction. In: ~~Tomida~~, Y. ~~Tomida~~, ~~Rich~~, T. H. ~~Rich~~, & ~~Vickers-Rich~~, P. ~~Vickers-Rich~~ (Eds.),
3
4 *Proceedings of the Second Gondwanan Dinosaur Symposium* (pp. 249–257). Tokyo: National
5
6 Science Museum Monographs, 15; ~~Tokyo~~, pp. 249–257.
7
8 Sereno, P. C., & Brusatte, S. L., (2008). Basal abelisaurid and carcharodontosaurid
9
10 theropods from the Lower Cretaceous Elrhaz Formation of Niger. *Acta Palaeontologica*
11
12 *Polonica*, 53, 15–46.
13
14 Sereno, P. C., Dutheil, D. B., Iarochene, M., Larsson, H. C. E., Lyon, G. H., Magwene, P. M.,
15
16 Sidor, C. A., Varricchio, D. J., & Wilson, J. A., (1996). Predatory dinosaurs from the Sahara
17
18 and Late Cretaceous faunal differentiation. *Science*, 272, 986–991.
19
20 Shen, Y. B., & Mateer, N. J., (1992). An outline of the Cretaceous system in northern
21
22 Xinjiang, western China. In: ~~Mateer~~, N. J. ~~Mateer~~, & ~~Chen~~, P. J. ~~Chen~~ (Eds.), *Aspects of*
23
24 *Nonmarine Cretaceous Geology* (pp. 49–77). ~~China Ocean Press~~, Beijing: ~~China Ocean~~
25
26 ~~Press~~, pp. 49–77.
27
28 Simionescu, I., (1906). Note sur l'age des calcaires de Cernavoda (Dobrogea). *Annales*
29
30 *Scientifiques de l'Université de Jassy*, 4(1), 1–3.
31
32 Simionescu, I., (1913). *Megalosaurus* aus der Unterkreide der Dobrogea. *Centralblatt für*
33
34 *Mineralogie, Geologie und Paläontologie*, 1913(20), 686–687.
35
36 Sissingh, W., (1977). Biostratigraphy of Cretaceous calcareous nannoplankton. *Geologie en*
37
38 *Mijnbouw*, 56, 37–65.
39
40 Smith, J. B., Vann, D. R., & Dodson, P., (2005). Dental morphology and variation in
41
42 theropod dinosaurs: implications for the taxonomic identification of isolated teeth. *The*
43
44 *Anatomical Record A*, 285A, 699–736.
45
46 Stein, K., Csiki, Z., Curry Rogers, K., Weishampel, D. B., Redelstorff, R., Carballido, J. L.,
47
48 & Sander, P. M., (2010). Small body size and extreme cortical bone remodeling indicate
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1577 phyletic dwarfism in *Magyarosaurus dacus* (Sauropoda: Titanosauria). *Proceedings of the*
1578 *National Academy of Sciences*, 107, 9258–9263.
1579 Stoica, M., & Csiki, Z. (2002). An earliest Cretaceous (Purbeckian) vertebrate fauna from
1580 Southern Dobrogea (southeastern Romania). In: Grigorescu, D. Grigorescu, & Csiki, Z. Csiki
1581 (Eds.), *7th European Workshop on Vertebrate Palaeontology, Sibiu, Romania*, (pp. 34).
1582 *Bucharest: Ars Docendi*.
1583 Stovall, J. W., & Langston, W., Jr. (1950). *Acrocanthosaurus atokensis*, a new genus and
1584 species of Lower Cretaceous Theropoda from Oklahoma. *American Midland Naturalist*, 43,
1585 686–728.
1586 Stromer, E. (1931). Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wüsten
1587 Ägyptens. II. Wirbeltier-Reste der Baharijstufe (unterstes Cenoman). 10. Ein Skelett-Rest
1588 von *Carcharodontosaurus* nov. gen. *Abhandlungen der Bayerischen Akademie der*
1589 *Wissenschaften, Mathematisch-naturwissenschaftliche Abteilung, Neue Folge* 9, 1–23.
1590 Therrien, F., & Henderson, D. M. (2007). My theropod is bigger than yours ... or not:
1591 estimating body size from skull length in theropods. *Journal of Vertebrate Paleontology*, 27,
1592 108–115.
1593 Tong, H., Claude, J., Suteethorn, V., Naksri, W., & Buffetaut, E. (2009). Turtle
1594 assemblages of the Khorat Group (Late Jurassic–Early Cretaceous) of NE Thailand and their
1595 palaeobiogeographical significance. In Buffetaut, E. Buffetaut, Cuny, G. Cuny, Le Loeuff, J.
1596 Le Loeuff, & Suteethorn, V. Suteethorn (Eds.), *Late Palaeozoic and Mesozoic Ecosystems in*
1597 *SE Asia* (pp. 141–151). The Geological Society, London, Special Publications, 315, pp. 141–
1598 151.
1599 Torcida Fernández-Baldor, F., Canudo, J. I., Huerta, P., Montero, D., Pereda Suberbiola, X.,
1600 & Salgado, L. (2011). *Demandasaurus darwini*, a new rebbachisaurid sauropod from the
1601 Early Cretaceous of the Iberian Peninsula. *Acta Palaeontologica Polonica*, 56, 535–552.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Turculeț, I., & Brânzilă, M. (2012). *Muzeul colecțiilor paleontologice originale de la Universitatea "Alexandru Ioan Cuza" Iași* (p. 173). Editura Universității "Alexandru Ioan Cuza" Iași, Iași; Editura Universității "Alexandru Ioan Cuza" Iași 173 pp.

Formatted: Font: Italic

Turner, A. H., Makovicky, P. J., & Norell, M. A. (2012). A review of dromaeosaurid systematics and paravian phylogeny. *Bulletin of the American Museum of Natural History*, 371, 1–206.

Formatted: Font: Italic

Formatted: Font: Italic

Turner, A. H., & Sertich, J. J. W. (2010). Phylogenetic history of *Simosuchus clarki* (Crocodyliformes: Notosuchia) from the Late Cretaceous of Madagascar. *Journal of Vertebrate Paleontology*, 30(Supplement 1), 177–236.

Formatted: Font: Italic

Formatted: Font: Italic

Turner, A.H., Makovicky, P.J., Norell, M.A., 2012. A review of dromaeosaurid systematics and paravian phylogeny. *Bulletin of the American Museum of Natural History* 371, 1–206.

Vasile, Ș., & Csiki, Z. (2011). New Maastrichtian microvertebrates from the Rusca Montană Basin (Romania). *Oltenia. Studii și comunicări. Științele Naturii*, 27(1), 221–230.

Formatted: Font: Italic

Formatted: Font: Italic

Vickers-Rich, P., Rich, T. H., Lanus, D. R., Rich, L. S. V., & Vacca, R. (1999). "Big Tooth" from the Early Cretaceous of Chubut Province, Patagonia: a possible

carcharodontosaurid. In: Tomida, Y., Tomida, Rieh, T. H., Rich, Vickers-Rich, & P. Vickers-Rich (Eds.), *Proceedings of the Second Gondwanan Dinosaur Symposium* (pp. 85–88). Tokyo: National Science Museum Monographs, 15, Tokyo, pp. 85–88.

Formatted: Font: Italic

Vremir, M. (2010). New faunal elements from the Late Cretaceous (Maastrichtian) continental deposits of Sebeș area (Transylvania). *Terra Sebus. Acta Musei Sabesiensis*, 2, 635–684.

Formatted: Font: Italic

Formatted: Font: Italic

Vullo, R., Néraudeau, D., & Lenglet, T. (2007). Dinosaur teeth from the Cenomanian of Charentes, western France: evidence for a mixed Laurasian-Gondwanan assemblage. *Journal of Vertebrate Paleontology*, 27, 931–943.

Formatted: Font: Italic

Formatted: Font: Italic

1
2 1626 Weishampel, D. B., (1990). Dinosaurian distribution. In: ~~Weishampel, D. B.~~ Weishampel,
3
4 1627 ~~Dodson, P. Dodson, Osmólska, & H. Osmólska~~ (Eds.), *The Dinosauria* (pp. 63–140).
5
6 1628 ~~California University Press, Berkeley, Los Angeles, Oxford~~ Berkeley, California University
7
8 1629 Press pp. 63–140.
9
10 1630 Weishampel, D. B., Barrett, P. M., Coria, R. A., Le Loeuff, J., Xu, X., Zhao, X-J., Sahni, A.,
11
12 1631 Gomani, E. M., & Noto, C. R. (2004). Dinosaur distribution. In D. B. Weishampel, P.
13
14 1632 Dodson, & H. Osmólska (Eds.), *The Dinosauria. Second Edition* (pp. 517–606). Berkeley,
15
16 1633 Los Angeles, London: University of California Press.
17
18 1634 Weishampel, D. B., Csiki, Z., Benton, M. J., Grigorescu, D., & Codrea, V. (2010).
19
20 Palaeobiogeographic relationships of the Hateg biota — Between isolation and innovation.
21 1635
22 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 293, 419–437.
23 1636
24
25 1637 Weishampel, D. B., Grigorescu, D., & Norman, D. B., (1991). The Dinosaurs of
26
27 1638 Transylvania. *National Geographic Research & Exploration*, 7(2), 196–215.
28
29 1639 Weishampel, D. B., Jianu, C. M., Csiki, Z., & Norman, D. B. (2003). Osteology and
30
31 1640 phylogeny of *Zalmoxes* (n. g.), an unusual euornithopod dinosaur from the latest Cretaceous
32
33 1641 of Romania. *Journal of Systematic Palaeontology*, 1, 65–123.
34
35 1642 Weishampel, D. B., Norman, D. B., & Grigorescu, D., (1993). *Telmatosaurus*
36
37 1643 *transsylvanicus* from the Late Cretaceous of Romania: the most basal hadrosaurid dinosaur.
38
39 1644 *Palaeontology*, 36, 361–385.
40
41 1645 ~~Weishampel, D. B., Jianu, C. M., Csiki, Z., Norman, D. B., 2003. Osteology and phylogeny of~~
42
43 1646 ~~*Zalmoxes* (n. g.), an unusual euornithopod dinosaur from the latest Cretaceous of Romania.~~
44
45 1647 ~~*Journal of Systematic Palaeontology* 1, 65–123.~~
46
47 1648 ~~Weishampel, D. B., Barrett, P. M., Coria, R. A., Le Loeuff, J., Xu, X., Zhao, X-J., Sahni, A.,~~
48
49 1649 ~~Gomani, E. M., Noto, C. R., 2004. Dinosaur distribution. In: Weishampel, D. B., Dodson, P.,~~
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

~~Osmólska, H. (Eds.), The Dinosauria. Second Edition. University of California Press,~~
~~Berkeley, Los Angeles, London, pp. 517–606.~~

~~Weishampel, D.B., Csiki, Z., Benton, M.J., Grigorescu, D., Codrea, V., 2010.~~
~~Palaeobiogeographic relationships of the Hateg biota—Between isolation and innovation.~~
~~Palaeogeography, Palaeoclimatology, Palaeoecology 293, 419–437.~~

Williamson, T. E., & Brusatte, S. L., (2014). Small theropod teeth from the Late Cretaceous
of the San Juan Basin, northwestern New Mexico and their implications for understanding
latest Cretaceous dinosaur evolution. *PLoS ONE*, 9(4), e93190.
doi:10.1371/journal.pone.0093190.

Zarcone, G., Cillari, F. M. P., Stefano, P. D., Guzzetta, D., & Nicosia, U., (2010). A
possible bridge between Adria and Africa: New palaeobiogeographic and stratigraphic
constraints on the Mesozoic palaeogeography of the Central Mediterranean area. *Earth-*
Science Reviews, 103, 154–162.

Figure captions

Figure 1. Simplified geological map of the Cernavodă-Cochirleni area; inset shows the
position of the study area within Romania. Legend: 1. Quaternary: a. Holocene alluvia, b.
Pleistocene–Holocene loessoid deposits; 2. Pre-Quaternary Cenozoic (Middle Eocene and
Miocene) deposits; Cretaceous: 3. Peștera Formation, Lower Cenomanian; 4. Cochirleni
Formation; uppermost Aptian–Lower Albian; 5. Gherghina Formation, Middle–Upper
Aptian; 6. Ostrov (= Ramadan) Formation; Barremian–Lower Aptian; 7. Cernavodă
Formation, Alimanu Member, Berriasian–Valanginian; 8. Water courses. (Redrawn after
Dragastan et al., 1998, 2014).

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Figure 2. Specimen UAIC (SCM1) 615, indeterminate carcharodontosaurid lateral tooth from Cochirleni, Southern Dobrogea. A. UAIC (SCM1) 615, as figured by Simionescu (1913); B. Current state of UAIC (SCM1) 615, mounted in a limestone holder.

Figure 3. Detailed morphology of UAIC (SCM1) 615, an indeterminate carcharodontosaurid lateral tooth from Cochirleni, Southern Dobrogea. UAIC (SCM1) 615 in A. labial? side; B., distal; C., lingual? side, and D., basal (mesial to the right) views. Details of the distal carina (marked with boxes in A, respectively C): apical part in E., labial? and F. distal views; basal part in G., lingual? and H., distal views. Scale bar: 1 cm (A–D), 5 mm (E–H).

Figure 4. Dental morphospace of the different theropod clades according to the results of the PCA analysis; UAIC (SCM1) 615 (red star) plots within the morphospace occupied by Carcharodontosauridae. See further details of this analysis, as well as other quantitative analyses used to identify the tooth that deliver similar results (cluster analysis, discriminant function analysis, phylogenetic analysis), in the Supplementary Material.

Figure 5. A. Palaeogeographic setting of the two early Early Cretaceous Romanian dinosaur occurrences: the Berriasian–Valanginian Cornet locality (orange star), located on a Neo-Tethyan archipelago island, and the Valanginian Cochirleni locality (red star), situated on the marginal areas of the Eastern European cratonic mainland. B. Global chronostratigraphic and palaeobiogeographic distribution of the Carcharodontosauridae, plotted on Middle Aptian (approx. 120 Mya) palaeogeographic map; red star marks the position of UAIC (SCM1) 615 from Southern Dobrogea. Legend: 1 – *Veterupristisaurus*, ‘*Megalosaurus*’ *ingens*, Carcharodontosauridae indet., Tanzania, Late Jurassic; 2 – *Concavenator*, Spain, Barremian; 3 – Carcharodontosauridae indet., Thailand, Barremian; 4 – *Acrocanthosaurus*, southeastern

1
 21700 United States, Aptian–Albian; 5 – Carcharodontosauridae indet., Spain, Aptian; 6 –
 3
 41701 *Eocarcharia*, Niger, Aptian–Albian; 7 – Carcharodontosauridae indet., Guangxi, China,
 5
 61702 Aptian; 8 – Carcharodontosauridae indet., Henan, China, Aptian; 9 – *Kelmayisaurus*,
 7
 81703 Xinjiang, China, Aptian–Albian; 10 – Carcharodontosauridae indet., France, Cenomanian; 11
 9
 101704 – *Sauroniops*, Morocco, Cenomanian; 12 – Carcharodontosauridae indet., Japan,
 11
 121705 Cenomanian–early Turonian; 13 – *Shaochilong*, Inner Mongolia, China, Turonian; 14 –
 13
 141706 Carcharodontosauridae indet., São Paulo, Brazil, Campanian–Maastrichtian (for relevant
 15
 161707 references, see text, 5.4.). Palaeogeographic maps, courtesy of Ron Blakey
 17
 181708 (<http://cpgeosystems.com/>).
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65

“*Megalosaurus cf. superbus*” from southeastern Romania: the oldest known Cretaceous
carcharodontosaurid (Dinosauria: Theropoda) and its implications for earliest Cretaceous
Europe-Gondwana connections

Zoltán Csiki-Sava^{1*}, Stephen L. Brusatte², Ștefan Vasile¹

¹ Department of Geology, Faculty of Geology and Geophysics, University of Bucharest, 1
Nicolae Bălcescu Boulevard, 010041 Bucharest, Romania

² School of GeoSciences, University of Edinburgh, Grant Institute, James Hutton Road,
Edinburgh, EH9 3FE, United Kingdom

* Corresponding author
zoltan.csiki@g.unibuc.ro

ABSTRACT

Some of the best records of continental vertebrates from the Cretaceous of Europe come from
Romania, particularly two well-known occurrences of dwarfed and morphologically aberrant
dinosaurs and other taxa that lived on islands (the Cornet and Hațeg Island faunas).
Substantially less is known about those vertebrates living in the more stable, cratonic regions
of Romania (and Eastern Europe as a whole), particularly during the earliest Cretaceous. We
describe one of the few early Early Cretaceous fossils that have ever been found from these
regions, the tooth of a large theropod dinosaur from Southern Dobrogea, which was
discovered over a century ago but whose age and identification have been controversial. We
identify the specimen as coming from the Valanginian stage of the Early Cretaceous, an
incredibly poorly sampled interval in global dinosaur evolution, and as belonging to
Carcharodontosauridae, a clade of derived, large-bodied apex predators whose earliest

Cretaceous history is poorly known. Quantitative analyses demonstrate that the Romanian tooth shows affinities with a derived carcharodontosaurid subgroup, the Carcharodontosaurinae, which until now has been known solely from Gondwana. Our results suggest that this subgroup of colossal predators did not evolved vicariantly as Laurasia split from Gondwana, but originated earlier, perhaps in Europe. The carcharodontosaurine diversification may have been tied to a north-to-south trans-Tethyan dispersal that took place sometime between the Valanginian and Aptian, illustrating the importance of palaeogeographic ties between these two realms during the largely mysterious early–mid Early Cretaceous.

Keywords

Southern Dobrogea; Valanginian; Carcharodontosauridae; cratonic Europe; palaeobiogeography

1. Introduction

Romania boasts one of the best records of continental vertebrate fossils from the Cretaceous of Europe (e.g., Grigorescu, 1992, 2003; Csiki-Sava et al., 2015). The vast majority of fossils come from two well-known occurrences: the Early Cretaceous bauxite accumulations of Cornet, in the northern Apuseni Mountains (e.g., Jurcsák, 1982; Benton et al., 1997; Posmoşanu, 2003; Dyke et al., 2011), and the famous latest Cretaceous beds of the Haţeg, Rusca Montană and western Transylvanian basins of Transylvania, which have yielded the dinosaur-dominated ‘Haţeg Island fauna’ (e.g, Nopcsa, 1923; Weishampel et al., 1991; Benton et al. 2010; Codrea et al., 2010, 2012; Grigorescu, 2010; Vremir, 2010; Vasile and Csiki, 2011; Csiki-Sava et al., 2015). Both of these faunas inhabited islands that were part of the vast Cretaceous European Archipelago of the Neo-Tethys Ocean. Based on their isolated

geological settings and the many dwarfed and morphologically aberrant taxa that make up the faunas, both have been interpreted as insular assemblages that give a unique window into how island environments affected the evolution of long-extinct organisms (e.g., Benton et al., 1997, 2010; Csiki-Sava et al., 2015).

The great volume of research on these assemblages over the past century, particularly the ‘Hațeg Island fauna’, has concealed an inconvenient bias: the stable, non-island, cratonic regions of Romania have yielded only extremely rare Mesozoic continental vertebrate remains (i.e., the Moldavian, Moesian and Scythian platforms; Săndulescu, 1984; Mutihac and Mutihac, 2010; Fig. 1). This is mostly because Mesozoic deposits are located in the subsurface in these regions, with only limited subaerial exposures available in the structurally highest-lying parts of the Moesian Platform, in Central and Southern Dobrogea (Middle Jurassic–Upper Cretaceous), as well as in the northeastern-most corner of the Moldavian Platform, along the Prut Valley (lower Upper Cretaceous) (see, e.g., Mutihac and Mutihac, 2010). This bias is unfortunate because fossils from these settings could lead to a better understanding of how mainland and island faunas differed during the Cretaceous, and because the cratonic portion of Europe was an important biogeographic stepping stone between the north and south as the continents fragmented and sea levels fluctuated.

Although the cratonic regions of Romania have yielded few Cretaceous terrestrial fossils, these deposits are not totally barren. In fact, one of the first Mesozoic continental vertebrates ever recorded from Romania comes from one of these deposits, the Lower Cretaceous shallow marine limestones of Southern Dobrogea (Fig. 1). This specimen—the isolated but well-preserved tooth of a large theropod dinosaur—has often been overlooked. It was described a little over a century ago by Simionescu (1913; Fig. 2A), and until a few recent discoveries of very rare isolated specimens (Stoica and Csiki, 2002; Csiki-Sava et al., 2013; Dragastan et al., 2014), it remained as the sole published record of Mesozoic terrestrial

vertebrates from the cratonic areas of Romania. It has never been comprehensively described and its precise age and taxonomic affinities have yet to be clarified, despite its potential importance as a well-preserved fossil from a poorly sampled area that could have critical evolutionary and biogeographic implications.

We here present a comprehensive description of the Dobrogea tooth and discuss its relevance for understanding dinosaur evolution and biogeography. We review the peculiar history of how this specimen was collected and curated, thoroughly document its morphology and age, identify it based on comparison to a broad range of theropods, and outline its importance. It turns out that this specimen, although only a single tooth, has wide-ranging implications. We identify it as coming from the Valanginian stage of the Early Cretaceous, which is incredibly poorly sampled both in Europe and globally (Weishampel et al., 2004), and as belonging to a carcharodontosaurid, a group of derived, large-bodied apex predators whose earliest Cretaceous history is poorly known. Carcharodontosaurids were once thought to be a uniquely Gondwanan group, but recent discoveries show that the basal members of the group were more widespread during the late Early-middle Cretaceous (e.g., Sereno et al., 1996; Brusatte and Sereno, 2008). The Romanian tooth shows affinities with a derived carcharodontosaurid subgroup, the Carcharodontosaurinae, that until now has been known only from Gondwana. It suggests that this subgroup of enormous predators did not evolve vicariantly as Pangaea split, but originated earlier, and perhaps in Europe, suggesting faunal interchange between Europe and Gondwana during the ‘dark ages’ of the early Early Cretaceous.

Abbreviations: UAIC – University “Alexandru Ioan Cuza”, Iași, Romania.

2. History of collecting and curation

Only two dinosaurian fossils are currently known from the cratonic areas of Romania: an isolated theropod tooth and an isolated caudal vertebral centrum. Both of these were reported from the Lower Cretaceous deposits of Southern Dobrogea (southeastern Romania; Csiki-Sava et al., 2013, see also below). Unfortunately, exact details of their discovery and places of origin are lost, a fact that can hinder an assessment of their age and interpretation of their phylogenetic and palaeobiogeographic significance. Our aim here is to gather and report all available information concerning the collecting of specimen UAIC (SCM1) 615, that is, the isolated theropod tooth reported by Simionescu (1913; Fig. 2A).

According to the existing information - unpublished museum labels and records, and the preliminary publication of Simionescu (1913) - specimen UAIC (SCM1) 615 was discovered in the surroundings of Cochirleni, a small village south of Cernavodă and close to the right bank of the Danube, in Southern Dobrogea, southeastern Romania (Fig. 1), probably shortly before 1913, the date of its publication by Simionescu (1913).

Although studied and preliminarily described by Simionescu, UAIC (SCM1) 615 was not collected by Simionescu personally. Instead, it was donated by a certain “de Tomas” (also mentioned as “de Thomas” in the registry of the Hârşova Museum) to V. Cotovu from Hârşova (Central Dobrogea), a local teacher, archaeology and natural history aficionado, and amateur fossil collector (see, e.g., Covacef, 1995). Cotovu, described by Simionescu himself as the “zélé fondateur et directeur du muséum de Hârşova” (*enthusiastic founder and director of the Hârşova Museum*; Simionescu, 1906: p. 2), had previously provided fossil specimens from Southern Dobrogea for study to Simionescu, a nationally acknowledged popular science writer and scientist, whom Cotovu knew personally (Brânzilă, 2010). These circumstances are supported by the fact that in the original description, Simionescu figures the specimen as being accessioned in the “Regional-Museum von Harschowa” (*Hârşova Regional Museum*; Simionescu, 1913: p. 687, fig.1), a designation he also used to refer to other Dobrogean

specimens not collected by him first-hand (e.g., a specimen of '*Nautilus' pseudoelegans* from Cernavodă, or a fragmentary tooth-bearing palatal fragment referred to as '*Coelodus*' sp., also originating from Cochirleni; see Simionescu, 1906). Confirming this deduction, an isolated tooth appears accessioned in the old registry book of the Hârşova Museum (under specimen number 200) as "*Megalosaurus cf. superbus*", with the mention that it was "described by Prof. Simionescu in the *Centralblatt f. min. etc.*". This is also the case of the '*Coelodus*' sp. specimen from Cochirleni (specimen number 86), similarly clearly identified as being described by Simionescu in the registry book.

Both of these vertebrate remains from Dobrogea that were formerly part of the Hârşova Museum collections are currently accessioned in the palaeontology collections of the UAIC (Turculeţ and Brânzilă, 2012), suggesting that, at one moment, several specimens were transferred there from the Hârşova Museum. Although no details are known about this transfer, it is probable that it took place right before (or when) the Hârşova Museum, including a part of its collections, was burned and largely destroyed during WWI, in 1916, a time when Simionescu still held a position at the UAIC.

After its original description, specimen UAIC (SCM1) 615 underwent a minor amount of damage (see below, Description). Also, at some point between its description in 1913 and the early 1960s (when the specimen was found in its present state in the collections of the UAIC by academic staff members who are still alive today and recall the discovery; I. Turculeţ, personal communication, May 2013) it was glued into a limestone matrix holder, while it was obviously completely freed of the surrounding matrix when it was described and figured in 1913 (Fig. 2). The circumstances under which these alterations took place are unclear. It is a distinct possibility that they occurred sometimes during WWII, when, in the spring of 1944, the frontline between the German-Romanian and Soviet armies reached the Iaşi–Chişinău line. At this moment, the geological-palaeontological collections of the UAIC

150 were packed in crates, and moved together with its personnel and other possessions to Zlatna,
151 in the Apuseni Mountains (western Romania), to safeguard them from any potential damage.
152 Mounting the specimen into the limestone stand would have been a quick way to stabilize it,
153 as it appears that packaging and transport of the specimens was done in haste (M. Brânzilă,
154 personal communication, April 2103). If that was indeed the case, the mounting would have
155 taken place without the knowledge of Simionescu, who left Iași and the UAIC in 1929, being
156 invited to become a professor of Palaeontology at the University of Bucharest (Brânzilă,
157 2010). Then again, however, Simionescu himself or staff of the Hârșova Museum might have
158 re-mounted the tooth after its original description, or else the mounting might have taken
159 place after the return of the collections to Iași, after WWII.

160 Unfortunately, it is not documented whether the mounting was made using the
161 original matrix, or if a trough corresponding to the tooth outline was carved into a randomly
162 chosen limestone block. The apparently excellent fit between the tooth and the depression
163 housing it (Fig. 2B, 3) suggests that this operation was completed carefully, and accurate
164 carving of a fake holder is difficult to reconcile with the rush accompanying the evacuation of
165 the Iași University, in 1944. Alternatively, the presence of a hand-written old registration
166 number on the specimen holder would support its early re-mounting, while still at the
167 Hârșova Museum. As noted previously, the original Hârșova Museum registration number of
168 the specimen was 200, which does not correspond to that currently written both on the
169 limestone holder and on a paper sticker (204). However, according to the old collection
170 registry of the Hârșova Museum, specimen numbers 201 through 225 were given to a series
171 of “indeterminate (fossil) bone fragments” from the “Cochirleni quarries”. Thus, these
172 specimens (now apparently lost) came from the same locality as the tooth, and they were
173 collected and donated by the same person to the Museum who donated UAIC (SCM1) 615.
174 There is, thus, a (albeit admittedly remote) possibility that the registration numbers were

175 mixed up during the re-mounting of the specimen, which in this case took place at an early
176 date in the Hârşova Museum. If this is indeed the case, the limestone used as holder could
177 have been the same as the original matrix of the specimen.

178 To conclude, the history of recovery and curation of the historically important
179 dinosaurian specimen UAIC (SCM1) 615 is rather convoluted and clouded by many
180 uncertainties. The exact date of discovery remains conjectural, and the exact place of the
181 discovery (thus also the original geological context of the tooth) is even more ambiguous.
182 The current state of the specimen, and especially its mounted status, suggest a curatorial
183 history that produced a moderate amount of damage to, but also partially obscured the
184 detailed morphology of the specimen. The convergence of such unfortunate events makes
185 deciphering the age, identity and evolutionary significance of the specimen troublesome,
186 although many lines of evidence, carefully considered, allow us to draw reasonable
187 conclusions (see below).

189 3. Geological setting

190 According to the available collecting information, the isolated theropod tooth UAIC (SCM1)
191 615 was discovered at Cochirleni (sometimes noted more specifically as the “*Cochirleni*
192 *quarry*” or “*Cokerleni quarry*”). Cochirleni is a small village in southwestern Dobrogea
193 situated close to the right bank of the Danube, and about 9 km south of the main urban center
194 of the region, Cernavodă (Fig. 1). The geology of the area has been well studied, because of
195 the unique outcropping conditions and rich fossiliferous nature of the Lower Cretaceous
196 deposits (reviewed in Avram et al., 1996; Neagu et al., 1997; Dragastan et al., 1998).

197 Southern Dobrogea is a cratonic area corresponding to the southeastern corner of
198 Romania. Whether it is considered part of the larger Moesian Platform (Săndulescu, 1984;
199 Ionesi, 1994), or a distinct craton (the South-Dobrogean Platform; Mutihac and Mutihac,

2010), researchers agree that it became integrated into the main European Craton towards the end of the Jurassic, at the latest, with the consolidation of the Cimmerian (Early Alpine: Triassic–earliest Cretaceous) North Dobrogean fold-and-thrust belt (Seghedi, 2001; Hyppolite, 2002). The age of its basement is also controversial, with estimates ranging from Archaic–Early Proterozoic (Mutihac and Mutihac, 2010) to latest Proterozoic (Ionesi, 1994).

The Precambrian basement of Southern Dobrogea is overlain by a flat-lying sedimentary cover that begins with the lowermost Palaeozoic and ends with the uppermost Neogene. The sedimentary succession is interrupted by a few major, as well as several less important, sedimentary hiatuses that separate 5 main sedimentary sequences corresponding to the Cambrian–Upper Carboniferous, the Permian–Triassic, the Middle Jurassic–Cretaceous, the Eocene–?Oligocene, and the middle Badenian (middle Miocene)–Upper Pliocene. The Palaeozoic and lower Mesozoic are known only from the subsurface of Southern Dobrogea, but Cretaceous and Cenozoic deposits have limited exposures along the main water courses of the region (Ionesi, 1994; Mutihac and Mutihac, 2010).

The outcropping Cretaceous in Southern Dobrogea is represented mainly by shallow marine, carbonate platform deposits in the lower part of the system, replaced by more open-water, chalky facies towards the later part of the period (e.g., Avram et al., 1993, 1996; Dragastan et al., 1998; Dinu et al., 2007); these crop out only as isolated patches along the main watercourses of the region (Fig. 1).

The Lower Cretaceous Series consists of several lithostratigraphic units with complex, partially overlapping and interfingering relationships (Dragastan et al., 1998, 2014). The lowest (and only artificially) outcropping unit is the Purbeck-type, siliciclastic-evaporitic Upper Kimmeridgian–Lower Berriasian Amara Formation that represents lagoonal to continental environments. This unit is covered by the shallow-marine, richly fossiliferous and locally reefal limestone-dominated Cernavodă Formation (restricted-open lagoonal to

carbonate platform, Upper Berriasian–Lower Hauterivian). A time-correlative unit of the Cernavodă Formation, the limestone-dolomitic Dumbrăveni Formation (Upper Berriasian–Lower Hauterivian), is restricted to the southeastern part of Southern Dobrogea. The Cernavodă and Dumbrăveni formations are covered unconformably by dominantly calcareous deposits with hippuritoid (‘pachyodont’) coquinas, small reefs and lens-like orbitolinid accumulations, referred to the Barremian–Lower Aptian Ostrov Formation by Dragastan et al. (1998), but to the Ramadan Formation (in part) by Avram et al. (1993, 1996). These deposits, formed in littoral to lagoonal and open reef terrace environments, are in turn capped by the fluvial-lacustrine, siliciclastic deposits of the Gherghina Formation, with Middle–Upper Aptian kaolinitic clays and thin coal intercalations. The Lower Cretaceous succession ends with the transgressive, glauconite-bearing, coastal to sublittoral siliciclastic deposits of the Cochirleni Formation (uppermost Aptian–Albian).

The Upper Cretaceous has a significantly more patchy development, mainly restricted to the eastern part of Southern Dobrogea, excepting the weakly glauconitic, chalky-sandy Peștera Formation (Lower Cenomanian) and the marly Dobromiru Formation (Upper Cenomanian) that cover the western-central parts of the area. The younger Cuza Vodă (Turonian), Murfatlar (Santonian–Lower-Middle Campanian), and Satu Nou (Upper Campanian) formations are dominantly chalky, suggesting the instalment of a relatively deeper, offshore depositional environment; neither of these units is known from western Southern Dobrogea.

In total, the Lower Cretaceous of Southern Dobrogea was deposited in a shallow marine, near-shore setting, fluctuating between carbonate platform, lagoonal, coastal-tidal flat, and continental environments (see Avram et al., 1996; Dragastan et al., 1998). Its main characteristic features, such as the observed lithological variability, the areal distribution of the different units, and the presence of several unconformities within the series, are all linked

to eustatic sea-level changes that affected the Southern Dobrogean territory during the Early Cretaceous (Dragastan et al., 1998). The main emergent land in the area was represented by the Central Dobrogean Massif, lying north of the study area, almost completely subaerially exposed and actively eroding during the Cretaceous. Consequently, shallow-marine to continental deposits are restricted mainly to the northern part of Southern Dobrogea, close to its boundary with the Central Dobrogean Massif (marked by the Capidava-Ovidiu Fault), and are replaced by more open marine deposits southward. As summarized above, several littoral, and even continental, sequences occur in this succession, including deposits in the Amara, Cernavodă, Ramadan (Avram et al., 1996) and Cochirleni formations, whereas the Gherghina Formation is purely continental, with occasional minor marine interbeds produced during short-term ingressions of the sea.

In the Cernavodă-Cochirleni area the outcropping Mesozoic is restricted to the Lower Cretaceous, and includes deposits belonging to the Cernavodă, Ostrov (or Ramadan), Gherghina, and Cochirleni formations. While the lower–middle part of the Cernavodă Formation is well exposed and widely distributed in this area, its upper part (the lower Hauterivian Vederoasa Member) is unevenly developed. This member is missing in the classical succession from Cernavodă-Hinog, on the right bank of the Danube (Dragastan et al., 1998), but was recently identified in the more eastern Cernavodă-lock section (Dragastan et al., 2014). Similarly, the Ostrov Formation is represented in the area only by its upper subunit (the Lower Aptian Lipnița Member; Dragastan et al., 1998), covering unconformably and transgressively the Valanginian Alimanu Member of the Cernavodă Formation in the southern end of the Cernavodă-Hinog section (Dragastan et al., 1998), and the lower Hauterivian Vederoasa Member in the Cernavodă-lock section (Dragastan et al., 2014).

Northward of the Hinog area, Valanginian deposits of the Alimanu Member are overlain directly by the Middle–Upper Aptian continental deposits of the Gherghina

Formation. These continental deposits also cover the *Orbitolina*-bearing calcareous-clayey deposits of the Lipnița Member towards the south, marking the advancement of emerged areas towards the central parts of Southern Dobrogea, including the Cernavodă-Cochirleni area, during this time interval (Avram et al., 1996). Marine conditions returned in the study area again in the latest Aptian, with a transgression marked by widespread deposition of the glauconitic, siliciclastic coastal to innermost shelf deposits of the Cochirleni Formation. These uppermost Aptian to Albian sands and sandstones cover transgressively all the underlying deposits, belonging to the Cernavodă, Ostrov, or Gherghina formations. Siliciclastic shallow-marine sedimentation continued into the Early Cenomanian, with the chalky-glauconitic deposits of the Peștera Formation.

4. Palaeontology

The isolated theropod tooth UAIC (SCM1) 615 (formerly in the collections of the Hârșova Museum, registered with no. 200; Fig. 2A) was described in a short note by Simionescu (1913), who referred it to *Megalosaurus* cf. *superbus*, a taxon erected by Sauvage (1882) from the Gault ('mid'-Cretaceous: Albian) of the Paris Basin, France. The Gault material described by Sauvage (1882; see also Sauvage, 1876) includes several isolated teeth that were deemed by Simionescu (1913) to be more similar to the Cochirleni tooth than are the teeth of *Megalosaurus bucklandi* (Buckland, 1824). Subsequently, the French Gault material was referred to the new genus *Erectopus* by Huene (1923), who also noted differences between it and the type species *M. bucklandi*.

The convoluted taxonomic history of *Erectopus superbus* was recently reviewed by Allain (2005), who established that both the isolated teeth first mentioned by Sauvage (1876) and the skeletal elements described by Sauvage (1882) belong to the same taxon, for which the name *Erectopus superbus* was retained. Allain (2005) regarded *Erectopus* as a member of

Carnosauria (= basal Tetanurae), an opinion also shared by Molnar (1990) and Holtz et al. (2004a), whereas the latest review of the Tetanurae (Carrano et al., 2012, p. 254) considered *Erectopus superbus* “a non-carcharodontosaurian allosauroid, possibly a metriacanthosaurid.” Accordingly, if we are following the original assessment of Simionescu (1913) but updating with contemporary taxonomy, the Cochirleni theropod tooth should now be considered referable to the basal tetanuran *Erectopus superbus*. However, the referral of this tooth to *Erectopus superbus* (or a close relative) was considered to be unsupported by positive evidence by Molnar (1990) and Holtz et al. (2004a). In order to re-assess this referral and to understand the exact taxonomic and phylogenetic affinities of UAIC (SCM1) 615 (Fig. 2B, 3), we provide here a detailed description of its morphology followed by a thorough comparative study of this tooth based on large datasets of theropod dental measurements and discrete characters compiled by Hendrickx and Mateus (2014) and Hendrickx et al. (2015a).

We note that in his review of Romanian dinosaurs, Grigorescu (2003) erroneously considered UAIC (SCM1) 615 as being referred by Simionescu to the taxon *Megalosaurus dunkeri* Kohen (sic; actually, *Megalosaurus dunkeri* Dames, 1884). This is clearly a simple misreading of Simionescu’s identification. Additionally, such a referral is also contradicted by the absence of mesial serrations in the holotype tooth of *M. dunkeri*, considered by Carrano et al. (2012) to represent an indeterminate theropod. The Dobrogea tooth, on the other hand, has mesial serrations (see below).

4.1. Age of UAIC (SCM1) 615

The age of UAIC (SCM1) 615 has been contentious, due to the uncertainties concerning its place of origin. Although it is often mentioned as originating from Cochirleni village (e.g., Grigorescu, 2003; Turculeț and Brânzilă, 2012), this has not been definitively established. According to the original report of Simionescu (1913), the tooth came from the upper part of

the Lower Cretaceous limestone succession exposed in the cliffs extending from Cernavodă to Cochirleni along the right bank of the Danube. The corresponding entry from the Hârşova Museum registry states that it was found in the ‘Cochirleni quarry’, a location that presently cannot be identified precisely. The only rocks to be quarried in the area are the calcareous deposits of either the Cernavodă or Ostrov formations, particularly the ones that crop out in the Danube bank cliffs between Cernavodă-Hinog-Cochirleni. Finally, although the mention ‘Cochirleni’ is usually considered to refer to Cochirleni village, it should be mentioned that the cliff-forming hill that extends between Cernavodă and Cochirleni is also known by the same name (Fig. 1). Taking all of this evidence into consideration, it is thus reasonable to conclude that the tooth was most likely found in the Lower Cretaceous limestone succession exposed in the Danube cliffs between Cernavodă and Cochirleni.

Based on the location of the discovery, in the upper part of the local limestone succession, and the age of the deposits from Cernavodă-Cochirleni known to him, Simionescu (1913) considered the tooth to be of Barremian age. Subsequently, the age of the tooth was given as Valanginian–Barremian (Weishampel, 1990; Weishampel et al., 2004) or Valanginian (e.g., Grigorescu, 2003), but without any supporting information.

New attempts have been made to more precisely constrain the age of UAIC (SCM1) 615. Dragastan et al. (2014) recently sampled the limestone matrix holder of the tooth, and reported from these samples an assemblage of foraminiferans, ostracods and microproblematicae (= *incertae sedis* microorganisms) that characterize their ‘Biozone IX with *Meandrospira favrei*’, of latest Valanginian age in the local lithostratigraphic scheme. In parallel, we also sampled the same limestone holder – a yellowish white, friable lime mudstone – that yielded a poor and badly preserved calcareous nannoplankton assemblage with *Watznaueria barnesiae*, *W. ovata*, *Nannoconus steinmanni*, *N. kamptneri*, *N. globulus*, *Calcicalathina* sp., *Speetonia colligata* and *Cyclagelosphaera deflandrei* (M. C. Melinte-

Dobrinescu, personal communication, November 2013), an assemblage that suggests a Berriasian–Hauterivian age of the limestone holder.

Since it is not clear if the limestone holder came from the same site as the tooth itself, we managed to take a second sample from the limestone matrix still partly filling the pulp cavity of the tooth, which must definitively be identical with the rocks the tooth was found in. This second, much smaller sample yielded only very scarce specimens of *Watznaueria barnesiae*, *Cyclagelosphaera margerelii* and *Diazomatolithus lehmanni* (M. C. Melinte-Dobrinescu, personal communication, November 2013), the latter two taxa having a peak in abundance during the Berriasian and, especially, the Valanginian.

In the nannoplankton succession reported previously by Avram et al. (1993) and derived from a systematic sampling of the Southern Dobrogean Lower Cretaceous, the concurrent presence of *Speetonia colligata*, *Calcicalathina oblongata*, *Diazomatolithus lehmanni* and *Nannoconus steinmanni* was noted in samples derived from the Alimanu Member of the Cernavodă Formation. These assemblages were interpreted to represent the nannoplankton zone CC3 of Sissingh (1977), of late Valanginian age. A comparable age was assigned to a roughly similar nannoplankton assemblage reported from the Lower Cretaceous of the Mecsek Mountains, Hungary, by Császár et al. (2000).

Together, all the available evidence (Simionescu's original account, geographic and geologic records, foraminifera, ostracods, microproblematicae, and calcareous nannoplankton) thus suggests that UAIC (SCM1) 615 originates from the Alimanu Member of the Cernavodă Formation, and it is most probably of late Valanginian age.

4.2. Description and comparisons

Specimen UAIC (SCM1) 615 is a large (total length, as preserved, is about 100 mm; Figs. 2, 3) lateral tooth of a theropod dinosaur, with a crown base length (CBL) of 29 mm, crown

base width (CBW) of 16.25 mm, crown height (CH) of 85.5 mm, and apical length (AL) of 91 mm (terminology following Smith et al., 2005 and Hendrickx et al., 2015b). It is remarkably well preserved, with the enamel in pristine condition. It preserves most of the crown and a small basal part of the root, but the crown tip is broken off, with an estimated 5 mm missing in the apical region.

In its present state, the mesial edge and part of the mesial third of the tooth are embedded in the limestone holder (Fig. 2B), although the tooth was once removed (see above, History of collecting and curation; Fig. 2A). Accordingly, it is exposed so that all faces of the tooth are widely visible, including the root region, except for the mesial surface.

Only the basal-most part of the root is preserved, and it is more complete near the mesial margin (Fig. 3B, C). Here, broken areas around the crown-root contact area (cervix) reveal details of the pulp cavity development, as well as the pattern of the dentine thickness variation (Fig. 3B–D). The crown also exhibits a transverse break at about two-thirds of its length (not present so obviously in the original figure of the specimen in Simionescu, 1913), and adjacent to it, the distal carina is also slightly chipped distal to mid-length. The labial face is superficially split near this break (Fig. 3A), while a more prominent region of damage appears on the lingual face, where a large (13 x 5 mm), slightly triangular wedge is broken off, exposing the deeper parts of the dentine (Fig. 3C). The damage to the lingual side apparently occurred after the original description of the tooth (Fig. 2), an observation that is concordant with the complex curatorial history of the specimen.

The basal-most, exposed part of the mesial face lacks the enamel cover (Fig. 3C, D), suggesting that this area already belongs to the root region. The mesial edge of the preserved crown base appears to be wider than the distal one, and is largely rounded transversely. Accordingly, the basal cross-section is teardrop-shaped (lanceolate); it is rounded mesially, but narrows distally into a small carina (Fig. 3D). As mentioned above, the pulp cavity is

exposed basally, being partly filled with a whitish-light gray limestone that is reminiscent of the matrix holder lithology. The pulp cavity narrows rapidly towards the cervix, as it is about 7.1 mm wide (labiolingually) at the apical-most part of the preserved root, but only about 4.5 mm wide at the base of the crown. In parallel, the enamel-dentine wall of the tooth becomes thicker: it is 3.5 mm thick in the apical-most part, 4.4 mm at the base of the crown, but thickens to 5.0–5.8 mm near the apical-most part of the basal break of the crown (Fig. 3B). Mirroring the outside cross-section, the contour of the pulp cavity is also teardrop-shaped (Fig. 3D).

The tooth is ziphodont and only very slightly recurved distally. The distal edge is nearly straight across its length, being very mildly concave in its basal half and slightly convex near its apex (Fig. 2, 3A). Thus, the apex is placed roughly at the distal margin of the tooth crown base. The mesial edge, as shown in the original publication of Simionescu (1913), is strongly convex across its entire length (Fig. 2A). The tooth is labiolingually compressed (Fig. 3B), with a crown base ratio ($CBR = CBW/CBL$) of 0.56, within the normal range of variation of most theropods. This differs from the thinner teeth of some, but not all, carcharodontosaurids ($CBR < 0.50$), and the much thicker incrassate teeth of derived tyrannosauroids and conical teeth of spinosaurids ($CBR > 0.75$) (Serenó et al., 1996; Brusatte et al., 2010a; Hendrickx and Mateus, 2014; Hendrickx et al., 2015a).

The crown cross-section is slightly asymmetrical labiolingually when it is seen in distal view. In this view, when the carina is facing directly distally, one side of the crown has a more pronounced bulge than its counterpart (about 8.5 mm wide, measured from the carina, vs. 6 mm on the other side; Fig. 3B); based on comparisons with the teeth of *Mapusaurus* (Coria and Currie, 2006), the more bulging side can be interpreted as the lingual one. This asymmetry diminishes apically, where both sides become about equally convex. The distal carina itself twists slightly sideways (labially) in apical direction, such that it is located closer

to the labial face where it terminates at the crown apex, and the lingual face of the denticles is exposed distally (Fig. 3B, F). This twist of the distal carina is accompanied by a similar outline of the lingual side; in distal view, this is somewhat convex basally, but becomes flat to slightly concave in the apical two-thirds of the crown. A similar S-shaped curvature of the crown, albeit more pronounced and different in details, was also reported in *Mapusaurus* and *Giganotosaurus* (Coria and Currie, 2006), and in indeterminate carcharodontosaurid teeth from Morocco (Richter et al., 2013).

The distal carina extends along the entire tooth height (Fig. 3A–C). It is covered with minute serrations across its entire preserved length; the denticles are proximodistally subrectangular, with a mesiodistal long axis that is greater than the apicobasal long axis (Fig. 3E–H). They are either roughly perpendicular to the tooth margin, or their long axes are oriented obliquely, such that they point slightly apically. The tip of the apex is broken off, so it is not possible to determine whether the serrations continued over the apex of the tooth. There are approximately 12.5 serrations (denticles) per 5 millimetres at the midpoint of the carina. Serration shape and size remain relatively constant across the carina, although the serrations near the midpoint and closer to the base of the carina (12 denticles per 5 mm; Fig. 3G, H) are slightly smaller than those near the apex (9 denticles per 5 mm; Fig. 3E, F). Changes in serration size are gradual across the carina, not sudden or sporadic.

Although they are all more or less rectangular in shape, the apical denticles are relatively shorter proximodistally than the more basal ones. Most of the denticles have slightly rounded, asymmetrically convex triangular tips, instead of being simply squarred-off, and they do not hook as in troodontids and to a lesser extent abelisaurids (Hendrickx and Mateus, 2014). Other denticles near the apex, however, show a faint concavity along their tips, giving them a bilobate aspect, although this is both less conspicuous and far less regularly developed than reported in *Tyrannotitan* (Novas et al., 2005). The denticles are

separated by simple, linear grooves (interdenticular slits or sulcae) along their entire length.

The interdenticular space between adjacent denticles is broad, measuring more than a third of the apicobasal width of a denticle (Fig. 3E, G). This space continues onto the surface of the crown as a very short interdenticular sulcus (“blood groove” of Currie et al., 1990). These sulci are so short and indistinct that they are only visible under low angle light.

Little can be said about the mesial carina, as it is not visible in the current state of the specimen, buried in the limestone matrix. Based on the description of Simionescu (1913), however, it is covered across its length with minute serrations; these decrease in size towards the base of the crown. Simionescu (1913) reported approximately 15 serrations (denticles) per 5 millimetres at the midpoint of the carina, meaning that the mesial denticles are slightly smaller than those on the distal carina. The denticle size difference index (DSDI: Rauhut and Werner, 1995) is 1.2, within the range of variation of most theropods (Hendrickx and Mateus, 2014). As Simionescu (1913) already pointed out, the presence of a mesial carina that extends towards the base of the crown sets apart UAIC (SCM1) 615 from *Megalosaurus bucklandii* where this stops well above the cervix (Benson et al., 2008), and it is instead similar to ‘*M. superbis*’ (Sauvage, 1876, 1882) in this respect.

The external enamel surface exhibits two forms of ornamentation. First, the majority of the labial and lingual faces are covered by relatively smooth enamel that exhibits a subtle form of braided texture visible under low angle light (Fig. 3A, C, E). This texture is made up of a series of very faint, apico-basally running ridges; these are of unequal lengths, starting at different points of the crown height, but none extends the whole length of the crown. The two longest ridges are placed near the distal carina. The enamel is also finely granulated.

Second, near the carinae on both labial and lingual surfaces there are marginal undulations: wrinkles in the enamel that stand out in bas relief (Brusatte et al., 2007). These are much better preserved and visible near the distal carina, where they are so pronounced

that they are clearly observable in normal light (Fig. 3A–C, G, H). Here, about 17 unevenly developed wrinkles are present along the crown height; in the basal half of the crown, the wrinkles extend about 6.5 mm onto the crown. These are elongate, such that they are longer than twice the space separating each undulation. The wrinkles project obliquely (in the mesiobasal direction) relative to the carina. They are apically concave, with a near-horizontal segment on the crown, and curve apically as they approach the carina (at about 45°) with a tendency to become tangential to the distal edge. The wrinkles are especially well developed, prominent and closely spaced in the basal part of the crown (about 7 wrinkles/16 mm; Fig. 3C, G)), but become more widely spaced and indistinct apically (about 3 wrinkles/16 mm). Apically, however, the wrinkles are somewhat wider and longer, extending over about half of the crown fore-aft length. Again, a slight asymmetry is present between the two sides of the crown in wrinkle development as well, these being better expressed on the more rounded, convex lingual face, but less well expressed on the flatter labial face (Fig. 3A, C, H). On the presumed labial face, only some of the basal-most wrinkles, particularly the second and third one, appear well defined.

Towards the base of the crown a few of the wrinkles continue across the labial and lingual surfaces as very subtle transverse undulations. Most conspicuous of these is a 3.5 mm wide horizontal swelling that crosses the crown, at the level of wrinkles 2 and 3; this swelling is clearly visible on both sides of the crown (Fig. 3. A, C). There are no lateral flutes, apico-basal ridges, or longitudinal grooves on the labial or lingual faces, either in the centre of the tooth or paralleling the carinae. Instead, the labial and lingual faces are uniformly convex, giving the tooth its teardrop-shaped outline in cross section.

5. Discussion

5.1. *Identification of UAIC (SCM1) 615*

The isolated tooth from Cochirleni can be referred to Theropoda based on its large size, recurved and labiolingually compressed morphology, and presence of a continuous series of well-defined serrations on the distal carina.

Besides theropods, certain derived crocodyliforms – the sebecosuchians of Colbert (1946; see also Turner and Sertich, 2010; Pol and Powell, 2011; Rabi and Sebők, 2015) – are also known to possess remarkably theropod-like, laterally compressed and serrated teeth, not unlike the morphology shown by UAIC (SCM1) 615. However, most sebecosuchian teeth are significantly smaller than the Southern Dobrogean specimen, especially in the case of the Cretaceous members of the clade (e.g. *Baurusuchus*; Carvalho et al., 2005). Even the largest, caniniform teeth of the largest representatives of Sebecosuchia, such as the Miocene *Barinasuchus* (Paolillo and Linares, 2007), are somewhat smaller than UAIC (SCM1) 615; moreover, these teeth are slightly conical and less laterally compressed than the Southern Dobrogean tooth. Finally, it should be noted that the oldest known members of Sebecosuchia appear beginning in the Late Cretaceous (e.g. Kellner et al., 2014), and are thus significantly younger than UAIC (SCM1) 615. Similarly, ziphodont crocodyliform teeth (i.e. with true denticles along their carinae) are reported in Europe only beginning in the Albian (Ösi et al., 2015), and these are both significantly smaller and different in morphology from the Dobrogean tooth. Taken together, these suggest that the hypothesis of sebecosuchian affinities of UAIC (SCM1) 615 can be discarded with confidence, and it indeed represents a theropod tooth.

We used four techniques to identify which type of theropod UAIC (SCM1) 615 likely belongs to (see also Supplementary Material).

First, we conducted a Principal Components Analysis (PCA) based on a large database that includes a broad and representative sample of theropod teeth. This dataset was compiled by Hendrickx et al. (2015a), which built upon the earlier studies of Smith et al.

(2005) and Larson and Currie (2013), and it or a similar version has been used in recent studies to identify isolated theropod teeth (e.g., Williamson and Brusatte, 2014; Brusatte and Clark, 2015). It comprises nearly 1000 theropod teeth scored for six measurements (CBL, CBW, CH, AL, MC, and DC, the latter two measuring the density of serrations per 5 mm at the midpoint of the mesial and distal carina, respectively). UAIC (SCM1) 615 was added to this dataset, the data were log-transformed prior to analysis, missing values for measurements were estimated with a mean value for that measurement from across the sample, and then a PCA was run using a correlation matrix. The analysis was conducted in PAST v2.17 (Hammer et al., 2001).

In the resulting two dimensional morphospace (Fig. 4), UAIC (SCM1) 615 plots close to many teeth belonging to carcharodontosaurids, along with some teeth belonging to spinosaurids and tyrannosauroids. It falls within the convex hull (maximum morphospace occupation area) of carcharodontosaurids only, although it is closely outside of the edges of spinosaurid and tyrannosauroid space. It also falls within the 95% confidence interval ellipse for carcharodontosaurids, but not within the ellipse of any other group (Supplementary Information). This exercise indicates that UAIC (SCM1) 615 is most similar to carcharodontosaurids.

Secondly, we used the log-transformed dataset that we also used for the PCA to conduct a clustering analysis. We performed the analysis in PAST v2.17, using the paired group algorithm and the correlation similarity measure. In the resulting dendrogram, UAIC (SCM1) 615 groups with a handful of teeth belonging to carcharodontosaurids, tyrannosauroids, and *Allosaurus* (Supplementary Information).

Third, we used the tooth measurement database to conduct a discriminant analysis in PAST v3.0 (Hammer et al., 2001). This analysis uses pre-determined groups (in this case, taxonomic clusters) to create a morphospace in which these groups are maximally separated.

This allows teeth of unknown affinities, such as UAIC (SCM1) 615, to be classified according to which taxonomic group it is most similar to in this discriminant morphospace. In total, 67.79% of other teeth are classified correctly when they are treated as having uncertain affinities and their measurements are used to classify them in discriminant space, indicating that this exercise returns reasonable results. Our analysis classifies the Romanian tooth as a carcharodontosaurid. Furthermore, the analysis places UAIC (SCM1) 615 within the convex hulls for carcharodontosaurids and tyrannosauroids, and the 95% confidence ellipses for carcharodontosaurids, coelophysoids, and neovenatorids.

Fourth, we ran a phylogenetic analysis by including UAIC (SCM1) 615 in the discrete character dataset of theropod dental features published by Hendrickx and Mateus (2014). The Romanian specimen was scored as a lateral tooth in this analysis. The analysis was conducted in TNT (Goloboff et al., 2008), and resulted in 224 most parsimonious trees (686 steps, consistency index of 0.338, retention index of 0.566). The strict consensus topology is moderately well resolved and places the Romanian tooth as the sister taxon to *Carcharodontosaurus* (Supplementary Material). This sister taxon pair is recovered as the sister clade to a grouping of the derived carcharodontosaurids *Mapusaurus* and *Giganotosaurus*.

Several synapomorphies support the carcharodontosaurid affinities of UAIC (SCM1) 615. The sister group relationship with *Carcharodontosaurus* is supported by two features: a roughly straight distal margin of the crown (character 68) and pronounced marginal undulations in the enamel that are well visible in normal light (character 112). The broader clade of UAIC (SCM1) 615, *Carcharodontosaurus*, *Mapusaurus*, and *Giganotosaurus* (= Carcharodontosaurinae, as defined by Brusatte and Sereno, 2008, and Carrano et al., 2012) is linked by numerous characters, including: large teeth with a crown height greater than 6 cm (character 65), a bowed or sigmoid distal carina in distal view (character 82), marginal

undulations that are at least twice as long mesiodistally as the space separating each undulation (character 111), and marginal undulations present on both mesial and distal sides of the crown (character 113).

The Romanian specimen also lacks many keystone dental synapomorphies of other theropod clades, based on the clade diagnoses of Hendrickx and Mateus (2014) and other cladistic studies that include dental characters. UAIC (SCM1) 615 does not possess the hooked distal denticles of some Abelisauridae, the strongly labially deflected distal carina and pronounced transverse enamel undulations extending across the labial and lingual tooth faces of Ceratosauridae, the incrassate teeth with apicobasal enamel flutes and deeply veined enamel surface texture of Spinosauridae, and the large transverse undulations of some basal allosauroids (Hendrickx and Mateus, 2014). It also lacks the thickened incrassate teeth of derived tyrannosauroids (Brusatte et al., 2010a) and the large and strongly hooked (or pointed) denticles of troodontids and therizinosauroids (e.g., Turner et al., 2012; Brusatte et al., 2014; Hendrickx and Mateus, 2014). The large size, as well as recurved and ziphodont shape of UAIC (SCM1) 615 is strikingly different from the non-ziphodont therizinosauroids, ornithomimosaur, alvarezsauroids, and most troodontids, which have conical, leaf-shaped, or peg-like teeth (when teeth are present) (e.g., Holtz et al., 2004a; Turner et al., 2012; Brusatte et al., 2014). Finally, besides its remarkably large size, the presence of serrations indicates that UAIC (SCM1) 615 does not belong to groups such as alvarezsauroids, oviraptorosaurs, basal troodontids, or avialans, which have unserrated crowns (e.g., Turner et al., 2012; Hendrickx and Mateus, 2014).

In summary, the four analyses all support carcharodontosaurid affinities for UAIC (SCM1) 615. Both overall tooth proportions and discrete phylogenetic characters point to a carcharodontosaurid identification, and the discriminant function analysis and phylogenetic analysis both explicitly recover the tooth as a carcharodontosaurid. For this reason we refer

1 this tooth to Carcharodontosauridae. Moreover, it appears to belong to a clade that unites very
2 derived and large-sized carcharodontosaurids (*Carcharodontosaurus*, *Giganotosaurus*, and
3 *Mapusaurus*), separated as such and named Carcharodontosaurinae by Brusatte and Sereno
4 (2008) and Carrano et al. (2012). The well-resolved internal topology of this clade, as
5 recovered in our analysis, is congruent with results of previous analyses based on larger sets
6 of characters from across the skeleton (e.g., Coria and Currie, 2006; Brusatte and Sereno,
7 2008; Brusatte et al., 2009; Ortega et al., 2010; Eddy and Clarke, 2011; Canale et al., 2015),
8 and offers some support for considering the Romanian carcharodontosaurid from Southern
9 Dobrogea as more closely related to the African *Carcharodontosaurus* than to the clade of
10 the South American giant carcharodontosaurids *Giganotosaurus* or *Mapusaurus*.

11 Two final notes are worth adding. First, our analyses also incorporated
12 carcharodontosaurids that are usually found to be basal within the clade, such as
13 *Acrocanthosaurus* and *Eocarcharia* (e.g., Harris, 1998; Sereno and Brusatte, 2008; Carrano
14 et al., 2012) as well as a host of other allosauroids, including members of Neovenatoridae
15 (*Neovenator*, *Australovenator* and *Fukuiraptor*), a clade that is often recovered as sister-
16 taxon to carcharodontosaurids within Carcharodontosauria (e.g., Benson et al., 2010; Carrano
17 et al., 2012; but see Novas et al., 2013; Porfiri et al., 2014, for an alternate placement of
18 neovenatorids in general). Both PCA and phylogenetic analysis clearly identified UAIC
19 (SCM1) 615 as more closely comparable morphologically to derived carcharodontosaurids
20 than to either basal carcharodontosaurids or to any other allosauroid subclade.

21 Second, our datasets also included teeth of *Erectopus*, the genus erected for
22 ‘*Megalosaurus*’ *superbus* to which UAIC (SCM1) 615 was originally referred. Again, our
23 analyses clearly indicate that there are no close morphological and morphometric similarities
24 between the two, which is in accordance with the suggestion of Carrano et al. (2012) that
25 *Erectopus* represents a non-carcharodontosaurid taxon, while our analysis identifies UAIC

(SCM1) 615 as a carcharodontosaurid. Instead, *Erectopus* groups with abelisauroids in the phylogenetic analysis. This is somewhat surprisingly, as Allain (2005) and Carrano et al. (2012) both identified *Erectopus* as a tetanuran. It should be noted, however, that Albian-aged abelisauroids are known from the same general area (eastern France) as that yielding the material referred to *Erectopus* (Accarie et al., 1995; Carrano and Sampson, 2008), raising the intriguing possibility that this taxon may represent an abelisauroid instead of an allosauroid tetanuran as suggested by Allain (2005) and Carrano et al. (2012). However, it must be remembered that this phylogenetic analysis is based on dental characters only, so it is probably more likely that *Erectopus* is a tetanuran with a dentition convergent to some extent with those of certain abelisauroids.

5.2. Body size of UAIC (SCM1) 615

One of the most salient and remarkable features of UAIC (SCM1) 615 is its large size. In the large and comprehensive sample of theropod teeth from our dataset, tooth size (estimated based on crown height – CH, and used as a rough proxy of body size) ranges from 2.2 mm (in the dromaeosaurid *Saurornitholestes* and the coelurosaur of uncertain affinities *Richardoestesia*) to 117.1 mm in the gigantic tyrannosauroid *Tyrannosaurus*. The Romanian specimen UAIC (SCM1) 615, with a CH of 85.5 mm, is ranked in the 60-80% maximum size (~ CH) range of the sample, and has a CH that is 73% of the largest tyrannosauroid teeth. Most of the teeth in the dataset (over 61% of the 966 measured teeth) are very small to small (less than 25 mm CH), and less than 10% of these fall in the 60-100% CH size categories. Teeth larger than UAIC (SCM1) 615 make up less than 5% of the total sample, and they represent only five taxa: the megalosaurid *Torvosaurus*, the tyrannosauroid *Tyrannosaurus*, the basal carcharodontosaurid *Acrocanthosaurus*, and the derived carcharodontosaurines *Carcharodontosaurus* and *Giganotosaurus*. Compared to other carcharodontosaurids, UAIC

(SCM1) 615 is smaller than the largest teeth of *Acrocanthosaurus* (9% difference), *Carcharodontosaurus* (20%), and *Giganotosaurus* (12.5%) in the dataset, but is 13% bigger than the largest tooth of *Mapusaurus*.

It is thus reasonable to conclude that UAIC (SCM1) 615 belonged to a large-sized carcharodontosaurid, comparable to, even if somewhat smaller than, the truly gigantic carcharodontosaurines *Giganotosaurus* and *Carcharodontosaurus* (Serenio et al., 1996; Calvo and Coria, 1998; Therrien and Henderson, 2007), taxa that were recovered as possible close relatives of the Romanian carcharodontosaurid by our phylogenetic analysis. This, in turn, corroborates growing evidence that very large body size was acquired very early in carcharodontosaurid history, since the earliest potential members of the clade are already of relatively large size (Rauhut, 2011). The oldest potential carcharodontosaurid is *Veterupristisaurus*, represented by isolated vertebrae that indicate an animal between 8.5 and 10 meters in total body length (compared to 11.5+ meters in *Acrocanthosaurus* and more derived carcharodontosaurids) (Rauhut, 2011). These specimens are known from the uppermost Jurassic of Tanzania, eastern Africa (Rauhut, 2011; Carrano et al., 2012; see below), predating at most ~18 million years (Mya) the occurrence of likely even larger-sized carcharodontosaurids in the Valanginian of Southern Dobrogea, Romania.

The inferred large body size of the South Dobrogean theropod is also remarkable as virtually all other dinosaur remains reported previously from Romania (both from the Early Cretaceous Cornet assemblage and the much later, end Cretaceous Hățeg Island fauna) are significantly smaller, and many have been interpreted as insular dwarfs (e.g., Weishampel et al., 1993, 2003; Benton et al., 2006, 2010; Stein et al., 2010; Ősi et al., 2014). Although other Romanian theropod dinosaurs were not particularly dwarfed (e.g. Brusatte et al., 2013), they were nonetheless small (Nopcsa, 1902; Csiki and Grigorescu, 1998; Csiki et al., 2010; Brusatte et al., 2013). This bias towards small bodied Romanian theropods was also

interpreted as a consequence of their insular habitat (Csiki and Grigorescu, 1998), as all previously reported theropod remains come from within the Carpathian Orogen, an area with an archipelago-type palaeogeography during the Cretaceous (Dercourt et al., 2000; Csontos and Vörös, 2004; Csiki-Sava et al., 2015). By contrast, UAIC (SCM1) 615 was found in shallow marine deposits bordering the emerged areas of Central Dobrogea, part of the stable cratonic areas of Europe and connected at least intermittently to the Ukrainian Shield since the Late Jurassic (Fig. 5A). Although cratonic Europe was also transformed into an archipelago of islands during much of the Cretaceous, these islands were often both larger in size and more stable in space and time than were the transient emerged areas of the Tethyan archipelagoes. As such, it is conceivable that the Southern Dobrogean carcharodontosaurid was less constrained by space or resource limitations than the Tethyan insular dinosaurs, allowing it to retain a large body size.

5.3. UAIC (SCM1) 615 and Valanginian dinosaur distribution

Besides documenting the presence of large-sized mainland carcharodontosaurids in the Lower Cretaceous of Romania, UAIC (SCM1) 615 is also important in that it fills a significant gap in our knowledge on the composition and distribution of the Early Cretaceous dinosaurs in Europe. In their review of dinosaur occurrences, Weishampel et al. (2004) listed 83 Early Cretaceous dinosaur localities spread throughout Europe, more than half of these being known from the later part (Barremian–Albian) of that epoch; only around a dozen localities were listed from each age of the early part of the Early Cretaceous (Berriasian, Valanginian, and Hauterivian). Even despite a significant increase in Early Cretaceous dinosaur discoveries in Europe in recent years (e.g., Royo-Torres et al., 2009; Cobos et al., 2010, 2014; Galton, 2009; Norman, 2010, 2013; Pereda-Suberbiola et al., 2011, 2012; Sachs and Hornung, 2013; Blows and Honeysett, 2014), these remain very strongly biased towards

western and southwestern Europe (especially the UK, France and Spain). Frustratingly, no occurrences are known from the entire central, eastern and southern Europe for the Berriasian–Hauterivian time interval except for two from Romania: the Berriasian–Valanginian locality of Cornet (e.g., Jurcsák and Popa, 1979, 1983; Jurcsák, 1982; Benton et al., 1997) in the northern Apuseni Mountains of northwestern Romania, and the carcharodontosaurid tooth (Simionescu, 1913) from the Valanginian of Cochirleni, in Southern Dobrogea, southeastern Romania we are describing here (Fig. 5A).

Our identification of the Romanian tooth as a carcharodontosaurid documents the presence of this clade in Europe in the very early Cretaceous. This is significant, as carcharodontosaurids were widely distributed tens of millions of years later, in the middle Cretaceous (Aptian to Cenomanian), in western Gondwana (Africa and South America, see below). Despite the recent discoveries documenting that the clade was also present in North America and Asia during the middle Cretaceous (e.g., Sereno et al., 1996; Currie and Carpenter, 2000; Brusatte et al., 2009, 2012), there has been only very few occurrences in Europe, most importantly the Barremian-aged *Concavenator* from Spain (Ortega et al., 2010; see below). The carcharodontosaurid tooth from Southern Dobrogea is substantially older than *Concavenator*, demonstrating that carcharodontosaurids appeared in Europe earlier than previously thought and were a long-term component of the European mainland Early Cretaceous faunas. It also suggests that habitat-related palaeobiological differentiation might have been already present between the cratonic, stable European mainland, with a dinosaur fauna made up of normal-sized (even very large) taxa, and the islands from the mobile Alpine areas of the Mediterranean Neo-Tethys, with by now dwarfed dinosaurs such as those described from the Berriasian–Valanginian Cornet assemblage in northwestern Romania (Benton et al., 2006).

This Valanginian carcharodontosaurid represents an important datapoint not only for the Romanian Lower Cretaceous, but also for that of wider Eurasia. The Valanginian is a poorly documented age in dinosaur evolution, with very few precisely dated fossil occurrences from anywhere in the world (e.g., Weishampel et al., 2004). The best record of Valanginian dinosaurs is from Europe, with fewer and less well dated occurrences known from Asia, some of which have debatable or controversial dates. These include sites in Japan (e.g., Manabe and Hasegawa, 1995; Matsukawa et al., 2006; but see Kusuhashi et al., 2009 and Evans and Matsumoto, 2015, supporting an alternative, younger age of these assemblages) and in Thailand (e.g., Buffetaut and Suteethorn, 1998, 2007, with age constraints according to Racey, 2009; Racey and Goodall, 2009). Occurrences of possible Valanginian age from China (e.g., Jerzykiewicz and Russell, 1991; Shen and Mateer, 1992; Lucas and Estep, 1998) are either poorly constrained as early Early Cretaceous, or were shown subsequently to be younger than Valanginian (Lucas, 2006; Tong et al., 2009). Rare dinosaur remains of possible Valanginian (or ‘Neocomian’) age were also reported from southern Africa (e.g., De Klerk et al., 2000) and, tentatively, from North America (e.g., Lucas, 1901; McDonald, 2011, with age assignments according to Sames et al., 2010; Cifelli et al., 2014).

As one of the two known reports of Valanginian dinosaurs in Europe east of France, the Southern Dobrogean dinosaur record fills a huge palaeogeographic gap between the western European and the eastern Asian dinosaur faunas. Moreover, none of these early Early Cretaceous dinosaur assemblages from outside Europe include carcharodontosaurids (see below), as theropods are represented by coelurosaurians interpreted either as compsognathids (Gishlick and Gauthier, 2007) or basal ornithomimosaurs (Choiniere et al., 2012) in southern Africa, metriacanthosaurid allosauroids (‘sinraptorids’) in Thailand (Buffetaut and Suteethorn, 2007), and indeterminate allosauroids (Pérez-Moreno et al., 1993), non-

carcharodontosaurid tetanurans (Carrano et al., 2012) or enantiornithine birds (Lacasa Ruiz, 1989), besides indeterminate taxa (Carrano et al., 2012), in western Europe. This may suggest that carcharodontosaurids had not achieved a wide geographic distribution by this point in time, and that their more cosmopolitan distribution came later, during the middle Cretaceous.

Finally, the presence of the Cochirleni carcharodontosaurid might hint at the presence of palaeobiogeographic provinciality between the western and the eastern parts of Europe, partly mirroring those reported from the later part of the Late Cretaceous (e.g., Le Loeuff and Buffetaut, 1995; Weishampel et al., 2010; Ősi et al., 2012; Csiki-Sava et al., 2015). In the reasonably well sampled, and significantly better known, western European dinosaur faunas, Valanginian large carnivorous dinosaurs include non-carcharodontosaurid tetanurans (*Becklespinax*), as well as indeterminate allosauroids or indeterminate theropods (often described as ‘*Megalosaurus*’ *dunkeri*, ‘*M.*’ *insignis* or ‘*M.*’ *oweni*), none of which can be referred positively to Carcharodontosauridae (Carrano et al., 2012). The apparently provincial geographic distribution of the large-bodied theropods suggests that some degree of faunal differentiation was occurring within the European mainland, most probably promoted by geographic distance. Notably, this intra-European differentiation in theropod assemblages appears to stand in contrast with the faunal homogeneity reported in the case of the ornithomimids from the UK and Romania (e.g., Galton, 2009). It is important, however, to re-emphasize at this point that the Valanginian dinosaur fossil record is both exceedingly poor and patchy, even in Europe. Accordingly, further discoveries are needed to verify and support (or contradict) the presence of such a distribution pattern pointing to palaeobiogeographic provinciality inside Europe, as the one suggested by our carcharodontosaurid identification for UAIC (SCM1) 615.

5.4. UAIC (SCM1) 615 and carcharodontosaurid evolution and palaeobiogeography

Carcharodontosauridae were long considered as an exclusively Gondwanan group of theropods (e.g., Allain, 2002; Novas et al., 2005) since their first discovery in northern Africa (e.g., Stromer, 1931), and subsequent description of a host of referred taxa from the Aptian–Cenomanian of Africa and South America (Coria and Salgado, 1995; Sereno et al., 1996; Novas et al., 2005; Coria and Currie, 2006; Brusatte and Sereno, 2007; Sereno and Brusatte, 2008; Cau et al., 2013). This view started to change with the identification of the Early Cretaceous (Aptian–Albian) *Acrocanthosaurus* from North America as a basal carcharodontosaurid (e.g., Sereno et al., 1996; Harris, 1998; Sereno 1999; Brusatte and Sereno, 2008), suggesting that the clade had a wider, Neopangean palaeobiogeographic distribution by the mid–late Early Cretaceous. Such a wide distribution, even a cosmopolitan one, was further supported by the discovery of definitive carcharodontosaurids in the Lower Cretaceous of Europe (Ortega et al., 2010), and in the upper Lower to lower Upper Cretaceous of China (Brusatte et al., 2009, 2010b, 2012; Mo et al., 2014; Lü et al., 2016).

Together, the available evidence pointed to an early, pre-mid Early Cretaceous origin of the carcharodontosaurids, followed by their dispersal across Laurasia and western Gondwana beginning at least by the Aptian (Fig. 5B), a scenario that is concordant with the tentatively suggested presence of early carcharodontosaurids in the Upper Jurassic of Tanzania, which are based on fragmentary specimens (Rauhut, 2011; Carrano et al., 2012). It is also concordant with the widespread appearance of carcharodontosaurids in the fossil record starting with the Aptian, when they are reported in Africa (*Eocarcharia*; Sereno and Brusatte, 2008), South America (Vickers-Rich et al., 1999), North America (*Acrocanthosaurus*; Stovall and Langston, 1950; Harris, 1998; Currie and Carpenter, 2000; Eddy and Clarke, 2011), Europe (Canudo and Ruiz-Omeñaca, 2003; Pereda-Suberbiola et al., 2012), and eastern Asia (*Kelmayisaurus*; Brusatte et al., 2012; Mo et al., 2014; Lü et al., 2016).

During the Albian–Turonian, carcharodontosaurids became especially abundant and diverse in Africa (*Carcharodontosaurus*, *Sauroniops*; Stromer, 1931; Sereno et al., 1996; Brusatte and Sereno, 2007; Le Loeuff et al., 2012; Cau et al., 2013; Richter et al., 2013) and South America (*Tyrannotitan*, *Giganotosaurus*, *Mapusaurus*, alongside with indeterminate carcharodontosaurids; Coria and Salgado, 1995; Calvo and Coria, 1998; Novas et al., 2005; Coria and Currie, 2006; Casal et al., 2009; Candeiro et al., 2011; Canale et al., 2015; Fig. 5B). They were still present during this time interval in other continents, as well: in North America with *Acrocanthosaurus* until the Albian (D’Emic et al., 2012), in Europe until the Cenomanian (Vullo et al., 2007; Csiki-Sava et al., 2015), and in Eastern Asia with *Shaochilong* until the Turonian (Brusatte et al., 2009, 2010b; see also Chure et al., 1999).

After dominating terrestrial ecosystems at least in Africa, South America and eastern Asia during the Albian–Turonian (Brusatte et al., 2009; Coria and Salgado, 2005; Novas et al., 2013), carcharodontosaurids were considered to disappear from the fossil record after the Turonian in both Asia (Brusatte et al., 2009) and South America (e.g., Coria and Salgado, 2005; Calvo et al., 2006; Novas et al., 2013), to be replaced by other groups of large theropods such as tyrannosaurids in parts of Laurasia and abelisaurids in parts of Gondwana. Canale et al. (2009) even cautioned against assigning isolated theropod teeth from post-Cenomanian deposits of South America to Carcharodontosauridae (e.g., Canudo et al., 2008; Casal et al., 2009; Salgado et al., 2009) due to their morphological similarity to those of the abelisaurid *Skorpiovenator*. Recently, however, more diagnostic cranial remains were reported to suggest the survival of carcharodontosaurids into the latest Cretaceous (Campanian–Maastrichtian) in Brazil (Azevedo et al., 2013).

Contrasting with this rich and relatively continuous fossil record of Carcharodontosauridae starting with the Aptian, the first half of its evolutionary history is very poorly documented (Fig. 5B). Prior to the identification of UAIC (SCM1) 615, only two

occurrences of pre-Aptian Cretaceous carcharodontosaurids were reported, one from the Barremian of Spain (Ortega et al., 2010; Gasca et al., 2014) and the other from the Barremian of Thailand (Buffetaut and Suteethorn, 2012). The Early Cretaceous *Kelmayisaurus* from Xinjiang, western China, was recognized as a carcharodontosaurid of possibly ?Valanginian to Aptian in age by Brusatte et al. (2012), but the deposits yielding these remains (the Lianmugui, or Lianmuxin, Formation of the Tugulu Group) were dated as Aptian–Albian by Eberth et al. (2001; see also Tong et al., 2009). An important temporal gap – of about 20 to 28 millions of years, according to the dates in Gradstein et al. (2012) – thus stretched between the oldest, tentatively assigned carcharodontosaurids from the Oxfordian–Tithonian of Tanzania, including the formally erected *Veterupristisaurus* (Rauhut, 2011; see also Carrano et al., 2012), and those that started to appear in the fossil record in the Barremian and then spread widely during the Aptian. Referral of UAIC (SCM1) 615 to Carcharodontosauridae partially fills this frustrating gap, effectively halving this shadowy period in the evolutionary history of the group.

Furthermore, our analyses tentatively cluster the Dobrogean theropod with the derived members of the Carcharodontosaurinae to the exclusion of the more basal, but significantly younger non-carcharodontosaurine carcharodontosaurids *Eocarcharia* and *Acrocanthosaurus*. If this placement is correct, then the Romanian tooth indicates that Carcharodontosaurinae diverged from other carcharodontosaurids considerably earlier than hitherto recognized.

The previously known fossil record of the clade suggested that Carcharodontosaurinae originated sometime between the Aptian and Albian, as basal carcharodontosaurids (*Acrocanthosaurus*, *Concavenator*, *Eocarcharia*) were moderately diverse in the Barremian–Aptian, followed by the appearance of many fossils of carcharodontosaurines beginning in the Albian (Fig. 5B). The proposed affinities of the oldest carcharodontosaurid material – including isolated teeth referred to as ‘*Megalosaurus*’ *ingens* – from the east African Upper

Jurassic, considered to be reminiscent of the Aptian–Albian *Acrocanthosaurus* (Rauhut, 2011), was also consistent with this evolutionary scenario. Now, our identification of UAIC (SCM1) 615 as a carcharodontosaurid dinosaur sharing important dental apomorphies with the derived Carcharodontosaurinae advocates the emergence of this clade (or at least the very large size and dental morphology characterizing it) well before the Albian, during or even before the Valanginian, and relegates taxa such as *Eocarcharia*, *Acrocanthosaurus* and *Concavenator* (the dentition of *Shaochilong* is unknown) as late-surviving members of the basal carcharodontosaurid radiation, with a relatively plesiomorphic dentition.

Besides shifting the emergence of the carcharodontosaurines earlier in time, identification of UAIC (SCM1) 615 as a carcharodontosaurid also has interesting palaeobiogeographic implications. As already noted, recent discoveries show that Carcharodontosauridae is not an endemic Gondwanan clade as was once proposed (e.g., Novas et al., 2005), with the identification of its widespread, Pangaeian distribution during the late Early Cretaceous (Sereno et al., 1996; Harris, 1998; Chure et al., 1999; Sereno, 1999; Brusatte and Sereno, 2008; Ortega et al., 2010; Brusatte et al., 2009, 2012; Mo et al., 2014). However, within Carcharodontosauridae itself, some palaeogeographic patterns have been widely accepted. For example, it has been widely acknowledged that Carcharodontosaurinae is a endemic subclade of Gondwanan carcharodontosaurids (e.g., Sereno 1999; Holtz et al., 2004b; Brusatte and Sereno, 2007; Sereno and Brusatte, 2008; Novas et al., 2013), as previously all its recognized members were restricted strictly to either Africa (Stromer, 1931; Sereno et al., 1996; Brusatte and Sereno, 2007) or South America (Coria and Salgado, 1995; Novas et al., 2005; Coria and Currie, 2006). Moreover, intra-clade relationships of Carcharodontosaurinae were still adhering to patterns of continental fragmentation and vicariant evolution, with a basal split between the Albian–Cenomanian African *Carcharodontosaurus* and the Giganotosaurini, uniting the similarly Albian–Cenomanian

southern South American *Giganotosaurus* and *Mapusaurus* (together with *Tyrannotitan*, if this taxon is also recovered within Carcharodontosaurinae; e.g., Novas et al., 2005, 2013).

This scenario is now challenged by our finding that the Southern Dobrogean carcharodontosaurid UAIC (SCM1) 615 may nest inside Carcharodontosaurinae. If true, such an affinity would suggest that the origin of Carcharodontosaurinae was not a southern, vicariant by-product of the Gondwana-Laurasia separation, a major palaeogeographic event that is considered to have been well underway by the end of the Jurassic, and essentially completed by the mid-Early Cretaceous (see Weishampel et al., 2010). Indeed, during this time palaeogeographic connections and faunal interactions were virtually non-existent between the northern Tethyan (European) and southern Tethyan (western Gondwanan, but essentially African) areas of the Mediterranean (e.g., Canudo et al., 2009), which makes a vicariant hypothesis intuitive. However, if the Romanian tooth represents a carcharodontosaurine, then it implies a much more complicated palaeogeographic history of the clade, which is not so clearly linked to continental breakup.

The palaeogeographic position of the Southern Dobrogean carcharodontosaurine in cratonic Europe, north of the Neo-Tethys, together with its significantly older age compared to other carcharodontosaurines, could indicate that separation of the carcharodontosaurine lineage took part in Europe and not in western Gondwana as previously assumed. This would also mean that representatives of this lineage were subsequently – after the Barremian – introduced to Africa and South America via trans-Tethyan dispersal, most probably at a time when faunal interactions between the southern and northern margins of the Mediterranean Tethys were resumed, after the early Barremian (Canudo et al., 2009).

Alternatively, it can be hypothesized that appearance of carcharodontosaurines in Southern Dobrogea is a consequence of southern immigration originating in western Gondwana, often considered the place of origin for this clade. However, this scenario has

several potential caveats. Although Europe has been considered as forming part of a larger Eurogondwanan palaeobioprovince during the early Early Cretaceous (Ezcurra and Agnolín, 2012), and occasional trans-Tethyan faunal connections have been recognized between Africa and Europe during Late Jurassic to Early Cretaceous times (e.g., Gheerbrant and Rage, 2006), these interchanges either pre-dated the Berriasian (e.g., Gardner et al., 2003; Knoll and Ruiz-Omeñaca, 2009), or post-dated the Barremian (Canudo et al., 2009; Torcida Fernández-Baldor et al., 2011), with no positive evidence for actual faunal exchanges taking place during the ‘Neocomian’ (Berriasian–Hauterivian) time interval.

More recently, some potential evidence has emerged for Gondwana-to-Europe interchange during the ‘Neocomian’. The presence of the basal rebbachisaurid *Histriasaurus* (Dalla Vecchia, 1998) in the upper Hauterivian–lower Barremian of Croatia has been cited as indicative of very early and very rapid northward dispersal of this clade from western Gondwana (southern South America; Carballido et al., 2012; Fanti et al., 2015). Timing of this particular dispersal event was even constrained to the Berriasian–Valanginian time interval (Fanti et al., 2015), which makes it roughly contemporaneous with the record of the Southern Dobrogean carcharodontosaurine. It was also suggested, however, that dispersal of the line leading to *Histriasaurus* was mediated by the northward drift of the Apulian Microplate (= Adria; see Bosselini, 2002), a continental sliver acting as a passive transportation mechanism (‘Noah’s Ark’; McKenna, 1973) for basal rebbachisaurids after its separation from mainland Africa (e.g., Torcida Fernández-Baldor et al., 2011). Furthermore, the palaeogeographical separation between Africa and Adria (and thus the effective movement of the presumed ark) is considered to be at most an incipient one during the Early Cretaceous by Bossellini (2002) and Zarcone et al. (2010), with spatial continuity still present between the two landmasses, while deep-water basins continued to separate Adria from the European Craton. Accordingly, although the presence of *Histriasaurus* can represent a case

of northward range extension of rebbachisaurids during the Berriasian–Valanginian, it took place not strictly speaking into Europe, but only reached the northernmost extremity of Adria, a northerly peninsular extension of the African mainland. It was only starting with the Barremian that rebbachisaurids dispersed as far north as the European cratonic areas, including Iberia and the British Isles (Mannion, 2009; Mannion et al., 2011; Torcida Fernández-Baldor et al., 2011), a time when faunal interchanges between Europe and Africa are considered to have been well underway (e.g., Gheerbrant and Rage, 2006; Canudo et al., 2009).

Unlike *Histriasaurus*, the taxon represented by UAIC (SCM1) 615 was an inhabitant of the European mainland. It is thus unclear to what extent the example of rebbachisaurid range extension into (present-day) Europe during the early Early Cretaceous, as potentially testified by the discovery of the Croatian taxon, would also be applicable for the Southern Dobrogean carcharodontosaurine. The available evidence suggests that these two cases are very different, and that faunal connections during this time interval are not documented between the African and European cratons as already pointed out by Gheerbrant and Rage (2006).

Absence of documented faunal interactions weakens support for a scenario of south-to-north immigration of derived carcharodontosaurines in Europe at the very beginning of the Cretaceous, and would argue instead for a local, European development to explain the presence of a Valanginian carcharodontosaurine in Southern Dobrogea. The pre-Barremian presence of carcharodontosaurids in Europe is also consistent with their appearance in the Barremian–Aptian fossil record of Eastern Asia, with Europe acting as a stepping stone in the eastward dispersal of the clade. Similarly, the presence of Aptian carcharodontosaurids in North America likely requires the presence of pre-Aptian members of the clade in Europe, since faunal exchanges between these two landmasses are known to have been halted before

the Aptian (e.g., Kirkland et al., 1999). Interestingly, it appears that only basal carcharodontosaurids were able to spread into the northern Laurasian landmasses, while the derived carcharodontosaurines dispersed exclusively across the Neo-Tethys, into western Gondwana. The causes of these distribution patterns remain as yet unknown, and further support – in the form on new carcharodontosaurid discoveries from the early-middle part of the Early Cretaceous – is required to better uphold such a scenario.

We finally reiterate that if the Romanian tooth does not belong to a carcharodontosaurine, but instead is artefactually grouping with them in the phylogenetic analysis because of the very incomplete nature of the material, then the traditional story of Carcharodontosaurinae as a product of vicariant evolution driven by the breakup of Pangea will remain strongly supported. However, even in such case UAIC (SCM1) 615 would still record the presence of early-occurring large carcharodontosaurid theropods with a very characteristic carcharodontosaurine-type dentition in the eastern part of the European craton, adding to known early Early Cretaceous theropod (and dinosaur) diversity, and potentially documenting dinosaur faunal provinciality in Europe and worldwide.

6. Conclusions

We re-describe and interpret the affinities of one of the most significant historical dinosaurian specimens of Romania, an isolated but well-preserved theropod tooth from Southern Dobrogea. Our extensive analyses suggest carcharodontosaurid relationships for this tooth, while the available evidence – including novel calcareous nannoplankton sampling – supports its Valanginian age. The Southern Dobrogean theropod tooth represents the oldest record of Carcharodontosauridae in the Cretaceous, and the second oldest globally, eclipsed only by a collection of isolated specimens from the Upper Jurassic of eastern Africa. As one of the only two known Valanginian dinosaurian occurrences from Central and Eastern Europe, this

record advances our understanding of European dinosaur distribution during the early Early Cretaceous, and also fills an important palaeogeographic gap between Western European and Eastern Asian dinosaurian assemblages of the Valanginian.

Based on dental apomorphies, our analyses further identify UAIC (SCM1) 615 as a possible member of Carcharodontosaurinae, a subclade of derived and gigantic carcharodontosaurids formerly known to be restricted to the Albian–Cenomanian of western Gondwana (Africa and South America). If this finding is correct, the Southern Dobrogean specimen documents the emergence of Carcharodontosaurinae earlier than previously recognized, thus also indicating an earlier acquisition of their characteristically large size.

Based on currently known palaeogeographic and chronostratigraphic constraints on the evolution of Carcharodontosauridae, it appears that not only did this clade have a wide distribution, but that crucial events of its evolutionary history such as the emergence of the derived carcharodontosaurines took place north of the Tethys, in cratonic Europe, instead of western Gondwana and as the result of vicariant evolution driven by the Gondwana-Laurasia split, as was formerly suggested. In such a case, instead of endemic evolution the emergence of the western Gondwanan mid-Cretaceous carcharodontosaurines was the result of a north-to-south trans-Tethyan dispersal that took place somewhere between the Valanginian and the Aptian. Recognizing a potential carcharodontosaurine dispersal event from Europe into western Gondwana adds further support for the presence of important palaeogeographic ties between the two realms during the second half of the Early Cretaceous.

Acknowledgements

This research was supported by the National Research Council of Romania (CNCS) grant PN-IIID-PCE-2011-3-0381 and a Sepkoski grant of the Paleontological Society for Z.Cs.-S.

S.L.B. is supported by a Marie Curie Career Integration Grant EC630652, the Division of Paleontology of the American Museum of Natural History, and the School of GeoSciences of the University of Edinburgh. He thanks Mátyás Vremir, Radu Totoianu, and Mark Norell for many hours of fun discussion on Romanian fossils, and for supporting his work and travel in Romania. We thank Mihai Brânzilă and Paul Țibuleac (UAIC) for access to the specimen, for allowing us to collect samples for the nannoplankton analyses, and for their help and collegiality during our visit to Iași, as well as Ilie Turculeț for sharing information about the history of the specimen. Mihaela C. Melinte-Dobrinescu has gracefully analyzed the nannoplankton samples derived from UAIC (SCM1) 615; her contribution was essential in assessing the age of the specimen. Finally, we thank the reviewers Eric W.A. Mulder (Denekamp, the Netherlands) and Xabier Pereda-Suberbiola (Bilbao, Spain), as well as Associated Editor Elena Jagt-Yazykova, for their useful comments and suggestions that helped improve previous versions of the manuscript.

References

- Accarie, H., Beaudoin, B., Dejax, J., Friès, G., Michard, J. G., & Taquet, P. (1995). Découverte d'un dinosaure théropode nouveau (*Genusaurus sisteronis* n. g., n. sp.) dans l'Albien marin de Sisteron (Alpes-de-Haute-Provence, France) et extension au Crétacé inférieur de la lignée cératosaurienne. *Comptes Rendus de l'Académie des Sciences Paris, IIa* 320, 327–334.
- Allain, R. (2002). Discovery of a megalosaur (Dinosauria, Theropoda) in the Middle Bathonian of Normandy (France) and its implications for the phylogeny of basal Tetanurae. *Journal of Vertebrate Paleontology*, 22, 548–563.

- Allain, R. (2005). The enigmatic theropod dinosaur *Erectopus superbus* (Sauvage, 1882) from the Lower Albian of Louppy-le-Chateau (Meuse, France). In K. Carpenter (Ed.), *The Carnivorous Dinosaurs* (pp. 72–86). Bloomington: Indiana University Press.
- Avram, E., Costea, I., Dragastan, O., Muşiu, R., Neagu, T., Şindilar, V., & Vinogradov, C. (1996). Distribution of the Middle-Upper Jurassic and Cretaceous facies in the Romanian eastern part of the Moesian Platform. *Revue Roumaine de Géologie*, 39-40, 3–33.
- Avram, E., Szasz, L., Antonescu, E., Baltreş, A., Iva, M., Melinte, M., Neagu, T., Rădan, S., & Tomescu, C. (1993). Cretaceous terrestrial and shallow marine deposits in northern South Dobrogea (SE Romania). *Cretaceous Research*, 14, 265–305.
- Azevedo, R. P. F. de, Simbras, F. M., Furtado, M. R., Candeiro, C. R. A., & Bergqvist, L. P. (2013). First Brazilian carcharodontosaurid and other new theropod dinosaur fossils from the Campanian-Maastrichtian Presidente Prudente Formation, São Paulo State, southeastern Brazil. *Cretaceous Research*, 40, 131–142.
- Benson, R. B. J., Barrett, P. M., Powell, H. P., & Norman, D. B. (2008). The taxonomic status of *Megalosaurus bucklandii* (Dinosauria, Theropoda) from the Middle Jurassic of Oxfordshire, UK. *Palaeontology*, 51, 419–424.
- Benson, R. B. J., Carrano, M. T., & Brusatte, S. L. (2010). A new clade of archaic large-bodied predatory dinosaurs (Theropoda: Allosauroidae) that survived to the latest Mesozoic. *Naturwissenschaften*, 97, 71–78.
- Benton, M. J., Cook, E., Grigorescu, D., Popa, E., & Tallódi, E. (1997). Dinosaurs and other tetrapods in an Early Cretaceous bauxite-filled fissure, northwestern Romania. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 130, 275–292.
- Benton, M. J., Csiki, Z., Grigorescu, D., Redelstorff, R., Sander, M., Stein, K., & Weishampel, D. B. (2010). Dinosaurs and the island rule: the dwarfed dinosaurs from Haţeg Island. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 293, 438–454.

- Benton, M. J., Minter, N. J., & Posmoşanu, E. (2006). Dwarfing in ornithopod dinosaurs from the Early Cretaceous of Romania. In Z. Csiki (Ed.), *Mesozoic and Cenozoic Vertebrates and Paleoenvironments; Tributes to the Career of Prof. Dan Grigorescu* (pp. 79–87). Bucharest: Ars Docendi.
- Blows, W. T., & Honeysett, K. (2014). First Valanginian *Polacanthus foxii* (Dinosauria, Ankylosauria) from England, from the Lower Cretaceous of Bexhill, Sussex. *Proceedings of the Geologists' Association*, 125, 233–251.
- Bosselini, A. (2002). Dinosaurs “re-write” the geodynamics of the eastern Mediterranean and the paleogeography of the Apulia Platform. *Earth-Science Reviews*, 59, 211–234.
- Brânzilă, M. (ed.) (2010). *Academicianul Ion Simionescu - savant şi dascăl al neamului* (p. 182). Iaşi: Ed. Universităţii "Alexandru Ioan Cuza".
- Brusatte, S., Benson, R. B. J., Carr, T. D., Williamson, T. E., & Sereno, P. C. (2007). The systematic utility of theropod enamel wrinkles. *Journal of Vertebrate Paleontology*, 27, 1052–1056.
- Brusatte, S. L., Benson, R. B. J., Chure, D. J., Xu, X., Sullivan, C., & Hone, D. W. E. (2009). The first definitive carcharodontosaurid (Dinosauria: Theropoda) from Asia and the delayed ascent of tyrannosaurids. *Naturwissenschaften*, 96, 1051–1058.
- Brusatte, S. L., Benson, R. B. J., & Xu, X. (2012). A reassessment of *Kelmayisaurus petrolicus*, a large theropod dinosaur from the Early Cretaceous of China. *Acta Palaeontologica Polonica*, 57, 65–72.
- Brusatte, S. L., & Clark, N. D. L. (2015). Theropod dinosaurs from the Middle Jurassic (Bajocian-Bathonian) of Skye, Scotland. *Scottish Journal of Geology*, 51, 157–164. doi: 10.1144/sjg2014-022

- Brusatte, S., Chure, D. J., Benson, R. B. J., & Xu, X. (2010b). The osteology of *Shaochilong maortuensis*, a carcharodontosaurid (Dinosauria: Theropoda) from the Late Cretaceous of Asia. *Zootaxa*, 2334, 1–46.
- Brusatte, S. L., Lloyd, G. T., Wang, S. C., & Norell, M. A. (2014). Gradual assembly of avian body plan culminated in rapid rates of evolution across the dinosaur-bird transition. *Current Biology*, 24, 2386–2392.
- Brusatte, S. L., Norell, M. A., Carr, T. D., Erickson, G. M., Hutchinson, J. R., Balanoff, A. M., Bever, G. S., Choiniere, J. N., Makovicky, P. J., & Xu, X. (2010a). Tyrannosaur paleobiology: new research on ancient exemplar organisms. *Science*, 329, 1481–1485.
- Brusatte, S. L., & Sereno, P. C. (2007). A new species of *Carcharodontosaurus* (Dinosauria: Theropoda) from the Cenomanian of Niger and a revision of the genus. *Journal of Vertebrate Paleontology*, 27, 902–916.
- Brusatte, S. L., & Sereno, P. C. (2008). Phylogeny of Allosauroidae (Dinosauria: Theropoda): comparative analysis and resolution. *Journal of Systematic Palaeontology*, 6, 155–182.
- Brusatte, S. L., Vremir, M., Csiki-Sava, Z., Turner, A. H., Watanabe, A., Erickson, G. M., & Norell, M. A. (2013). The osteology of *Balaur bondoc*, an island-dwelling dromaeosaurid (Dinosauria: Theropoda) from the Late Cretaceous of Romania. *Bulletin of the American Museum of Natural History*, 374, 3–100. doi: 10.1206/798.1
- Buckland, W. (1824). Notice on the *Megalosaurus* or great fossil lizard of Stonesfield. *Transactions of the Geological Society*, 21, 390–397.
- Buffetaut, E., & Suteethorn, V. (1998). Early Cretaceous dinosaurs from Thailand and their bearing on the early evolution and biogeographical history of some groups of Cretaceous dinosaurs. In S. G. Lucas, J. I. Kirkland, & J. W. Estep (Eds.), *Lower and Middle Cretaceous*

- 1093 *Terrestrial Ecosystems* (pp. 205–210), New Mexico Museum of Natural History and Science
1
2 1094 Bulletin, 14.
3
4
5 1095 Buffetaut, E., & Suteethorn, V. (2007). A sinraptorid theropod (Dinosauria: Saurischia) from
6
7 1096 the Phu Kradung Formation of northeastern Thailand. *Bulletin de la Société Géologique de*
8
9 1097 *France*, 178, 497–502.
10
11
12 1098 Buffetaut, E., & Suteethorn, V. (2012). A carcharodontid theropod (Dinosauria, Saurischia)
13
14 1099 from the Sao Khua Formation (Early Cretaceous, Barremian) of Thailand. In R. Royo-Torres,
15
16 F. Gascó, & L. Alcalá (Eds.), *10th Annual Meeting of the European Association of Vertebrate*
17 1100 *Palaeontologists* (pp. 27–30). Teruel: ¡Fundamental!, 20.
18
19 1101
20
21 1102 Calvo, J. O., & Coria, A. (1998). New specimen of *Giganotosaurus carolinii* (Coria &
22
23 Salgado, 1995), supports it as the largest theropod ever found. *Gaia*, 15, 117–122.
24 1103
25
26 1104 Calvo, J. O., Gandossi, P., & Porfiri, J. D. (2006). Dinosaur faunal replacement during
27
28 Cenomanian times in Patagonia, Argentina. In: S. E. Evans, & P. M. Barrett (Eds.), *9th*
29 1105 *Mesozoic Terrestrial Ecosystems and Biota* (pp. 17–20). Manchester, UK.
30
31 1106
32
33 1107 Canale, J. I., Scanferla, C. A., Agnolin, F. L., & Novas, F. E. (2009). New carnivorous
34
35 dinosaur from the Late Cretaceous of NW Patagonia and the evolution of abelisaurid
36 1108
37 theropods. *Naturwissenschaften*, 96, 409–414.
38
39 1109
40
41 1110 Canale, J. I., Novas, F. E., & Pol, D. (2015). Osteology and phylogenetic relationships of
42
43 *Tyrannotitan chubutensis* Novas, de Valais, Vickers-Rich and Rich, 2005 (Theropoda:
44 1111
45 Carcharodontosauridae) from the Lower Cretaceous of Patagonia, Argentina. *Historical*
46 1112
47 *Biology*, 27, 1–32. doi: 10.1080/08912963.2013.861830.
48
49 1113
50
51 1114 Candeiro, C. R. A., Fanti, F., Therrien, F., & Lamanna, M. C. (2011). Continental fossil
52
53 vertebrates from the mid-Cretaceous (Albian-Cenomanian) Alcântara Formation, Brazil, and
54 1115
55 their relationship with contemporaneous faunas from North Africa. *Journal of African Earth*
56 1116
57 *Sciences*, 60, 79–92.
58 1117
59
60
61
62
63
64
65

- Canudo, J. I. Barco, J. L., Pereda Suberbiola, X., Ruiz-Omeñaca, J. I., Salgado, L., Torcida
- Fernández-Baldor, F., & Gasulla, J. M. (2009). What Iberian dinosaurs reveal about the
bridge said to exist between Gondwana and Laurasia in the Early Cretaceous. *Bulletin de la
Société Géologique de France*, 180, 5–11.
- Canudo, J. I., Filippi, L. S., Salgado, L., Garrido, A. C., Cerda, I. A., Garcia, R., & Otero, A.
(2008). Theropod teeth associated with a sauropod carcass in the Upper Cretaceous (Plottier
Formation) of Rincón de los Sauces. In Colectivo Arqueológico y Paleontológico de Salas de
los Infantes (Ed.), *Actas de las IV Jornadas Internacionales sobre Paleontología de
Dinosaurios y su Entorno* (pp. 321–330). Salas de los Infantes, Burgos.
- Canudo, J. I., & Ruiz-Omeñaca, J. I. (2003). Los restos directos de dinosaurios terópodos
(excluyendo Aves) en España. *Ciencias de la Tierra* 26, 347–373.
- Carballido, J. L., Salgado, L., Pol, D., Canudo, J. I., & Garrido, A. (2012). A new basal
rebbachisaurid (Sauropoda, Diplodocoidea) from the Early Cretaceous of the Neuquén Basin;
evolution and biogeography of the group. *Historical Biology*, 24, 631–654.
- Carrano, M. T., Benson, R. B. J., & Sampson, S. D. (2012). The phylogeny of Tetanurae
(Dinosauria: Theropoda). *Journal of Systematic Palaeontology*, 10, 211–300.
- Carrano, M. T., & Sampson, S. D. (2008). The phylogeny of Ceratosauria (Dinosauria:
Theropoda). *Journal of Systematic Palaeontology*, 6, 183–236.
- Carvalho, I. S., Campos, A. C. A., & Nobre, P. H. (2005). *Baurusuchus salgadoensis*, a new
Crocodylomorpha from the Bauru Basin (Cretaceous), Brazil. *Gondwana Research*, 8, 11–30.
- Casal, G., Candeiro, C. R. A., Martinez, R., Ivany, E., & Ibiricu, L. (2009). Dientes de
Theropoda (Dinosauria: Saurischia) de la Formación Bajo Barreal, Cretácico Superior,
Provincia del Chubut, Argentina. *Géobios*, 42, 553–560.

- 1141 Cau, A., Dalla Vecchia, F. M., & Fabbri, M. (2013). A thick-skulled theropod (Dinosauria,
1
2 1142 Saurischia) from the Upper Cretaceous of Morocco with implications for carcharodontosaurid
3
4
5 1143 cranial evolution. *Cretaceous Research*, 40, 251–260.
6
7 1144 Choiniere, J. N., Forster, C. A., & De Klerk, W. J. (2012). New information on
8
9
10 1145 *Nqwebasaurus thwazi*, a coelurosaurian theropod from the Early Cretaceous Kirkwood
11
12 1146 Formation in South Africa. *Journal of African Earth Sciences*, 71–72, 1–17.
13
14
15 1147 Chure, D. J., Manabe, M., Tanimoto, M., & Tomida, Y. (1999). An unusual theropod tooth
16
17 1148 from the Mifune Group (Late Cenomanian to Early Turonian), Kumamoto, Japan. In Y.
18
19 1149 Tomida, T. H. Rich, & P. Vickers-Rich (Eds.), *Proceedings of the Second Gondwanan*
20
21
22 1150 *Dinosaur Symposium* (pp. 291–296). Tokyo: National Science Museum Monographs, 15.
23
24 1151 Cifelli, R. L., Davis, B. M., & Sames, B. (2014). Earliest Cretaceous mammals from the
25
26 1152 western United States. *Acta Palaeontologica Polonica*, 59, 31–52.
27
28
29 1153 doi:10.4202/app.2012.0089.
30
31
32 1154 Cobos, A., Lockley, M. G., Gascó, F., Royo-Torres, R., & Alcalá, L. (2014). Megatheropods
33
34 1155 as apex predators in the typically Jurassic ecosystems of the Villar del Arzobispo Formation
35
36 1156 (Iberian Range, Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 399, 31–41.
37
38
39 1157 doi: 10.1016/j.palaeo.2014.02.008.
40
41 1158 Cobos, A., Royo-Torres, R., Luque, L., Alcalá, L., & Mampel, L. (2010). An Iberian
42
43
44 1159 stegosaurs paradise: The Villar del Arzobispo Formation (Tithonian–Berriasian) in Teruel
45
46 1160 (Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 293, 223–236.
47
48
49 1161 Codrea, V., Godefroit, P., & Smith, T. (2012). First discovery of Maastrichtian (latest
50
51 1162 Cretaceous) terrestrial vertebrates in Rusca Montană Basin (Romania). In P. Godefroit (Ed.)
52
53 1163 *Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems* (pp. 570–581).
54
55
56 1164 Bloomington: Indiana University Press.
57
58
59
60
61
62
63
64
65

- 1165 Codrea, V., Vremir, M., Jipa, C., Godefroit, P., Csiki, Z., Smith, T., & Fărcaș, C. (2010).
1166 More than just Nopcsa's Transylvanian dinosaurs: A look outside the Hațeg Basin.
1167 *Palaeogeography, Palaeoclimatology, Palaeoecology* 293, 391–405. doi:
1168 10.1016/j.palaeo.2009.10.027.
- 1169 Colbert, E. H. (1946). *Sebecus*, representative of a peculiar suborder of fossil Crocodilia from
1170 Patagonia. *Bulletin of the American Museum of Natural History*, 87(4), 217–270.
- 1171 Coria, R. A., & Currie, P. J. (2006). A new carcharodontosaurid (Dinosauria, Theropoda)
1172 from the Upper Cretaceous of Argentina. *Geodiversitas*, 28, 71–118.
- 1173 Coria, R. A., & Salgado, L. (1995). A new giant carnivorous dinosaur from the Cretaceous of
1174 Patagonia. *Nature*, 377, 224–226.
- 1175 Coria, R. A., & Salgado, L. (2005). Mid-Cretaceous turnover of saurischian dinosaur
1176 communities: evidence from the Neuquén Basin. In G. D. Veiga, L. A. Spalletti, J. A.
1177 Howell, & E. Schwartz (Eds.), *The Neuquén Basin, Argentina: a case study in sequence*
1178 *stratigraphy and basin dynamics* (pp. 317–327). Geological Society, London, Special
1179 Publications, 252.
- 1180 Covacef, Z. (1995). Pionieri ai culturii românești în Dobrogea; Ioan Cotovu și Vasile Cotovu.
1181 *Analele Dobrogei, I(1)*, 127–134.
- 1182 Császár, G., Kollányi, K., Lantos, M., Lelkes, G., & Tardine Filácz, E. (2000). A
1183 Hidasivölgyi Márga Formáció kora és képződési környezete. *Földtani Közlöny*, 130, 695–
1184 723.
- 1185 Csiki, Z., & Grigorescu, D. (1998). Small theropods of the Late Cretaceous of the Hațeg
1186 Basin (Western Romania) - an unexpected diversity at the top of the food chain. *Oryctos*, 1,
1187 87–104.

- 1188 Csiki, Z., Vremir, M., Brusatte, S. L., & Norell, M. A. (2010). An aberrant island-dwelling
theropod dinosaur from the Late Cretaceous of Romania. *Proceedings of the National
Academy of Sciences*, 107, 15357–15361.
- 1191 Csiki-Sava, Z., Buffetaut, E., Ősi, A., Pereda-Suberbiola, X., & Brusatte, S. L. (2015). Island
life in the Cretaceous - faunal composition, biogeography, evolution, and extinction of land-
living vertebrates on the Late Cretaceous European archipelago. *Zookeys*, 469, 1–161. doi:
10.3897/zookeys.469.8439.
- 1195 Csiki-Sava, Z., Codrea, V., & Vasile, Ș. (2013). Early Cretaceous dinosaur remains from
Dobrogea (southeastern Romania). In L. Picot (Ed.), *Abstracts, 11th Annual Meeting of the
European Association of Vertebrate Palaeontologists* (pp. 28). Villers-sur-Mer, France.
- 1198 Csontos, L., & Vörös, A. (2004). Mesozoic plate tectonic reconstruction of the Carpathian
region. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 210, 1–56. doi:
10.1016/j.palaeo.2004.02.033.
- 1201 Currie, P. J., & Carpenter, K. (2000). A new specimen of *Acrocanthosaurus atokensis*
(Theropoda, Dinosauria) from the Lower Cretaceous Antlers Formation (Lower Cretaceous,
Aptian) of Oklahoma, USA. *Geodiversitas*, 22, 207–246.
- 1204 Currie, P. J., Rigby, J. K., & Sloan, R. E. (1990). Theropod teeth from the Judith River
Formation of southern Alberta, Canada. In K. Carpenter, & P. J. Currie (Eds.), *Dinosaur
Systematics: Approaches and Perspectives* (pp. 107–125). Cambridge, Boston: Cambridge
University Press.
- 1208 Dalla Vecchia, F. M. (1998). Remains of Sauropoda (Reptilia, Saurischia) in the Lower
Cretaceous (Upper Hauterivian/Lower Barremian) limestones of SW Istria (Croatia).
Geologia Croatica, 5, 105–134.
- 1211 Dames, W. (1884). *Megalosaurus dunkeri*. *Sitzungsberichte Gesellschaft Naturforschender
Freunde zu Berlin*, 1884, 186–188.

- De Klerk, W. J., Forster, C. A., Sampson, S. D., Chinsamy, A., & Ross, C. F. (2000). A new coelurosaurian dinosaur from the Early Cretaceous of South Africa. *Journal of Vertebrate Paleontology*, 20, 324–332.
- D’Emic, M. D., Melstrom, K. M., & Eddy, D. R. (2012). Paleobiology and geographic range of the large-bodied Cretaceous theropod dinosaur *Acrocanthosaurus atokensis*. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 333–334, 13–23.
- Dercourt, J., Gaetani, M., Vrielynck, B., Barrier, E., Biju-Duval, B., Brunet, M., Cadet, J. P., Crasquin, S., & Săndulescu, M. (Eds.) (2000). *Atlas Peri-Tethys Palaeogeographical Maps* (269 p). Paris: CCGM/CGMW .
- Dinu, C., Grădinaru, E., Stoica, M., & Diaconescu, V. (2007). *Dobrogea 2007 Field Trip Preparation and Assistance* (123 p.). University of Bucharest.
- Dragastan, O. N., Antoniadu, C., & Stoica, M. (2014). Biostratigraphy and zonation of the Lower Cretaceous succession from Cernavodă-lock section, South Dobrogea, eastern part of the Moesian Platform (Romania). *Carpathian Journal of Earth and Environmental Sciences*, 9, 231–260.
- Dragastan, O., Neagu, T., Bărbulescu, A., & Pană, I. (1998). *Jurassicul și Cretacicul din Dobrogea Centrală și de Sud* (249 p.). Bucharest.
- Dyke, G. J., Benton, M. J., Posmoșanu, E., & Naish, D. (2011). Early Cretaceous (Berriasian) birds and pterosaurs from the Cornet bauxite mine, Romania. *Palaeontology*, 54, 79–95.
- Eberth, D. A., Brinkman, D. B., Chen, P.-J., Yuan, F.-T., Wu, X.-C., Li, G., & Cheng, X.-S. (2001). Sequence stratigraphy, paleoclimate patterns, and vertebrate fossil preservation in Jurassic–Cretaceous strata of the Juggar Basin, Xinjiang Autonomous Region, People’s Republic of China. *Canadian Journal of Earth Sciences*, 38, 1627–1644.

- Eddy, D. R., & Clarke, J. A. (2011). New information on the cranial anatomy of *Acrocanthosaurus atokensis* and its implications for the phylogeny of Allosauroidae (Dinosauria: Theropoda). *PLoS ONE*, 6(3), e17932. doi:10.1371/journal.pone.0017932.
- Evans, S. E., & Matsumoto, R. (2015). An assemblage of lizards from the Early Cretaceous of Japan. *Palaeontologia Electronica*, 18.2.36A, 1–36.
- Ezcurra, M. D., & Agnolín, F. L. (2012). A new global palaeobiogeographical model for the Late Mesozoic and Early Tertiary. *Systematic Biology*, 61, 553–566. doi:10.1093/sysbio/syr115.
- Fanti, F., Cau, A., Cantelli, L., Hassine, M., & Audatore, M. (2015). New information on *Tataouinea hannibalis* from the Early Cretaceous of Tunisia and implications for the tempo and mode of rebbachisaurid sauropod evolution. *PLoS ONE*, 10(4), e0123475. doi:10.1371/journal.pone.0123475.
- Galton, P. M. (2009). Notes on Neocomian (Lower Cretaceous) ornithomimid dinosaurs from England - *Hypsilophodon*, *Valdosaurus*, “*Camptosaurus*”, “*Iguanodon*” - and referred specimens from Romania and elsewhere. *Revue de Paléobiologie*, 28, 211–273.
- Gardner, J. D., Evans, S. E., & Sigogneau-Russell, D. (2003). New albanerpetontid amphibians from the Early Cretaceous of Morocco and Middle Jurassic of England. *Acta Palaeontologica Polonica*, 48, 301–319.
- Gasca, J. M., Canudo, J. I., & Moreno-Azanza, M. (2014). A large-bodied theropod (Tetanurae: Carcharodontosauria) from the Mirambel Formation (Barremian) of Spain. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 273, 13–23. doi: 10.1127/0077-7749/2014/0413.
- Gheerbrant, E., & Rage, J.-C. (2006). Paleobiogeography of Africa: How distinct from Gondwana and Laurasia? *Palaeogeography, Palaeoclimatology, Palaeoecology*, 241, 224–246. doi:10.1016/j.palaeo.2006.03.016.

- Gishlick, A. D., & Gauthier, J. A. (2007). On the manual morphology of *Compsognathus longipes* and its bearing on the diagnosis of Compsognathidae. *Zoological Journal of the Linnean Society*, 149, 569–581.
- Goloboff, P. A., Farris, J. S., & Nixon, K. C. (2008). TNT, a free program for phylogenetic analysis. *Cladistics*, 24, 774–786.
- Gradstein, F. M., Ogg, J. G., Schmitz, M. D., & Ogg, G. M. (Eds.) (2012). *The Geologic Time Scale 2012* (1144 p.). Amsterdam: Elsevier.
- Grigorescu, D. (1992). Nonmarine Cretaceous Formations of Romania. In N. J. Matter, & P.-J. Chen (Eds.), *Aspects of Nonmarine Cretaceous Geology* (pp. 142–164). Beijing: China Ocean Press .
- Grigorescu, D. (2003). Dinosaurs of Romania. *Comptes Rendus Palevol*, 2, 97–101.
- Grigorescu, D. (2010). The Latest Cretaceous fauna with dinosaurs and mammals from the Hațeg Basin — A historical overview. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 293, 271–282.
- Hammer, O., Harper, D. A. T., & Ryan, P. D. (2001). Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica*, 4, 1–9.
- Harris, J. D. (1998). A reanalysis of *Acrocanthosaurus atokensis*, its phylogenetic status, and paleobiogeographic implications, based on a new specimen from Texas. *New Mexico Museum of Natural History and Science Bulletin*, 13, 1–75.
- Hendrickx, C., & Mateus, O. (2014). Abelisauridae (Dinosauria: Theropoda) from the Late Jurassic of Portugal and dentition-based phylogeny as a contribution for the identification of isolated theropod teeth. *Zootaxa*, 3751(1), 1–74.
- Hendrickx, C., Mateus, O., & Araújo, R. (2015a). The dentition of megalosaurid theropods. *Acta Palaeontologica Polonica*, 60, 627–642. doi:10.4202/app.00056.2013.

- Hendrickx, C., Mateus, O., & Araújo, R. (2015b). A proposed terminology of theropod teeth (Dinosauria, Saurischia). *Journal of Vertebrate Paleontology*, 35(5), e982797. doi: 10.1080/02724634.2015.982797.
- Holtz, T. R., Chapman, R. E., & Lamanna, M. C. (2004b). Mesozoic biogeography of Dinosauria. In D. B. Weishampel, P. Dodson, & H. Osmólska (Eds.), *The Dinosauria. Second Edition* (pp. 627–642). Berkeley, Los Angeles, London: University of California Press.
- Holtz, T. R., Jr., Molnar, R. E., & Currie, P. J. (2004a). Basal Tetanurae. In D. B. Weishampel, P. Dodson, & H. Osmólska (Eds.), *The Dinosauria. Second Edition* (pp. 71–110). Berkeley, Los Angeles, London: University of California Press.
- Huene, F. von (1923). Carnivorous Saurischia in Europe since the Triassic. *Bulletin of the Geological Society of America*, 34, 449–458.
- Hippolyte, J.-C. (2002). Geodynamics of Dobrogea (Romania): new constraints on the evolution of the Tornquist–Teisseyre Line, the Black Sea and the Carpathians. *Tectonophysics*, 357, 33–53.
- Ionesi, L. (1994). *Geologia unităților de platformă și a orogenului Nord-Dobrogean* (p. 280). Bucharest: Ed. Tehnică.
- Jerzykiewicz, T., & Russell, D. A. (1991). Late Mesozoic stratigraphy and vertebrates of the Gobi Basin. *Cretaceous Research*, 12, 345–377.
- Jurcsák, T. (1982). Occurrences nouvelles des Sauriens mésozoïques de Roumanie. *Vertebrata Hungarica*, 21, 175–184.
- Jurcsák, T., & Popa, E. (1979). Dinozaurieni ornithopozi din bauxitele de la Cornet (Munții Pădurea Craiului). *Nymphaea*, 7, 37–75.

- Jurcsák, T., & Popa, E. (1983). La faune de dinosauriens du Bihor (Roumanie). In E. Buffetaut, J. M. Mazin, E. Salmon (Eds.), *Actes du Symposium Paléontologique Georges Cuvier* (pp. 325–335). Montbéliard: Le Serpentaire.
- Kellner, A. W. A., Pinheiro, A. E. P., & Campos, D. A. (2014). A new sebecid from the Paleogene of Brazil and the crocodyliform radiation after the K–Pg boundary. *PLoS ONE*, 9(1), e81386. doi:10.1371/journal.pone.0081386.
- Kirkland, J. I., Cifelli, R. L., Britt, B. B., Burge, D. L., DeCourten, F. L., Eaton, J. G., & Parrish, J. M. (1999). Distribution of vertebrate faunas in the Cedar Mountain Formation, east-central Utah. *Utah Geological Survey Miscellaneous Publication*, 99-1, 201–217.
- Knoll, F., & Ruiz-Omeñaca, J. I. (2009). Theropod teeth from the basalmost Cretaceous of Anoual (Morocco) and their palaeobiogeographical significance. *Geological Magazine*, 146, 602–616.
- Kusuhashi, N., Matsumoto, A., Murakami, M., Tagami, T., Hirata, T., Iizuka, T., Handa, T., & Matsuoka, H. (2006). Zircon U–Pb ages from tuff beds of the upper Mesozoic Tetori Group in the Shokawa district, Gifu Prefecture, central Japan. *The Island Arc*, 15, 378–390.
- Lacasa Ruiz, A. (1989). Nuevo genero de ave fosil del yacimiento Neocomiense del Montsec (Provincia de Lerida, España). *Estudios geológicos*, 45(5-6), 417–425.
- Larson, D. W., & Currie, P. J. (2013). Multivariate analyses of small theropod dinosaur teeth and implications for paleoecological turnover through time. *PLoS ONE*, 8(1), e54329. doi:10.1371/journal.pone.0054329.
- Le Loeuff, J., & Buffetaut, E. (1995). The evolution of Late Cretaceous non-marine vertebrate faunas in Europe. In A.-L. Sun, & Y.-Q. Wang (Eds.), *Sixth Symposium on Mesozoic Terrestrial Ecosystems and Biota, Short Papers* (pp. 181–184). Beijing: China Ocean Press.

- 1332 Le Loeuff, J., Lang, E., Cavin, L., & Buffetaut, E. (2012). Between Tendaguru and Bahariya:
1333 on the age of the Early Cretaceous dinosaur sites from the Continental Intercalaire and other
1334 African formations. *Journal of Stratigraphy*, 36, 486–502.
- 1335 Lucas, F. A. (1901). A new dinosaur, *Stegosaurus marshi*, from the Lower Cretaceous of
1336 South Dakota. *Proceedings of the United States National Museum*, 23(1224), 591–592.
- 1337 Lucas, S. G. (2006). The *Psittacosaurus* biochron, Early Cretaceous of Asia. *Cretaceous*
1338 *Research*, 27, 189–198.
- 1339 Lucas, S. G., & Estep, J. W. (1998). Vertebrate biostratigraphy and biochronology of the
1340 Cretaceous of China. In S. G. Lucas, J. I. Kirkland, J. W. Estep (Eds.), *Lower and Middle*
1341 *Cretaceous Terrestrial Ecosystems* (pp. 1–20). New Mexico Museum of Natural History and
1342 Science Bulletin, 14.
- 1343 Lü, J.-C., Xu, L., Pu, H.-Y., Jia, S.-H., Azuma, Y., Chang, H.-L., & Zhang, J.-M. (2016).
1344 Paleogeographical significance of carcharodontosaurid teeth from the late Early Cretaceous
1345 of Ruyang, Henan Province of central China. *Historical Biology*, 28, 8–13. doi:
1346 10.1080/08912963.2014.947287.
- 1347 Manabe, M., & Hasegawa, Y. (1995). Diapsid fauna and its paleobiogeographical
1348 implication, the Neocomian section of the Tetori Group. In A. Sun, & Y. Wang (Eds.), *Sixth*
1349 *Symposium on Mesozoic Terrestrial Ecosystems and Biota, Short Papers* (pp. 179). Beijing:
1350 China Ocean Press.
- 1351 Mannion, P. D. (2009). A rebbachisaurid sauropod from the Lower Cretaceous of the Isle of
1352 Wight, England. *Cretaceous Research*, 30, 521–526.
- 1353 Mannion, P. D., Upchurch, P., & Hutt, S. (2011). New rebbachisaurid (Dinosauria:
1354 Sauropoda) material from the Wessex Formation (Barremian, Early Cretaceous), Isle of
1355 Wight, United Kingdom. *Cretaceous Research*, 32, 774–780.

- Matsukawa, M., Ito, M., Nishida, N., Koarai, K., Lockley, M. G., Nichols, D. J. (2006). The Cretaceous Tetori biota in Japan and its evolutionary significance for terrestrial ecosystems in Asia. *Cretaceous Research*, 27, 199–225.
- McDonald, A. T. (2011). The taxonomy of species assigned to *Camptosaurus* (Dinosauria: Ornithopoda). *Zootaxa*, 2783, 52–68.
- McKenna, M. C. (1973). Sweepstakes, filters, corridors, Noah's Arks, and Beached Viking Funeral Ships in palaeogeography. In D. H. Tarling, S. K. Runcorn (Eds.), *Implications of Continental Drift to the Earth Sciences* (pp. 295–308). New York: Academic Press.
- Mo, J.-Y., Huang, C.-L., Xie, S.-W., & Buffetaut, E. (2014). A megatheropod tooth from the Early Cretaceous of Fusui, Guangxi, Southern China. *Acta Geologica Sinica (English Edition)*, 88, 6–12.
- Molnar, R. E. (1990). Problematic Theropoda: "Carnosaurs". In D. B. Weishampel, P. Dodson, & H. Osmólska (Eds.), *The Dinosauria* (pp. 306–317). Berkeley, Los Angeles, Oxford: University of California Press.
- Mutihac, V., & Mutihac, G. (2010). *The geology of Romania, within the Central East European geostructural context* (p. 690). Bucharest: Ed. Didactică și Pedagogică.
- Neagu, T., Dragastan, O., & Csiki, Z. (1997). Early Cretaceous shelf paleocommunities of Cernavodă (South Dobrogea, SE Romania). *Acta Palaeontologica Romaniae*, 1, 28–36.
- Nopcsa, F. (1902). Notizen über cretacische Dinosaurier. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe*, 111, 93–114.
- Nopcsa, F. (1923). On the geological importance of the primitive reptilian fauna of the uppermost Cretaceous of Hungary; with a description of a new tortoise (*Kallokibotium*). *Quarterly Journal of the Geological Society of London*, 79, 100–116.
- Norman, D. B. (2010). A taxonomy of iguanodontians (Dinosauria: Ornithopoda) from the lower Wealden Group (Cretaceous: Valanginian) of southern England. *Zootaxa*, 2489, 47–66.

- Norman, D. B. (2013). On the taxonomy and diversity of Wealden iguanodontian dinosaurs (Ornithischia: Ornithopoda). *Revue de Paléobiologie*, 32, 385–404.
- Novas, F. E., Agnolín, F. L., Ezcurra, M. D., Porfiri, J., & Canale, J. I. (2013). Evolution of the carnivorous dinosaurs during the Cretaceous: The evidence from Patagonia. *Cretaceous Research*, 45, 174–215.
- Novas, F. E., de Valais, S., Vickers-Rich, P. A., & Rich, T. H. (2005). A large Cretaceous theropod from Patagonia, Argentina, and the evolution of carcharodontosaurids. *Naturwissenschaften*, 92, 226–230.
- Ortega, F., Escaso, F., & Sanz, J. L. (2010). A bizarre, humped Carcharodontosauria (Theropoda) from the Lower Cretaceous of Spain. *Nature*, 467, 203–206.
- Ősi, A., Codrea, V., Prondvai, E., & Csiki-Sava, Z. (2014). New ankylosaurian material from the Upper Cretaceous of Transylvania. *Annales de Paléontologie*, 100, 257–271. doi: 10.1016/j.annpal.2014.02.001.
- Ősi, A., Rabi, M., Makádi, L., Szentesi, Z., Botfalvai, G., & Gulyás, P. (2012). The Late Cretaceous continental vertebrate fauna from Iharkút (western Hungary): a review. In P. Godefroit (Ed.), *Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems* (pp. 533–569). Bloomington: Indiana University Press.
- Ősi, A., Rabi, M., & Makádi, L. (2015). An enigmatic crocodyliform tooth from the bauxites of western Hungary suggests hidden mesoeucrocodylian diversity in the Early Cretaceous European archipelago. *PeerJ*, 3, e1160. doi:10.7717/peerj.1160.
- Paolillo, A., & Linares, O. J. (2007). Nuevos cocodrilos Sebecosuchia del Cenozoico Suramericano (Mesosuchia: Crocodylia). *Paleobiologia Neotropical*, 3, 1–25.
- Pereda-Suberbiola, X., Ruiz-Omeñaca, J. I., Canudo, J. I., Torcida, F., & Sanz, J. L. (2012). Dinosaur faunas from the Early Cretaceous (Valanginian–Albian) of Spain. In P. Godefroit

(Ed.), *Bernissart Dinosaurs and Early Cretaceous Terrestrial Ecosystems* (pp. 378–407).
 Bloomington: Indiana University Press.

Pereda-Suberbiola, X., Ruiz-Omeñaca, J. I., Fernandez-Baldor, F. T., Maisch, M. W., Huerta,
 P., Contreras, R., Izquierdo, L. A., Huerta, D. M., Montero, V. U., & Welle, J. (2011). A tall-
 spined ornithopod dinosaur from the Early Cretaceous of Salas de los Infantes (Burgos,
 Spain). *Comptes Rendus Palevol*, 10, 551–558.

Pérez-Moreno, B. P., Sanz, J. L., Sudre, J., & Sigé, B. (1993). A theropod dinosaur from the
 Lower Cretaceous of southern France. *Revue de Paléobiologie, Volume spéciale* 7, 173–188.

Pol, D., & Powell, J. E. (2011). A new sebecid mesoeucrocodylian from the Rio Loro
 Formation (Palaeocene) of north-western Argentina. *Zoological Journal of the Linnean
 Society*, 163, S7–S36.

Porfiri, J. D., Novas, F. E., Calvo, J. O., Agnolín, F. L., Ezcurra, M. D., & Cerda, I. A.
 (2014). Juvenile specimen of *Megaraptor* (Dinosauria, Theropoda) sheds light about
 tyrannosauroid radiation. *Cretaceous Research*, 51, 35–55. doi:10.1016/j.cretres.2014.04.007.

Posmoşanu, E. (2003). Iguanodontian dinosaurs from the Lower Cretaceous bauxite site from
 Romania. *Acta Palaeontologica Romaniae*, 4, 431–439.

Rabi, M., & Sebök, N. (2015). A revised Eurogondwana model: Late Cretaceous notosuchian
 crocodyliforms and other vertebrate taxa suggest the retention of episodic faunal links
 between Europe and Gondwana during most of the Cretaceous. *Gondwana Research*, 28,
 1197–1211. doi:10.1016/j.gr.2014.09.015.

Racey, A. (2009). Mesozoic red bed sequences from SE Asia and the significance of the
 Khorat Group of NE Thailand. In E. Buffetaut, G. Cuny, J. Le Loeuff, & V. Suteethorn
 (Eds.), *Late Palaeozoic and Mesozoic Ecosystems in SE Asia* (pp. 41–67). Geological
 Society, London, Special Publications, 315. doi: 10.1144/SP315.5.

- 1429 Racey, A., & Goodall, J. G. S. (2009). Palynology and stratigraphy of the Mesozoic Khorat
1
2 1430 Group red bed sequences from Thailand. In E. Buffetaut, G. Cuny, J. Le Loeuff, & V.
3
4 1431 Suteethorn (Eds.), *Late Palaeozoic and Mesozoic Ecosystems in SE Asia* (pp. 69–83).
5
6
7 1432 Geological Society, London, Special Publications, 315. doi: 10.1144/SP315.6.
8
9
10 1433 Rauhut, O. W. M. (2011). Theropod dinosaurs from the Late Jurassic of Tendaguru
11
12 1434 (Tanzania). *Special Papers in Palaeontology*, 86, 195–239.
13
14 1435 Rauhut, O. W. M., & Werner, C. (1995). First record of the family Dromaeosauridae
15
16
17 1436 (Dinosauria: Theropoda) in the Cretaceous of Gondwana (Wadi Milk Formation, northern
18
19 1437 Sudan. *Paläontologische Zeitschrift*, 69, 475–489.
20
21
22 1438 Richter, U., Mudroch, A., & Buckley, L. G. (2013). Isolated theropod teeth from the Kem
23
24 1439 Kem Beds (Early Cenomanian) near Taouz, Morocco. *Paläontologische Zeitschrift*, 87, 291–
25
26
27 1440 309.
28
29 1441 Royo-Torres, R., Cobos, A., Luque, L., Aberasturi, A., Espilez, E., Fierro, I., Gonzales, A.,
30
31 1442 Mampel, L., & Alcalá, L. (2009). High European sauropod dinosaur diversity during
32
33
34 1443 Jurassic–Cretaceous transition in Riodeva (Teruel, Spain). *Palaeontology*, 52, 1009–1027.
35
36 1444 Sachs, S., & Hornung, J. J. (2013). Ankylosaur remains from the Early Cretaceous
37
38
39 1445 (Valanginian) of Northwestern Germany. *PLoS ONE*, 8(4), e60571.
40
41 1446 doi:10.1371/journal.pone.0060571.
42
43
44 1447 Salgado, L., Canudo, J. I., Garrido, A. C., Ruiz-Omeñaca, J. I., Garcia, R. A., de la Fuente,
45
46 1448 M. S., Barco, J. L., & Bollati, R. (2009). Upper Cretaceous vertebrates from El Anfiteatro
47
48
49 1449 area, Río Negro, Patagonia, Argentina. *Cretaceous Research*, 30, 767–784.
50
51 1450 Sames, B., Cifelli, R. L., & Schudack, M. E. (2010). The nonmarine Lower Cretaceous of the
52
53
54 1451 North American Western Interior foreland basin: New biostratigraphic results from ostracod
55
56 1452 correlations and early mammals, and their implications for paleontology and geology of the
57
58
59 1453 basin—An overview. *Earth-Science Reviews*, 101, 207–224.
60
61
62
63
64
65

- 1454 Sauvage, H. E. (1876). Notes sur les reptiles fossiles. *Bulletin de la Société Géologique de*
1455 *France*, 4, 435–442.
- 1456 Sauvage, H. E. (1882). Recherches sur les reptiles trouves dans le Gault de l'est du bassin de
1457 Paris. *Mémoires de la Société Géologique de France*, 2, 1–42.
- 1458 Săndulescu, M. (1984). *Geotectonica României* (p. 329). Bucharest: Ed. Tehnică.
- 1459 Seghedi, A. (2001). The North Dobrogea orogenic belt (Romania): a review. In P. A. Ziegler,
1460 W. Cavazza, A. H. F. Robertson, & S. Crasquin-Soleau (Eds.), *Peri-Tethys Memoir 6: Peri-*
1461 *Tethyan Rift/Wrench Basins and Passive Margins* (pp. 237–257). Paris: Mémoires de la
1462 Musée National d'Histoire Naturelle .
- 1463 Sereno, P. C. (1999). Dinosaurian biogeography: vicariance, dispersal and regional
1464 extinction. In Y. Tomida, T. H. Rich, & P. Vickers-Rich (Eds.), *Proceedings of the Second*
1465 *Gondwanan Dinosaur Symposium* (pp. 249–257). Tokyo: National Science Museum
1466 Monographs, 15 .
- 1467 Sereno, P. C., & Brusatte, S. L. (2008). Basal abelisaurid and carcharodontosaurid theropods
1468 from the Lower Cretaceous Elrhaz Formation of Niger. *Acta Palaeontologica Polonica*, 53,
1469 15–46.
- 1470 Sereno, P. C., Dutheil, D. B., Iarochene, M., Larsson, H. C. E., Lyon, G. H., Magwene, P. M.,
1471 Sidor, C. A., Varricchio, D. J., & Wilson, J. A. (1996). Predatory dinosaurs from the Sahara
1472 and Late Cretaceous faunal differentiation. *Science*, 272, 986–991.
- 1473 Shen, Y. B., & Mateer, N. J. (1992). An outline of the Cretaceous system in northern
1474 Xinjiang, western China. In N. J. Mateer, & P. J. Chen (Eds.), *Aspects of Nonmarine*
1475 *Cretaceous Geology* (pp. 49–77). Beijing: China Ocean Press.
- 1476 Simionescu, I. (1906). Note sur l'age des calcaires de Cernavoda (Dobrogea). *Annales*
1477 *Scientifiques de l'Université de Jassy*, 4(1), 1–3.

- 1478 Simionescu, I. (1913). *Megalosaurus* aus der Unterkreide der Dobrogea. *Centralblatt für*
1479 *Mineralogie, Geologie und Paläontologie*, 1913(20), 686–687.
- 1480 Sissingh, W. (1977). Biostratigraphy of Cretaceous calcareous nannoplankton. *Geologie en*
1481 *Mijnbouw*, 56, 37–65.
- 1482 Smith, J. B., Vann, D. R., & Dodson, P. (2005). Dental morphology and variation in theropod
1483 dinosaurs: implications for the taxonomic identification of isolated teeth. *The Anatomical*
1484 *Record A*, 285A, 699–736.
- 1485 Stein, K., Csiki, Z., Curry Rogers, K., Weishampel, D. B., Redelstorff, R., Carballido, J. L.,
1486 & Sander, P. M. (2010). Small body size and extreme cortical bone remodeling indicate
1487 phyletic dwarfism in *Magyarosaurus dacus* (Sauropoda: Titanosauria). *Proceedings of the*
1488 *National Academy of Sciences*, 107, 9258–9263.
- 1489 Stoica, M., & Csiki, Z. (2002). An earliest Cretaceous (Purbeckian) vertebrate fauna from
1490 Southern Dobrogea (southeastern Romania). In D. Grigorescu, & Z. Csiki (Eds.), 7th
1491 *European Workshop on Vertebrate Palaeontology, Sibiu, Romania* (pp. 34). Bucharest: Ars
1492 Docendi.
- 1493 Stovall, J. W., & Langston, W., Jr. (1950). *Acrocanthosaurus atokensis*, a new genus and
1494 species of Lower Cretaceous Theropoda from Oklahoma. *American Midland Naturalist*, 43,
1495 686–728.
- 1496 Stromer, E. (1931). Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wüsten
1497 Ägyptens. II. Wirbeltier-Reste der Baharijstufe (unterstes Cenoman). 10. Ein Skelett-Rest
1498 von *Carcharodontosaurus* nov. gen. *Abhandlungen der Bayerischen Akademie der*
1499 *Wissenschaften, Mathematisch-naturwissenschaftliche Abteilung, Neue Folge* 9, 1–23.
- 1500 Therrien, F., & Henderson, D. M. (2007). My theropod is bigger than yours ... or not:
1501 estimating body size from skull length in theropods. *Journal of Vertebrate Paleontology*, 27,
1502 108–115.

- 1503 Tong, H., Claude, J., Suteethorn, V., Naksri, W., & Buffetaut, E. (2009). Turtle assemblages
1504 of the Khorat Group (Late Jurassic–Early Cretaceous) of NE Thailand and their
1505 palaeobiogeographical significance. In E. Buffetaut, G. Cuny, J. Le Loeuff, & V. Suteethorn
1506 (Eds.), *Late Palaeozoic and Mesozoic Ecosystems in SE Asia* (pp. 141–151). Geological
1507 Society, London, Special Publications, 315.
- 1508 Torcida Fernández-Baldor, F., Canudo, J. I., Huerta, P., Montero, D., Pereda Suberbiola, X.,
1509 & Salgado, L. (2011). *Demandasaurus darwini*, a new rebbachisaurid sauropod from the
1510 Early Cretaceous of the Iberian Peninsula. *Acta Palaeontologica Polonica*, 56, 535–552.
- 1511 Turculeț, I., & Brânzilă, M. (2012). *Muzeul colecțiilor paleontologice originale de la*
1512 *Universitatea "Alexandru Ioan Cuza" Iași* (p. 173). Iași: Editura Universității "Alexandru
1513 Ioan Cuza" Iași.
- 1514 Turner, A. H., Makovicky, P. J., & Norell, M. A. (2012). A review of dromaeosaurid
1515 systematics and paravian phylogeny. *Bulletin of the American Museum of Natural History*,
1516 371, 1–206.
- 1517 Turner, A. H., & Sertich, J. J. W. (2010). Phylogenetic history of *Simosuchus clarki*
1518 (Crocodyliformes: Notosuchia) from the Late Cretaceous of Madagascar. *Journal of*
1519 *Vertebrate Paleontology*, 30(Supplement 1), 177–236.
- 1520 Vasile, Ș., & Csiki, Z. (2011). New Maastrichtian microvertebrates from the Rusca Montană
1521 Basin (Romania). *Oltenia. Studii și comunicări. Științele Naturii*, 27(1), 221–230.
- 1522 Vickers-Rich, P., Rich, T. H., Lanus, D. R., Rich, L. S. V., & Vacca, R. (1999). "Big Tooth"
1523 from the Early Cretaceous of Chubut Province, Patagonia: a possible carcharodontosaurid. In
1524 Y. Tomida, T. H. Rich, & P. Vickers-Rich (Eds.), *Proceedings of the Second Gondwanan*
1525 *Dinosaur Symposium* (pp. 85–88). Tokyo: National Science Museum Monographs, 15.

- Vremir, M. (2010). New faunal elements from the Late Cretaceous (Maastrichtian) continental deposits of Sebeş area (Transylvania). *Terra Sebus. Acta Musei Sabesiensis*, 2, 635–684.
- Vullo, R., Néraudeau, D., & Lenglet, T. (2007). Dinosaur teeth from the Cenomanian of Charentes, western France: evidence for a mixed Laurasian-Gondwanan assemblage. *Journal of Vertebrate Paleontology*, 27, 931–943.
- Weishampel, D. B. (1990). Dinosaurian distribution. In D. B. Weishampel, P. Dodson, & H. Osmólska (Eds.), *The Dinosauria* (pp. 63–140). Berkeley, Los Angeles, Oxford: California University Press.
- Weishampel, D. B., Barrett, P. M., Coria, R. A., Le Loeuff, J., Xu, X., Zhao, X-J., Sahni, A., Gomani, E. M., & Noto, C. R. (2004). Dinosaur distribution. In D. B. Weishampel, P. Dodson, & H. Osmólska (Eds.), *The Dinosauria. Second Edition* (pp. 517–606). Berkeley, Los Angeles, London: University of California Press.
- Weishampel, D. B., Csiki, Z., Benton, M. J., Grigorescu, D., & Codrea, V. (2010). Palaeobiogeographic relationships of the Haţeg biota — Between isolation and innovation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 293, 419–437.
- Weishampel, D. B., Grigorescu, D., & Norman, D. B. (1991). The Dinosaurs of Transylvania. *National Geographic Research & Exploration*, 7(2), 196–215.
- Weishampel, D. B., Jianu, C. M., Csiki, Z., & Norman, D. B. (2003). Osteology and phylogeny of *Zalmoxes* (n. g.), an unusual euornithopod dinosaur from the latest Cretaceous of Romania. *Journal of Systematic Palaeontology*, 1, 65–123.
- Weishampel, D. B., Norman, D. B., & Grigorescu, D. (1993). *Telmatosaurus transsylvanicus* from the Late Cretaceous of Romania: the most basal hadrosaurid dinosaur. *Palaeontology*, 36, 361–385.

Williamson, T. E., & Brusatte, S. L. (2014). Small theropod teeth from the Late Cretaceous of the San Juan Basin, northwestern New Mexico and their implications for understanding latest Cretaceous dinosaur evolution. *PLoS ONE*, 9(4), e93190.

doi:10.1371/journal.pone.0093190.

Zarcone, G., Cillari, F. M. P., Stefano, P. D., Guzzetta, D., & Nicosia, U. (2010). A possible bridge between Adria and Africa: New palaeobiogeographic and stratigraphic constraints on the Mesozoic palaeogeography of the Central Mediterranean area. *Earth-Science Reviews*, 103, 154–162.

Figure captions

Figure 1. Simplified geological map of the Cernavodă-Cochirleni area; inset shows the position of the study area within Romania. Legend: 1. Quaternary: a. Holocene alluvia, b. Pleistocene–Holocene loessoid deposits; 2. Pre-Quaternary Cenozoic (Middle Eocene and Miocene) deposits; Cretaceous: 3. Peștera Formation, Lower Cenomanian; 4. Cochirleni Formation; uppermost Aptian–Lower Albian; 5. Gherghina Formation, Middle–Upper Aptian; 6. Ostrov (= Ramadan) Formation; Barremian–Lower Aptian; 7. Cernavodă Formation, Alimanu Member, Berriasian–Valanginian; 8. Water courses. (Redrawn after Dragastan et al., 1998, 2014).

Figure 2. Specimen UAIC (SCM1) 615, indeterminate carcharodontosaurid lateral tooth from Cochirleni, Southern Dobrogea. A. UAIC (SCM1) 615, as figured by Simionescu (1913); B. Current state of UAIC (SCM1) 615, mounted in a limestone holder.

Figure 3. Detailed morphology of UAIC (SCM1) 615, an indeterminate carcharodontosaurid lateral tooth from Cochirleni, Southern Dobrogea. UAIC (SCM1) 615 in A. labial? side; B., distal; C., lingual? side, and D., basal (mesial to the right) views. Details of the distal carina (marked with boxes in A, respectively C): apical part in E., labial? and F. distal views; basal part in G., lingual? and H., distal views. Scale bar: 1 cm (A–D), 5 mm (E–H).

Figure 4. Dental morphospace of the different theropod clades according to the results of the PCA analysis; UAIC (SCM1) 615 (red star) plots within the morphospace occupied by Carcharodontosauridae. See further details of this analysis, as well as other quantitative analyses used to identify the tooth that deliver similar results (cluster analysis, discriminant function analysis, phylogenetic analysis), in the Supplementary Material.

Figure 5. A. Palaeogeographic setting of the two early Early Cretaceous Romanian dinosaur occurrences: the Berriasian–Valanginian Cornet locality (orange star), located on a Neotethyan archipelago island, and the Valanginian Cochirleni locality (red star), situated on the marginal areas of the Eastern European cratonic mainland. B. Global chronostratigraphic and palaeobiogeographic distribution of the Carcharodontosauridae, plotted on Middle Aptian (approx. 120 Mya) palaeogeographic map; red star marks the position of UAIC (SCM1) 615 from Southern Dobrogea. Legend: 1 – *Veterupristisaurus*, ‘*Megalosaurus*’ *ingens*, Carcharodontosauridae indet., Tanzania, Late Jurassic; 2 – *Concavenator*, Spain, Barremian; 3 – Carcharodontosauridae indet., Thailand, Barremian; 4 – *Acrocanthosaurus*, southeastern United States, Aptian–Albian; 5 – Carcharodontosauridae indet., Spain, Aptian; 6 – *Eocarcharia*, Niger, Aptian–Albian; 7 – Carcharodontosauridae indet., Guangxi, China, Aptian; 8 – Carcharodontosauridae indet., Henan, China, Aptian; 9 – *Kelmaysaurus*, Xinjiang, China, Aptian–Albian; 10 – Carcharodontosauridae indet., France, Cenomanian; 11

1599 – *Sauroniops*, Morocco, Cenomanian; 12 – Carcharodontosauridae indet., Japan,
1
2 1600 Cenomanian–early Turonian; 13 – *Shaochilong*, Inner Mongolia, China, Turonian; 14 –
3
4
5 1601 Carcharodontosauridae indet., São Paulo, Brazil, Campanian–Maastrichtian (for relevant
6
7 1602 references, see text, 5.4.). Palaeogeographic maps, courtesy of Ron Blakey
8
9
10 1603 (<http://cpgeosystems.com/>).
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1 color
[Click here to download high resolution image](#)

Figure 2 color
[Click here to download high resolution image](#)

Figure 3 color
[Click here to download high resolution image](#)

Figure 4 color
[Click here to download high resolution image](#)

Figure 5 color

[Click here to download high resolution image](#)

○ basal / indeterminate
Carcharodontosauridae

☆ UAIC (SCM1) 615

Carcharodontosaurinae

△ Carcharodontosaurus

▽ Giganotosaurus

△ Mapusaurus

▽ Tyrannotitan

1								
2								
3	Taxon	Side	Position	Specimen	Source from CBL	CBW	CH	
4	ROMANIAN TOOTH			SCM1 615		29	16.25	85.5
5	Eoraptor	Left	pmx2	PVSJ 512	Pers. Obsei	2.5	1.62	6.74
6	Eoraptor	Left	pmx3	PVSJ 512	Pers. Obsei	1.97	2.35	5.92
7	Eoraptor	Left	pmx4	PVSJ 512	Pers. Obsei	2.19	1.74	6.52
8	Eoraptor	Right	pmx2	PVSJ 512	Pers. Obsei	2.17	1.56	5.01
9	Eoraptor	Right	pmx4	PVSJ 512	Pers. Obsei	2.08	1.61	4.17
10	Eoraptor	Left	mx2	PVSJ 512	Pers. Obsei	2.69	1.82	5.55
11	Eoraptor	Left	mx4	PVSJ 512	Pers. Obsei	3.03	1.48	5.65
12	Eoraptor	Left	mx5	PVSJ 512	Pers. Obsei	3.56	1.69	5.48
13	Eoraptor	Left	mx9	PVSJ 512	Pers. Obsei	2.49	1.75	5.11
14	Eoraptor	Left	mx10	PVSJ 512	Pers. Obsei	2.7	1.22	4.58
15	Eoraptor	Left	mx11	PVSJ 512	Pers. Obsei	2.32	1.42	2.34
16	Eoraptor	Left	pm02	PVSJ 512	Smith & La	2.88	1.85	7.15
17	Eoraptor	Left	pm03	PVSJ 512	Smith & La	1.98	1.48	5.73
18	Eoraptor	Left	pm04	PVSJ 512	Smith & La	1.89	1.55	6.12
19	Eoraptor	Right	pm02	PVSJ 512	Smith & La	2.32	1.8	5.19
20	Eoraptor	Left	mx02	PVSJ 512	Smith & La	2.11	1.51	5.71
21	Eoraptor	Left	mx04	PVSJ 512	Smith & La	3.04	1.97	6.58
22	Eoraptor	Left	mx06	PVSJ 512	Smith & La	2.9	1.74	5.44
23	Eoraptor	Left	mx07	PVSJ 512	Smith & La	2.71	1.58	6.17
24	Eoraptor	Left	mx09	PVSJ 512	Smith & La	2.67	1.82	4.99
25	Eoraptor	Left	mx10	PVSJ 512	Smith & La	2.56	1.69	4.72
26	Eoraptor	Right	mx02	PVSJ 512	Smith & La	2.94	1.87	5.32
27	Eoraptor	Right	mx04	PVSJ 512	Smith & La	2.54	1.55	6.5
28	Eoraptor	Right	mx05	PVSJ 512	Smith & La	3.33	1.82	6.76
29	Eoraptor	Right	mx07	PVSJ 512	Smith & La	2.86	1.6	4.83
30	Eoraptor	Right	mx08	PVSJ 512	Smith & La	2.8	1.5	4.71
31	Ischisaurus	Right	pmx1	MACN 18.C	Pers. Obsei	8.16	4.7	14.62
32	Ischisaurus	Right	pmx2	MACN 18.C	Pers. Obsei	7.48	4.43	14.5
33	Eodromaeus	Left	mx3	PVSJ 561	Pers. Obsei	3.61	1.59	9.67
34	Coelophysis	Left	pmx2	CM 82931	Pers. Obsei	1.7	0.54	4
35	Coelophysis	Left	pmx3	CM 82931	Pers. Obsei	1.8	1.03	6.8
36	Coelophysis	Left	mx1	CM 81765	Pers. Obsei	3	1.49	8.2
37	Coelophysis	Left	mx2	CM 81765	Pers. Obsei	4.1	1.37	8.9
38	Coelophysis	Left	mx4	CM 81765	Pers. Obsei	4.4	1.63	11.6
39	Coelophysis	Left	mx6	CM 81765	Pers. Obsei	5.5	1.71	9.7
40	Coelophysis	Left	mx8	CM 81765	Pers. Obsei	5.9	1.79	9.3
41	Coelophysis	Left	mx9	CM 81765	Pers. Obsei	5.4	1.74	8.6
42	Coelophysis	Left	mx11	CM 81765	Pers. Obsei	5.4	1.85	7.5
43	Coelophysis	Left	mx13	CM 81765	Pers. Obsei	4.7	1.58	5.5
44	Coelophysis	Left	mx14	CM 81765	Pers. Obsei	5.2	1.7	6.2
45	Coelophysis	Left	mx15	CM 81765	Pers. Obsei	3.9	1.49	5.4
46	Coelophysis	Left	mx16	CM 81765	Pers. Obsei	3.3	1.14	5
47	Coelophysis	Left	mx17	CM 81765	Pers. Obsei	3.5	1.63	4.1
48	Coelophysis	Left	mx19	CM 81765	Pers. Obsei	3.35	1.86	3.3
49	Coelophysis	Left	mx21	CM 81765	Pers. Obsei	3.54	0.93	3.45
50								
51								
52								
53								
54								
55								
56								
57								
58								
59								
60								
61								
62								
63								
64								
65								