THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Regulation of the avian central melanocortin system and the role of leptin

Citation for published version:

Boswell, T & Dunn, IC 2015, 'Regulation of the avian central melanocortin system and the role of leptin', *General And Comparative Endocrinology*, vol. 221, pp. 278-283.
<https://doi.org/10.1016/j.ygcen.2014.12.009>

Digital Object Identifier (DOI):

[10.1016/j.ygcen.2014.12.009](https://doi.org/10.1016/j.ygcen.2014.12.009)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

General And Comparative Endocrinology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

1 Mini-review for special issue of GCE - 27th Conference of European Comparative
2 Endocrinologists

3

4 Regulation of the avian central melanocortin system and the role of leptin

5

6 Timothy Boswell^{a*}, Ian C. Dunn^b

7

8 ^aSchool of Biology, Institute of Neuroscience, and Centre for Behaviour and Evolution,
9 Newcastle University, England, United Kingdom

10

11 ^bRoyal (Dick) School of Veterinary Studies, Roslin Institute, University of Edinburgh,
12 Easter Bush, Scotland, United Kingdom

13

14 *Corresponding Author

15 Dr. Tim Boswell

16 School of Biology,

17 Ridley Building

18 Newcastle University,

19 Newcastle upon Tyne

20 NE1 7RU

21 United Kingdom

22 timothy.boswell@ncl.ac.uk

23 +44(0)1912088502

Abstract

The avian central melanocortin system is well conserved between birds and mammals in terms of the component genes, the localisation of their expression in the hypothalamic arcuate nucleus, the effects on feeding behaviour of their encoded peptides and the sensitivity of agouti-related protein (AGRP) and pro-opiomelanocortin (POMC) gene expression to changes in energy status. Our recent research has demonstrated that AGRP gene expression precisely differentiates between broiler breeder hens with different histories of chronic food restriction and refeeding. We have also shown that the sensitivity of AGRP gene expression to loss of energy stores is maintained even when food intake has been voluntarily reduced in chickens during incubation and in response to a stressor. However, the similarity between birds and mammals does not appear to extend to the way AGRP and POMC gene expression are regulated. In particular, the preliminary evidence from the discovery of the first avian leptin genes suggests that leptin is more pleiotrophic in birds and may not even be involved in regulating energy balance. Similarly, ghrelin exerts inhibitory, rather than stimulatory, effects on food intake. The fact that the importance of these important long-term regulators of AGRP and POMC expression in mammals appears diminished in birds suggests that the balance of regulatory inputs in birds may have shifted to more short-term influences such as the tone of cholecystokinin (CCK) signalling. This is likely to be related to the different metabolic fuelling required to support flight.

Keywords: AGRP; POMC; AMPK; leptin; arcuate nucleus; chicken

1. Introduction

The avian melanocortin system genes were all cloned and characterised 14 years ago. They comprise five melanocortin receptors (Takeuchi et al., 1996; Takeuchi et al., 1998; Takeuchi and Takahashi, 1998; Takeuchi and Takahashi, 1999), and the pro-opiomelanocortin (POMC) and agouti-related protein (AGRP) genes encoding their endogenous agonists and antagonists, respectively (Berghman et al., 1998; Takeuchi et al., 1999; Takeuchi et al., 2000). The central melanocortin system shows evolutionarily functional and neuroanatomical conservation in comparison with mammals. Thus, melanocortin signalling is regulated by opposing, agonistic and antagonistic actions of endogenous α -melanocyte-stimulating hormone (α -MSH) and agouti-related protein (AGRP) that, respectively, increase and decrease food intake after central injection (Tachibana et al., 2001; Strader and Buntin, 2003; Strader et al., 2003). Also, the AGRP and POMC genes are expressed in the arcuate nucleus of the hypothalamus where AGRP is co-expressed with neuropeptide Y (NPY) in individual arcuate nucleus neurons (Boswell et al., 2002). In this mini-review we focus on the regulation of the AGRP and POMC genes in birds with emphasis on our recent work on chronic food restriction and natural models of voluntarily reduced food intake. We also consider the implications of the identification and preliminary characterisation in 2014 of the first avian leptin genes.

2. Effects of manipulating energy status

The functional evolutionary conservation of the avian central melanocortin system extends to its response to manipulation of energy status. Most investigations have focused on short-term food deprivation, which increases AGRP mRNA (Phillips-Singh et al.,

2003; Higgins et al., 2010; Song et al., 2012). Changes in mRNA appear to reflect parallel changes in peptide because food deprivation increased the number of AGRP-immunoreactive cells in the arcuate nucleus of the ring dove (*Streptopelia risoria*) which were also elevated during the parental phase of the breeding cycle when the birds are in negative energy balance (Strader and Buntin, 2003; Strader et al., 2003). For POMC, Higgins et al. (2010) observed a significant decrease in mRNA after food deprivation and two studies have reported a non-significant decrease (Phillips-Singh et al., 2003; Song et al., 2012). Thus the changes in AGRP and POMC expression during fasting are in the direction predicted from the behavioural effects of their encoded peptides as part of a coordinated counter-regulatory response of hypothalamic appetite control peptides to restore lost energy stores (Phillips-Singh et al., 2003). Higgins et al. (2010) used microarray analysis to explore patterns of gene expression in the hypothalamus of domestic chicks in relation to food deprivation. Pathway analysis software predicted that POMC lies within a network of six interrelated differentially-expressed genes and experimental evidence for the existence of the network was provided from *in vitro* experiments. In addition to POMC, the network contained genes encoding the neuropeptide relaxin-3, the neuropeptide receptors NPY Y5 and somatostatin receptor 5, and the β 2 adrenergic and metabotropic glutamate receptor 8 neurotransmitter receptors (Fig. 1). POMC was the only gene downregulated by fasting while the others in the network were upregulated. This suggests a role for these pathways in the control of appetite within the hypothalamus. There is also suggestive evidence for a role of hypothalamic energy sensing pathways in regulating central melanocortin signalling. Proszkowiec-Weglarz observed strong immunostaining for the energy sensor AMP-

93 activated protein kinase (AMPK) in the arcuate nucleus and Song et al. (2012) reported
94 changes in AMPK activity indicated by phosphorylation during food deprivation and
95 refeeding that paralleled changes in AGRP gene expression.

96 In comparison to food deprivation, the effects of chronic food restriction on the
97 central melanocortin system have received less attention. We recently measured AGRP
98 and POMC gene expression in broiler breeder hens subjected to different levels of
99 chronic food restriction and refeeding (Dunn et al., 2013a). Food restriction for six weeks
100 at a level routinely used in the poultry industry to improve reproductive performance
101 strongly stimulated AGRP gene expression. We also made comparisons between hens of
102 the same body mass that had experienced different feeding histories. Thus, we were able
103 to distinguish in terms of significant changes in AGRP mRNA between birds killed at the
104 same body mass that had been either been maintained on an intermediate level of food
105 restriction or more severely restricted and then refed for two weeks. We also observed a
106 difference between restricted birds compared to restricted hens that had been allowed to
107 refeed for two days, with refeeding causing a pronounced decrease in AGRP mRNA.
108 Overall we found that AGRP mRNA provided an integrated and sensitive measure of
109 feeding history and has potential as an objective measure of hunger in animal welfare
110 research. As has been observed in food deprivation studies, there appears to be a
111 tendency for POMC gene expression to be decreased during food restriction. Hen et al.
112 (2006) reported a significant decrease in POMC mRNA after seven days' food restriction
113 in broiler and layer chickens Although we did not observe a significant effect of
114 restriction on POMC expression in our study (Dunn et al. 2013a), we have subsequently
115 observed a significant decrease in chronically restricted broiler chickens in a recent study

with greater sample sizes and statistical power (Dunn et al., unpublished).

3. Seasonal and stress-induced changes in body mass

Seasonal cycles of food intake and body mass have been well studied in birds, particularly in relation to migration, but little is known about the regulatory role of the central melanocortin system (Cornelius et al., 2013). Experiments in white-crowned sparrows (*Zonotrichia leucophrys gambelii*) have suggested that increases in body mass during the migratory period are associated with a regulated change in the level around which body mass is defended, a phenomenon known as a ‘sliding set point’ or rheostasis (King et al., 1963; Mrosovsky, 1990). One hypothesis is that seasonal changes in appetite and body weight are driven by an altered basal expression of AGRP and POMC in a direction reflecting the behavioural effects of the peptides. For example, to generate a seasonal body mass increase, the expression of AGRP and POMC would be respectively increased and decreased. However preliminary evidence from experiments in the European quail (*Coturnix coturnix*) suggested that AGRP and POMC gene expression were unchanged at a time when food intake and body mass were increasing during the migratory period (Cornelius et al., 2013). These findings are reminiscent of those from studies of seasonal mammals where the rheostatic mechanisms appear to lie in brain areas outside the arcuate nucleus (Ebling, 2014).

Another example of rheostasis that has been well defined in birds is the voluntary loss of appetite and body mass that occurs during incubation, one of several examples of naturally occurring ‘animal anorexias’ (Savory, 1979; Sherry et al., 1980; Mrosovsky and Sherry, 1980). We have recently examined the expression of AGRP and POMC during

incubation in the domestic chicken (Dunn et al., submitted) where hens show reduced feeding behaviour for a three-week period, even though food is freely available. We hypothesised that gene expression would be unchanged because the decreased body mass represents a rheostatic change in the defended level. However, unexpectedly, AGRP was increased in incubating birds compared to laying hens. To control for the possibly confounding fact that the reproductive system is regressed in incubating birds, we imposed food restriction on laying hens to induce a mass loss comparable to that in incubating birds. AGRP expression was similarly high in the incubating and food restricted groups and was significantly reduced in restricted birds allowed to refeed. These findings suggest that AGRP expression remains sensitive to nutritional state during incubation, which may be of adaptive significance in allowing a hen to terminate incubation if her energy stores fall too low. However, the normally anabolic effects associated with increased AGRP mRNA must be overridden by inhibitory inputs. POMC is a candidate because its mRNA was increased in incubating birds on the boundary of significance, although hypothalamic vasoactive intestinal polypeptide (VIP) may also be involved.

To consider another example of voluntarily reduced food intake in birds, we examined AGRP and POMC gene expression six days after transferring domestic chickens from individual housing in cages to social housing in pens, a stressor that induced a 70% reduction in food intake (Dunn et al., submitted). The results were comparable to incubation, with AGRP mRNA being significantly elevated in transferred birds together with significantly increased POMC mRNA. Again, the results suggested that AGRP gene expression is sensitive to energy status and this may be important to

allow the birds to recover from the effects of the stressor. Similar to incubation, it is possible that increased POMC expression plays a role in stress-induced inhibition of food intake as has been suggested in some mammalian studies (Liu et al., 2007).

4. Regulation of AGRP and POMC gene expression by leptin

As noted in Section 3, a period of two days' refeeding following prolonged food restriction in broiler chickens results in a rapid decrease in AGRP gene expression. The endocrine signals underlying this response, together with those responsible for increased AGRP expression after energy deprivation or restriction, are currently uncertain. In mammals, leptin is well established as an important regulator of AGRP and POMC gene expression, upregulating POMC and downregulating AGRP (Friedman, 2009). However progress in understanding the role of leptin in regulating the central melanocortin system in birds has been slow because the existence of leptin genes in avian genomes has been controversial for twenty years. Early claims in the literature that bird leptin genes had been cloned were challenged and there was no evidence for their existence from sequenced avian genomes and other genomic resources (Friedman-Einat et al., 1999; Pitel et al., 2010). However, in 2013, a leptin-like sequence was released into public databases during annotation of the zebra finch (*Taeniopygia guttata*) genome that has led to leptin-like genes being identified in, currently, ten bird genomes (Prokop et al., 2014; Friedman-Einat et al., 2014; Huang et al., 2014; Friedman-Einat and Seroussi, 2014). These show a similar syntenic relationship with surrounding genes as other vertebrate leptins. The fact that avian leptin proved difficult to clone by traditional methods is explained by the approximately 30% amino acid identity to mouse and human leptin, together with a high

(up to 80%) GC content in the coding region. Recombinant peregrine falcon (*Falco peregrinus*), zebra finch, and rock dove (*Columba livia*) leptins activate the leptin receptor (LEPR) in avian LEPR reporter gene cell lines, and also the JAK-STAT pathway through phosphorylation of STAT3 (Prokop et al., 2014; Huang et al., 2014; Friedman-Einat et al., 2014). The tissue distribution of expression of the newly discovered leptin genes has only been studied in two species to date. In the zebra finch real time PCR showed expression of leptin primarily in the brain and the pituitary gland, of which the latter was also the most abundant site of expression of the LEPR (Huang et al., 2014). The pituitary was also the main site of expression of the rock dove LEPR while leptin was largely expressed in the liver and gonads (Friedman-Einat et al., 2014). In both species, leptin expression in adipose tissue was low or undetectable. Thus, although limited, the expression data suggest that leptin cannot be regarded as being predominantly an adipose tissue hormone in birds to the extent it is in mammals and that it shares a more widespread distribution comparable to that observed in other non-mammalian vertebrates (Crespi and Denver, 2006; Rønnestad et al., 2010). This led Millar (2014) in a commentary on the discovery of avian leptin-like genes to suggest that they may primarily be involved in pituitary-gonadal function rather than energy balance regulation, although studies investigating avian leptin expression in response to nutritional manipulation have yet to be performed. Previous investigations in birds had suggested that putative leptin protein and LEPR binding activity are very low or undetectable in the blood in several avian species (Richards et al., 2000; Adachi et al., 2008; Hen et al., 2008; Yosefi et al., 2010). Also, although short isoforms of the avian LEPR have been identified (Liu et al., 2007) the most abundant avian LEPR isoform is orthologous to the

mammalian Rb long isoform and none has so far been shown to be equivalent to the mammalian Ra short isoform thought to be involved in the transport of leptin into the brain. Collectively, these data may indicate a more paracrine role for leptin in birds, although rock dove serum was recently shown to activate a LEPR reporter bioassay (Friedman-Einat et al., 2014).

Much of the investigation into the role of leptin in regulating energy balance in birds has focused on the domestic chicken. However, the existence of functional leptin genes in chickens, Japanese quail and the mallard duck has yet to be proven (Friedman-Einat and Seroussi, 2014). Friedman-Einat and Seroussi (2014) suggest that this may be associated with changes in appetite regulation linked to domestication, and that it is in line with the observation that a pegylated leptin antagonist that inhibited activation of a LEPR bioassay reporter system by rock dove leptin was ineffective in altering food intake and body mass in chickens (Gertler et al., 2014). However, that conclusion is dependent on leptin having a physiological role in energy balance regulation in the chicken, which may not be the case. Also, if chicken leptin is absent from the genome, it might be expected that the LEPR would also be missing or degenerate as appears to be the case with kisspeptin in birds (Pasquier, 2014). It is therefore possible that leptin may yet be identified in chickens, and the recent discoveries in other species have provided a new impetus and scope for comparative analysis. It is certainly the case that mammalian leptin inhibits food intake in chickens and other birds after central or peripheral administration (Denbow et al., 2000; Cerasale et al., 2011). Similar inhibitory results have been obtained after administration of the ‘chicken leptin’ protein that shares 97% amino acid sequence identity to mouse leptin and is evidently artifactual (Raver et al., 1998; Löhmus et al.,

2003). Given the new information about the low sequence identity of avian leptins with mammalian leptins, these results need to be reevaluated using recombinant avian leptin proteins. This applies to the only study in which the effects of mammalian leptin ('chicken leptin') were investigated in relation to AGRP and POMC expression (Dridi et al., 2005). Here, leptin was administered peripherally to young broiler chickens for six hours after which hypothalamic AGRP and POMC gene expression were measured. No change in expression was detected for either gene, in contrast to what would be expected from mammalian studies. If leptin does regulate AGRP and POMC expression in birds, it would be expected that the LEPR is co-expressed with those genes in arcuate nucleus neurons. However, in our experience, the chicken LEPR long isoform is difficult to detect by *in situ* hybridisation because although detectable by PCR it is not highly expressed in the hypothalamus. Although we were able to increase the sensitivity of radioactive oligonucleotide *in situ* hybridisation sufficiently to detect expression of low-abundance prolactin receptor in the chicken hypothalamus, we were unable to detect expression of the LEPR at the same time using the same methodology (T. Boswell, unpublished).

5. Regulation of AGRP and POMC gene expression by other neuroendocrine signals

In the absence of information regarding leptin's possible role in regulating AGRP and POMC gene expression, a small number of investigations have focused on other candidate signals. Several studies have been performed in the domestic chick in relation to insulin, the circulating concentrations of which decrease with fasting in birds as well as in mammals (Christensen et al., 2013). Insulin was established in mammals as a regulator of arcuate nucleus appetite peptide gene expression before the discovery of leptin. It

stimulates POMC gene expression and inhibits that of NPY and AGRP, corresponding to an inhibitory effect of centrally-administered insulin on food intake (Porte et al., 2002). Insulin also inhibited food intake after central injections in domestic chicks, an effect blocked by melanocortin receptor antagonists (Honda et al., 2007; Shiraishi et al., 2008). Chick NPY neurons were demonstrated to co-express the insulin receptor and it is likely, but was not determined, that those same individual neurons expressed AGRP given the co-expression of NPY and AGRP previously observed in birds (Boswell et al., 2002; Shiraishi et al., 2011). Co-expression of the insulin receptor with α -MSH was also observed in the arcuate nucleus (Shiraishi et al., 2011). Two independent studies in chicks reported increased gene expression of POMC in the chick brain after central insulin injection but no effect on AGRP mRNA in contrast to an inhibitory effect on NPY mRNA (Honda et al., 2007; Shiraishi et al., 2008). The apparent absence of an inhibitory effect of insulin on AGRP gene expression may point to a regulatory difference between birds and mammals although observations need to be extended to adult chickens and to other species.

With regard to other neuroendocrine signals, Byerly et al. (2009) combined *in vivo* and *in vitro* studies on chicken hypothalami from fat and lean chicken lines to investigate the possible regulation of appetite peptide gene expression by brain-derived neurotrophic factor (BDNF), triiodothyronine (T3) and corticosterone. BDNF did not affect AGRP and POMC gene expression but T3 was found both *in vitro* and *in vivo* to upregulate AGRP mRNA and downregulate POMC mRNA. A specific effect of T3 on AGRP gene expression would not be predicted from the fact that circulating T3 concentrations are decreased by fasting in chickens (Harvey and Klandorf, 1983) at the time AGRP mRNA

is increased. However it is possible that T3 concentrations within the hypothalamus may differ from those in the blood as a result of local conversion of thyroxine. Fasting increases plasma corticosterone concentrations in birds as in mammals (Harvey and Klandorf, 1983) and it would therefore be predicted that corticosterone would increase AGRP expression and decrease POMC expression in line with mammalian studies (Makimura et al., 2003). However Byerly et al. (2009) reported that while corticosterone administration *in vitro* and *in vivo* decreased POMC mRNA, it did not affect AGRP gene expression. Liu et al. (2012) replicated the suppressive effect of corticosterone on POMC expression after peripheral administration in laying hens, but also reported decreased AGRP mRNA. However, central injection of the glucocorticoid receptor agonist dexamethasone in broiler chicks produced no change in AGRP or POMC gene expression (Liu et al., 2014). The lack of consensus may be because adrenalectomy and glucocorticoid replacement together with consideration of corticosterone and insulin interactions are needed to demonstrate a stimulatory response for AGRP mRNA unambiguously (Makimura et al., 2003). Such studies have not been performed in birds.

An additional candidate neuroendocrine feedback signal to the arcuate nucleus that we have investigated recently is tonic signalling by cholecystokinin (CCK). Investigation into the genetic basis of growth rate between lines of fast- and slow-growing chickens revealed the importance of an apparently cis-acting regulatory region downstream of the CCKAR (CCK1 receptor) that explained 19% of the difference in body mass between the lines. Birds with the high-growth haplotype at that locus showed lower expression of CCKAR throughout the body together with decreased sensitivity to the inhibitory effects of exogenous CCK administration (Dunn et al., 2013b). Thus high growth rate was

associated with a lower overall tone of CCK signalling. When we investigated AGRP and POMC gene expression, we found that AGRP mRNA was significantly higher in the high-growth haplotype compared to heterozygotes and the low-growth haplotype but there was no difference in POMC mRNA. These data suggest that AGRP gene expression is sensitive to the tone of CCK signalling. The effect of CCK on the central melanocortin system has not been widely investigated but sensitivity of the feeding response to CCK injections was reduced in MC4R-knockout mice and MC4R neurons were observed to project to hindbrain regions influenced by CCK signalling (Blevins et al., 2009). It is, of course, possible that the increased AGRP gene expression we observed in the high-growth birds was a secondary consequence of altered CCKAR expression, and we are carrying out further studies to test for a direct regulatory influence of CCK.

6. Future perspectives

It is now well established that the avian central melanocortin system shows conservation with its mammalian counterpart in terms of the genes involved and the neuroanatomical localisation of the neurons expressing the genes within the arcuate nucleus. There is functional conservation with respect to the behavioural effects of the peptides and the response of gene expression to changes in energy status. The precision with which we were able to distinguish between hens with different histories of chronic food restriction and refeeding is remarkable (Dunn et al., 2013a) and from our work that we report in Section 3, sensitivity of AGRP expression to energy status appears to be maintained even in situations where food intake is voluntarily decreased, indicating the functional importance of this mechanism. However a similar degree of conservation does

not appear to extend to the way the AGRP and POMC genes are regulated. There are striking differences in the regulatory effects of the two hormones principally associated with the long-term regulation of energy balance in mammals, leptin and ghrelin. As we review in Section 4, a 20-year search for leptin in birds has finally resulted in the identification of leptin-like genes in several avian orders. However, the encoded proteins show low sequence identity to mammalian leptins and the preliminary evidence from the tissue distribution of gene expression suggests that leptin is more pleiotrophic in birds and may not be primarily involved in regulating energy balance. Ghrelin research in birds has been more straightforward in that there has been no lengthy controversy about its existence. However the behavioural effect of ghrelin on feeding is fundamentally different to the situation in mammals in that there is consistent evidence that ghrelin acts in the brain to inhibit food intake rather than stimulate it (Kaiya et al., 2013). The inhibitory effect appears to be mediated by hypothalamic corticotrophin-releasing factor (Saito et al., 2005) rather than through the central melanocortin system. Two other hormones that influence long-term regulation of arcuate nucleus peptide signalling in mammals, insulin and corticosterone, share more similarity with regulatory effects observed in mammals, but with some differences (Fig. 1). Overall, the influence of the neuroendocrine signals involved in long-term energy balance regulation appear diminished in birds compared to mammals, perhaps indicating that regulation of AGRP and POMC synthesis and activity is under more short-term control. Our preliminary observations suggesting a regulatory influence of CCK support this idea. It is also possible that direct energy sensing within the hypothalamus plays a relatively greater role in regulating the arcuate nucleus neural circuitry in birds. The application of genomic

methods has provided a foundation for understanding how changes in the hypothalamic environment may influence central melanocortin signalling (Fig. 1). It should not be surprising that the regulation of energy balance is likely to differ between vertebrate taxa. Birds in particular face very different metabolic challenges from most mammals and from ectothermic vertebrates imposed by the requirements of the intense aerobic activity of flight. Fundamental differences in metabolism such as the relative circulating concentrations of insulin and glucagon, present even in domesticated birds like the chicken, are likely to reflect this. It would be beneficial for future investigations to draw more on the established metabolic differences between birds and mammals, particularly on the relatively greater reliance in birds on lipid metabolism as a source of metabolic fuels. We hope that the impetus to the field provided by the identification of avian leptins will encourage a new approach.

Acknowledgments

We thank Drs. Hélène Volkoff and Ivar Rønnestad for organising the Symposium on Endocrine regulation of feeding, digestion and metabolism at the 27th Conference of European Comparative Endocrinologists. Our work has been supported by DEFRA grant AW1141 'Quantifying the subjective state of feed-restricted broiler chickens using behavioural and neurochemical measures' and by BBSRC grant BB/L000571/1 'Investigating how the type and quantity of food affect foraging behaviour and the neural circuits controlling feeding in broiler breeder chickens'. The Roslin Institute is funded by the BBSRC through Institute Strategic Grant funding.

References

- Adachi, H., Takemoto, Y., Bungo, T., Ohkubo, T., 2008. Chicken leptin receptor is functional in activating JAK-STAT pathway in vitro. *J. Endocrinol.* 197, 335-342.
- Berghman L.R., Devreese, B., Verhaert, P., Gerets, H., Arckens, L., Vanden Broeck, J., Van Beeumen, J., Vaudry, H., Vandesande, F., 1998. The molecular characterisation of chicken pituitary N-terminal pro-opiomelanocortin (POMC). *Mol. Cell. Endocrinol.* 142, 119-130.
- Blevins, J.E., Morton, G.J., Williams, D.L., Caldwell, D.W., Bastian, L.S., Wisse, B.E., Schwartz, M.W., Baskin, D.G., 2009. Forebrain melanocortin signaling enhances the hindbrain satiety response to CCK-8. *Am. J. Physiol.* 296, R476-R484.
- Boswell, T., Li, Q., Takeuchi, S., 2002. Neurons expressing neuropeptide Y mRNA in the infundibular hypothalamus of Japanese quail are activated by fasting and co-express agouti-related protein mRNA. *Mol. Brain Res.* 100, 31-42.
- Byerly, M.S., Simon, J., Lebihan-Duval, E., Duclos, M.J., Cogburn, L.A., Porter, T.E., 2009. Effects of BDNF, T3, and corticosterone on expression of the hypothalamic obesity gene network *in vivo* and *in vitro*. *Am. J. Physiol.* 296, R1180-R1189.
- Cerasale, D.J., Zajac, D.M., Guglielmo, C.G., 2011. Behavioral and physiological effects of photoperiod-induced migratory state and leptin on a migratory bird, *Zonotrichia albicollis*: I. Anorectic effects of leptin administration. *Gen. Comp. Endocrinol.* 174, 276-286.
- Christensen, K., McMurtry, J.P., Thaxton, Y.V., Corzo, A., McDaniel, C., Scanes, C.G., 2013. Metabolic and hormonal responses of growing modern meat-type chickens to fasting. *Brit. Poultry Sci.* 54, 199-205.
- Cornelius, J.M., Boswell, T., Jenni-Eiermann, S., Breuner, C.W., Ramenofsky, M., 2013. Contributions of endocrinology to the migration life history of birds. *Gen. Comp. Endocrinol.* 190, 47-60.
- Crespi E.J., Denver, R.J., 2006. Leptin (*ob* gene) of the South African clawed frog *Xenopus laevis*. *Proc. Nat. Acad. Sci. USA* 103, 10092-10097.
- Denbow, D.M., Meade, S., Robertson, A., McMurtry, J.P., Richards, M., Ashwell, C., 2000. Leptin-induced decrease in food intake in chickens. *Physiol. Behav.* 69, 359-362.
- Dridi, S., Swennen, Q., Decuypere, E., Buyse, J., 2005. Mode of leptin action in chicken hypothalamus. *Brain Res.* 1047, 214-223.
- Dunn, I.C., Wilson, P.W., Smulders, T.V., Sandilands, V., D'Eath, R.B., Boswell, T., 2013a. Hypothalamic agouti-related protein expression is affected by both acute and chronic experience of food restriction and re-feeding in chickens. *Journal of Neuroendocrinology* 25, 920-928.
- Dunn, I.C., Meddle, S.L., Wilson, P.W., Wardle, C.A., Law, A.S., Bishop, V.R., Hindar, C., Robertson, G.W., Burt, D.W., Ellison, S.J.H., Morrice, D.M., Hocking, P.M., 2013b. Decreased expression of the satiety signal receptor CCKAR is responsible for increased growth and body weight during the domestication of chickens. *Am. J. Physiol.* 304, E909-E921.
- Ebling, F.J.P., 2014. On the value of seasonal mammals for identifying mechanisms underlying the control of food intake and body weight. *Horm. Behav.* 66, 56-65.
- Friedman, J.M., 2009. Leptin at 14 y of age: an ongoing story. *Am. J. Clin. Nutr.* 89, 973S-979S.
- Friedman-Einat, M., Seroussi, E., 2014. Quack leptin. *BMC Genomics* 15:551.

416 Friedman-Einat, M., Boswell, T., Horev, G., Girishvarma, G., Dunn, I.C., Talbot, R.T.,
 417 Sharp, P.J., 1999. The chicken leptin gene: has it been cloned? Gen. Comp.
 418 Endocrinol. 115, 354-363.
 419 Friedman-Einat, M., Cogburn, L.A., Yosefi, S., Hen, G., Shinder, D., Shirak, A., Seroussi,
 420 E., 2014. Discovery and characterization of the first genuine avian leptin gene in
 421 the rock dove (*Columba livia*). Endocrinology 155, 3376-3384.
 422 Gertler, A., Shindler, D., Yosefi, S., Shpilman, M., Rosenblum, C.I., Ruzal, M., Seroussi,
 423 E., Friedman-Einat, M., 2014. Pegylated leptin antagonist with strong orexigenic
 424 activity in mice is not effective in chickens. J. Exp. Biol. 217, 180-184.
 425 Harvey, S., Klandorf, H., 1983. Reduced adrenocortical function and increased thyroid
 426 function in fasted and refed chickens. J. Endocrinol. 98, 129-135.
 427 Hen, G., Yosefi, G., Simchaev, V., Shinder, D., Hruby, V.J., Friedman-Einat, M., 2006.
 428 The melanocortin circuit in obese and lean strains of chicks. J. Endocrinol. 190,
 429 527-535.
 430 Hen, G., Yosefi, G., Ronin, A., Einat, P., Rosenblum, C.I., Denver, R.J., 2008.
 431 Monitoring leptin activity using the chicken leptin receptor. J. Endocrinol. 197,
 432 325-333.
 433 Higgins, S.E., Ellestad, L.E., Trakooljul, N., McCarthy, F., Saliba, J., Cogburn, L.A.,
 434 Porter, T.E., 2010. Transcriptional and pathway analysis in the hypothalamus of
 435 newly hatched chicks during fasting and delayed feeding. BMC Genomics 11, 162.
 436 Honda, K., Kamisoyama, H., Saneyasu, T., Sugahara, K., Hasegawa, S., 2007. Central
 437 administration of insulin suppresses food intake in chicks. Neurosci. Lett. 423, 153-
 438 157.
 439 Huang, G., Li, J., Wang, H., Lan, X., Wang, Y., 2014. Discovery of a novel functional
 440 leptin protein (LEP) in zebra finches: evidence for the existence of an authentic
 441 avian leptin gene predominantly expressed in the brain and pituitary.
 442 Endocrinology 155, 3389-3396.
 443 Kaiya, H., Kangawa, K., Miyazato, M., 2013. Update on ghrelin biology in birds. Gen.
 444 Comp. Endocrinol. 190, 170-175.
 445 King, J.R., Barker, S., Farner, D.S., 1963. A comparison of energy reserves during
 446 autumnal and vernal migratory periods in the white-crowned sparrow, *Zonotrichia*
 447 *leucophrys gambelii*. Ecology 44, 513-521.
 448 Liu, J., Garza, J.C., Truong, H.V., Henschel, J., Zhang, W., Lu, X.Y., 2007. The
 449 melanocortinergic pathway is rapidly recruited by emotional stress and contributes
 450 to stress-induced anorexia and anxiety-like behavior. Endocrinology 148, 5531-
 451 5540.
 452 Liu, X., Dunn, I.C., Sharp, P.J., Boswell, T., 2007. Molecular cloning and tissue
 453 distribution of a short form chicken leptin receptor mRNA. Domest. Anim.
 454 Endocrinol. 32, 155-166.
 455 Liu, L., Song, Z., Sheikahmadi, A., Jiao, H., Lin, H., 2012. Effect of corticosterone on
 456 gene expression of feed intake regulatory peptides in laying hens. Comp. Biochem.
 457 Physiol. B 162, 81-87.
 458 Löhms, M., Sundström, L.F., El Halawani, M., Silverin, B., 2003. Leptin depresses food
 459 intake in great tits (*Parus major*). Gen. Comp. Endocrinol. 131, 57-61.

460 Makimura, H., Mizuno, T., Isoda, F., Beasley, J., Silverstein, J., Mobbs, C., 2003. Role of
 461 glucocorticoids in mediating effects of fasting and diabetes on hypothalamic gene
 462 expression. *BMC Physiol.* 3, 5.
 463 Millar, R.P., 2014. Identification of genuine/authentic avian leptin: some answers and
 464 more questions. *Endocrinology* 155, 3203-3205.
 465 Mrosovsky, N., 1990. *Rheostasis: The Physiology of Change*. Oxford University Press,
 466 New York.
 467 Mrosovsky, N., Sherry, D.F., 1980. Animal anorexias. *Science* 207, 837-842.
 468 Pasquier, J., Lafont, A.G., Rousseau, K., Quérat, B., Chemineau, P., Dufour, S., 2014.
 469 Looking for the bird Kiss: evolutionary scenario in sauropsids. *BMC Evol. Biol.* 14,
 470 30.
 471 Phillips-Singh, D., Li, Q., Takeuchi, S., Ohkubo, T., Sharp, P.J., Boswell, T., 2003.
 472 Fasting differentially regulates expression of agouti-related peptide, pro-
 473 opiomelanocortin, prepro-orexin, and vasoactive intestinal polypeptide mRNAs in
 474 the hypothalamus of Japanese quail. *Cell Tissue Res.* 313, 217-225.
 475 Pitel, F., Faraut, T., Bruneau, G., Monget, P., 2010. Is there a leptin gene in the chicken
 476 genome? Lessons from phylogenetics, bioinformatics and genomics. *Gen. Comp.*
 477 *Endocrinol.* 167, 1-5.
 478 Porte, D. Jr., Baskin, D.G., Schwartz, M.W., 2002. Leptin and insulin action in the central
 479 nervous system. *Nutr. Rev.* 60, S20-S29.
 480 Prokop, J.W., Schmidt, C., Gasper, D., Duff, R.J., Milsted, A., Ohkubo, T., Ball, H.C.,
 481 Shawkey, M.D., Mays Jr., H.L., Cogburn, L.A., Londraville, R.L., 2014. Discovery
 482 of the elusive leptin in birds: identification of several 'missing links' in the
 483 evolution of leptin and its receptor. *PLoS ONE* 9(3), e92751.
 484 Proszkowiec-Weglarz, M., Richards, M.P., Ramachandran, R., McMurtry, J.P., 2006.
 485 Characterization of the AMP-activated protein kinase pathway in chickens. *Comp.*
 486 *Biochem. Physiol. B* 143, 92-106.
 487 Raver, N., Taouis, M., Dridi, S., Derouet, M., Simon, J., Robinzon, B., Djiane, J., Gertler,
 488 A., 1998. Large-scale preparation of biologically active recombinant chicken obese
 489 protein (leptin). *Protein Expr. Purif.* 14, 403-408.
 490 Richards, M.P., Caperna, T.J., Elsasser, T.H., Ashwell, C.M., McMurtry, J.P., 2000.
 491 Design and application of a polyclonal peptide antiserum for the universal detection
 492 of leptin protein. *J. Biochem. Biophys. Methods* 45, 147-156.
 493 Rønnestad, I., Nilsen, T.O., Murashita, K., Angotzi, A.R., Gamst Moen, A.-G.,
 494 Steffenson, S.O., Kling, P., Thrandur Björnsson, B., Kurokawa, T., 2010. Leptin
 495 and leptin receptor genes in Atlantic salmon: cloning, phylogeny, tissue distribution
 496 and expression correlated to long-term feeding status. *Gen. Comp. Endocrinol.* 168,
 497 55-70.
 498 Saito, E.S., Kaiya, H., Tachibana, T., Tomonaga, S., Denbow, D.M., Kangawa, K.,
 499 Furuse, M., 2005. Inhibitory effect of ghrelin on food intake is mediated by the
 500 corticotropin-releasing factor system in neonatal chicks. *Regul. Pept.* 15, 201-208.
 501 Savory, C.J., 1979. Changes in food intake and body weight of bantam hens during
 502 breeding. *Appl. Anim. Ethol.* 5, 283-288.
 503 Sherry, D.F., Mrosovsky, N., Hogan, J.A., 1980. Weight loss and anorexia during
 504 incubation in birds. *J. Comp. Physiol. Psychol.* 94, 89-98.

- Shiraishi, J., Yanagita, K., Fujita, M., Bungo, T., 2008. Central insulin suppresses feeding behavior via melanocortins in chicks. *Domest. Anim. Endocrinol.* 34, 223-228.
- Shiraishi, J., Tanizawa, H., Fujita, M., Kawakami, S., Bungo, T. 2011. Localization of hypothalamic insulin receptor in neonatal chicks: evidence for insulinergic system control of feeding behavior. *Neurosci. Lett.* 491, 177-180.
- Song, Z., Liu, L., Yue, Y., Jiao, H., Lin, H., Sheikahmadi, A., Everaert, N., Decuypere, E., Buyse, J., 2012. Fasting alters protein expression of AMP-activated protein kinase in the hypothalamus of broiler chicks (*Gallus gallus domesticus*). *Gen. Comp. Endocrinol.* 178, 546-555.
- Strader, A.D., Buntin, J.D., 2003. Changes in agouti-related peptide during the ring dove breeding cycle in relation to prolactin and parental hyperphagia. *J. Neuroendocrinol.* 15, 1046-1053.
- Strader, A.D., Schiöth, H.B., Buntin, J.D., 2003. The role of the melanocortin system and the melanocortin-4 receptor in ring dove (*Streptopelia risoria*) feeding behavior. *Brain Res.* 960, 112-121.
- Tachibana, T., Sugahara, K., Ohgushi, A., Ando, R., Kawakami, S., Yoshimatsu, T., Furuse, M., 2001. Intracerebroventricular injection of agouti-related protein attenuates the anorexigenic effect of alpha-melanocyte stimulating hormone in neonatal chicks. *Neurosci. Lett.* 305, 131-134.
- Takeuchi, S., Suzuki, S., Hirose, S., Yabuuchi, M., Sato, C., Yamamoto, H., Takahashi, S., 1996. Molecular cloning and sequence analysis of the chick melanocortin 1-receptor gene. *Biochim. Biophys. Acta* 1306, 122-126.
- Takeuchi, S., Kudo, T., Takahashi, S., 1998. Molecular cloning of the chicken melanocortin 2 (ACTH)- receptor gene. *Biochim. Biophys. Acta.* 1403, 102-108.
- Takeuchi, S., Takahashi, S., 1998. Melanocortin receptor genes in the chicken – tissue distributions. *Gen. Comp. Endocrinol.* 112, 220-231.
- Takeuchi, S., Takahashi, S., 1999. A possible involvement of melanocortin 3 receptor in the regulation of adrenal gland function in the chicken. *Biochim. Biophys. Acta* 1448, 512-518.
- Takeuchi S., Teshigawara, K., Takahashi, S., 1999. Molecular cloning and characterization of the chicken pro-opiomelanocortin (POMC) gene. *Biochim. Biophys. Acta* 1450, 452-459.
- Takeuchi, S., Teshigawara, K., Takahashi, S., 2000. Widespread expression of agouti-related protein (AGRP) in the chicken: a possible involvement of AGRP in regulating peripheral melanocortin systems in the chicken. *Biochim. Biophys. Acta* 1496, 261-269.
- Yosefi, S., Hen, G., Rosenblum, C.I., Cerasale, D.J., Beaulieu, M., Criscuolo, F., Friedman-Einat, M., 2010. Lack of leptin activity in blood samples of Adelie penguin and bar-tailed godwit. *J. Endocrinol.* 207, 113-122.

Figure legend

Figure 1. Summary of current knowledge of regulatory inputs (positive and negative influences are indicated) on agouti-related protein (AGRP) and pro-opiomelanocortin (POMC) gene expression in the arcuate nucleus of the avian hypothalamus. Signalling within the hypothalamus is shown in the box. Abbreviations: ADRB2 - β 2 adrenergic receptor; AMPK – AMP-activated protein kinase; GRM8 – metabotropic glutamate receptor 8; NPY5R – neuropeptide Y receptor Y5; RLN3 – relaxin-3; SSTR5 – somatostatin receptor 5.