THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Using satellite estimates of aboveground biomass to assess carbon stocks in a mixed-management, semi-deciduous tropical forest in the Yucatan Peninsula

Citation for published version:

George-Chacón, SP, Milodowski, D, Dupuy, JM, Max, J-F, Williams, M, Castillo-Santiago, M & Hernández-Stefanoni, JL 2021, 'Using satellite estimates of aboveground biomass to assess carbon stocks in a mixed-management, semi-deciduous tropical forest in the Yucatan Peninsula', *Geocarto International*.
<https://doi.org/10.1080/10106049.2021.1980619>

Digital Object Identifier (DOI):

[10.1080/10106049.2021.1980619](https://doi.org/10.1080/10106049.2021.1980619)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Geocarto International

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Using satellite estimates of aboveground biomass to assess carbon stocks in a mixed-management, semi-deciduous tropical forest in the Yucatan Peninsula

Stephanie P. George-Chacón¹, David T. Milodowski², Juan Manuel Dupuy¹, Jean-François Mas³, Mathew Williams^{2,4}, Miguel Castillo-Santiago⁵, José Luis Hernández-Stefanoni^{1*}.

1. Centro de Investigación Científica de Yucatán A.C. Unidad de Recursos Naturales, Calle 43 # 130. Colonia Chuburná de Hidalgo. C.P. 97200, Mérida, Yucatán. México.

2. School of GeoSciences, University of Edinburgh, Edinburgh EH9 3FF, United Kingdom

3. Laboratorio de análisis espacial, Centro de Investigaciones en Geografía Ambiental, Universidad Nacional Autónoma de México, 5 Campus Morelia, Antigua Carretera a Pátzcuaro 8701, Col. Ex-Hacienda de San José de La Huerta, C.P. 58190 Morelia, México.

4. National Centre for Earth Observation, University of Edinburgh, Edinburgh EH9 3FF, United Kingdom

5. El Colegio de la Frontera Sur, Laboratorio de Análisis de Información Geográfica y Estadística, Carretera Panamericana y Periférico sur s/n., San Cristóbal de las Casas, CP 29290 Chiapas, Mexico.

* Corresponding author, e-mail: jl_stefanoni@cicy.mx

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding

This work was funded by Ecometrica LTD and the UK space agency under project Forests 2020.

27 **Keywords**

28 Remote sensing, machine learning, LiDAR, error propagation, Sentinel-2, ALOS PALSAR

29

30 **Abstract**

31 Information on the spatial distribution of forest aboveground biomass (AGB) and its uncertainty is important to
32 evaluate management and conservation policies in tropical forests. However, the scarcity of field data and robust
33 protocols to propagate uncertainty prevent a robust estimation through remote sensing. We upscaled AGB from
34 field data to LiDAR, and to landscape scale using Sentinel-2 and ALOS-PALSAR through machine learning,
35 propagated uncertainty using a Monte Carlo framework and explored the relative contributions of each sensor.
36 Sentinel-2 outperformed ALOS-PALSAR ($R^2 = 0.66$, vs 0.50), however, the combination provided the best fit (R^2
37 $= 0.70$). The combined model explained 49% of the variation comparing against plots within the calibration area,
38 and 17% outside, however, 94% of observations outside calibration area fell within the 95% confidence intervals.
39 Finally, we partitioned the distribution of AGB in different management and conservation categories for evaluating
40 the potential of different strategies for conserving carbon stock.

Introduction

Tropical forests hold large stocks of carbon and play a key role in the global carbon cycle and its interactions with climate (Bonan et al., 2008; Pan et al. 2011; Mitchard, 2018). Carbon contained in aboveground biomass (AGB) is most susceptible to be emitted through deforestation and degradation, which are important sources of emissions in tropical forests (Houghton et al., 2005; Houghton, 2012; 2013). Accurate estimation of the spatial distribution of AGB and its uncertainty is an important part of the implementation of strategies aimed at reducing emissions through deforestation and degradation throughout the tropics, such as REDD+. However, previous research has underestimated the uncertainty due to inadequate methods for estimating and propagating errors throughout the estimation process (Yunai et al., 2020).

The Yucatan Peninsula, located in the south-eastern part of Mexico, holds one of the largest extents of continuous tropical dry forest in Latin America (Dupuy et al., 2015). Mexico is an active participant in the REDD+ initiative, and particularly, the Yucatan Peninsula is a part of the REDD+ early actions priority areas, due to increasing pressure of permanent conversion to urban areas and the expansion of agriculture (Ellis et al., 2017). Several protected areas targeting conservation of forest resources are located within the Yucatan Peninsula (CONANP, 2017) encompassing old-growth forests. However, the extensive use of traditional agricultural practices, such as slash-and-burn agriculture, as well as other land use such as agricultural areas and pastures for cattle ranching, shapes the landscape outside the protected areas into a mosaic of forest areas in diverse stages of natural regeneration, with biodiversity and forest biomass gradually recovering after abandonment (Dupuy et al., 2012). In line with global policy efforts to restore forests across the tropics, significant areas of the Yucatan Peninsula have been allocated for restoration (CONANP, 2017). Whether this restoration will lead to significant carbon sequestration, and thus help mitigate climate change, will depend on the balance between forest loss through deforestation and degradation (including the exploitation of forest resources) and forest gain from forest regrowth from conservation and natural regeneration of disturbed areas (Houghton, 2013; Chazdon et al., 2016; Lewis et al., 2019).

The balance between forest (re)growth and disturbance determines the distribution of AGB (Williams et al., 2013). Post-disturbance, forest ecosystems can aggrade, accumulating carbon until they reach a quasi-steady state, where gains through growth and recruitment become balanced by mortality losses. At steady state the distribution of AGB within the landscape tends towards a normal distribution (Williams et al., 2013). In disturbed forests, on the other hand, repeated removal of AGB results in a skewed distribution of AGB, resulting in a long tail of low AGB values. Therefore, information on the distribution of AGB can be used to assess the state of AGB stocks in areas under different management strategies.

Reliable estimates of the spatial distribution of forest AGB are essential for effective forest management, to detect areas of loss and assess the success of conservation efforts. To date, AGB across Mexico has been mapped in a number of National (Cartus et al., 2014; Rodriguez-Veiga et al., 2016; Urbazev et al., 2018) and pan-tropical (Saatchi et al., 2011; Baccini et al., 2012., Avitabile et al., 2016) products. However, there are large and systematic uncertainties (Mitchard et al., 2013) with existing maps, which tend to underestimate the AGB in the Yucatan Peninsula (Rodriguez-Veiga et al., 2019; Hernández-Stefanoni et al., 2020), leading to potential underestimation of carbon emissions from deforestation and degradation.

Production of regional AGB maps typically relies on upscaling field estimates of AGB based on a relationship between a network of field inventory plots (Chave et al., 2004; Réjou-Méchain et al., 2019) and remotely sensed data (Goetz et al., 2015). A number of passive sensors (e.g. multispectral optical imagery from Sentinel 2) and active sensors (e.g. L-band Synthetic Aperture Radar (SAR) from Advanced Land Observation Satellite) (ALOS PALSAR; Shimada, 2010) are available that offer frequent coverage at global scales. Each sensor has its own limitations. Optical data are limited by cloud and smoke, both common in tropical forests (Asner et al., 2001); SAR penetrates clouds, and polarized backscatter has been shown to be sensitive to AGB (Mermoz et al., 2015, Thapa et al., 2015, Mitchard et al., 2009), however, optical and L-band saturate at $\sim 150 \text{ Mg ha}^{-1}$ (Lu et al., 2006; Mitchard et al., 2009; Joshi et al., 2017). Compared to tropical wet forests, old-growth tropical dry forest canopies are generally shorter and simpler, and AGB correspondingly lower (Murphy and Lugo 1986). Therefore, the AGB range occupied by tropical dry forests is potentially still within the sensitivity range of L-band systems. Multi-sensor approaches can leverage the strengths of these various data sources to improve AGB estimates (Bispo et al., 2020).

Generating maps of AGB based on satellite data requires calibration against estimations of AGB typically taken from field inventories (e.g. Rodriguez-Veiga et al., 2016; Saatchi et al., 2011; McNicol et al., 2018). High-resolution airborne LiDAR surveys offer the potential to bridge the scale gap between inventory plots and satellite data and enhance the range of training sites over which to calibrate models (Urbazev et al., 2016; Wulder et al., 2012; Asner et al., 2018; Bispo et al., 2020). LiDAR is particularly powerful as it captures precise information on forest structure without signal saturation in dense tropical forests (Lefsky et al., 1999; Asner et al., 2014). However, the cost of obtaining airborne LiDAR data through on-demand surveys is high. Consequently, publicly available data are typically scarce over many tropical forests. The GEDI mission offers global open, spatially distributed waveform LiDAR (Dubayah et al., 2020), which will undoubtedly facilitate calibration of satellite-based biomass products (e.g. Qi and Dubayah, 2016). However, GEDI has a nominal mission lifetime of two years from its on-orbit checkout in April 2019, thus limiting its scope for future and past monitoring of change in tropical forests. Therefore, it is important to develop methods that utilize spatially limited airborne surveys inside

upscaling frameworks and quantify their predictive uncertainty with robust error estimation (Zhao et al., 2020). In developing upscaling frameworks, particularly when working with spatially limited data, it is critical to account for spatial autocorrelation to avoid overfitting and thus greatly overstating the predictive power of upscaled models (Roberts et al., 2017; Ploton et al., 2020).

This research has three core aims: (i) to produce accurate spatially explicit estimations of AGB and its uncertainty in a semi-deciduous tropical dry forest of the Yucatan Peninsula; (ii) to quantify the effectiveness of active and passive sensors and their combination for achieving (i); (iii) to use the spatial distribution of AGB to inform on the state of carbon stock of forest areas under different management and conservation conditions. We develop an upscaling framework that uses airborne LiDAR surveys as an intermediate step to link field inventory AGB estimates to Sentinel 2 and ALOS PALSAR data. First, we generate a LiDAR AGB model, AGB_{LiDAR} , calibrated using field inventory data. Subsequently, we use a machine-learning framework to upscale these AGB_{LiDAR} maps with satellite data from Sentinel 2 and ALOS PALSAR to generate a satellite-based model for AGB, AGB_{SAT} . Previous studies suggest image texture metrics can improve estimates of AGB in dense forests (Castillo et al., 2005; Wood et al., 2012; Thapa et al., 2015; Hernández-Stefanoni et al., 2020). We therefore explore the potential for texture variables to improve the predictive power of our machine-learning models. We assess the effect of spatial resolution in the calibration of the LiDAR-to-satellite model and explore the improvement in performance of multi-sensor models over single-sensor models. We propagate uncertainty through the analysis using a Monte Carlo framework, including a spatially independent cross-validation strategy for robust estimates of errors arising during upscaling (e.g. Roberts et al., 2017). Finally, we use the AGB_{SAT} map to gain insight into the impact of forest management (production vs. protection) on forest biomass, and thus the likely carbon sequestration potential for areas set aside for restoration in this region.

Methods

Study Area

The study area comprises 3600 km² of tropical dry forest in the centre of the Yucatan Peninsula, Mexico, located between 20° 09' 39" and 19° 37' 08" N latitude and 89° 16' and 89° 50' 36" W longitude (Figure 1). The vegetation at this site is predominately semi-deciduous tropical dry forest, sitting in the transition zone between deciduous tropical dry forest in the drier northern part of the Peninsula and semi-evergreen tropical forest in the south-west (Rzedowski 2006). Trees in this region are typically 8–15 m tall, and 50–75 % of trees drop their leaves during the dry season, which typically falls between November and April (Carnevali et al., 2003). The limestone terrain underlying this region is characterized by a mixture of low hills (elevation range: 16–216 m) and flat areas. Three protected natural reserves exist within the study area: [Kaxil Kiuic Biocultural Reserve \(Reserva Biocultural Kaxil Kiuic\)](#) (1,800

ha) a private reserve located inside a state protected area: [del Puuc Biocultural reserve \(Reserva Estatal Biocultural del Puuc\)](#) (135,849 ha), and a small fraction (~ 5,000 ha) of the Bala'an K'aax national protected area (128,390 ha) (CONANP 2017) (Figure 1). Several low impact subsistence activities occur in the adjacent forest surrounding the Kaxil Kiuic reserve (swidden agriculture, with some selective logging and cattle grazing) and agricultural fields. Unprotected forest areas are subdivided into areas suitable for production of forest species and areas suitable for forest restoration. These areas were designated according to structural characteristics such various degrees of degradation in the restoration forest and tree cover for production forest. For a detailed description refer to CONAFOR (2013).

[insert Figure 1 around here]

Field inventory data

Field data were taken from two surveys: (i) The Intensive Carbon Monitoring (ICM) site; (ii) a sparser, spatially more extensive dataset across the region, obtained from the Mexican National Forest Inventory (NFI). The majority of plots (20) from the ICM are located within the Kaxil Kiuic Biological Reserve, and 12 are placed outside the reserve boundary in a chronosequence in several ages of abandonment. (Figure 1). [In both cases, plots are composed of clusters of four GPS-located circular subplots of 400 m².](#) The plots are distributed systematically with one central plot surrounded by three peripheral plots at 90°, 120° and 240° azimuths within a 1 ha sampling area (CONAFOR 2013). Within each plot, height and Diameter at Breast Height (DBH Diameter at 1.30 m) were recorded for all woody plants with DBH > 7.5 cm and each individual was identified to species level. In addition, small stems (2.5 cm ≤ DBH < 7.5 cm) were also measured at the ICM plots, within a central subplot of 80 m² (Caamal-Sosa et al., 2016). AGB was calculated for each tree using the allometric equation of Chave et al. (2005) for trees with DBH ≥ 10 cm, and that of Ramirez et al., (2017), for trees with DBH < 10 cm, based on DBH and height from the above-mentioned datasets. Wood densities were taken from Sanaphre-Villanueva et al. (2016) where species were present in the database, otherwise a mean value of wood density at the genus or the plot level were used. Plot AGB was estimated based as the sum of the AGB of all individual trees. In the ICM plots, the contribution of small stems averaged 24.4 ± 13.5 Mg ha⁻¹. This contribution was added to the NFI plots to standardize the two datasets. [In total, 33 plots \(132 subplots\)](#) fell within the LiDAR survey. A further [435 subplots](#) fell outside the survey, providing independent validation of the final upscaled map outside the LiDAR survey.

[The workflow of methods applied in this research is displayed in supplementary material 1 and described in more details in the following sections.](#)

LiDAR data

We obtained LiDAR data from NASA's Goddard's LiDAR, Hyperspectral and Thermal (G-LiHT) airborne imager (Cook et al., 2013) available for the study area (Figure 1). The LiDAR point cloud was pre-processed using the USFS FUSION software (McGaughey et al., 2012) resulting in two 1-m resolution raster representing the top of canopy elevation and the underlying topography. The difference in elevation between these surfaces provides a direct estimate of canopy height.

ALOS PALSAR and Sentinel-2 data processing

Two scenes of Advanced Land Observation Satellite Phased Array L-Band Synthetic Aperture Radar (ALOS PALSAR) yearly mosaics at 25 m spatial resolution for the year 2015 were merged to cover the extent of the study area. The images, obtained in digital numbers, were converted to backscatter coefficient by means of the formula provided by Shimada et al. (2010). Afterward, they were pre-processed to obtain gridded, topographically corrected backscatter amplitudes for HH and HV polarizations (Mitchard et al., 2009). The ALOS PALSAR backscatter was processed to remove "speckle" (Woodhouse, 2017) using the standard enhanced Lee filter (Lee 1980), as implemented in the GIS software package ENVI 5.0 (Hernández-Stefanoni et al., 2020).

Two Sentinel 2A scenes corresponding to April 2017 were mosaicked using linear normalization in order to produce a seamless mosaic of the study area. We used the following bands: blue (492.4 nm, hereafter named as Band 1), green (559.8 nm, Band 2), red (664.6 nm, Band 3) and near infrared (832.8 nm, NIR, Band 4) with a spatial resolution (pixel size) of 10 m. Also, we calculated the Normalized difference vegetation index (NDVI).

We also used image texture metrics such as Gray Level Co-occurrence Matrix (GLCM Haralick et al., 1979), since they are able to capture the spatial variability in the spectral response of different elements in the landscape and have been related by previous work to variability in forest structure (Gallardo-Cruz et al., 2012, Wood et al., 2012). These statistics can be categorized into homogeneity and heterogeneity metrics. Higher values in metrics such as contrast and dissimilarity indicate a higher variability in the elements in an area, whereas metrics such as homogeneity, second moment and correlation, indicate similarity within an area. The mean and variance of the surface reflectance of bands in addition to the aforementioned GLCM measures (hereby texture measures) were calculated at the spatial resolution of the LiDAR-satellite upscaling step for all individual bands and for NDVI using scikit-image, a collection of algorithms for image processing in python 3.6 (Van der Walt et al., 2014).

Upscaling field inventory to regional AGB

In order to estimate the spatial distribution of AGB we carried out a two-step process: (1) creation of the LiDAR AGB map, AGB_{LiDAR} at 20 m resolution, corresponding to the resolution of the

individual 0.04 ha inventory plots (AGB_{Field}); (2) upscaling AGB_{LiDAR} across the study area using machine learning models based on data from Sentinel 2 and/or ALOS PALSAR to produce AGB_{SAT} .

Spatial mapping of AGB with LiDAR

The first step in upscaling the field inventory AGB estimates (AGB_{Field}) was to extrapolate these across the LiDAR survey extent. To do this we fitted a power law relationship between the AGB of the 0.04 ha inventory plots and the mean top of canopy height (TCH) measured by the LiDAR sensor within the footprint of each 0.04 plot (Figure 2). This follows from the allometric expectation of power law scaling of AGB with tree height, and therefore stand height (e.g. Asner and Mascaro, 2014). To reduce the risk of bias in canopy height estimates from areas of low point density (Roussel et al., 2017), we filtered out areas of the survey with less than 6 pts m^2 . We also investigated alternative variants and canopy metrics, including gap fraction (e.g. Jucker et al., 2017), but these did not lead to significant overall improvement in the model under leave-one-out (LOO) cross validation.

To model the power law relationship, we fitted a linear mixed effects model in log-transformed space to account for the hierarchical structure of the inventory data (i.e. four 0.04 ha plots within each plot cluster):

$$[ln(AGB_{LiDAR})]_{i,j} = \alpha + \beta * [ln(TCH)]_{i,j} + u_i + \varepsilon_{i,j},$$

where i represents the plot cluster, j represents the plot within the cluster, α is the intercept term, β is the fixed effect for $ln(TCH)$, u_i represents a random effect associated with the plot cluster i , and $\varepsilon_{i,j}$ represents the residuals for each plot. Finally, after back-transformation of the final estimates, we applied the necessary correction factor (Baskerville, 1972):

$$CF = \exp\left(\frac{\sigma^2}{2}\right).$$

where σ^2 is the RMSE of the model fit in log-space. The RMSE under Leave-One-Out (LOO) cross validation (Supplementary 2) was 46.14 $Mg\ ha^{-1}$ and the R^2 was 0.40, for a spatial resolution of 0.04 ha. Relatively high RMSE values are in line with expectations for small plot sizes (e.g. Mascaro et al., 2011), but relative errors should drop considerably when aggregating across larger regions (Gonzalez et al., 2010).

Upscaling AGB with satellite data

In order to produce spatially explicit estimations of AGB in the Kiuc landscape we upscaled the AGB_{LiDAR} map with the satellite data using random forest regression (Breiman, 2001), with a bootstrap bias correction (Hooker and Mentch, 2018; Xu et al., 2016). Random forest regression is a flexible, non-parametric machine learning algorithm that has previously been employed to fuse LiDAR and satellite data and produce maps of AGB and other structural parameters (e.g. Luther et al., 2019;

Mascaro et al., 2014; Urbazaev et al., 2018; Wulder et al., 2012). Random forest models were fitted using the implementation of scikit-learn in Python (Pedregosa et al., 2011). To optimize the random forest regression models, we employed a Bayesian hyperparameter search seeded with 100 random trials, followed by a further 350 iterations (Bergstra et al., 2011). To determine the best spatial resolution at which to undertake the LiDAR-satellite upscaling, we tested the effect of aggregating to three different spatial resolutions (20 m, 50 m and 100 m). The relative importance of the sensors and textures to explain the variation in AGB in the fitted models was explored based on the drop in R^2 following permutation of each variable (permutation importance e.g. Strobl et al., 2007). Given the strong collinearities between texture metrics for different bands, we permuted all variables associated with (i) each sensor, and (ii) each texture index, to capture their contributions more concisely. Finally, we compared the performance of the combined Sentinel 2/ALOS PALSAR model against single-sensor models to investigate the improvement in predictive power provided by the complementary attributes of these sensors.

Error propagation

Robust characterization of uncertainty is critical to understanding the utility and limitations of remotely sensed maps of AGB (Ploton et al., 2020). Uncertainty arises from a multitude of factors. Uncertainties in the field AGB estimates (Chave et al., 2004), combined with spatial registration errors (Hernández-Stefanoni et al., 2018), crown overlap at plot boundaries (Mascaro et al., 2011); and temporal lags (Babcock et al., 2016; Clark and Kellner, 2012) lead to uncertainties in AGB_{LiDAR} . These uncertainties are compounded by unexplained variance in the subsequent LiDAR-satellite upscaling model. In addition, geospatial data are frequently spatially autocorrelated. In scenarios like this one, the clustered geometry of the available LiDAR data survey precludes the robust inclusion of a spatial effect into the random forest models through additional spatial covariates (e.g. Mascaro et al., 2014). Spatial autocorrelation, if not accounted for, can lead to overfitting resulting in significant underestimation in predictive error during cross-validation and misleading diagnostic analyses regarding feature importance (Roberts et al., 2017; Ploton et al., 2020).

In order to propagate uncertainty in the upscaling process we employed Monte-Carlo simulations to propagate errors across every step of the upscaling framework. Uncertainty in AGB_{field} was estimated based on estimates of uncertainty in the biomass of individual trees, assumed to be 47% of tree AGB (see Chave et al., 2004). Uncertainties between trees were assumed to be independent and thus they were aggregated at the plot-level by adding in quadrature (the square root of the sum of squares), a standard procedure for combining uncorrelated errors (Yanai et al., 2020). Relative errors at the plot level were therefore significantly lower and tended to be dominated by the largest trees. To characterize the uncertainty in AGB_{LiDAR} , we fitted the mixed effects model 100 times. In each iteration we resampled the biomass of AGB_{field} assuming normally distributed uncertainties. We accounted for

spatial registration errors by shifting the plot location randomly assuming a standard deviation of 5 m in the plot coordinates. Corresponding uncertainties in TCH were strongly non-normal in some cases, particularly close to forest edges. We did not attempt to account for canopy overlap, or temporal lags. Fitting the model 100 times produced 100 candidate AGB_{LiDAR} maps for upscaling with the satellite data.

To propagate uncertainty across the LiDAR-satellite step, the 100 AGB_{LiDAR} maps were used as the target for an ensemble of 100 random forest models. To account for predictive uncertainty of these models, we also fitted a model to predict the median AGB_{LiDAR} using a 16-fold buffered, blocked cross validation procedure, whereby the training data were split into square blocks (block width 1 km), and randomly allocated to one of the folds. In each iteration, we buffered the validation set by a distance of 500 m to reduce the impact of spatial autocorrelation and therefore minimize overfitting (Note that in the optimization and feature importance calculations, only five folds were used to reduce processing time). This spatial cross-validation was undertaken for the three tested upscaling resolutions (20 m, 50 m and 100 m) to determine the best option for upscaling (Figure 4). Errors in predicted AGB resulting from fitted spatial correlations were modelled by resampling from the residuals from the results of this cross-validation (using median AGB_{LiDAR}). As the residuals were not uniformly distributed along the range of predicted AGB (AGB_{upscaled}), residuals were resampled from a 20 Mg ha⁻¹ window around the AGB estimate for each pixel. Thus, the 100 x 100 iterations of the upscaling procedure capture both uncertainty in AGB_{LiDAR} propagated through the random forest models, and the predictive uncertainty associated with fitting models with spatially autocorrelated data. We present the median and 95% confidence intervals as our best estimates and uncertainty in the upscaled AGB maps (AGB_{upscaled}).

Comparison with other work

The AGB map obtained in this study was compared with previous AGB maps generated by Santoro et al., (2018), Rodriguez-Veiga et al., (2016) and Cartus et al., (2014). We performed a validation between field AGB data used in this study for validation and estimated AGB values from our AGB map and the previously mentioned maps. We also calculated the root mean square error (RMSE) and the relative root mean square error (%RMSE) obtained as the RMSE divided by mean AGB observed values for comparisons.

Relative contributions by sensor

In order to obtain the relative contributions by sensor, we partitioned the information provided by (a) Sentinel 2 reflectance and texture; (b), ALOS PALSAR backscatter and texture; and (c) shared variation, which is the variance in AGB that can be explained by either sensor. Total variation explained by the full model using information from both sensors can be summarized as: $Y = (a + b + c) + \varepsilon$, where ε is variation that cannot be accounted for by the predictor variables. The relative contribution of the two

sensors and the shared variation can then be partitioned by comparison against the variance explained by single sensor models using only Sentinel-2 (a + c) and only ALOS PALSAR (b + c).

Results

Calibrating LiDAR biomass estimates at the plot scale

Validation of the AGB TCH model had an R^2 of 0.40, RMSE of 46.14 Mg ha⁻¹ between AGB measured by Top of Canopy Height (TCH) and our field calculated AGB (in 400 m² plots) (Supplementary 2). Due to spatial uncertainty, heterogeneous canopies can result in large uncertainties in plot TCH, particularly where plots are located at or close to sharp transitions between short, secondary vegetation and old-growth forest. In this case, three field plots showed large residuals in the validation of the AGB TCH model (Figure 2). This derives from the presence of very large trees inside these plots which increase the field calculated biomass considerably, without a corresponding increase in height or TCH. Nevertheless, at 20 m resolution, estimations of AGB using LiDAR TCH show a good fit with the power law relationship (Figure 2).

[insert figure 2 around here]

Upscaling AGB using single sensor and combined models

Models upscaled at 100 m resolution provided greater explanatory power ($R^2 = 0.70$, RMSE = 27.9%) than either models upscaled at 50 m ($R^2 = 0.67$, RMSE = 29.8%) or 20 m resolution ($R^2 = 0.62$, RMSE = 31.8%), after aggregation post-upscaling to the same resolution grid (i.e. 100 m). This highlights that the reduction in noise by averaging spatially prior to upscaling led to a more robust upscaling model. Therefore, we only consider the 100 m resolution models from now onwards.

The upscaled models were clearly able of distinguishing forest from non-forest cover (Figure 3). However, sensitivity to AGB variations within the forest area was limited, especially for models reliant only on ALOS PALSAR, which had very little explanatory power regarding AGB variations above 100 Mg ha⁻¹ (Figure 3). The best upscaling model combined both Sentinel 2 and ALOS PALSAR ($R^2 = 0.70$; RMSE = 27.8%). In comparison, the Sentinel 2-only model had slightly lower predictive power ($R^2 = 0.66$; RMSE = 29.5%), while the model solely reliant on ALOS PALSAR performed worst ($R^2 = 0.50$; RMSE = 36.2%). Sentinel 2 explained a greater amount of variation of AGB (20%) solely compared to ALOS PALSAR (4 %). The majority of the explained variation (46% of the total variance) was shared between both sensors. Uncertainties in the combined model and in the Sentinel 2 model were highest in the mid-range of AGB < 100 Mg ha⁻¹. Conversely, ALOS PALSAR showed higher uncertainty above 100 Mg ha⁻¹, as its sensitivity saturated (Figure 3).

341 [insert figure 3 around here]

342 ***Relative contributions by sensor and variable importance***

343 Sentinel 2 explained a greater amount of variation of AGB (20 %) by itself, compared to ALOS PALSAR
344 (4 %), although a considerable amount of variation was shared between both sensors (46%). Sentinel 2
345 on its own was able to provide reasonable estimations of AGB in the study area, explaining 66 % in the
346 single sensor model, whereas ALOS PALSAR proved to be less effective explaining 50%, while the
347 combination of sensors provided the best fit (70 %).

348 The results of the permutation importance under spatial cross-validation highlighted the relative
349 importance of Sentinel 2 reflectance and texture measures over ALOS PALSAR in the random forest
350 model (Figure 4). Moreover, of the texture metrics, only the mean of AGB showed a high importance in
351 the model. Variables relating to heterogeneity (variance, contrast, dissimilarity) had marginal importance.
352 Variables relating to homogeneity (correlation, angular second moment 'ASM') were not important
353 indicated by the low values in permutation importance (Figure 4).

354 [insert figure 4 around here]

355 ***Validation of the AGB random forest model inside vs. outside the LiDAR survey area***

356 AGB showed a much higher fit ($R^2 = 0.49$) and a much lower error (relative RMSE = 24.6%)
357 inside the LiDAR survey extent compared to outside the LiDAR survey area ($R^2 = 0.17$ and relative
358 RMSE = 39.3%) (Figure 5). Importantly, the uncertainty estimates appear to be robust as estimates for
359 all plots inside the LiDAR survey area and 94% of plots outside of the LiDAR survey fell under the 95%
360 confidence intervals for AGB_{Field} and $AGB_{\text{satellite}}$. Outside the LiDAR survey extent there is one plot with
361 unusually large trees and exceptionally high AGB_{field} ($>300 \text{ Mg ha}^{-1}$), considerably higher than any of
362 the other plots in the inventory. Excluding this plot leads to a significant improvement in the fit outside
363 of the LiDAR area ($R^2 = 0.22$, relative RMSE = 36%).

364 The validation analysis to compare the AGB maps with previous studies revealed that the RMSE
365 and %RMSE obtained in this study were the lowest compared to the other maps (RMSE= 42.5 Mg ha^{-1}
366 and %RMSE = 35.0 in this study, RMSE= 51.2 Mg ha^{-1} and %RMSE = 42.0 for Santoro et al (2018),
367 RMSE= 57.5 Mg ha^{-1} and %RMSE = 47.0 for Cartus et al. (2016) and RMSE= 90.59 Mg ha^{-1}
368 and %RMSE = 90 in that of Rodriguez-Veiga et al (2014)) (Figure 6).

369 [insert figure 5 around here]

370 [insert figure 6 around here]

371 ***Spatial distribution of AGB and its uncertainty in the study area***

372 The spatial distribution of AGB (Figure 7) indicates that the higher biomass areas are located in
373 the north-east portion of the window, coinciding with the distribution of the state reserve Reserva Estatal
374 Biocultural del Puuc. Lower biomass areas are distributed around non-forest urban or agricultural areas,

where forests are likely to be more degraded. The largest uncertainties are associated to areas with intermediate ranges (50 – 75 Mg ha⁻¹) of AGB (Figure 8).

[insert figure 7 around here]

[insert figure 8 around here]

Land management appears to have a significant effect on forest AGB stocks (Tables 1 and 2). The highest AGB densities by management class were located in the protected reserves of Kaxil Kiuic and Del Puuc Biocultural reserve. Conversely, the small portion of the Bala'an Kaax reserve contained within our study area showed similar AGB to unprotected forest.

Moreover, we found greater areas of high biomass and smaller areas of low AGB in protected areas. Forest areas suitable for production and restoration showed large areas of both low and high AGB.

Comparing the distributions of the median AGB estimates from the Monte Carlo upscaling process there are marked differences between the protected and unprotected areas (Figure 9). Kaxil Kiuic and Reserva Estatal Biocultural del Puuc have higher AGB, with very low frequencies with AGB < 100 Mg ha⁻¹. These distributions contrast with the potential production and restoration areas, which both show much lower frequencies in the upper end of the AGB distributions, and a long tail of AGB < 100 Mg ha⁻¹. This is consistent with these areas of forest being subject to high levels of disturbance (Williams et al., 2013). The portion of the Reserva Bala'an Kaax within the study area has a similar distribution of AGB to forest production and restoration areas, suggesting this area of the reserve may have been subjected to similar degradation pressures.

[insert figure 9 around here]

Discussion

This study provides a spatially explicit estimation of AGB and its uncertainty in a semi-deciduous tropical dry forest of Yucatan using LiDAR data and a combination of information from passive and active sensors. As a first step, LiDAR data was used to estimate AGB using field plot information. The effectiveness of using LiDAR-derived AGB for upscaling plot-based estimations to continuous landscape level estimations has been demonstrated in various forests worldwide (Mascaro et al., 2011, Wulder et al., 2012, Asner et al., 2018). Random Forest models using information from a combination of Sentinel 2 and ALOS PALSAR were able to upscale AGB estimates based on a locally calibrated map of AGB based on LiDAR top-of-canopy height. Several studies have shown that tropical forest AGB can be estimated using ALOS PALSAR backscatter (Mitchard et al., 2013; Hernández-Stefanoni et al., 2020) and Sentinel 2 reflectance (Pandit et al., 2018), however, the combination of both sensors has been little explored (but see Vafaei et al., 2017). To assess the improvement on the precision of estimates by combining active and passive sensors we tested each sensor individually then produced a combined model using information from both sensors. Our results suggest that the estimation of AGB

in the semi deciduous tropical forest of Yucatan can be improved through a combination of ALOS PALSAR backscatter information and Sentinel 2 reflectance and texture variables, increasing the variance explained by the best single sensor model from 66% to 70% and reducing the RMSE from 29.5% to 27.8%. This improvement in AGB estimation is similar to the results found in Vafaei et al., (2017) in a subtropical forest in Iran also combining ALOS PALSAR backscatter and Sentinel 2. Furthermore, we tested the contribution of each sensor to explain AGB and found that Sentinel 2 on its own explained a greater amount of variation of AGB, compared to ALOS PALSAR, although the majority of the explained variation was shared between both sensors. One of the main caveats in the sensor combination approach is the difference in spatial resolution between the ALOS PALSAR backscatter (25 m) and Sentinel 2 (10 m). It is possible that this difference has an impact on the amount of variability that can be captured by each sensor at the plot level. Given its higher spatial resolution, Sentinel 2 could capture a greater range of variability of AGB within the plots than ALOS PALSAR. Coarser resolutions may not reflect the variability of structure as they contain averaged information from varying heights and may include reflectance from non-forest areas or canopy gaps within the same pixel (Lu, 2006).

We tested the use texture information as a way to quantify the variability of reflectance and backscatter within the plots and related this to LiDAR-estimated AGB. In this case, the upscaled models were principally reliant on the mean with limited additional contributions to the predictive power added by texture information. Other studies that have used texture information from ALOS PALSAR backscatter (Thapa et al., 2015; Hernández-Stefanoni et al., 2020) and Sentinel 2 reflectance (Pandit et al., 2019) have found large improvements in estimations of AGB by capturing the spatial variability and minimizing sensor saturation. To test the effect of spatial resolution in the upscaling process we compared models with different resolutions and found that an upscaling resolution of 100 m increased the fit of the best model by 8% and decreased the errors by 3.9%, compared to upscaling at 20 m resolution (Supplement 3). This suggests that the aggregation of information prior to upscaling might improve models and reduce the overall errors. However, as there is a trade-off between the information lost and the reduction of error when aggregating information (Camel, 2003), we chose not to aggregate further than 100 m, as this would reduce the spatial information gained from Sentinel's 10 m resolution. [The comparison with the work of Santoro et al., \(2018\), Rodriguez-Veiga et al., \(2016\), and Cartus et al., 2014 suggests that by performing a bias-corrected upscaling procedure we were able to reduce the error, thus, improving upon previous AGB mapping efforts in the dry forests of Yucatan. Such procedures can be used to produce AGB maps to inform regional and national strategies for reducing greenhouse gas emissions such as REDD+.](#)

Furthermore, by propagating errors through each step of the upscaling process and applying a spatially independent validation procedure, we were able to produce a robust estimation of errors (94%

of field AGB estimates for aggregated plot clusters overlap within the estimated 95% confidence interval outside of the LiDAR survey area). While the error propagation estimates appear to be robust, it is evident from the distribution of residuals (Figure 8) that there remains a trend in the residuals highlighting a tendency to underpredict the AGB of higher biomass field plots and overpredict the AGB at low biomass field plots. This suggests that the bootstrap bias correction was not sufficient to fully remove the bias in the random forest models, possibly a consequence of spatial correlations. Given that degradation and deforestation act to lower AGB, this outstanding source bias will likely lead to conservative estimates of the AGB differences between protected and unprotected forests, and therefore conservative estimates of restoration potential. This result suggests an improvement of previous efforts to estimate AGB in semi-deciduous dry forests of the Yucatan Peninsula using active sensors such as ALOS PALSAR (Hernandez-Stefanoni et al., 2020) and national scale efforts (Cartus et al., 2014; Rodriguez-Veiga et al., 2016). Previous attempts to map AGB across Mexico have found a wide range of AGB values in the Yucatan Peninsula reaching 150 Mg ha^{-1} (Hernandez-Stefanoni et al., 2020; Rodriguez-Veiga et al., 2016; Cartus et al., 2014) and greatest uncertainties in the lower end of the AGB distribution (Rodriguez-Veiga et al., 2016). The spatial distribution of uncertainty showed that the largest uncertainties were associated to the middle range of AGB distribution (Figures 7 and 8) and it is derived from the underrepresentation of areas with this range of AGB values ranging between 25 and 75 Mg ha^{-1} in the calibration data (Figure 8). However, in accordance with Hernandez-Stefanoni et al. (2020), estimates were also found to be constrained by the range of AGB variation captured by LiDAR data available across the calibration landscape. In particular, the predictability of the upper bounds of the biomass ranges was severely affected by the lack of LiDAR coverage in the very high biomass forest ($> 200 \text{ Mg ha}^{-1}$). Therefore, areas with high biomass, located in the north-east of the window area, in the protected area of “del Puuc Biocultural reserve”, are underrepresented in the LiDAR survey with only a portion of the area, corresponding to the location of “Kaxil Kiuic Biocultural reserve”, represented by both field and LiDAR data (Figure 1). To estimate AGB in tropical forests where forest protection areas and areas where disturbances such as slash-and-burn agriculture shape the spatial variability of forest AGB, the accuracy of estimates will depend on the distribution of LiDAR and field data available across all the possible ranges of AGB. As it has been previously cautioned, the range of variability in AGB captured by both the LiDAR data and the forest inventory constrained the next stages of the analysis (Hernandez-Stefanoni et al., 2020), limiting the predictability in the lower and upper ranges of our estimated AGB. In order to reduce the uncertainty in AGB mapping, future upscaling efforts could aim for a more thoroughly distributed airborne sampling campaign that better characterizes the full range of AGB values in the landscape. Moreover, uncertainty in the upper and lower ranges of AGB was reduced when combining information from both sensors, suggesting that the combination of these sensors is an effective way to improve AGB mapping.

Within the study region, larger areas with high biomass were found in the protected areas of “del Puuc Biocultural reserve” and “Kaxil Kiuic Biocultural Reserve”, which were created for the conservation of forests and their environmental services (Table 2). In particular, the Kaxil Kiuic protected area shows a more symmetric distribution with the highest mean AGB (Figure 9; Table 2) indicating a large proportion of this area may be approaching a steady state condition (Williams et al., 2013). However, several other low impact activities such as extraction of woody species for fuel, and agricultural and pastures for cattle ranching take place inside del Puuc Biocultural reserve. This is reflected in the tail of low AGB values in this area, although significantly less prevalent than outside the forest reserves. The study region has a long history of land use, mainly for slash-and-burn agriculture, also practiced presently in the area (Ellis et al., 2017). The effect of the more recent repeated disturbance is reflected in the AGB distributions of the production forest, which have skewed AGB distributions with a long tail of low AGB (Figure 9). Critically, regions allocated for restoration have large areas with low AGB (Table 2) and similar AGB distributions to existing production forest (Figure 9). Therefore, while there is potential for substantial gains in aboveground carbon stocks through restoration, whether these gains are realized is likely to be dependent on these restored forests being protected and allowed to develop into high biomass old-growth systems (Lewis et al., 2019; Chazdon et al., 2016).

Conclusions

LiDAR data proved a useful upscaling tool for calibrating and validating satellite models of AGB, however, the reliability of these estimates is constrained by the degree to which the sampled areas represent the range of AGB values found in the whole landscape, to avoid potential biases when upscaling outside the training area. The sensitivity to within-forest variation in AGB was more limited particularly in the upper end of the AGB range, thus limiting our ability to predict AGB in high biomass forest areas.

We found that the information from active (ALOS PALSAR backscatter) and passive (Sentinel 2 reflectance) sensors can be combined to improve spatially explicit estimations of AGB in semi-deciduous tropical forest. However, Sentinel 2 explained a higher proportion of the variance in the combined model and performs better than ALOS PALSAR when considered separately. We believe the methods described in this study can be used to improve estimations of AGB and its uncertainty in tropical forests. Using a combination of LiDAR and satellite data, we upscaled LiDAR estimates of AGB across a landscape of semi-deciduous forest in the Yucatan peninsula to gain insights on the distribution of AGB in different categories of forest protection. The frequency distributions of AGB obtained from our maps highlighted the benefits of protected areas for maintaining forest carbon stocks. On the other hand, a significantly greater portion of the areas designated for restoration currently have low AGB, comparable to the distribution of AGB in existing production forest. The similarity in the distributions of these categories suggests areas of restoration should be effectively protected for carbon sequestration, biodiversity

514 conservation and for other important ecosystem services, which can take several decades to reach old-
515 growth forest values.

516 We believe the information obtained can provide insights on the state of the AGB stock in
517 different management or protection categories in the region and thus aid conservation, restoration, and
518 sustainable management policies in the semi-deciduous forests of the Yucatan Peninsula.

519

520 **Acknowledgements**

521 The authors would like to thank Fernando Tun Dzul and Reserva Biocultural Kaxil Kiuic, for their
522 support during fieldwork. The first author would like to thank Consejo Nacional de Ciencia y
523 Tecnologia (CONACyT) for the Ph D. scholarship awarded.

524

525 **Availability of data and materials**

526 The ALOS PALSAR data used in this study was downloaded from
527 (https://www.eorc.jaxa.jp/ALOS/en/top/obs_top.htm). The LiDAR data can be accessed at
528 (<https://gliht.gsfc.nasa.gov/>). Data from national forest inventory in Mexico can be obtained by request
529 to CONAFOR (Comisión Nacional Forestal, <https://www.gob.mx/conafor>).

530 **References**

- 531 Asner, G.P., Clark, J.K., Mascaro, J., Galindo Garcia, G.A., Chadwick, K.D., Navarette Encinales, D.A.,
532 Paez-Acosta, G., Montenegro, C., Kennedy-Bowdoin, T., Duque, A. (2012). High-resolution
533 mapping of forest carbon stocks in the Colombian Amazon. *Biogeosciences* 2012, 9, 2683–2696.
- 534 Asner, G.P., Mascaro, J., Muller-Landau, H.C., Vieilledent, G., Vaudry, R., Rasamoelina, M., Hall, J.S.,
535 van Breugel, M. (2012). A universal airborne LiDAR approach for tropical forest carbon
536 mapping. *Oecologia* 168, 1147–1160. <https://doi.org/10.1007/s00442-011-2165-z>
- 537 Asner, G.P., Mascaro, J. (2014). Mapping tropical forest carbon: Calibrating plot estimates to a simple
538 LiDAR metric. *Remote Sensing of Environment* 140, 614–624.
539 <https://doi.org/10.1016/j.rse.2013.09.023>
- 540 Asner, G. P., Brodrick, P. G., Philipson, C., Vaughn, N. R., Martin, R. E., Knapp, D. E., ... & Stark, D. J.
541 (2018). Mapped aboveground carbon stocks to advance forest conservation and recovery in
542 Malaysian Borneo. *Biological Conservation*, 217, 289-310.
- 543 Babcock, C., Finley, A.O., Cook, B.D., Weiskittel, A., Woodall, C.W. (2016). Modeling forest biomass
544 and growth: Coupling long-term inventory and LiDAR data. *Remote Sensing of Environment* 182,
545 1–12. <https://doi.org/10.1016/j.rse.2016.04.014>
- 546 Baskerville, G.L. (1972). Use of Logarithmic Regression in the Estimation of Plant Biomass. *Canadian*
547 *Journal of Forest Research*. 2, 49–53. <https://doi.org/10.1139/x72-009>
- 548 Bergstra, J., Yamins, D., Cox, D.D. (2013). Hyperopt: A python library for optimizing the
549 hyperparameters of machine learning algorithms, in: Proceedings of the 12th Python in Science
550 Conference. Citeseer, pp. 13–20.
- 551 Bergstra, J.S., Bardenet, R., Bengio, Y., Kégl, B. (2011). Algorithms for Hyper-Parameter Optimization,
552 in: Shawe-Taylor, J., Zemel, R.S., Bartlett, P.L., Pereira, F., Weinberger, K.Q. (Eds.), Advances
553 in Neural Information Processing Systems 24. Curran Associates, Inc., pp. 2546–2554.
- 554 Bispo, P.C., Rodríguez-Veiga, P., Zimbres, B., do Couto de Miranda, S., Henrique Giusti Cezare, C.,
555 Fleming, S., ... & Balzter, H. (2020). Woody aboveground biomass mapping of the brazilian
556 savanna with a multi-sensor and machine learning approach. *Remote Sensing*, 12(17), 2685.
- 557 Breiman, L. (2001). Random Forests. *Machine Learning* 45, 5–32.
558 <https://doi.org/10.1023/A:1010933404324>
- 559 Bonan, G. B. (2008). Forests and climate change: forcings, feedbacks, and the climate benefits of forests,
560 *Science*, 320, 1444–1449, 2008.

561 Caamal-Sosa, J. P., Dupuy, J. M., Torres, J.L.A., Luis, J., Stefafoni, H., Ruíz, A.H.H., Chim, M.T.,
562 Wayson, C., Álvarez, M.O., Merlín, D.L. and Montero, V.M., (2016) Estudio de caso del sitio de
563 monitoreo intensivo del carbono, Kaxil Kiuic, Yucatán *Programa Mexicano del Carbono*.

564 Carnevali, G., Ramírez, I.M., González–Iturbe, J.A. Flora y vegetación de la Península de Yucatán. In:
565 Colunga–García–Marín, P. and Larqué–Saavedra, A. Eds. (2003) Naturaleza y Sociedad en el
566 Área Maya, Pasado, Presente y Futuro, *Academia Mexicana de Ciencias y Centro de*
567 *Investigación Científica de Yucatán*, México, D. F. (2003); pp. 53-68.

568 Cartus O., Kelldorfer J., Walker W., Franco C., Bishop J., Santos L., Fuentes J. (2014) A national,
569 detailed map of forest aboveground carbon stocks in Mexico. *Remote Sensing*. 2014; 6(6):5559-
570 5588.

571 Castillo-Santiago, M. A., Ricker, M., & de Jong, B. H. (2010). Estimation of tropical forest structure
572 from SPOT-5 satellite images. *International Journal of Remote Sensing*, 31(10), 2767-2782.

573 Chave, J., Condit, R., Aguilar, S., Hernandez, A., Lao, S., Perez, R. (2004). Error propagation for tropical
574 forest biomass estimates. *Philosophical Transactions of the Royal Society of London B*, 359, 409–
575 420.

576 Chave, J., Andalo, C., Brown, S., Cairns, M.A., Chambers, J.Q., Eamus, D., Fölster, H., Fromard, F.,
577 Higuchi, N., Kira, T. and Lescure, J.P. (2005). Tree allometry and improved estimation of carbon
578 stocks and balance in tropical forests. *Oecologia* 145(1), 87-99.

579 Chazdon, R. L., E. N. Broadbent, D. M. A. Rozendaal, F. Bongers, A. M. A. Zambrano, T. M. Aide, P.
580 Balvanera, J. M. Becknell, V. Boukili, P. H. S. Brancalion, D. Craven, J. S. Almeida-Cortez, G.
581 A. L. Cabral, B. de Jong, J. S. Denslow, D. H. Dent, S. J. DeWalt, J. M. Dupuy, S. M. Durán, M.
582 M. Espírito-Santo, M. C. Fandino, R. G. César, J. S. Hall, J. Luis Hernández-Stefanoni, C. C.
583 Jakovac, A. B. Junqueira, D. Kennard, S. G. Letcher, M. Lohbeck, M. Martínez-Ramos, P.
584 Massoca, J. A. Meave, R. Mesquita, F. Mora, R. Muñoz, R. Muscarella, Y. R. F. Nunes, S. Ochoa-
585 Gaona, E. Orihuela-Belmonte, M. Peña-Claros, E. A. Pérez-García, D. Piotto, J. S. Powers, J.
586 Rodríguez-Velazquez, I. E. Romero-Pérez, J. Ruíz, J. G. Saldarriaga, A. Sanchez-Azofeifa, N. B.
587 Schwartz, M. K. Steininger, N. G. Swenson, M. Uriarte, M. van Breugel, H. van der Wal, M. D.
588 M. Veloso, H. Vester, I. C. G. Vieira, T. Vizcarra Bentos, G. Bruce Williamson & L. Poorter.
589 (2016) Carbon sequestration potential of second-growth forest regeneration in the Latin American
590 tropics. *Science Advances* 2: e1501639.

591 Clark D. B., Kellner, J. R. (2012). Tropical forest biomass estimation and the fallacy of misplaced
592 concreteness. *Journal of Vegetation Science*. 23(2012): 1191 - 1196 doi: 10.1111/j.1654-
593 1103.2012.01471.x

594 Comisión Nacional de Áreas Protegidas (CONANP) (2017). Áreas Naturales Protegidas Federales de la
595 República Mexicana. Shapefile. México CONANP.

596 Comisión Nacional Forestal (CONAFOR) (2013). Inventario Nacional Forestal y de Suelos,
 597 Procedimientos de muestreo, CONAFOR, México.2013.

598 Comisión Nacional Forestal (CONAFOR) (2015). Áreas de importancia para la producción, restauración
 599 y conservación Shapefile. CONAFOR México 2015.

600 Cook, B. D., Nelson, R. F., Middleton, E. M., Morton, D. C., McCorkel, J. T., Masek, J. G., Ranson K.
 601 J., Montesano, P. M. (2013). NASA Goddard's LiDAR, hyperspectral and thermal (G-LiHT)
 602 airborne imager. *Remote Sensing*, 5(8), 4045-4066.

603 Cook-Patton, S. C., Leavitt, S. M., Gibbs, D., Harris, N. L., Lister, K., Anderson-Teixeira, K. J., Briggs,
 604 R. D., Chazdon, R. L., Crowther, T. W., Ellis, P. W. and Griscom, H. P. (2020). Mapping carbon
 605 accumulation potential from global natural forest regrowth. *Nature*, 585(7826), 545-550.

606 Dubayah, R., Blair, J. B., Goetz, S., Fatoyinbo, L., Hansen, M., Healey, S., ... Silva, C. (2020). The Global
 607 Ecosystem Dynamics Investigation: High-resolution laser ranging of the Earth's forests and
 608 topography. *Science of Remote Sensing*, 1, 100002. <https://doi.org/10.1016/j.srs.2020.100002>

609 Dupuy J.M., Durán García, R., García Contreras, G., Arellano Morin, J., Acosta Lugo, E., Méndez
 610 González, M.E., Andrade Hernandez, M. (2015) Chapter 8: Conservation and Use in Islebe, G.
 611 A., Schmook, B., Calmé, S., & León-Cortés, J. L. (2015). Introduction: biodiversity and
 612 conservation of the Yucatán Peninsula, Mexico. In *Biodiversity and Conservation of the Yucatán*
 613 *Peninsula* (pp. 1-5). Springer, Cham.

614 Dupuy, J. M., Hernández-Stefanoni, J. L., Hernández-Juárez, R. A., Tetetla-Rangel, E., López-Martínez,
 615 J. O., Leyequién-Abarca, E., Fernando J. Tun-Dzul, and Filogonio May-Pat. & May-Pat, F.
 616 (2012). Patterns and correlates of tropical dry forest structure and composition in a highly
 617 replicated chronosequence in Yucatan, Mexico. *Biotropica*, 44(2), 151-162.

618 Ellis, E. A., Montero, J. A. R., Gómez, I. U. H., Porter-Bolland, L., & Ellis, P. W. (2017). Private property
 619 and Mennonites are major drivers of forest cover loss in central Yucatan Peninsula, Mexico. *Land*
 620 *Use Policy*, 69, 474-484.

621 Gallardo-Cruz J. A., Meave J. A., González E. J., Lebrija-Trejos E. E., Romero-Romero M. A., Pérez-
 622 García E. A., et al. (2012) Predicting Tropical Dry Forest Successional Attributes from Space: Is
 623 the Key Hidden in Image Texture? *PLoS ONE* 7(2): e30506.
 624 <https://doi.org/10.1371/journal.pone.0030506>.

625 Gonzalez, P., Asner, G.P., Battles, J.J., Lefsky, M.A., Waring, K.M., Palace, M. (2010). Forest carbon
 626 densities and uncertainties from Lidar, QuickBird, and field measurements in California. *Remote*
 627 *Sensing of Environment* 114, 1561–1575. <https://doi.org/10.1016/j.rse.2010.02.011>

628 Haralick, R. M. (1979). Statistical and structural approaches to texture. *Proceedings of the IEEE*, 67(5),
 629 786-804.

630 Hernández-Stefanoni J. L., Reyes-Palomeque G, Castillo-Santiago M, George-Chacón S, Huechacona-
631 Ruiz A, Tun-Dzul F, Rondon-Rivera D, Dupuy J. M. (2018). Effects of sample plot size and gps
632 location errors on AGB estimates from LiDAR in tropical dry forests. *Remote Sensing*
633 10(10):1586.

634 Hernández-Stefanoni, J. L., Castillo-Santiago, M. Á., Mas, J. F., Wheeler, C. E., Andres-Mauricio, J.,
635 Tun-Dzul, F., ... & Dupuy, J. M. (2020). Improving AGB maps of tropical dry forests by
636 integrating LiDAR, ALOS PALSAR, climate and field data. *Carbon Balance and Management*,
637 15(1), 1-17.

638 Hooker, G. and Mentch, L. (2018). Bootstrap bias corrections for ensemble methods *Statistics and*
639 *Computing*. 28 77–86.

640 Houghton, R. A. (2005). Tropical deforestation as a source of greenhouse gas emissions. *Tropical*
641 *deforestation and climate change*, 13.

642 Houghton, R. A. (2012). Carbon emissions and the drivers of deforestation and forest degradation in the
643 tropics. *Current Opinion in Environmental Sustainability*, 4(6), 597-603.

644 Houghton, R. A. (2013). The emissions of carbon from deforestation and degradation in the tropics: past
645 trends and future potential. *Carbon Management*, 4(5), 539-546.

646 Joshi, N., Mitchard, E. T., Brolly, M., Schumacher, J., Fernández-Landa, A., Johannsen, V. K., ... &
647 Fensholt, R. (2017). Understanding ‘saturation’ of radar signals over forests. *Scientific reports*,
648 7(1), 1-11.

649 Jucker, T., Asner, G. P., Dalponte, M., Brodrick, P., Philipson, C. D., Vaughn, N., Brelsford, C., Burslem,
650 D. F. R. P., Deere, N. J., Ewers, R. M., Kvasnica, J., Lewis, S. L., Malhi, Y., Milne, S., Nilus, R.,
651 Pfeifer, M., Phillips, O., Qie, L., Renneboog, N., Reynolds, G., Riutta, T., Struebig, M. J., Svátek,
652 M., Teh, Y. A., Turner, E. C., Coomes, D. A. (2017). A regional model for estimating the
653 aboveground carbon density of Borneo’s tropical forests from airborne laser scanning.
654 *arXiv:1705.09242 [q-bio]*.

655 Lee J. S. (1980) Digital image enhancement and noise filtering by use of local statistics. *IEEE Trans.*
656 *Pattern Anal. Mach. Intel.* 1980; 2(2):165-186.

657 Lewis, S. L., Wheeler, C. E.; Mitchard, E. T. A.; Koch, A. (2019). Restoring natural forests is the best
658 way to remove atmospheric carbon. *Nature* 568, 25–28, doi:[10.1038/d41586-019-01026-8](https://doi.org/10.1038/d41586-019-01026-8).

659 Lu, D. (2006). The potential and challenge of remote sensing-based biomass estimation. *International*
660 *Journal of Remote Sensing*, 27(7), 1297-1328.

661 Luther, J. E., Fournier, R. A., van Lier, O. R., Bujold, M. (2019). Extending ALS-Based Mapping of
662 Forest Attributes with Medium Resolution Satellite and Environmental Data. *Remote Sensing* 11,
663 1092. <https://doi.org/10.3390/rs11091092>

664 Mascaro, J., Detto, M., Asner, G., Muller-Landau, H. C. (2011). Evaluating uncertainty in mapping forest
665 carbon with airborne LiDAR. *Remote Sensing of Environment* 115 (2011) 3770-3774.

666 Mascaro, J., Asner, G.P., Knapp, D.E., Kennedy-Bowdoin, T., Martin, R.E., Anderson, C., Higgins, M.,
667 Chadwick, K.D. (2014). A Tale of Two “Forests”: Random Forest Machine Learning Aids
668 Tropical Forest Carbon Mapping. *PLoS ONE* 9, e85993.
669 <https://doi.org/10.1371/journal.pone.0085993>

670 McGaughey R. J. FUSION/LDV: Software for LIDAR data analysis and visualization. United States
671 Department of Agriculture, Forest Service, Pacific Northwest Research Station (2012); p 154.
672 Available online: http://forsys.cfr.washington.edu/fusion/fusion_overview.html (accessed on
673 December 11, 2019).

674 McNicol, I. M., Ryan, C. M., & Mitchard, E. T. A. (2018). Carbon losses from deforestation and
675 widespread degradation offset by extensive growth in African woodlands. *Nature*
676 *Communications*, 9(1). <https://doi.org/10.1038/s41467-018-05386-z>

677 Mitchard, E. T. A. (2018). The tropical forest carbon cycle and climate change. *Nature*, 559(7715), 527-
678 534. doi:10.1038/s41586-018-0300-2

679 Mermoz, S., Réjou-Méchain, M., Villard, L., Le Toan, T., Rossi, V., and Gourlet-Fleury, S. (2015).
680 Decrease of L-band SAR backscatter with biomass of dense forests. *Remote Sensing of*
681 *Environment*, 159, 307-317.

682 Mitchard, E. T. A., Saasan, S., Baccini A., Asner G. P., Goetz S. J., Harris, N. L., Brown, S. (2013)
683 Uncertainty in the spatial distribution of aboveground forest biomass: a comparison of pan
684 tropical methods. *Carbon Balance and Management* 2013, 8:10
685 <http://www.cbmjournal.com/content/8/1/10>

686 Mitchard, E. T. A., Saatchi S. S., Woodhouse I. H., Feldpausch T. R., Lewis S. L., Sonke B., Rowland
687 C., and P. Meir (2009), Measuring biomass changes due to woody encroachment and
688 deforestation/degradation in a forest-savanna boundary region of central Africa using multi-
689 temporal L-band radar backscatter, *Remote Sensing of Environment*. 115 11(2009): 2861 – 2873.

690 Murphy, P. G., Lugo, A. E. (1986). Ecology of tropical dry forest. *Annual review of ecology and*
691 *systematics*, 17(1), 67-88.

692 Pandit, S., Tsuyuki, S., Dube, T. (2019). Exploring the inclusion of Sentinel-2 MSI texture metrics in
693 above-ground biomass estimation in the community forest of Nepal. *Geocarto International*, 1-
694 18.

695 Pan, Y., R. A. Birdsey, J. Fang, R. Houghton, P. E. Kauppi, W.A. Kurz, O.A. Phillips, A. Shvidenko, S.L.
696 Lewis, J.G. Canadell, P. Ciais, R.B. Jackson, S.W. Pacala, A.D. McGuire, S. Piao, A. Rautiainen,
697 S. Sitch y D. Hayes (2011). A large and persistent carbon sink in the world’s forests. *Science*, 333,
698 988–993.

699 Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer,
700 P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M.,
701 Duchesnay, É., (2011). Scikit-learn: Machine Learning in Python. *Journal of Machine Learning*
702 *Research* 12, 2825–2830.

703 Ploton, P., Mortier, F., Réjou-Méchain, M., Barbier, N., Picard, N., Rossi, V., ... & Pélissier, R. (2020).
704 Spatial validation reveals poor predictive performance of large-scale ecological mapping models.
705 *Nature communications*, 11(1), 1-11.

706 Qi, W.; Dubayah, R.O. (2016) Combining Tandem-X InSAR and simulated GEDI lidar observations for
707 forest structure mapping. *Remote Sensing of Environment* , 187, 253–266,
708 doi:10.1016/j.rse.2016.10.018.

709 R Development Core Team (2018). A Language and Environment for Statistical Computing. R
710 Foundation for Statistical Computing, Vienna, Austria, ISBN 3-900051-07-0.
711 <http://www.Rproject.org>.

712 Ramírez Ramírez, G., Dupuy Rada, J. M., Ramírez y Avilés, L., & Solorio Sánchez, F. J. (2017).
713 Evaluación de ecuaciones alométricas de biomasa epigea en una selva mediana subcaducifolia de
714 Yucatán. *Madera y bosques*, 23(2), 163-179.

715 Réjou-Méchain, M., Barbier, N., Couteron, P., Ploton, P., Grégoire, V., Herold, M., Mermoz, Stéphane,
716 Saatchi, A., Chave, J., de Boissieu, F., Féret, J.B., Takoudjou, S.M., Pélissier, R. (2019).
717 Upscaling Forest Biomass from Field to Satellite Measurements: Sources of Errors and Ways to
718 Reduce Them. *Surveys in Geophysics* <https://doi.org/10.1007/s10712-019-09532-0>

719 Roberts D. R., Bahn V., Ciuti S., Boyce M. S., Elith J., Guillerá-Arroita G., Hauenstein S., Lahoz-Monfort
720 J. J., Schröder B., Thuiller W., Warton D. I., Wintle B. A., Hartig F. and Dormann C. F. (2017).
721 Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure
722 *Ecography* 40 913–29

723 Rodríguez-Veiga, P., Saatchi, S., Tansey, K., Balzter, H., (2016). Magnitude, spatial distribution and
724 uncertainty of forest biomass stocks in Mexico. *Remote Sensing of Environment* 183, 265–281.
725 <https://doi.org/10.1016/j.rse.2016.06.004>

726 Rodríguez-Veiga, P., Quegan, S., Carreiras, J., Persson, H. J., Fransson, J. E., Hosillo, A., ... & Berninger,
727 A. (2019). Forest biomass retrieval approaches from earth observation in different biomes.
728 *International Journal of Applied Earth Observation and Geoinformation*, 77, 53-68.

729 Roussel, J.-R., Caspersen, J., Béland, M., Thomas, S., Achim, A., (2017). Removing bias from LiDAR-
730 based estimates of canopy height: Accounting for the effects of pulse density and footprint size.
731 *Remote Sensing of Environment* 198, 1–16. <https://doi.org/10.1016/j.rse.2017.05.032>

732 Rzedowski, J. (2006). Vegetación de México. 1ra. Edición digital, *Comisión Nacional para el*
733 *Conocimiento y Uso de la Biodiversidad*. México.

734 Sanaphre-Villanueva, L., Dupuy, J. M., Andrade, J. L., Reyes-García, C., Paz, H., Jackson, P. C. (2016).
 735 Functional diversity of small and large trees along secondary succession in a tropical dry forest.
 736 *Forests*, 7(8), 163.

737 Santoro, M., Cartus, O., Mermoz, S., Bouvet, A., Le Toan, T., Carvalhais, N., Rozendaal, D., Herold, M.,
 738 Avitabile, V., Quegan, S., Carreiras, J., Rauste, Y., Balzter, H., Schmullius, C., Seifert, F.M.,
 739 (2018), GlobBiomass global above-ground biomass and growing stock volume datasets, available
 740 on-line at <http://globbiomass.org/products/global-mapping>

741 Shimada M., Ohtaki T. (2010). Generating large-scale high-quality sar mosaic datasets: application to
 742 PALSAR data for global monitoring. *IEEE Journal of Selected Topics in Applied Earth*
 743 *Observation and Remote Sensing*. 2010; 3 (4): 637–656.

744 Strobl, C., Boulesteix, A.-L., Zeileis, A., Hothorn, T. (2007). Bias in random forest variable importance
 745 measures: Illustrations, sources and a solution. *BMC Bioinformatics* 8, 25.
 746 <https://doi.org/10.1186/1471-2105-8-25>

747 Swinfield, T., Both, S., Riutta, T., Bongalov, B., Elias, D., Majalap-Lee, N., Ostle, N., Svátek, M.,
 748 Kvasnica, J., Milodowski, D., Jucker, T., Ewers, R.M., Zhang, Y., Johnson, D., Teh, Y.A.,
 749 Burslem, D.F.R.P., Malhi, Y., Coomes, D. (2020). Imaging spectroscopy reveals the effects of
 750 topography and logging on the leaf chemistry of tropical forest canopy trees. *Global Change*
 751 *Biology* 26, 989–1002. <https://doi.org/10.1111/gcb.14903>

752 Thapa R.B., Watanabe M., Motohka T., Shimada M. (2015). Potential of high-resolution ALOS PALSAR
 753 mosaic texture for aboveground forest carbon tracking in tropical region. *Remote Sensing of*
 754 *Environment*. 2015;160:122–33.

755 Urbazaev M, Thiel C, Cremer F, Dubayah R, Migliavacca M, Reichstein M and Schmullius C. (2018)
 756 Estimation of forest AGB and uncertainties by integration of field measurements, airborne
 757 LiDAR, and SAR and optical satellite data in Mexico *Carbon Balance and Management* 13 5

758 Uriarte M., Muscarella, R., Zimmerman, J.K. (2018). Environmental heterogeneity and biotic
 759 interactions mediate climate impacts on tropical forest regeneration. *Global Change Biology* 24:2

760 Vafaei S., Soosani J., Adeli K., Fadaei H., Naghavi H., Pham T.D., Tien Bui D. (2018). Improving
 761 Accuracy Estimation of Forest AGB Based on Incorporation of ALOS PALSAR-and Sentinel-2A
 762 Imagery and Machine Learning: A Case Study of the Hyrcanian Forest Area (Iran). *Remote*
 763 *Sensing*. 10, 172.

764 van der Walt, S., Schönberger, J. L., Nunez-Iglesias, J., Boulogne, F., Warner, J. D., Yager, N., Guillard,
 765 E., Yu, T., & contributors. (2014) scikit-image: Image processing in Python. *PeerJ*, 2, e453.
 766 <https://doi.org/10.7717/peerj.453>

767 Williams M., Hill, T. C., & Ryan, C. M. (2013). Using biomass distributions to determine probability and
 768 intensity of tropical forest disturbance. *Plant Ecology & Diversity*, 6(1), 87-99.

769 Wood E. M., Pidgeon, A. M., Radeloff, V. C., & Keuler, N. S. (2012). Image texture as a remotely sensed
770 measure of vegetation structure. *Remote Sensing of Environment*, 121, 516-526.

771 Woodhouse I. H. (2017) Introduction to Microwave Remote Sensing (CRC Press) Online:
772 <https://www.taylorfrancis.com/books/9781315272573>.

773 Wulder M. A., White J. C., Nelson R. F., Næsset E., Ørka H. O., Coops N. C., Hilker T., Bater C. W. and
774 Gobakken T. (2012). Lidar sampling for large-area forest characterization: A review *Remote*
775 *Sensing of Environment* **121** 196–209

776 Xu, L., Saatchi, S.S., Yang, Y., Yu, Y., White, L. (2016). Performance of non-parametric algorithms for
777 spatial mapping of tropical forest structure. *Carbon Balance and Management*, 11, 18.
778 <https://doi.org/10.1186/s13021-016-0062-9>

779 Yanai, R. D., Wayson, C., Lee, D., Espejo, A. B., Campbell, J. L., Green, M. B., ... & Gamarra, J. G.
780 (2020). Improving uncertainty in forest carbon accounting for REDD+ mitigation efforts.
781 *Environmental Research Letters*, 15(12), 124002.

782

783 Table 1. Mean AGB and confidence intervals (CI) [Mg ha⁻¹] for protected areas and areas without
784 protection in the Kiuic area.

Management condition	Site	AGB	CI
Protected	Kaxil Kiuic	129.14	125 - 134
	Reserva Estatal Biocultural del Puuc	126.13	122 - 132
	Bala'an Kaax	100.64	97 - 104
Without protection	Restoration	106.63	103 - 110
	Production	99.23	96 - 103

785

786

787   Table 2. Summary of the area occupied by different AGB classes for different management
788   conditions, with 95 confidence intervals provided in parentheses. Area in size classes is expressed as
789   percentage relative to total area (last column).

AGB class (Mg ha ⁻¹)	Area by AGB Class (%)							Total Area (km ²)
	0-25	25-50	50-75	75-100	100-125	125-150	>150	
Kaxil Kiuiç (protected)	0.2 (0.0/0.3)	1.0 (0.5/1.6)	4.5 (3.2/5.8)	11.8 (9.5/14.0)	26.5 (23.1/29.7)	30.3 (27.8/32.6)	25.7 (20.8/32.3)	18.5
Reserva Estatat Biocultural Del Puuc (protected)	2.3 (2.0/2.6)	3.1 (2.6/3.5)	5.3 (4.5/6.0)	10.6 (9.2/11.9)	22.1 (19.1/24.7)	27.4 (25.1/28.9)	28.7 (23.8/35.7)	697.0
Bala'an kaax (protected)	7.2 (5.9/8.3)	9.4 (8.3/10.4)	10.5 (9.3/11.7)	15.2 (12.7/16.7)	25.0 (23.3/26.9)	20.0 (18.5/21.8)	12.1 (9.9/14.9)	53.3
Production forest	10.2 (9.6/10.7)	6.6 (6.1/7.1)	8.0 (7.1/8.8)	13.5 (12.1/14.5)	24.2 (22.5/25.5)	21.3 (20.1/22.9)	13.2 (11.0/16.5)	2154.2
Restoration forest	7.6 (6.8/8.2)	6.9 (6.2/7.5)	8.1 (7.2/8.8)	12.8 (11.4/14.0)	23.1 (21.0/24.9)	22.4 (21.3/23.7)	17.6 (14.7/21.7)	216.8

790

791

792

793 **Figure captions**

794

795 Figure 1. Location of study area in Mexico (upper-right box) and location of protected areas within study
796 area, LiDAR and field data used in this study. National Protected Area (Bala'an K'aax), State Protected
797 Area (del Puuc Biocultural Reserve) (CONANP 2017) and private protected area (Kaxil Kiuic
798 Biocultural Reserve). Areas without protection are subdivided into areas suitable for production and those
799 suitable for restoration (CONAFOR 2015).

800 Figure 2. Comparison of field inventory AGB and LiDAR TCH for the 0.04 ha inventory plots, shown
801 with a series of example plots (numbers in blue) highlighting variations in TCH across the range of AGB
802 spanned by the plot network. In the first panel, the line is the fitted relationship between field AGB and
803 plot TCH. Error bars (horizontal and vertical lines) represent the uncertainty in plot field AGB (points),
804 and the uncertainty (both 50% CI and 95% CI shown) in plot TCH, modelled by randomly sampling the
805 TCH with positional uncertainty.

806 Figure 3. Regression lines, R^2 , RMSE and relative %RMSE based on a five-fold buffered-blocked cross-
807 validation between LiDAR estimated AGB (AGB_{lidar}) and upscaled AGB ($AGB_{satellite}$) for models using
808 both sensors a), Sentinel 2 reflectance and textures b), and ALOS PALSAR and textures c). The dashed
809 line represents the 1:1 relationship, the solid and dotted lines represent the median estimate and 95%
810 confidence interval for a 20 Mg ha⁻¹ moving window across the predicted AGB range ($AGB_{satellite}$).

811 Figure 4. Permutation importance based on permutation of different aggregated input variables
812 corresponding to specific sensors (green) and texture measures (grey).

813 Figure 5. Regression lines of the validation of the upscaled AGB against field inventory data inside and
814 outside the LiDAR survey area. Points represent clusters of four 400 m² plots (1600 m²), uncertainty is
815 shown as vertical and horizontal lines.

816 Figure 6. Comparison of observed AGB (obtained with field data used for validation)) and predicted
817 AGB values (mapped AGB of different studies). The predicted values were obtained from Santoro et al.
818 (2018), Rodriguez-Veiga et al. (2016), and Cartus et al. (2014). Solid lines indicate the regression
819 between observed and predicted AGB, while dashed gray line shows a 1:1 relationship.

820

821 Figure 7. Spatial distribution of AGB (left pane) and its uncertainty (right pane) in the study area. Grid
822 lines are spaced 10 km. Letters correspond to the location of officially designated protected areas within
823 the study landscape: A) Kaxil Kiuic Biocultural Reserve, B) del Puuc Reserva Biocultural reserve C)
824 National protected area Bala'an K'aax. Dark blue color corresponds to non forest areas such as urban
825 settlements, agriculture, and water bodies.

826 Figure 8. Residuals from field-calculated AGB (inventory) - upscaled AGB (satellite) in Mg ha^{-1}
827 distributed by categories of AGB.

828 Figure 9. Kernel-Density Estimation (KDE) plots showing the frequency distribution of AGB [Mg ha^{-1}]
829 ¹] in protected areas ('Kaxil Kiuic' Kaxil Kiuic Biocultural reserve, 'del Puuc' del Puuc Biocultural
830 reserve, 'Bala'an Kaax', Bala'an Kaax protected area) vs unprotected areas (areas designated for
831 restoration and production), based on the median AGB per pixel from the Monte Carlo upscaling
832 process.