THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Concurrent Transitional Meningioma and Ceruminous Gland Adenocarcinoma in a Scottish Wildcat Hybrid (*Felis silvestris*)

Citation for published version:

Drew, SJ, Perpiñán, D & Baily, J 2016, 'Concurrent Transitional Meningioma and Ceruminous Gland Adenocarcinoma in a Scottish Wildcat Hybrid (*Felis silvestris*)', *Journal of Comparative Pathology*, vol. 154, no. 2-3, pp. 253-257. <https://doi.org/10.1016/j.jcpa.2015.11.005>

Digital Object Identifier (DOI):

[10.1016/j.jcpa.2015.11.005](https://doi.org/10.1016/j.jcpa.2015.11.005)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Comparative Pathology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

1 SHORT PAPER

2
3 Concurrent Transitional Meningioma and Ceruminous Gland Adenocarcinoma in a Scottish
4 wildcat hybrid (*Felis silvestris*)

5 S. J. Drew[†] D. Perpiñán[‡] and J. Baily[†]

6 [†]Easter Bush Pathology and [‡]Exotic Animal and Wildlife Service, Royal (Dick) School of
7 Veterinary Studies, The University of Edinburgh, Roslin, Midlothian, UK.

8
9 Corresponding author: Stephen J Drew, Easter Bush Pathology, Royal (Dick) School of
10 Veterinary Studies, The University of Edinburgh, Roslin, Midlothian, UK. EH25 9RG; Tel:
11 0131 651 7455; Stephen.Drew@ed.ac.uk

12
13 Summary

14 The Scottish wildcat (*Felis silvestris*) is an iconic and endangered, sub-population of the
15 European wildcat (*Felis silvestris silvestris*). As such, there is much research devoted to its'
16 ecology, genetics and conservation but little published information on pathology and disease.
17 The investigation and reporting of such information is vital to furthering our understanding of
18 the effects of hybridisation, a factor which is crucial if we are to secure a future for the
19 Scottish wildcat. This report describes the clinical presentation, gross post-mortem and
20 histological findings, in an elderly Scottish wildcat hybrid with concurrent transitional
21 meningioma and ceruminous gland adenocarcinoma. To the authors' knowledge there have
22 been no previous reports of meningioma or ceruminous gland adenocarcinoma in the
23 European wildcat (*Felis silvestris silvestris*) and there are only isolated reports of primary
24 central nervous system neoplasia in other non-domestic felid species.

Keywords: Meningioma; Ceruminous gland adenocarcinoma; European wildcat; *Felis silvestris silvestris*.

The Scottish wildcat is an isolated sub-population of the European wildcat (*Felis silvestris silvestris*) and is also considered by some scientists to be a distinct sub-species (*Felis silvestris grampia*) (Miller, 1912; Beaumont et al., 2001; Kilshaw et al., 2010). The future of the European wildcat, and particularly the Scottish wildcat, is uncertain due to habitat loss, hunting and persecution throughout the 18th and 19th centuries (McOrist and Kitchener, 1994), and more recently the effects of hybridisation, as a result of inter-breeding with domestic cats (*Felis silvestris catus*), (Daniels et al., 2001; Davis and Gray, 2010; Kilshaw et al., 2010), along with the potential transmission of common infectious diseases from domestic cats (McOrist et al., 1991). Now classified as endangered, they are legally protected under both UK and European legislation (Kilshaw et al., 2010). As such, the majority of published literature is focused on its ecology, genetics and conservation.

There have been a limited number of publications regarding pathology and disease of the European wildcat. Most publications are serological studies focusing on the prevalence of common feline viruses (McOrist et al., 1991; Watt et al., 1993; Daniels et al., 1999; Leutenegger et al., 1999; Millán and Rodriguez, 2009; Wasieri et al., 2009), although there are isolated reports dealing with other pathogens such as *Chlamydophila* sp. (Millán and Rodriguez, 2009), *Toxoplasma gondii* (McOrist et al., 1991; Herrmann et al., 2013), lungworm (Falsone et al., 2014), hemoplasmas (Willi et al., 2007), and other endoparasites (Burt et al., 1980; Krone et al., 2008). Neoplasia has been rarely reported in the European wildcat; from 79 necropsy examinations reported in several articles (Jefferies, 1991; McOrist et al., 1991; Watt et al., 1993; Krone et al., 2008; Wasieri et al., 2009; Hermann et al., 2013;

Falsone et al., 2014), only one tumour (a pulmonary lymphoma) was found (Hermann et al., 2013).

This report describes the presentation, gross post-mortem, and histological findings in a Scottish wildcat hybrid with concurrent transitional meningioma and ceruminous gland adenocarcinoma. To the authors' knowledge, there have been no previous reports of meningioma or ceruminous gland adenocarcinoma in the European wildcat (*Felis silvestris silvestris*).

A 16.5 year old, neutered male, Scottish wildcat hybrid from a zoological collection reportedly suffered from sporadic episodes of incoordination and mild lethargy over a four week period. Although this individual was deemed to be a hybrid (*Felis silvestris silvestris* x *Felis silvestris catus*), rather than a true wildcat, according to the records of the zoological institution, a sample of skeletal muscle was tested using a 35 Single Nucleotide

Polymorphism (SNP) marker test developed by the Royal Zoological Society of Scotland from a panel of markers published by Nussberger et al., (2013) in an attempt to confirm this and investigate the degree of hybridisation. Two extracts of DNA were conducted and three replicates of the assay were run but the assay failed to prove or disprove the hybrid status of this individual due to insufficient DNA quality, presumably due to sample degradation. The cat was transferred from a wildlife centre when it was 9 years old, and had since lived with a female Scottish wildcat in an outdoor enclosure. Prior medical history was unremarkable.

Husbandry and nutrition were considered appropriate for the species. Physical examination under general anaesthesia revealed loose skin, thought to be consistent with poor hydration or recent weight loss, a round bony proliferation on the right stifle joint and a hard mass behind the right ear. Body condition score was considered acceptable (4/9) with a weight of 4.5kg, teeth were in an excellent condition for an old cat and vital parameters (heart and respiratory rate, rectal temperature) were unremarkable. The cat initially appeared to recover well from

anaesthesia, but remained recumbent and died 2 hours later. The animal was subsequently submitted for post-mortem examination at the Royal (Dick) School of Veterinary Studies (The University of Edinburgh, Roslin, Midlothian, UK).

Post-mortem examination revealed the entrance to both external ear canals to be obscured by dark grey to blue black, well demarcated, multi-lobular, firm, occasionally cystic, exophytic, polypoid masses ranging from 1 to 5mm in diameter and a cream, well demarcated, 60x30x20mm, multi-lobular, firm, subcutaneous mass at the base of the right ear which contained a dark red, central, well circumscribed, depressed area 3mm in diameter on cut section. The right pre-scapular lymph node was moderately enlarged and the brain contained a cream to yellow, well demarcated, multi-lobular, expansile and compressive mass, approximately 15mm in diameter, between the cerebral hemispheres in the region of the falx cerebri (Fig.1). Samples of all gross lesions along with representative samples from all tissues and internal organs were collected, fixed in 10% buffered formalin and routinely processed according to current histological methods. Sections 5µm thick were stained with haematoxylin and eosin.

Histopathological examination of the external ear canal masses (Fig.2) revealed a moderately acanthotic, stratified squamous keratinising epithelium elevated by numerous, multi-focal, dilated, cystic, glandular structures lined by a flattened to low cuboidal epithelium. Many cysts contained amorphous, tan to pale brown, granular material (cerumen) and low numbers of large foamy macrophages also contained this granular brown material. Within a mass from the right external ear canal, the superficial dermis was focally expanded by more dense proliferations of polygonal to cuboidal cells forming tubules, acini and fronds. Cells had variably well-defined cell borders, a moderate amount of eosinophilic, finely granular cytoplasm, open-faced, oval to round nuclei with 1-2 prominent nucleoli. Anisocytosis and anisokaryosis were mild with an average of 1 mitotic figure per high power field (x400). Foci

99 of necrosis expanded the centre of these cellular proliferations. Examination of the right ear
100 base mass (Fig.3) was consistent with the pre-existing architecture of a lymph node, 80% of
101 which was effaced by tubules and acini of cells, similar to those described in the mass from
102 the right external ear canal, with evidence of lymphatic invasion. Similar cell proliferation
103 was present in the right pre-scapular lymph node.

104 The histopathological appearance of the auricular lesions was consistent with a diagnosis of
105 bilateral ceruminous gland dilatation and hyperplasia (feline ceruminous cystomatosis) with
106 unilateral (right side) ceruminous gland adenocarcinoma and metastasis to cervical and pre-
107 scapular lymph nodes.

108 Sections of cerebral cortex, lateral ventricle and meninges showed the leptomeninges of the
109 cingulate sulcus to be expanded by a large, densely cellular, well demarcated, finely
110 encapsulated, compressive, nodular mass (Fig.4). The mass consisted predominantly of
111 lobules of loosely to densely packed cells forming concentric whorls, occasionally
112 surrounding blood vessels or containing central areas of necrosis. Adjacent cells formed long,
113 interlacing fascicles streaming around the whorls and were supported by a loosely arranged,
114 eosinophilic, fibrillar stroma. Cells were fusiform to polygonal with indistinct cell borders
115 and variable amounts of eosinophilic, wispy, fibrillar cytoplasm. Within the centre of
116 concentric whorls, cells adopted a more epithelioid morphology. Nuclei were round to
117 elongated with loosely packed, finely stippled chromatin and 1 to 2 nucleoli. Anisocytosis
118 and anisokaryosis were moderate with occasional multi-nucleate forms. Mitoses were rare (1
119 figure per 10 high power fields (x400)). Moderate numbers of neutrophils, occasional
120 lymphocytes, plasma cells, pyknotic cells, small clusters of foamy macrophages and
121 occasional acicular clear spaces (cholesterol clefts) were present within the mass. Both the
122 grey matter and white matter tracts of the adjacent cerebral cortex contained mild diffuse
123 vacuolation with mild perivascular clearing and spacing (oedema).

The histopathological findings of the cerebral mass were consistent with a diagnosis of transitional (mixed) meningioma containing features of both meningotheial (characterised by moderately cellular lobules of polygonal cells) and fibrous tumours (long interlacing fascicles of fusiform cells).

Also identified following post-mortem and histological examination were bilateral stifle joint osteoarthritis, nodular hyperplasia of both thyroid glands, mild hepatic lipidosis, mild, multi-focal, chronic cholangiohepatitis and mild, multi-focal, chronic interstitial nephritis with a focal, chronic, renal infarct.

Ceruminous gland adenocarcinomas are the most commonly diagnosed tumour of the external acoustic meatus in cats, accounting for up to 2% of all feline neoplasms (Njaa and Wilcock, 2012) and are more frequently diagnosed than adenomas in domestic cats (Moisan and Watson, 1996; London et al., 1996). They exhibit locally invasive behaviour (London et al., 1996) and metastasis to regional lymph nodes, lungs and viscera can occur in up to 50% of cases (Njaa and Wilcock, 2012). No evidence of pulmonary or visceral metastasis was found in this case.

Differentiation between ceruminous adenoma and adenocarcinoma can be challenging unless there is evidence of local invasion or metastatic disease, as in this case (Wilcock et al., 2002).

Meningiomas are the most common primary central nervous system (CNS) neoplasm of domestic cats (Koestner & Higgins, 2002; Troxel et al., 2003; Tomek et al., 2006; Motta et al., 2012) typically occurring in cats older than 9 years (Troxel et al., 2003; Tomek et al., 2006) with an increasing incidence with age. Domestic shorthaired cats seem to be predisposed but no significant sex predilection has been found (Troxel et al., 2003; Tomek et al., 2006). Transitional meningiomas, as reported here, and fibrous subtypes are most frequently encountered in domestic cats. Meningiomas are typically slow growing, with the exception of the uncommon anaplastic (malignant form), rarely metastatic and approximately

50% of cases do not exhibit any clinical signs. To the authors' knowledge this is the first report of meningioma in a European wildcat. There are only isolated additional reports in the literature of central nervous system neoplasia in non-domestic felids, such as meningioma in a Bengal tiger (*Panthera tigris tigris*) (Akin et al., 2013) and intracranial oligodendroglioma in a lion (*Panthera leo*) (Tucker et al., 2008).

The findings reported here pose the question for future studies to determine the incidence of common neoplasms of domestic felines in pure-bred wildcats and their hybrids. This may help to elucidate whether their occurrence may be an associated effect of hybridisation, or purely a reflection of the increasing age of individuals living in zoological collections in comparison to their free living relatives.

Acknowledgement: We thank Roisin Campbell-Palmer, Conservation Projects Manager, Jennifer Kaden, Senior Technician and Helen Senn, Research Scientist, The Royal Zoological Society of Scotland, Edinburgh, EH12 6TS for their assistance with attempts at genetic identification. Thanks also, to the staff of Five Sisters Zoo, Gavieside, West Calder, West Lothian, EH55 8PT.

Conflict of Interest Statement: The authors declare no conflict of interest.

References:

- Akin EY, Baumgartner WA, Lee JK, Beasley MJ (2013) Meningioma in a bengal tiger (*Panthera tigris tigris*). Journal of Zoo and Wildlife Medicine, **44**(3), 761-764.
- Beaumont M, Barratt EM, Gottelli D, Kitchener AC, Daniels MJ et al. (2001) Genetic diversity and introgression in the Scottish wildcat. Molecular Ecology, **10**, 319-336.

173 Burt MDB, Pike AW, Corbett LK (1980) Helminth parasites of wildcats in north-east
174 Scotland. Journal of Helminthology, **54**, 303-308.

175 Daniels MJ, Golder MC, Jarrett O, MacDonald DW (1999) Feline viruses in wildcats from
176 Scotland. Journal of Wildlife Diseases, **35(1)**, 121-124.

177 Daniels MJ, Beaumont MA, Johnson PJ, Balharry D, Macdonald DW et al. (2001) Ecology
178 and genetics of wild-living cats in the north-east of Scotland and the implications for
179 the conservation of the wildcat. Journal of Applied Ecology, **38**, 146-161.

180 Davis AR, Gray D (2010) The distribution of Scottish wildcats (*Felis silvestris*) in Scotland
181 (2006-2008). Scottish Natural Heritage Commissioned Report No. 360.

182 Falsone L, Brianti E, Gaglio G, Napoli E, Anile S et al. (2014) The European wildcats (*Felis*
183 *silvestris silvestris*) as reservoir hosts of *Troglostrongylus brevior* (Strongylida:
184 Crenosomatidae) lungworms. Veterinary Parasitology, **205(1-2)**, 193-198.

185 Herrmann DC, Wibbelt G, Götz M, Conraths FJ, Schares G (2013) Genetic characterisation
186 of *Toxoplasma gondii* isolates from European beavers (*Castor fiber*) and European
187 wildcats (*Felis silvestris silvestris*). Veterinary Parasitology, **191**, 108-111.

188 Jefferies DJ (1991) Some observations on Scottish wildcats (*Felis silvestris*) based on the
189 results of autopsies. The Glasgow Naturalist, **22(1)**, 11-19.

190 Kilshaw K, Drake A, Macdonald DW, Kitchener AC (2010) The Scottish wildcat: a
191 comparison of genetic and pelage characteristics. Scottish Natural Heritage
192 Commissioned Report No.356.

193 Koestner A, Higgins RJ (2002) Tumors of the nervous system. In: Tumors in Domestic
194 Animals, 4th Edit., DJ Meuten, Ed., Iowa State Press, Iowa, pp. 697-738.

195 Krone O, Guminsky O, Meinig H, Herrmann M, Trinzen M et al. (2008) Endoparasite
196 spectrum of wild cats (*Felis silvestris* Schreber,1777) and domestic cats (*Felis catus*

197 L.) from the Eifel, Pfalz region and Sarland, Germany. European Journal of Wildlife
198 Research, **54**(1), 95-100.

199 Leutenegger CM, Hofmann-Lehmann R, Riols C, Liberek M, Worel G et al. (1999) Viral
200 infections in free-living populations of the European wildcat. Journal of Wildlife
201 Diseases, **35**, 678-686.

202 London CA, Dubilzeig RR, Vail DM, Ogilvie GK, Hahn KA et al. (1996) Evaluation of dogs
203 and cats with tumors of the ear canal: 145 cases (1978-1992). Journal of the American
204 Veterinary Medical Association, **208**(9), 1413-1418.

205 McOrist S, Bold R, Jones TW, Easterbee N, Hubbard L et al. (1991) Some viral and
206 protozool diseases in the European wildcat (*Felis silvestris*). Journal of Wildlife
207 Diseases, **27**(4), 693-696.

208 McOrist S, Kitchener AC (1994) Current threats to the European wildcat, *Felis silvestris*, in
209 Scotland. Ambio, **23**(4/5), 243-245.

210 Millán J, Rodriguez A (2009) A serological survey of common feline pathogens in free-living
211 European Wildcats (*Felis silvestris*) in central Spain. European Journal of Wildlife
212 Research, **55**, 285-291.

213 Miller GS (1912) Catalogue of the mammals of western Europe (Europe exclusive of Russia)
214 in the collection of the British Museum (Natural History), London.

215 Moisan PG, Watson GL (1996) Ceruminous gland tumors in dogs and cats: a review of 124
216 cases. Journal of the American Animal Hospital Association, **32**, 449-453.

217 Motta L, Mandara MT, Skerritt GC (2012) Canine and feline intracranial meningiomas: an
218 updated review. The Veterinary Journal, **192**, 153-165.

219 Njaa BL, Wilcock BP (2012) The ear and eye. In: Pathologic Basis of Veterinary Disease, 5th
220 Edit., JF Zachary, MD McGavin Eds., Elsevier, Missouri, pp. 1153-1244.

221 Nussberger B, Greminger MP, Grossen C, Keller LF, Wandeler P (2013) Development of
 222 SNP markers identifying European wildcats, domestic cats and their admixed
 223 progeny. *Molecular Ecology Resources*, **13**, 447-460.

224 Tomek A, Cizinauskas S, Doherr M, Gandini G, Jaggy A (2006) Intracranial neoplasia in 61
 225 cats: localisation, tumour types and seizure patterns. *Journal of Feline Medicine and*
 226 *Surgery*, **8**, 243-253.

227 Troxel MT, Vite CH, Van Winkle TJ, Newton AL, Tiches D et al. (2003) Feline intracranial
 228 neoplasia: retrospective review of 160 cases (1985–2001). *Journal of Veterinary*
 229 *Internal Medicine*, **17**, 850-859.

230 Tucker AR, Ramsay EC, Donnell RL (2008) Oligodendroglioma in an African lion (*Panthera*
 231 *leo*). *Journal of Zoo and Wildlife Medicine*, **39**, 650-654.

232 Wasieri J, Schmiedeknecht G, Förster C, König M, Reinacher M (2009) Parvovirus infection
 233 in a Eurasian Lynx (*Lynx lynx*) and in a European Wildcat (*Felis silvestris silvestris*).
 234 *Journal of Comparative Pathology*, **140**, 203-207.

235 Watt NJ, MacIntyre NJ, McOrist S (1993) An extended outbreak of infectious peritonitis in a
 236 closed colony of European wildcats (*Felis silvestris*). *Journal of Comparative*
 237 *Pathology*, **108**, 73-79.

238 Wilcock B, Dubielzig RR, Render R (2002) Tumors of the external ear canal. In: *Histological*
 239 *Classification of Ocular and Otic Tumors of Domestic Animals* WHO International
 240 *Histological Classification of Tumors of Domestic Animals*, Series 2, Vol.9, FY
 241 Schulman Ed., Armed Forces Institute of Pathology, Washington DC, USA, p. 36.

242 Willi B, Filoni C, Catão-Dias JL, Cattori V, Meli ML et al. (2007) Worldwide occurrence of
 243 feline hemoplasma infections in wild felid species. *Journal of Clinical Microbiology*,
 244 **45(4)**, 1159-1166.

Legend for Figures:

Fig 1. Brain, showing the presence of a nodular mass between the cerebral hemispheres.

Fig 2. Section of tissue from the right external ear canal showing acinar arrangement of neoplastic epithelial cells with central necrosis. Epithelial cells are present within the lumen of the thin-walled vessel at the top of the image. HE.

Fig 3. Section of the right cervical lymph node showing replacement of normal architecture by metastatic neoplastic epithelia forming acinar structures. HE.

Fig 4. Section from the brain mass showing whorls and bundles of neoplastic spindle cells consistent with the transitional form of meningioma. HE.