



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Aging-related tau astroglipathy (ARTAG)

Citation for published version:

Kovacs, GG, Ferrer, I, Grinberg, LT, Alafuzoff, I, Attems, J, Budka, H, Cairns, NJ, Crary, JF, Duyckaerts, C, Ghetti, B, Halliday, GM, Ironside, JW, Love, S, Mackenzie, IR, Munoz, DG, Murray, ME, Nelson, PT, Takahashi, H, Trojanowski, JQ, Ansorge, O, Arzberger, T, Baborie, A, Beach, TG, Bieniek, KF, Bigio, EH, Bodi, I, Dugger, BN, Feany, M, Gelpi, E, Gentleman, SM, Giaccone, G, Hatanpaa, KJ, Heale, R, Hof, PR, Hofer, M, Hortobágyi, T, Jellinger, K, Jicha, GA, Ince, P, Kofler, J, Kövari, E, Kril, JJ, Mann, DM, Matej, R, McKee, AC, McLean, C, Milenkovic, I, Montine, TJ, Murayama, S, Lee, EB, Rahimi, J, Rodriguez, RD, Rozemüller, A, Schneider, JA, Schultz, C, Seeley, W, Selkoe, D, Smith, C, Tagliavini, F, Takao, M, Thal, DR, Toledo, JB, Tolnay, M, Troncoso, JC, Vinters, HV, Weis, S, Wharton, SB, White, CL, Wisniewski, T, Woulfe, JM, Yamada, M & Dickson, DW 2016, 'Aging-related tau astroglipathy (ARTAG): harmonized evaluation strategy', *Acta Neuropathologica*, vol. 131, no. 1, pp. 87-102. <https://doi.org/10.1007/s00401-015-1509-x>

Digital Object Identifier (DOI):

[10.1007/s00401-015-1509-x](https://doi.org/10.1007/s00401-015-1509-x)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Acta Neuropathologica

Publisher Rights Statement:

This is the author's a final peer-reviewed manuscript as accepted for publication.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Aging-related tau astroglialopathy (ARTAG): harmonized evaluation strategy

Gabor G. Kovacs¹, Isidro Ferrer², Lea T. Grinberg^{3,4}, Irina Alafuzoff⁵, Johannes Attems⁶, Herbert Budka⁷, Nigel J. Cairns⁸, John F. Crary⁹, Charles Duyckaerts¹⁰, Bernardino Ghetti¹¹, Glenda M. Halliday¹², James W. Ironside¹³, Seth Love¹⁴, Ian R. Mackenzie¹⁵, David G. Munoz¹⁶, Melissa E. Murray¹⁷, Peter T. Nelson¹⁸, Hitoshi Takahashi¹⁹, John Q. Trojanowski²⁰, Olaf Ansorge²¹, Thomas Arzberger²², Atik Baborie²³, Thomas G. Beach²⁴, Kevin F. Bieniek¹⁷, Eileen H. Bigio²⁵, Istvan Bodi²⁶, Brittany N. Dugger^{24,27}, Mel Feany²⁸, Ellen Gelpi²⁹, Stephen M. Gentleman³⁰, Giorgio Giaccone³¹, Kimmo J. Hatanpää³², Richard Heale⁶, Patrick R. Hof³³, Monika Hofer²¹, Tibor Hortobágyi³⁴, Kurt Jellinger³⁵, Gregory A. Jicha³⁶, Paul Ince³⁷, Julia Kofler³⁸, Enikő Kövari³⁹, Jillian J. Kril⁴⁰, David M. Mann⁴¹, Radoslav Matej⁴², Ann C. McKee⁴³, Catriona McLean⁴⁴, Ivan Milenkovic^{1,45}, Thomas J. Montine⁴⁶, Shigeo Murayama⁴⁷, Edward B. Lee²⁰, Jasmin Rahimi¹, Roberta D. Rodriguez⁴⁸, Annemieke Rozemüller⁴⁹, Julie A. Schneider⁵⁰, Christian Schultz⁵¹, William Seeley³, Danielle Seilhean¹⁰, Colin Smith¹³, Fabrizio Tagliavini³¹, Masaki Takao⁵², Dietmar Rudolf Thal^{53,54}, Jon B. Toledo²⁰, Markus Tolnay⁵⁵, Juan C. Troncoso⁵⁶, Harry V. Vinters⁵⁷, Serge Weis⁵⁸, Stephen B. Wharton³⁷, Charles L. White, III³², Thomas Wisniewski⁵⁹, John M. Wolfe⁶⁰, Masahito Yamada⁶¹, and Dennis W. Dickson¹⁷

Affiliations:

- 1: Institute of Neurology, Medical University of Vienna, Vienna, Austria;
- 2: Institute of Neuropathology, Bellvitge University Hospital, University of Barcelona, Hospitalet de Llobregat, Barcelona, Spain;
- 3: Memory and Aging Center, Department of Neurology, University of California, San Francisco
- 4: Department of Pathology, LIM-22, University of Sao Paulo Medical School, Sao Paulo, Brazil
- 5: Department of Immunology, Genetics and Pathology, Uppsala University, Uppsala, Sweden
- 6: Institute of Neuroscience, Newcastle University, Newcastle upon Tyne, UK
- 7: Institute of Neuropathology, University Hospital Zürich, Switzerland
- 8: Department of Pathology and Immunology, Washington University School of Medicine, St. Louis, Missouri, USA
- 9: Department of Pathology, Fishberg Department of Neuroscience, Friedman Brain Institute, and the Ronald M. Loeb Center for Alzheimer's Disease, Icahn School of Medicine at Mount Sinai
- 10: Neuropathology Department, Hopital de La Salpêtrière, AP-HP, UPMC-Sorbonne-University, Paris, France
- 11: Indiana University School of Medicine Department of Pathology and Laboratory Medicine, Indianapolis, IN, USA
- 12: GMH - Neuroscience Research Australia and the University of New South Wales, Sydney, NSW, Australia
- 13: Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh UK
- 14: Institute of Clinical Neurosciences, University of Bristol, Learning & Research level 2, Southmead Hospital, Bristol, UK
- 15: Department of Pathology and Laboratory Medicine, University of British Columbia, Vancouver, Canada
- 16: Division of Pathology, St. Michael's Hospital 30 Bond St, Toronto, ON, Canada
- 17: Department of Neuroscience, Mayo Clinic, 4500 San Pablo Road, Jacksonville, FL 32224, USA
- 18: Department of Pathology and Sanders-Brown Center on Aging, University of Kentucky, Lexington, KY USA 40536, USA

- 19: Department of Pathology, Brain Research Institute, Niigata University, Niigata 951-8585, Japan
- 20: Center for Neurodegenerative Disease Research, Institute on Aging and Department of Pathology & Laboratory Medicine of the Perelman School of Medicine at the University of Pennsylvania
- 21: Department of Neuropathology, John Radcliffe Hospital, Oxford, UK
- 22: Department of Psychiatry and Psychotherapy, and Centre for Neuropathology and Prion Research, Ludwig-Maximilians-University Munich, Germany
- 23: Department of Neuropathology, Walton Centre, Liverpool, UK
- 24: Civin Laboratory for Neuropathology, Banner Sun Health Research Institute, Sun City, AZ 85351, USA
- 25: Northwestern ADC Neuropathology Core, Northwestern University Feinberg School of Medicine, Chicago, IL, USA
- 26: Clinical Neuropathology, King's College Hospital and London Neurodegenerative Brain Bank, London, UK
- 27: University of California San Francisco, Institute for Neurodegenerative Diseases, San Francisco, CA, USA
- 28: Department of Pathology, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA
- 29: Neurological Tissue Bank of the Biobank-Hospital Clinic-IDIBAPS, Institut d'Investigacions Biomediques August Pi i Sunyer, Barcelona, Spain
- 30: Department of Medicine, Imperial College London, London, UK
- 31: IRCCS Foundation "Carlo Besta" Neurological Institute, Milan, Italy
- 32: Department of Pathology, University of Texas Southwestern Medical Center, Dallas, TX, USA
- 33: Fishberg Department of Neuroscience, Friedman Brain Institute, and Ronald M. Loeb Center for Alzheimer's Disease, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA
- 34: Department of Neuropathology, Institute of Pathology, Faculty of Medicine, University of Debrecen, Nagyerdei krt. 98, H-4032, Debrecen, Hungary.
- 35: Institute of Clinical Neurobiology; Alberichgasse 5/13; A-1150 Vienna
- 36: Department of Neurology and Sanders-Brown Center on Aging, University of Kentucky, Lexington, KY USA 40536
- 37: Sheffield Institute for Translational Neuroscience, University of Sheffield, UK
- 38: Department of Pathology, University of Pittsburgh, Pittsburgh, PA, USA
- 39: Department of Mental Health and Psychiatry, University Hospitals and University of Geneva School of Medicine, Geneva, Switzerland
- 40: Discipline of Pathology, Sydney Medical School, The University of Sydney, Sydney NSW 2006
- 41: Institute of Brain, Behaviour and Mental Health, Manchester University Faculty of Medical and Health Sciences, Manchester, UK
- 42: Department of Pathology and Molecular Medicine, Thomayer Hospital, Prague 4, Czech Republic
- 43: Department of Neurology and Pathology, Boston University School of Medicine and VA Healthcare system
- 44: Department of Anatomical Pathology, Alfred Hospital, Prahran, Victoria, 3004, Australia
- 45: Department of Neurology, Medical University of Vienna, Vienna, Austria
- 46: Department of Pathology, University of Washington, Seattle, WA, USA
- 47: Department of Neuropathology (the Brain Bank for Aging Research), Tokyo Metropolitan Geriatric Hospital & Institute of Gerontology, Tokyo, Japan
- 48: Physiopathology in Aging Lab/Brazilian Aging Brain Study Group-LIM22, University

of Sao Paulo Medical School, Sao Paulo, Brazil
49: Netherlands Brainbank and Dept. of Pathology, VU University Medical Center, Amsterdam, The Netherlands
50: Departments of Pathology and Neurological Sciences, Rush University Medical Center, Chicago IL, USA
51: Institute of Neuroanatomy, Centre for Biomedicine and Medical Technology Mannheim, Medical Faculty Mannheim, Heidelberg University, Germany
52: Department of Neurology, Saitama Medical University International Medical Center, Saitama, Japan
53: Institute of Pathology – Laboratory of Neuropathology, University of Ulm, D-89081 Ulm, Germany
54: Department of Neuroscience, KU-Leuven, B-3000 Leuven, Belgium
55: Institute of Pathology, University Hospital Basel, Basel, Switzerland.
56: Johns Hopkins University School of Medicine, Department of Pathology, Division of Neuropathology, Baltimore, MD, USA
57: Section of Neuropathology, Department of Pathology and Laboratory Medicine, and Department of Neurology, Brain Research Institute, University of California, Los Angeles (UCLA) Medical Center and David Geffen School of Medicine, Los Angeles, California, USA
58: Laboratory of Neuropathology, Department of Pathology and Neuropathology, State Neuropsychiatric Hospital Wagner-Jauregg, Medical School, Johannes Kepler University Linz, Linz, Austria
59: Center for Cognitive Neurology, Departments of Neurology, Pathology and Psychiatry, New York University School of Medicine, ERSP, 450 East 29th Street, NY, NY, USA 10016
60: Centre for Cancer Therapeutics, Ottawa Hospital Research Institute, Department of Pathology and Laboratory Medicine, University of Ottawa
61: Department of Neurology & Neurobiology of Aging, Kanazawa University Graduate School of Medical Sciences, Kanazawa, Japan

Word count abstract: 299

Word count text: 4817

Number of Figures: 4

Number of tables: 3

Corresponding authors:

Gabor G. Kovacs MD PhD

Institute of Neurology, AKH 4J, Währinger Gürtel 18-20, A-1097 Vienna, Austria

Phone: +43-1-40400-55070; Fax +43-1-40400-55110;

Email: gabor.kovacs@meduniwien.ac.at

and

Dennis W. Dickson MD

Department of Neuroscience, Mayo Clinic, 4500 San Pablo Road, Jacksonville, FL 32224,

USA

Email: dickson.dennis@mayo.edu

Abstract

Pathological accumulation of abnormally phosphorylated tau protein in astrocytes is a frequent,

but poorly characterized feature of the aging brain. Its etiology is uncertain, but its presence is sufficiently ubiquitous to merit further characterization and classification, which may stimulate clinicopathological studies and research into its pathobiology. This paper aims to harmonize evaluation and nomenclature of aging-related tau astroglial pathology (ARTAG), a term that refers to a morphological spectrum of astroglial pathology detected by tau immunohistochemistry, especially with phosphorylation-dependent and 4R isoform specific antibodies. ARTAG occurs mainly, but not exclusively, in individuals over 60 years of age. Tau-immunoreactive astrocytes in ARTAG include thorn-shaped astrocytes at the glia limitans and in white matter, as well as solitary or clustered astrocytes with perinuclear cytoplasmic tau immunoreactivity that extends into the astroglial processes as fine fibrillar or granular immunopositivity, typically in gray matter. Various forms of ARTAG may co-exist in the same brain and might reflect different pathogenetic processes. Based on morphology and anatomical distribution, ARTAG can be distinguished from primary tauopathies but may be concurrent with primary tauopathies or other disorders. We recommend four steps for evaluation of ARTAG: 1) identification of five types based on the location of either of the two morphologies of tau astroglial pathology: subpial, subependymal, gray matter, white matter, perivascular; 2) documentation of the regional involvement: medial temporal lobe, lobar (frontal, parietal, occipital, lateral temporal), subcortical, brainstem; 3) documentation of the severity of tau astroglial pathology; and 4) description of subregional involvement. Some types of ARTAG may underlie neurological symptoms; however, the clinical significance of ARTAG is currently uncertain and awaits further studies. The goal of this proposal is to raise awareness of astroglial tau pathology in the aged brain, facilitating communication among neuropathologists and researchers, and informing interpretation of clinical biomarkers and imaging studies that focus on tau-related indicators.

Key words: aging; ARTAG; tau astroglial pathology; tau;

Introduction

Tau is a microtubule-associated protein that binds to tubulin and promotes its polymerization and stabilization into microtubules. Tau isoforms, ranging from 352 to 441 amino acids, are generated by the alternative splicing of exons 2, 3, and 10 of the *MAPT* gene. The six isoforms differ from each other by the presence or absence of 29- or 58-amino acid inserts in the N-terminus domain and by the presence of either three (3R tau isoforms) or four (4R tau isoforms) tandem repeat sequences of 31 or 32 amino acids [24]. Mutations in the tau gene (*MAPT*) can cause hereditary frontotemporal dementia and associate with frontotemporal lobar degeneration (FTLD) [23, 26, 51, 63]. Following the description of a disorder in one family named ‘multiple system tauopathy with presenile dementia’ [62], the term tauopathy was introduced to refer to disorders in which tau protein deposition is the predominant feature [23]. Tauopathies are characterized by the accumulation of abnormal and hyper-phosphorylated tau protein in the brain and are also classified as primary or secondary [32, 37]. Tau pathology is characterized as 3R or 4R predominant or mixed 3R+4R type [12, 30, 32]. Primary tauopathies are grouped also as FTLD-tau and comprise Pick disease (PiD), progressive supranuclear palsy (PSP), corticobasal degeneration (CBD), argyrophilic grain disease (AGD), neurofibrillary tangle (NFT) predominant senile dementia (NFT-dementia or “tangle-only” dementia; now included in the category of PART, see below), and globular glial tauopathy (GGT) [32, 47]. In addition, many other diseases or conditions with diverse etiology, including Alzheimer disease (AD), may be associated with tau pathology [32]. The recently introduced term ‘primary age-related tauopathy’ (PART) encompasses neuronal changes previously considered as “normal aging” as well as NFT-dementia [14]. PART is distinguished from AD, largely by the absence or scarcity of amyloid (A β) plaques [14]. In aged individuals sex-dependent tau pathology, developing independently from AD has been also described in the hypothalamus [16, 54, 56]. Furthermore, chronic traumatic encephalopathy (CTE) is associated with a distinctive pattern of progressive neuronal and glial tau pathology [40-42].

The introduction of the Gallyas silver stain and particularly diagnostic tau immunohistochemistry led to the identification of astroglial tau pathology in the aging brain in people with or without AD-related changes, cognitive decline or movement disorders [5, 7, 11, 21, 25, 31, 34-36, 38, 46, 57]. There have been attempts to classify these tau astroglipathies [34], but there is lack of consensus as to how best to describe and categorize them. We recommend the term *aging-related tau astroglipathy* (ARTAG) to describe the spectrum of otherwise unclassified tau immunoreactivity in astrocytes (i.e., distinct from tufted astrocytes, astrocytic plaques, ramified astrocytes, or globular astroglial inclusions) mostly in aged

individuals detected by tau immunohistochemistry using phosphorylation-specific, conformation-specific, or isoform-specific (4R) anti-tau antibodies. Both ARTAG and PART affect predominantly the elderly, but PART is characterized by neurofibrillary degeneration that is largely restricted to the medial temporal lobe (MTL), basal forebrain, brainstem, and olfactory bulb and cortex [14]. PART thus describes neuronal tau pathology, while ARTAG focuses on astrocytic tau pathology. Whether PART and ARTAG have separate or shared pathogenic pathways is unknown.

We propose a four-step approach for the morphological classification of ARTAG. We anticipate that harmonizing the nomenclature and improving consistency in documentation of ARTAG is a necessary first step for defining diagnostic guidelines that will result in progress in clinicopathological correlation and investigation of the pathogenesis of ARTAG.

Morphology of tau-immunoreactive astrocytes in primary tauopathies and CTE

The defining lesions of tauopathies are intracellular aggregates of abnormal conformers of tau, consistently detectable by immunohistochemistry for phosphoepitopes (e.g., PHF1, CP13, and AT8), as well as epitopes to conformational epitopes (e.g., Alz50 and MC1) and tau isoform specific epitopes (e.g., 4R tau isoforms) [32]. Regardless of the tauopathy, astroglial tau inclusions are mostly 4R tau-immunopositive, although ramified astrocytes in PiD as well as occasional protoplasmic astrocytes in PSP may show 3R tau immunoreactivity [21]. Tufted astrocytes are characteristic of PSP, and astrocytic plaques are signature lesions of CBD, while so-called ramified astrocytes have been described in PiD [15, 32]. In addition, astroglial, argyrophilic, and intracytoplasmic flame or thorn-shaped inclusions were described by Nishimura et al. in PSP [49]. Phosphorylation-dependent anti-tau antibodies are highly sensitive and label lesions that are not consistently detectable by silver impregnation methods, but may show variable ubiquitin or p62/sequestosome immunopositivity, such as the globular astroglial inclusions (GAI) of GGT [1], or the fine granular tau immunopositivity (some with ‘bush-like’ appearance) of the astrocytes of AGD [11]. Some of the variation in the morphology of the immunolabeled structures was interpreted as representing stages of a process of aggregation and fibrillation, analogous to progression from pretangles to neurofibrillary tangles in AD [6, 11]. The concept of early-stage tau accumulation in astrocytes has also been discussed in relation to the changes in the basal ganglia in PSP [53]. Finally, subpial and subependymal clusters of astrocytic tangles have been described in CTE [41].

Overview of astrocytic tau pathologies in the aging brain

Both neuronal and glial tau pathology increases in frequency with age. The most frequent neuronal tau inclusions are neurofibrillary tangles, threads, and argyrophilic grains. Neuronal and glial inclusions resembling PSP pathology can be seen in the elderly, even without clinical evidence of PSP [17, 18, 34], but these lack the typical multisystem degeneration seen with PSP. Furthermore, tuft-shaped astrocytes have been described in a subgroup of elderly individuals, especially in association with Lewy body pathology, in a distribution resembling that of PSP [25]. Nevertheless, converging data emphasize the presence of a tau astroglial pathology that differs from tufted astrocytes or astrocytic plaques as a common finding in the elderly. Despite its high prevalence, there is a lack of consensus on whether these astroglial tau pathologies in the elderly are clinically relevant, even as a concomitant pathology that might lower an individual's threshold for the development of clinical symptoms. Research in this field has been hampered by the variation in staining methods, tau antibodies, and the inconsistent nomenclature for astroglial tau pathologies. Importantly, hypertrophic astrocytes, as revealed by haematoxylin and eosin staining and immunohistochemistry for glial fibrillary acidic protein and excitatory amino acid transporter 2 (EAAT2), are common in the elderly, and presumably represent a reaction to multiple types of injury. The location of such reactive astrocytes varies considerably among individuals [8, 59]. Colocalization studies have indicated that glial fibrillary acidic protein-immunoreactive reactive astrocytes are not necessarily those that are also immunoreactive for tau pathology [19].

Ikeda and colleagues were the first to describe tau-positive thorn-shaped astrocytes (TSA), which were similar in morphology to tau-positive astrocytes described by Nishimura et al. in PSP [49]) in the subpial or subependymal regions of the gray and white matter and frequently in the depths of gyri, as well as in the basal forebrain and brainstem, in aged individuals [27-29]. TSA may occur in multiple conditions [13]. In comparison to the tufted astrocytes of PSP, TSA showed more voluminous perinuclear cytoplasm and their processes are often thicker and shorter [27]. TSA were only occasionally found in the deep cortical layers. The authors noted that anti-ubiquitin antibodies do not label TSA. They interpreted TSA as a non-specific finding and found no relationship between the number of TSA and the severity of neurofibrillary changes. Argyrophilic, tau-positive subpial and perivascular structures were also described as common TSA [27-29].

Schultz et al. reported a high prevalence of TSA in the aged human MTL, particularly the

anterior MTL, at the level of the amygdala [57]. TSA were absent in individuals under 60 years, but affected almost half of brains from those over 75 years [57]. Indeed, another study also failed to find this type of tau astroglia in younger individuals [33]. Schultz et al. [57] commented that tau immunopositivity was not confined to the thorn-shaped proximal processes of astroglia, but also presented as thread-like processes in the neuropil. They found that immunolabeling with AT8 was the most sensitive for demonstrating the TSA, while silver staining was less consistent [57]. They also speculated that the preferential subpial and perivascular location could be a result of astrocytic reaction to exposure to extravasated plasma proteins from the cerebrospinal fluid due to defects in blood-brain barrier permeability commonly seen in aging and neurodegeneration [36, 57]. Interestingly, a similar distribution of TSA was reported in aged baboons [55]. A study by the MRC-CFAS group confirmed the findings of Schultz et al. and added that the TSA could be less commonly observed in the vicinity of neuronal cell bodies in gray matter areas such as amygdala and dentate gyrus [36]. Also, this study documented the 4R tau nature of TSA [36]. Variable staining for Gallyas and p62 suggest that some of these astrocytes accumulate tau in a fibrillar state [34-36, 38, 46]. Uchikado et al. also reported that the frequency of TSA increased with age and was independent of AGD [66]. All studies agree that the burden of TSA is independent of AD pathology, AGD, coiled bodies, dementia status at death, or presence of the *APOE* ϵ 4 allele [27, 28, 36, 57, 66]. These studies, however, were limited to evaluation of the MTL and did not take account of cortical and subcortical tau astroglia. Few studies reported tau immunopositivity in glial cells in AD cases with prolonged duration of the disease [5, 48, 68]. Finally, a report on NFT-dementia mentioned the presence of astrocytic tau pathology in white matter and cortex [31].

The possibility that TSA may have clinical significance was first raised by Munoz and colleagues [46]. They used the term "argyrophilic thorny astrocyte clusters (ATACs)" and observed them in the fronto-temporo-parietal cortices and in subcortical white matter in a cohort of patients with nonfluent variant of primary progressive aphasia associated with AD pathology [46]. Subsequent reports also linked TSA to symptomatology, although not all found an association of ATACs with focal neurological syndromes [9, 43]. Munoz and colleagues noted ATACs in the white matter without discernible changes in sections stained for myelin. ATACs did not show a consistent topographic relationship to amyloid deposits, NFTs, or reactive astrogliosis [46]. Interestingly, focal glial tauopathy, interpreted as PSP-type, associated with progressive aphasia was reported by Wakabayashi et al. [67]. These observations suggested that TSA-like astrocytes might be detected not only in a subependymal or subpial location.

A peculiar constellation of tau pathology was documented in a subset of patients with dementia [35]. The most characteristic feature was a tau astrogliopathy, which was described as diffuse granular tau immunoreactivity in astrocytic processes [35]. The study emphasized additional neuronal pathologies, including threads and diffuse neuronal cytoplasmic tau immunoreactivity (pretangle-like). A further study distinguished four different patterns based on the anatomical distribution of the tau astrogliopathy and its combination with neuronal tau pathology, characterized mostly by pretangles and scattered threads [34]: 1) medial temporal lobe type (*Group I*); 2) amygdala type (*Group II*); 3) limbic-basal ganglia-nigral type with neuronal tauopathy (*Group III*); and 4) hippocampus-dentate gyrus-amygdala type with neuronal tauopathy (*Group IV*). Some of these might represent stages of the same process whereas other might be different. Nevertheless, evaluation of tau astrogliopathy in several anatomical regions indicated that in some cases astroglial tau pathology in the elderly extends beyond the MTL to involve the frontal and parietal cortices, striatum, substantia nigra, and medulla [34]. The morphology of tau astrogliopathy in these studies, was reminiscent of that reported by Munoz et al. as ATACs [46], although extension of the immunoreactivity into the astrocytic processes was emphasized [34, 35]. Distinct accumulation of TSA in the dentate gyrus of the hippocampus was also recognized [34, 36]. Mathematical modeling of hippocampal tau immunolabeling patterns suggested that some forms of tau astrogliopathy in the elderly involve hippocampal subregions in a different pattern from that of primary tauopathies [44]. Ferrer et al. [21, 38] showed that the biochemical signature of astroglial tau pathology in the elderly in both white and gray matter differed from that of other astrocytic tau pathologies in primary tauopathies. Specifically, astroglial tau pathologies in the white matter and gray matter in aging brains were not consistently detectable using phospho-specific anti-tau antibody Ser262 or conformational tau modifications at amino acids 312 to 322 (MC1), or tau truncated at aspartic acid 421 (tau-C3) [21, 38].

In addition, isolated tufted astrocytes were reported in the occipitotemporal gyrus in an elderly, population-representative cohort [36], and a tauopathy with tufted, thorny, fibrous, and protoplasmic forms of astrocytic pathology was described by Beach et al. [7], in a series of cases with hippocampal sclerosis and also in a community-based study [34]. Sakai et al. [52] reported prominent subcortical white matter astrocytic tau pathology in brains from two elderly patients in whom CBD was considered. In a study on cervical spondylotic myelopathy, AT8 immunohistochemistry revealed tau-positive, neuropil threads, astrocytic foot-like perivascular or subpial structures, and glial cells with short and thick processes, which the authors termed

TSA [58]. Interestingly, prominent tau astrogliopathy may be seen in familial disorders without *MAPT* mutation [20].

In summary, tau-immunoreactive astrogliopathy in the elderly represent a spectrum of morphological abnormalities including those originally described as TSA (plump, perinuclear cytoplasmic immunoreactivity) and additional fine granular immunoreactivity extending into the astrocytic processes in the gray matter. These two morphologies can be present in the same brain. TSA may be seen in the subpial, subependymal, or perivascular areas, and in the white but also in the gray matter, while the fine granular immunoreactivity is seen in the gray matter. Most likely, the different tau-immunoreactive astroglial morphologies in different locations in the aging brain, with or without clinical correlation, reflect different pathogenetic events. We propose the umbrella term ARTAG to encompass all of these, *with or without* accompanying morphological features of other neurodegenerative disorders, including PSP, CBD, PiD, GGT, PART, AD, AGD, and Lewy body pathology. Some clinicopathological studies suggest that ARTAG may present clinically with focal symptoms like aphasia when circumscribed to a smaller number of regions [46], whereas, in cases with widespread pathology dementia with or without parkinsonism might be the clinical presentation [34, 35]; on the other hand, studies focusing only on the MTL have found no relationship between ARTAG and cognitive impairment or dementia [36].

Differential diagnosis

We provide the following operational criteria for the six well-defined tau immunoreactive astrocytic cytopathologies seen in primary tauopathies and ageing brain as follows (see comparison in **Table 1** and **Fig 1**):

- 1) Tufted astrocytes: starlike tufts of tau positive radiating fibers. The abnormal densely tau immunoreactive tufts are detected in the proximal part of the astrocytic processes, often usually in a symmetrical fashion. They are localized to the gray matter (mostly basal ganglia and neocortex).
- 2) Astrocytic plaques: focal and densely tau immunoreactive stubby dilatations of distal processes of astrocytes giving a senile-plaque-like appearance without amyloid core. They are localized to the gray matter (mostly basal ganglia and neocortex).
- 3) Ramified astrocytes: tau immunoreactivity occupying mostly the perikarya and ramifying into the cell processes usually localized to one side of the cell giving the appearance of eccentric nuclei of the astrocyte. They are localized to the gray matter and to the white matter in

neocortices with severe neuronal loss.

4) Globular astroglial inclusions: tau immunoreactive distinct globules (up to the size of the astroglial nucleus; 1-5 μm) and dots (1 μm) in the perikarya and proximal parts of astrocytic processes, found in gray matter. In contrast to tufted astrocytes and astrocytic plaques, GAI are not argyrophilic with Gallyas silver stain.

5) Thorn-shaped astrocytes (TSA): tau immunoreactivity is localized in astrocytic perikarya with extension into the proximal parts of the astrocytic processes, with inclusions also in the astrocytic endfeet at the glia limitans around blood vessels and at the pial surface. The processes are thick and short and reminiscent of thorns. They are localized mostly to the subpial and perivascular locations, but also can be found in white matter and less often as clusters in the gray matter.

6) Granular or fuzzy tau immunoreactivity in processes of astrocytes (GFA): fine granular immunoreactivity of branching processes with a few dilatations of gray matter astrocytes. The perinuclear soma is densely immunoreactive in most of these astrocytes.

The two major cytomorphologies of ARTAG (i.e. TSA and GFA) may accompany tauopathies or other neurodegenerative disorders, but ARTAG should be distinguished from the more specific astrocytic lesions that are characteristic of primary tauopathies. To understand the frequency and relevance of ARTAG it is recommended to document ARTAG as an additional feature in primary tauopathies. It must be noted, that the astroglial tau immunoreactivity described by Botez et al. in the amygdala of AGD [11] fit best with the granular fine or fuzzy tau immunoreactivity in astroglia (GFA) now included as a form of ARTAG. Indeed, astrocytic tau pathology is variably seen in AGD [22]. Therefore, it is helpful to comment whether in a case of AGD additional ARTAG is present. Furthermore, there are other tau-related disorders with astrocytic tau pathology. For instance, astrocytic tau pathology is also a component of CTE [40-42]. CTE is associated with a history of repetitive concussive or subconcussive brain trauma and is characterized by widespread accumulation of hyperphosphorylated tau in NFTs and astrocytes, which have similarity to TSA seen in ARTAG [41]. ARTAG has features that overlap those of CTE, including the accentuation of tau pathology around small cerebral vessels and in subpial and periventricular areas. On the other hand, tau pathology in CTE is more abundant in the depths of the convexity cerebral sulci, especially in early stages [41], an aspect that has not been reported in tau astroglionopathy in the aging brain [29, 34-36, 46, 57]. It is possible that CTE pathology has been considered to be age-related astroglionopathy prior to recognition of ARTAG, especially for lesions in the MTL, which can be severely affected in

more advanced stages of CTE [42]. The characteristic patchy lesions at depths of cerebral sulci were not recognized as a specific morphological feature of CTE in earlier studies. Finally, tufted astrocytes in PSP, astrocytic plaques in CBD, globular astrocytic inclusions in GGT, and ramified astrocytes in PiD are distinct from tau-immunoreactive astrocytes in the gray matter in ARTAG (see **Table 1**).

These observations raise the possibility that ARTAG affects distinct astrocytic populations to those in established primary tauopathies. Ikeda and colleagues noted that the distribution of TSA was coexistent with prominent subpial and subependymal gliosis [29]. Corpora amylacea, which are heavily invested by reactive astrocytes, also share this distribution. Importantly, these astroglial populations of the “glia limitans” share common features with fibrous astrocytes, which predominate in the white matter and subpial zone [8] and with a subset of astrocytes in the gray matter [61], where ARTAG can be also observed. In contrast, astrocytic tau pathologies in CBD or PSP involve protoplasmic astrocytes and are independent of reactive astrogliosis [19, 65]. A few studies report association of glial fibrillary acidic protein and AT8 immunoreactivity in subpial but also in gray matter localization of tau astrogliopathy in elderly brains [35, 36]. Protoplasmic and fibrous astrocytes differ substantially in their glutamate uptake capabilities and capacity and have very different degrees of coupling, which are important with regard to their respective calcium wave signals, resting membrane potentials, potassium buffering, glutamate metabolism, exchange of second messengers, metabolites, and other signaling intermediates between cells [50]. In addition to these differences, reaction of astrocytes varies considerably between distinct diseases of the nervous system [60]. It is these differences that may be of pathogenetic relevance to the morphologic diversity of astrogliopathy in ARTAG.

Evaluation of ARTAG

Inconsistency in assessing, describing and documenting ARTAG has impeded research and limited our understanding of the significance of this pathology. It is not clear whether the different patterns of anatomical involvement represent a continuum or distinct abnormalities with different causes. Most previous studies have focused on the MTL, but more widespread involvement is possible [34, 35]. The relative frequency of ARTAG limited to MTL as opposed

to more widespread tau astroglial pathology remains unclear. Potential etiologies are not known, although defective function of the blood-brain barrier [57], metabolic encephalopathy, neurodegenerative pathologies, hypoperfusion associated with aging, AD, or vascular dementia [39, 64], and even repeated minor trauma with possible genetic risk factors may play a role. Clinical, imaging and neuropathological data related to these aspects need to be documented precisely to allow a better understanding of the pathogenesis of ARTAG [47]. A method is needed to describe morphologies that can be widely accepted and reproducible. As silver impregnation methods are difficult to standardize and immunohistochemistry for ubiquitin and p62 does not demonstrate all forms of tau cytopathology, optimal characterization of ARTAG requires the use of immunohistochemistry for phosphorylated tau. The most widely used phosphorylation-dependent anti-tau antibodies that have allowed characterization of ARTAG to date include: AT8 (pSer202/Thr205; available from different commercial sources), CP13 (Ser202; Peter Davies, Litwin-Zucker Research Center for The Study of Alzheimer's Disease and Memory, Manhasset, NY, USA) and PHF-1 or AD2 (Ser396/Ser404; Peter Davies, NY, USA for PHF-1 or commercial sources for AD2s). Other antibodies that may prove useful in the characterization of ARTAG include those specific for tau phosphorylated at Thr181, Ser202, Ser214, Ser396, Ser422, N-terminus region epitope-specific, 4R tau isoform-specific, and some conformation-dependent antibodies such as Alz50 (but not MC-1) [21, 35, 38].

Recommendations for sampling and staining are as follows:

- Preliminary screening for ARTAG should include tau-immunohistochemistry (antibodies AT8, CP13, AD2 or PHF-1 are recommended) on two sections representative of the MTL (i.e., amygdala and hippocampus at the level of the lateral geniculate body). These regions are vulnerable to TSA and GFA.
- If tau astroglial pathology is noted in the screening section, a systematic characterization of ARTAG will require analysis involving additional areas of the frontal, parietal, lateral temporal, and occipital cortices, as well as anterior and posterior portion of the basal ganglia, thalamus, midbrain at the level of substantia nigra, pons at the level of locus coeruleus, and medulla oblongata.
- In cases where focal cortical symptoms are reported, further cortical areas corresponding to the clinical symptoms or signal alterations detected in MRI should also be evaluated.

ARTAG should be considered when detecting either or both of the two cytomorphologies: TSA or GFA. Following this, we propose the following four-step characterization TReSS algorithm (**Table 2**):

Type? Regional involvement? Severity? Subregional involvement?

- First: Identify the morphologic and distribution types of ARTAG based on parenchymal localization of TSA and GFA (note that combination of these types is generally the rule):
 - 1) Laminar subpial TSA (**Fig 2a**), or
 - 2) Subependymal TSA: plump perinuclear cytoplasmic tau immunoreactivity in astrocytes in subpial or subependymal locations (**Fig 2b**). If subpial, it is important to note whether this is more pronounced in the sulcal depths in the convexity cerebral cortices, as in CTE.
 - 3) Gray matter GFA (or rarely TSA): solitary (one or two/20x field) (**Fig 2c, e**) or clustered (**Fig 2d, f**) GFA (three or more/20x field) with fine granular immunopositivity of the cytoplasmic processes (GFA), with plump perinuclear cytoplasmic tau immunoreactivity.
 - 4) White matter TSA: astrocytes in the subcortical white matter that show plump cytoplasmic immunoreactivity (**Fig 2g**). Note that in the white matter these usually form small clusters (>3 astrocytes) and that it may extend into the adjacent gray matter as described for ATAC [46].
 - 5) Perivascular TSA: plump cytoplasmic immunoreactivity with tau-immunoreactive astrocytic processes around vessels (**Fig 2h**) in the gray or white matter.

[It must be noted that tau immunohistochemistry occasionally decorates astrocytes at the border of chronic traumatic or vascular lesions in young and aged individuals. Therefore, this lesion-associated tau astrogliaopathy is important to document, but is not considered an aging related astrogliaopathy.]

- Second: Identify involvement of gross anatomical regions:
 - A) MTL
 - B) Lobar
 - C) Subcortical
 - D) Brainstem

[Although the most frequently involved region is the MTL, involvement of further regions should be recognized. Moreover, MTL is important for comparison with neuroimaging data on MTL atrophy.]

- Third: Document the severity of ARTAG pathology in the region or subregion (see step four) examined. ARTAG may appear in focal clusters or in a widespread distribution. We propose documentation as to whether ARTAG involves only 1) occasional or 2) numerous astrocytes. If the latter, the focal clusters or widespread distribution should be noted. Semiquantitative scoring for ARTAG will need to be refined.

- Fourth: Map subregional involvement to promote future exploration and scientific discovery related to ARTAG. These are the subdivisions within the gross anatomical regions of the second step and include the following (**Table 3**; examples are shown in **Fig 3**):
 - amygdala and hippocampus, inferior temporal gyrus for MTL
 - frontal, parietal, occipital, lateral temporal (middle and superior gyrus) for lobar
 - caudate nucleus, putamen, nucleus accumbens, globus pallidus, thalamus, basal forebrain for subcortical
 - mesencephalon, pons, medulla oblongata for brainstem

It should be noted whether ARTAG is associated with features of a particular neurodegenerative disorder, or with other disease (cerebrovascular, inflammatory, metabolic, etc.) followed by the description of the type and major regional involvement.

Some examples for the diagnostic reporting are provided as follows:

1) Examples for pure types:

a. ARTAG subpial type;

Region: MTL;

Subregion: hippocampus, inferior temporal cortex;

Extent: numerous astrocytes and widespread distribution

b. ARTAG subependymal type;

Region: Subcortical;

Subregion: lateral ventricle;

Extent: occasional

2) Examples for combinations:

a. ARTAG gray matter type;

Region: MTL and subcortical;

Subregion: inferior temporal cortex and nucleus accumbens;

Extent: numerous astrocytes in focal clusters

plus

ARTAG white matter type;

Region: MTL;

Subregion: hippocampus, periamygdala white matter, and temporal;

Extent: numerous astrocytes with widespread distribution;

b. ARTAG perivascular type;

Region: lobar and subcortical;

Subregion: frontal cortex and striatum;

Extent: occasional

plus

ARTAG white matter type;

Region: MTL and lobar;

Subregion: lateral temporal, frontal, and parietal lobes;

Extent: numerous astrocytes in focal clusters.

For example, the cases described by Munoz et al. [46] would be summarized in the diagnostic report as: ARTAG gray and white matter type; region: MTL and lobar; extent: numerous in focal clusters. For research purposes, the subregional involvement should be added as: gyrus ambiens, parahippocampal gyrus, fusiform gyrus, inferior, middle, and superior temporal gyri, frontal dorsolateral and orbitofrontal cortices, cingulate gyrus, and inferior parietal lobe. The cases described by Kovacs et al. [35] could be summarized as ARTAG gray matter type; region: MTL, lobar, subcortical, and brainstem; extent: numerous in focal clusters; *plus* white matter type; region: MTL; extent: numerous and widespread; with further details on the subregional involvement. The cases discussed by Santpere and Ferrer [53] as early PSP-like astrocytic changes also represent ARTAG with additional features of concomitant PSP-type pathology (i.e. cases 4 and 5).

Summary

ARTAG describes a spectrum of astroglial tau pathologies detected mainly in the elderly represented by TSA and GFA, which are distinct from astroglial lesions of primary tauopathies (i.e. tufted astrocytes, astrocytic plaques, ramified astrocytes or globular astroglial inclusions). The frequency of ARTAG varies depending on the type: subpial, subependymal, and perivascular types are more frequent, while gray matter and cerebral white matter types might be less common. The etiology of different types might be different; however, all appear mostly associated with aging. Although documented in several publications, there is a lack of consensus on how ARTAG should be recorded and interpreted. Here we propose steps for a systematic characterization with the expectation that this will improve communication about and understanding of this condition, including its relation to other brain pathologies and clinical symptoms. This approach has the potential to help in several respects:

- 1) It will facilitate communication between neuropathologists and researchers. Revisiting and standardizing the terminology should help to move the field forward. It will also increase awareness of this pathology, which is under-recognized and under-studied.
- 2) A better differentiation of ARTAG types may help with assessing their relationship to other tauopathies. This may be particularly important in the context of CTE-related tau pathologies. Furthermore, this may help better understanding of differences in the pathogenesis of ARTAG types.
- 3) A regular system for typing and grading of ARTAG should facilitate comparisons between different centers, and the pooling of information in harmonized clinicopathological studies. These will potentially pave the way towards mechanistic insights and genetic studies into their pathogenesis.
- 4) Understanding the nature of ARTAG may help in the interpretation of clinical biomarker and imaging studies.

Development of such a common concept (**Fig 4**) and nomenclature that allows comparisons across studies and aggregation of data for large scale multi-institutional analyses is imperative in order to understand the phenomena and clinical implications of ARTAG. Future studies should also aim to re-evaluate these observations to validate this approach and to develop a concise classification of ARTAG for diagnostic neuropathology. To reach this goal, paradigms

will be need to be designed for ARTAG along the lines used to standardize evaluation and diagnostic criteria for tau, amyloid and α -synuclein and other major pathologies [2-4, 10, 45]. Subsequently, it will be possible to evaluate inter-rater reliability of the proposed evaluation and to eventually merge clinical and pathologic data from multiple centers to determine practical significance of ARTAG. At this preliminary stage, however, we limit recommendation to a common nomenclature and classification of aging related astroglipathy and an evaluation strategy.

Acknowledgements

We are extremely grateful to the patients, clinicians, and fellow researchers that made this effort possible. We also acknowledge the following funding sources: FP7 EU Project Develage No. 278486 (GGK); Grant "NIH P30 AG10133 (BG); NIA grants P50 AG05681, P01 AG03991 (NJC), NIH R01 AG040311, institutional grants NIH P01 AG019724-03 and P50 AG023501, and the tau consortium (LTG); the Nelson Family Foundation and NIH grants AG010124, AG017586 (MEM); NIH grant P50 AG005138 (PRH); Alzheimer's Research UK (ARUK), Alzheimer's Society, National Institute for Health Research (NIHR), and UK Medical Research Council (MRC; G0400074)(JA); GMH is a National Health and Medical Research Council of Australia Senior Principal Research Fellow (#630434); grant IGA NT12094-5 from Grant Agency of Ministry of Health of Czech Republic (RM); NIH grant # AG028383 (PN); UK Medical Research Council (MRC; MR/L016400/1) (CS); NIA P50 AG005133 (J.K); National Institute of Neurological Disorders and Stroke (1U01NS086659-01), Department of Veterans Affairs,), the National Institute of Aging Boston University Alzheimer's Disease Center (P30AG13846; supplement 0572063345-5) (ACM); UK Medical Research Council (MC-PC-13044) (JWI and CS); National Brain Research Program, Hungary (KTIA_13_NAP-A-II/7) and Grant-in-Aid (KAKEN 26250017)(HT); NIH grant P30AG12300 (KH, CLW); DFG Grant (SFB 1134/A03)(CS); Johns Hopkins Alzheimer's Disease Research Center NIH grant #P50AG05146 (JCT); Alzheimer's Disease Core Center grant P30AG008051-26 (TW); Grant AG13854 (EHB); JSPS KAKENHI Grant Number 26430060 (MT); Italian Ministry of Health (GG and FT); National Institute of Health grants P50 AG05136 and P50 NS062684 (TJM); National Brain Research Program, Hungary (KTIA_13_NAP-A-II/7 for TH). The help of Brain Banks in collecting tissue is also highly acknowledged: Vienna KIN-Neurobiobank and VITA-study (GGK); GIE NeuroCEB (funded by the patients associations France Alzheimer, France Parkinson, Fondation ARSEP and CSC)(CD); Sydney Brain Bank (funded by Neuroscience Research Australia and the University of New South Wales)(GMH); the Sheffield and Cambridge Brain Banks (CFAS)(PI, SW); Parkinson's UK Tissue Bank at Imperial College, funded by Parkinson's UK, a charity registered in England and Wales (948776) and Scotland (SC037554)(SG); The Edinburgh Brain Bank is supported by the UK Medical Research Council (MR/L016400/1)(CS, JWI).

References

- 1 Ahmed Z, Bigio EH, Budka H, Dickson DW, Ferrer I, Ghetti B, Giaccone G, Hatanpaa KJ, Holton JL, Josephs KA et al (2013) Globular glial tauopathies (GGT): consensus recommendations. *Acta Neuropathol* 126: 537-544
- 2 Alafuzoff I, Arzberger T, Al-Sarraj S, Bodi I, Bogdanovic N, Braak H, Bugiani O, Del-Tredici K, Ferrer I, Gelpi E et al (2008) Staging of neurofibrillary pathology in Alzheimer's disease: a study of the BrainNet Europe Consortium. *Brain Pathol* 18: 484-496
- 3 Alafuzoff I, Ince PG, Arzberger T, Al-Sarraj S, Bell J, Bodi I, Bogdanovic N, Bugiani O, Ferrer I, Gelpi E et al (2009) Staging/typing of Lewy body related alpha-synuclein pathology: a study of the BrainNet Europe Consortium. *Acta Neuropathol* 117: 635-652
- 4 Alafuzoff I, Thal DR, Arzberger T, Bogdanovic N, Al-Sarraj S, Bodi I, Boluda S, Bugiani O, Duyckaerts C, Gelpi E et al (2009) Assessment of beta-amyloid deposits in human brain: a study of the BrainNet Europe Consortium. *Acta Neuropathol* 117: 309-320
- 5 Arima K, Izumiyama Y, Nakamura M, Nakayama H, Kimura M, Ando S, Ikeda K, Takahashi K (1998) Argyrophilic tau-positive twisted and non-twisted tubules in astrocytic processes in brains of Alzheimer-type dementia: an electron microscopical study. *Acta Neuropathol* 95: 28-39
- 6 Bancher C, Brunner C, Lassmann H, Budka H, Jellinger K, Wiche G, Seitelberger F, Grundke-Iqbal I, Iqbal K, Wisniewski HM (1989) Accumulation of abnormally phosphorylated tau precedes the formation of neurofibrillary tangles in Alzheimer's disease. *Brain Res* 477: 90-99
- 7 Beach TG, Sue L, Scott S, Layne K, Newell A, Walker D, Baker M, Sahara N, Yen SH, Hutton M et al (2003) Hippocampal sclerosis dementia with tauopathy. *Brain Pathol* 13: 263-278
- 8 Beach TG, Walker R, McGeer EG (1989) Patterns of gliosis in Alzheimer's disease and aging cerebrum. *Glia* 2: 420-436
- 9 Bigio EH, Mishra M, Hatanpaa KJ, White CL, 3rd, Johnson N, Rademaker A, Weitner BB, Deng HX, Dubner SD, Weintraub S et al (2010) TDP-43 pathology in primary progressive aphasia and frontotemporal dementia with pathologic Alzheimer disease. *Acta Neuropathol* 120: 43-54
- 10 Boluda S, Toledo JB, Irwin DJ, Raible KM, Byrne MD, Lee EB, Lee VM, Trojanowski JQ (2014) A comparison of Abeta amyloid pathology staging systems and correlation with clinical diagnosis. *Acta Neuropathol* 128: 543-550
- 11 Botez G, Probst A, Ipsen S, Tolnay M (1999) Astrocytes expressing hyperphosphorylated tau protein without glial fibrillary tangles in argyrophilic grain disease. *Acta Neuropathol* 98: 251-256
- 12 Cairns NJ, Bigio EH, Mackenzie IR, Neumann M, Lee VM, Hatanpaa KJ, White CL, 3rd, Schneider JA, Grinberg LT, Halliday G et al (2007) Neuropathologic diagnostic and nosologic criteria for frontotemporal lobar degeneration: consensus of the Consortium for Frontotemporal Lobar Degeneration. *Acta Neuropathol* 114: 5-22
- 13 Chin SS, Goldman JE (1996) Glial inclusions in CNS degenerative diseases. *J Neuropathol Exp Neurol* 55: 499-508
- 14 Crary JF, Trojanowski JQ, Schneider JA, Abisambra JF, Abner EL, Alafuzoff I, Arnold SE, Attems J, Beach TG, Bigio EH et al (2014) Primary age-related tauopathy (PART): a common pathology associated with human aging. *Acta Neuropathol* 128: 755-766
- 15 Dickson DW, Kouri N, Murray ME, Josephs KA (2011) Neuropathology of frontotemporal lobar degeneration-tau (FTLD-tau). *J Mol Neurosci* 45: 384-389
- 16 Dugger B, Uchikado H, Ahmed Z, Dickson DW (2008) Sex differences in perivascular tauopathy in the mediobasal tuberal hypothalamus in neurodegenerative diseases in humans. *Alzheimers Dementia* 4 Suppl.: T715
- 17 Dugger BN, Hentz JG, Adler CH, Sabbagh MN, Shill HA, Jacobson S, Caviness JN, Belden C, Driver-Dunckley E, Davis KJ et al (2014) Clinicopathological outcomes of prospectively followed normal elderly brain bank volunteers. *J Neuropathol Exp Neurol* 73: 244-252

- 18 Evidente VG, Adler CH, Sabbagh MN, Connor DJ, Hentz JG, Caviness JN, Sue LI, Beach TG (2011) Neuropathological findings of PSP in the elderly without clinical PSP: possible incidental PSP? *Parkinsonism Relat Disord* 17: 365-371
- 19 Feany MB, Dickson DW (1995) Widespread cytoskeletal pathology characterizes corticobasal degeneration. *Am J Pathol* 146: 1388-1396
- 20 Ferrer I, Legati A, Garcia-Monco JC, Gomez-Beldarrain M, Carmona M, Blanco R, Seeley WW, Coppola G (2015) Familial behavioral variant frontotemporal dementia associated with astrocyte-predominant tauopathy. *J Neuropathol Exp Neurol* 74: 370-379
- 21 Ferrer I, Lopez-Gonzalez I, Carmona M, Arregui L, Dalfo E, Torrejon-Escribano B, Diehl R, Kovacs GG (2014) Glial and neuronal tau pathology in tauopathies: characterization of disease-specific phenotypes and tau pathology progression. *J Neuropathol Exp Neurol* 73: 81-97
- 22 Ferrer I, Santpere G, van Leeuwen FW (2008) Argyrophilic grain disease. *Brain* 131: 1416-1432
- 23 Ghetti B, Oblak AL, Boeve BF, Johnson KA, Dickerson BC, Goedert M (2015) Invited review: Frontotemporal dementia caused by microtubule-associated protein tau gene (MAPT) mutations: a chameleon for neuropathology and neuroimaging. *Neuropathol Appl Neurobiol* 41: 24-46
- 24 Goedert M, Spillantini MG, Jakes R, Rutherford D, Crowther RA (1989) Multiple isoforms of human microtubule-associated protein tau: sequences and localization in neurofibrillary tangles of Alzheimer's disease. *Neuron* 3: 519-526
- 25 Hishikawa N, Hashizume Y, Yoshida M, Niwa J, Tanaka F, Sobue G (2005) Tuft-shaped astrocytes in Lewy body disease. *Acta Neuropathol* 109: 373-380
- 26 Hutton M, Lendon CL, Rizzu P, Baker M, Froelich S, Houlden H, Pickering-Brown S, Chakraverty S, Isaacs A, Grover A et al (1998) Association of missense and 5'-splice-site mutations in tau with the inherited dementia FTDP-17. *Nature* 393: 702-705
- 27 Ikeda K (1996) Glial fibrillary tangles and argyrophilic threads: Classification and disease specificity. *Neuropathology* 16: 71-77
- 28 Ikeda K, Akiyama H, Arai T, Nishimura T (1998) Glial tau pathology in neurodegenerative diseases: their nature and comparison with neuronal tangles. *Neurobiol Aging* 19: S85-91
- 29 Ikeda K, Akiyama H, Kondo H, Haga C, Tanno E, Tokuda T, Ikeda S (1995) Thorn-shaped astrocytes: possibly secondarily induced tau-positive glial fibrillary tangles. *Acta Neuropathol* 90: 620-625
- 30 Irwin DJ, Cairns NJ, Grossman M, McMillan CT, Lee EB, Van Deerlin VM, Lee VM, Trojanowski JQ (2015) Frontotemporal lobar degeneration: defining phenotypic diversity through personalized medicine. *Acta Neuropathol* 129: 469-491
- 31 Jellinger KA, Attems J (2007) Neurofibrillary tangle-predominant dementia: comparison with classical Alzheimer disease. *Acta Neuropathol* 113: 107-117
- 32 Kovacs GG (2015) Invited review: Neuropathology of tauopathies: principles and practice. *Neuropathol Appl Neurobiol* 41: 3-23
- 33 Kovacs GG, Horvath MC, Majtenyi K, Lutz MI, Hurd YL, Keller E (2015) Heroin abuse exaggerates age-related deposition of hyperphosphorylated tau and p62-positive inclusions. *Neurobiol Aging* 36: 3100-3107
- 34 Kovacs GG, Milenkovic I, Wohrer A, Hofberger R, Gelpi E, Haberler C, Honigschnabl S, Reiner-Concin A, Heinzl H, Jungwirth S et al (2013) Non-Alzheimer neurodegenerative pathologies and their combinations are more frequent than commonly believed in the elderly brain: a community-based autopsy series. *Acta Neuropathol* 126: 365-384
- 35 Kovacs GG, Molnar K, Laszlo L, Strobel T, Botond G, Honigschnabl S, Reiner-Concin A, Palkovits M, Fischer P, Budka H (2011) A peculiar constellation of tau pathology defines a subset of dementia in the elderly. *Acta Neuropathol* 122: 205-222
- 36 Lace G, Ince PG, Brayne C, Savva GM, Matthews FE, de Silva R, Simpson JE, Wharton SB (2012) Mesial temporal astrocyte tau pathology in the MRC-CFAS ageing brain cohort. *Dement Geriatr Cogn Disord* 34: 15-24

- 37 Lee VM, Goedert M, Trojanowski JQ (2001) Neurodegenerative tauopathies. *Annu Rev Neurosci* 24: 1121-1159
- 38 Lopez-Gonzalez I, Carmona M, Blanco R, Luna-Munoz J, Martinez-Mandonado A, Mena R, Ferrer I (2013) Characterization of thorn-shaped astrocytes in white matter of temporal lobe in Alzheimer's disease brains. *Brain Pathol* 23: 144-153
- 39 Love S, Miners JS (2015) White matter hypoperfusion and damage in dementia: post-mortem assessment. *Brain Pathol* 25: 99-107
- 40 McKee AC, Cantu RC, Nowinski CJ, Hedley-Whyte ET, Gavett BE, Budson AE, Santini VE, Lee HS, Kubilus CA, Stern RA (2009) Chronic traumatic encephalopathy in athletes: progressive tauopathy after repetitive head injury. *J Neuropathol Exp Neurol* 68: 709-735
- 41 McKee AC, Stein TD, Kiernan PT, Alvarez VE (2015) The neuropathology of chronic traumatic encephalopathy. *Brain Pathol* 25: 350-364
- 42 McKee AC, Stern RA, Nowinski CJ, Stein TD, Alvarez VE, Daneshvar DH, Lee HS, Wojtowicz SM, Hall G, Baugh CM et al (2013) The spectrum of disease in chronic traumatic encephalopathy. *Brain* 136: 43-64
- 43 Mesulam M, Wicklund A, Johnson N, Rogalski E, Leger GC, Rademaker A, Weintraub S, Bigio EH (2008) Alzheimer and frontotemporal pathology in subsets of primary progressive aphasia. *Ann Neurol* 63: 709-719
- 44 Milenkovic I, Petrov T, Kovacs GG (2014) Patterns of hippocampal tau pathology differentiate neurodegenerative dementias. *Dement Geriatr Cogn Disord* 38: 375-388
- 45 Montine TJ, Monsell SE, Beach TG, Bigio EH, Yunqi B, Cairns NJ, Frosch M, Henriksen J, Julia K, Kukull WA et al (2015) Multisite assessment of NIA-AA guidelines for the neuropathologic evaluation of Alzheimer's disease. *Alzheimers Dementia* doi: 10.1016/j.jalz.2015.07.492.
- 46 Munoz DG, Woulfe J, Kertesz A (2007) Argyrophilic thorny astrocyte clusters in association with Alzheimer's disease pathology in possible primary progressive aphasia. *Acta Neuropathol* 114: 347-357
- 47 Murray ME, Kouri N, Lin WL, Jack CR, Jr., Dickson DW, Vemuri P (2014) Clinicopathologic assessment and imaging of tauopathies in neurodegenerative dementias. *Alz Res Ther* 6: 1
- 48 Nakano I, Iwatsubo T, Otsuka N, Kamei M, Matsumura K, Mannen T (1992) Paired helical filaments in astrocytes: electron microscopy and immunohistochemistry in a case of atypical Alzheimer's disease. *Acta Neuropathol* 83: 228-232
- 49 Nishimura M, Namba Y, Ikeda K, Oda M (1992) Glial fibrillary tangles with straight tubules in the brains of patients with progressive supranuclear palsy. *Neurosci Lett* 143: 35-38
- 50 Oberheim NA, Goldman SA, Nedergaard M (2012) Heterogeneity of astrocytic form and function. *Methods Mol Biol* 814: 23-45
- 51 Poorkaj P, Bird TD, Wijsman E, Nemens E, Garruto RM, Anderson L, Andreadis A, Wiederholt WC, Raskind M, Schellenberg GD (1998) Tau is a candidate gene for chromosome 17 frontotemporal dementia. *Ann Neurol* 43: 815-825
- 52 Sakai K, Piao YS, Kikugawa K, Ohara S, Hasegawa M, Takano H, Fukase M, Nishizawa M, Kakita A, Takahashi H (2006) Corticobasal degeneration with focal, massive tau accumulation in the subcortical white matter astrocytes. *Acta Neuropathol* 112: 341-348
- 53 Santpere G, Ferrer I (2009) Delineation of early changes in cases with progressive supranuclear palsy-like pathology. Astrocytes in striatum are primary targets of tau phosphorylation and GFAP oxidation. *Brain Pathol* 19: 177-187
- 54 Schultz C, Braak H, Braak E (1996) A sex difference in neurodegeneration of the human hypothalamus. *Neurosci Lett* 212: 103-106
- 55 Schultz C, Deghani F, Hubbard GB, Thal DR, Struckhoff G, Braak E, Braak H (2000) Filamentous tau pathology in nerve cells, astrocytes, and oligodendrocytes of aged baboons. *J Neuropathol Exp Neurol* 59: 39-52

- 56 Schultz C, Ghebremedhin E, Braak E, Braak H (1999) Sex-dependent cytoskeletal changes of the human hypothalamus develop independently of Alzheimer's disease. *Exp Neurol* 160: 186-193
- 57 Schultz C, Ghebremedhin E, Del Tredici K, Rub U, Braak H (2004) High prevalence of thorn-shaped astrocytes in the aged human medial temporal lobe. *Neurobiol Aging* 25: 397-405
- 58 Shimizu H, Kakita A, Takahashi H (2008) Spinal cord tau pathology in cervical spondylotic myelopathy. *Acta Neuropathol* 115: 185-192
- 59 Simpson JE, Ince PG, Lace G, Forster G, Shaw PJ, Matthews F, Savva G, Brayne C, Wharton SB, Function MRCC et al (2010) Astrocyte phenotype in relation to Alzheimer-type pathology in the ageing brain. *Neurobiol Aging* 31: 578-590
- 60 Sofroniew MV, Vinters HV (2010) Astrocytes: biology and pathology. *Acta Neuropathol* 119: 7-35
- 61 Sosunov AA, Wu X, Tsankova NM, Guilfoyle E, McKhann GM, 2nd, Goldman JE (2014) Phenotypic heterogeneity and plasticity of isocortical and hippocampal astrocytes in the human brain. *J Neurosci* 34: 2285-2298
- 62 Spillantini MG, Goedert M, Crowther RA, Murrell JR, Farlow MR, Ghetti B (1997) Familial multiple system tauopathy with presenile dementia: a disease with abundant neuronal and glial tau filaments. *Proc Natl Acad Sci U S A* 94: 4113-4118
- 63 Spillantini MG, Murrell JR, Goedert M, Farlow MR, Klug A, Ghetti B (1998) Mutation in the tau gene in familial multiple system tauopathy with presenile dementia. *Proc Natl Acad Sci U S A* 95: 7737-7741
- 64 Thomas T, Miners S, Love S (2015) Post-mortem assessment of hypoperfusion of cerebral cortex in Alzheimer's disease and vascular dementia. *Brain* 138: 1059-1069
- 65 Togo T, Dickson DW (2002) Tau accumulation in astrocytes in progressive supranuclear palsy is a degenerative rather than a reactive process. *Acta Neuropathol* 104: 398-402
- 66 Uchikado H, Fujino Y, Lin W, Dickson D (2008) Frequency and Relation of Argyrophilic Grain Disease and Thorn-Shaped Astrocytes in Alzheimer's Disease. *Advances in Alzheimer's and Parkinson's Disease: Insights, Progress, and Perspectives*. New York, pp 375-379
- 67 Wakabayashi K, Shibasaki Y, Hasegawa M, Horikawa Y, Soma Y, Hayashi S, Morita T, Iwatsubo T, Takahashi H (2000) Primary progressive aphasia with focal glial tauopathy. *Neuropathol Appl Neurobiol* 26: 477-481
- 68 Yamazaki M, Nakano I, Imazu O, Terashi A (1995) Paired helical filaments and straight tubules in astrocytes: an electron microscopic study in dementia of the Alzheimer type. *Acta Neuropathol* 90: 31-36

Table 1. Operational criteria for the six well-defined tau immunoreactive astrocytic cytopathologies seen in primary tauopathies and aging-related tau-astrogliopathy (ARTAG). GM: gray matter; WM: white matter; SP: subpial; SE: subependymal; PV: perivascular. In the column "Reported" those diseases are mentioned, where there are literature reports of the specific astrocytic tau pathologies. PSP: progressive supranuclear palsy; CBD: corticobasal

degeneration; PID: Pick disease; GGT: globular glial tauopathy; AGD: argyrophilic grain disease.

Name	Characteristic immunoreactivity							
	Soma	Astrocytic processes			Gallyas	Reported	Location	ARTAG
		Proximal	Distal					
Tufted astrocytes	relatively spared	dense tufts: usually symmetric	no	+	PSP	GM	no	
Astrocytic plaques	spared	no	stubby dilatations	+	CBD	GM	no	
Ramified astrocytes	dense: localized to one side	dense ramifications: usually asymmetric	no	+	PID	GM, WM/GM junction	no	
Globular astroglial inclusions	dense: filled with globules	1-5 μ m globules	1-2 μ m globules	-	GGT	GM	no	
Thorn-shaped astrocytes (TSA)	dense: thorn or flame shaped	short and thick	no	+/-	<i>with or without</i> PSP, AGD, AD, CTE, other	GM, WM, SP, SE, PV	yes	
Granular or fuzzy astrocytes (GFA)	dense: perinuclear accumulation	fine or fuzzy granular	fine granular	-	<i>with or without</i> AGD	GM	yes	

Table 2. Evaluation of aging-related tau astroglial pathology (ARTAG). TSA: thorn-shaped astrocyte; GFA: granular /fuzzy astrocyte; AP: astrocytic plaque; TA: tufted astrocyte; RA: ramified astrocyte; GAI: globular astroglial inclusions.

Requires:

Presence of thorn-shaped astrocytes (TSA) and/or solitary or clustered astrocytes with plump cytoplasmic tau immunoreactivity that extend into the astroglial processes as fine granular immunopositivity (GFA) distinguishable from AP, TA, RA, and GAI.

Four-step characterization of ARTAG:**Step 1: Distinguish types according to the location:**

- Subpial
- Subependymal
- Gray matter
- White matter
- Perivascular

Step 2: Describe major anatomical distribution

- Medial temporal lobe
- Lobar
- Subcortical
- Brainstem

Step 3: Document the severity of ARTAG

- Occasional
- Numerous
 - Focal
 - Widespread

Step 4: Map subregional involvement and extent (see Table 2)**Ancillary studies:**

- Description of additional tau pathologies in specific anatomical regions:
 - Neurofibrillary degeneration
 - Neuropil threads
 - Diffuse cytoplasmic neuronal tau immunoreactivity (“pretangles”)
 - Dystrophic neurites around or within amyloid plaques
 - Oligodendroglial tau immunoreactivity
 - Characterization of tau phosphorylation, conformation, truncation, nitration, ubiquitination, immunohistochemistry for 4R and 3R tau; ultrastructural study; genetic studies (*MAPT* and other genes associated with neurodegeneration)
 - Description of concomitant neurodegenerative and non-neurodegenerative pathologies
 - Description of relation to lesions (“perilesional” astrocytic tau immunoreactivity), to corpora amylacea, and Rosenthal fibers
-

Table 3. Description of aging-related tau astrogliopathy (ARTAG) based on the type and distribution of astrocytic tau pathology.

MTL: medial temporal lobe; Gy: gyrus; GP: globus pallidus; Caud/Put: Caudate and Putamen; Dent Gyr: dentate gyrus; Medulla obl.: medulla oblongata; Aq: Aqueduct; LV: lateral ventricle; 3V: 3rd Ventricle. *: in the case of focal cortical symptoms the anatomical area with clinical relevance should be noted additionally. Combinations of subtypes should be expected and described.

Diagnostic screening				Clinicopathological correlation and studies on pathogenesis (research)						
STEP 1		STEP 2		STEP 3		STEP 4				
Type	→	Major anatomical involvement	→	Severity	Detailed regional distribution and extent of ARTAG					
Subpial	→	MTL	→	occasional or numerous (focal/widespread)	→	<input type="checkbox"/>	Inf. Temporal Gy	Hippocampus	Amygdala	
		Lobar	→		→	<input type="checkbox"/>	Frontal	Parietal	Occipital	Lateral temporal
		Subcortical	→		→	<input type="checkbox"/>	Basal forebrain	-	-	
		Brainstem	→		→	<input type="checkbox"/>	Mesencephalon	Pons	Medulla obl.	
Subependymal	→	MTL	→		→	<input type="checkbox"/>	Inferior horn			
		Lobar	→		→	<input type="checkbox"/>	LV: Frontal horn	LV:Posterior horn		
		Subcortical	→		→	<input type="checkbox"/>	LV: Caud	3V: Thalamus		
		Brainstem	→		→	<input type="checkbox"/>	Aq/Mesencephalon	Aq/Pons	Aq/Medulla obl.	
Gray matter	→	MTL	→		→	<input type="checkbox"/>	Inf. Temporal Gy	Hippocampus	Amygdala	
		Lobar*	→		→	<input type="checkbox"/>	Frontal	Parietal	Occipital	Lateral temporal
		Subcortical	→		→	<input type="checkbox"/>	Accumbens	Caud/Put	GP	Basal forebrain
		Brainstem	→		→	<input type="checkbox"/>	Substantia nigra	Pons	Medulla obl.	
White matter	→	MTL	→		→	<input type="checkbox"/>	Inf. Temporal Gy	Hippocampus	Amygdala	
		Lobar*	→		→	<input type="checkbox"/>	Frontal	Parietal	Occipital	Lateral temporal
		Subcortical	→		→	<input type="checkbox"/>	Internal capsule	Subinsular	Pencil fibers	
		Brainstem	→		→	<input type="checkbox"/>	Cerebral pedunculi	Pyramids	Midline	
Perivascular	→	MTL	→	→	<input type="checkbox"/>	Inf. Temporal Gy	Hippocampus	Amygdala		
		Lobar	→	→	<input type="checkbox"/>	Frontal	Parietal	Occipital	Lateral temporal	
		Subcortical	→	→	<input type="checkbox"/>	Accumbens	Caud/Put	GP	Basal forebrain	
		Brainstem	→	→	<input type="checkbox"/>	Mesencephalon	Pons	Medulla obl.		

Figure legends

Figure 1. Comparison of tau (here using AT8 antibody) immunoreactivities (IRs) seen in primary tauopathies with those observed in aging-related tau astrogliopathy (ARTAG).

Figure 2. Representative photomicrographs of ARTAG types.

Plump cytoplasmic tau immunoreactivity of astrocytes and tau-positive lining in subpial (**a**) and subependymal (**b**) location. Gray matter (GM)-type: Single-appearing (**c, d**) or clusters (**e, f**) of astrocytes with fine granular tau immunoreactivity in the processes without (**c**) or with (**d**) plump perinuclear cytoplasmic tau immunoreactivity. White matter (WM)-type: Astrocytes in the subcortical white matter with plump cytoplasmic immunoreactivity (**g**). Perivascular-type: Tau-immunoreactive astrocytic processes arranged around vessels (**h**).

The bar shown in “a” represents 30 μm for a, b, d; 50 μm for c, f, g; and 100 μm for e, h.

Figure 3. Representative images of different anatomical regions showing ARTAG.

a: Temporal cortex and white matter (WM); **b:** Dentate gyrus (Gray matter-type cluster enlarged in the right); **c:** Amygdala; **d:** Frontal cortex (Gray matter-type single); **e:** Accumbens nucleus (Gray matter-type clusters and single forms); **f:** Substantia nigra; **g** and **h:** medulla oblongata (IO: inferior olive; ML: medial lemniscus; n. XII: hypoglossal nucleus).

The bar shown in “a” represents 150 μm for a, b; 100 μm for the right inset in b, and c-h.

Figure 4. Summary of the concept of ARTAG.

Four distinct astroglial tau pathologies are seen in primary tauopathies: tufted astrocytes (TA), astrocytic plaques (AP), ramified astrocytes (RA) and globular astroglial inclusions (GAI). ARTAG is characterized by two different morphologies: thorn-shaped astrocytes (TSA) and fine granular immunoreactivity in astrocytic processes (granular/fuzzy astrocytes: GFA); these are seen in the subpial (SP), subependymal (SE), perivascular (PV) areas, and in the white (WM) and gray matter (GM). TSA and GFA may be present in the same brain together. Other neurodegenerative diseases (NDDs) may co-exist with ARTAG or with primary tauopathies.