THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Sibling competition does not exacerbate inbreeding depression in the burying beetle *Nicrophorus vespilloides*

Citation for published version:

Pilakouta, N, Sieber, D & Smiseth, PT 2016, 'Sibling competition does not exacerbate inbreeding depression in the burying beetle *Nicrophorus vespilloides*', *Journal of Evolutionary Biology*, vol. 29, no. 4, pp. 704–710. <https://doi.org/10.1111/jeb.12816>

Digital Object Identifier (DOI):

[10.1111/jeb.12816](https://doi.org/10.1111/jeb.12816)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Evolutionary Biology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

**Sibling competition does not exacerbate inbreeding depression in the
burying beetle *Nicrophorus vespilloides***

Natalie Pilakouta¹, Daniel J. Sieber, and Per T. Smiseth

Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh,
Edinburgh, UK

¹Corresponding author (e-mail: n.pilakouta@gmail.com, telephone: +44(0)1316505462)

Running title: *Inbreeding depression and sibling competition*

Word count:

Main text (excluding references): 3,052 words; Abstract: 250 words; 34 references, 2 tables,
1 figure, 1 supplementary table

Abstract

Inbreeding results from matings between relatives and can cause a reduction in offspring fitness, known as inbreeding depression. Previous work has shown that a wide range of environmental stresses, such as extreme temperatures, starvation, and parasitism, can exacerbate inbreeding depression. It has recently been argued that stresses due to intraspecific competition should have a stronger effect on the severity of inbreeding depression than stresses due to harsh physical conditions. Here, we tested whether an increase in the intensity of sibling competition can exacerbate inbreeding depression in the burying beetle *Nicrophorus vespilloides*. We used a 2×3 factorial design with offspring inbreeding status (outbred or inbred) and brood size (5, 20, or 40 larvae) as the two factors. We found a main effect of inbreeding status, as inbred larvae had lower survival than outbred larvae, and a main effect of brood size, as larvae in large broods had lower survival and mass than larvae in medium-sized broods. However, there was no effect of the interaction between inbreeding status and brood size, suggesting that sibling competition did not influence the severity of inbreeding depression. Since we focused on sibling competition within homogeneous broods of either inbred or outbred larvae, we cannot rule out possible effects of sibling competition on inbreeding depression in mixed paternity broods comprising of both inbred and outbred offspring. More information on whether and when sibling competition might influence the expression of inbreeding depression can help advance our understanding of the causes underlying variation in the severity of inbreeding depression.

Keywords: burying beetle, environmental stress, fitness, inbreeding depression, *Nicrophorus vespilloides*, sibling competition

Introduction

Inbreeding results from matings between relatives and can cause a reduction in offspring fitness, known as inbreeding depression (Crnokrak & Roff, 1999). These negative fitness effects are due to the higher degree of homozygosity associated with inbreeding, which increases the risk that deleterious recessive alleles are expressed (Charlesworth & Charlesworth, 1987). Although there is widespread evidence for inbreeding depression in a range of taxa (Crnokrak & Roff, 1999), there is substantial variation both among and within species with respect to the severity of inbreeding depression (Crnokrak & Roff, 1999; Keller & Waller, 2002; Meunier & Kölliker, 2013; Moorad & Wade, 2005). This variation may be in part driven by differences in the physical and social environment, which can have profound effects on inbreeding depression (Fox & Reed, 2011; Reed *et al.*, 2012). For example, environmental stresses, such as extreme temperatures, starvation, parasitism, and competition, can exacerbate inbreeding depression (Fox & Reed, 2011; Meunier & Kölliker, 2013), while parental care, which evolved to neutralize environmental stresses to the offspring, can buffer against inbreeding depression (Avilés & Bukowski, 2006; Pilakouta *et al.*, 2015a). Although there is growing evidence that physical and social stresses can alter the severity of inbreeding depression (Fox & Reed, 2011; Reed *et al.*, 2012), little is known about the mechanisms by which particular environmental stresses influence its expression.

A recent study suggested that stresses due to intense intraspecific competition over limited resources should have a stronger effect on the severity of inbreeding depression than stresses due to harsh physical conditions, such as extreme temperatures (Yun & Agrawal, 2014). Intraspecific competition over limited resources can take several forms and can occur at different stages of the life cycle, including competition with siblings during development (Mock & Parker, 1997). Sibling competition for resources provided by the parents occurs because parents usually produce an optimistic brood size, thereby creating a mismatch

between the supply of resources from the parents and the demand of resources by the offspring (Mock & Parker, 1997). Sibling competition is an important determinant of the offspring's growth and survival in many species and may therefore be a key source of environmental stress to the offspring (Mock & Parker, 1997; Roulin & Dreiss, 2012; Meunier & Kölliker, 2013). Thus, sibling competition should be associated with an increase in this mismatch between supply and demand of resources, which in turn may exacerbate inbreeding depression. To our knowledge, the only experimental study to test this hypothesis was conducted on the European earwig, *Forficula auricularia* (Meunier & Kölliker, 2013). This study found no effect of the interaction between the intensity of sibling competition and the offspring's inbreeding status on offspring fitness, suggesting that sibling competition did not influence the severity of inbreeding depression (Meunier & Kölliker, 2013). However, the absence of such an interaction effect may reflect that there was no evidence for a main effect of inbreeding status on offspring fitness during the early life stages (Meunier & Kölliker, 2013). In order to advance our understanding of whether sibling competition can exacerbate inbreeding depression, it is now essential to focus on species in which inbred offspring suffer a significant reduction in fitness and sibling competition negatively affects offspring fitness.

Here, we tested whether sibling competition influences the severity of inbreeding depression in the burying beetle *Nicrophorus vespilloides*. This species is well suited for addressing this question, because previous work has shown that inbred offspring suffer significant fitness costs during the larval stage (Mattey *et al.*, 2013; Pilakouta *et al.*, 2015a) and that larvae in experimentally enlarged broods suffer a reduction in fitness due to the increased mismatch between supply and demand for resources (Smiseth *et al.*, 2007a). In this species, which breeds on carcasses of small vertebrates, larvae compete for resources by begging for predigested carrion from the parents and by self-feeding directly from the carcass (Smiseth *et al.*, 2003). Earlier work has also shown that sibling competition reduces offspring

fitness only in the presence of the parents, reflecting interference competition due to successful larvae excluding their siblings from getting access to the parents (Smiseth *et al.*, 2007a,b). Although the independent effects of inbreeding status and sibling competition on offspring fitness are well established in this species, there is no prior information on the effect of their interaction.

To test for such an interaction, we used a 2×3 factorial design with offspring inbreeding status and brood size as the two factors. We assessed the joint effects of inbreeding status and sibling competition on fitness traits previously shown to be affected by these two factors (Smiseth *et al.*, 2007a; Matthey *et al.*, 2013; Pilakouta *et al.*, 2015a): (i) average larval mass at dispersal; (ii) survival from hatching to dispersal; (iii) survival from dispersal to eclosion; and (iv) total survival from hatching to eclosion. If sibling competition exacerbates inbreeding depression, we would expect offspring in larger broods to incur higher fitness costs if they are inbred than if they are outbred.

Materials and methods

Study species

Like all burying beetles of the genus *Nicrophorus*, *N. vespilloides* breeds on carcasses of small vertebrates and has highly elaborate forms of parental care (Scott 1998). Once a carcass is found, parents bury it into the soil, remove any fur or feathers, deposit antimicrobial secretions to its surface, and lay eggs around it 24-48 hours after mating (Eggert, 1992; Scott, 1998). When the eggs hatch approximately 60 hours later (Smiseth *et al.*, 2006), the larvae crawl to the carcass and start feeding in a crater created by the parents on the top of the carcass. The larvae can self-feed, but the parents also provision larvae with predigested carrion (Smiseth *et al.*, 2003). Although both parents typically provide care, females often stay on the carcass for longer than males and spend more time provisioning food to the larvae

(Fetherston *et al.*, 1994; Eggert *et al.*, 1998; Smiseth & Moore, 2002; Rauter & Moore, 2004; Smiseth *et al.*, 2005; Pilakouta *et al.*, 2015b). Larvae disperse from the carcass about five days after hatching, which corresponds to the end of the parental care period. They eclose as adults about 20 days later.

Experimental design

We used virgin beetles from an outbred laboratory population maintained at The University of Edinburgh. The beetles used in this study comprised of sixth-, seventh-, and eighth-generation beetles from lines originally collected in Edinburgh, UK and Warmond, The Netherlands. Beetles were housed individually in transparent plastic containers (12×8×2 cm) filled with moist soil and kept at 20°C and constant light. Non-breeding adults were fed raw organic beef twice a week.

To test for a causal effect of sibling competition on the severity of inbreeding depression, we used a 2×3 factorial design with offspring inbreeding status (outbred or inbred) and brood size (5, 20, or 40 larvae) as the two factors. Inbred larvae were produced by pairing males and females that were full siblings ($n=186$), while outbred larvae were produced by pairing unrelated males and females that shared no common ancestors for at least two generations ($n=187$). We only used outbred parents in this experiment as inbreeding in the parents has a negative effect on offspring survival (Mattey *et al.*, 2013). These breeding pairs ($n=373$) were transferred to transparent plastic containers (17cm×12cm×6cm) filled with 1 cm of moist soil and provided with a previously frozen mouse carcass (Livefoods Direct Ltd, Sheffield, UK) of a standardized size (22–25 g). Immediately after the eggs were laid, we removed the male and moved the female and the carcass to a new container with fresh, moist soil. We removed the males because the amount of care provided by the male is highly variable and male removal has no effect on offspring fitness under

laboratory conditions (Smiseth *et al.*, 2005). We left the females to provide care for the brood because previous work on this species showed that sibling competition reduces offspring fitness only when larvae compete by begging for food from a parent (Smiseth *et al.*, 2007a,b).

When the eggs started hatching, we used the newly hatched larvae to generate inbred and outbred broods comprising of 5, 20, or 40 larvae. All experimental broods included larvae of mixed maternity in accordance with established protocols (Smiseth *et al.*, 2007; Pilakouta *et al.*, 2015b). This brood size manipulation is within the natural variation of brood size in *N. vespilloides* (mean $\pm$ SD: 21 ± 10 larvae, range: 2–47 larvae; Smiseth & Moore, 2002) and corresponds to small, average, and large broods (i.e., low, medium, and high level of sibling competition), respectively. We used a design with more than two levels of stress, because the relationship between stress and inbreeding depression may be nonlinear (Fox & Reed, 2011).

Each experimental brood (outbred or inbred) was randomly assigned to an unrelated female who had been mated either to their full-sib brother or to an unrelated male. To account for potential effects of relatedness between the female and her male partner (who was always removed before the female was provided with a foster brood), we added this information as a factor in all of our models (see below). In this species, parents cannot distinguish between unrelated foster broods and their own broods, as long as the larvae are at the same developmental stage (Müller & Eggert, 1990). Since parents kill any larvae that arrive on the carcass before their eggs are expected to hatch (Müller & Eggert, 1990), we only provided females with a brood once their own eggs had hatched.

Females were left to care for their brood until the larvae dispersed from the carcass about five days later. At dispersal from the carcass, we recorded the number of larvae and total brood mass to calculate larval survival rate and average larval mass. Lastly, to assess survival after independence (i.e., from dispersal to eclosion), we placed all dispersing larvae

from each brood into a large transparent container (17cm×12cm×6cm) filled with moist soil. About 20 days after dispersal, we recorded the number of individuals that eclosed successfully from each brood and calculated the survival rate from dispersal to eclosion.

The total sample size in the experiment was $n=166$ broods. The sample sizes for the different treatments were as follows: $n=31$ for outbred broods with 5 larvae, $n=32$ for outbred broods with 20 larvae, $n=22$ for outbred broods with 40 larvae, $n=31$ for inbred broods with 5 larvae, $n=30$ for inbred broods with 20 larvae, and $n=20$ for inbred broods with 40 larvae.

Statistical analysis

Data were analyzed using R version 3.2.0. Larval mass at dispersal was analyzed using a linear model. Proportion data (i.e., survival rates from hatching to dispersal, from dispersal to eclosion, and from hatching to eclosion) were analyzed using generalized linear models fitted with a binomial distribution corrected for overdispersion. Proportion data were entered into the models using the 'cbind' function.

All models included offspring inbreeding status (outbred or inbred), brood size (small, medium, or large), and the interaction between these two factors. As additional factors, we included carcass mass, the relatedness between the foster mother and her removed male partner (i.e., whether the female rearing the brood had been mated to a full sibling or an unrelated male), the interaction between foster parent relatedness and offspring inbreeding status, and the interaction between foster parent relatedness and brood size. Decisions on which factors to include in the final models were based on AIC scores. For models where brood size was found to have a statistically significant effect, we used the 'glht' function in the 'multcomp' package (Hothorn *et al.* 2008) to perform a Tukey test for post-hoc pairwise comparisons.

Results

We found evidence for a main effect of offspring inbreeding status on survival to dispersal, survival to eclosion, and total survival, as inbred larvae suffered lower survival than outbred larvae (Table 1; Figure 1b-d). There was no significant difference in average larval mass between inbred and outbred larvae (Table 1; Figure 1a). In addition, we found evidence for a main effect of sibling competition on offspring fitness: larvae in large broods were smaller and had a lower rate of survival to dispersal than larvae in medium-sized broods (Tables 1 and 2; Figure 1a-b). Sibling competition also had a non-significant effect on total survival (Tables 1 and 2; Figure 1d). In contrast, larvae in small broods had a lower rate of survival to eclosion than larvae in medium-sized or large broods (Table 2; Figure 1c). We found no evidence that sibling competition exacerbated inbreeding depression, as there was no effect of the interaction between the offspring's inbreeding status and the size of the brood on any component of offspring fitness (Table 1). In addition, there was no effect of foster parent relatedness (i.e. whether the foster mother had been mated to a brother or an unrelated male) or the interaction between foster parent relatedness and offspring inbreeding status on offspring fitness (Table S1). There was an effect of the interaction between foster parent relatedness and brood size on survival to eclosion but not on larval mass, survival to dispersal, or total survival (Table S1). Lastly, larvae reared on larger carcasses had a higher larval mass at dispersal and higher overall survival (Table S1).

Discussion

We find no evidence for an effect of the interaction between sibling competition and inbreeding status, suggesting that sibling competition does not exacerbate inbreeding depression in *N. vespilloides*. We show that inbreeding status negatively affected offspring fitness, as inbred larvae suffered lower survival during both the larval and pupal stages.

215 These results demonstrate that there is significant inbreeding depression in this species as
216 reported in previous studies (Mattey *et al.*, 2013; Pilakouta *et al.*, 2015a). Furthermore, we
217 show that sibling competition reduced offspring fitness, as larvae in large broods were
218 smaller and suffered lower survival to dispersal than larvae in medium-sized broods. These
219 results confirm that the intensity of sibling competition has a negative effect on larval fitness
220 as previously reported by Smiseth *et al.* (2007a). Even though there were significant main
221 effects of both inbreeding status and sibling competition, there was no effect of the
222 interaction between these two factors on offspring fitness. To our knowledge, the only other
223 study to investigate this question (conducted on the European earwig) found no evidence for
224 a main effect of inbreeding status on offspring fitness in the early life stages (Meunier &
225 Kölliker, 2013). Given the absence of inbreeding depression, it was not possible to test
226 whether inbreeding depression becomes more severe when sibling competition increases.
227 Thus, our study is the first to show that sibling competition for resources provided by the
228 parents does not exacerbate inbreeding depression in a species where inbred offspring do
229 suffer substantial fitness costs.

230 There is accumulating evidence that the negative effects of inbreeding can be
231 intensified by a wide range of environmental stresses, such as parasitism, starvation,
232 population density, extreme temperatures, and exposure to chemicals (Armbruster & Reed,
233 2005; Waller *et al.*, 2008; Fox & Reed, 2011; Reed *et al.*, 2012). Nevertheless, we still have a
234 limited understanding of the mechanisms by which particular stresses exacerbate inbreeding
235 depression. Yun and Agrawal (2014) argue that density-dependent stresses caused by intense
236 competition among conspecifics should have a stronger effect on the severity of inbreeding
237 depression than density-independent stresses caused by harsh physical conditions. In their
238 study on *Drosophila melanogaster*, they found a moderate correlation between environmental

stress and density dependence, but inbreeding depression was significantly more correlated with density dependence than environmental stress *per se* (Yun & Agrawal, 2014).

We expected that an increase in sibling competition should affect the severity of inbreeding depression because sibling competition is a density-dependent source of environmental stress caused by a mismatch between the supply and demand for resources (Mock & Parker, 1997; Roulin & Dreiss, 2012). Yet, we find that sibling competition does not exacerbate inbreeding depression in *N. vespilloides*, which appears to contradict the argument made by Yun and Agrawal (2014). One potential explanation for this discrepancy is the difference in experimental designs between our study and that of Yun and Agrawal (2014). We studied the effects of sibling competition within homogeneous broods comprised of either outbred or inbred larvae, while Yun and Agrawal (2014) studied the effects of competition within heterogeneous groups of unrelated inbred and outbred fruit flies. Thus, in our study, any effect of sibling competition on the severity of inbreeding depression would be mediated through an increase in the level of stress. In contrast, in the latter study, such effects would be mediated through direct competitive interactions between inbred individuals (inferior competitors) and outbred individuals (superior competitors). Homogeneous broods comprised of either outbred or inbred larvae are likely to be the norm in *N. vespilloides* given that caring males sire over 90% of the offspring in their brood (Müller & Eggert, 1989). In such homogeneous broods, all larvae may be equally disadvantaged by stresses caused by a shortage of resources. However, we cannot rule out possible effects of sibling competition on the severity of inbreeding depression in mixed paternity broods comprising of both inbred and outbred larvae. Under those conditions, higher-quality (i.e., outbred) offspring may outcompete their lower-quality (i.e., inbred) half-siblings, thus magnifying differences in fitness between them. We encourage future studies to investigate this question in the context of family groups that comprise of both outbred and inbred offspring, as in socially

monogamous birds where the female is closely related to her social partner and has extra-pair matings with non-relatives (e.g., Blomqvist *et al.*, 2002; Foerster *et al.*, 2003; Brouwer *et al.*, 2011; Reid *et al.*, 2015).

Even though understanding the factors that drive the observed variation in the severity of inbreeding depression across species and across environments could have important implications for the conservation of many endangered populations, these dynamics are still not well understood. Our findings suggest that the intensity of sibling competition may not contribute towards variation in the severity of inbreeding depression, at least within homogeneous broods of inbred offspring. Determining whether and when sibling competition might play a role in the expression of inbreeding depression may help us better understand the causes underlying temporal and spatial patterns of variation in inbreeding depression in natural populations.

Acknowledgments

We thank the Edinburgh Countryside Rangers for permission to collect beetles at Craiglockhart Park and Daniel Rozen for supplying beetles from the Netherlands. The study was funded by the Institute of Evolutionary Biology and the School of Biological Sciences, University of Edinburgh.

References

- Armbruster, P. & Reed, D.H. 2005. Inbreeding depression in benign and stressful environments. *Heredity* **95**: 235–242.
- Avilés, L. & Bukowski, T.C. 2006. Group living and inbreeding depression in a subsocial spider. *Proc. Biol. Sci.* **273**: 157–163.

- Blomqvist, D., Andersson, M., Kupper, C., Cuthill, I.C., Kis, J., Lanctot, R.B., Sandercock, B.K., Székely, T., Wallander, J. & Kempenaers, B. 2002 Genetic similarity between mates and extra-pair parentage in three species of shorebird. *Nature* **419**: 613–615.
- Brouwer, L., van de Pol, M., Atema, E. & Cockburn, A. 2011. Strategic promiscuity helps avoid inbreeding at multiple levels in a cooperative breeder where both sexes are philopatric. *Mol. Ecol.* **20**: 4796–4807.
- Charlesworth, D. & Charlesworth, B. 1987. Inbreeding depression and its evolutionary consequences. *Annu. Rev. Ecol. Syst.* **18**: 237–268.
- Crnokrak, P. & Roff, D. 1999. Inbreeding depression in the wild. *Heredity* **83**: 260–270.
- Eggert, A.-K. 1992. Alternative male mate-finding tactics in burying beetles. *Behav. Ecol.* **3**: 243–254.
- Eggert, A.-K., Reinking, M. & Müller, J.K. 1998. Parental care improves offspring survival and growth in burying beetles. *Anim. Behav.* **55**: 97–107.
- Fetherston, I.A., Scott, M.P. & Traniello, J.F.A. 1994. Behavioural compensation for mate loss in the burying beetle *Nicrophorus orbicollis*. *Anim. Behav.* **47**: 777–785.
- Foerster, K., Delhey, K., Johnsen, A., Lifjeld, J.T. & Kempenaers, B. 2003. Females increase offspring heterozygosity and fitness through extra-pair matings. *Nature* **425**: 714–717.
- Fox, C.W. & Reed, D.H. 2011. Inbreeding depression increases with environmental stress: an experimental study and meta-analysis. *Evolution* **65**: 246–258.
- Hothorn, T., Bretz F. & Westfall, P. 2008. Simultaneous inference in general parametric models. *Biometrical Journal* **50**: 346–363.
- Keller, L. & Waller, D. 2002. Inbreeding effects in wild populations. *Trends Ecol. Evol.* **17**: 230–241.

- Mattey, S.N., Strutt, L. & Smiseth, P.T. 2013. Intergenerational effects of inbreeding in *Nicrophorus vespilloides*: offspring suffer fitness costs when either they or their parents are inbred. *J. Evol. Biol.* **26**: 843–853.
- Meunier, J. & Kölliker, M. 2013. Inbreeding depression in an insect with maternal care: influences of family interactions, life stage and offspring sex. *J. Evol. Biol.* **26**: 2209–2220.
- Mock, D.W. & Parker, G.A. 1997. The evolution of sibling rivalry. Oxford University Press.
- Moorad, J.A. & Wade, M.J. 2005. A genetic interpretation of the variation in inbreeding depression. *Genetics* **170**: 1373–1384.
- Müller, J.K. & Eggert, A.-K. 1989. Paternity assurance by "helpful" males: adaptations to sperm competition in burying beetles. *Behav. Ecol. Sociobiol.* **24**: 245–249.
- Müller, J.K. & Eggert, A.-K. 1990. Time-dependent shifts between infanticidal and parental behavior in female burying beetles: a mechanism of indirect mother-offspring recognition. *Behav. Ecol. Sociobiol.* **27**: 11–16.
- Pilakouta, N., Jamieson, S., Moorad, J.A. & Smiseth, P.T.S. 2015. Parental care buffers against inbreeding depression in burying beetles. *Proc. Natl. Acad. Sci. U.S.A.* **112**: 8031–8035.
- Pilakouta, N., Richardson, J. & Smiseth, P.T.S. 2015. State-dependent cooperation in burying beetles: parents adjust their contribution towards care based on both their own and their partner's size. *J. Evol. Biol.* (doi: 10.1111/jeb.12712)
- Rauter, C.M. & Moore, A.J. 2004. Time constraints and trade-offs among parental care behaviours: effects of brood size, sex and loss of mate. *Anim. Behav.* **68**: 695–702.
- Reed, D.H., Fox, C.W., Enders, L.S. & Kristensen, T.N. 2012. Inbreeding-stress interactions: evolutionary and conservation consequences. *Ann. N.Y. Acad. Sci.* **1256**: 33–48.

Reid, J.M., Arcese, P., Keller, L.K., Germain, R.R., Duthie, A.B., Losdat, S., Wolak, M.E. & Nietlisbach, P. 2015. Quantifying inbreeding avoidance through extra-pair reproduction. *Evolution* **69**: 59-74.

Roulin, A. & Dreiss, A.N. 2012. Sibling competition and cooperation over parental care. In: The Evolution of Parental Care (N.J. Royle, P.T. Smiseth, M. Kolliker, eds), pp. 133–149. Oxford University Press, Oxford.

Scott, M.P. 1998. The ecology and behavior of burying beetles. *Annu. Rev. Entomol.* **43**: 595–618.

Smiseth, P.T., Darwell, C. & Moore, A.J. 2003. Partial begging: an empirical model for the early evolution of offspring signalling. *Proc. R. Soc. B* **270**: 1773–1777.

Smiseth, P.T., Dawson, C., Varley, E. & Moore, A.J. 2005. How do caring parents respond to mate loss? Differential response by males and females. *Anim. Behav.* **69**: 551–559.

Smiseth, P.T., Lennox, L. & Moore, A.J. 2007a. Interaction between parental care and sibling competition: parents enhance offspring growth and exacerbate sibling competition. *Evolution* **61**: 2331-2339.

Smiseth, P.T. & Moore, A.J. 2002. Does resource availability affect offspring begging and parental provisioning in a partially begging species? *Anim. Behav.* **63**: 577–585.

Smiseth, P.T., Ward, R.S.J. & Moore, A.J. 2006. Asynchronous hatching in *Nicrophorus vespilloides*, an insect in which parents provide food for their offspring. *Funct Ecol* **20**: 151–156.

Smiseth, P.T., Ward, R.S.J. & Moore, A.J. 2007b. Parents influence asymmetric sibling competition: experimental evidence with partially dependent young. *Ecology* **88**: 3174–3182.

Waller, D.M., Dole, J. & Bersch, A.J. 2008. Effects of stress and phenotypic variation on inbreeding depression in *Brassica rapa*. *Evolution* **62**: 917–931.

360 Yun, L. & Agrawal, A.F. 2014. Variation in the strength of inbreeding depression across
361 environments: effects of stress and density dependence. *Evolution* **68-12**: 3599–3606.
362

Table 1. Effects of offspring inbreeding status (inbred or outbred) and sibling competition (small, medium-sized, or large brood) on offspring fitness traits: average larval mass at dispersal (mg), survival rate from hatching to dispersal, survival rate from dispersal to eclosion, and survival rate from hatching to eclosion.

	Larval mass (mg)		Survival to dispersal		Survival to eclosion		Total survival	
	<i>F</i> -value	<i>P</i> -value	LR χ^2	<i>P</i> -value	LR χ^2	<i>P</i> -value	LR χ^2	<i>P</i> -value
Inbreeding status	2.27	0.13	4.54	0.03	37.79	<0.0001	17.17	<0.0001
Brood size	4.93	<0.01	6.72	0.03	12.23	<0.01	6.07	0.048
Interaction	0.09	0.91	0.31	0.86	4.00	0.14	0.38	0.83

Larval mass data were analyzed using a linear model. Survival rate data were analyzed using generalized linear models fitted with a binomial distribution corrected for overdispersion. LR refers to likelihood ratio. Statistically significant *P*-values are in bold.

Table 2. Post-hoc pairwise comparisons (Tukey test) for the effect of sibling competition (small, medium-sized, or large brood) on offspring fitness traits: average larval mass at dispersal (mg), survival rate from hatching to dispersal, survival rate from dispersal to eclosion, and survival rate from hatching to eclosion.

Brood size	Larval mass (mg)				Survival to dispersal				Survival to eclosion				Total survival			
	Est	SE	<i>t</i>	<i>P</i>	Est	SE	<i>z</i>	<i>P</i>	Est	SE	<i>z</i>	<i>P</i>	Est	SE	<i>z</i>	<i>P</i>
Small-Medium	-12.37	5.75	-2.15	0.08	-0.10	0.26	-0.38	0.92	-1.15	0.34	-3.36	<0.01	-0.49	0.25	-1.97	0.11
Small-Large	6.65	6.38	1.04	0.55	0.29	0.25	1.13	0.49	-1.13	0.33	-3.40	<0.01	-0.18	0.24	-0.73	0.74
Medium-Large	19.02	6.39	2.98	<0.01	0.39	0.15	2.52	0.03	0.02	0.27	-0.06	>0.99	0.31	0.15	2.08	0.09

We provide information on the parameter estimates (Est), standard errors (SE), test statistics (*t*- and *z*-values), and *P*-values. Parameter estimates represent the difference when subtracting the mean fitness of larvae in the second group listed from the mean fitness of larvae in the first group on each row. Statistically significant *P*-values are in bold.