



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Nitric oxide function in plant abiotic stress

Citation for published version:

Fancy, NN, Bahlmann, AK & Loake, GJ 2016, 'Nitric oxide function in plant abiotic stress', *Plant, Cell and Environment*, vol. 40, no. 4, pp. 462-472. <https://doi.org/10.1111/pce.12707>

Digital Object Identifier (DOI):

[10.1111/pce.12707](https://doi.org/10.1111/pce.12707)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Plant, Cell and Environment

Publisher Rights Statement:

© 2016 JohnWiley & Sons Ltd

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Table 2: Differential S-nitrosylation of individual proteins related to abiotic stress.

S-nitrosylated proteins related to abiotic stress	Activity change after S-nitrosylation	Cellular localization of the protein	Reference
AHb1	ND	Cytoplasm	(Perazzolli <i>et al.</i> 2004)
APX	+	Cytoplasm	(Bai <i>et al.</i> 2011; Begara-Morales <i>et al.</i> 2011)
GAPDH (<i>in vitro</i> only)	-	Cytoplasm	(Wawer <i>et al.</i> 2010; Lin <i>et al.</i> 2012)
OST1/SnRK2.6	-	Cytoplasm	(Wang <i>et al.</i> 2015)
PAL	-	Cytoplasm	(Vanzo <i>et al.</i> 2014)
Trx	ND	Cytoplasm	(Lin <i>et al.</i> 2012)
GR	+/=	Cytoplasm & Chloroplast	(Bai <i>et al.</i> 2011; Begara-Morales <i>et al.</i> 2011)
Rubisco	-	Chloroplast	(Abat & Deswal 2009)
PrxII F	-	Mitochondria	(Camejo <i>et al.</i> 2013)
CAT	-	Peroxisome	(Ortega-Galisteo <i>et al.</i> 2010)
DHAR	+	Peroxisome	(Bai <i>et al.</i> 2011)
GOX	-	Peroxisome	(Ortega-Galisteo <i>et al.</i> 2010)
MDAR	-	Peroxisome	(Begara-Morales <i>et al.</i> 2011)
MDH	-	Peroxisome	(Ortega-Galisteo <i>et al.</i> 2010)

Key: ND, Not Detected; +, increased; -, decreased; =, not changed.

Figure Legends

Figure 1. Regulation of S-nitrosylation. S-nitrosylation is the attachment of an NO group to a cysteine (Cys) thiol (SH) to form an S-nitrosothiol (SNO). The SNO bond is reversible due to its redox sensitivity: i.e. ascorbate, reduced metal ions and GSH can reduce SNO to SH. S-nitrosoglutathione (GSNO), which acts as a stable NO reservoir, can transfer NO to SH group via transnitrosylation. Turnover of GSNO occurs via the enzyme GSNO reductase (GSNOR), generating oxidized glutathione (GSSG) and ammonia (NH₃).

Figure 2. Regulation of ascorbate-glutathione cycle by S-nitrosylation during salt stress. During salt stress the ascorbate–glutathione cycle is modulated via S-nitrosylation of APX, MDAR and GR proteins. MDAR activity is reduced after S-nitrosylation while GR activity is not significantly affected by this modification. However, APX activity is enhanced by S-nitrosylation. This regulation suggests that APX works to detoxify hydrogen peroxide while GR supports the regeneration of GSH, thereby maintaining the antioxidant capability of the ascorbate-glutathione cycle under stress conditions. APX, ascorbate peroxidase; GR, Glutathione reductase; MDAR, monodehydroascorbate reductase.

Figure 3. Model showing the role of NO and S-nitrosylation during drought stress signalling. Drought stress triggers ABA accumulation which in turn can induce the production of H₂O₂ leading to the generation of NO, which results in stomatal closure via the activation of a MAPK pathway. On the other hand, OST1/SnRK2.6 is a central component that regulates ABA signalling via a phosphorylation cascade. NO can regulate ABA signaling through S-nitrosylation of OST1/SnRK2.6, which inhibits its kinase activity, thus acting as a negative regulator of ABA signalling.

Figure 4. S-nitrosylation regulates different nodes of abiotic stress signalling. During abiotic stress, proteins related to different nodes of stress signalling become targets for S-nitrosylation/denitrosylation. These proteins include antioxidant proteins, proteins involved in RNS/ROS, cellular metabolism and also proteins that regulate programmed cell death (PCD). These proteins change their S-nitrosylation pattern during stress to control cell fate in two ways: either

stress acclimation or PCD. Rubisco, ribulose-1,5-bisphosphate carboxylase/oxygenase; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; Trx, thioredoxin; AHb1, haemoglobin 1; APX, ascorbate peroxidase; GR, Glutathione reductase; DHAR, dehydroascobate reductase; GOX, glycolate oxidase; PrxII F, peroxiredoxin II F; MDAR, monodehydroascorbate reductase; CAT, catalase.




Figure 1. Regulation of S-nitrosylation.




Figure 2. Regulation of ascorbate-glutathione cycle by S-nitrosylation during salt stress.




Figure 3. Model showing the role of NO and S-nitrosylation during drought stress signalling.




Figure 4. S-nitrosylation regulates different nodes of abiotic stress signalling.