



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Brain state dependent stimulus information in the auditory thalamocortical system

Citation for published version:

Bamber, J, Sakata, S & Herrmann, JM 2015, 'Brain state dependent stimulus information in the auditory thalamocortical system' BMC Neuroscience, vol 16, no. 1, pp. P99. DOI: 10.1186/1471-2202-16-S1-P99

Digital Object Identifier (DOI):

[10.1186/1471-2202-16-S1-P99](https://doi.org/10.1186/1471-2202-16-S1-P99)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

BMC Neuroscience

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



POSTER PRESENTATION

Open Access

Brain state dependent stimulus information in the auditory thalamocortical system

Jon Bamber^{1*}, Shuzo Sakata², J Michael Herrmann³

From 24th Annual Computational Neuroscience Meeting: CNS*2015
Prague, Czech Republic. 18-23 July 2015

Activity in the absence of stimuli is ubiquitous across the thalamocortical system (TS), with patterns of spontaneous activity reflecting ongoing behavioural state. Under anaesthesia and during deep sleep the TS operates in an inactivated state (characterised by low frequency high amplitude oscillations in local field potential (LFP)) in which neurons collectively alternate between periods of local silence and high synaptic activity. During wakefulness and REM sleep, however, the TS operates in an activated state (characterised by high frequency low amplitude oscillations in LFP) in which neurons fire in a sustained desynchronised manner. Such brain states may be indicative of different modes of neural processing, but the effect of brain state on neural processing remains unclear.

Here we analyse data recorded in the auditory TS of urethane anaesthetised rats subjected to single-click acoustic stimulation over a range of intensities, presented in both the inactivated state (natural under the anaesthesia) and the activated state (induced through electrical stimulation of the basal forebrain). Evoked spike trains were identified for single units in the auditory thalamus and across depths of the primary auditory cortex. Mutual information (MI) between stimulus and response was then computed for spike probability, counts and timing.

Evoked spiking activity was observed to last around 300ms, consisting of distinct initial and secondary (*rebound*) activity. Analysis of responses of single units over a full 300ms window showed that spike count and timing measures gave little more MI than response probability, suggesting that stimulus intensity is primarily encoded probabilistically, at least at the level of the single

unit. Moreover, whilst many single units are uninformative of stimulus intensity, the most informative thalamocortical (TC) and infragranular (IF) single units show increased MI in the activated state. Additionally, upon temporally partitioning data according to initial or rebound activity, we see that TC units are more informative in the activated state during initial activity whereas IF units are more informative in the activated state during rebound activity, suggesting that information loss in the inactivated state may be cumulative in time as sensory signals propagate through neural circuits.

Authors' details

¹Institute for Adaptive and Neural Computation, University of Edinburgh, Edinburgh, EH8 9AB, UK. ²Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, G4 0RE, UK. ³Institute of Perception, Action and Behaviour, University of Edinburgh, Edinburgh, EH8 9AB, UK.

Published: 18 December 2015

doi:10.1186/1471-2202-16-S1-P99

Cite this article as: Bamber et al.: Brain state dependent stimulus information in the auditory thalamocortical system. *BMC Neuroscience* 2015 **16**(Suppl 1):P99.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

 **BioMed Central**

* Correspondence: jon.bamber@ed.ac.uk

¹Institute for Adaptive and Neural Computation, University of Edinburgh, Edinburgh, EH8 9AB, UK

Full list of author information is available at the end of the article