

THE UNIVERSITY of EDINBURGH

Edinburgh Research Explorer

Plumbing the depths: extracellular water storage in specialized leaf structures and its functional expression in a three-domain pressure-volume relationship

Citation for published version:

Nguyen, HT, Meir, P, Wolfe, J, Mencuccini, M & Ball, MC 2016, 'Plumbing the depths: extracellular water storage in specialized leaf structures and its functional expression in a three-domain pressure-volume relationship', *Plant, Cell and Environment*. https://doi.org/10.1111/pce.12788

Digital Object Identifier (DOI):

10.1111/pce.12788

Link:

Link to publication record in Edinburgh Research Explorer

Document Version: Peer reviewed version

Published In: Plant, Cell and Environment

Publisher Rights Statement:

This article is protected by copyright. All rights reserved.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Édinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1	Plumbing the depths: extracellular water storage in specialized leaf structures and its
2	functional expression in a three-domain pressure-volume relationship
3	
4	
5	Running title: Three-domain pressure-volume relationship reflects leaf structure
6	
7	
8	
9	¹ Hoa T. Nguyen, ^{1,3} Patrick Meir, ² Joe Wolfe, ^{3,4} Maurizio Mencuccini and ¹ Marilyn C. Ball
10	¹ Plant Science Division, Research School of Biology, The Australian National University, Acton, ACT
11	2601, Australia
12	² School of Physics, University of New South Wales, Sydney, NSW 2052, Australia
13	³ School of GeoSciences, University of Edinburgh, Crew Building, West Mains Road, Edinburgh EH9
14	3JN, UK
15	⁴ ICREA at CREAF, Universidad Autonoma de Barcelona, Cerdanyola del Valles, Barcelona, Spain
16	
17	
18	Authors for correspondence:
19	Hoa T. Nguyen
20	Email: <u>hoa-thi.nguyen@anu.edu.au</u>
21	
22	Marilyn C. Ball
23	Email: marilyn.ball@anu.edu.au
24	
25	
26	

27 Abstract:

28 A three-domain pressure - volume relationship (PV curve) was studied in relation to leaf anatomical 29 structure during dehydration in the grey mangrove, Avicennia marina. In domain 1, relative water 30 content (RWC) declined 13% with 0.85 MPa decrease in leaf water potential, reflecting a decrease in 31 extracellular water stored primarily in trichomes and petiolar cisternae. In domain 2, RWC decreased 32 by another 12% with further reduction in leaf water potential to -5.1 MPa, the turgor loss point. Given 33 the osmotic potential at full turgor (-4.2 MPa) and the effective modulus of elasticity (~40 MPa), 34 domain 2 emphasized the role of cell wall elasticity in conserving cellular hydration during leaf water 35 loss. Domain 3 was dominated by osmotic effects and characterized by plasmolysis in most tissues 36 and cell types without cell wall collapse. Extracellular and cellular water storage could support an evaporation rate of 1 mmol m⁻²s⁻¹ for up to 54 and 50 min, respectively, before turgor loss was 37 38 reached. This study emphasizes the importance of leaf anatomy for the interpretation of PV curves, 39 and identifies extracellular water storage sites that enable transient water use without substantive 40 turgor loss when other factors, such as high soil salinity, constrain rates of water transport.

41

42 Key words: mangrove, PV curve, leaf structure, dehydration, plasmolysis, extracellular water,
43 cisternae, trichomes, leaf water capacitance.

44 Introduction

45 Increasing vulnerability to drought is a major global concern as rise in average temperature is 46 associated with increase in the frequency and intensity of drought in many locations (Reichstein et al., 47 2013). Drought - induced tree mortality has been recorded worldwide in diverse ecosystems and 48 climatic zones in the past few decades (Allen et al., 2015). These reports range across a continuum of 49 drought tolerance from species as drought sensitive as those growing in tropical rainforests (Rowland 50 et al., 2015) to the highly drought and salinity tolerant species of mangrove forests (Lovelock et al., 51 2009). Drought-induced mortality has been linked with stem hydraulic deterioration (Anderegg et al., 52 2012, Rowland et al., 2015). Determination of stem vulnerability to drought can be difficult, especially 53 in trees. However, Zimmermann (1983) suggested that leaves should be more vulnerable to hydraulic 54 dysfunction than stems, as shown, for example, in sugar maple (Choat et al., 2005). Indeed, leaf 55 traits, such as the turgor loss point, can provide a powerful means of identifying species that are most 56 vulnerable to drought (Bartlett et al., 2012).

57

58 There is an urgent need to better understand the interplay of leaf water relations and leaf structure. 59 Most recent studies have focussed on correlations between hydraulic anatomy and hydraulic 60 conductance that inform our understanding of morphological constraints on carbon gain in relation to 61 water loss by leaves under different environmental conditions (Sack & Scoffoni, 2013). For example, 62 interspecific differences in vein and stomatal densities are associated with differences in leaf hydraulic 63 conductance, and hence also in capacities for photosynthetic gas exchange under conditions of both 64 high and low water availability (Brodribb & Holbrook, 2003). However, much less is known about the 65 influence of other aspects of leaf anatomy on leaf water relations as measured by pressure/volume 66 relationships.

67

The pressure-volume method is used to analyse the water content of a leaf in terms of water potential and its components (Scholander *et al.*, 1964, Tyree & Hammel, 1972, Cheung *et al.*, 1975). For a leaf or tissue in a pressure chamber, with the petiole protruding through a seal into air at atmospheric pressure, a plot of the pressure in the chamber vs. the volume of sap expressed from the petiole or the residual weight of the leaf (hereafter called the PV curve) yields a relationship which is usually reported as having two domains with differing behaviours. In these two-domain examples, when the

74 leaf water potentials are relatively high, decline in water content with increasing pressure in the 75 chamber is dominated by quasi-elastic shrinkage of the cells as the turgor pressure is reduced. This 76 first domain ends when turgor falls to near zero (i.e. intracellular osmotic pressure approximately 77 equals the air pressure in the chamber). Further reduction in cell water content is determined mainly 78 by the osmotic behaviour of the flaccid cell and is often approximated by the van't Hoff law. Thus the 79 PV curve is information-rich, enabling the calculation and functional evaluation of parameters such as 80 leaf capacitance for water storage (Q) (Tyree & Ewers, 1991), the osmotic potentials at full turgor (Ψ_{π}^{100}) and at the turgor loss point (Ψ_{π}^{0}) , and an effective volumetric modulus of elasticity ($\mathbf{\epsilon}$), all of 81 82 which are strongly associated with interspecific variation in drought tolerance (Bartlett et al., 2012).

83

84 The theory underpinning interpretation of PV curves was developed with the approximation that 85 leaves behave like osmometers (Scholander et al., 1965) and that an average behaviour can be 86 described for all cells in a leaf. However, two aspects of leaf structural complexity affect interpretation 87 of PV curves. First, leaves are composed of a diverse range of cells specialised in different aspects of 88 function. These cell types are structurally distinct, displaying, for example, differences in cell size and 89 in the thickness and composition of cell walls, each of which could affect their modulus of elasticity, 90 and hence maintenance of turgor during desiccation. In addition, even within a given cell type, 91 variation in solute composition among adjacent hydrated cells (McCully et al., 2010) could affect the 92 osmotic potentials of individual cells at full turgor, and hence also affect the water potentials at which 93 turgor is lost (Bartlett et al., 2012). Such differences between cell types and between cells of a given 94 type could explain the curvilinear transition between the linear elastic and osmotic domains of the PV 95 curve (Cheung et al., 1975). Indeed, Melkonian et al. (1982) suggested that the departure from 96 linearity of the elastic domain of the PV curve signals the onset of turgor loss in the most vulnerable 97 cells within a leaf; similarly, Cheung et al. (1975) suggested that the turgor loss point estimated from 98 the onset of osmotic behaviour in the PV curve represents the extreme condition when all cells have 99 lost turgor.

100

A second complication to interpretation of PV curves arises from the distribution of water in symplastic
 and apoplastic spaces within leaves. The symplastic space consists of the intracellular components of
 living cells bounded by the plasma membranes. The apoplastic space is more difficult to define.

Canny (1995) recognized four different appoplastic spaces: the lumina of xylem conduits and fibers,
 the gaseous intercellular space, and the water and Donnan free spaces of cell walls. However, more
 spaces may be identified as functional leaf anatomy becomes better understood. Differentiation of
 symplastic and apoplastic water is important in the estimation of PV parameters, such as elasticity.

109 In some species, PV curves with more than two domains have also been reported. The temporary 110 accumulation of water in apoplastic spaces has given rise in some studies to a third domain at the 111 highest levels of hydration in a PV curve. For example, in sphagnum moss, an initial domain largely 112 corresponded to the loss of water stored in the specialised, dead, hollow hyaline cells (Hajek & 113 Beckett, 2008). In this domain, sphagnum moss shoots lost up to 70% of water content without a 114 measurable change in shoot water potential. Once this extracellular water was depleted, then the PV 115 curves of drying shoots showed the two successive domains dominated by turgor reduction and 116 osmotic relationships (Hajek & Beckett, 2008). Similar patterns were observed in other poikilohydric 117 species of moss, lichen, liverworts, and a filmy fern (Beckett, 1997). A similar domain has also been 118 reported in more complex homoiohydric species, such as white oak (Parker & Pallardy, 1987), 119 Pseudotsuga menziesii (Mirb.) Franco (Kubiske & Abrams, 1991), and olive (Dichio et al., 2003), but 120 interpreted as an artefact of rehydration, also known as "the plateau effect" (Parker & Pallardy, 1987).

121

122 In the present study, the nature of the PV curve was related to leaf anatomy in Avicennia marina 123 subsp. australasica (Walp.) J. Everett, a highly salt-tolerant mangrove species (Naidoo et al., 2011, 124 Nguyen et al., 2015). Leaves of A. marina naturally endure enormous variability in hydration with leaf 125 water potentials ranging from -0.1 MPa during rainfall events to -6.0 MPa during low tides under hot, 126 dry weather (Constable, 2014, Walker, 2014). The leaf structure of *A. marina* is typical of C₃ species 127 in having layers of palisade and spongy mesophyll. In addition, A. marina has salt secretion glands, a 128 trichome layer on the abaxial surface, and a hypodermal water storage layer, similar to many other 129 halophytes (Hutchings & Saenger, 1987). Thus A. marina is ideal for our study because it is a widely 130 distributed species with leaves that are both highly tolerant of drought and possess complex cellular 131 structures.

132

133 Materials and Methods

134 Plant materials

One branch was collected from each of five co-occurring trees of *A. marina* subsp. *australasica* growing naturally along the Clyde River, Batemans Bay, New South Wales, Australia, and brought back to the lab in a black plastic bag stored in an insulated box to prevent water loss. A hand-held refractometer (A.S.T. Co. Ltd., Japan) was used to measure salinity of surface water collected at high tide and soil water collected at low tide from a soil depth of 30 cm. Soil water was extracted with a suction device (McKee, 1993) from three locations around each tree and measurements of these three salinities were averaged to give one value per tree (n = 5 trees).

142

143 **PV curve**

144 Upon return to the lab, the cut ends of all branches were recut under a solution of 25 mM NaCl and 145 transferred to a bucket, keeping the cut ends under solution and enclosing the bucket to keep the 146 humidity high around the leaves. The branches were allowed to rehydrate overnight in the dark in a 147 cold room at 5°C. The composition of this perfusion solution matched average salt concentrations measured in the xylem sap of stems and leaves from Avicennia marina grown in seawater salinity 148 149 under laboratory (Ball, 1988) and natural field conditions (Stuart et al., 2007). Before PV curves were 150 constructed, leaves were cut from branches near the petiole - stem junctions while the petioles were 151 submerged in perfusion solution. Leaves were then transferred to a beaker, keeping the petioles 152 under solution. The beaker was covered with plastic wrap to maintain a high humidity around the 153 leaves and they were allowed to fully hydrate in the dark at room temperature for at least 30 min. The 154 leaves were then dabbed dry with tissue paper to remove excess water before measuring leaf fresh 155 mass (XP 205 Metter Toledo balance, Mettler - Toledo Ltd., Greifensee, Switzerland) and leaf water 156 potential (Pressure chamber model 1050D, PMS Instrument, Albany, USA).

157

Five PV curves were constructed, each made with one leaf from one tree. Leaf water potential was measured at intervals corresponding to decreases in leaf fresh mass of 5-10 mg as the leaves airdried at room temperature. Time intervals between measurements ranged from 1 min at the start of dehydration to 40 min for the final measurements, made when leaf water potentials reached values between -5 and -6 MPa, corresponding to the minimal values Constable (2014) measured during diurnal periods under field conditions in the same population of trees as in the present study. The leaf perimeter with petiole excluded was traced on paper for leaf area measurement with a LI 3100 area
 meter (LICOR Inc., Lincoln, Nebraska, USA). Leaf dry mass was recorded after 48 h oven-drying to
 constant mass at 70°C. PV curve parameters were calculated as in Table 2 and other particulars are
 given in the results.

168

169 **Leaf anatomy**

170 Light microscopy

171 Two opposite, mature leaves were selected for anatomical analyses from the same branches that provided leaves for measurement of PV curves. One leaf in each pair was used for measurements of 172 173 the tissue composition and trichome volume. Trichomes were approximated as cylinders, and 174 trichome volumes were estimated based on height and internal lumen diameter. Transverse sections 175 (20 - 30 µm thickness) were cut from the petiole and leaf blade with a sledge microtome (GSL 1, S. 176 Lucchinetti, Schenkung Dapples, Zürich, Switzerland) and stained with either Toluidine blue (0.05% 177 w/v in water) or a 50:50 mixture of Alcian Blue (1% w/v in 50% v/v alcohol) and Safranin O (1% w/v in 178 water). The matching leaf in each pair was used to estimate trichome density, which required an 179 inside-out epidermal preparation. The leaf epidermis was loosened by incubating the leaf in 180 maceration solution (five parts 30% v/v hydrogen peroxide to one part glacial acetic acid) at 70°C for 181 at least 2 h. Then the abaxial epidermis was removed with forceps, washed in water, and stained as 182 above. Micrographs were taken at a range of magnifications using an upright microscope (DM 6000, 183 Leica, Wetzlar, Germany) equipped with a digital camera (SPOT Flex 64MP Color FireWire 15.2, 184 Diagnostic instruments.inc, USA) and analysed using ImageJ (Rasband, 1997 - 2014).

185

186 **Cryo- Scanning Electron Microscopy (cryo-SEM)**

PV curves were measured, and then leaf anatomical features, including the distribution of gas, liquid water, and cell structure, were visualized at different states of hydration using cryo – Scanning Electron Microscopy (cryo-SEM) according to McCully *et al.* (2009). One branch with four twigs similar in size was selected from each of three of the five study trees, then cut under perfusion solution (25 mM NaCl) and left for a few hours until fully hydrated. Then the four twigs, each with at least three pairs of fully expanded leaves, were cut under perfusion solution from each branch. One twig from each branch was sampled immediately upon removal from perfusion solution and the

- 194 remaining three twigs of each branch were then air-dried for differing periods of time before sampling
- 195 to capture change in leaf structure associated with key features of the previously measured PV
- 196 curves. At each sampling time, one leaf in each pair was used to measure leaf water potential while
- the other leaf was frozen rapidly (less than 0.1 s) with cryo-pliers cooled with liquid N₂ (LN2) (McCully
- 198 *et al.*, 2010). Frozen pieces (1 2 x 0.5 cm) were cut from the desired region under LN2, placed in
- 199 cryo-vials and stored in LN2 until processed for cryo-SEM. Then, frozen leaf segments were trimmed
- 200 to smaller pieces (3 mm length) under LN2, planed with a cryo-microtome (Leica EM FC7, Leica,
- 201 Wetzlar, Germany) and visualised by cryo-SEM (Hitachi 4300SE/N, Hitachi High Technologies,
- 202 Japan) following McCully *et al.* (2010).
- 203

204 Mucilage Analysis

- 205 Sample preparation
- Gel extruded from petioles of *A. marina* was collected and dispersed in 5% EtOH/water. The solution
 was centrifuged to remove insoluble extraneous matter and the supernatant freeze-dried.
- 208

209 Polysaccharide analysis (% composition)

- 210 The reductions of the constituent uronic acid methyl esters and the free uronic acids were carried out
- following the protocol of Kim and Carpita (1992) and of Pettolino *et al.* (2012). The reduced
- 212 polysaccharides were then hydrolysed, reduced, acetylated and subject to gas chromatography mass
- 213 spectrometry (GC/MS) analysis as described by Peng *et al.* (2000).
- 214

215 Water uptake by trichomes

216 Three twigs, each with a fully mature leaf pair, were harvested from three saplings of A. marina that 217 had been hydroponically grown (nutrient solution salinized with addition of 250 mM NaCI) from 218 propagules that originated from the same field populations used for PV curve analyses. Leaves were 219 rehydrated and then air-dried, with water potential determined at intervals during drying as described 220 above for PV curves. Following each measurement of leaf water potential, leaves were tested for 221 uptake of water by the trichome layer that covered the abaxial leaf surface. Upon removal from the 222 pressure chamber, a leaf was positioned to view the trichome layer under an epi-fluorescence 223 microscope (Zeiss Axiostar plus, Carl Zeiss, Germany). A drop of 0.1% fluorescein sodium salt

224 solution (British Drug Houses, Poole, England) was applied to the trichome surface and fluorescence 225 from the dye was observed by standard methods under blue-exciting light. A threshold for water 226 uptake by trichomes was determined by the highest leaf water potential at which dye applied to the 227 abaxial leaf surface disappeared into the trichomes. This threshold was used to estimate the onset of 228 drainage from trichomes as well-hydrated leaves were air-dried. 229 230 **Data analysis** 231 Data analyses using simple linear regression or one-way ANOVA were performed in Genstat version 232 16 (Payne, 2014). Data did not require transformation prior to analysis. Where relationships were 233 significant, differences between treatment means were tested post hoc for significance (here 234 considered P ≤ 0.05) using Fisher's Least Significant Difference test. Unless otherwise stated, all 235 results given in the text are mean values \pm standard error (se), n = 5. 236 237 Results 238 239 **PV curve analysis** 240 All measurements were made on leaves collected from mature trees that grew naturally where soil 241 water salinity at a depth of 30 cm averaged 49 ± 0.5 ppt (parts per thousand by weight) and surface 242 water at high tide averaged 35 ppt. For reference, standard seawater has a salinity of 35 ppt, and an 243 osmotic potential of -2.4 MPa. Thus the roots were exposed to soil and surface water with an osmotic 244 potential ranging from -3.4 to -2.4 MPa, respectively. However, surface water salinity can become 245 higher during low tide when drying concentrates salt at the soil surface. 246 247 Characteristics of the leaves used for PV curve measurements are summarised in Table 3. Fully expanded, sun leaves of A. marina had an average area of 14.8 ± 1.1 cm² and contained 0.66 ± 0.05 248 249 g of water when fully hydrated. Relative water content (RWC) was measured on these leaves as a 250 function of leaf water potential (Ψ_{leaf}) during air-drying. Variation in RWC with Ψ_{leaf} followed a 251 complicated relationship with three domains as shown in Fig. 1a. 252

253 Water relations parameters of A. marina leaves were calculated from plots relating water content to 254 water potential (commonly called PV curves) and are summarised in Table 4. Domain 1 of the curve 255 was characterised by rapid decline in RWC produced by relatively small application of pressure in the 256 chamber. On average, RWC declined 13 ± 1% in this domain, for which there was less than 1 MPa 257 decrease in the inferred leaf water potential. Water potential at the transition from the first to the 258 second domain (Ψ_x) averaged -0.85 ± 0.06 MPa (point **B**, Fig. 1a). Domain 2 covered the largest 259 variation in Ψ_{leaf} : a decrease from -0.85 to - 5.1 MPa, five times larger than that of domain 1. Over the 260 second domain, RWC decreased by an average of $12 \pm 3\%$, almost as much as that of domain 1. 261 Finally, domain 3 was characterized by more rapid rate of leaf water loss for a given decrease in leaf 262 water potential than the previous domain, domain 2. The transition between domains 2 and 3 (point C, 263 Fig.1a) averaged -5.1 \pm 0.1 MPa. Given the salinities of pore and surface water, Ψ_{leaf} at this transition was inferred to be the leaf water potential at turgor loss point (Ψ_{π}^{0}), which was supported by 264 265 calculations from the fitted straight line (dashes and points) in Fig. 1b as shown below.

266

267 The data set used in Fig. 1a was replotted in Fig. 1b with the reciprocal of Ψ_{leaf} , (|1/ Ψ_{leaf}) plotted as a 268 function of relative water content deficit, calculated with total leaf water content (RWD+, %). The 269 starting point, **c**, of the linear relationship between $|1/\Psi_{leaf}|$ and RWD+ indicated the reciprocal of leaf water potential at the turgor loss point ($|1/\Psi_{\pi}^{0}|$); further, the intercept **d**_x is given by the coordinates 270 RWD+ = RWD_x+, i.e. the percentage of water content decrease in domain 1, and $|1/\Psi_{leaf}| = |1/\Psi_{\pi}^{100}|$, 271 272 i.e. the reciprocal of the osmotic potential at full turgor. According to these calculations, leaf water 273 potential at the turgor loss point and osmotic potential at full turgor averaged -5.1 ± 0.1 MPa and -4.2 274 ± 0.1 MPa, respectively, in A. marina leaves. These calculations excluded extracellular water implied in domain 1. If that water was included, then Ψ_{π}^{100} would be calculated from the $1/\Psi_{\text{leaf}}$ intercept, **d**₀, 275 276 as shown in Fig. 1b and would average -3.5 ± 0.1 MPa.

277

The PV curves were replotted without domain 1, as suggested by Beckett (1997) and Dichio *et al.* (2003) to account for effects of extracellular water (the plateau effect) on calculation of leaf water relations parameters from the PV curve (Supplement S1). Calculation of Ψ_{π}^{0} was unaffected by removal of domain 1, but Ψ_{π}^{100} was changed because the method of Beckett (1997) and Dichio et al. (2003) redefined the status of a fully hydrated leaf. Specifically, calculation of maximum leaf water

283 content (WC_{max}) changed from (FM_{max} - DM) to (FM_{max} - WC_{ex} - DM), where FM_{max} is the maximum 284 leaf fresh mass, DM is the leaf dry mass, and WCex is the extracellular water content. As WCmax 285 decreased, RWC increased accordingly (see Table 2 for RWC calculation). Upon the removal of domain 1, Ψ_{π}^{100} was calculated at the $1/\Psi_{leaf}$ intercept of the linear regression between inverse leaf 286 water potential $(|1/\Psi_{leaf}|, MPa^{-1})$ and relative water deficit, calculated without extracellular water, 287 288 (RWD-, %), i.e. point d (Supplement S1b), and averaged -4.2 ± 0.1 MPa. This value was equal to that 289 calculated at d_x (-4.2 ± 0.1 MPa, P = 0.968) but significantly different from that calculated at d_0 (-3.5 ± 290 0.1 MPa, P < 0.001).

291

292 Calculation of water storage capacitance and bulk modulus of elasticity

293 The area-specific water storage capacitance (called water storage capacitance hereafter) was 294 calculated as the mass of water lost per unit area per unit decrease in water potential (Table 2). Two 295 values of water storage capacitance were calculated from different points on the PV curve (Table 4). 296 Water storage capacitance of domain 1 (Q_x) was calculated from points **A** to **B** (Fig. 1a) and the capacitance of domain 2 (Q_e) was calculated from points **B** to **C** (Fig. 1a). Q_x averaged 69 ± 6 g m⁻² 297 MPa⁻¹, and \mathbf{Q}_{e} averaged 13 ± 4 g m⁻² MPa⁻¹. In other words, application of a chamber pressure of 1.0 298 MPa during dehydration in domain 1 would release approximately 69 g water per m² leaf area 299 whereas similar application of 1.0 MPa in domain 2 would release about 13 g water per m² leaf area. 300 The average storage capacitance over both domains (Q_g , points A to C, Fig. 1a) was 22 ± 4.8 g m⁻² 301 MPa⁻¹. 302

303

```
304 The effective bulk modulus of elasticity, E, is defined as the change in chamber pressure or water
```

- 305 potential required for a fractional decrease in a cell's water volume ($\boldsymbol{\varepsilon} = \Delta P/(\Delta V/V)$). The
- 306 compressibility of water over this range of pressures is negligible, so $\boldsymbol{\xi} = \Delta P / \Delta RWC$. The values of $\boldsymbol{\xi}$

307 thus estimated (Table 4) averaged 7 ± 1 and 37 ± 10 MPa for domains 1 ($\boldsymbol{\xi}_{x}$) and 2 ($\boldsymbol{\xi}_{e}$), respectively,

308 with an average value, $\mathbf{\xi}_{g}$, of 21 ± 2 MPa calculated over both domains.

- 310 Water storage capacitance and elasticity calculated upon the removal of domain 1 averaged 13 ± 2 g
- m^{-2} MPa⁻¹ and 38 ± 4 MPa, respectively, and were not significantly different from those calculated
- from domain 2, i.e. Q_e and ε_e , (P = 0.383, and P = 0.841, respectively).
- 313

314 Leaf structure

315 Petiole

316 The basic features of petioles and leaves were characterised by bright field microscopy (Fig. 2). 317 Petioles (Fig. 2a) of A. marina leaves were covered with two types of multicellular trichomes. 318 Trichomes of the first type were clustered together with mucus secretion glands in a groove at the 319 petiole - stem junction. Trichomes of the second type covered the rest of the petiole and leaf abaxial 320 surfaces where the trichomes coexisted with salt secretion glands. These trichomes had a rivet-like 321 shape with a cap cell on top, one to two stalk cells in the middle, and a basal cell at the bottom. The 322 cap and stalk cells were dead with a hollow lumen. Trichomes of both types were transparent; 323 however, they appeared black if cyanobacteria were present.

324

325 The cortex was the largest tissue within the petiole and contained two distinct layers. A layer of 326 collenchyma (6 -7 cells thick) occurred beneath the epidermis, and was underlain by a layer of 327 parenchyma (15 - 16 cells thick). Substantial gas spaces occurred between parenchymal cells. The 328 gas spaces in the petiole cortex extended into the bundle sheath extension of the midvein but not to 329 the higher vein orders. The remaining space in the petiole was occupied by the two fimbrial veins and 330 the central vascular cylinder which contained a circular array of vascular bundles as commonly found 331 in eudicots. However, xylem vessels were not evenly distributed among the vascular bundles, with 332 more vessels occurring in bundles near the abaxial than adaxial sides of the petiole. Finally, the pith 333 occupied the most central space, surrounded by vascular tissue. The pith consisted mainly of tightly 334 packed parenchymal cells interspersed with small phloem bundles.

335

336 Leaf lamina

Leaves had a reticulate venation system with a prominent midvein as shown in Fig. 2b. A thick cuticle
 covered the adaxial leaf surface except where salt secretion glands occurred at the base of scattered
 depressions in the leaf surface. These salt secretion glands were in contact with underlying epidermal

and hypodermal cells. The midvein vascular structure was similar to that of the petiole. Nevertheless,
a ring of fibers surrounding the central vascular cylinder was more developed in the midvein than in
the petiole. In place of the cortex, the midvein was bounded by a bundle sheath extension (BSE) that
connected the vascular tissue with the upper and lower epidermis. Leaves of *A. marina* were
heterobaric but bundle sheath extensions only occurred at vein orders ranging from 1st to 4th. The
midvein was also linked with adjacent mesophyll and hypodermal tissues (Fig. 2c).

346

347 Four major tissues contributed most of the leaf lamina thickness (Table 3). The structure of A. marina 348 leaves was typical of C_3 species, with veins embedded between the palisade and spongy mesophyll 349 (Fig. 2c). Spongy mesophyll thickness was about two thirds that of palisade mesophyll. Together, they 350 accounted for 43 ± 1% of total leaf thickness. Avicennia marina leaves had an additional hypodermal 351 layer located below the upper epidermis, accounting for about 36 ± 1% of leaf thickness. Trichomes 352 on the abaxial leaf surface were similar in structure to the rivet-like trichomes on the petiole but 353 longer, and accounted for 19 ± 1% of total leaf thickness. Salt secretion glands occurred on both leaf 354 surfaces, but stomata were only on the abaxial surface.

355

356 **Change in leaf structures during drying**

357 Leaves were cryo-preserved during drying to fix the cell structure and the spatial distribution of gas 358 and liquid. Changes in these three components of leaf structure were assessed in relation to the three 359 domains of the PV curve using cryo-SEM. Domain 1 was characterized by the occurrence of 360 extracellular water in the gas spaces of the rivet-shaped trichomes, petiole, bundle sheath extension, 361 and leaf lamina. Extracellular water was not observed in samples taken in domain 2. Changes in cell 362 RWC could not be determined from the images because of variation in cell size and shape between 363 leaves. However, during domain 3, plasmolysis was apparent in most cells, but no cell wall collapse 364 was observed except in the trichomes.

365

Trichome water status changed with leaf water status. For reference, general trichome structure is
shown in Fig. 3a. Trichome lumina were full of liquid water when leaves were fully hydrated (Fig. 3b).
During domain 1, this extracellular water was replaced with gas (Fig. 3d) while the stalk cells of

trichomes retained their shape; however, the cap cells were collapsed (Fig. 3c, d). In domains 2 and
3, both cap and upper stalk cells were shrivelled and collapsed (Fig. 3e).

371

372 Additional measurements using fluorescence microscopy on fresh materials demonstrated the onset 373 of trichome drainage with dehydration (Fig. 4). When viewed under white light the cap cells of the 374 trichomes formed a continuous surface (Fig. 4a). The hollow stalk cells appeared like dark circular 375 areas centred beneath the cap cells. No structure was visible under blue light showing the absence of 376 detectable auto-fluorescence (Fig. 4b). In contrast, the presence of fluorescein was detected on the 377 leaf surface by bright green fluorescence under blue light (Fig. 4c - f). When trichomes were filled with 378 water ($\Psi_{\text{leaf}} > 0.1 \text{ MPa}$), fluorescein quickly spread over the leaf surface without loss in fluorescence 379 (Fig. 4c, d and movie 1(Supplement S2)). Drainage of water-filled trichomes was detected when Ψ_{leaf} 380 declined to -0.25 ± 0.03 MPa (n = 3). Under these conditions, fluorescein drops remained where 381 applied for approximately a minute before rapidly disappearing into the trichomes (Fig. 4e, f, and 382 movie 2 (Supplement S3)). The estimated density and volume of trichomes (Table 3) indicated that 383 they could potentially hold up to 10% of total leaf water.

384

385 Petiolar gas spaces also functioned in temporary water storage of well-hydrated leaves, and, 386 therefore, were named cisternae. Due to the complex three-dimensional structure, it was not possible 387 to obtain a reliable estimate of water storage in cisternae from the micrographs. When viewed by 388 bright-field microscopy, longitudinal sections through the petiole revealed greater surface structure 389 delimiting these cisternae than expected (Fig. 5a, b). Closer inspection under cryo-SEM revealed that 390 the walls of these cisternae had a rough surface that appeared coated with a mucus-like substance 391 that formed globular structure, hereafter called droplets, with drying. Analysis of a gel extruded 392 naturally from petioles revealed a composition consistent with mucilaginous polysaccharides (Table 393 5). When leaves were well hydrated ($\Psi_{\text{leaf}} = -0.1 \text{ MPa}$), most cisternae were full of water but some 394 contained large droplets with diameters averaging $6.8 \pm 0.8 \mu m$ (n = 20) (Fig. 5c, d). Where the 395 planing knife cut through the ice of a water-filled cistern, the eutectic domains of the ice indicated 396 water of high solute content. At the end of domain 1, no cisternae were filled with water and droplet 397 diameter averaged 4.9 \pm 0.3 μ m (n = 20) (Fig. 5e), smaller than that observed at higher leaf water

398 potentials. As leaf water content decreased over domain 2, very few droplets were found and their 399 average diameter was $2.2 \pm 0.3 \mu m$ (n = 20) (Fig. 5f).

400

401 In the leaf lamina and midvein, small gas spaces between pith parenchyma and between mesophyll 402 cells were also filled with liquid water but only when leaf water potential was less negative than -0.1 403 MPa (Fig. 6). Where the planing knife cut through the ice of a water-filled intercellular gas space, the 404 general absence of eutectic domains in the ice indicated water of very low solute content, in contrast 405 to that of the cisternae. The boundary of the lamina spaces was not specially coated like that of the 406 cisternae. No gas space or external water was found between hypodermal cells. Due to variability in 407 cell sizes within and between leaves, it was not possible to quantify changes in size of the 408 hypodermal cells over domains 1 and 2. 409 410 In domain 3, a wide variety of cell types, including collenchyma and parenchyma cells in the petiole 411 and the hypodermal, mesophyll cells, and epidermal cells in the leaf lamina, showed similar 412 responses to decreases in leaf water potential below the turgor loss point. Gaps appeared between 413 cell walls and the enclosed cells, as is characteristic of plasmolysis (Fig. 7). 414 415 416 Discussion 417 The results of the present study revealed a new twist in an old curve and novel insights into the 418 interpretation of the PV curve in Avicennia marina. This PV curve exhibited three domains over the 419 range of hydration in which the leaves naturally function under field conditions, even in a single day 420 (Constable, 2014, Walker, 2014). The three domains of leaf dehydration were characterized by (1) the 421 presence and consumption of extracellular water, (2) decline in turgor, and (3) the occurrence of 422 plasmolysis. Each domain, as illustrated in the summary diagram (Fig. 8), and its implications for leaf 423 function, are discussed in detail below.

424

425 **Domain 1: dominated by variation in extracellular water**

426 The domain of highest hydration was characterised by the presence of extracellular water and its

427 disappearance as RWC decreased by 13% while leaf water potential only declined from -0.1 to -0.85

MPa (Fig. 1a, Fig. 8). At the highest hydration level, water filled extracellular spaces, including those between mesophyll cells (Fig. 6c), the hollow cores of trichomes (Fig. 3b), and the petiolar cisternae (Fig. 5c). As the leaf dried, water was replaced by gas in these distinct spaces (Figs. 3d, 5e, f, 6d) over different ranges of water potentials, implying that filling of the spaces occurred under different conditions and depended on different processes.

433

434 Liquid water was only observed in some mesophyll extracellular gas spaces when water potential was 435 less negative than -0.1 MPa. Water storage between mesophyll cells would have a negative impact 436 on gas exchange because diffusion of CO₂ is about 10,000 times slower in water than in air, but the 437 extent of this storage was small and short-lived. The lack of eutectic domains in this extracellular 438 water that had been cryo-preserved indicated that the water had very low solute concentrations. How 439 can we account for movement of liquid water from cells to extracellular gas spaces in the absence of 440 an osmotic gradient? The answer may be similar to processes driving refilling of embolized 441 protoxylem vessels in well-hydrated tissues (Rolland et al., 2015). Once the cells had achieved high 442 water potentials, capillary forces due to the combination of hydrophilic walls and close contact 443 between cells may have driven extracellular water accumulation in confined spaces. This limited filling 444 of extracellular gas spaces in excised leaves was not an artefact as similar observations have been 445 made on A. marina leaves under natural conditions when they were collected predawn following 446 exposure to nocturnal wetting events (Constable, 2014). Such accumulation of extracellular water in 447 these well-hydrated leaves presumably occurred through foliar absorption of atmospheric water, even 448 though roots grew in seawater; in other words, the water potential gradient during leaf wetting events 449 was reversed from the atmosphere to the leaves to the soil.

450

The presence of low solute concentrations in the extracellular water has important implications for salt tolerance of halophytes. During symplastic water uptake at the roots (Moon *et al.*, 1986), *A. marina* excludes approximately 90 - 95% of salt in soil water from entry to the transpiration stream (Ball, 1988). When grown in seawater, *A. marina* typically has 25 mM NaCl in the transpiration stream (Ball, 1988, Stuart *et al.*, 2007). In *A. marina*, water and associated salt are distributed across a leaf lamina through the vascular system to vein endings and subsequently dispersed to client cells by symplastic pathways (Fitzgerald & Allaway, 1991). This symplastic control prevents catastrophic salt

accumulation in apoplastic spaces and facilitates salt management within a leaf. In *A. marina*, part of
the salt transported in the transpiration stream is accommodated in growing cells for osmotic
adjustment of the vacuoles while the remainder is exported from the leaves via either the phloem or
epidermal salt secretion glands, thereby maintaining a favourable salt balance (Ball, 1988). Thus the
observation of low solute concentrations in the extracellular water (Fig 5d) is consistent with strong
symplastic control of both water and associated salt distribution within the leaves (Fitzgerald &
Allaway, 1991).

465

466 Water-filled trichomes began to empty when Ψ_{leaf} declined below -0.25 MPa (Fig. 4e, f and movie 2 467 (Supplement S3)). In contrast to intercellular spaces in the mesophyll, the trichome lumina appeared 468 to contain a mucilaginous substance. This was suggested by the similarity of the ice texture in water-469 filled trichomes that had been cryo-preserved (Fig 1b) to that observed in cryo-preserved 470 mucilaginous secretions of root tips (McCully et al., 2009). Such a hydrophilic substance in 471 combination with the highly hydrophilic walls and the small diameter of the hairs could have facilitated 472 water uptake at relatively high water potentials, as might occur with exposure to rain or dew. Indeed, 473 under natural field conditions, Constable (2014) observed trichomes to be full of water when leaves of 474 A. marina were wet in the morning. Trichomes seemed to serve as one of the main water storage 475 sites, which could hold up to 10% of total leaf RWC. This storage water could be released to the leaf 476 as leaf water potentials declined with dehydration if water transport into the leaf followed a symplastic 477 pathway through the living basal cells of the trichomes to the underlying cells (Fig. 3a). Similarly, 478 water absorbed by trichomes contributed to the water status of underlying leaf cells in oak leaves 479 (Fernández et al., 2014). Thus, identification of trichomes as a major water storage site in A. marina 480 also implies that the trichomes may provide a means for rehydration via foliar water uptake.

481

Finally, the present study identified a novel water storage structure, here named the cisternae, which occurred mainly in the petiole and extended into the bundle sheath extension of the midvein near its junction with the petiole. Extracellular water occurred in cisternae until leaf water potentials declined to approximately -0.8 MPa. Unlike the extracellular spaces in the mesophyll and the trichomes, the cisternae were relatively large, completely bounded by the walls of cells comprising the surrounding tissues (Fig. 5a,b), and appeared to contain a combination of mucilage, mostly arabinose, and a

attract water from the surrounding cells into the cisternae when leaf water potentials rose above -0.8
MPa. Extension of the cisternae into BSE of the midvein would enhance the reach of their influence
on leaf water relations as the BSEs are known to play a role in extravascular water dispersal (Canny,
1986, Buckley *et al.*, 2011, Sommerville *et al.*, 2012, Zsogon *et al.*, 2015). Conversely, because the
osmotic potential of the cisternae would vary with water content, differential drainage of the cisternae

relatively high concentration of solutes (Table 5, Fig. 5d, f). The latter must have been sufficient to

494 during leaf drying could contribute to the curvature linking domains 1 and 2 in the PV curve.

495

488

496 **Domain 2: Dominated by variable turgor**

The second domain was characterized by 12% loss in leaf RWC as leaf water potentials declined from -0.85 to -5.1 MPa. Decrease in cell volume, except for that in trichomes, was not obvious in our micrographs, possibly due to the small scale of total loss in RWC (i.e. 12% across domain 2, Fig. 1a), combined with high variability in cell sizes (Fig. 8). For example, a 12% change in leaf water content would be equivalent to a 4% change in each dimension of an isotropic cell.

502

503 Moreover, there were no obvious changes in gas spaces between mesophyll cells with decreasing 504 Ψ_{leaf} , consistent with observations of desiccating sunflower leaves (Fellows & Boyer, 1978). Tissue 505 connections in the leaf lamina were maintained as cell walls and membranes also maintained contact 506 during shrinkage (i.e. cytorrhysis). Given the large osmotic potential at full turgor ($\Psi_{\pi}^{100} = -4.2 \text{ MPa}$) 507 and the high modulus of elasticity, most of the decline in leaf water potential with decrease in RWC in 508 the second domain was associated with decreasing turgor pressure.

509

510 The curvature between domains 2 and 3 may reflect differential turgor loss in different cell types with 511 the turgor loss point representing the extreme case where all cells have lost turgor (Cheung et al., 512 1975). Some cells might lose water more rapidly than others due to their properties or their positions 513 within the leaf. In addition, **£** is likely to be itself a function of turgor. For example, the high and 514 constant E region could represent cell wall stretching, and the lower and variable modulus region 515 could represent change in shape. Thus, the curvature shown linking domains 2 and 3 (Fig. 1) likely 516 reflects the large contribution of different cell types to water relations in A. marina. Similar curvature 517 was found in many studies where PV curves were constructed from repeated measurements on a

single leaf or shoot (Cheung *et al.*, 1975, Meinzer *et al.*, 1986, Parker & Pallardy, 1987). In contrast, a
sharp transition between domains 1 and 2 has been observed when the PV curve is constructed from
a composite of measurements made on many leaves of a species, such as those from woody tropical
rainforest (Brodribb & Holbrook, 2003) or used in theoretical analyses, e.g. Suarez and Sobrado
(2000). The convexity of the transition may make a useful tool for comparing hydraulic complexity in
leaves grown under different conditions or between different species.

524

525 **Domain 3: Dominated by osmotic behaviour of flaccid cells**

526 Domain 3 was characterised by plasmolysis (Figs. 7, 8) as leaf water potentials declined below the 527 turgor loss point once RWC decreased below approximately 75% (or below about 86% if the 528 extracellular water was not included in the accounting) (Table 4). In most living cell types, connections 529 between tissues or between cells comprising a tissue were maintained during desiccation-induced 530 shrinkage. No collapse of cell walls was observed. However, separation between cell walls and plasma membranes developed when Ψ_{leaf} was about 1 MPa lower than Ψ_{π}^{0} . Plasmolysis in *A. marina* 531 532 was also noticed in the field when Ψ_{leaf} was as low as -6 MPa, and the cells recovered following 533 nocturnal rehydration (Constable, 2014). The fact that A. marina leaves could recover after reaching 534 the turgor loss point was consistent with a study by Brodribb and Holbrook (2003) which showed that 535 irreversible damage to PSII system only occurred in woody tropical rainforest species at Ψ_{leaf} much lower than Ψ_{π}^{0} . 536

537

538 Ecophysiological implications of the new PV curve to *A. marina* function

The results of the current study emphasized the importance of leaf anatomy to leaf water relations with far reaching implications for leaf function under field conditions. Previous studies have referred to the first domain as "the plateau effect" (Parker & Pallardy, 1987, Dichio *et al.*, 2003), attributed to artefactual rehydration. The presence of domain 1 due to extracellular water does not change values of PV curve parameters, including Ψ_{π}^{0} , Ψ_{π}^{100} , $\boldsymbol{\epsilon}$, and \mathbf{Q} , as long as those parameters are calculated using data from appropriate domains as shown in Fig. 1 and indicated in Table 4, consistent with recommendations to account for "the plateau effect" (Parker & Pallardy, 1987). However, dismissal of

546 domain 1 as an artefact of rehydration would lead to false assumptions about leaf functions in *A*.

547 *marina* that are inconsistent with field observations and leaf anatomy.

548

The PV curves revealed a high modulus of elasticity in leaves of *A. marina*, which plays a major role in cell function over the whole range of water potentials naturally experienced by these leaves. At maximum hydration, leaf water potentials were very high (i.e. -0.1 MPa) while osmotic potential at full turgor was -4.2 MPa (Fig. 1b). The turgor pressure would therefore have been around 4.1 MPa, but there was no evidence that cells burst under these conditions. Presumably, the high modulus of elasticity, i.e. 37 MPa (Table 4), contributed to cell survival when the leaves were fully hydrated.

556 Previous studies have interpreted the role of a high **E** on leaf function under low water potentials in 557 two ways. First, high E was proposed to enhance water uptake into a transpiring leaf by increasing 558 the water potential gradient between the leaf and the soil (Bowman & Roberts, 1985). However, 559 Bartlett et al. (2012) argued that this proposal disregards the role of hydraulic conductance in 560 determining the water potential gradient. Indeed, studies of diurnal gas exchange in the same 561 population of A. marina as the present study showed that reduction in leaf hydraulic conductance 562 began when leaf water potentials declined below -3 MPa (Walker, 2014). Consequently, decline in 563 Ψ_{leaf} with decreasing turgor would not necessarily enhance water uptake into leaves as they dried. 564 Indeed, a low E would allow the concentration of the cytoplasm to vary significantly over the range of 565 positive turgor, whereas a high E would keep the cytoplasmic concentrations similar, and also reduce 566 the possibly deleterious effects of changes in cell geometry. Thus, we agree with the second 567 suggestion that a high modulus of elasticity coupled with a low solute potential enables leaves to 568 maintain the levels of hydration required for cellular function during drying to the turgor loss point 569 (Cheung et al., 1975, Bartlett et al., 2012).

570

571 The extent of water storage in domains 1 and 2 revealed by PV curve could play an important role in 572 the maintenance of diurnal photosynthetic activities in leaves of *A. marina*. Specifically, extracellular 573 water storage capacitance of domain 1 ($\mathbf{Q}_{\mathbf{x}}$) was about five times greater than the intracellular water 574 storage capacitance of domain 2 ($\mathbf{Q}_{\mathbf{e}}$). However, $\mathbf{Q}_{\mathbf{x}}$ was functional over a range of Ψ_{leaf} approximately 575 five times less than that of Qe. Consequently, extra and intracellular water storage was about equal. In 576 other words, storage of water in extracellular spaces doubled the amount of water available for use in gas exchange without loss in turgor. For example, area-specific evaporation rates measured for A. 577 marina grown at the same location as those in the present study ranged from 1 to 2 mmol H₂O m⁻² s⁻¹ 578 (Martin *et al.*, 2010). So a leaf with extracellular water storage capacitance of 69 g $H_2O \text{ m}^{-2} \text{ MPa}^{-1}$ 579 (Table 4) or 3800 mM H₂O m⁻² MPa⁻¹ as estimated from domain 1, when fully charged, could alone 580 581 supply the water loss needed to support photosynthesis at the above evaporation rates for 3230 to 582 1615 s, that is for 54 to 27 min with only 0.85 MPa decrease in Ψ_{leaf} . Once extracellular water is exhausted, then the water storage capacitance of domain 2 (i.e. 13 g H_2O m⁻² MPa⁻¹ or 700 mM H_2O 583 m⁻² MPa⁻¹) could alone support the same rates of gas exchange for another 50 to 25 min until the 584 585 turgor loss point (i.e. -5.1 MPa) is reached. Given the relative proportions of space occupied by 586 different tissue layers in the leaf lamina (Table 3), the hypodermis alone could have buffered water 587 loss from the mesophyll by contributing almost half of the water available for photosynthesis during 588 dehydration over domain 2 (all in the absence of water flux from the roots). Thus, water stored within 589 a leaf could supplement that provided by the roots, thereby extending the duration of gas exchange 590 activities for perhaps about two hours longer than otherwise, which could be critically important for a 591 species when soil salinity constrains the supply of water to the leaves.

592

593 Acknowledgments

594 HTN was supported by an Australia Awards PhD scholarship and the research was supported by 595 Australian Research Council Discovery Grant (DP150104437) to MCB and MM; PM was supported by 596 ARC grant FT110100457. We thank New South Wales Department of Primary Industries for 597 permission to work in the Clyde River mangrove system (S37 Permit P13/0066-1.0), the Centre for 598 Advanced Microscopy (Australian National University) for access to bright field microscopes and 599 assistance with cryo-SEM, and Dr. Charles Hocart (RSB Mass Spectrometry Facility, ANU) for 600 mucilage analysis. We thank the two anonymous reviewers for their insightful comments and 601 Professor John Evans (ANU) and Dr. Rafael Coopman (Universidad Austral de Chile) for their 602 thoughtful comments on the diagrammatic figure.

603

605 References

- Allen C.D., Breshears D.D. & McDowell N.G. (2015) On underestimation of global vulnerability to tree
 mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere*, 6, 1-55.
- 608Anderegg W.R.L., Berry J.A., Smith D.D., Sperry J.S., Anderegg L.D.L. & Field C.B. (2012) The roles609of hydraulic and carbon stress in a widespread climate-induced forest die-off. *Proceedings of*

610 the National Academy of Sciences, **109**, 233-237.

- Ball M.C. (1988) Salinity tolerance in the mangroves *Aegiceras corniculatum* and *Avicennia marina* I.
 Water use in relation to growth, carbon partitioning, and salt balance. *Australian Journal of Plant Physiology*, **15**, 447-464.
- Bartlett M.K., Scoffoni C. & Sack L. (2012) The determinants of leaf turgor loss point and prediction of
 drought tolerance of species and biomes: a global meta-analysis. *Ecology Letters*, **15**, 393405.
- Beckett R.P. (1997) Pressure–volume analysis of a range of poikilohydric plants implies the existence
 of negative turgor in vegetative cells. *Annals of Botany*, **79**, 145-152.
- 619 Bowman W.D. & Roberts S.W. (1985) Seasonal changes in tissue elasticity in chaparral shrubs.

620 *Physiologia Plantarum*, **65**, 233-236.

- Brodribb T.J. & Holbrook N.M. (2003) Stomatal closure during leaf dehydration, correlation with other
 leaf physiological traits. *Plant Physiology*, **132**, 2166-2173.
- Buckley T.N., Sack L. & Gilbert M.E. (2011) The role of bundle sheath extensions and life form in
 stomatal responses to leaf water status. *Plant Physiology*, **156**, 962-973.
- 625 Canny M.J. (1986) Water pathways in wheat leaves. III. The passage of the mestome sheath and the
 626 function of the suberised lamellae. *Physiologia Plantarum*, **66**, 637-647.
- 627 Canny M.J. (1995) Apoplastic water and solute movement: new rules for an old space. *Annual* 628 *Review of Plant Physiology and Plant Molecular Biology*, **46**, 215-236.

629 Cheung Y.N.S., Tyree M.T. & Dainty J. (1975) Water relations parameters on single leaves obtained

- 630 in a pressure bomb and some ecological interpretations. *Canadian Journal of Botany*, **53**,
 631 1342-1346.
- 632 Choat B., Lahr E.C., Melcher P.J., Zwieniecki M.A. & Holbrook N.M. (2005) The spatial pattern of air
 633 seeding thresholds in mature sugar maple trees. *Plant, Cell and Environment*, **28**, 1082-1089.

- Constable L. (2014) Nocturnal top-down rehydration: contribution of atmospheric moisture to water
 status of the mangrove, Avicennia marina. Honours Thesis, The Australian National
 University.
- 637 Dichio B., Xiloyannis C., Angelopoulos K., Nuzzo V., Bufo S.A. & Celano G. (2003) Drought-induced
 638 variations of water relations parameters in *Olea europaea*. *Plant and Soil*, **257**, 381-389.
- Fellows R.J. & Boyer J.S. (1978) Altered ultrastructure of cells of sunflower leaves having low water
 potentials. *Protoplasma*, 93, 381-395.
- Fernández V., Sancho-Knapik D., Guzmán P., Peguero-Pina J.J., Gil L., Karabourniotis G., . . . GilPelegrín E. (2014) Wettability, polarity, and water absorption of holm oak leaves: effect of leaf
 side and age. *Plant Physiology*, **166**, 168-180.
- Fitzgerald M.A. & Allaway W.G. (1991) Apoplastic and symplastic pathways in the leaf of the grey
 mangrove *Avicennia marina* (Forsk.) Vierh. *New Phytologist*, **119**, 217-226.
- Hajek T. & Beckett R.P. (2008) Effect of water content components on desiccation and recovery in
 Sphagnum mosses. *Annals of Botany*, **101**, 165-173.
- 648 Hutchings P. & Saenger P. (1987) *Ecology of mangrove*. University of Queensland Press.
- Kim J.B. & Carpita N.C. (1992) Changes in esterification of the uronic acid groups of cell wall
 polysaccharides during elongation of maize coleoptiles. *Plant Physiology*, **98**, 646-653.
- Kubiske M.E. & Abrams M.D. (1991) Seasonal, diurnal and rehydration-induced variation of pressurevolume relationships in *Pseudotsuga menziesii*. *Physiologia Plantarum*, **83**, 107-116.
- Lovelock C.E., Ball M.C., Martin K.C. & Feller I.C. (2009) Nutrient enrichment increases mortality of
 mangroves. *PLoS One*, 4, e5600.
- Martin K.C., Bruhn D., Lovelock C.E., Feller I.C., Evans J.R. & Ball M.C. (2010) Nitrogen fertilization
 enhances water-use efficiency in a saline environment. *Plant, Cell and Environment*, **33**, 344357.
- 658 McCully M.E., Canny M.J. & Huang C.X. (2009) Cryo-scanning electron microscopy (CSEM) in the
- advancement of functional plant biology. Morphological and anatomical applications. *Functional Plant Biology*, **36**, 97-124.
- McCully M.E., Canny M.J., Huang C.X., Miller C. & Brink F. (2010) Cryo-scanning electron
- 662 microscopy (CSEM) in the advancement of functional plant biology: energy dispersive X-ray
- 663 microanalysis (CEDX) applications. *Functional Plant Biology*, **37**, 1011-1040.

- McKee K.L. (1993) Soil physicochemical patterns and mangrove species distribution--reciprocal
 effects? *Journal of Ecology*, **81**, 477-487.
- Meinzer F.C., Rundel P.W., Sharifi M.R. & Nilsen E.T. (1986) Turgor and osmotic relations of the
 desert shrub Larrea tridentata. *Plant, Cell and Environment*, 9, 467-475.
- 668 Melkonian J.J., Wolfe J. & Steponkus P.L. (1982) Determination of the volumetric modulus of
- elasticity of wheat leaves by pressure-volume relations and the effect of drought conditioning. *Crop Science*, **22**, 116-123.
- Moon G., Clough B., Peterson C. & Allaway W. (1986) Apoplastic and symplastic pathways in *Avicennia marina* (Forsk.) Vierh. Roots revealed by fluorescent tracer dyes. *Functional Plant Biology*, 13, 637-648.
- Naidoo G., Hiralal O. & Naidoo Y. (2011) Hypersalinity effects on leaf ultrastructure and physiology in
 the mangrove *Avicennia marina*. *Flora*, **206**, 814-820.
- Nguyen H.T., Stanton D.E., Schmitz N., Farquhar G.D. & Ball M.C. (2015) Growth responses of the
 mangrove *Avicennia marina* to salinity: development and function of shoot hydraulic systems
 require saline conditions. *Annals of Botany*, **115**, 397-407.
- Parker W.C. & Pallardy S.G. (1987) The influence of resaturation method and tissue type on
 pressure- volume analysis of *Quercus alba* L. seedlings. *Journal of Experimental Botany*, 38,
 535-549.
- Payne R.W., Murray, D.A., Harding, S.A., Baird, D.B. & Soutar, D.M. (2014) GenStat for Windows
 16th Edition. VSN International, Hemel Hempstead, UK.
- 684 Peng L., Hocart C.H., Redmond J.W. & Williamson R.E. (2000) Fractionation of carbohydrates in
- 685 Arabidopsis root cell walls shows that three radial swelling loci are specifically involved in 686 cellulose production. *Planta*, **211**, 406-414.
- 687 Pettolino F.A., Walsh C., Fincher G.B. & Bacic A. (2012) Determining the polysaccharide composition
 688 of plant cell walls. *Nature Protocols*, **7**, 1590-1607.
- Rasband W.S. (1997 2014) ImageJ, U. S. National Institutes of Health, Bethesda, Maryland, USA.
- Reichstein M., Bahn M., Ciais P., Frank D., Mahecha M.D., Seneviratne S.I., ... Wattenbach M.
- 691 (2013) Climate extremes and the carbon cycle. *Nature*, **500**, 287-295.

- Rolland V., Bergstrom D.M., Lenne T., Bryant G., Chen H., Wolfe J., . . . Ball M.C. (2015) Easy come,
 easy go: capillary forces enable rapid refilling of embolized primary xylem vessels. *Plant Physiology*, **168**, 1636-1647.
- Rowland L., da Costa A.C.L., Galbraith D.R., Oliveira R.S., Binks O.J., Oliveira A.A.R., ... Meir P.
- 696 (2015) Death from drought in tropical forests is triggered by hydraulics not carbon starvation.
 697 *Nature*, **528**, 119-122.
- Sack L. & Scoffoni C. (2013) Leaf venation: structure, function, development, evolution, ecology and
 applications in the past, present and future. *New Phytologist*, **198**, 983-1000.
- Scholander P.F., Bradstreet E.D., Hemmingsen E.A. & Hammel H.T. (1965) Sap pressure in vascular
 plants: Negative hydrostatic pressure can be measured in plants. *Science*, **148**, 339-346.
- Scholander P.F., Hammel H.T., Hemmingsen E.A. & Bradstreet E.D. (1964) Hydrostatic pressure and
 osmotic potential in leaves of mangroves and some other plants. *Proceedings of the National Academy of Sciences*, 52, 119-125.
- Sommerville K.E., Sack L. & Ball M.C. (2012) Hydraulic conductance of Acacia phyllodes (foliage) is
 driven by primary nerve (vein) conductance and density. *Plant, Cell and Environment*, 35,
 158-168.
- Stuart S.A., Choat B., Martin K.C., Holbrook N.M. & Ball M.C. (2007) The role of freezing in setting the
 latitudinal limits of mangrove forests. *New Phytologist*, **173**, 576-583.
- 710 Suarez N. & Sobrado M.A. (2000) Adjustments in leaf water relations of mangrove (*Avicennia*

711 *germinans*) seedlings grown in a salinity gradient. *Tree Physiology*, **20**, 277-282.

- Turner N. (1988) Measurement of plant water status by the pressure chamber technique. *Irrigation Science*, 9, 289-308.
- Tyree M.T. & Ewers F.W. (1991) The hydraulic architecture of trees and other woody plants. *New Phytologist*, **119**, 345-360.
- Tyree M.T. & Hammel H.T. (1972) The measurement of the turgor pressure and the water relations of
- plants by the Pressure-bomb technique. *Journal of Experimental Botany*, **23**, 267-282.
- 718 Walker W. (2014) The role of leaf hydraulics in the photosynthetic performance of the mangrove
- 719 Avicennia marina growing in seawater and hyper-saline habitats. Honours Thesis, The

720 Australian National University.

721 Zimmermann M.H. (1983) *Xylem structure and the ascent of Sap*. Springer-Verlag.

- 722 Zsogon A., Negrini A.C., Peres L.E., Nguyen H.T. & Ball M.C. (2015) A mutation that eliminates
- 523 bundle sheath extensions reduces leaf hydraulic conductance, stomatal conductance and
- assimilation rates in tomato (*Solanum lycopersicum*). *New Phytologist*, **205**, 618-626.

Table 1. Abbreviations

Parameter	Symbol	Unit
Calculation with extracellular water included	+	
Calculation without extracellular water	-	
Water mass difference	Δw	q
Leaf water potential gradient	ΔΨ	MPa
Leaf dry mass	DM	g
Leaf fresh mass	FM	g
Maximum leaf fresh mass	FM _{max}	g
Water storage capacitance	Q	g MP ⁻¹ m ⁻²
Water storage capacitance at domain 2	Q	g MP⁻¹m⁻²
General water storage capacitance	Qg	g MP ⁻¹ m ⁻²
Water storage capacitance at domain 1	Q _x	g MP ⁻¹ m ⁻²
Relative water content	RWC	%
RWC at turgor loss point	RWC _0	%
RWC at full turgor	RWC 100	%
RWC at the transition between domains 1 and 2	RWC _x	%
Relative water deficit	RWD	%
RWD at turgor loss point	RWD _0	%
RWD during domain 2	RWDe	%
RWD during domain 1	RWDx	%
Leafarea	S	cm ² , m ²
Maximum leaf water content	WC _{max}	g
Extracellular water content	WC _x	g
Pressure gradient	ΔP	MPa
RWC difference	ΔRWC	%
Fractional decrease in a cell's water volume	$\Delta V/V$	
Volumetric modulus of elasticity	٤	MPa
Bulk modulus of elasticity at domain 2	٤e	MPa
General bulk modulus of elasticity	٤g	MPa
Bulk modulus of elasticity at domain 1	٤x	MPa
Leaf water potential	Ψ_{leaf}	MPa
Leaf water potential at the transition between domains 1 and 2	Ψx	MPa
Leaf water potential at full hydration	Ψ_{100}	MPa
Leaf water potential at turgor loss point	Ψ_{π^0}	MPa
Osmotic potential at full turgor	$\Psi_{\pi}{}^{100}$	MPa

Parameter	Symbol	Unit	Calculation
Water storage capacitance	Q	g MP ⁻¹ m ⁻²	$Q = \frac{\Delta w - 1}{\Delta \Psi - S}$
			$=\frac{\Delta RWC. WC_{max} 1}{\Delta \Psi S}$
Bulk modulus of elasticity	٤	MPa	$\boldsymbol{\varepsilon} = \frac{\Delta P}{\frac{\Delta V}{V}} = \frac{\Delta \Psi}{\Delta RWC}$
Maximum water content	WC _{max}	g	WC _{max} = FM _{max} - DM
Relative water content	RWC	%	RWC = (FM - DM)/WC _{max}
Leaf water potential gradient	ΔΨ	MPa	$\Delta \Psi = \Psi_{t p} - \Psi_{100} $
Relative water content deficit	RWD	%	RWD = 100 - RWC

Table 2. Summary of PV curve components and their calculations as previously described(Scholander *et al.*, 1964, Turner, 1988, Tyree & Ewers, 1991, Bartlett *et al.*, 2012).

Parameter	Symbol	Unit Mean		se (n = 5)	
Leaf area	S	cm ²	14.8	1.1	
Maximum fresh	mass	FM _{max}	g	1.04	0.07
Dry mass	DM	g	0.38	0.02	
Maximum water	WC _{max}	g	0.66	0.05	
Leaf mass per a	LMA	a m ⁻²	256.3	13.9	
Maximum water	WC _{max} DM ⁻¹	g g⁻¹	1.76	0.08	
Leafthickness		μm	588	10	
	Upper epidermis		%	3	0
	Hypodermis		%	36	1
Contribution to	Palisade mesophyll		%	30	1
	Spongy mesophyll		%	13	0
	Trichome		%	19	1
Trichome	Trichome Density		cm ⁻²	218,758	8,783
	Volume*		μm³	20,460	3,288 (n = 15)

Table 3. Properties of fully expanded, field-grown, sun leaves of *A. marina* used for measurements of PV curves and anatomical analyses. Values are mean \pm se (n = 5, except for * where n = 15)

Table 4. Leaf water relations of *A. marina* derived from three – domain PV curves as shown in Fig. 1. Values are mean \pm se, n = 5. Note that symbols + or – denote whether extracellular water was included or excluded, respectively, in the calculations.

Parameter	Parameter Symbol			se (n = 5)
Relatively water deficit during domain 1	RWD _x +	%	13	1
Leaf water potential at RWC _x +	Ψx	MPa	-0.85	-0.03
Relative water deficit during domain 2	RWD _e +	%	12	1
Osmotic potential at full turgor	Ψ_{π}^{100}	MPa	-4.2	-0.1
Water potential at turgor loss point	Vater potential at turgor loss point u_0			
RWC at full turgor	BWC ¹⁰⁰	%	87	1
RWC at turgor loss point, including extracellular water	RWC ⁰ +	%	75	3
RWC at turgor loss point, excluding extracellular water	RWC _π ⁰-	%	86	2
General bulk modulus of elasticity	٤g	MPa	21	2
Modulus of elasticity for domain 1	٤e	MPa	7	1
Modulus of elasticity for domain 2	٤x	MPa	37	4
General water capacitance	Qg	g m ⁻² MPa ⁻¹	22	2
Water storage capacitance for domain 1	Q _x	g m ⁻² MPa ⁻¹	69	6
Water storage capacitance for domain 2	Qe	g m ⁻² MPa ⁻¹	13	2

Table 5. Composition of A. marina petiolar gel. Values are mean percent dry mass of the total drymass of the polysaccharides named in the table below \pm se, n = 3. Abbreviations: Rha: Rhamnose,Ara: Arabinose, Xyl: Xylose, Man: Mannose, ManUA: Mannuronic acid, Gal: Galactose, GalUA:Galacturonic acid, Glc: Glucose, GlcUA: Glucuronic acid. A: esterified uronic acid analysis, B:unesterified uronic acid analysis. NA: not applicable.

	Rha	Ara	Xyl	Man	ManUA	Gal	GalUA	Glc	GIcUA
A (%)	2.07	83.47	0.97	0.23	NA	10.60	NA	2.67	NA
se	0.19	0.32	0.09	0.03	NA	0.29	NA	0.22	NA
B (%)	2.23	83.17	0.80	0.23	0.00	10.53	0.00	0.53	2.50
se	0.19	0.50	0.06	0.03	NA	0.20	NA	0.15	0.15

Figure legends:

Fig. 1. A representative pressure volume (PV) curve of an Avicennia marina leaf grown at Batemans Bay, New South Wales, Australia: (a) Relative water content, calculated with total leaf water content, (RWC+, %) as a function of leaf water potential in absolute value ($|\Psi_{\text{leaf}}|$, MPa). Key points and their x, y coordinates are indicated by open symbols. In domain 1, A (open circle) shows the point of full hydration at which $|\Psi_{\text{leaf}}|$ was 0.1 MPa and leaf RWC was set at 100%; the transition between domains 1 and 2 is indicated by **B** (open diamond); **C** (open triangle) is the turgor loss point at the transition between domains 2 and 3. (b) Inverse leaf water potential in absolute value ($|1/\Psi_{\text{leaf}}|$, MPa⁻¹) as a function of relative water deficit, calculated with total leaf water content, (RWD+, %). The full data set is plotted in the inset graph, whereas points representing extracellular water (points A to B in (a)) were excluded from the data set plotted in the main graph. The main graph shows relationships among calculated values of Ψ_{leaf} at the turgor loss point (point **c**, open circle), the y intercept (**d**₀), and the intercept (d_x) at RWD+ = RWD_x+, i.e. relative water deficit during domain 1. At point **c**, $|1/\Psi_{\pi}^{0}| = 0.2$ so $|\Psi_{\pi}^{0}| = 5$ MPa. The linear regression between $|1/\Psi_{leaf}|$ and RWD+ after the turgor loss point was reached followed the equation: y = -0.0034x + 0.2682 ($r^2 = 0.98$). Therefore, $|1/\Psi_{leaf}|$ at $d_0 = 0.2682$, and at $d_x = 0.2376$ as RWD+ at $d_x = 11$. Accordingly, $|\Psi_{\text{leaf}}|$ at $d_0 = 3.7$ MPa and at $d_x (\Psi_{\pi}^{100}) = 4.3$ MPa.

Fig. 2. Transverse sections of (a) petiole, (b) leaf midvein, and (c) leaf lamina of *A. marina*. Symbols: B: bundle sheath extension; C: cisternae; F: fimbrial vein; H: hypodermis; P: palisade mesophyll; S: spongy mesophyll; T: trichomes; V: vascular bundle. Yellow arrows indicate mucus secretion glands in (a) and salt secretion glands in (b) and (c). Black arrow indicates stomate. Bars are 100 μm.

Fig. 3. Trichomes from the abaxial surface of *A. marina* leaves at different hydration states viewed with (a) bright-field microscopy and (b-e) cryo-SEM. (a) Trichome general structure. Symbols: ca: cap cell, st: stalk cell, ba: basal cell. (b) Both cap and stalk cells were filled with water at $\Psi_{\text{leaf}} = -0.1$ MPa. (c, d) Cap cells were collapsed but stalk cells maintained their shapes and were filled with gas at $\Psi_{\text{leaf}} = -0.1$ MPa. (c, d) The lower stalk cell maintained its shape while the cap and upper stalk cell were shrivelled and collapsed at $\Psi_{\text{leaf}} = -4.3$ MPa. Bars are 20 µm. (f) Diagrammatic representation of changes in trichomes with dehydration, showing a trichome fully expanded and filled with water (blue) at $\Psi_{\text{leaf}} = -0.1$ MPa, shrunken, filled with gas (white) and with a collapsed cap cell at $\Psi_{\text{leaf}} = -0.25$ MPa, and shrunken, filled with the cap and upper stalk cells collapsed at $\Psi_{\text{leaf}} = -4.3$ MPa.

Fig. 4. Detection of Ψ_{leaf} threshold for draining of water-filled trichomes during dehydration in A. marina leaves. (a) Trichome layer covering abaxial leaf surface viewed under white light showing the cap cells (grey) and underline stalk cells (circular back area) of the trichomes and (b) under blue light showing the absence of detectable auto-fluorescence which could interfere with identification of the green fluorescence emitted by fluorescein under blue-exciting light. (c, d and movie 1 (Supplement S2)) The spread of fluorescein over the wet cap surface of water – filled trichomes (Ψ_{leaf} = -0.1 MPa). The images were collected at 10s (c) and 2 min (d) after fluorescein was applied to the cap surface at a distance approximately 1 mm from the observed area. Fluorescence intensity was initially weak (c) and increased with time (d) as more fluorescein diffused across the observed area. (e-f and movie 2 (Supplement S3)) Uptake of fluorescein applied to the cap surface of trichomes when Ψ_{leaf} = -0.25 MPa. The observed area partly included the place where fluorescein was applied. Fluorescence from the dye drop was saturating (white area in e), and reflected by surrounding trichomes (green area in e). The drop of fluorescein initially maintained its shape on the cap surface and then rapidly disappeared a minute later when the drop was absorbed by underlying trichomes (f). Note that some stalk cells were filled with dye (arrow in f) while in others, dye remained in the wall of cap and stalk cells and the surrounding area appeared black. Bars are 0.05 mm.

Fig. 5. Petiolar cisternae in leaves of *A. marina*. Distribution of cisternae (\star) in (a) transverse and (b) longitudinal sections as visualized by bright- field microscopy. Cryo-SEM micrographs of cisternae

filled with water when $\Psi_{\text{leaf}} = -0.1 \text{ MPa}$ (c), or air and droplets when $\Psi_{\text{leaf}} = -0.1 \text{ MPa}$ (c, d), $\Psi_{\text{leaf}} = -0.7 \text{ MPa}$ (e) and $\Psi_{\text{leaf}} = -6 \text{ MPa}$ (f). Bars are 50 μ m in (a, b), and 25 μ m in (c-f).

Fig. 6. Cryo-SEM micrographs of transverse sections through (a, b) pith parenchyma in the midvein and (c, d) paradermal sections through spongy mesophyll cells in the lamina of leaves of *A. marina* differing in leaf water status. Arrows point at extracellular spaces filled with ice, indicating the presence of liquid water (a, c) at $\Psi_{\text{leaf}} = -0.06$ MPa and with gas (b,d) at $\Psi_{\text{leaf}} = -1$ MPa. Bars equal 20 µm in (a, b) and 10 µm in (c, d).

Fig. 7. Plasmolysis in different cell types of leaves of *A. marina* after turgor had been lost ($\Psi_{\text{leaf}} \approx -6$ MPa). (a) Petiolar collenchyma, (b) petiolar parenchyma surrounding gas-filled cisternae, (c) hypodermis, and (d) palisade mesophyll. Arrows indicate gaps between cell walls and cell membranes. Bars are 25 µm in (a, b), and 20 µm in (c, d).

Fig. 8. Diagrammatic summary of changes in the leaf lamina during dehydration through domains 1, 2, and 3 of the PV curve shown in Fig. 2. The major tissue layers of hypodermis (H), palisade (P), spongy mesophyll (S), and trichomes (T) are indicated as in Fig. 3. Salt secretion glands (purple) occurred on both leaf surfaces while stomata (pink dots) occurred only on the abaxial surface. At maximum hydration in Domain 1, all cells were turgid. Extracellular water (blue) occurred in a few extracellular spaces of the mesophyll where cells were in close apposition as in Fig. 6c. Trichomes were filled with water as in Fig. 4b, f. At the beginning of Domain 2, extracellular water was absent from spongy mesophyll as in Fig. 6d and trichomes as in Fig. 4c, d, f. Finally, after turgor was lost at -5 MPa in Domain 3, plasmolysis was evident with further dehydration to -6 MPa in most cell types, including hypodermis (Fig. 7c), palisade (Fig. 7d) and spongy mesophyll cells (data not shown). Plasmolysed cells are indicated by thin white gaps between cell walls and cell membranes. While the leaves declined in thickness during dehydration, cellular connections were maintained with no sign of cell collapse, except in the shrivelled, hollow cap and stalk cells of trichomes (Fig. 4e, f). The variation in panel size and colours indicate progressive shrinkage of the leaf lamina during dehydration from Domain 1 through Domain 3.

Supplement S1. Reanalysis of the data set from Fig.1, assuming domain 1 represented a "plateau effect". (a) Relative water content, calculated without extracellular water, (RWC-, %) as a function of leaf water potential in absolute value ($|\Psi_{leaf}|$, MPa) with the fully hydrated status (100% RWC) set at *B* (open diamond). Open dots indicate extracellular water (RWD_x-). (b) Inverse leaf water potential in absolute value ($|1/\Psi_{leaf}|$, MPa⁻¹) as a function of relative water deficit, calculated without extracellular water (RWD-,%). For reference, the full data set in Fig.1b, including the extracellular water, is plotted in the inset graph. Point **c** (open circle) is the turgor loss point and $|\Psi_{\pi}^{0}| = 5$ MPa as in Fig. 1b. With the changes excluding extracellular water in RWD calculation, the equation for the linear regression between $|1/\Psi_{leaf}|$ and RWD- after **c** was reached was y = -0.003x + 0.2313 (r² = 0.98). Thus, the y intercept at point **d**, $|1/\Psi_{\pi}^{100}| = 0.2314$ so $|\Psi_{\pi}^{100}| = 4.32$ MPa.

Supplement S2: Movie showing diffusion of fluorescein across an abaxial leaf surface when $\Psi_{\text{leaf}} = -0.1$ MPa and trichomes were full of water (see Fig. 4 for further details).

Supplement S3: Movie showing absorption of fluorescein by draining trichomes when $\Psi_{\text{leaf}} = -0.25$ MPa (see Fig. 4 for further details).