



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Maladaptive innate immune training of myelopoiesis links inflammatory comorbidities

Citation for published version:

Li, X, Wang, H, Yu, X, Saha, G, Kalafati, L, Ioannidis, C, Mitroulis, I, Netea, MG, Chavakis, T & Hajishengallis, G 2022, 'Maladaptive innate immune training of myelopoiesis links inflammatory comorbidities', *Cell*. <https://doi.org/10.1016/j.cell.2022.03.043>

Digital Object Identifier (DOI):

[10.1016/j.cell.2022.03.043](https://doi.org/10.1016/j.cell.2022.03.043)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Cell

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.


Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Maladaptive innate immune training of myelopoiesis links inflammatory comorbidities

Graphical abstract



Authors

Xiaofei Li, Hui Wang, Xiang Yu, ..., Mihai G. Netea, Triantafyllos Chavakis, George Hajishengallis

Correspondence

triantafyllos.chavakis@uniklinikum-dresden.de (T.C.), geoh@upenn.edu (G.H.)

In brief

Trained innate immune responses contribute to pathology of a comorbid condition, as seen with arthritis after periodontitis in animal models.

Highlights

- Experimental periodontitis (EP) induces maladaptive trained myelopoiesis
- EP-induced trained phenotype is transmissible by bone marrow transplantation
- IL-1 signaling in hematopoietic progenitors mediates maladaptive training by EP
- Maladaptively trained myelopoiesis links the periodontitis-arthritis comorbidity

Article

Maladaptive innate immune training of myelopoiesis links inflammatory comorbidities

Xiaofei Li,^{1,10} Hui Wang,^{1,10} Xiang Yu,^{2,3} Gundappa Saha,¹ Lydia Kalafati,⁴ Charalampos Ioannidis,⁴ Ioannis Mitroulis,^{4,5} Mihai G. Netea,^{6,7} Triantafyllos Chavakis,^{4,8,9,*} and George Hajishengallis^{1,9,11,*}

¹Department of Basic and Translational Sciences, Laboratory of Innate Immunity and Inflammation, Penn Dental Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

²Joint International Research Laboratory of Metabolic and Developmental Sciences, School of Life Sciences and Biotechnology, Shanghai Jiao Tong University, Shanghai 200240, China

³Department of Biology, University of Pennsylvania, Philadelphia, PA 19104, USA

⁴Institute for Clinical Chemistry and Laboratory Medicine, Faculty of Medicine, Technische Universität Dresden, 01307 Dresden, Germany

⁵First Department of Internal Medicine and Department of Haematology, Democritus University of Thrace, 681 00 Alexandroupolis, Greece

⁶Department of Internal Medicine and Radboud Center for Infectious Diseases, Radboud University Medical Center, Nijmegen 6525 XZ, the Netherlands

⁷Department of Immunology and Metabolism, Life and Medical Science Institute, University of Bonn, 53115 Bonn, Germany

⁸Centre for Cardiovascular Science, University of Edinburgh, Edinburgh EH16 4TJ, UK

⁹Senior author

¹⁰These authors contributed equally

¹¹Lead contact

*Correspondence: triantafyllos.chavakis@uniklinikum-dresden.de (T.C.), geoh@upenn.edu (G.H.)
<https://doi.org/10.1016/j.cell.2022.03.043>

SUMMARY

Bone marrow (BM)-mediated trained innate immunity (TII) is a state of heightened immune responsiveness of hematopoietic stem and progenitor cells (HSPC) and their myeloid progeny. We show here that maladaptive BM-mediated TII underlies inflammatory comorbidities, as exemplified by the periodontitis-arthritis axis. Experimental-periodontitis-related systemic inflammation in mice induced epigenetic rewiring of HSPC and led to sustained enhancement of production of myeloid cells with increased inflammatory preparedness. The periodontitis-induced trained phenotype was transmissible by BM transplantation to naive recipients, which exhibited increased inflammatory responsiveness and disease severity when subjected to inflammatory arthritis. IL-1 signaling in HSPC was essential for their maladaptive training by periodontitis. Therefore, maladaptive innate immune training of myelopoiesis underlies inflammatory comorbidities and may be pharmacologically targeted to treat them via a holistic approach.


INTRODUCTION

Innate immune cells are now appreciated to retain heterologous memory of earlier microbial or inflammatory challenges that enables them to respond stronger upon future challenge with the same or unrelated stimuli. This state of heightened responsiveness, which is based on epigenetic innate immune memory, is designated as “trained innate immunity” (TII) (Netea et al., 2020; Penkov et al., 2019). We, and others, have shown that systemic inflammatory stimuli can initiate TII in the mouse and human bone marrow (BM) through long-lasting adaptations in hematopoietic stem and progenitor cells (HSPC), associated with a long-term myeloid bias (“trained myelopoiesis”) (Chavakis et al., 2019; Cirovic et al., 2020; de Laval et al., 2020; Kalafati et al., 2020; Kaufmann et al., 2018; Mitroulis et al., 2018). Experimental studies show that TII can be protective against infections and tumors (Ciarlo et al., 2020; Kalafati et al., 2020; Moorlag

et al., 2020a). However, TII is potentially detrimental, hence maladaptive, in chronic inflammatory pathologies (Chavakis et al., 2019).

By expressing Toll-like receptors and receptors for cytokines and growth factors, HSPC sense peripheral infection/inflammation and adapt via expansion and myeloid skewing (Chavakis et al., 2019). Thus, inflammation-adapted or trained HSPC could act as a central hub that perpetuates inflammation through a positive-feedback loop between the BM and peripheral tissues affected by inflammatory disorders (Chavakis et al., 2019), although experimental evidence for this hypothesis is scarce.

Here, we explored the concept that maladaptive TII in BM hematopoietic progenitors may provide a mechanistic link for inflammatory comorbidities, such as the enhanced risk of systemic diseases (e.g., cardiometabolic disease and arthritis) in periodontitis patients (D’Aiuto et al., 2018; Hajishengallis and Chavakis, 2021; Potempa et al., 2017). Periodontitis is a prevalent



(legend on next page)

inflammatory disease of the soft and bone tissues that support the dentition and poses a significant public health burden (Hajishengallis, 2015; Kassebaum et al., 2014; Listl et al., 2015; Peres et al., 2019). Conceivably, the low-grade systemic inflammation caused by periodontitis may contribute to the periodontitis-systemic disease connection (Genco and Sanz, 2020; Hajishengallis and Chavakis, 2021). Indeed, compared with healthy controls, patients with severe periodontitis have elevated inflammatory mediators and neutrophil numbers in the blood (D'Aiuto et al., 2013; Schenkein et al., 2020). Conversely, successful local treatment of periodontitis attenuates systemic inflammatory markers (Bajaj et al., 2018; D'Aiuto et al., 2018; Schenkein et al., 2020). The relationship between periodontitis and linked comorbidities is often bidirectional as systemic diseases can promote susceptibility to periodontitis (Genco and Sanz, 2020; Winning and Linden, 2017). However, there is no documented unifying causal mechanism of how periodontitis affects and is affected by comorbidities.

Since many chronic inflammatory diseases are driven by inflammatory myeloid cells, we hypothesized that inflammation-driven alterations in their BM progenitors, as may occur in the context of maladaptive TII (Chavakis et al., 2019; Netea et al., 2020), could influence distinct inflammatory disorders that emerge as comorbidities. We addressed this hypothesis in the context of the comorbidity of periodontitis and arthritis, which leads to progressive erosion of cartilage and bone in the joints (Smolen et al., 2018). An epidemiological association between these two inflammatory diseases remains even after adjusting for common risk factors (de Pablo et al., 2008; Scher et al., 2014). Our present findings show that maladaptive BM-mediated TII underlies the comorbid connection between periodontitis and arthritis, as modeled by ligature-induced periodontitis (LIP) and collagen antibody-induced arthritis (CAIA) in mice.

RESULTS

LIP induces a sustained increase in myelopoiesis

We first investigated whether experimental periodontitis induces inflammatory adaptation of HSPC in the BM. To this end, mice were subjected, or not, to LIP by ligature placement around the left and right maxillary second molar teeth. After 7 days, LIP-

subjected mice displayed significantly higher frequency and numbers of hematopoietic progenitors (LSK; Lin[−]cKit⁺Sca1⁺) and multipotent progenitors (MPP; CD48⁺CD150[−]LSK) in the BM, as compared with unligated controls (Figure S1A). No difference was found in long-term (LT)-hematopoietic stem cells (HSC) (CD48[−]CD150⁺LSK), whereas the frequency and numbers of short-term (ST)-HSC (CD48[−]CD150[−]LSK) were decreased (Figure S1A). The LIP-induced increase in MPP was associated with an increase in the frequency of the myeloid-biased MPP3 subset (FIt3[−]CD48⁺CD150[−]LSK) (Pietras et al., 2015) (Figure S1B). No significant difference was observed in the lymphoid-biased MPP4 (FIt3[−]CD48⁺CD150[−]LSK) or the erythro-megakaryocytic-biased MPP2 subset (FIt3[−]CD48⁺CD150[−]LSK) (Pietras et al., 2015) (Figure S1B). Although the total numbers of LT-HSC remained unaltered (Figure S1A), LIP led to increased frequencies of CD41⁺ and CD61⁺ LT-HSC subpopulations (Figure S1C); CD41⁺ LT-HSC harbor myeloid-biased cells (Gekas and Graf, 2013) and CD61⁺ LT-HSC are responsive to inflammatory stimuli (Mann et al., 2018).

Although LIP causes substantial bone loss within a few days, maintaining the ligatures for 21 days mimics the chronic phase of human periodontitis. Mice subjected to LIP for 21 days maintained the changes seen at day 7 (Figure S1), such as, increased frequency and numbers of LSK and MPP (Figure 1A). The elevated frequency of myeloid-biased MPP3 was now associated with a decreased frequency of lymphoid-biased MPP4 in LIP-subjected mice (Figure 1B). There was an even more pronounced difference on day 21 (than on day 7) between LIP-subjected and control mice regarding the elevated frequencies of CD41⁺ and CD61⁺ LT-HSC in the former group (Figure 1C). Analysis of myeloid progenitors (MyP; Lin[−]cKit⁺Sca1[−]) revealed that LIP led to elevated absolute numbers of granulocyte macrophage progenitors (GMP; Lin[−]cKit⁺Sca1[−]CD16/32⁺CD34⁺) and increased proportion of GMP within the MyP population, associated with a corresponding reduction in the relative abundance of common myeloid progenitors (CMP; Lin[−]cKit⁺Sca1[−]CD16/32[−]CD34⁺) (Figure 1D). Consistently, LIP-subjected mice exhibited an increased proportion of Gr1^{hi}CD11b⁺ granulocytes and Gr1^{int}CD11b⁺ myeloid cells with a corresponding reduction in

Figure 1. LIP causes a sustained increase in myelopoiesis

Mice were subjected, or not (control), to LIP followed by BM analysis after 21 days.

(A) Frequencies (top) of LSK, LT-HSC, ST-HSC, and MPP in total BM cells and absolute cell numbers (bottom) of the same populations.

(B) Frequencies of MPP subsets in LSK in the BM of mice.

(C) Frequency of CD41⁺ and CD61⁺ LT-HSC in total LT-HSC cells in the BM.

(D) (Top) Representative fluorescence-activated cell sorting (FACS) plots for GMP and CMP in the BM; (bottom) absolute cell numbers of GMP (left) and frequency within the MyP pool of GMP (middle) and CMP (right) in the BM.

(E) (Top) Representative FACS plots to identify Gr1^{hi}CD11b⁺ granulocytes, Gr1^{int}CD11b⁺ myeloid cells, CD19⁺ B cells, and CD3⁺ T cells and (bottom) frequencies of these populations in CD45⁺ cells in the BM.

(F–I) Mice were subjected, or not (control, NL), to LIP, and BM cells were harvested on day 7. FACS-sorted LSK were subjected to RNA-sequencing analysis.

(F) Differential gene expression in LSK from LIP-subjected mice versus non-ligated controls. Volcano plot showing the distribution of the adjusted p values (−log₂(FDR)) and the fold changes (log₂ fold change). Significant changes are shown in blue (downregulated) or orange (upregulated) (FDR < 0.05), n = 3.

(G) PCA plot of LSK samples from ligated and non-ligated (NL) groups.

(H and I) Top overrepresented GO terms and KEGG pathways including upregulated (orange) (H) or downregulated (blue) genes (I) in LSK from LIP-subjected mice versus NL controls.

(A–D) Control, n = 6 mice/group; LIP, n = 9 mice/group and (E) n = 6 mice/group.

Data are means ± SD. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, NS, not significant versus control mice; two-tailed Student's t test.

See Figure S1.




Figure 2. LIP-experienced mice display increased myelopoiesis after LPS challenge

(A–C) BM LSK were sorted from mice subjected to 21dL/14dR or not (NL), and scRNA-seq was performed.

(A and B) Two-dimensional UMAP representation of 15,257 cells, according to (A) sample origin and (B) results of clustering.

(C) Heat map visualization of the distribution of cells within each of the identified clusters, normalized for the number of cells per sample in the dataset.

(D) Experimental design.

(E) Frequencies (top) of LSK, LT-HSC, ST-HSC, and MPP in total BM cells and absolute cell numbers (bottom) of the same populations.

(F) Frequency (left) within the MyP pool of GMP and absolute numbers (right) of GMP in the BM.

(legend continued on next page)

the proportion of CD19⁺ B cells and CD3⁺ T cells, compared with the proportions of these populations in control mice (Figure 1E). Hence, chronic LIP leads to sustained enhancement of myelopoiesis.

LIP induces transcriptomic changes in hematopoietic progenitors

To study the mechanisms underlying the LIP-induced myeloid bias in HSPCs, we performed transcriptome analysis of LSK from LIP-subjected and control mice. We identified 2,139 differentially expressed genes of which 979 and 1,160 genes were significantly up- or downregulated, respectively, in LSK from LIP-subjected mice, compared with those from control mice (false discovery rate [FDR] < 0.05) (Figure 1F). Principal component analysis (PCA) showed a clear separation between the two groups (Figure 1G). Gene Ontology (GO) enrichment analysis revealed that several GO terms, including “myeloid leukocyte migration,” “innate immune response,” and “neutrophil-mediated immunity,” were overrepresented in the significantly upregulated genes, whereas terms such as “cellular response to interleukin-7,” “erythrocyte differentiation,” or “lymphocyte proliferation” were overrepresented in the significantly downregulated genes (Figures 1H and 1I). Significant upregulation of myeloid lineage markers, including *Cd68*, *Csf1r*, *kit*, *Ccr2*, *Csf3r*, and *Mpo* (Figure S1D), and downregulation of lymphoid lineage and erythrocyte markers were observed in LSK of LIP-subjected mice (Figure S1E). Moreover, the GO terms “oxidative phosphorylation” and “mitochondrial acetyl-CoA biosynthetic process from pyruvate” were overrepresented in the significantly downregulated genes, which was consistent with the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis (Figure 1I). Accordingly, the expression of genes encoding NADH dehydrogenase (e.g., *Ndufc2* and *Ndufs6*) and ATP synthase (e.g., *Atp5d* and *Atp5o*) were significantly downregulated (Figure S1E). The GO terms “interleukin-1-mediated signaling pathway” and the KEGG pathway “osteoclast differentiation” were also significantly overrepresented (Figures 1H and S1). Also, the GO terms “negative regulation of histone H3-K9 methylation” (GO:0051573), “histone H3-K4 trimethylation” (GO:0080182), “positive regulation of DNA demethylation” (GO:1901537), and “histone H3-K9 demethylation” (GO:0033169) were overrepresented in the significantly upregulated genes, whereas the GO term “positive regulation of histone H3-K27 methylation” (GO:0061087) was significantly overrepresented in the significantly downregulated genes (Figures 1H and 1I). Hence, the expression of *Kdm1a* (Perillo et al., 2008), *Kdm3b* (Kim et al., 2012), and *Pax5* (Johnson

et al., 2004) (critical for H3-K9 demethylation) as well as *Tet3* (Deplus et al., 2013) and *Ncoa6* (Qing et al., 2014) (critical for H3-K4 methylation) was upregulated, whereas the expression of *Mtf2* (Perino et al., 2018) and *Eed* (Margueron et al., 2009) (critical for H3-K27 methylation) was downregulated (Figure S1F).

Ingenuity pathway analysis (IPA) for upstream regulators revealed that several myeloid transcription factors (TF) were predicted to act as upstream regulators in LSK from LIP-subjected mice, compared with those of control mice, including CEBPA, CEBPE, ID1, ID2, and SPI1 (PU.1) (Figure S1G), which mediates IL-1 β -induced HSC myeloid differentiation (Pietras et al., 2016). Moreover, IL-1 β /IL-1 and growth factors (CSF2 and CSF3) involved in myelopoiesis were also predicted to act on LSK from LIP-subjected mice (Figure S1G). These findings suggest transcriptomic rewiring of BM progenitors associated with inflammatory signaling, such as IL-1.

LIP induces trained innate immunity

We then investigated if hematopoietic progenitors of mice with previous LIP exposure, but without active disease, can elicit enhanced myelopoiesis responses and generate hyper-responsive myeloid cells to a future challenge. Ligature removal abrogates the microbial challenge that drives inflammation, thereby leading to periodontitis resolution (Kourtzelis et al., 2019; Yuh et al., 2020). Mice were thus subjected to LIP for 21 days followed by ligature removal. After 14 days (when local periodontal inflammation was resolved; Figure S2A), analysis of BM myelopoiesis revealed no significant differences between LIP-subjected and control mice (Figures S2B–S2F), except for slight differences between the two groups regarding the proportion of GMP within the MyP (Figure S2E). Thus, inflammation resolution in the periodontium quantitatively restores phenotypic myelopoiesis, as assessed by flow cytometry, to basal levels.

Next, we performed scRNA-seq in sorted LSK from mice subjected to 21 days LIP followed by 14 days resolution (“21dL/14dR” mice) and naive controls (rested for 35 days; “NL” mice). Two-dimensional uniform manifold approximation and projection (UMAP) of 15,257 (NL: 7,051; 21dL/14dR: 8,206) LSK and clustering, partitioned LSK into 8 clusters (C1–C8) (Figures 2A and 2B). Although most clusters were comparably represented in the 21dL/14dR and NL groups, C7 was modestly enriched with cells from the 21dL/14dR group (Figure 2C). However, C7 comprised a tiny portion of LSK (Figure 2B). GO enrichment analysis of C7-specific markers showed that the GO terms “leukocyte activation” and “regulation of myeloid cell differentiation” were in the top 20 significantly enriched GO terms (Figure S2G). There were no significant differences in myeloid marker expression in

(G) Frequency of Gr1^{hi}CD11b⁺ granulocytes and Gr1^{int}CD11b⁺ myeloid cells in CD45⁺ cells (left) and cell numbers (right) of the same populations in the BM.

(H) Total white blood cell (WBC) count (left) and Gr1⁺CD11b⁺ cell counts (right) in the peripheral blood.

(I) Representative FACS plots (left) to identify Gr1^{hi}CD11b⁺ granulocytes and Gr1^{int}CD11b⁺ myeloid cells and frequency (right) of the same populations in CD45⁺ cells in the lungs of mice.

(J) Experimental design.

(K) Isolated splenic monocytes and neutrophils were re-stimulated *ex vivo* with LPS (10 ng/mL) for 24 h. The supernatant was collected for measuring IL-6 and TNF.

Data are means \pm SD (n = 6 mice/group) (E–I and K). *p < 0.05, **p < 0.01, NS, not significant versus untrained mice; two-tailed Student's t test. U, untrained; T, trained.

See Figure S2.




Figure 3. Epigenetic rewiring of trained LSK and GMP

BM LSK and GMP were sorted from mice subjected to 21dL/14dR or not (NL), and scATAC-seq was performed.

(A and B) Two-dimensional UMAP representation of 37,903 cells, according to (A) sample origin and (B) results of clustering.

(C) UMAP (up) of the distribution of cells from the four different samples (LSK and GMP from mice subjected to 21dL/14dR or not) and heat map visualization (bottom) within each of the identified clusters, normalized for the number of cells per sample in the dataset, are shown.

(legend continued on next page)

the overall LSK population from “21dL/14dR” mice compared with those from NL mice (Figure S2H).

We then addressed possible qualitative alterations (associated with trained myelopoiesis) in hematopoietic progenitors of LIP-subjected mice currently under LIP resolution (“LIP-experienced” mice). We investigated if the LIP-induced modulation of HSPC could promote a higher response of hematopoietic progenitors to future systemic challenge with lipopolysaccharide (LPS) (Mitroulis et al., 2017), which would be indicative of inflammatory memory in the BM. To this end, mice, previously subjected to LIP for 21 days followed by ligature removal for 28 days, received a single LPS dose i.p.; BM analysis was performed after 72 h (Figure 2D). LIP-experienced mice exhibited a stronger response to LPS than naive controls, as shown by significantly increased frequency and numbers of the LSK and MPP pools (Figure 2E), GMP (Figure 2F), as well as Gr1^{hi}CD11b⁺ granulocytes and Gr1^{int}CD11b⁺ myeloid cells (Figure 2G). Consistently, the LIP-experienced group showed significantly higher white blood cell counts and Gr1⁺CD11b⁺ cells in peripheral blood (Figure 2H). Upon LPS challenge, LIP-experienced mice also exhibited significantly higher frequencies of Gr1^{hi}CD11b⁺ and Gr1^{int}CD11b⁺ cells in the lungs (Figure 2I), compared with LPS-challenged controls. Therefore, BM hematopoietic progenitors of LIP-experienced mice maintain a myeloid differentiation bias and respond with enhanced myelopoiesis to a future systemic challenge. LIP-experienced mice are thereafter referred to as “LIP-trained” mice. Consistent with the notion of LIP-induced inflammatory memory, mature myeloid progeny (spleen-derived neutrophils and monocytes) of LIP-trained mice responded with higher IL-6 and tumor necrosis factor (TNF) production to secondary (ex vivo) LPS challenge than their counterparts from naive controls (Figures 2J and 2K).

LIP induces epigenetic rewiring of BM progenitors

The ability of hematopoietic progenitors and mature myeloid cells from LIP-trained mice to respond stronger to future LPS challenges (Figures 2D–2K) suggested the emergence of an epigenetically-based innate immune memory. To establish LIP-induced innate immune memory, we tested if LIP can indeed induce epigenetic rewiring of BM progenitors.

We thus performed single-cell assay for transposase-accessible chromatin using sequencing (scATAC-seq) analysis of hematopoietic progenitors LSK and GMP in mice that were trained by subjecting them to LIP for 21 days followed by ligature removal for 14 days (“21dL/14dR” mice) or were not ligated during the entire period (“NL” controls) (Figure 3). Two-dimensional UMAP of 12,758 (NL: 5,341; 21dL/14dR: 7,417) LSK and 25,145 (NL: 12,944; 21dL/14dR: 12,201) GMP

(Figure 3A) and clustering, partitioned LSK and GMP into 15 clusters (C1–C15) (Figure 3B). C2, C7, C8, C10, and C12–C14 comprised LSK; C1, C3–C6 comprised GMP, and C9, C11, and C15 included both LSK and GMP (Figure 3C). The LSK cluster C2 displayed enhanced chromatin accessibility in genes indicating myeloid differentiation, *Itga2b* (*Cd41*) and *Gata2* (Figure S3A). The GMP cluster C3 showed enhanced chromatin accessibility in genes with an inflammatory signature: *Csf1r*, encodes CD115 and promotes myelopoiesis; and *Nlrp3*, encodes Nod-like receptor (NLR) family pyrin domain containing 3 (NLRP3) and acts as an inflammasome component contributing to IL-1 β secretion (Figure S3B).

In line with a LIP-induced increased myeloid bias in LSK (Figures 1 and S1), differential accessibility analysis revealed differentially accessible regions (DARs) in LSK from “21dL/14dR”-trained mice as compared with LSK from “NL” mice (Figure 3D). The top 20 significantly enriched GO terms in LSK, identified on the basis of genes annotated to regions more accessible due to LIP-induced training, revealed terms such as “regulation of cell differentiation,” “regulation of metabolic process,” and “cell migration” (Figure 3E), consistent with the LSK RNA-seq results (Figure 1H) and with previous findings implicating metabolic reprogramming in T11 (Hajishengallis et al., 2021). GO enrichment analysis of the enriched TF motifs in LSK revealed that the GO terms “myeloid cell differentiation,” “granulocyte differentiation,” “NIK/NF-kappaB signaling,” and “response to interleukin-1” were significantly enriched by LIP-induced T11 (Figure 3F). The top enriched TF motifs in the GO term “myeloid cell differentiation” includes SPI1 (PU.1), KLF2 (Krüppel-like transcription factor 2, a potent regulator of myeloid cell activation) (Maha-beleshwar et al., 2011), and the GO term “granulocyte differentiation” includes the TF motifs CEBPA, CEBPE, SP3, SPI1, and GATA2 (Figure 3G). The GO term “NIK/NF-kappaB signaling” includes REL, RELB, and NFkB2, indicating that the NF- κ B binding motifs are more accessible in LSK from “21dL/14dR” mice than in their counterparts from NL mice (Figure 3G); in the GO term “response to interleukin-1,” the enrichment of TF motifs IRF1, CEBPB, and EGR1 implied the activation of IL-1 signaling pathway (Figure 3G), consistent with the transcriptomic analysis of LSK (Figures 1F–1I and S1D–S1G). A genome browser track displays a DAR in the *Il6* gene locus and the RELA binding motifs within this region (Figure 3H, top), as well as DARs near the promoter regions of *Tlr4* and *Myd88* gene locus in LSK from “21dL/14dR” (trained) and control (NL) mice (Figure 3H, bottom).

Differentiation trajectories analysis showed that C4 and C5 existed at the end of the pseudotime trajectories from LSK to GMP (Figure S3C), suggesting that they consist of the most differentiated myeloid precursors within GMP, in which C5 was

(D) Volcano plots displaying differential accessibility analysis results (blue, lower versus red, greater differential chromatin accessibility) for LSK from mice subjected to 21dL/14dR versus their counterparts from control mice ($\text{abs}(\log_2\text{FC}) \geq 0.2$, $p < 0.05$).


(E) GO enrichment results of top 20 significantly enriched GO terms sorted by Protein Analysis Through Evolutionary Relationships (PANTHER) based on genes annotated to regions more accessible due to 21dL/14dR treatment (Bonferroni-corrected, $p < 0.05$).

(F) GO enrichment analysis of enriched TF binding motifs in LSK from mice subjected to 21dL/14dR versus control (NL) group and the top 20 significantly enriched GO terms (Bonferroni-corrected, $p < 0.05$).

(G) Visualization of top enriched TF binding motifs in the indicated GO terms for the LIP specifically accessible regions in LSK using the homer TF motif database.

(H) Genome browser track showing DAR in the *Il6* gene locus and the RELA binding motifs within this region, as well as DARs close to the promoter regions of *Tlr4* and *Myd88* gene locus.

See Figure S3.



(legend on next page)

significantly enriched by LIP-induced TII (Figure 3C). GO enrichment analysis of C5-specific markers revealed that the GO terms “myeloid cell differentiation,” “positive regulation of ROS metabolic process,” “regulation of phagocytosis,” and “positive regulation of MAPK cascade” were significantly enriched by LIP-induced TII (Figure S3D). C5 also showed enhanced chromatin accessibility in *Tlr4*, as well as *Camk1d*, which encodes calcium/calmodulin-dependent protein kinase 1D that promotes reactive oxygen species (ROS) production and phagocytosis by granulocytes (Verploegen et al., 2005); *Itga1*, encoding CD49a, which promotes neutrophil migration (Ridger et al., 2001); and *Map2k4*, encoding MAP2K4, which phosphorylates mitogen-activated protein (MAP) kinases (MAPK)8/JNK1, MAPK9/JNK2, and MAPK14/p38 in response to environmental stresses or stimuli (Figure S3E).

Therefore, scATAC-seq suggests that LIP-induced training is associated with a sustained myeloid lineage bias in LSK, allowing for an enhanced myelopoiesis response upon secondary stimulation, as well as pro-inflammatory signatures in both LSK and GMP. Moreover, the results of the scATAC-seq analysis reveal that, upon LIP resolution, HSPC retain an epigenetic myeloid differentiation bias.

LIP-induced trained myelopoiesis underlies the periodontitis-arthritis comorbidity

We next determined whether the above-documented inflammatory memory renders LIP-trained mice more susceptible to arthritis, as modeled by CAIA (Khachigian, 2006). To this end, mice subjected to LIP for 21 days followed by 14 days of ligature removal were challenged with CAIA. Mice without previous exposure to LIP were also challenged with CAIA and served as controls (Figure S4A). LIP-trained mice were more susceptible to CAIA than untrained controls, as evidenced by significantly higher clinical arthritis score and ankle joint thickness (Figure S4B). Whether this increased susceptibility to arthritis is mediated by an inflammatory LIP-BM axis was addressed by BM transplantation (BMT) experiments from LIP-trained mice to naive recipients.

To this end, BM cells were isolated from CD45.2⁺ mice, which were either trained by subjecting them to 21-day LIP followed by 14-day resolution (“21dL/14dR” mice), or were left un-

trained, and transferred to lethally irradiated congenic B6.SJL CD45.1⁺ mice. Next, 12 weeks post-BMT, groups of recipient CD45.1⁺ mice were subjected to LIP (for 5 days) or CAIA (14 days) (Figure 4A). CD45.1⁺ mice transplanted with BM cells from “21dL/14dR” mice developed increased bone loss (Figure 4B) and inflammatory gene expression (Figure 4C), as well as a higher abundance of monocytes and neutrophils and total CD45⁺ immune cells in the gingiva (Figure 4D), compared with CD45.1⁺ mice that received BM cells from untrained CD45.2⁺ mice. Similarly, upon induction of CAIA in CD45.1⁺ recipient mice, we observed more severe arthritis in mice transplanted with BM cells from 21dL/14dR mice than in mice transplanted with BM cells from untrained controls, as shown by increased clinical arthritis score and ankle joint thickness (Figure 4E) and aggravated histopathology in ankle joints (Figure 4F). Moreover, compared with the control group, CD45.1⁺ mice receiving BM cells from “21dL/14dR” CD45.2⁺ mice showed a higher abundance of monocytes and neutrophils and total CD45⁺ leukocytes in the synovium (Figure 4G). Mature splenic neutrophils and monocytes from CD45.1⁺ mice receiving BM cells from 21dL/14dR CD45.2⁺ donor mice responded with significantly higher secretion of IL-6 and TNF to secondary LPS challenge than their counterparts from CD45.1⁺ mice receiving BM cells from untrained CD45.2⁺ donors (Figures S4C and S4D). Thus, LIP-induced maladaptive TII confers increased susceptibility to not only periodontitis but also a comorbid condition, namely arthritis.

To exclude the possibility that the differences in the abundance of myeloid cells infiltrating the synovium of recipient groups were due to potential differences in the reconstitution potential of HSPC from the “21dL/14dR” donors versus those from untrained donors, we performed an analysis of BM and peripheral chimerism in recipient mice. Using the CD45.1/CD45.2 congenic system (Figure 4H), 100 LT-HSC isolated from “21dL/14dR” or control CD45.2⁺ mice were transplanted together with 5×10^5 CD45.1⁺ competitors to lethally irradiated C57BL/6.SJL CD45.1⁺ mice. At 16 weeks post-BMT, no differences were observed in the frequencies of HSPC derived from “21dL/14dR” or control donors (Figures S4E and S4F). Peripheral blood analysis at 4–16 weeks post-BMT showed no difference in white blood cell counts and in the percentages of

Figure 4. LIP-induced trained myelopoiesis contributes to the periodontitis-arthritis comorbidity

(A) CD45.2⁺ mice were either trained (T) by LIP for 21 days followed by 14-day resolution or left untrained (U). BM cells were isolated from trained and untrained mice and transferred to lethally irradiated congenic B6.SJL CD45.1⁺ mice.

(B–D) At 12 weeks post-BMT, groups of recipient CD45.1⁺ mice were subjected to LIP for 5 days (B–D) or CAIA for 14 days (E–G).

(B) Bone loss, (C) relative gingival mRNA expression of indicated cytokines, and (D) FACS analysis of gingival monocytes (live CD45⁺CD11c⁺CD11b⁺Ly6G⁺Ly6C⁺), neutrophils (live CD45⁺CD11c⁺CD11b⁺Ly6G⁺Ly6C⁺), and total CD45⁺ cells in recipient CD45.1⁺ mice subjected to 5-day LIP.

(E–G) (E) Clinical arthritis score (left) and hind ankle joint thickness (right), (F) representative images of H&E (left) and Safranin-O staining (right) of tissue sections from knee joints harvested on day 7 (scale bars, 500 μ m), and (G) quantification of monocytes (live CD45⁺CD11c⁺CD11b⁺Ly6G⁺Ly6C⁺), neutrophils (live CD45⁺CD11c⁺CD11b⁺Ly6G⁺Ly6C⁺), and total CD45⁺ cells in the synovium of knee joints harvested on day 7.

(H) 100 LT-HSC sorted from CD45.2⁺ mice subjected, or not (control), to LIP for 21 days followed by 14-day resolution were co-transplanted with 5×10^5 CD45.1⁺ BM competitors.

(I) WBC count, donor-derived percentage, and lineage output (% of indicated cell types in donor-derived cells) in peripheral blood of mice receiving CD45.2⁺ LT-HSC.

Data are means \pm SD (B and C, n = 5 mice/group; D, G, and I, n = 6 mice/group; E, n = 7 mice/group). *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, NS, not significant versus untrained mice. Two-tailed Student's t test (B–D and G) except for *Rankl* in (C) (two-tailed Mann-Whitney U test); two-way repeated measures ANOVA and Sidak's multiple comparisons test (E and I). W, weeks.

See Figure S4.




Figure 5. CAIA- or LIP-induced trained myelopoiesis contributes to the bidirectional arthritis-periodontitis comorbidity

(A) CD45.2⁺ mice were either trained (T), by subjecting them to CAIA as indicated, or were left untrained (U; naive controls). BM cells were isolated from trained and untrained mice and transferred to lethally irradiated congenic B6.SJL CD45.1⁺ mice.

(B–G) At 12 weeks post-BMT, groups of recipient CD45.1⁺ mice were subjected to LIP for 5 days (B–D) or CAIA for 14 days (E–G).

(legend continued on next page)

“21dL/14dR” or control donor-derived cells (Figure 4I). However, LT-HSC from “21dL/14dR” mice gave rise to an elevated proportion of Gr1⁺CD11b⁺ myeloid cells with a corresponding decrease in the proportion of CD19⁺ B cells and CD3⁺ T cells, as compared with LT-HSC from control mice (Figure 4I). Thus, although there was no difference in the overall reconstitution potential, there was an inherent myeloid differentiation bias of the LT-HSC from the “21dL/14dR” group (versus those of controls), resulting from LIP-induced trained myelopoiesis in donor mice.

CAIA-induced trained myelopoiesis underlies the arthritis-periodontitis comorbidity

To demonstrate the bidirectional association of periodontitis with arthritis, as suggested by observations in humans (Lee and Choi, 2020; Mikuls et al., 2014; Potempa et al., 2017; Rodríguez-Lozano et al., 2019), we tested whether CAIA causes maladaptive training of HSPC leading to enhanced susceptibility to periodontitis. We first showed that CAIA significantly increased the frequency and numbers of LSK, LT-HSC, and MPP (versus controls; Figure S5A), as well as the frequency of myeloid-biased MPP3 (Figure S5B) and CD41⁺ LT-HSC (Figure S5C). These changes were associated with decreased frequency of erythro-megakaryocytic-biased MPP2 and lymphoid-biased MPP4 (Figure S5B), thus signifying a myelopoiesis bias. Indeed, CAIA also resulted in increased numbers and frequency of GMP (Figure S5D) and of Gr1^{hi}CD11b⁺ granulocytes in the BM (Figure S5E).

We next performed BMT experiments to test if CAIA-induced trained myelopoiesis underlies the comorbid connection between arthritis and periodontitis. To this end, BM cells were isolated from CD45.2⁺ mice 36 days after CAIA. This interval was selected based on observations that 22 days were required for resolution of arthritis (Figure S5F) and another 14 days were required for phenotypic restoration of myelopoiesis to steady-state levels (Figure S5G). BM cells from CAIA-experienced or untreated control CD45.2⁺ mice were transferred to lethally irradiated CD45.1⁺ recipients. Thereafter, 12 weeks post-BMT, groups of CD45.1⁺ recipient mice were subjected to LIP or CAIA (Figure 5A). CD45.1⁺ mice receiving BM cells from CAIA-subjected CD45.2⁺ mice developed increased periodontal bone loss (Figure 5B) and inflammatory gene expression

(Figure 5C) and a higher gingival abundance of monocytes, neutrophils, and total CD45⁺ immune cells (Figure 5D), compared with CD45.1⁺ mice receiving BM cells from control CD45.2⁺ mice. Moreover, mice that received BM cells from CAIA-trained mice also developed more severe arthritis (Figure 5E), increased infiltration of monocytes, neutrophils, and total leukocytes in the joints (Figure 5F), and aggravated histopathology of ankle joints (Figure 5G), compared with the recipients of BM cells from untrained controls. Mature splenic neutrophils and monocytes from CD45.1⁺ mice receiving BM cells from CAIA-trained CD45.2⁺ mice responded with significantly higher production of IL-6 and TNF to secondary LPS challenge than their counterparts from CD45.1⁺ mice receiving BM cells from untrained CD45.2⁺ mice (Figures S5H and S5I). Therefore, LIP-induced (or CAIA-induced) inflammatory maladaptive training of BM progenitors confers, in a bidirectional fashion, increased susceptibility to inflammatory comorbid conditions, namely, periodontitis and arthritis.

Maladaptively trained LT-HSC link comorbid inflammatory conditions

To obtain evidence that maladaptive TII is initiated at the level of LT-HSC, we performed competitive BMT with the CD45.1/CD45.2 congenic system and sorted LT-HSC from LIP (“21dL/14dR”) trained mice and untrained donors. Chimerism analysis in the BM at 12 weeks post-BMT showed no significant differences in the frequencies of LIP-trained or untrained control donor-derived cells (Figure 5H). Next, 12 weeks post-BMT, additional recipient CD45.1⁺ mice were subjected to CAIA. Compared with mice transplanted with LT-HSC from untrained CD45.2⁺ mice (“untrained” LT-HSC), mice receiving LT-HSC from trained CD45.2⁺ mice (“trained” LT-HSC) exhibited more severe arthritis (Figure 5I), a higher abundance of monocytes, neutrophils, and total CD45⁺ leukocytes in the synovium (Figure 5J), as well as increased percentage and counts of CD45.2⁺ monocytes, neutrophils, and leukocytes (Figure 5K). Mature splenic neutrophils and monocytes from mice that received trained LT-HSC responded with significantly higher secretion of IL-6 and TNF to secondary LPS challenge than their counterparts from mice transplanted with untrained LT-HSC (Figure 5L). Thus, periodontitis-trained LT-HSC lead to exacerbated inflammation and disease in transplanted mice in the context of a comorbid condition.

(B) Bone loss, (C) relative gingival mRNA expression of indicated cytokines, and (D) FACS analysis of gingival monocytes, neutrophils, and total CD45⁺ cells in recipient CD45.1⁺ mice subjected to 5-day LIP.

(E) Clinical arthritis score (left) and hind ankle joint thickness (right), (F) quantification of monocytes, neutrophils, and total CD45⁺ cells in the synovium of knee joints harvested on day 7, and (G) representative images of H&E (left) and Safranin-O staining (right) of tissue sections from knee joints harvested on day 7 (scale bars, 500 μ m).

(H) CD45.1⁺ mice were transplanted with 200 CD45.2⁺ LT-HSC from LIP-trained or untrained control mice together with 8×10^5 CD45.1⁺ BM competitor cells (left). Frequency of donor-derived cells in the BM of recipient mice (chimerism) 12 weeks post-BMT (right).

(I–L) 12 weeks post-BMT, additional recipient CD45.1⁺ mice were subjected to CAIA (I–K) or left untreated (L).

(I) Clinical arthritis score (left) and hind ankle joint thickness (right).

(J) Quantification of total monocytes, neutrophils, and CD45⁺ leukocytes, and (K) percentages of CD45.2⁺ cells in monocytes, neutrophils, and CD45⁺ cells (top panel) and absolute numbers of CD45.2⁺ monocytes, neutrophils, and total CD45.2⁺ cells (bottom panel) in the synovium of knee joints harvested on day 7.

(L) Isolated splenic monocytes and neutrophils were stimulated with LPS (10 ng/mL) for 24 h, and IL-6 and TNF concentration in the supernatant was measured. Data are means \pm SD (B–D, F, and I–K, n = 6 mice/group; E, n = 7 mice/group; H, right, L, n = 4 mice/group). *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, NS, not significant versus untrained mice. Two-tailed Student's t test (B–D, F, H, and J–L); two-way repeated measures ANOVA and Sidak's multiple comparisons test (E and I).

See Figures S5 and S6.




Figure 6. Analysis of pro-inflammatory mediators in LIP-trained BM

(A–E) Mice were subjected, or not (control), to LIP. After 14 days, indicated cytokines were analyzed in BM extracellular fluid (A) and serum (B) ($n = 4$ –10 mice/group). (C) After 21 days, relative mRNA expression (normalized to *Gapdh*) of indicated cytokines was analyzed in the gingiva ($n = 8$ –10 mice/group). (D and E) After 14 days, BM cells were harvested, and the indicated cell types were FACS sorted (gating strategies of LSK, CMP, and GMP in Figures 1 and S1; gating strategies for pre-neutrophils, CD11b $^{+}$ CD115 $^{-}$ Gr-1 $^{+}$ cKit $^{+}$ CXCR4 $^{+}$; immature neutrophils, CD11b $^{+}$ CD115 $^{-}$ Gr-1 $^{+}$ cKit $^{-}$ CXCR4 $^{-}$ Ly6G $^{+}$ CXCR2 $^{-}$; mature neutrophils, CD11b $^{+}$ CD115 $^{-}$ Gr-1 $^{+}$ cKit $^{-}$ CXCR4 $^{-}$ Ly6G $^{+}$ CXCR2 $^{+}$; monocytes, CD11b $^{+}$ Ly6G $^{-}$ Ly6C $^{+}$) and examined for *Csf3r*, *Il1b*, and *Il1r1* expression (D). Mature neutrophils isolated from the BM of LIP-trained or untrained controls were stimulated with recombinant mouse G-CSF for 24 h, and IL-1 β was measured in culture supernatants ($n = 6$ mice/group) (E).

Data are means \pm SD. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, NS, not significant versus control (A–C, E) or mature neutrophils (D); two-tailed Student's t test (A–C), except for IL-12p70 in (A) (two-tailed Mann-Whitney U test); one-way ANOVA with Dunnett's multiple comparisons test (D and E). ND, not detected.

To determine if LIP-induced maladaptive TII was restricted to myeloid-biased LT-HSC, we performed a similar competitive BMT using sorted CD41 $^{+}$ (myeloid-biased) or CD41 $^{-}$ (non-biased) LT-HSC from LIP-trained or untrained CD45.2 $^{+}$ mice

(Figure S6A). Peripheral blood analysis in mice receiving CD41 $^{+}$ LT-HSC showed no difference in white blood cell counts and percentage of LIP-trained or untrained donor-derived cells; however, LT-HSC from LIP-trained mice gave rise to an elevated

proportion of Gr1⁺CD11b⁺ myeloid cells with a corresponding decrease in the proportion of CD19⁺ B cells and CD3⁺ T cells, relative to LT-HSC from control donors (Figure S6B). No differences were seen in the frequencies of HSPC in the BM derived from LIP-trained or control donors (Figure S6D), suggesting comparable engraftment. Similar observations were made in mice transplanted with CD41⁺ LT-HSC (Figures S6C and S6E). These findings reveal an inherent myeloid bias in both CD41⁺ and CD41⁺ LT-HSC from LIP-trained donors leading to higher myelopoiesis in recipient mice. Consistently, when recipient mice were subjected to CAIA, those transplanted with trained LT-HSC, regardless of CD41 phenotype, had exacerbated arthritis (Figures S6F and S6G) and joint inflammation (Figures S6H and S6I). Mature splenic myeloid cells from mice that received trained CD41⁺ or CD41⁺ LT-HSC responded with higher secretion of IL-6 and TNF to secondary LPS challenge than their counterparts from mice transplanted with untrained LT-HSC (Figures S6J and S6K). Thus, LIP can induce myeloid differentiation bias and a maladaptive trained phenotype even in LT-HSC that are not originally myeloid biased, at least phenotypically, based on CD41 expression.

IL-1 signaling in HSPC mediates LIP-induced maladaptive training of myelopoiesis

To better understand how LIP modulates HSPC for enhanced myelopoiesis, we analyzed the BM extracellular fluid for cytokines implicated in the inflammatory modulation of HSPC (Chavakis et al., 2019), collected 14 days post-LIP (between 7 and 21 days when significant changes in HSPC modulation have occurred; Figures 1 and S1). We detected significantly increased concentrations of IL-1 β and G-CSF (but not of other cytokines tested) in the BM extracellular fluid in LIP-subjected mice, relative to controls (Figure 6A). Parallel analysis of the same cytokines in the serum of LIP-subjected mice indicated an elevated concentration of G-CSF but not of IL-1 β , relative to controls (Figure 6B). Analysis of cytokines upregulated in periodontitis (Dutzan et al., 2018; Shin et al., 2015) revealed high abundance of IL-1 β and G-CSF in the periodontal tissues of LIP-subjected mice, although the local induction of G-CSF (6.8-fold versus control) was more pronounced than that of IL-1 β (2.3-fold versus control) (Figure 6C). These data suggested that LIP-induced serum G-CSF might access the BM, where it might induce IL-1 β production.

To test this notion, HSPC, MyP, and distinct types of myeloid cells were sorted from the BM of mice subjected to LIP for 14 days. Analysis of G-CSF receptor (colony-stimulating factor 3 receptor, *Csf3r*) mRNA expression showed that mature neutrophils expressed the highest levels of *Csf3r* (Figure 6D). Analysis of *Il1b* expression in the same cell populations revealed that mature neutrophils were also a major source of *Il1b* (Figure 6D). Thus, mature neutrophils might be the cells that secrete IL-1 β upon G-CSF stimulation in the BM. In support of this notion, mature neutrophils isolated from the BM of LIP-subjected (but not control) mice responded to recombinant G-CSF stimulation with dose-dependent increase of IL-1 β production (Figure 6E). To identify possible target cells of IL-1 β in the BM, we examined the same cell populations from the BM of LIP-subjected mice and found that LSK had the most prominent *Il1r* expression

(Figure 6D). Thus, IL-1 β secretion in the BM might link LIP-induced inflammation and activation of HSPC during trained myelopoiesis. Consistently, IL-1, IL-1 β , and the myeloid transcription factor SPI1 (PU.1) were predicted as activated upstream regulators in LSK of LIP-subjected mice (Figure S1G). Moreover, the TF binding motifs enrichment analysis of the scATAC-seq using LSK revealed that the DNA binding sites of TFs responsive to IL-1 signaling pathway were significantly enriched in the LIP group, compared with control (Figures 3F and 3G).

To directly link IL-1 signaling to LIP-induced maladaptive trained myelopoiesis, we used mice with inducible deletion of IL-1 receptor specifically in HSPC, generated by breeding HSC-SCL-Cre-ERT mice (Schoedel et al., 2016) with *Il1r1*^{fl/fl} mice (HSC-SCL-Cre-ERT/*Il1r1*^{fl/fl}), hereafter designated *Il1r1*^{HSPC-KO} mice. Efficient *Il1r1* deletion was achieved by administering tamoxifen by oral gavage, as described (Schoedel et al., 2016) and detailed in STAR Methods. After induction of deletion, *Il1r1*^{HSPC-KO} and littermate controls with intact IL1-R1 expression in HSPC were subjected to 21-day LIP. The HSPC-specific deletion of IL1-R1 resulted in reduced LIP-induced myelopoiesis, as evidenced by decreased frequencies of myeloid-biased HSPC subsets (MPP3 and CD41⁺ LT-HSC), GMP, Gr1^{hi}CD11b⁺ granulocytes, and Gr1^{int}CD11b⁺ myeloid cells (Figure S7A).

Using *Il1r1*^{HSPC-KO} mice and BMT, we next tested the hypothesis that IL-1 signaling in HSPC mediates LIP-induced trained myelopoiesis and increased inflammatory disease activity. To this end, groups of CD45.2⁺ *Il1r1*^{HSPC-KO} and littermate controls with intact IL1-R1 expression in HSPC were trained by subjecting them to 21-day LIP and 14-day resolution and then used as donors for BMT to groups of lethally irradiated congenic B6.SJL (CD45.1⁺) mice (Figure 7A). Subsequently, 12 weeks post-BMT, the recipient mice were euthanized for BM and peripheral chimerism analysis or were subjected to LIP for 5 days.

At 12 weeks post-BMT, CD45.1⁺ recipients (without any further challenge) that had received *Il1r1*^{HSPC-KO} BM cells from LIP-trained donor mice exhibited a reduced proportion of Gr1⁺CD11b⁺ myeloid cells with a corresponding increase in the proportion of CD19⁺ B cells and CD3⁺ T cells in the BM (Figure 7B) and peripheral blood (Figure 7C), as compared with mice that received wild-type (WT) BM cells. Consistently, analysis of MPP subsets and downstream progenitors in the BM showed that CD45.1⁺ mice receiving *Il1r1*^{HSPC-KO} BM cells had reduced frequency of the myeloid-biased MPP3 and GMP (Figure 7D). These findings indicate a reduced myeloid bias in HSPC due to IL1-R1 deficiency, especially, as the two groups of recipient mice displayed similar frequencies of donor-derived HSPC (Figure S7B). Mature splenic neutrophils and monocytes isolated from CD45.1⁺ mice transplanted with BM cells from *Il1r1*^{HSPC-KO} mice responded with significantly lower production of IL-6 and TNF to secondary LPS challenge than their counterparts from CD45.1⁺ mice that received BM cells from littermate controls bearing IL1-R1-sufficient HSPC (Figure 7E). These data suggest a critical role for HSPC-specific IL-1R signaling in LIP-induced trained myelopoiesis, associated with a transmissible myeloid differentiation bias and production of mature myeloid cells with increased inflammatory responsiveness.

Consistent with the demonstrated trained phenotype, LIP-subjected recipient CD45.1⁺ mice that received *Il1r1*^{HSPC-KO}




Figure 7. IL-1 signaling in HSPC mediates LIP-induced maladaptive training of myelopoiesis

(A) CD45.2⁺ *Il1r1*^{HSPC-KO} and littermate controls with intact HSPC IL-1R1 expression were trained (21-day LIP and 14-day resolution) and used as donors for BMT to lethally irradiated congenic B6.SJL (CD45.1⁺) mice.

(B–D) 12 weeks post-BMT, the frequencies of indicated myeloid cells and lymphocytes in CD45⁺ cells in BM (B) and peripheral blood (C) of recipients, and the frequencies of MPP3 (in LSK) and GMP (in MyP) (D), were determined by FACS.

(E) Splenic monocytes (left) and neutrophils (right) isolated from recipient mice were stimulated with LPS (10 ng/mL) for 24 h, and IL-6 and TNF concentrations in the supernatant were measured.

(F and G) The recipient mice were subjected to 5-day LIP, as shown in (A) and assayed for bone loss (F) and abundance of gingival monocytes, neutrophils, and total CD45⁺ cells by FACS (G).

Data are means ± SD (n = 4–6 mice/group). *p < 0.05, **p < 0.01, NS, not significant versus WT littermate controls; two-tailed Student's t test. T, trained; U, untrained.

See Figure S7.

BM cells from LIP-trained donor mice showed significantly decreased bone loss (Figure 7F) and reduced infiltration of the gingival tissue with monocytes, neutrophils, and total leukocytes, relative to control CD45.1⁺ mice that received BM cells from LIP-trained WT donor mice (Figure 7G). The decreased bone loss could not be attributed to a lack of IL-1R signaling in myeloid cells in the gingival tissue, since even mice with global deletion of IL1-R1 were equally susceptible to LIP as WT littermates (Figure S7C), as shown earlier (Dutzan et al., 2018). Therefore, IL-1R signaling acts on HSPC to mediate LIP-induced maladaptive training of myelopoiesis, which exacerbates periodontal inflammation and bone loss.

DISCUSSION

We demonstrated that maladaptive training of myelopoiesis underlies the emergence of inflammatory comorbidities, exemplified here with the periodontitis-arthritis axis. Experimental periodontitis-associated systemic inflammation induced long-lasting myeloid differentiation bias in HSPC that was retained predominantly at the epigenetic level, indicating prolonged readiness for myelopoiesis induction upon future challenges. The periodontitis-induced trained phenotype was transmissible by transplantation of sorted LT-HSC to naive recipients, which displayed increased severity of arthritis upon CAIA challenge. In line with the bidirectional association of periodontitis and rheumatoid arthritis (Fuggle et al., 2016; Potempa et al., 2017), CAIA-induced alterations to BM HSPC toward a maladaptive inflammatory phenotype, which exacerbated experimental periodontitis in transplanted mice. Thus, induction of central BM-mediated TII due to an inflammatory disease increases susceptibility to another inflammatory condition (comorbidity).

Our present study results are consistent with recent clinical imaging studies based on ¹⁸F-fluorodeoxyglucose positron emission tomography/computed tomography (¹⁸F-FDG-PET/CT) (Fifer et al., 2011; Ishai et al., 2019; Van Dyke et al., 2021). The use of ¹⁸F-FDG-PET/CT revealed a correlation between metabolic activity within the periodontal tissue (surrogate of periodontal inflammation) and hematopoietic tissue activity (reflecting stimulated hematopoiesis) (Ishai et al., 2019). Although this clinical study is correlative, it suggests an inflammatory periodontitis-BM axis, resembling the one described in our preclinical model.

Similar to the hyper-responsive phenotype of myeloid cells from trained mice, upon *ex vivo* stimulation, peripheral blood neutrophils or monocytes from individuals with periodontitis respond with a higher inflammatory cytokine production than the same cells from healthy controls; this hyper-responsiveness often persists post-therapy for at least 2 months (Ling et al., 2015; Radvar et al., 2008). This apparent trained phenotype could predispose individuals with periodontitis to inflammatory comorbidities. As human periodontitis influences hematopoietic tissue activity (Ishai et al., 2019), the hyper-responsiveness of peripheral myeloid cells in periodontitis might result from epigenetically imprinted immune memory in inflammation-adapted BM HSPC, a concept that we, and others, have recently described in mice (de Laval et al., 2020; Kalafati et al., 2020; Kaufmann et al., 2018) (and this study) and humans (Cirovic et al., 2020; Moorlag et al., 2020b).

As with individuals with periodontitis, a subset of individuals with rheumatoid arthritis under clinical remission display elevated BM metabolic activity (by ¹⁸F-FDG-PET/CT) and a pro-inflammatory phenotype of circulating monocytes (Bernelot Moens et al., 2016). These findings imply that remission of arthritis does not necessarily reduce the individual's risk of a comorbid condition. A common underlying pathophysiology, involving inflammatory memory in the BM that sustains trained myelopoiesis, might thus be an overlooked factor contributing to the connection between distinct comorbid inflammatory disorders.

Studies in rodents have shown that experimental periodontitis promotes experimental arthritis and vice-versa; in these studies, the same animals were subjected to both disease models (Cantley et al., 2011; Flak et al., 2019; Ramamurthy et al., 2005; Sato et al., 2017). This approach would not allow dissecting BM-dependent mechanisms from other mechanisms contributing to the periodontitis-arthritis relationship, such as the ability of certain periodontal pathogens to cause a breakdown in immune tolerance to citrullinated epitopes, leading to the generation of arthritogenic antibodies (Konig et al., 2016; Maresz et al., 2013; Potempa et al., 2017). However, such non-mutually exclusive mechanisms do not explain the bidirectional association of periodontitis and arthritis, in contrast to the mechanism reported here.

Enduring epigenetic modifications that unfold chromatin and render promoter and enhancer regions accessible to transcription factors, constitute a major pillar of the TII concept (Fanucchi et al., 2021). In this study, TII could be transferred via BMT from trained donors to naive recipient mice, whose mature myeloid cells—12 weeks post-BMT—displayed enhanced inflammatory responsiveness. Although the transmission of the trained state by BMT could involve concurrent transmission of epigenetic modifications, this has not been formally shown in the literature. Indirectly, however, several studies have suggested a key role of epigenetic memory in TII induction and maintenance. LT-HSC retain epigenetic memory of previous inflammatory challenge and this underlies their sustained myeloid bias, i.e., persistent changes in the accessibility of specific myeloid lineage enhancers, which augments the responsiveness of the respective immune genes to secondary stimuli (de Laval et al., 2020). It is currently thought that, unlike repressive modifications (in particular DNA methylation), accessibility of enhancers and active histone modifications are not self-maintained in mammals; indeed, studies in dividing cell populations showing long-term and mitotically inheritable epigenetic changes have so far been attributed to DNA methylation changes and not histone modifications (Sun and Barreiro, 2020). Future research in dividing cells may clarify whether histone modifications are also epigenetically transmissible. In this regard, certain chromatin accessibility changes are transmitted from HSPC to progeny cells along differentiation trajectories (Buenrostro et al., 2018). Moreover, epigenetic changes in the form of histone modifications were shown to be passed on to the mouse offspring (Siklenka et al., 2015). Transgenerational transmission of various immune traits including TII, attributable to sustained epigenetic memory, has been shown recently in mice (Bomans et al., 2018; Katzmarski et al., 2021; Lim et al., 2021) and humans (Berendsen et al., 2021; Gee et al., 2021).

Based on our findings, an inflammatory disease could modulate TII in the BM in a manner that not only aggravates the pre-existing

disease but can also increase susceptibility to a distinct inflammatory condition. Such unified conceptual framework could also provide a platform for therapeutic interventions targeting inflammatory comorbidities. Our data suggest that systemic inhibition of IL-1-induced signaling may potentially block the maladaptive training of BM progenitors and thereby disrupt a common mechanism for inflammatory comorbidities. In this regard, it might be argued that the successful application of IL-1 β blockade in the CANTOS trial for the treatment of atherosclerosis (Ridker et al., 2017) might, in part, have resulted from the inhibition of TII in the BM. In conclusion, our findings establish the principle that maladaptive innate immune training of myelopoiesis underlies inflammatory comorbidities, paving the way for their treatment in a holistic manner.

Limitations of the study

Our findings have linked experimental periodontitis to maladaptive BM-mediated TII, which underlies inflammatory comorbidities. Nevertheless, our study has several limitations. Despite initial evidence for an inflammatory periodontitis-BM axis in humans and for a trained phenotype of myeloid cells from individuals with periodontitis, formal evidence for periodontitis-induced maladaptive TII in humans may require studying hematopoietic stem cell transplant (HSCT) recipients. Thus, future investigations of the transmissibility of inflammatory memory to HSCT recipients from donors with or without periodontitis would confer increased clinical relevance to our study. Such study would prompt clinicians to take inflammatory memory into consideration when selecting appropriate donors for BMT. In most of the experiments (and all functional assays), LIP lasted for 21 days to mimic chronic periodontitis in humans, although shorter durations (e.g., 7 days) were also used to determine early effects of LIP on modulating myelopoiesis. We cannot rule out the possibility that epigenetic changes seen after 21 days of LIP and 14 days resolution might differ from those occurring at earlier time points and whether such earlier changes might also contribute to a trained phenotype. Our findings that a maladaptive trained phenotype was transmissible via BMT in a HSPC-specific and IL-1-dependent manner indicated that IL-1 is critical for induction of a long-lasting trained phenotype with detrimental consequences in inflammatory disease. We have not determined if IL-1 signaling shapes the epigenetically imprinted inflammatory memory associated with a myeloid differentiation bias in trained LT-HSC; this question should be studied in the future. Moreover, we cannot exclude the possibility that IL-1 β may cooperate (or even synergize) with other as yet unidentified inflammatory factors, thus warranting additional mechanistic investigations. Despite the *de facto* transmission of the TII phenotype via HSC transplantation, the underlying molecular epigenetic mechanisms remain incompletely understood. Finally, at this point, we do not know how long a maladaptive trained phenotype would last, a question that must be carefully addressed in the future.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**
- **RESOURCE AVAILABILITY**
 - Lead contact
 - Materials availability
 - Data and code availability
- **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
 - Mice
- **METHOD DETAILS**
 - Ligature-induced periodontitis
 - Collagen antibody-induced arthritis
 - Cells preparations and sample collection
 - Flow cytometry and sorting
 - Bone marrow transplantation
 - Immunoassays
 - Quantitative real-time PCR
 - Histological analysis
 - Bulk RNA sequencing
 - Single-cell RNA sequencing
 - Single-cell ATAC sequencing
- **QUANTIFICATION AND STATISTICAL ANALYSIS**
 - Bioinformatic analysis of bulk RNA-seq
 - Single-cell RNA sequencing analysis
 - Single cell ATAC sequencing analysis
 - Statistical analysis

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.cell.2022.03.043>.

ACKNOWLEDGMENTS

This work was supported by NIH grants (DE029436 and DE031206 to G.H.; DE028561 to G.H. and T.C.) and the Deutsche Forschungsgemeinschaft (SFB-TR 127, A3 and SFB1181, C7 to T.C.). T.C. was also supported by the British Heart Foundation Centre for Research Excellence at The University of Edinburgh (RE/18/5/34216) and the “Sonderzuweisung zur Unterstützung profilbestimmender Struktureinheiten 2021” by the SMWK. M.G.N. is supported by a Spinoza grant from the Netherlands Organization for Scientific Research and a European Research Council Advanced Grant (#833247). We thank Sylvia Grossklaus (Institute for Clinical Chemistry and Laboratory Medicine, Technische Universität Dresden) for technical assistance. The graphical abstract was created using BioRender.com.

AUTHOR CONTRIBUTIONS

X.L. designed and performed research, analyzed and interpreted data, and co-wrote the manuscript; H.W. designed and performed research, analyzed and interpreted data, and contributed to writing; X.Y. performed experiments and analyzed data; G.S. performed experiments; L.K. and C.I. generated critical reagents; I.M. interpreted data; M.G.N. interpreted data and edited the manuscript; T.C. conceived and designed the study, interpreted data, and edited the manuscript; G.H. conceived and designed the study, supervised research, interpreted data, and wrote the manuscript.

DECLARATION OF INTERESTS

M.G.N. is the scientific founder and member of the Scientific Advisory Board of Trained Therapeutics and Discovery (TTxD). M.G.N. has two patents: US18/61935 (Targeted nanoimmunotherapy to increase trained immunity) and US18/61939 (Targeted nanoimmunotherapy for inhibition of trained immunity). The other authors declare no competing interests.

Received: August 11, 2021

Revised: February 22, 2022

Accepted: March 30, 2022

Published: April 27, 2022

REFERENCES

- Abe, T., and Hajishengallis, G. (2013). Optimization of the ligature-induced periodontitis model in mice. *J. Immunol. Methods* 394, 49–54.
- Bajaj, J.S., Matin, P., White, M.B., Fagan, A., Deeb, J.G., Acharya, C., Dalmat, S.S., Sikaroodi, M., Gillevet, P.M., and Sahingur, S.E. (2018). Periodontal therapy favorably modulates the oral-gut-hepatic axis in cirrhosis. *Am. J. Physiol. Gastrointest. Liver Physiol.* 315, G824–G837.
- Berendsen, M.L.T., Scholtz-Buchholzer, F., Bles, P., Biering-Sørensen, S., Jensen, K.J., Monteiro, I., Silva, I., Aaby, P., and Benn, C.S. (2021). Parental *Bacillus Calmette-Guérin* vaccine scars decrease infant mortality in the first six weeks of life: a retrospective cohort study. *EClinicalmedicine* 39, 101049.
- Bernelot Moens, S.J., van der Valk, F.M., Strang, A.C., Kroon, J., Smits, L.P., Kneepkens, E.L., Verberne, H.J., van Buul, J.D., Nurmohamed, M.T., and Stroes, E.S. (2016). Unexpected arterial wall and cellular inflammation in patients with rheumatoid arthritis in remission using biological therapy: a cross-sectional study. *Arthritis Res. Ther.* 18, 115.
- Bomans, K., Schenz, J., Tamulyte, S., Schaack, D., Weigand, M.A., and Uhle, F. (2018). Paternal sepsis induces alterations of the sperm methylome and dampens offspring immune responses – an animal study. *Clin. Epigenetics* 10, 89.
- Buenrostro, J.D., Corces, M.R., Lareau, C.A., Wu, B., Schep, A.N., Aryee, M.J., Majeti, R., Chang, H.Y., and Greenleaf, W.J. (2018). Integrated single-cell analysis maps the continuous regulatory landscape of human hematopoietic differentiation. *Cell* 173, 1535–1548.e16.
- Cantley, M.D., Haynes, D.R., Marino, V., and Bartold, P.M. (2011). Pre-existing periodontitis exacerbates experimental arthritis in a mouse model. *J. Clin. Periodontol.* 38, 532–541.
- Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D.M., Hill, A.J., Zhang, F., Mundlos, S., Christiansen, L., Steemers, F.J., et al. (2019). The single-cell transcriptional landscape of mammalian organogenesis. *Nature* 566, 496–502.
- Chavakis, T., Mitroulis, I., and Hajishengallis, G. (2019). Hematopoietic progenitor cells as integrative hubs for adaptation to and fine-tuning of inflammation. *Nat. Immunol.* 20, 802–811.
- Ciarlo, E., Heinonen, T., Théroude, C., Asgari, F., Le Roy, D., Netea, M.G., and Roger, T. (2020). Trained immunity confers broad-spectrum protection against bacterial infections. *J. Infect. Dis.* 222, 1869–1881.
- Cirovic, B., de Bree, L.C.J., Groh, L., Blok, B.A., Chan, J., van der Velden, W.J.F.M., Bremmers, M.E.J., van Crevel, R., Händler, K., Picelli, S., et al. (2020). BCG vaccination in humans elicits trained immunity via the hematopoietic progenitor compartment. *Cell Host Microbe* 28, 322–334.e5.
- D'Aiuto, F., Gkraniats, N., Bhowruth, D., Khan, T., Orlandi, M., Suvan, J., Masi, S., Tsakos, G., Hurel, S., Hingorani, A.D., et al. (2018). Systemic effects of periodontitis treatment in patients with type 2 diabetes: a 12 month, single-centre, investigator-masked, randomised trial. *Lancet Diabetes Endocrinol.* 6, 954–965.
- D'Aiuto, F., Orlandi, M., and Gunsolley, J.C. (2013). Evidence that periodontal treatment improves biomarkers and CVD outcomes. *J. Clin. Periodontol.* 40, S85–S105.
- de Laval, B., Maurizio, J., Kandalla, P.K., Brisou, G., Simonnet, L., Huber, C., Gimenez, G., Matcovitch-Natan, O., Reinhardt, S., David, E., et al. (2020). C/EBP β -dependent epigenetic memory induces trained immunity in hematopoietic stem cells. *Cell Stem Cell* 26, 657–674.E658.
- de Pablo, P., Dietrich, T., and McAlindon, T.E. (2008). Association of periodontal disease and tooth loss with rheumatoid arthritis in the US population. *J. Rheumatol.* 35, 70–76.
- Deplus, R., Delatte, B., Schwinn, M.K., Defrance, M., Méndez, J., Murphy, N., Dawson, M.A., Volkmar, M., Putmans, P., Calonne, E., et al. (2013). TET2 and TET3 regulate GlcNAcylation and H3K4 methylation through OGT and SET1/COMPASS. *EMBO J.* 32, 645–655.
- Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29, 15–21.
- Dutzan, N., Kajikawa, T., Abusleme, L., Greenwell-Wild, T., Zuazo, C.E., Ikeuchi, T., Brenchley, L., Abe, T., Hurabielle, C., Martin, D., et al. (2018). A dysbiotic microbiome triggers TH17 cells to mediate oral mucosal immunopathology in mice and humans. *Sci. Transl. Med.* 10, eaat0797.
- Fanucchi, S., Domínguez-Andrés, J., Joosten, L.A.B., Netea, M.G., and Mhlanga, M.M. (2021). The intersection of epigenetics and metabolism in trained immunity. *Immunity* 54, 32–43.
- Fifer, K.M., Qadir, S., Subramanian, S., Vijayakumar, J., Figueroa, A.L., Truong, Q.A., Hoffmann, U., Brady, T.J., and Tawakol, A. (2011). Positron emission tomography measurement of periodontal 18F-fluorodeoxyglucose uptake is associated with histologically determined carotid plaque inflammation. *J. Am. Coll. Cardiol.* 57, 971–976.
- Flak, M.B., Colas, R.A., Muñoz-Atienza, E., Curtis, M.A., Dalli, J., and Pitzalis, C. (2019). Inflammatory arthritis disrupts gut resolution mechanisms, promoting barrier breakdown by *Porphyromonas gingivalis*. *JCI Insight* 4, e125191.
- Fuggle, N.R., Smith, T.O., Kaul, A., and Sofat, N. (2016). Hand to mouth: a systematic review and meta-analysis of the association between rheumatoid arthritis and periodontitis. *Front. Immunol.* 7, 80.
- Gee, S., Chandiramani, M., Seow, J., Pollock, E., Modestini, C., Das, A., Tree, T., Doores, K.J., Tribe, R.M., and Gibbons, D.L. (2021). The legacy of maternal SARS-CoV-2 infection on the immunology of the neonate. *Nat. Immunol.* 22, 1490–1502.
- Gekas, C., and Graf, T. (2013). CD41 expression marks myeloid-biased adult hematopoietic stem cells and increases with age. *Blood* 121, 4463–4472.
- Genco, R.J., and Sanz, M. (2020). Clinical and public health implications of periodontal and systemic diseases: an overview. *Periodontol.* 2000 83, 7–13.
- Göthert, J.R., Gustin, S.E., Hall, M.A., Green, A.R., Göttgens, B., Izon, D.J., and Begley, C.G. (2005). In vivo fate-tracing studies using the Scl stem cell enhancer: embryonic hematopoietic stem cells significantly contribute to adult hematopoiesis. *Blood* 105, 2724–2732.
- Granja, J.M., Corces, M.R., Pierce, S.E., Bagdatli, S.T., Choudhry, H., Chang, H.Y., and Greenleaf, W.J. (2021). ArchR is a scalable software package for integrative single-cell chromatin accessibility analysis. *Nat. Genet.* 53, 403–411.
- Hajishengallis, G. (2015). Periodontitis: from microbial immune subversion to systemic inflammation. *Nat. Rev. Immunol.* 15, 30–44.
- Hajishengallis, G., and Chavakis, T. (2021). Local and systemic mechanisms linking periodontal disease and inflammatory comorbidities. *Nat. Rev. Immunol.* 21, 426–440.
- Hajishengallis, G., Li, X., and Chavakis, T. (2021). Immunometabolic control of hematopoiesis. *Mol. Aspects Med.* 77, 100923.
- Ishai, A., Osborne, M.T., El Kholy, K., Takx, R.A.P., Ali, A., Yuan, N., Hsue, P., Van Dyke, T.E., and Tawakol, A. (2019). Periodontal disease associates with arterial inflammation via potentiation of a hematopoietic-arterial axis. *JACC Cardiovasc. Imaging* 12, 2271–2273.
- Johnson, K., Pflugh, D.L., Yu, D., Hesslein, D.G., Lin, K.I., Bothwell, A.L., Thomas-Tikhonenko, A., Schatz, D.G., and Calame, K. (2004). B cell-specific loss of histone 3 lysine 9 methylation in the V(H) locus depends on Pax5. *Nat. Immunol.* 5, 853–861.
- Kalafati, L., Kourtzelis, I., Schulte-Schrepping, J., Li, X., Hatzioannou, A., Gri-nenko, T., Hagag, E., Sinha, A., Has, C., Dietz, S., et al. (2020). Innate immune training of granulopoiesis promotes anti-tumor activity. *Cell* 183, 771–785.e12.
- Kassebaum, N.J., Bernabé, E., Dahiya, M., Bhandari, B., Murray, C.J., and Marcenes, W. (2014). Global burden of severe periodontitis in 1990–2010: a systematic review and meta-regression. *J. Dent. Res.* 93, 1045–1053.
- Katzmarski, N., Domínguez-Andrés, J., Cirovic, B., Renieris, G., Ciarlo, E., Le Roy, D., Lepikhov, K., Kattler, K., Gasparoni, G., Händler, K., et al. (2021).

- Transmission of trained immunity and heterologous resistance to infections across generations. *Nat. Immunol.* 22, 1382–1390.
- Kaufmann, E., Sanz, J., Dunn, J.L., Khan, N., Mendonça, L.E., Pacis, A., Tzelepis, F., Pernet, E., Dumaine, A., Grenier, J.C., et al. (2018). BCG educates hematopoietic stem cells to generate protective innate immunity against tuberculosis. *Cell* 172, 176–190.e19.
- Khachigian, L.M. (2006). Collagen antibody-induced arthritis. *Nat. Protoc.* 1, 2512–2516.
- Kim, J.Y., Kim, K.B., Eom, G.H., Choe, N., Kee, H.J., Son, H.J., Oh, S.T., Kim, D.W., Pak, J.H., Baek, H.J., et al. (2012). KDM3B is the H3K9 demethylase involved in transcriptional activation of *Imo2* in leukemia. *Mol. Cell. Biol.* 32, 2917–2933.
- Kitamoto, S., Nagao-Kitamoto, H., Jiao, Y., Gilliland, M.G., III, Hayashi, A., Imai, J., Sugihara, K., Miyoshi, M., Brazil, J.C., Kuffa, P., et al. (2020). The intermucosal connection between the mouth and gut in commensal pathobiont-driven colitis. *Cell* 182, 447–462.e14.
- König, M.F., Abusleme, L., Reinholdt, J., Palmer, R.J., Teles, R.P., Sampson, K., Rosen, A., Nigrovic, P.A., Sokolove, J., Giles, J.T., et al. (2016). *Aggregatibacter actinomycetemcomitans*-induced hypercitrullination links periodontal infection to autoimmunity in rheumatoid arthritis. *Sci. Transl. Med.* 8, 369ra176.
- Kourtzelis, I., Li, X., Mitroulis, I., Grosser, D., Kajikawa, T., Wang, B., Grzybek, M., von Renesse, J., Czogalla, A., Troullinaki, M., et al. (2019). DEL-1 promotes macrophage efferocytosis and clearance of inflammation. *Nat. Immunol.* 20, 40–49.
- Lee, K.H., and Choi, Y.Y. (2020). Rheumatoid arthritis and periodontitis in adults: using the Korean National Health Insurance Service – National Sample Cohort. *J. Periodontol.* 91, 1186–1193.
- Li, X., Colamatteo, A., Kalafati, L., Kajikawa, T., Wang, H., Lim, J.-H., Bdeir, K., Chung, K.-J., Yu, X., Fusco, C., et al. (2020). The DEL-1/ β 3 integrin axis promotes regulatory T cell responses during inflammation resolution. *J. Clin. Invest.* 130, 6261–6277.
- Lim, A.I., McFadden, T., Link, V.M., Han, S.-J., Karlsson, R.-M., Stacy, A., Farley, T.K., Lima-Junior, D.S., Harrison, O.J., Desai, J.V., et al. (2021). Prenatal maternal infection promotes tissue-specific immunity and inflammation in offspring. *Science* 373, eabf3002.
- Ling, M.R., Chapple, I.L., and Matthews, J.B. (2015). Peripheral blood neutrophil cytokine hyper-reactivity in chronic periodontitis. *Innate Immun.* 21, 714–725.
- Listl, S., Galloway, J., Mossey, P.A., and Marcenes, W. (2015). Global economic impact of dental diseases. *J. Dent. Res.* 94, 1355–1361.
- Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 550.
- Mahabeleshwar, G.H., Kawanami, D., Sharma, N., Takami, Y., Zhou, G., Shi, H., Nayak, L., Jeyaraj, D., Grealy, R., White, M., et al. (2011). The myeloid transcription factor KLF2 regulates the host response to polymicrobial infection and endotoxic shock. *Immunity* 34, 715–728.
- Mann, M., Mehta, A., de Boer, C.G., Kowalczyk, M.S., Lee, K., Haldeman, P., Rogel, N., Knecht, A.R., Farouq, D., Regev, A., and Baltimore, D. (2018). Heterogeneous responses of hematopoietic stem cells to inflammatory stimuli are altered with age. *Cell Rep.* 25, 2992–3005.e5.
- Maresz, K.J., Hellvard, A., Sroka, A., Adamowicz, K., Bielecka, E., Koziel, J., Gawron, K., Mizgalska, D., Marcinska, K.A., Benedyk, M., et al. (2013). *Porphyromonas gingivalis* facilitates the development and progression of destructive arthritis through its unique bacterial peptidylarginine deiminase (PAD). *PLOS Pathog.* 9, e1003627.
- Margueron, R., Justin, N., Ohno, K., Sharpe, M.L., Son, J., Drury, W.J., 3rd, Voigt, P., Martin, S.R., Taylor, W.R., De Marco, V., et al. (2009). Role of the polycomb protein EED in the propagation of repressive histone marks. *Nature* 461, 762–767.
- Mikuls, T.R., Payne, J.B., Yu, F., Thiele, G.M., Reynolds, R.J., Cannon, G.W., Markt, J., McGowan, D., Kerr, G.S., Redman, R.S., et al. (2014). Periodontitis and *Porphyromonas gingivalis* in patients with rheumatoid arthritis. *Arthritis Rheumatol.* 66, 1090–1100.
- Mitroulis, I., Chen, L.-S., Singh, R.P., Kourtzelis, I., Economopoulou, M., Kajikawa, T., Troullinaki, M., Ziogas, A., Ruppova, K., Hosur, K., et al. (2017). Secreted protein Del-1 regulates myelopoiesis in the hematopoietic stem cell niche. *J. Clin. Invest.* 127, 3624–3639.
- Mitroulis, I., Ruppova, K., Wang, B., Chen, L.S., Grzybek, M., Grinenko, T., Eugster, A., Troullinaki, M., Palladini, A., Kourtzelis, I., et al. (2018). Modulation of myelopoiesis progenitors is an integral component of trained immunity. *Cell* 172, 147–161.e12.
- Moorlag, S.J.C.F.M., Khan, N., Novakovic, B., Kaufmann, E., Jansen, T., van Crevel, R., Divangahi, M., and Netea, M.G. (2020a). β -glucan induces protective trained immunity against *Mycobacterium tuberculosis* infection: a key role for IL-1. *Cell Rep.* 31, 107634.
- Moorlag, S.J.C.F.M., Rodriguez-Rosales, Y.A., Gillard, J., Fanucchi, S., Theunissen, K., Novakovic, B., de Bont, C.M., Negishi, Y., Fok, E.T., Kalafati, L., et al. (2020b). BCG vaccination induces long-term functional reprogramming of human neutrophils. *Cell Rep.* 33, 108387.
- Netea, M.G., Domínguez-Andrés, J., Barreiro, L.B., Chavakis, T., Divangahi, M., Fuchs, E., Joosten, L.A.B., van der Meer, J.W.M., Mhlanga, M.M., Mulder, W.J.M., et al. (2020). Defining trained immunity and its role in health and disease. *Nat. Rev. Immunol.* 20, 375–388.
- Penkov, S., Mitroulis, I., Hajishengallis, G., and Chavakis, T. (2019). Immunometabolic crosstalk: an ancestral principle of trained immunity? *Trends Immunol.* 40, 1–11.
- Peres, M.A., Macpherson, L.M.D., Weyant, R.J., Daly, B., Venturelli, R., Mathur, M.R., Listl, S., Celeste, R.K., Guarnizo-Herreño, C.C., Kearns, C., et al. (2019). Oral diseases: a global public health challenge. *Lancet* 394, 249–260.
- Perillo, B., Ombra, M.N., Bertoni, A., Cuozzo, C., Sacchetti, S., Sasso, A., Chiariotti, L., Malorni, A., Abbondanza, C., and Avvedimento, E.V. (2008). DNA oxidation as triggered by H3K9me2 demethylation drives estrogen-induced gene expression. *Science* 319, 202–206.
- Perino, M., van Mierlo, G., Karemaker, I.D., van Genesen, S., Vermeulen, M., Marks, H., van Heeringen, S.J., and Veenstra, G.J.C. (2018). MTF2 recruits polycomb repressive complex 2 by helical-shape-selective DNA binding. *Nat. Genet.* 50, 1002–1010.
- Pietras, E.M., Mirantes-Barbeito, C., Fong, S., Loeffler, D., Kovtonyuk, L.V., Zhang, S., Lakshminarasimhan, R., Chin, C.P., Techner, J.M., Will, B., et al. (2016). Chronic interleukin-1 exposure drives haematopoietic stem cells towards precocious myeloid differentiation at the expense of self-renewal. *Nat. Cell Biol.* 18, 607–618.
- Pietras, E.M., Reynaud, D., Kang, Y.A., Carlin, D., Calero-Nieto, F.J., Leavitt, A.D., Stuart, J.M., Göttgens, B., and Passegué, E. (2015). Functionally distinct subsets of lineage-biased multipotent progenitors control blood production in normal and regenerative conditions. *Cell Stem Cell* 17, 35–46.
- Pliner, H.A., Packer, J.S., McFaline-Figueroa, J.L., Cusanovich, D.A., Daza, R.M., Aghamirzaie, D., Srivatsan, S., Qiu, X., Jackson, D., Minkina, A., et al. (2018). Cicero predicts cis-regulatory DNA interactions from single-cell chromatin accessibility data. *Mol. Cell* 71, 858–871.e8.
- Potempa, J., Mydel, P., and Koziel, J. (2017). The case for periodontitis in the pathogenesis of rheumatoid arthritis. *Nat. Rev. Rheumatol.* 13, 606–620.
- Qing, Y., Yin, F., Wang, W., Zheng, Y., Guo, P., Schozer, F., Deng, H., and Pan, D. (2014). The Hippo effector Yorkie activates transcription by interacting with a histone methyltransferase complex through NcoA6. *eLife* 3.
- Radvar, M., Tavakkol-Afshari, J., Bajestan, M.N., Naseh, M.R., and Arab, H.R. (2008). The effect of periodontal treatment on IL-6 production of peripheral blood monocytes in aggressive periodontitis and chronic periodontitis patients. *Iran. J. Immunol.* 5, 100–106.
- Ramamurthy, N.S., Greenwald, R.A., Celiker, M.Y., and Shi, E.Y. (2005). Experimental arthritis in rats induces biomarkers of periodontitis which are ameliorated by gene therapy with tissue inhibitor of matrix metalloproteinases. *J. Periodontol.* 76, 229–233.

- Ridger, V.C., Wagner, B.E., Wallace, W.A., and Hellewell, P.G. (2001). Differential effects of CD18, CD29, and CD49 integrin subunit inhibition on neutrophil migration in pulmonary inflammation. *J. Immunol.* 166, 3484–3490.
- Ridker, P.M., Everett, B.M., Thuren, T., MacFadyen, J.G., Chang, W.H., Ballantyne, C., Fonseca, F., Nicolau, J., Koenig, W., Anker, S.D., et al. (2017). Antiinflammatory therapy with canakinumab for atherosclerotic disease. *N. Engl. J. Med.* 377, 1119–1131.
- Rodríguez-Lozano, B., González-Febles, J., Garnier-Rodríguez, J.L., Dadlani, S., Bustabad-Reyes, S., Sanz, M., Sánchez-Alonso, F., Sánchez-Piedra, C., González-Dávila, E., and Díaz-González, F. (2019). Association between severity of periodontitis and clinical activity in rheumatoid arthritis patients: a case-control study. *Arthritis Res. Ther.* 21, 27.
- Sato, K., Takahashi, N., Kato, T., Matsuda, Y., Yokoji, M., Yamada, M., Nakajima, T., Kondo, N., Endo, N., Yamamoto, R., et al. (2017). Aggravation of collagen-induced arthritis by orally administered *Porphyromonas gingivalis* through modulation of the gut microbiota and gut immune system. *Sci. Rep.* 7, 6955.
- Schenkein, H.A., Papapanou, P.N., Genco, R., and Sanz, M. (2020). Mechanisms underlying the association between periodontitis and atherosclerotic disease. *Periodontol.* 2000 83, 90–106.
- Scher, J.U., Bretz, W.A., and Abramson, S.B. (2014). Periodontal disease and subgingival microbiota as contributors for rheumatoid arthritis pathogenesis: modifiable risk factors? *Curr. Opin. Rheumatol.* 26, 424–429.
- Schoedel, K.B., Morcos, M.N.F., Zerjatke, T., Roeder, I., Grinenko, T., Voehringer, D., Göthert, J.R., Waskow, C., Roers, A., and Gerbaulet, A. (2016). The bulk of the hematopoietic stem cell population is dispensable for murine steady-state and stress hematopoiesis. *Blood* 128, 2285–2296.
- Shin, J., Maekawa, T., Abe, T., Hajishengallis, E., Hosur, K., Pyaram, K., Mitroulis, I., Chavakis, T., and Hajishengallis, G. (2015). DEL-1 restrains osteoclastogenesis and inhibits inflammatory bone loss in nonhuman primates. *Sci. Transl. Med.* 7, 307ra155.
- Siklenka, K., Erkek, S., Godmann, M., Lambrot, R., McGraw, S., Lafleur, C., Cohen, T., Xia, J., Suderman, M., Hallett, M., et al. (2015). Disruption of histone methylation in developing sperm impairs offspring health transgenerationally. *Science* 350, aab2006.
- Smolen, J.S., Aletaha, D., Barton, A., Burmester, G.R., Emery, P., Firestein, G.S., Kavanaugh, A., McInnes, I.B., Solomon, D.H., Strand, V., and Yamamoto, K. (2018). Rheumatoid arthritis. *Nat. Rev. Dis. Primers* 4, 18001.
- Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive integration of single-cell data. *Cell* 177, 1888–1902.e21.
- Stuart, T., Srivastava, A., Lareau, C., and Satija, R. (2020). Multimodal single-cell chromatin analysis with Signac. *bioRxiv*. <https://doi.org/10.1101/2020.11.09.373613>.
- Sun, S., and Barreiro, L.B. (2020). The epigenetically-encoded memory of the innate immune system. *Curr. Opin. Immunol.* 65, 7–13.
- Tsukasaki, M., Komatsu, N., Nagashima, K., Nitta, T., Pluemsakunthai, W., Shukunami, C., Iwakura, Y., Nakashima, T., Okamoto, K., and Takayanagi, H. (2018). Host defense against oral microbiota by bone-damaging T cells. *Nat. Commun.* 9, 701.
- Van Dyke, T.E., Kholy, K.E., Ishai, A., Takx, R.A.P., Mezue, K., Abohashem, S.M., Ali, A., Yuan, N., Hsue, P., Osborne, M.T., and Tawakol, A. (2021). Inflammation of the periodontium associates with risk of future cardiovascular events. *J. Periodontol.* 92, 348–358.
- Verploegen, S., Ulfman, L., van Deutekom, H.W., van Aalst, C., Honing, H., Lammers, J.W., Koenderman, L., and Coffey, P.J. (2005). Characterization of the role of CaMKI-like kinase (CKLiK) in human granulocyte function. *Blood* 106, 1076–1083.
- Wang, H., Li, X., Kajikawa, T., Shin, J., Lim, J.H., Kourtzelis, I., Nagai, K., Korostoff, J.M., Grossklaus, S., Naumann, R., et al. (2021). Stromal cell-derived DEL-1 inhibits Tfh cell activation and inflammatory arthritis. *J. Clin. Invest.* 131, e150578.
- Winning, L., and Linden, G.J. (2017). Periodontitis and systemic disease: association or causality? *Curr. Oral Health Rep.* 4, 1–7.
- Yuh, D.Y., Maekawa, T., Li, X., Kajikawa, T., Bdeir, K., Chavakis, T., and Hajishengallis, G. (2020). The secreted protein DEL-1 activates a $\beta 3$ integrin-FAK-ERK1/2-RUNX2 pathway and promotes osteogenic differentiation and bone regeneration. *J. Biol. Chem.* 295, 7261–7273.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
FITC anti-mouse Lineage Cocktail	Biolegend	Cat#133301
Biotin anti-mouse Lineage Panel	Biolegend	Cat#133307
Rat anti-mouse cKit (CD117)	Biolegend	Cat#105808; RRID: AB_313217
Rat anti-mouse Sca1 (Ly6-A/E)	Biolegend	Cat#122514; RRID: AB_756199
Armenian Hamster anti-mouse CD48	Biolegend	Cat#103426; RRID: AB_10612755
Rat anti-mouse CD150	Biolegend	Cat#115922; RRID: AB_2303663
Rat anti-mouse CD16/CD32	Biolegend	Cat#101324; RRID: AB_1877267
Armenian Hamster anti-mouse CD34	Biolegend	Cat#128612; RRID: AB_10553896
Rat anti-mouse CD135	Biolegend	Cat#135310, 135315; RRID: AB_2107050, AB_2571919
Rat anti-mouse CD41	Biolegend	Cat#133912; RRID: AB_2650893
Armenian Hamster anti-mouse CD61	Biolegend	Cat#104316; RRID: AB_2561734
Rat anti-mouse CD11b	BD Pharmingen™	Cat# 552850; RRID: AB_394491
Rat anti-mouse Gr-1 (Ly-6G/C)	eBioscience™	Cat#48-5931-82; RRID: AB_1548788
Rat anti-mouse CD45	Biolegend	Cat#103132; RRID: AB_893340
Rat anti-mouse CD19	Biolegend	Cat#115508; RRID: AB_313643
Rat anti-mouse CD3	Biolegend	Cat#100236; RRID: AB_2561456
Mouse anti-mouse CD45.1	Biolegend	Cat#110728; RRID: AB_893346
Mouse anti-mouse CD45.2	Biolegend	Cat#109828; RRID: AB_893350
Rat anti-mouse Ly6G	Biolegend	Cat#127614, 127604; RRID: AB_2227348, AB_1186108
Rat anti-mouse Ly6C	Biolegend	Cat#128008, 128004; RRID: AB_1186132, AB_1236553
Armenian Hamster anti-mouse CD11c	Biolegend	Cat#117306; RRID: AB_313775
Rat anti-mouse CD115	Biolegend	Cat#135526; RRID: AB_2566462
Rat anti-mouse CXCR4	Biolegend	Cat#146508; RRID: AB_2562785
Rat anti-mouse CXCR2	Biolegend	Cat#149610; RRID: AB_2565690
Arthritogenic monoclonal antibodies (5-clone collagen antibody cocktail)	Chondrex	Cat#53010
Chemicals, Enzymes, Reagents and recombinant proteins		
Collagenase IV	Gibco	Cat#17104019; CAS: 9001-12-1
ACK Lysing Buffer	Gibco	Cat#A1049201
Percoll gradient	GE Healthcare	Cat#17-0891-01
Tamoxifen	Sigma-Aldrich	Cat#T5648
Tamoxifen-containing (0.5 mg/g) diet	Envigo	Cat#TD.130857
Nonidet P40 Substrate	Sigma-Aldrich	Cat#74385
Nuclei Buffer	10x Genomics	Cat#2000153
Streptavidin microBeads	Miltenyi Biotec	Cat# 130-048-101
Anti-biotin microbeads	Miltenyi Biotec	Cat# 130-090-485
LPS-EB (LPS from <i>E. coli</i> O111:B4)	Invivogen	Cat# tlr1-ebmps
DAPI (4',6-Diamidino-2-Phenylindole, Dihydrochloride)	ThermoFisher Scientific	Cat# D1306
TRIzol™ Reagent	Invitrogen	Cat# 15596026
Mouse recombinant G-CSF	R&D Systems	Cat# 414-CS-005/CF

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Critical commercial assays		
IL-6 mouse ELISA kit	Invitrogen	Cat# 88-7064-88
IL-1 β mouse ELISA kit	Invitrogen	Cat# 88-7013-88
IFN- γ mouse ELISA kit	Invitrogen	Cat# 88-7314-88
IL-12p70 mouse ELISA kit	Invitrogen	Cat# 88-7121-88
IL-10 mouse ELISA kit	Invitrogen	Cat# 88-7105-88
Mouse G-CSF ELISA	RayBiotech	Cat# ELM-GCSF-1
High-Capacity RNA-to-cDNA Kit	Applied Biosystems	Cat# 4387406
TURBO DNA-free Kit	Invitrogen	Cat# AM1907
Chromium Next GEM Chip H Single Cell Kit v1.1	10x Genomics	Cat#1000162
Chromium Next GEM Single Cell ATAC Library & Gel Bead Kit v1.1	10x Genomics	Cat#1000176
NGS High Sensitivity Fragment Analysis Kit	Agilent	Cat#DNF-474
NovaSeq 6000 SP Reagent Kit v1.5 (100 cycles)	Illumina	Cat#20028401
Chromium Next GEM Single Cell 3' Kit v3.1, 16 rxns	10x Genomics	Cat# PN-1000268
ATAC-Seq Kit	Active Motif	Cat#53150
Deposited data		
RNA sequencing data	This paper	GEO: GSE180002
Single cell ATAC sequencing data	This paper	GEO: GSE180025
Single cell RNA sequencing data	This paper	GEO: GSE196808
Experimental models: Organisms/strains		
Mouse: C57BL/6	The Jackson Laboratory	Stock#000664
Mouse: C57BL/6-CD45.1 B6.SJL-Ptprca Pepcb/BoyJ	The Jackson Laboratory	Stock#002014
Mouse: B6.129(Cg)-Il1r1tm1.1Rbl/J (Il1r1 ^{fl/fl} mice)	The Jackson Laboratory	Stock#028398
Mouse: HSC-SCL-Cre-ERT	Donated by Dr. Joachim R. Göthert, University Hospital Essen	N/A
Software and algorithms		
GraphPad Prism 8	Graphpad Software	N/A
STAR	Dobin et al. (2013)	http://code.google.com/p/rna-star/
Ingenuity Pathway Analysis	QIAGEN	https://www.qiagenbioinformatics.com/products/ingenuity-pathway-analysis/
DESeq2	Love et al. (2014)	https://bioconductor.org/packages/release/bioc/html/DESeq2.html
NovoExpress software	ACEA Biosciences	https://www.aceabio.com
Cell Ranger ATAC version 1.2.0	10x Genomics	https://support.10xgenomics.com/single-cell-atac/software/overview/welcome
R package ArchR v1.0.1	Granja et al., 2021	https://github.com/GreenleafLab/ArchR/releases/tag/v1.0.1
R package Seurat	Stuart et al., 2019	https://github.com/satijalab/seurat/releases/tag/v3.0.0
R package Signac	Stuart et al., 2020	https://github.com/timoast/signac/releases/tag/1.2.1
Monocle 3 v1.0.0	Cao et al., 2019	https://github.com/cole-trapnell-lab/monocle3/releases/tag/1.0.0
Cicero v1.4	Pliner et al., 2018	https://github.com/stjude/CICERO/releases/tag/v1.4.0

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact George Hajishengallis (geoh@upenn.edu).

Materials availability

This study did not generate any unique reagents.

Data and code availability

Data are available upon request to the Lead Contact. Sequencing data are available at the Gene Expression Omnibus database (<http://www.ncbi.nlm.nih.gov/geo/>) under the accession number GSE180032.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mice

C57BL/6 and congenic C57BL/6.SJL CD45.1⁺ male mice (B6.SJL-*Ptprca*^a*Peprc*^b/BoyJ) were purchased from the Jackson Laboratory. Mice with tamoxifen-inducible deletion of the IL-1 receptor specifically in HSPC, designated HSC-SCL-Cre-ERT/*Il1r1*^{fl/fl} (hereafter referred to as *Il1r1*^{HSPC-KO} mice) were generated by crossing using *Il1r1*^{fl/fl} mice (Stock # 028398; Jackson Laboratory) and HSC-SCL-Cre-ERT mice (Göthert et al., 2005; Schoedel et al., 2016) (donated by Dr. Joachim R. Göthert, University Hospital Essen). To induce *Il1r1* deletion, *Il1r1*^{HSPC-KO} mice were administered tamoxifen (0.3 mg/g body weight by oral gavage) two times within 72h, followed by feeding the mice with tamoxifen-containing (0.5 mg/g) diet for 18 days. Mice were maintained in individually ventilated cages under specific pathogen-free conditions on a standard 12-h light/dark cycle. Food and water were provided ad libitum. The mice were 8- to 10-week-old at the start of the experiments. Animal experiments were approved by the Institutional Animal Care and Use Committee (IACUC) of the University of Pennsylvania and were performed in compliance with institutional, state, and federal policies.

METHOD DETAILS

Ligature-induced periodontitis

Ligature-induced periodontitis (LIP) simulates human periodontitis by generating a local biofilm-retentive milieu leading to inflammation and bone loss (Abe and Hajishengallis, 2013; Dutzan et al., 2018; Kitamoto et al., 2020; Kourtzelis et al., 2019; Tsukasaki et al., 2018). To study the effect of experimental periodontitis on BM hematopoietic progenitor cells, LIP was performed in mice as previously described (Abe and Hajishengallis, 2013). Briefly, 5-0 silk ligatures were tied around the left and right maxillary second molar teeth for time intervals specified in the figure legends (up to 21 days). The same teeth were left unligated in control mice. In some experiments, the placement of ligatures was followed by their removal (for 14 days in most experiments) to enable inflammation resolution (Kourtzelis et al., 2019; Li et al., 2020) prior to BM analysis. In a subset of these experiments, mice during resolution were subjected to a secondary challenge, namely *Escherichia coli* O111:B4 LPS (Invivogen) which was injected i.p. at 1.5 mg/kg body weight, modeling bacteremia. The mice were euthanized 72h after the LPS injection for analysis. To determine ligature-induced bone loss, the 5-day standard LIP model was used, in which only the left maxillary second molar was ligated, whereas the contralateral (right) molar tooth was left unligated to serve as baseline control for measuring bone loss as we previously described (Abe and Hajishengallis, 2013; Dutzan et al., 2018; Kourtzelis et al., 2019).

Collagen antibody-induced arthritis

Collagen antibody-induced arthritis was induced in mice by i.v. (retro-orbital) injection of 1.5 mg arthritogenic monoclonal antibodies (5-clone collagen antibody cocktail; Chondrex) (Khachigian, 2006; Wang et al., 2021). Three days later, mice were injected i.p. with 50 µg of LPS. Clinical symptoms of arthritis were daily evaluated visually for each paw using a semiquantitative scoring system graded on a scale of 0–4 per paw (Khachigian, 2006) by a blinded procedure: 0 for normal; 1 for mild redness, slight swelling of ankle or wrist; 2 for moderate swelling of ankle or wrist; 3 for severe swelling, including some digits, ankle and foot; 4 for maximally inflamed joint. The clinical score for each mouse was the sum of the 4 paw scores for a maximum score of 16. Hind ankle joint thickness was measured by using J 15 pocket dial thickness gauge (Käfer).

Cells preparations and sample collection

For BM single-cell suspension preparation, femoral bones of C57BL/6 mice were flushed with ice-cold PBS (Gibco) supplemented with 5% FBS (Gibco). Cells were forced through 70-µm nylon cell strainer to get single-cell suspension for further flow cytometric analysis and FACS cell sorting. To isolate BM mature neutrophils, BM cells were flushed from femurs of C57BL/6 mice subjected to LIP for 14 days. Upon lysis of red blood cells with ACK Lysing Buffer (Gibco), the cells were centrifuged in 62% Percoll gradient (GE Healthcare). The sharp interface atop the 62% Percoll (containing immature cells and non-granulocytic lineages) was carefully removed and

discarded, whereas the pellet (mature neutrophils) was collected. To collect BM extracellular fluid, mice femurs were flushed with 500 μ l ice-cold PBS (Gibco) and the supernatant was harvested after centrifugation at 500 x g for 5 min at 4°C. Serum was collected after retrobulbar bleeding. Specifically, collected whole blood was left undisturbed at room temperature for 30 mins and the clot was removed by centrifugation at 2,000 x g for 10 minutes at 4°C followed by collection of the supernatant serum.

Mice subjected to LIP and a secondary challenge with LPS (see above) were euthanized and lung lobes were dissected after tracheal and intracardial perfusion with cold PBS. Lung tissues were then cut into 1-mm² pieces with scissor and digested in freshly prepared digestion medium consisting of 2 mg/ml collagenase IV (Invitrogen) in RPMI 1640 medium (Invitrogen) supplemented with 1% penicillin-streptomycin (Invitrogen). The digestion was performed at 37°C with shaking at 100 rpm for 1h and was stopped by addition of EDTA (Invitrogen) at a final concentration of 5mM. Cells were forced through 100- μ m nylon cell strainer and subjected to 40% Percoll density gradient centrifugation (GE Healthcare) to remove debris. The cell pellet was resuspended in ACK Lysing Buffer (Gibco) for lysis of red blood cells. Cells were washed and resuspended for immunofluorescence staining and flow cytometric analysis.

To isolate splenic neutrophils and monocytes, splenocytes were incubated with biotinylated anti-mouse Ly6G antibody (clone 1A8; Biolegend) followed by anti-biotin microbeads from Miltenyi Biotec. Neutrophils were positively selected using LS columns on the magnetic field of QuadroMACS™ Separator according to the manufacturer's instructions (Miltenyi Biotec). For isolating splenic monocytes, neutrophils were first removed by negative selection for Ly6G⁺ cells and then monocytes were obtained by positive selection for Ly6C⁺ (clone HK1.4; Biolegend) cells as we previously described (Kalafati et al., 2020).

Flow cytometry and sorting

Flow cytometric analysis was performed on a NovoCyte flow cytometer (ACEA Biosciences). For cell surface phenotypic analysis, a lineage (Lin) cocktail, including the following monoclonal antibodies was used: CD3e (clone 145-2C11), CD11b (clone M1/70), Gr1 (clone RB6-8C5), B220 (clone RA3-6B2) and TER119 (clone TER-119). Other antibody reagents used in experiments included anti-Sca1 (clone E13-161.7), anti-cKit (clone 2B8), anti-CD135 (clone A2F10), anti-CD48 (clone HM48-1), anti-CD150 (clone TC15-12F12.2), anti-CD41 (clone MWR30), anti-CD61 (clone 2C9.G2 [HM β 3-1]), anti-CD16/CD32 (clone 93), anti-CD34 (clone HM34), anti-CD45.1 (clone A20), anti-CD45.2 (clone 104), anti-CD3 (clone 17A2), anti-CD19 (clone 6D5), anti-CD11b (clone M1/70), anti-Gr1 (clone RB6-8C5), anti-CD11c (clone N418), anti-Ly6C (clone HK1.4), anti-Ly6G (clone 1A8), anti-CD115 (clone AFS98), anti-CXCR2 (clone SA045E1) and anti-CXCR4 (clone L276F12) were used. Data were analyzed with NovoExpress® software (ACEA Biosciences). Gating strategies for hematopoietic stem and progenitor cells were as follows: LSK, Lin⁻Sca-1⁺cKit⁺; LS⁻K (MyP), Lin⁻Sca-1⁻cKit⁺; LT-HSC, CD48⁻CD150⁺LSK; CD41⁻LT-HSC, CD48⁻CD150⁺CD41⁻LSK; CD41⁺LT-HSC, CD48⁻CD150⁺CD41⁺LSK; ST-HSC, CD48⁻CD150⁻LSK; MPP, CD48⁺CD150⁻LSK; MPP2, Flt3⁻CD48⁺CD150⁺LSK; MPP3, Flt3⁻CD48⁺CD150⁻LSK; MPP4, Flt3⁺CD48⁺CD150⁻LSK; GMP, Lin⁻Sca-1⁻cKit⁺CD16/32⁺CD34⁺; CMP, Lin⁻Sca-1⁻cKit⁺CD16/32⁻CD34⁺. Gating strategies for neutrophils and monocytes in mouse gingiva and knee joints were as follows: neutrophils, live CD45⁺CD11c⁻CD11b⁺Ly6G⁺Ly6C⁻; monocytes, live CD45⁺CD11c⁻CD11b⁺Ly6G⁻Ly6C⁺. For BM cell sorting, the following gating strategies were used: for pre-neutrophils, CD11b⁺CD115⁻Gr-1⁺cKit⁺CXCR4⁺; immature neutrophils, CD11b⁺CD115⁻Gr-1⁺cKit⁺CXCR4⁻Ly6G⁺CXCR2⁻; mature neutrophils, CD11b⁺CD115⁻Gr-1⁺cKit⁺CXCR4⁻Ly6G⁺CXCR2⁺; monocytes, CD11b⁺Ly6G⁻Ly6C⁺. Cells sorting was performed on a FACSARIA™ instrument (Becton Dickinson Immunocytometry Systems, USA).

Bone marrow transplantation

To generate BM chimeras, a total of 2x10⁶ BM cells from C57BL/6 (CD45.2⁺) mice, or from CD45.2⁺ *Il1r1*^{HSPC-KO} and littermate controls were transplanted (via a single retro-orbital injection) into lethally (9.5 Gy) irradiated B6/SJL (CD45.1⁺) mice. Twelve weeks after BM transplantation, the recipient mice were subjected to treatments and/or analyses as described in results and Figure legends. The CD45.1/CD45.2 congenic system was also used in competitive BM chimeras, e.g., to assess lineage output of LT-HSC in transplanted mice. Sorted LT-HSC (100 cells per recipient) isolated from the BM of CD45.2⁺ mice (which were subjected to 21-day LIP and 14-day resolution or were left untreated for 35 days) were retro-orbitally transferred into lethally (9.5 Gy) irradiated naïve B6/SJL (CD45.1) recipients along with 5x10⁵ CD45.1⁺ competitor cells. The percentage of different CD45.2⁺ cell populations was assessed at 4, 8, 12 and 16 weeks post transplantation in the blood of recipient mice, and at 16 weeks post transplantation (end point) in the BM of recipient mice.

In additional competitive BM chimera experiments, sorted LT-HSC (total or further sorted into CD41⁺LT-HSC and CD41⁻LT-HSC) isolated from the BM of CD45.2⁺ mice (which were subjected to 21-day LIP and 14-day resolution or were left untreated for 35 days) were retro-orbitally transferred (200 cells per recipient) into lethally (9.5 Gy) irradiated naïve B6/SJL (CD45.1) recipients along with 8x10⁵ CD45.1⁺ competitor cells. In all BMT experiments, irradiated recipient mice were kept on antibiotics-containing water for 2 weeks after irradiation. Complete blood count (CBC) test was performed using Sysmex XT-2000iV Hematology Analyzer.

Immunoassays

The concentrations of mouse IL-6, IL-1 β , IFN- γ , IL-12p70 and IL-10 in BM fluid and serum were measured using mouse ELISA kits (Invitrogen) according to the manufacturer's instructions. Mouse G-CSF was measured using ELISA kit from RayBiotech, according to the manufacturer's instructions. For *in vitro* stimulation of mature neutrophils from the BM, isolated mature neutrophils were stimulated with different concentration of mouse recombinant G-CSF (R&D Systems) for 24h. The supernatant was collected for

measuring IL-1 β concentration using mouse ELISA kit (Invitrogen). For *in vitro* re-stimulation of splenic monocytes and neutrophils with LPS, isolated splenic monocytes and neutrophils were seeded into 24-wells plates and stimulated with 10ng/ml of *E. coli* O111:B4 LPS (Invivogen) for 24h. The supernatant was collected for measuring IL-6 and TNF concentration using mouse ELISA kit (Invitrogen).

Quantitative real-time PCR

Total cellular RNA was isolated from mouse tissues and sorted BM cells using Trizol (Invitrogen). For real-time PCR, 500 ng of total RNA was reverse-transcribed using High-Capacity RNA-to-cDNA Kit (Applied Biosystems) and real-time PCR with cDNA was performed using the Applied Biosystems 7500 Fast Real-Time PCR System, according to the manufacturer's protocol (Applied Biosystems). TaqMan probes and gene-specific primers for detection and quantification of murine genes investigated in this study were purchased from Thermo-Fisher Scientific (Table S1). Data were analyzed using the comparative ($\Delta\Delta C_t$) method and were normalized to *Gapdh* mRNA. In LIP experiments assessing gingival cytokine mRNA expression in ligated sites, the data are shown as fold change relative to the contralateral unligated control sites (baseline), set as 1.

Histological analysis

Harvested joints were fixed in 4% buffered formaldehyde. The bones were subsequently decalcified, embedded in Optimal Cutting Temperature (OCT) media and sectioned at 10- μ m thickness for hematoxylin and eosin (H&E) or Safranin-O staining (ScienCell) (Wang et al., 2021).

Bulk RNA sequencing

LSK from the BM of mice subjected to 7-day LIP or unligated control mice were sorted using a FACSARIA™ sorter. Three mice were pooled for each replicate. Total RNA was extracted by Trizol (Invitrogen) and DNA was removed by using TURBO DNA-free Kit (Invitrogen). PolyA-selected mRNA libraries were generated following the manufacturer's protocols (BGI). Samples were sequenced on BGISEQ platform to generate 100 bp paired-end reads with an average depth of 45.0 M reads per sample. Clean reads were mapped to the mouse genome (Ensembl assembly GRCm38) using Spliced Transcripts Alignment to a Reference (STAR) (Dobin et al., 2013) with default settings after filtering low-quality, adaptor-polluted and high content of unknown base (N) reads. The average mapping ratio with reference genome was 96.46%, the average mapping ratio with gene was 80.83%. A total of 16,403 genes were detected. Most transcripts were completely covered, and reads were evenly distributed throughout the transcript.

Single-cell RNA sequencing

LSK from the BM of mice which were subjected to 21-day LIP and 14-day resolution, or were left untreated for 35 days, were sorted using a FACSARIA™ sorter. The cells were resuspended in 1X PBS containing 0.04% BSA (400 μ g/ml) and processed with Chromium Next GEM Single Cell 3' Kit v3.1, 16 rxns (10x Genomics) at the CAG Sequencing Core of Children's Hospital of Philadelphia, USA. Cell suspensions contained more than 90% viable cells as determined by microscopy. The cell numbers were adjusted to 700–1200 cells per μ l and added to 10x Chromium RT mix to achieve loading target numbers of around 20000 cells. cDNA synthesis was performed per the manufacturer's instructions, and library preparation and sequencing (Novaseq SP reagent kit; 100 cycles) were performed using the Novaseq 6000 platform (Illumina) per the manufacturer's instructions.

Single-cell ATAC sequencing

Mouse LSK and GMP were sorted as described above. Nuclei isolation for single-cell ATAC sequencing was performed according to the "Nuclei Isolation for Single Cell ATAC Sequencing demonstrated protocol" (10x Genomics). Briefly, after sorting, cells were centrifuged at 300 rcf for 5 min at 4°C and washed once with PBS with 0.04% BSA. One hundred μ L of chilled Lysis buffer (10mM Tris-HCl (pH 7.4), 10mM NaCl, 3mM MgCl₂, 0.1% Tween-20, 0.1% Nonidet P40 Substrate, 0.01% Digitonin, 1% BSA in nuclease free water) was added to the pellet and cells were incubated on ice for 2 min for LSK and 3 min for GMP. Cells were washed once with 1 ml of wash buffer (10mM Tris-HCl (pH 7.4), 10mM NaCl, 3mM MgCl₂, 0.1% Tween-20, 1% BSA in nuclease free water) and concentrated by centrifugation at 500 rcf for 5 min at 4°C. Nuclei in pellet were suspended in 1x Nuclei Buffer (10x Genomics); nuclei quality and concentration were determined using a microscope and a Countess II FL Automated Cell Counter (Thermo Fisher). Nuclei suspension was then diluted and used according to the Chromium Single Cell ATAC Reagent Kits protocol (10x Genomics). Briefly, nuclei suspension was mixed with the tagmentation mix and incubated for 1 hr at 37°C. After mixing with a barcoding mix, the nuclei were loaded into a 10x chip H together with barcoded beads and partitioning oil (Chromium Next GEM Chip H Single Cell Kit v1.1, Chromium Next GEM Single Cell ATAC Library & Gel Bead Kit v1.1, 10x Genomics) and encapsulated using the Chromium controller (10x Genomics). The Gel Bead-In EMulsions (GEMs) was transferred into a PCR tube and amplified for 12 cycles in a thermocycler. The barcoded DNA was purified and subjected to an index PCR for 11 cycles. The library amplification was assessed using fragment analyzer (Agilent, NGS High Sensitivity Fragment Analysis Kit) and sequenced on an Illumina NovaSeq 6000 system using S1 Reagent Kit v1.5 (100 cycles) in PE mode (50x8x16x50 read lengths), at a median depth of 7500 fragments/cell.

QUANTIFICATION AND STATISTICAL ANALYSIS

Bioinformatic analysis of bulk RNA-seq

Reads count in each gene were calculated by htseq-count package. Subsequently, differential gene expression was analyzed using DESeq2 (Love et al., 2014) and significantly differentially expressed genes (DEGs) were defined by FDR-adjusted P value < 0.05 in the LIP-subjected group relative to their expression in the control group. Significantly up- or down-regulated DEGs were subjected to Gene Ontology (GO) enrichment analyses using PANTHER (Protein Analysis THrough Evolutionary Relationships) (<http://www.pantherdb.org>) and KEGG pathway analysis (<https://www.genome.jp/kegg/kaas/>) with default background and default threshold. Significantly enriched Biological Process GO terms and KEGG pathways were defined by FDR-adjusted P value < 0.05 . DEGs were also subjected to upstream regulator analysis using Ingenuity Pathway Analysis (IPA, QIAGEN Redwood City <http://qiagen.com/ingenuity>). Top significantly activated transcription factors and upstream regulators are shown. Volcano plot, principal component analysis (PCA) plot, bar plots and heat maps were generated by customized R script. Overlapping genes, e.g., *Csfr1* and *Itgb3*, in heatmaps of two different GO terms are shown in both terms.

Single-cell RNA sequencing analysis

Cell Ranger 3.0.2 (<https://support.10xgenomics.com/single-cell-gene-expression/software/overview/welcome>) was used to process raw paired-end sequencing data. Briefly, Cell Ranger mkfastq pipeline was used to demultiplex sample index reads to generate FASTQ files for each sequencing library, and then raw reads were aligned to the UCSC mouse reference genome (mm10) using STAR aligner with default parameters. Subsequently, data filtering, normalization, scaling, and Principal component analysis (PCA) were performed using the R package Seurat version 4.0.3 (Stuart et al., 2019) and dimensionality reduction and clustering were further done by Uniform Manifold Approximation and Projection (UMAP) analysis. Low-quality cells and possible cell doublets were filtered out using the following criteria: (i) number of detected genes between 200 and 2500, and (ii) percentage of UMIs derived from mitochondrial genes below 5%. The UMI counts were normalized by library size factors. As a result, Cell Ranger recovered total 7051 (NL) and 8206 (LSK-21dL/14dR) LSK. Genes differentially expressed across clusters were identified using likelihood ratio test. GO enrichment analysis of cluster 7-specific markers were performed using PANTHER. Heat maps and bubble plots were generated by customized R script.


Single cell ATAC sequencing analysis

Cell Ranger ATAC 1.2.0 (<https://support.10xgenomics.com/single-cell-atac/software/overview/welcome>) was used to process raw sequencing data. Cell Ranger ATAC's pipelines was used to align reads, generate peak matrix with single-cell accessibility counts and fragment file with all unique fragments across all single cells, using 'refdata-cellranger-atac-mm10-1.2.0' from the 10X Genomics website (<https://support.10xgenomics.com/single-cell-atac/software/downloads/>) as reference genome file. The following analysis of the scATAC-seq data was performed using the R package Seurat and Signac (Stuart et al., 2019, 2020). Briefly, a chromatin accessibility matrix was created by merging fragment files from 4 samples and filtering out peaks with peak width larger than 10000 bp or smaller than 500 bp. Next, Uniform Manifold Approximation and Projection (UMAP) based on latent semantic indexing (LSI) was generated to visualize the data structure in the two-dimensional space (designated LSK-NL, LSK-21dL/14dR, GMP-NL and GMP-21dL/14dR). Smart local moving (SLM) approach implemented in the R package Seurat (Stuart et al., 2019) was used to cluster the single-cell accessibility profiles. A logistic regression framework was used for differentially accessible regions (DAR) analysis ($\text{abs}(\text{Log}_2\text{FC}) \geq 0.2$, $P < 0.05$). GO enrichment analysis of the treatment-specific DAR of LSK were performed using PANTHER (Protein Analysis THrough Evolutionary Relationships) (<http://www.pantherdb.org>). Additionally, a gene activity matrix was created to quantify the activity of each gene by assessing the chromatin accessibility using Signac package. For motif enrichment analysis, an ArchR project was created using ArchR v1.0.1 (Granja et al., 2021), and peak regions were annotated with homer motif set. Moreover, the *peakAnnoEnrichment* function implemented in ArchR was used to identify the enriched transcription factor binding motifs based on the treatment-specific DARs in LSK with $\text{FDR} < 0.01$ and $\text{log}_2\text{FC} \geq 1$, followed by GO enrichment analysis using PANTHER. A cellular trajectory was constructed spanning from LSK over cluster C9 to GMP using Monocle 3 and Cicero tool (Cao et al., 2019; Pliner et al., 2018), based on the chromatin accessibility.

Statistical analysis

After confirming normality, data were analyzed with two-tailed unpaired Student's t test (comparison of only two groups) or one-way ANOVA followed by Dunnett's multiple-comparisons test (when comparing more than two groups). In a few instances (comparison of only two groups) where data did not follow normal distribution, the non-parametric two-tailed Mann-Whitney U-test was used. Two-way repeated measures ANOVA and Sidak's multiple-comparisons test was used to analyze data in repeated-measures designs. All statistical analyses were performed using GraphPad Prism software (version 8.4.3; GraphPad Inc). P values < 0.05 were considered to be statistically significant.

Supplemental figures



(legend on next page)

Figure S1. LIP activates bone marrow and enhances myelopoiesis, related to Figure 1

Groups of mice were subjected, or not (control), to ligature-induced periodontitis (LIP) followed by BM analysis after 7 days.

(A) (Left) Representative FACS plots for identification of hematopoietic stem and progenitor cells. After gating for Lin⁻ cells, LSK were identified as cKit⁺Sca1⁺ cells. LSK subpopulations were further characterized as MPP (CD48⁺CD150⁻LSK), ST-HSC (CD48⁻CD150⁻LSK), and LT-HSC (CD48⁻CD150⁺LSK) cells. (Right) Frequencies of LSK, LT-HSC, ST-HSC, and MPP in total BM cells (top) and absolute cell numbers (bottom) of the same populations.

(B) (Top) Representative FACS plots of MPP subsets. After gating for LSK, MPP4 were defined as CD48⁺Flt3⁺CD150⁻LSK, MPP3 as CD48⁺Flt3⁻CD150⁻LSK, and MPP2 cells as CD48⁺Flt3⁻CD150⁺LSK. (Bottom) Frequencies of MPP subsets in LSK in the BM.

(C) (Left) Representative FACS plots for the identification of CD41⁺ and CD61⁺ LT-HSC cells and (right) frequency of the same populations in total LT-HSC cells in the BM of mice.


(D–G) Groups of mice were subjected, or not (control, NL), to ligature-induced periodontitis (LIP), and BM cells were harvested on day 7. FACS-sorted LSK were subjected to RNA-sequencing and transcriptome analysis.

(D) Heatmaps of myeloid lineage, osteoclast differentiation.

(E) Heatmaps of erythroid, lymphoid lineage, and oxidative phosphorylation-related genes.

(F) Heatmap of significantly regulated epigenetic regulators by LIP.

(G) Significant upstream transcription factors (left) and upstream regulators (right) predicted by ingenuity pathway analysis (IPA). Data in (A)–(C) are means \pm SD (n = 6 mice/group). *p < 0.05, **p < 0.01, NS, not significant versus control mice; two-tailed Student's t test.



(legend on next page)

Figure S2. Inflammation resolution in the periodontium phenotypically restores myelopoiesis to steady-state levels in the BM, related to Figure 2

Groups of mice were subjected, or not (control), to ligature-induced periodontitis (LIP) for 21 days and subsequently the ligatures were removed (to enable inflammation resolution) for 14 days, at which time BM analysis was performed.

(A) Relative mRNA expression of indicated cytokines in the gingival tissue of mice after 14 days resolution.

(B) Frequencies in total BM cells (top) and absolute cell numbers (bottom) of LSK, LT-HSC, ST-HSC, and MPP.

(C) Frequencies of MPP subsets in LSK in the BM.

(D) Frequencies of CD41⁺ LT-HSC in total LT-HSC cells in the BM.

(E) Frequency within the MyP pool of GMP and CMP (left and middle, respectively) and cell numbers of GMP (right) in the BM.

(F) Frequency of Gr1^{hi}CD11b⁺ granulocytes, Gr1^{int}CD11b⁺ myeloid cells, and CD19⁺ B cells in CD45⁺ cells in the BM.

(G and H) BM LSK were sorted from mice that were subjected to 21dL/14dR or not (NL) and scRNA-seq was performed.

(G) GO enrichment results of top 20 significantly enriched GO terms sorted by PANTHER on the basis of marker genes specific for cluster 7 shown in Figures 2A–2C (Bonferroni-corrected, $p < 0.05$).

(H) Two-dimensional UMAP showing the expression of myeloid marker genes (gray denotes minimal expression, purple intermediate and blue high).

Data are means \pm SD ($n = 6$ mice/group) (A–F). * $p < 0.05$, NS, not significant versus control mice; two-tailed Student's t test.




Figure S3. Single-cell ATACseq analysis, related to Figure 3

- (A) Violin plots showing gene activity scores of *Itga2b* and *Gata2* across cluster 2.
- (B) Violin plots showing gene activity scores of *Csf1r* and *Nlrp3* across cluster 3.
- (C) A cellular trajectory spanning from LSK to GMP.
- (D) Bubble plot of GO enrichment analysis of the cluster 5-specific marker genes.
- (E) Violin plots showing gene activity scores of *Tlr4*, *Camk1d*, *Itga1*, and *Map2k4* across cluster 5.




Figure S4. LIP-subjected mice are more susceptible to CAIA than non-ligated (NL) control mice, related to Figure 4

(A) Groups of mice were subjected, or not (control), to ligature-induced periodontitis (LIP) for 21 days followed by a 14-day resolution. Mice were then subjected to CAIA model and monitored for 14 days.


(B) Clinical arthritis score and hind ankle joint thickness were measured daily.

(legend continued on next page)

(C) CD45.2⁺ mice were either trained (T) by subjecting them to 21-day LIP followed by 14-day resolution ("21dL/14dR"-trained mice), or were left untrained (U; naive controls). BM cells were isolated from trained and untrained mice and transferred to lethally irradiated congenic B6.SJL CD45.1⁺ mice. Twelve weeks after receiving BM transplantation from trained or untrained donor mice, splenic monocytes and neutrophils were isolated.

(D) Isolated splenic monocytes and neutrophils were stimulated with LPS (10 ng/mL) for 24 h. The supernatant was collected for measuring IL-6 and TNF concentration by ELISA.

(E and F) Frequency of donor-derived cells in the BM of recipient mice (E) and frequency of donor-derived (CD45.2⁺) LSK, LT-HSC, ST-HSC, and MPP cells in respective populations in the BM of recipient mice (F) at 16 weeks post-BMT, as described in [Figure 4H](#). Data are means \pm SD (n = 6 recipient mice/group). p* < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, NS, not significant versus untrained mice; two-way repeated measures ANOVA and Sidak's multiple comparisons test (B); two-tailed Student's t test (D–F). T, trained; U, untrained.



(legend on next page)

Figure S5. CAIA activates bone marrow and enhances myelopoiesis, related to Figure 5

(A–E) Mice were subjected, or not (control), to CAIA, and BM analysis was performed after 14 days.

(A) Frequencies (top) of LSK, LT-HSC, ST-HSC, and MPP in total BM cells and absolute cell numbers (bottom) of the same populations.

(B) Frequency of MPP subsets in LSK in the BM.

(C) Frequency of CD41⁺ LT-HSC in total LT-HSC cells in the BM.


(D) Absolute cell numbers of GMP (left) and frequency within the MyP pool of GMP (right).

(E) Absolute cell numbers and frequency of Gr1^{hi}CD11b⁺ granulocytes in CD45⁺ cells in the BM.

(F and G) Mice were subjected, or not (control), to CAIA and monitored daily for ankle joint thickness (F). BM analysis was performed on day 36, i.e., 14 days after complete arthritis resolution. (G) Frequencies of LSK, LT-HSC, ST-HSC, and MPP cells in total BM cells.

(H) CD45.2⁺ mice were either trained (T) by subjecting them to CAIA (disease activity returned to baseline on day 22, and the mice were rested post-resolution for 14 days; total 36 days), or were left untrained (U; naive controls). 36 days post-CAIA, BM cells were isolated and transferred to lethally irradiated congenic B6.SJL CD45.1⁺ mice. Twelve weeks after receiving BM transplantation from donor mice, splenic monocytes and neutrophils from recipient mice were isolated.

(I) Isolated splenic monocytes and neutrophils were stimulated with LPS (10 ng/mL) for 24 h. The supernatant was collected for measuring IL-6 and TNF concentration by ELISA. Data are means \pm SD (n = 6 mice/group). *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, NS, not significant versus control or untrained mice; two-tailed Student's t test. T, trained; U, untrained.



(legend on next page)

Figure S6. Effects of competitive BM transplantation using sorted CD41⁺ and CD41⁻ LT-HSC, related to Figure 5

(A) CD45.1⁺ mice were transplanted with 200 CD41⁺ or CD41⁻ LT-HSC sorted from LIP-trained or untrained CD45.2⁺ mice together with 8×10^5 CD45.1⁺ BM competitor cells.

(B and C) WBC count, donor-derived percentage, and lineage output (% of indicated cell types in donor-derived cells at the indicated time points) in the peripheral blood of mice transplanted with CD41⁺ (B) or CD41⁻ (C) LT-HSC.

(D and E) Frequency of donor-derived cells in the BM of recipient mice and frequency of donor-derived (CD45.2⁺) LSK, LT-HSC, ST-HSC, and MPP cells in respective populations in the BM of mice transplanted with CD41⁺ (D) or CD41⁻ (E) LT-HSC, at 12 weeks post-BMT.

(F–I) 12 weeks post-BMT, the recipient CD45.1⁺ mice were subjected to CAIA. Clinical arthritis score (left) and hind ankle joint thickness (right) in mice transplanted with CD41⁺ (F) or CD41⁻ (G) LT-HSC.

(H and I) Cell suspensions prepared from the synovium of knee joints (harvested on day 7) were stained for live/dead, CD45, CD45.1, CD45.2, CD11b, CD11c, Ly6C, and Ly6G followed by FACS to identify and quantify monocytes (live CD45⁺CD11c⁻CD11b⁺Ly6G⁻Ly6C⁺), neutrophils (live CD45⁺CD11c⁻CD11b⁺Ly6G⁺Ly6C⁻), and CD45⁺ or CD45.2⁺ leukocytes. Total numbers of monocytes, neutrophils, and CD45⁺ cells (top), percentages of CD45.2⁺ cells in monocytes, neutrophils, and CD45⁺ cells (middle), and absolute numbers of CD45.2⁺ monocytes, neutrophils, and total CD45.2⁺ cells (bottom) in mice transplanted with CD41⁺ (H) or CD41⁻ (I) LT-HSC.

(J and K) Splenic monocytes and neutrophils isolated 12 weeks post-BMT from recipient CD45.1⁺ mice not subjected to CAIA were stimulated with LPS (10 ng/mL) for 24 h. The supernatant was collected for measuring IL-6 and TNF concentration by ELISA in mice transplanted with CD41⁺ (J) or CD41⁻ (K) LT-HSC. Data are means \pm SD (B, C, F–I, n = 6 mice/group; D, E, J, and K, n = 4 mice/group). *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, NS, not significant versus untrained mice. Two-tailed Student's t test (D, E, H–K) except for (I), 2nd row, 1st left (two-tailed Mann-Whitney U test); two-way repeated measures ANOVA and Sidak's multiple comparisons test (B, C, F, and G). T, trained; U, untrained.




Figure S7. Role of IL-1 signaling in LIP and LIP-induced BM myelopoiesis, related to Figure 7

(A) *Il1r1*^{HSPC-KO} mice and WT littermate controls were subjected to tamoxifen treatment to induce *Il1r1* deletion in HSPC. Mice were then subjected to ligature-induced periodontitis (LIP) for 21 days. BM cells were harvested, and the frequencies of LSK, ST-HSC, and MPP in total BM cells, CD41⁺ cells in total LT-HSC, MPP3 in LSK, GMP in MyP pool, as well as Gr1^{hi}CD11b⁺ granulocytes and Gr1^{int}CD11b⁺ myeloid cells in CD45⁺ cells, were analyzed by FACS.

(B) CD45.2 *Il1r1*^{HSPC-KO} and littermate controls with intact HSPC IL1-R1 expression were trained, by subjecting them to 21-day LIP and 14-day resolution, and then used as donors for BMT to groups of lethally irradiated congenic B6.SJL (CD45.1⁺) mice. Frequency of donor-derived LSK, LT-HSC, ST-HSC, and MPP cells in CD45.2⁺ BM cells of recipient mice at 12 weeks post-BMT.

(C) Mice with global IL1-R1 deletion (*Il1r1*^{KO}) mice and WT littermate controls were subjected to 5-day ligature-induced periodontitis and bone loss was measured. Data are means ± SD (A, n = 5 mice/group; B and C, n = 6 mice/group). *p < 0.05, **p < 0.01, ***p < 0.001, NS, not significant versus WT littermate control mice; two-tailed Student's t test.