



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Purinergic Signaling in Kidney Disease

**Citation for published version:**

Menzies, R, Tam, FWK, Unwin, RJ & Bailey, M 2016, 'Purinergic Signaling in Kidney Disease', *Kidney International*. <https://doi.org/10.1016/j.kint.2016.08.029>

**Digital Object Identifier (DOI):**

[10.1016/j.kint.2016.08.029](https://doi.org/10.1016/j.kint.2016.08.029)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Kidney International

**Publisher Rights Statement:**

This is the authors accepted manuscript as accepted by Elsevier on 15 August 2016

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## Purinergic Signaling in Kidney Disease

*Robert I. Menzies<sup>1</sup>, Frederick W Tam<sup>2</sup>, Robert J Unwin<sup>3,4</sup> and Matthew A. Bailey<sup>1</sup>*

<sup>1</sup>British Heart Foundation Centre for Cardiovascular Science, The University of Edinburgh, <sup>2</sup>Imperial College Renal and Transplant Centre, Department of Medicine, Imperial College London, <sup>3</sup>Cardiovascular and Metabolic Diseases (CVMD iMed) Biotech Unit, AstraZeneca Gothenburg, Sweden; <sup>4</sup>UCL Centre for Nephrology, University College London, UK.

Running title: Renal purinoceptors

Keywords: ATP, adenosine, P2X, P2Y, adenosine, kidney, renal tubule, vasculature, inflammation

Correspondence:

Robert Unwin  
UCL Centre for Nephrology,  
UCL Medical School  
Royal Free Campus  
Rowland Hill Street,  
London NW3 2PF,  
United Kingdom  
Email: robert.unwin@ucl.ac.uk

## 1 **Abstract**

2 Nucleotides are key subunits for nucleic acids and provide energy for intracellular metabolism.  
3 They can also be released from cells to act physiologically as extracellular messengers or  
4 pathologically as danger signals. Extracellular nucleotides stimulate membrane receptors in the  
5 P2 and P1 family. P2X are ATP-activated cation channels; P2Y and P1 are G-protein coupled  
6 receptors activated by ATP, ADP, UTP and UDP or adenosine, respectively. Renal P2 receptors  
7 influence both vascular contractility and tubular function. Renal cells also express  
8 ectonucleotidases that rapidly hydrolyze extracellular nucleotides. These enzymes integrate this  
9 multi-receptor purinergic-signaling complex by determining the nucleotide milieu, as well as  
10 titrating receptor activation.

11 Purinergic signaling also regulates immune cell function by modulating the synthesis and release  
12 of various cytokines such as IL1- $\beta$  and IL-18 as part of inflammasome activation. Abnormal or  
13 excessive stimulation of this intricate paracrine system can be pro- or anti-inflammatory, and is  
14 also linked to necrosis and apoptosis. Kidney tissue injury causes a localized increase in ATP  
15 concentration, and sustained activation of P2 receptors can lead to renal glomerular, tubular  
16 and vascular cell damage. Purinergic receptors also regulate the activity and proliferation of  
17 fibroblasts, promoting both inflammation and fibrosis in chronic disease.

18 In this short review we summarize some of the recent findings related to purinergic signaling in  
19 the kidney. We focus predominantly on the P2X7 receptor, discussing why antagonists have so  
20 far disappointed in clinical trials and how advances in our understanding of purinergic signaling  
21 might help to reposition these compounds as potential treatments for renal disease.

## 22 Introduction

23 Since their discovery in the 1970s, P2 purinergic receptors (P2R) have evolved from an initially  
24 contentious biological concept <sup>1</sup>, through to a progressive understanding of their complex  
25 physiological actions, emerging now as attractive and 'druggable' targets for disease <sup>2,3</sup>. To date,  
26 the most advanced potential therapeutic P2R targets are antagonists for P2Y<sub>12</sub>R to inhibit  
27 thrombosis <sup>4</sup>, and P2X<sub>7</sub>R for the treatment of chronic inflammatory diseases such as  
28 rheumatoid arthritis <sup>5</sup> and COPD <sup>6</sup>. Several P2X<sub>7</sub>R antagonists have completed Phase 2 clinical  
29 trials, but despite pre-clinical promise, these compounds have failed to deliver the expected  
30 benefit and so interest in P2X<sub>7</sub>R has declined. In this concise review we cover purinergic  
31 signaling in the kidney and explore the contribution of this system to renal physiology and  
32 disease. The main focus is on the role of P2X receptors, particularly P2X<sub>7</sub>R, in renal injury and  
33 disease. P2X<sub>7</sub>R can orchestrate interactions between the immune and vascular systems, and  
34 defining this complex interaction as inflammation and injury develop may help us unlock the  
35 potential of P2X<sub>7</sub>R antagonists as renal therapeutics.

36

## 37 P2 receptors and purinergic signaling in the kidney

38 Purinergic receptors are sub-classified as P1R that bind adenosine and P2R that are activated by  
39 purine/pyrimidine nucleotides; P2R are in turn subdivided into P2YR and P2XR. The 8 P2YRs are  
40 coupled to G-proteins and are activated with differing selectivity by adenosine triphosphate  
41 (ATP), adenosine diphosphate (ADP), uridine triphosphate (UTP) and uridine diphosphate (UDP).  
42 The 7 P2XRs are trimeric ligand-gated ion channels activated by ATP, but not, or only weakly, by  
43 ADP or adenosine monophosphate (AMP). The molecular properties of these receptors and  
44 their ligands are described in detail in the *IUPHAR/BPS Guide to Pharmacology*:  
45 <http://www.guidetopharmacology.org>.

46 P2 receptors are expressed in all segments of the nephron and renal cells often express multiple  
47 receptor subtypes at both the apical and basolateral cell membranes <sup>7,8</sup>. Renal cells can also  
48 release ATP and UTP into the extracellular space. This release is likely to be regulated and is  
49 facilitated by several transport systems that involve vesicular or lysosomal exocytosis, or  
50 channel-mediated release via connexins <sup>9</sup> or pannexins <sup>10</sup>. Extracellular ATP and UTP have short  
51 half-lives due to rapid catabolism by ectonucleotidases (**Figure 1**) that are also expressed by  
52 renal cells <sup>11,12</sup>. Their immediate breakdown products, ADP and UDP, are potent agonists at  
53 P2Y1R,12R,13R, and P2Y6R,14R, respectively. Further metabolism of ADP produces the 5'AMP  
54 (through CD39) and eventually adenosine (through CD73), the agonist at P1R (A1,2A,2B,3) that  
55 are also present in renal epithelia. Thus, the kidney has complex and regulated machinery for  
56 hierarchical purinergic signaling integrated by the action of ectonucleotidases. Ascribing specific  
57 physiological functions to a given receptor subtype has been challenging: available receptor  
58 agonists are not sufficiently selective and are often unstable <sup>11</sup>. In contrast, selective and  
59 specific receptor antagonists are providing a pharmacological means of assessing the function(s)  
60 of this system *in vivo*.

61 Extracellular nucleotides can influence a range of physiological functions, from cell-proliferation  
62 and growth, through to energy metabolism and transepithelial solute flux. These functions have  
63 been reviewed in depth recently <sup>13</sup> and we can provide only a brief overview. It is evident that  
64 abnormal P2R activity can occur in various inflammatory and non-inflammatory disease states  
65 ranging from hypertension <sup>14</sup> to transplant rejection, to polycystic kidney disease <sup>15</sup>. However,  
66 more beguiling is the therapeutic potential for P2XR antagonists in chronic kidney disease (CKD).

67

68

69

## 70 **P2 receptors control renal vascular and microvascular function**

71 P2 receptors are expressed throughout the vasculature and microvasculature (**Figure 2**) and  
72 strongly influence vessel function <sup>16</sup>. The renal vasculature and microvasculature also expresses  
73 NTPDase1 (CD39) that hydrolyses ATP to ADP and AMP, and thereby rapidly curtail purinergic  
74 signaling <sup>17</sup>. P2X1R is the dominant receptor in vascular smooth muscle and application of ATP  
75 to the adventitia evokes contraction in the pre-glomerular vasculature <sup>18,19</sup>. P2X1R null mice  
76 display an attenuated pressure-induced constriction of the afferent arteriole <sup>20</sup> and targeted  
77 deletion of NTPDase1 prolongs the half-life of extracellular ATP, enhancing the vascular  
78 response to increased pressure <sup>21</sup>.

79 Direct renal artery infusion of ATP increases blood flow, causing vasodilation due to production  
80 of nitric oxide (NO) by the endothelium <sup>22</sup> and also NO-independent vasodilatation induced by  
81 intra-renal prostanoids <sup>23</sup>. The P2 receptor subtype(s) that mediates the vasodilatory response  
82 to ATP is unknown. In human arterial endothelial cells and endothelial cells cultured from the  
83 mouse pulmonary artery, P2X4R is the most abundantly expressed receptor, followed by P2X7R  
84 <sup>24-26</sup>. P2X4R mediates the release of NO in response to increased shear stress <sup>24</sup>. This response is  
85 lost in P2X4R null mice, which have endothelial dysfunction and hypertension <sup>25</sup>. P2X7R  
86 activation seems to promote a tonic vasoconstriction of both the pre-glomerular arteries and  
87 medullary microcirculation <sup>14</sup>, which is discussed more below. Other P2 receptors can influence  
88 endothelial function, for example, vasodilatation caused by UDP is abolished in P2Y6R null mice  
89 <sup>27</sup>. The descending *vasa recta* are also affected by extracellular nucleotides, since infusion of  
90 ATP into the renal artery reduces medullary blood flow as a result of P2X1R activation <sup>23</sup>, and  
91 ATP released from sympathetic nerves causes constriction of *vasa recta* pericytes <sup>28</sup>.

92

93 Multiple P2R subtypes are expressed in glomerular cells (**Figure 2**). Under normal conditions,  
94 P2YR predominate <sup>29</sup> and extracellular nucleotides influence mesangial proliferation and  
95 contraction, as well as contraction of the parietal sheet <sup>29</sup>. In podocytes, P2Y1R is the dominant  
96 functional receptor as demonstrated by comprehensive pharmacological profiling and  
97 immunolocalization <sup>30</sup>; however, recently P2X4R has been shown to have a mechano-sensitive  
98 role affecting the podocyte actin cytoskeleton <sup>31</sup>, although P2X4R knockout mice, while  
99 hypertensive, have no obvious gross glomerular phenotype and are not known to be proteinuric.  
100 In contrast, P2Y1R null mice are protected from acute nephrotoxic injury, showing preserved  
101 renal function, reduced capillary rarefaction and fibrosis, and enhanced survival <sup>32</sup>. P2Y1R  
102 activation may, therefore, contribute to glomerular injury. P2X7R expression also seems to be  
103 associated with glomerular injury, since it is increased in multiple glomerular cells types,  
104 including inflammatory cells, in models of severe hypertension, type 1 diabetes <sup>33</sup>, and acute  
105 inflammatory glomerulonephritis <sup>34</sup>. Uncovering the primary role of this increased glomerular  
106 P2X7R expression remains an active area of research.

107

## 108 **P2 receptors and renal tubular physiology**

109 P2R exert a largely inhibitory effect on tubular electrolyte transport and this, together with  
110 expression in specific nephron segments, has been reviewed extensively elsewhere <sup>35</sup> and is  
111 summarized in **Figure 2**. The processes are best defined for sodium flux, which is tonically  
112 suppressed by P2R activation in several nephron segments <sup>36</sup>. It is likely that such paracrine  
113 control by extracellular nucleotides provides a route for rapid modulation of tubular transport  
114 that can link solute and fluid delivery to adaptive transport capacity, for example adenosine-  
115 mediated tubuloglomerular feedback is impaired in CD73<sup>-/-</sup> mice <sup>37</sup>. This form of control can  
116 integrate with more slowly adapting hormonal systems, for example the renin-angiotensin-

117 aldosterone system (RAAS) to regulate the phenomenon of aldosterone escape <sup>38</sup>. Indeed, ATP  
118 release by tubular cells, stimulated by increased flow, contributes to the control of extracellular  
119 fluid volume by the kidney, and blood pressure regulation, as discussed below.

120

### 121 **Proximal tubule**

122 The proximal tubule, which expresses apical P2Y1R and P2X5R, and basolateral P2Y4R and  
123 P2Y6R <sup>39, 40</sup>, accounts for reabsorption of ~65% of the filtered sodium load. Extracellular  
124 nucleotides inhibit the major sodium transporters in this segment, NHE3 <sup>41</sup>, NaPi2 <sup>42</sup> and Na,K-  
125 ATPase <sup>43</sup>, and inhibition of transepithelial flux has been confirmed *in vivo* <sup>44</sup>. The ATP  
126 concentration in tubular fluid is unknown, although measurements in bulk fluid collected from  
127 the end of the proximal convoluted tubule (PCT) report concentrations of 100-300nmol/l <sup>45</sup>. The  
128 brush border membrane expresses ENPP3 (ectonucleotide pyrophosphatase/ phosphodiesterase  
129 3) and ecto-5'-nucleotidase (NT5E; CD73) <sup>12</sup> that should terminate physiological signaling.  
130 Microperfusion studies using nucleotide scavengers suggest that the 'ambient' concentration of  
131 the physiological purinergic ligand, most probably ADP, is ~10μmol/l, exerting a tonic inhibitory  
132 effect that may help to balance tubular sodium reabsorption with glomerular filtration <sup>44</sup>.

133

### 134 **The distal nephron**

135 Increased fluid flow or changes in osmolality of the tubular fluid promotes nucleotide secretion  
136 in both the thick limb of Henle <sup>46</sup> and collecting duct <sup>47</sup>, inhibiting transport in downstream  
137 nephron segments. In the thick ascending limb of Henle (TALH), ATP release is dependent on  
138 activation of the transient receptor potential cation channel TRPV4 osmosensor <sup>48</sup>. These  
139 nucleotides activate endothelial NO synthase (NOS3) in thick limb cells, and P2R signaling  
140 underpins the flow-dependent increase in NO production <sup>49</sup> and subsequent inhibition of apical



141 NKCC2 and basolateral Na,K-ATPase activity<sup>50</sup>. Studies in knockout mice suggest P2X4R and  
142 P2Y2R contribute to this signaling arc<sup>51,52</sup>.

143 Extracellular ATP has long been known to inhibit the epithelial sodium channel (ENaC), the rate-  
144 limiting step for sodium transport in the connecting tubule and collecting duct<sup>53</sup>. Studies in  
145 isolated segments show that ATP activates P2Y2R to reduce the open probability of ENaC<sup>54-56</sup>.  
146 P2yr2 null mice lack the tonic suppression of ENaC and are hypertensive<sup>54</sup>. Studies *in vivo*  
147 suggest that P2X4R activation also inhibits ENaC<sup>53,57</sup> and our own pilot studies in a P2X4R null  
148 mouse suggest that this receptor may be important in the modulation of sodium transport by  
149 aldosterone (Craigie et al, unpublished).

150

#### 151 **P2R and blood pressure regulation**

152 Hypertension is a major modifiable risk factor for cardiovascular and renal disease and is highly  
153 prevalent<sup>58</sup>. Human genetic studies have found an association between SNPs in P2XR encoding  
154 genes and blood pressure or cardiovascular disease. The loss of function variant rs28360472 in  
155 P2RX4 associates with increased pulse pressure<sup>59</sup>, itself an important cardiovascular risk factor.  
156 An intronic SNP (rs591874) in the gene encoding P2X7R is associated with elevated blood  
157 pressure<sup>60</sup>. The loss of function variant rs3751143 is common (25% heterozygosity and up to 3%  
158 homozygosity) and protects against ischemic stroke<sup>61</sup>. The physiology of P2RX7 genetic  
159 variation is almost certainly subtle, if not complex. For example, rs3751143 does not associate  
160 with impaired endothelial dysfunction or vascular stiffness in essential hypertensives<sup>62</sup>, but  
161 does confer a significantly reduced sensitivity to P2X7R antagonism<sup>63</sup>.

162 Pressure-natriuresis is an important mechanism of long-term blood pressure control<sup>64</sup> and is  
163 modulated by paracrine factors that inhibit sodium transport in the renal proximal tubule,  
164 including extracellular nucleotides. Microdialysis experiments reveal a direct relationship  
165 between renal artery perfusion pressure and the concentration of ATP in the interstitial fluid of

166 the kidney cortex <sup>65</sup>. As mentioned earlier, extracellular nucleotides inhibit the key transporters  
167 in the proximal tubule <sup>41-43</sup>. This natriuretic effect is buttressed by inhibition of sodium transport  
168 in the distal nephron. Increased flow through the collecting duct promotes ATP secretion to  
169 inhibit ENaC. This ATP release is abolished in connexin 30 knockout mice, severely attenuating  
170 the pressure-natriuresis response <sup>9</sup>. Consistent with this, mice over-expressing human  
171 NTPDase1 (CD39), a cell surface enzyme that scavenges extracellular nucleotides, display a small  
172 impairment of the natriuretic response to a high sodium diet and concomitant aldosterone  
173 infusion <sup>66</sup>. It is assumed that P2Y2R mediates the inhibitory effect of ATP on distal tubule  
174 sodium transport. Receptor agonists have been considered as potential antihypertensives.  
175 P2yr2 null mice display enhanced ENaC activity and are hypertensive. Surprisingly, blood  
176 pressure is salt resistant <sup>67</sup> and endothelial dysfunction with impaired NO release may be causal.  
177 Recent studies also suggest that ATP can inhibit ENaC indirectly: in IMCD cells, activation of  
178 P2X7R promotes synthesis of endothelin-1, which is pro-natriuretic due to ETB-mediated  
179 inhibition of ENaC <sup>68</sup>. However, the significance of this cell line-based study is not clear, since  
180 acute P2X7R antagonism *in vivo* improves the pressure-natriuresis relationship <sup>14</sup>.

181 Although P2X7R activation contributes to the physiological control of blood pressure by the  
182 kidney, sustained activation of the receptor, which does not de-sensitize with repeated  
183 exposure to ATP, promotes hypertensive renal injury. Thus, prophylactic P2X7R antagonism <sup>69</sup> or  
184 'knock-out' of the murine P2X7k transcript <sup>70</sup>, which leaves several functional P2RX7 transcripts  
185 intact <sup>71</sup>, protects against the injury associated with salt-sensitive hypertension. P2X7R  
186 antagonism/deletion reduced albuminuria and interstitial fibrosis, lowered blood pressure and  
187 reduced the infiltration of T and B cells, macrophages and leucocytes. The mechanisms  
188 underpinning these effects are not known, as discussed further below. Our data suggest that  
189 P2X7R in the renal vasculature and microvasculature may impair blood pressure regulation by  
190 the kidney <sup>14</sup>. We identified elevated renal expression of P2X7R (and P2X4R) as a candidate gene

191 for hypertensive renal vascular injury in rats <sup>72</sup>. P2X7R localized to the vascular and  
192 microvascular endothelium down to afferent arterioles. The selective P2X7R antagonist  
193 AZ11657312 increased renal medullary perfusion and improved tissue oxygenation in  
194 angiotensin II-treated rats <sup>14</sup>; these beneficial effects were partially dependent on NO synthesis.  
195 Overall, activation of P2X7R induces microvascular dysfunction and regional hypoxia,  
196 particularly under high angiotensin II tone. These effects are pro-inflammatory and may  
197 contribute to progression of renal injury. In the next section, we discuss the role of P2X7R in  
198 renal injury and disease and assess the potential for antagonists as renal therapeutics.

199

## 200 **P2XR and renal injury**

201 There is consistent pre-clinical evidence supporting a role for P2X7R in inflammation (**Figure 3**),  
202 and, as already mentioned, P2X7R antagonists have been explored as a treatment target in  
203 rheumatoid arthritis <sup>5</sup>, COPD <sup>6</sup>, and IBD <sup>73</sup>, but with mixed or generally disappointing results. This  
204 has caused interest in the receptor to wax and wane. However, it is likely that an improved  
205 understanding of the biological roles of P2X7R, including its unique two-stage ability to induce  
206 membrane permeability to large (>900 Da) molecules, rather than cations alone, as well as the  
207 regulation and function of the main splice variants, will provide a fresh impetus to the clinical  
208 testing of antagonists.

209 In the normal kidney P2X7R is typically only present at low levels, often undetectable by RNA  
210 analysis in whole kidney extracts. The receptor is normally localized to certain compartments,  
211 particularly the vasculature and microvasculature, at least in the rat <sup>7, 14, 72</sup>. A wealth of data  
212 shows that injury/inflammation increases expression in renal cells. For example, TNF $\alpha$  can  
213 induce expression of P2X7R in cultured mesangial cells <sup>74</sup>. In renal biopsy material from patients  
214 with lupus nephritis, increased expression of P2X7R protein has been found <sup>75</sup>. Nevertheless, it

215 remains to be investigated whether the extent of P2X7R expression correlates with the severity  
216 of clinical disease and a more detailed study with larger patient numbers is needed.

217

### 218 *Glomerulonephritis*

219 A more detailed characterization of the expression and potential function of P2X7R have been  
220 carried out in rodent models of nephrotoxic nephritis (NTN)<sup>75</sup>. In a mouse model of accelerated  
221 NTN, increased expression of P2X7R was co-localized to glomerular macrophages as well as  
222 intrinsic glomerular cells. In NTN in WKY rats, onset P2X7R expression coincided with onset of  
223 proteinuria. The inflamed glomeruli are infiltrated by macrophages showing the NLRP3  
224 inflammasome activation<sup>76</sup>. The WKY strain of rat is known to be more susceptible to  
225 developing severe and progressive glomerulonephritis when compared with the resistant LEW  
226 rat strain. WKY and LEW rats have identical MHC genes, but have distinct genetic differences  
227 and differences in their expression of P2X7R and the NLRP3 inflammasome<sup>76</sup>. More specifically,  
228 bone marrow derived (BMD) macrophages from WKY rats have increased expression of P2X7R  
229 protein and mRNA associated with increased expression of multiple genes of the NLRP3  
230 inflammasome pathway, even in their basal state *in vitro*, again when compared with BMD  
231 macrophages from LEW rats. Following priming with endotoxin and stimulation with  
232 extracellular ATP, compared with LEW rats, macrophages from WKY rats have higher levels of  
233 caspase-1 activation and secretion of more mature IL-1 $\beta$  and IL-18. Thus, strain differences in  
234 expression of P2X7R and subsequent downstream activation of the inflammasome may be  
235 responsible for the difference in susceptibility to experimental glomerulonephritis.

236 The functional importance of P2X7R was investigated in gene knockout mice and with systemic  
237 treatment by a small molecule P2X7R antagonist<sup>34</sup>. Using the model of accelerated NTN, the  
238 P2X7R knockout mice had lower urinary monocyte chemoattract-1 (CCL2), fewer infiltrating

239 glomerular macrophages, less glomerular fibrin deposition and less proteinuria than in wild-type  
240 mice. In NTN rats, treatment with the P2X7R antagonist A438079 significantly reduced  
241 glomerular expression of CCL2, glomerular macrophage infiltration, glomerular fibrinoid  
242 necrosis and proteinuria compared with vehicle-treated rats. However, exactly how P2X7R is  
243 involved in antibody-mediated glomerulonephritis is unclear. Typically, extracellular ATP binds  
244 to P2X7R in endotoxin-primed macrophages, resulting in inflammasome activation and release  
245 of mature IL-1 $\beta$  and IL-18<sup>77</sup>, yet endotoxin or other bacterial products are not involved in the  
246 induction of NTN in WKY rats<sup>34</sup>. The interaction between immune complex stimulation and  
247 P2X7R needs further investigation and to ascertain whether treatment with the P2X7R  
248 antagonist after the onset of disease is effective in reducing the severity of glomerulonephritis.  
249 There is also recent evidence in lupus prone mice that treatment with a P2X7R antagonist can  
250 decrease the severity of renal injury and levels of dsDNA antibodies<sup>78</sup>.

251

### 252 *Acute kidney injury*

253 Renal ischemia-reperfusion injury (IRI) is a common occurrence in many clinical settings from  
254 sepsis to major surgery, including renal transplantation. There is increased expression of P2X7R,  
255 mainly in the renal tubules, in a mouse model of renal IRI; treatment with A438079 reduced  
256 renal expression of chemokines (MCP-1 and RANTES), p-ERK, NGAL, renal tubular injury and cell  
257 death<sup>79</sup>.

258 As well as the mentioned increase in P2X7R in a rat model of type 1 diabetes<sup>33</sup>, in a mouse  
259 model of high fat diet-induced metabolic disease, proteinuria and albuminuria developed in the  
260 wild-type mice, but not in P2X7a variant knockout mice<sup>80</sup>. In the high fat diet fed mice there  
261 was also increased renal expression of P2X7R and components of the NLRP3 inflammasome that  
262 were attenuated in the high fat diet fed P2X7R knockout mice, as was renal expression of

263 chemokine CCL2, macrophage infiltration and expression of extracellular matrix protein.  
264 Moreover, increased expression of P2X7R and inflammasome components were found in renal  
265 tissue from patients with glomerulonephritis <sup>75</sup>.

266

### 267 *Fibrosis*

268 Purinergic signaling is involved in tissue remodeling (**Figure 3**) and several studies in various  
269 tissues suggest that these pathways may also drive tissue fibrosis in chronic injury, one feature  
270 of which is a sustained increase in ambient concentrations of ATP, ADP, UTP and UDP <sup>81</sup>. Tissue  
271 fibroblasts express multiple P2R subtypes and respond to extracellular nucleotides by activating  
272 key pathways for the production of extracellular matrix. In cardiac fibroblasts, for example,  
273 P2Y2R activation is strongly pro-fibrotic <sup>82</sup>, and activation of P2X4R and P2X7R promotes  
274 ERK1/2-dependent fibroblast proliferation <sup>83</sup>. This cluster of P2Rs is also relevant to the kidney  
275 in which fibroblasts and mesangial cells mainly determine ECM deposition. In this context,  
276 P2Y2R activation increases mesangial cell proliferation <sup>74</sup> and P2X7R activation increases matrix  
277 production by mesangial cells <sup>84</sup>.

278 The role of P2 receptors in renal fibrosis has been investigated in the unilateral ureteral  
279 obstruction (UUO) model <sup>85</sup>. Transient expression of P2X7R was detected only in tubular  
280 epithelial cells 7 days after induction of UUO in wild-type mice. The renal tubular expression of  
281 TGF- $\beta$ 1, macrophage infiltration, tubular apoptosis and tubulointerstitial fibrosis were reduced  
282 in P2X7R knockout mice compared with wild-type mice by day 14. The role of the  
283 inflammasome in this model has also been investigated. Knockout of apoptosis-associated  
284 speck-like protein containing a caspase recruitment domain (ASC) in mice results in reduced  
285 UUO-mediated tubulointerstitial fibrosis, together with fewer infiltrating inflammatory cells and  
286 reduced renal expression of mRNA for IL-1 $\beta$ , CCL2, TGF $\beta$ 1 and collagen I; however, it is not clear

287 how P2X7R may regulate TGF- $\beta$ 1 expression <sup>86</sup>. While there is a well-established relationship  
288 between stimulation of P2X7R and activation of the inflammasome, it is not known what the  
289 priming signal is in the sterile UUO model and what triggers fibrogenesis.

290 P2X4R is closely related to P2X7R and there has been ongoing controversy over whether P2X4R  
291 and P2X7R can form heterotrimers <sup>87, 88</sup>. The potential importance of P2X4R in renal fibrosis has  
292 been investigated in the UUO model. Surprisingly, the P2X4R knockout mice showed increased  
293 renal fibrosis following induction of UUO associated with increased expression of TGF $\beta$ 1 and  
294 connective tissue growth factor (CTGF, also known as CCN2), and increased amounts of type I  
295 collagen <sup>89</sup>. These results suggest that P2X7R is pro-fibrotic in this model and that P2X4R may  
296 have an anti-fibrotic role through its regulation of pro-fibrotic growth factors.

297 More recent studies show that nucleotidases may also contribute to fibrosis by regulating the  
298 half-life of ATP. ENTPD1 (CD39)-null mice are more sensitive to ischemic tissue injury than wild-  
299 type mice <sup>90</sup>, because ATP persists and its hydrolysis to protective adenosine is blunted.  
300 Similarly, these null mice have more pronounced renal injury in the IRI model <sup>91, 92</sup>; although in  
301 this setting the role of adenosine is less certain, since the deletion of CD73, the enzyme that  
302 converts AMP to adenosine, was also protective <sup>93</sup>. Overall, these data suggest that enzymes  
303 involved in terminating P2R signaling may be less tractable as therapeutic targets than the  
304 receptors themselves. Recent studies indicate that CD39 expression by T-reg lymphocytes is  
305 essential for their pro-reparative role in response to chronic renal injury <sup>94</sup>.

306

### 307 **What now for P2X7R antagonists?**

308 P2X7R antagonists may have failed because of significant gaps in our knowledge about the  
309 complex processing and diverse roles of *P2XR7* gene products and the implications this may  
310 have for P2X7R in disease. Single nucleotide polymorphisms (SNPs) such as rs3751143 (causing

311 Glu496Ala) can impair P2X7R function <sup>95,96</sup>: ATP-dependent IL-1 $\beta$  release from lymphocytes is  
312 significantly suppressed in individuals carrying this SNP <sup>97</sup>. Alternative splicing can produce novel  
313 protein isoforms that are emerging as important factors in disease pathogenesis, as well as in  
314 determining the right treatment target <sup>98</sup>.

315 Human P2X7R has at least 10 splice isoforms, the functions of which have not been unraveled;  
316 however, in rodents, the common 'k variant' of P2X7R is much more sensitive to ATP than the  
317 original full-length 'a variant' <sup>99</sup>. Pre-clinical data suggest that genetic variation in P2X7R will  
318 increase the population wide variance of both agonist and antagonist binding affinities,  
319 suggesting that we need to re-evaluate or redefine clinical trials on the basis of the P2X7R  
320 "fingerprint". The tissue distribution, regulation and function of these splice isoforms in the  
321 healthy kidney is just beginning to be explored; the pharmacogenomics of P2X7R and the impact  
322 of disease is largely unknown. The next phase of research will define these key biological  
323 processes involving P2X7R, which may not all be 'bad' <sup>100</sup>, and provide a better understanding of  
324 how isoform-specific receptor antagonists should be deployed in kidney disease. Is this *P2X7R*  
325 *Redux*?

326



327 **References:**

- 328 1. Burnstock G. Purinergic nerves. *Pharmacol Rev* 1972; **24**: 509-581.  
329
- 330 2. Burnstock G. Introduction: P2 receptors. *Curr Top Med Chem* 2004; **4**: 793-803.  
331
- 332 3. Erlinge D, Burnstock G. P2 receptors in cardiovascular regulation and disease. *Purinergic*  
333 *Signal* 2008; **4**: 1-20.  
334
- 335 4. van Giezen JJ, Humphries RG. Preclinical and clinical studies with selective reversible  
336 direct P2Y12 antagonists. *Semin Thromb Hemost* 2005; **31**: 195-204.  
337
- 338 5. Keystone EC, Wang MM, Layton M, *et al.* Clinical evaluation of the efficacy of the P2X7  
339 purinergic receptor antagonist AZD9056 on the signs and symptoms of rheumatoid  
340 arthritis in patients with active disease despite treatment with methotrexate or  
341 sulphasalazine. *Ann Rheum Dis* 2012; **71**: 1630-1635.  
342
- 343 6. Eltom S, Stevenson CS, Rastrick J, *et al.* P2X7 receptor and caspase 1 activation are  
344 central to airway inflammation observed after exposure to tobacco smoke. *PLoS One*  
345 2011; **6**: e24097.  
346
- 347 7. Menzies RI, Unwin RJ, Bailey MA. Renal P2 receptors and hypertension. *Acta Physiol*  
348 *(Oxf)* 2015; **213**: 232-241.  
349
- 350 8. Bailey MA, Hillman KA, Unwin RJ. P2 receptors in the kidney. *J Auton Nerv Syst* 2000; **81**:  
351 264-270.  
352
- 353 9. Sipos A, Vargas SL, Toma I, *et al.* Connexin 30 deficiency impairs renal tubular ATP  
354 release and pressure natriuresis. *J Am Soc Nephrol* 2009; **20**: 1724-1732.  
355
- 356 10. Jackson MF. Interdependence of ATP signalling and pannexin channels; the servant was  
357 really the master all along? *Biochem J* 2015; **472**: e27-30.  
358
- 359 11. Shirley DG, Vekaria RM, Sevigny J. Ectonucleotidases in the kidney. *Purinergic Signal*  
360 2009; **5**: 501-511.  
361
- 362 12. Vekaria RM, Shirley DG, Sevigny J, *et al.* Immunolocalization of ectonucleotidases along  
363 the rat nephron. *Am J Physiol Renal Physiol* 2006; **290**: F550-560.  
364
- 365 13. Burnstock G. Purinergic signalling: from discovery to current developments. *Exp Physiol*  
366 2014; **99**: 16-34.  
367
- 368 14. Menzies RI, Howarth AR, Unwin RJ, *et al.* Inhibition of the purinergic P2X7 receptor  
369 improves renal perfusion in angiotensin-II-infused rats. *Kidney Int* 2015; **88**: 1079-1087.  
370
- 371 15. Hillman KA, Woolf AS, Johnson TM, *et al.* The P2X7 ATP receptor modulates renal cyst  
372 development in vitro. *Biochem Biophys Res Commun* 2004; **322**: 434-439.  
373

- 374 16. Inscho EW. ATP, P2 receptors and the renal microcirculation. *Purinergic Signal* 2009; **5**:  
375 447-460.  
376
- 377 17. Zhang Y, Morris KL, Sparrow SK, *et al.* Defective renal water handling in transgenic mice  
378 over-expressing human CD39/NTPDase1. *Am J Physiol Renal Physiol* 2012; **303**: F420-430.  
379
- 380 18. Chan CM, Unwin RJ, Bardini M, *et al.* Localization of P2X1 purinoceptors by  
381 autoradiography and immunohistochemistry in rat kidneys. *Am J Physiol* 1998; **274**:  
382 F799-804.  
383
- 384 19. Inscho EW, Cook AK, Imig JD, *et al.* Physiological role for P2X1 receptors in renal  
385 microvascular autoregulatory behavior. *J Clin Invest* 2003; **112**: 1895-1905.  
386
- 387 20. Inscho EW, Cook AK, Imig JD, *et al.* Renal autoregulation in P2X1 knockout mice. *Acta*  
388 *Physiol Scand* 2004; **181**: 445-453.  
389
- 390 21. Kauffenstein G, Drouin A, Thorin-Trescases N, *et al.* NTPDase1 (CD39) controls  
391 nucleotide-dependent vasoconstriction in mouse. *Cardiovasc Res* 2010; **85**: 204-213.  
392
- 393 22. Eltze M, Ullrich B. Characterization of vascular P2 purinoceptors in the rat isolated  
394 perfused kidney. *Eur J Pharmacol* 1996; **306**: 139-152.  
395
- 396 23. Eppel GA, Ventura S, Evans RG. Regional vascular responses to ATP and ATP analogues in  
397 the rabbit kidney in vivo: roles for adenosine receptors and prostanoids. *Br J Pharmacol*  
398 2006; **149**: 523-531.  
399
- 400 24. Yamamoto K, Korenaga R, Kamiya A, *et al.* P2X(4) receptors mediate ATP-induced  
401 calcium influx in human vascular endothelial cells. *Am J Physiol Heart Circ Physiol* 2000;  
402 **279**: H285-292.  
403
- 404 25. Yamamoto K, Sokabe T, Matsumoto T, *et al.* Impaired flow-dependent control of vascular  
405 tone and remodeling in P2X<sub>4</sub>-deficient mice. *Nat Med* 2006; **12**: 133-137.  
406
- 407 26. Ray FR, Huang W, Slater M, *et al.* Purinergic receptor distribution in endothelial cells in  
408 blood vessels: a basis for selection of coronary artery grafts. *Atherosclerosis* 2002; **162**:  
409 55-61.  
410
- 411 27. Bar I, Guns PJ, Metallo J, *et al.* Knockout mice reveal a role for P2Y6 receptor in  
412 macrophages, endothelial cells, and vascular smooth muscle cells. *Mol Pharmacol* 2008;  
413 **74**: 777-784.  
414
- 415 28. Crawford C, Kennedy-Lydon TM, Callaghan H, *et al.* Extracellular nucleotides affect  
416 pericyte-mediated regulation of rat in situ vasa recta diameter. *Acta Physiol (Oxf)* 2011;  
417 **202**: 241-251.  
418
- 419 29. Bailey MA, Turner CM, Hus-Citharel A, *et al.* P2Y receptors present in the native and  
420 isolated rat glomerulus. *Nephron Physiol* 2004; **96**: p79-90.  
421

- 422 30. Ilatovskaya DV, Palygin O, Levchenko V, *et al.* Pharmacological characterization of the P2  
423 receptors profile in the podocytes of the freshly isolated rat glomeruli. *Am J Physiol Cell*  
424 *Physiol* 2013; **305**: C1050-1059.  
425
- 426 31. Forst AL, Olteanu VS, Mollet G, *et al.* Podocyte Purinergic P2X4 Channels Are  
427 Mechanotransducers That Mediate Cytoskeletal Disorganization. *J Am Soc Nephrol* 2016;  
428 **27**: 848-862.  
429
- 430 32. Hohenstein B, Renk S, Lang K, *et al.* P2Y1 gene deficiency protects from renal disease  
431 progression and capillary rarefaction during passive crescentic glomerulonephritis. *J Am*  
432 *Soc Nephrol* 2007; **18**: 494-505.  
433
- 434 33. Vonend O, Turner CM, Chan CM, *et al.* Glomerular expression of the ATP-sensitive P2X  
435 receptor in diabetic and hypertensive rat models. *Kidney Int* 2004; **66**: 157-166.  
436
- 437 34. Taylor SR, Turner CM, Elliott JI, *et al.* P2X7 deficiency attenuates renal injury in  
438 experimental glomerulonephritis. *J Am Soc Nephrol* 2009; **20**: 1275-1281.  
439
- 440 35. Burnstock G, Evans LC, Bailey MA. Purinergic signalling in the kidney in health and  
441 disease. *Purinergic Signal* 2014; **10**: 71-101.  
442
- 443 36. Praetorius HA, Leipziger J. Intrarenal purinergic signaling in the control of renal tubular  
444 transport. *Annu Rev Physiol* 2010; **72**: 377-393.  
445
- 446 37. Castrop H, Huang Y, Hashimoto S, *et al.* Impairment of tubuloglomerular feedback  
447 regulation of GFR in ecto-5'-nucleotidase/CD73-deficient mice. *J Clin Invest* 2004; **114**:  
448 634-642.  
449
- 450 38. Stockand JD, Mironova E, Bugaj V, *et al.* Purinergic inhibition of ENaC produces  
451 aldosterone escape. *J Am Soc Nephrol* 2010; **21**: 1903-1911.  
452
- 453 39. Bailey MA, Imbert-Teboul M, Turner C, *et al.* Evidence for basolateral P2Y(6) receptors  
454 along the rat proximal tubule: functional and molecular characterization. *J Am Soc*  
455 *Nephrol* 2001; **12**: 1640-1647.  
456
- 457 40. Bailey MA, Shirley DG. Effects of extracellular nucleotides on renal tubular solute  
458 transport. *Purinergic Signal* 2009; **5**: 473-480.  
459
- 460 41. Bagorda A, Guerra L, Di Sole F, *et al.* Extracellular Adenine Nucleotides Regulate Na<sup>+</sup>/H<sup>+</sup>  
461 Exchanger NHE3 Activity in A6-NHE3 Transfectants by a cAMP/PKA-dependent  
462 Mechanism. *J Membr Biol* 2002; **188**: 249-259.  
463
- 464 42. Lee YJ, Park SH, Jeung TO, *et al.* Effect of adenosine triphosphate on phosphate uptake in  
465 renal proximal tubule cells: involvement of PKC and p38 MAPK. *J Cell Physiol* 2005; **205**:  
466 68-76.  
467
- 468 43. Jin W, Hopfer U. Purinergic-mediated inhibition of Na<sup>+</sup>-K<sup>+</sup>-ATPase in proximal tubule  
469 cells: elevated cytosolic Ca<sup>2+</sup> is not required. *Am J Physiol* 1997; **272**: C1169-1177.  
470

- 471 44. Bailey MA. Inhibition of bicarbonate reabsorption in the rat proximal tubule by  
472 activation of luminal P2Y1 receptors. *Am J Physiol Renal Physiol* 2004; **287**: F789-796.  
473
- 474 45. Vekaria RM, Unwin RJ, Shirley DG. Intraluminal ATP concentrations in rat renal tubules. *J*  
475 *Am Soc Nephrol* 2006; **17**: 1841-1847.  
476
- 477 46. Jensen ME, Odgaard E, Christensen MH, *et al.* Flow-induced  $[Ca^{2+}]_i$  increase depends on  
478 nucleotide release and subsequent purinergic signaling in the intact nephron. *J Am Soc*  
479 *Nephrol* 2007; **18**: 2062-2070.  
480
- 481 47. Bjaelde RG, Arnadottir SS, Overgaard MT, *et al.* Renal epithelial cells can release ATP by  
482 vesicular fusion. *Front Physiol* 2013; **4**: 238.  
483
- 484 48. Silva GB, Garvin JL. TRPV4 mediates hypotonicity-induced ATP release by the thick  
485 ascending limb. *Am J Physiol Renal Physiol* 2008; **295**: F1090-1095.  
486
- 487 49. Silva G, Beierwaltes WH, Garvin JL. Extracellular ATP stimulates NO production in rat  
488 thick ascending limb. *Hypertension* 2006; **47**: 563-567.  
489
- 490 50. Silva GB, Garvin JL. Extracellular ATP inhibits transport in medullary thick ascending  
491 limbs: role of P2X receptors. *Am J Physiol Renal Physiol* 2009; **297**: F1168-1173.  
492
- 493 51. Marques RD, de Bruijn PI, Sorensen MV, *et al.* Basolateral P2X receptors mediate  
494 inhibition of NaCl transport in mouse medullary thick ascending limb (mTAL). *Am J*  
495 *Physiol Renal Physiol* 2012; **302**: F487-494.  
496
- 497 52. Marques RD, Praetorius HA, Leipziger J. P2Y2 receptor knock-out mice display normal  
498 NaCl absorption in medullary thick ascending limb. *Front Physiol* 2013; **4**: 280.  
499
- 500 53. Shirley DG, Bailey MA, Unwin RJ. In vivo stimulation of apical P2 receptors in collecting  
501 ducts: evidence for inhibition of sodium reabsorption. *Am J Physiol Renal Physiol* 2005.  
502
- 503 54. Pochynyuk O, Bugaj V, Rieg T, *et al.* Paracrine regulation of the epithelial Na<sup>+</sup> channel in  
504 the mammalian collecting duct by purinergic P2Y2 receptor tone. *J Biol Chem* 2008; **283**:  
505 36599-36607.  
506
- 507 55. Deetjen P, Thomas J, Lehrmann H, *et al.* The luminal P2Y receptor in the isolated  
508 perfused mouse cortical collecting duct. *J Am Soc Nephrol* 2000; **11**: 1798-1806.  
509
- 510 56. Lehrmann H, Thomas J, Kim SJ, *et al.* Luminal P2Y<sub>2</sub> receptor-mediated inhibition of Na<sup>+</sup>  
511 absorption in isolated perfused mouse CCD. *J Am Soc Nephrol* 2002; **13**: 10-18.  
512
- 513 57. Wildman SS, Marks J, Turner CM, *et al.* Sodium-dependent regulation of renal amiloride-  
514 sensitive currents by apical P2 receptors. *J Am Soc Nephrol* 2008; **19**: 731-742.  
515
- 516 58. Kearney PM, Whelton M, Reynolds K, *et al.* Global burden of hypertension: analysis of  
517 worldwide data. *Lancet* 2005; **365**: 217-223.  
518

- 519 59. Stokes L, Scurrah K, Ellis JA, *et al.* A loss-of-function polymorphism in the human P2X4  
520 receptor is associated with increased pulse pressure. *Hypertension* 2011; **58**: 1086-1092.  
521
- 522 60. Palomino-Doza J, Rahman TJ, Avery PJ, *et al.* Ambulatory blood pressure is associated  
523 with polymorphic variation in P2X receptor genes. *Hypertension* 2008; **52**: 980-985.  
524
- 525 61. Gidlof O, Smith JG, Melander O, *et al.* A common missense variant in the ATP receptor  
526 P2X7 is associated with reduced risk of cardiovascular events. *PLoS One* 2012; **7**: e37491.  
527
- 528 62. Ghiadoni L, Rossi C, Duranti E, *et al.* P2X7 receptor polymorphisms do not influence  
529 endothelial function and vascular tone in neo-diagnosed, treatment-naive essential  
530 hypertensive patients. *J Hypertens* 2013; **31**: 2362-2369.  
531
- 532 63. McHugh SM, Roman S, Davis B, *et al.* Effects of genetic variation in the P2RX7 gene on  
533 pharmacodynamics of a P2X(7) receptor antagonist: a prospective genotyping approach.  
534 *Br J Clin Pharmacol* 2012; **74**: 376-380.  
535
- 536 64. Ivy JR, Bailey MA. Pressure natriuresis and the renal control of arterial blood pressure. *J*  
537 *Physiol* 2014; **592**: 3955-3967.  
538
- 539 65. Nishiyama A, Majid DS, Taher KA, *et al.* Relation between renal interstitial ATP  
540 concentrations and autoregulation-mediated changes in renal vascular resistance. *Circ*  
541 *Res* 2000; **86**: 656-662.  
542
- 543 66. Zhang Y, Robson SC, Morris KL, *et al.* Impaired natriuretic response to high-NaCl diet plus  
544 aldosterone infusion in mice overexpressing human CD39, an ectonucleotidase  
545 (NTPDase1). *Am J Physiol Renal Physiol* 2015; **308**: F1398-1408.  
546
- 547 67. Rieg T, Bunday RA, Chen Y, *et al.* Mice lacking P2Y<sub>2</sub> receptors have salt-resistant  
548 hypertension and facilitated renal Na<sup>+</sup> and water reabsorption. *FASEB J* 2007.  
549
- 550 68. Pandit MM, Inscho EW, Zhang S, *et al.* Flow regulation of endothelin-1 production in the  
551 inner medullary collecting duct. *Am J Physiol Renal Physiol* 2015; **308**: F541-552.  
552
- 553 69. Ji X, Naito Y, Hirokawa G, *et al.* P2X(7) receptor antagonism attenuates the hypertension  
554 and renal injury in Dahl salt-sensitive rats. *Hypertens Res* 2012; **35**: 173-179.  
555
- 556 70. Ji X, Naito Y, Weng H, *et al.* P2X7 deficiency attenuates hypertension and renal injury in  
557 deoxycorticosterone acetate-salt hypertension. *Am J Physiol Renal Physiol* 2012; **303**:  
558 F1207-1215.  
559
- 560 71. Masin M, Young C, Lim K, *et al.* Expression, assembly and function of novel C-terminal  
561 truncated variants of the mouse P2X7 receptor: re-evaluation of P2X7 knockouts. *Br J*  
562 *Pharmacol* 2012; **165**: 978-993.  
563
- 564 72. Menzies RI, Unwin RJ, Dash RK, *et al.* Effect of P2X4 and P2X7 receptor antagonism on  
565 the pressure diuresis relationship in rats. *Front Physiol* 2013; **4**: 305.  
566

- 567 73. Eser A, Colombel JF, Rutgeerts P, *et al.* Safety and Efficacy of an Oral Inhibitor of the  
568 Purinergic Receptor P2X7 in Adult Patients with Moderately to Severely Active Crohn's  
569 Disease: A Randomized Placebo-controlled, Double-blind, Phase IIa Study. *Inflamm*  
570 *Bowel Dis* 2015; **21**: 2247-2253.
- 571  
572 74. Harada H, Chan CM, Loesch A, *et al.* Induction of proliferation and apoptotic cell death  
573 via P2Y and P2X receptors, respectively, in rat glomerular mesangial cells. *Kidney Int*  
574 2000; **57**: 949-958.
- 575  
576 75. Turner CM, Tam FW, Lai PC, *et al.* Increased expression of the pro-apoptotic ATP-  
577 sensitive P2X7 receptor in experimental and human glomerulonephritis. *Nephrol Dial*  
578 *Transplant* 2007; **22**: 386-395.
- 579  
580 76. Deplano S, Cook HT, Russell R, *et al.* P2X7 receptor-mediated Nlrp3-inflammasome  
581 activation is a genetic determinant of macrophage-dependent crescentic  
582 glomerulonephritis. *J Leukoc Biol* 2013; **93**: 127-134.
- 583  
584 77. Ferrari D, Pizzirani C, Adinolfi E, *et al.* The P2X7 receptor: a key player in IL-1 processing  
585 and release. *J Immunol* 2006; **176**: 3877-3883.
- 586  
587 78. Zhao J, Wang H, Dai C, *et al.* P2X7 blockade attenuates murine lupus nephritis by  
588 inhibiting activation of the NLRP3/ASC/caspase 1 pathway. *Arthritis Rheum* 2013; **65**:  
589 3176-3185.
- 590  
591 79. Yan Y, Bai J, Zhou X, *et al.* P2X7 receptor inhibition protects against ischemic acute  
592 kidney injury in mice. *Am J Physiol Cell Physiol* 2015; **308**: C463-472.
- 593  
594 80. Solini A, Menini S, Rossi C, *et al.* The purinergic 2X7 receptor participates in renal  
595 inflammation and injury induced by high-fat diet: possible role of NLRP3 inflammasome  
596 activation. *J Pathol* 2013; **231**: 342-353.
- 597  
598 81. Lu D, Insel PA. Cellular mechanisms of tissue fibrosis. 6. Purinergic signaling and response  
599 in fibroblasts and tissue fibrosis. *Am J Physiol Cell Physiol* 2014; **306**: C779-788.
- 600  
601 82. Lu D, Insel PA. Hydrolysis of extracellular ATP by ectonucleoside triphosphate  
602 diphosphohydrolase (ENTPD) establishes the set point for fibrotic activity of cardiac  
603 fibroblasts. *J Biol Chem* 2013; **288**: 19040-19049.
- 604  
605 83. Chen JB, Liu WJ, Che H, *et al.* Adenosine-5'-triphosphate up-regulates proliferation of  
606 human cardiac fibroblasts. *Br J Pharmacol* 2012; **166**: 1140-1150.
- 607  
608 84. Solini A, Iacobini C, Ricci C, *et al.* Purinergic modulation of mesangial extracellular matrix  
609 production: role in diabetic and other glomerular diseases. *Kidney Int* 2005; **67**: 875-885.
- 610  
611 85. Goncalves RG, Gabrich L, Rosario A, Jr., *et al.* The role of purinergic P2X7 receptors in the  
612 inflammation and fibrosis of unilateral ureteral obstruction in mice. *Kidney Int* 2006; **70**:  
613 1599-1606.
- 614

- 615 86. Komada T, Usui F, Shirasuna K, *et al.* ASC in renal collecting duct epithelial cells  
616 contributes to inflammation and injury after unilateral ureteral obstruction. *Am J Pathol*  
617 2014; **184**: 1287-1298.  
618
- 619 87. Guo C, Masin M, Qureshi OS, *et al.* Evidence for functional P2X4/P2X7 heteromeric  
620 receptors. *Mol Pharmacol* 2007; **72**: 1447-1456.  
621
- 622 88. Antonio LS, Stewart AP, Xu XJ, *et al.* P2X4 receptors interact with both P2X2 and P2X7  
623 receptors in the form of homotrimers. *British Journal of Pharmacology* 2011; **163**: 1069-  
624 1077.  
625
- 626 89. Kim MJ, Turner CM, Hewitt R, *et al.* Exaggerated renal fibrosis in P2X4 receptor-deficient  
627 mice following unilateral ureteric obstruction. *Nephrol Dial Transplant* 2014; **29**: 1350-  
628 1361.  
629
- 630 90. Kohler D, Eckle T, Faigle M, *et al.* CD39/ectonucleoside triphosphate diphosphohydrolase  
631 1 provides myocardial protection during cardiac ischemia/reperfusion injury. *Circulation*  
632 2007; **116**: 1784-1794.  
633
- 634 91. Grenz A, Zhang H, Hermes M, *et al.* Contribution of E-NTPDase1 (CD39) to renal  
635 protection from ischemia-reperfusion injury. *FASEB J* 2007; **21**: 2863-2873.  
636
- 637 92. Roberts V, Lu B, Rajakumar S, *et al.* The CD39-adenosinergic axis in the pathogenesis of  
638 renal ischemia-reperfusion injury. *Purinergic Signal* 2013; **9**: 135-143.  
639
- 640 93. Rajakumar SV, Lu B, Crikis S, *et al.* Deficiency or inhibition of CD73 protects in mild  
641 kidney ischemia-reperfusion injury. *Transplantation* 2010; **90**: 1260-1264.  
642
- 643 94. Wang YM, McRae JL, Robson SC, *et al.* Regulatory T cells participate in CD39-mediated  
644 protection from renal injury. *Eur J Immunol* 2012; **42**: 2441-2451.  
645
- 646 95. Gu BJ, Zhang W, Worthington RA, *et al.* A Glu-496 to Ala polymorphism leads to loss of  
647 function of the human P2X7 receptor. *J Biol Chem* 2001; **276**: 11135-11142.  
648
- 649 96. Sluyter R, Shemon AN, Wiley JS. Glu496 to Ala polymorphism in the P2X7 receptor  
650 impairs ATP-induced IL-1 beta release from human monocytes. *J Immunol* 2004; **172**:  
651 3399-3405.  
652
- 653 97. Ali Z, Laurijssens B, Ostefeld T, *et al.* Pharmacokinetic and pharmacodynamic profiling  
654 of a P2X7 receptor allosteric modulator GSK1482160 in healthy human subjects. *Br J Clin*  
655 *Pharmacol* 2013; **75**: 197-207.  
656
- 657 98. Stevens M, Oltean S. Alternative Splicing in CKD. *J Am Soc Nephrol* 2016; **27**: 1596-1603.  
658
- 659 99. Bartlett R, Stokes L, Sluyter R. The P2X7 receptor channel: recent developments and the  
660 use of P2X7 antagonists in models of disease. *Pharmacol Rev* 2014; **66**: 638-675.  
661

- 662 100. de Torre-Minguela C, Barbera-Cremades M, Gomez AI, *et al.* Macrophage activation and  
663 polarization modify P2X7 receptor secretome influencing the inflammatory process. *Sci*  
664 *Rep* 2016; **6** :22586.

665

666

667



668 **Acknowledgements**

669 Research in the authors' laboratories was funded by The British Heart Foundation and Kidney  
670 Research UK.

671 **Disclosures**

672 RJU is currently on secondment as Chief Scientist to Cardiovascular and Metabolic Diseases  
673 (iMed CVMD) R&D, AstraZeneca, Mölndal, Sweden. FWKT and MAB have received research  
674 funding from AstraZeneca.

675

676

677

678

679 **Figure 1: The autocrine / paracrine purinoceptor system**

680 A range of stimuli including cellular stretch, trauma, or agonist binding triggers ATP release into  
681 the extracellular space. Ectonucleotidases located on the plasma membrane catalyse sequential  
682 hydrolysis of ATP to ADP, 5'AMP and adenosine. P1 receptors recognize adenosine while P2  
683 receptors bind di- and tri-phosphate nucleotide molecules. P2X receptors are non-selective  
684 cation channels with 3 protein subunits that may form homo- or heteromeric arrangements; all  
685 bind ATP. P2Y receptors are 7 transmembrane-spanning domain G-protein-coupled receptors;  
686 agonist preferences span adenosine and uracil di- and tri- nucleotides. NTPDase: ectonucleoside  
687 triphosphate diphosphohydrolase.

688

689 **Figure 2: P2 Receptors in the kidney**

690 P2Y and P2X receptor expression along the nephron: vasculature, glomeruli and tubules.

691

692 **Figure 3: P2XR related inflammation in (diabetic) kidney disease**

693 Local production of chemokines, adhesion molecules and inflammatory cytokines are  
694 upregulated under chronic stimulation of factors including hyperglycemia. Macrophages are the  
695 main infiltrating inflammatory cell type (expressing P2X7R) in both the glomerular and  
696 tubulointerstitial compartments where they contribute to extracellular matrix (ECM) secretion,  
697 amplification of the inflammatory cascade and eventually fibrosis.

698

699


700

701




703

704 **Figure 1**



705


706

707 **Figure 2**

708

709

710 **Figure 3**



711