

# THE UNIVERSITY of EDINBURGH

# Edinburgh Research Explorer

## Mode of resistance to viral lysis affects host growth across multiple environments in the marine picoeukaryote Ostreococcus tauri

#### Citation for published version:

Heath, S & Collins, S 2016, 'Mode of resistance to viral lysis affects host growth across multiple environments in the marine picoeukaryote Ostreococcus tauri', Environmental Microbiology, vol. 18, no. 12, pp. 4628-4639. https://doi.org/10.1111/1462-2920.13586

#### **Digital Object Identifier (DOI):**

10.1111/1462-2920.13586

#### Link: Link to publication record in Edinburgh Research Explorer

**Document Version:** Peer reviewed version

**Published In: Environmental Microbiology** 

#### **General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1	Mode of resistance to viral lysis affects host growth across multiple environments in		
2	the marine picoeukaryote Ostreococcus tauri		
3			
4	Sarah E. Heath & Sinead Collins		
5			
6	Institute of Evolutionary Biology, School of Biological Sciences, University of		
7	Edinburgh, Ashworth Laboratories, The King's Buildings, West Mains Road,		
8	Edinburgh EH9 3FL, UK.		
9			
10	Corresponding author: Sarah E. Heath.		
10 11	Corresponding author: Sarah E. Heath. Institute of Evolutionary Biology, School of Biological Sciences, University of		
10 11 12	Corresponding author: Sarah E. Heath. Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, Ashworth Laboratories, The King's Buildings, West Mains Road,		
10 11 12 13	Corresponding author: Sarah E. Heath. Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, Ashworth Laboratories, The King's Buildings, West Mains Road, Edinburgh EH9 3FL, UK.		
10 11 12 13 14	Corresponding author: Sarah E. Heath. Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, Ashworth Laboratories, The King's Buildings, West Mains Road, Edinburgh EH9 3FL, UK. +44(0)131 650 8657		
10 11 12 13 14 15	Corresponding author: Sarah E. Heath. Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, Ashworth Laboratories, The King's Buildings, West Mains Road, Edinburgh EH9 3FL, UK. +44(0)131 650 8657 s.heath-2@sms.ed.ac.uk		

17 Running title: *O. tauri* virus resistance in multiple environments

#### 18 SUMMARY

19

20 Viruses play important roles in population dynamics and as drivers of evolution in 21 single-celled marine phytoplankton. Viral infection of Ostreococcus tauri often 22 causes cell lysis, but two spontaneously arising resistance mechanisms occur: resistant 23 cells that cannot become infected and resistant producer cells that are infected but not 24 lysed, and which may slowly release viruses. As of yet, little is known about how 25 consistent the effects of viruses on their hosts are across different environments. To 26 measure the effect of host resistance on host growth, and to determine whether this 27 effect is environmentally dependent, we compared the growth and survival of 28 susceptible, resistant and resistant producer O. tauri cells under five environmental 29 conditions with and without exposure to O. tauri virus. While the effects of exposure 30 to virus on growth rates did not show a consistent pattern in populations of resistant 31 cells, there were several cases where exposure to virus affected growth in resistant 32 hosts, sometimes positively. In the absence of virus, there was no detectable cost of 33 resistance in any environment, as measured by growth rate. In fact, the opposite was 34 the case, with populations of resistant producer cells having the highest growth rates 35 across four of the five environments.

36

#### **37 INTRODUCTION**

38

39 Marine viruses play a large role in nutrient and energy cycling in the oceans. Viral 40 lysis of single celled organisms releases large quantities of organic matter into the 41 environment, making nutrients available for use by bacteria and algae. This process 42 has been termed the viral shunt (Wilhelm and Suttle, 1999). Studies on marine viruses 43 typically focus on the importance of viruses in nutrient cycling and the release of 44 organic matter through cell lysis. Despite the important role of marine viruses in 45 ecosystem function across many environments, from nutrient rich coastal waters to 46 more oligotrophic regions of the open ocean (Bruussard, 2004), host-virus interactions 47 are typically studied in single environments. Here, we use the Ostreococcus 48 tauri/Ostreococcus tauri virus model system to investigate variation in host-virus 49 interactions across environments to understand (1) whether susceptibility/resistance to 50 viruses changes with environmental change, and (2) whether the growth effect of host 51 resistance depends on environmental context or resistance type.

52

53 We explore the relationship between host responses to environmental change and the 54 resistance strategies of those hosts using the marine picoeukaryote Ostreococcus tauri 55 (order Mamiellales). O. tauri is commonly isolated from Mediterranean lagoons that 56 are connected to the open ocean via narrow channels (Clerissi et al., 2014). These 57 channels limit the exchange of seawater between the lagoon and ocean, making 58 variations in the environmental salinity, pH, temperature and nutrients more extreme 59 than in the open ocean (Bellec et al., 2010; Clerissi et al., 2014). Ostreococcus tauri 60 viruses (OtVs) have been sampled frequently in water samples collected from lagoon 61 and coastal waters where O. tauri is found. OtVs have strict host specificity (Clerissi 62 et al., 2012), and the three OtVs sequenced to date have all been described as lytic 63 viruses (Derelle et al., 2008; Weynberg et al., 2009, 2011). Virus infection of O. tauri 64 usually causes cell lysis in susceptible (S) cells, though two mechanisms of resistance 65 have been observed (Thomas et al., 2011). In the first case, viruses are unable to 66 infect the host, and these cells are referred to here as resistant (R). In the second case, 67 hosts are tolerant to viral infection and are able to slowly release them without 68 damage to the host cell. These cells are termed resistant producers (RP). In this paper, 69 we refer to the three cell types as resistance types.

70

71 Resistance type could have consequences for growth and other cell properties, such as 72 size and chlorophyll content. For example, a trade-off of acquiring resistance to viral 73 lysis may come as a fitness cost. This often occurs as reduced competitive ability 74 (Lenski, 1988; Bohannan et al., 2002) and sometimes reduced growth rate (Lennon et 75 al., 2007; Frickel et al., 2016). A modification in cell surface receptors to limit virus 76 attachment could also result in a loss of the original function of the protein, such as 77 metabolism or being able to target the host immune system. In several bacteria 78 species, loss of a bacteriophage receptor results in lower virulence of the bacteria in 79 its host, thereby lowering the fitness of resistant compared to non-resistant strains 80 (Seed *et al.*, 2012; León and Bastías, 2015). Lastly, strong resistance to one specific 81 virus strain may lead to increased susceptibility to lysis by other strains, as has been 82 observed in O. tauri (Clerissi et al., 2012) and cyanobacteria (Marston et al., 2012; 83 Avrani and Lindell, 2015).

84

The group of viruses that infects phytoplankton is the Phycodnaviruses. These viruses have been studied under environmental conditions that differ from a benign control

87 environment in a single driver, such as increases in temperature (Nagasaki and 88 Yamaguchi, 1998; Wells and Deming, 2006), nutrient (Bratbak et al., 1993, 1998; 89 Bellec et al., 2010; Clerissi et al., 2014), light (Bratbak et al., 1998; Weinbauer, 90 2004), UV (Jacquet and Bratbak, 2003), CO<sub>2</sub> (Larsen et al., 2007; Chen et al., 2014; 91 Maat et al., 2014) and pH levels (Weinbauer, 2004). When environmental conditions 92 are stressful, one consequence can be inactivation of the virus particle. This affects 93 host-virus interactions by preventing infection through structural degradation, the 94 inability of the virus to inject its genome into the host or the inability of the virus to 95 replicate (Børsheim, 1993; Jacquet and Bratbak, 2003). Additionally, since viral 96 replication and life cycle are often closely linked to host metabolism, environmental 97 changes such as increased temperature or nutrients will often have an indirect effect 98 on responses to viral attack (Weinbauer, 2004; Danovaro et al., 2011). Understanding 99 the role of viruses in marine communities requires investigating their activity across 100 environments. Here, we focus on the environmental changes of increased temperature, 101 decreased nutrients, decreased light and decreased salinity levels.

102

103 Previous studies of resistance in O. tauri found that when each resistance type was 104 maintained separately there was no significant difference in growth rates, such that a 105 cost of resistance was too low to be detected by differences in growth alone. 106 However, when resistant types were competed against each other, a competitive 107 hierarchy was observed in which S had the fastest growth rate, followed by R and 108 then by RP (Thomas et al., 2011). Since the three resistance types share the same 109 starting genotype, it is possible to make direct comparisons between them. In this 110 study, an experiment was performed in which three populations of each O. tauri 111 resistance type (S, R and RP) derived from a common ancestor were grown for one 112 week in the following environments in both the absence and presence of OtV5: high 113 temperature, low light, low phosphate, and low salt. These environments were 114 selected to represent relatively small variations from the control environment in which 115 the populations are normally maintained in the laboratory, so that the cells responded, 116 but were still able to grow at a rate that was measureable. The average number of cell 117 divisions per day over a single transfer cycle (7 days), cell size and cell chlorophyll 118 content were measured in the novel environments in the absence of OtV5. Offspring 119 production over a fixed period of time is a proxy for fitness in single celled organisms 120 in batch culture experiments (Brennan and Collins, 2015). Cell size and chlorophyll 121 content were measured as additional phenotypes, to examine effects on organismal 122 function other than cell division rates, since only small differences in growth were 123 detected previously (Thomas et al., 2011). After one week of growth in the novel 124 environment, all populations were inoculated with OtV5 and cell densities were 125 measured three days after inoculation to test for susceptibility to viral lysis.

126

127

129

130 *The effect of viral exposure on cell division rates depends on resistance type* 

131

132 After one week of growth in a novel environment, all populations were inoculated 133 with OtV5 and cell densities were measured three days later. Supporting Information 134 Tables S1 and S2 provide all statistical outputs in this study. Susceptibility of O. tauri 135 to OtV5 was driven by resistance type, as expected (ANOVA, resistance type  $\times$  virus treatment,  $F_{2,234} = 360.14$ , p <0.0001). After inoculation of O. tauri with OtV5. all R 136 137 and RP cells remained resistant to lysis and S cells remained susceptible (Figure 1). 138 Thus, OtV5 inoculation had a significant effect on cell density (ANOVA effect of 139 virus treatment on growth,  $F_{1,234} = 361.62$ , p < 0.0001), since populations of S cells 140 fell to almost zero (Figure 1). No difference was observed in resistance between R 141 and RP populations (t=0.46, p=0.66).

142

143 Counter to our expectation, the effect of virus inoculation did not vary with environment (ANOVA, environment × virus treatment,  $F_{4,234} = 0.89$ , p < 0.46). 144 145 However, environment alone had a significant effect on growth ( $F_{4,234} = 26.01$ , p 146 <0.0001), because of the S cell lysis in all environments. Additionally, an interaction 147 was identified between resistance type and environment (ANOVA, environment  $\times$ 148 resistance type,  $F_{8,234} = 6.09$ , p < 0.0001). For both R and RP cells, there were cases 149 where virus inoculation resulted in higher growth rates than the non-inoculated 150 controls (Figure 1). Cell densities were repeatedly higher in one inoculated population 151 (NG'13) than the control in the low salt environment in R cells and in the low light 152 environment for one population (NG27) in RP cells. This indicates that cell growth

153 can increase in response to viruses in resistant populations. This occurs consistently in 154 all replicates of a given population when it happens, but does not occur in all 155 populations of a resistance type. We also see cases where lysis in some populations of S cells is incomplete, notably in the low light (NG'2) and low salt (NG'3) 156 157 environments. Again, this does not occur in all populations, but it occurs reliably in 158 replicates of the same population. While these effects of environment on lysis are not 159 statistically significant because they do not occur over all populations within a 160 resistance type, it could have evolutionary and ecological effects on the occasions 161 when it does occur, which we discuss below.

162

#### 163 Growth rate varied across environments regardless of resistance type

164

165 All populations were grown in a novel environment in the absence of OtV5 for one 166 week, over which growth rate was measured. The response of O. tauri growth to the environment depended on resistance type (effect of environment  $\times$  resistance type,  $F_{8}$ . 167 114=4.45, p=0.0001). Additionally, regardless of resistance type, population growth 168 rates differed between environments (effect of environment on growth  $F_{4.114} = 231.39$ , 169 170 P < 0.0001) (Figure 2). Growth rates were higher in the control environment except for 171 a single RP population, NG'10, which divided rapidly in the low salt environment 172 (Figure 2). Populations grown in the low phosphate environment all had reduced 173 growth rates and showed less variation in growth than in all other environments.

174

### 175 The effect of resistance type on growth depends on environment

176

177 Resistance type alone did not significantly affect the growth rate of O. tauri 178 ( $F_{2,6}=2.88$ , p=0.1328). This is because S and R cells had similar population growth 179 rates in all environments (Figure 2). In contrast, some populations of RP had different 180 growth rates than both R and S cells. There was variation in growth rates between 181 replicate populations of RP cells, with some populations consistently showing 182 elevated growth rates. Two out of the three RP populations (NG'10 (shown by circle 183 in Fig 2) and NG'16 (shown by cross in Fig 2) had higher growth rates than S and R 184 cells in four out of the five environments ( $F_{3,5}=17.19$ , p=0.046). The single exception 185 was the low phosphate environment, where all resistance types had similar low 186 growth rates. These data indicate that there is either no cost or an undetectable cost of 187 resistance in terms of growth to either infection or lysis over a range of environments, 188 and that there can be a growth benefit of being resistant to lysis in some 189 environments, as evidenced by the rapid growth of some RP populations. The low or 190 absent cost of resistance is consistent with previous studies in single environments, 191 which have reported costs of resistance detectable in competitions, but too low to be 192 detectable by comparing growth rates (Thomas et al., 2011).

193

194 Populations resistant to lysis can have a growth advantage in some environments

195

In order to assess whether the S, R, and RP resistance types responded similarly to the different environments, environments were ranked from best to worst, based on population growth rates. All resistance types displayed highest growth rates in the control environment (See Supporting Information Table S3). R cells had the same rank order of environments as the S cells. Since the growth rates of the RP cells were highly variable relative to the other resistance types, containing two populations that grew quickly, the RP populations were grouped into fast growing (NG'10 and NG'16) and normal growing (NG27). RP cells showed the same rank order of environments for both the fast and normal growing populations, except in low salt for the fast growing populations. This was due to one population (NG'10) displaying exceptionally high growth. Growth rate was the same in the low salt and low light environments for the normal growing RP population. Fast growing RP cells had higher cell growth in all environments except low phosphate.

209

To measure how sensitive growth rates were to environmental change, the slopes of 210 211 the ranked environments were compared (Figure 3). The two fast growing RP 212 populations had a higher intercept (ANOVA effect of rank on growth, F 213  $_{1,125}$ =1112.56, p <0.0001), demonstrating the increase in growth rate compared to the 214 other populations. These data show that faster growing populations had a stronger 215 preference for environments in which they can grow more quickly, however in the 216 lowest ranking environment (which was low phosphate for all resistance types), these 217 populations grew equally badly.

218

219 Size and chlorophyll content vary between cells with different resistance types in
220 response to environment

221

After one week of growth in a novel environment without viruses, cell size and relative cell chlorophyll content were measured. Response of resistance type on cell size depended on environment (effect of environment × resistance type,  $F_{8,114}$ =5.48, p<0.0001). Regardless of resistance type, environment had a significant effect on cell size ( $F_{4,114}$ =77.93, p <0.0001). Cells were larger under low light (t=3.83, p=0.0002) and low phosphate conditions (t=7.49, p<0.0001), compared to the control environment (Figure 4). No significant effect of resistance type was observed on cell size ( $F_{2,6}$ =0.01, p=0.9945). However, under low phosphate, there was a large variation in cell size between the fast and normal growing RP populations.

231

232 The two fast-growing RP populations had smaller cells than the normal growing RP 233 population in the low phosphate environment. The RP population with normal growth 234 had cells that were similar in size to the S populations (Figure 4). To examine whether 235 fast growing RP populations had different cell sizes than did populations with normal 236 growth rates, *post hoc* models were used to analyse the two fast growing populations 237 separately. Overall, no significant effect of resistance type was observed on cell size 238 when normal and fast growing RP populations were analysed separately (ANOVA 239  $F_{2.6}=0.22$ , p=0.8812). Additionally, a model examining growth rate as a fixed effect was also performed. This showed a significant effect of growth rate ( $F_{1,99}$ =54.23, 240 241 p < 0.0001) and an interaction between resistance type and growth rate ( $F_{2.99} = 4.64$ , 242 p=0.01), although no effect of resistance type alone was detected ( $F_{2.6}=0.001$ , 243 p=0.99). However, the statistical power in this data set, which contained only one 244 population of normal growing RP cells and two populations of fast growing RP cells, 245 was low, such that the chances of detecting an effect of resistance type on cell size is 246 unlikely here even if one exists (power=0.142).

247

The effect of resistance type on chlorophyll content per cell volume depended on environment (effect of environment × resistance type,  $F_{8,114}$ =10.68, p < 0.0001). In addition, environment alone had a significant effect on relative chlorophyll per cell volume ( $F_{4,114}$ =120.45, p < 0.0001), however resistance type alone did not ( $F_{2.6}$ =1.61, p = 0.2757). Under low light, chlorophyll varied little between the three resistance types. In the other environments, S and R strategies usually displayed similar chlorophyll content levels with RP displaying lower chlorophyll levels in all environments except low phosphate.

256

257 By inspection, we see that the fast growing RP populations have less chlorophyll per 258 cell volume than the normal growing RP population in all environments except low 259 phosphate (Figure 5). We used a *post hoc* model with growth rate as a fixed effect to 260 investigate whether the fast growing RP populations also had different chlorophyll 261 contents. Growth rate had a significant effect on chlorophyll content ( $F_{1,99}$ =57.86, p 262 <0.0001), with fast growing RP populations having lower chlorophyll content, and the 263 effect of growth rate was dependent on environment ( $F_{4,99}=3.85$ , p=0.01) and 264 resistance type ( $F_{2,99}$ =6.27, p =0.003). Furthermore, when growth rate was considered 265 in the model, resistance type alone had a significant effect on chlorophyll content 266 ( $F_{2.6}$ =5.49, p =0.04), suggesting that the growth rate of the fast growing RP populations reduced chlorophyll content. 267

268

```
269 DISCUSSION
```

270

#### 271 *Effect of environment on host resistance*

272

We observed no differences in susceptibility of any of the populations to OtV5 over the environments tested. While the ability of the virus to lyse host cells did not depend on the environment, R and RP cells had different growth responses to viral exposure. There were two cases in which a resistant population repeatedly had a higher cell density after exposure to OtV5 than its paired control culture that was not inoculated.
We speculate that this may be a response to the virus, which causes the phytoplankton
cells to divide more rapidly. This would be advantageous if, for example, a population
that was made up of mixed susceptible and resistant cells were exposed to virus – any
resistant cell lineages that could increase their growth rate would then take over the
population by overgrowing any remaining resistant cells whose growth rate was
unaffected by exposure to virus.

284

285 We did not detect a growth cost of resistance when R and RP populations were grown 286 in the absence or presence of OtV5 after exposure to a novel environment. We expect 287 to see a trade-off for being resistant to viral infection, because if there were no cost 288 there should be a strong selection pressure for all cells to become resistant, yet we still 289 find susceptible populations both in the laboratory and in the ocean (Thomas et al., 290 2011; Clerissi et al., 2012). Previous work shows that that susceptible cells can have a 291 competitive advantage against resistant cells (Lenski, 1988). Additionally, we 292 speculate that resistance to one virus strain could make these cells susceptible to other 293 OtVs. Clerissi et al., (2012) showed that OtVs are mainly intraspecies-specific and 294 that hosts that are the most resistant to infection can often be infected by more 295 generalist viruses. This specificity could be caused by proteins involved in adaptive 296 behavior (Clerissi et al., 2012). Thus, we suggest that in addition to the abiotic 297 environment, biotic environment could play a large role in O. tauri resistance 298 strategy.

299

300 Since viruses are responsible for a large proportion of microbial death, there is strong301 selection on hosts for resistance or tolerance to viral infection. There are several

302 suggestions to explain the paradox of how susceptible algal cells and their viruses are 303 able to co-exist in marine environments without extinction of the host. One theory as 304 to how viruses and their hosts are able to coexist is that there must be a cost to being 305 resistant to infection. This is often expected to be a reduction in growth (Weinbauer, 306 2004), as has been observed in Synechococcus, in which there was a 20% reduction in 307 fitness compared to the ancestor in resistant strains (Lennon et al., 2007). Thus in the 308 absence of viruses, resistant cells often have a lower fitness. This could lead to 309 decreased numbers in the absence of viruses. An evolutionary "arms race" may occur 310 when viruses and their hosts adapt reciprocally to overcome resistance and infection, 311 respectively. We find little evidence for a cost of resistance in our study, but this may 312 be because the laboratory environments used are missing a key aspect of the natural 313 environment that, if present, results in a cost of resistance in O.tauri. Alternatively, 314 although deviating from the standard control environment, none of the environments 315 in this study were severely stressful, with even the low phosphate environment 316 allowing reasonable growth. Thus, it is possible that we did not detect a growth cost 317 because the changes to the environments used were relatively modest.

318

Various strategies for virus resistance have been reported in algae, including activation of programmed cell death (Bidle *et al.*, 2007), absence of metacaspase (caspase orthologues) protein expression (Bidle *et al.*, 2007), stage of the life cycle (Frada *et al.*, 2008), changes to cell surface receptor proteins (Tarutani *et al.*, 2006), colony formation (Brussaard *et al.*, 2007) and genetic mutations (Stoddard *et al.*, 2007). However, it is still unknown how *O. tauri* cells acquire their two resistance strategies. We found that short-term exposure to novel environments does not affect resistance type and we did not observe any cost of resistance leading to cells losingtheir resistance to OtV5.

328

329 *Effect of resistance type on population growth and other phenotypic traits* 

330

331 We found that after one week in a novel environment, growth rate of O. tauri, as 332 measured by the average number of cell divisions per day over seven days, varied 333 across environments for all resistance types. RP populations had the fastest average 334 rates of cell division in most environments. All resistance types showed the same 335 environmental preferences, with average cell division rates highest in the control 336 environment. The only exception was one RP population that divided rapidly in the 337 low salt environment. The lowest growth rates were observed in the low phosphate 338 environment, which was expected since these cells were deprived of a key nutrient.

339

340 Two of the three RP populations divided more rapidly than all of the S and R 341 populations. These populations were fast growing in many environments, including 342 the control environment, suggesting that the rapid growth is a general character of 343 these two RP populations, rather than a response to stress or novelty. This faster 344 growth rate in RP populations relative to S and R populations is in contrast with 345 previous studies on O. tauri. Thomas et al (2011) detected no difference in growth 346 rate between S, R and RP cells, although competition experiments revealed a small 347 reduction of fitness in RP compared to R, and R compared to S. Our results suggest 348 that the opposite can be true. Similarly to Thomas et al. (2011), we did not observe a 349 fitness cost in terms of growth rate for the remaining populations since S, R and 350 normal growing RP populations had similar growth rates across environments. This 351 was expected, at least in the control environment, where previous studies have only 352 been able to detect a minimal cost of resistance by using direct competitions. 353 Surprisingly, the two fast growing RP populations could not be detected as having 354 more rapid growth under low phosphate, however these populations responded 355 differently in their size and chlorophyll contents.

356

357 Reduced growth rate is often observed as a cost of resistance in microbes and has 358 been measured in several species (Lennon et al., 2007; Haaber and Middelboe, 2009). 359 Ecologically, a cost of resistance is part of "kill the winner" dynamics, where, it is 360 hypothesized that viruses kill the faster growing (susceptible) cells, and thus provide 361 an opportunity for slower growing (resistant) cells (Mojica and Brussaard, 2014). This 362 role for viruses requires that there be a cost of resistance. However, here we did not 363 detect a cost of resistance in terms of growth rate, since there was no environment in 364 which resistant cell types grew at slower rates than S cells. In fact, we observed the 365 opposite in two out of the three RP populations, where resistant cells grew faster than 366 the S populations across all environments except low phosphate. In cases where 367 resistant cells (R or RP populations) did not divide faster than susceptible ones, they 368 divided at the same rate. Taken together, this suggests that the cost of resistance to 369 OtVs is likely to be small or absent, and may not play into kill the winner dynamics. 370 This opens the question of how the appearance of resistance to OtVs affects both host 371 and viral ecology.

372

Environment affected cell size, whereas generally, resistance type did not. However,
under low phosphate, the two fast growing RP populations were smaller than the
normal growing RP populations, suggesting that under nutrient limitation these cells

376 were able to divide at a smaller cell size. Smaller phytoplankton cell size is often 377 selected for in nutrient limited environments since smaller cells have a larger surface area to volume ratio and a thinner diffusion boundary layer, thus facilitating nutrient 378 379 uptake (Finkel et al., 2010; Peter and Sommer, 2015). Although fast growing RP 380 populations in this selection environment were smaller than the normal growing RP 381 population, their cell size was not different from the fast growing RP populations in 382 the other environments. The control was the only environment in which fast growing 383 RP populations were larger than the normal growing populations, indicating that there may be a (direct or indirect) fitness benefit associated with the increased size of the 384 385 RP type under control conditions.

386

387 In contrast to previous studies, all populations in the low phosphate environment, 388 except fast growing RP, increased in cell size. Cell division of larger phytoplankton 389 cells requires greater nutrient concentrations, which can decrease the division rate. 390 Since cells in the low phosphate environment had a reduced growth rate in terms of 391 cell divisions, this could have resulted in cells that reached a larger volume even 392 though the environment was phosphate-poor. It has previously been suggested that 393 increasing algal cell size, and thus the volume to surface area ratio, can facilitate 394 reduced phosphorus uptake under phosphate-limited conditions, and that this 395 adaptation response may be more favourable than decreasing cell size (Supraha *et al.*, 396 2015). A common response of coccolithophores to phosphate limitation is reduced 397 growth rate and increased cell size (Šupraha et al., 2015).

398

399 Smaller phytoplankton cells have often been observed growing at higher temperatures400 in natural environments, which is thought to arise from the temperature-size rule (e.g.

401 Atkinson *et al.*, 2003; Morán *et al.*, 2010). These studies used large temperature 402 ranges, but there was no effect of the modest increase in temperature on cell size in 403 this study. Smaller cells have also been reported to cope better with both light 404 limitation and light saturation compared to larger cells due to a reduction in internal 405 shading (Geider *et al.*, 1986; Raven, 1998; Finkel *et al.*, 2010). We found no 406 significant difference in cell size under low light, although there was a non-significant 407 trend for cells to be slightly larger in low than under control light.

408

409 Environment was found to have a significant effect on chlorophyll content per cell 410 volume, whereas resistance type alone had no effect. We observed lower chlorophyll 411 per cell volume in all environments compared to the control except high temperature. 412 Although resistance type alone did not have an effect, growth rate had a significant 413 effect on chlorophyll content per cell volume when included in the model and normal 414 growing S, R and RP cells in the control and high temperature environments had the 415 highest chlorophyll levels across all environments. In contrast, fast growing RP 416 populations showed no significant difference in chlorophyll content per cell volume 417 across all five environments. All populations had their lowest growth rates in the low 418 phosphate environment and cells in this environment had the lowest chlorophyll 419 content, except for fast growing RP populations. Fast growing RP populations had 420 lower chlorophyll content than the normal growing RP population in all environments 421 except for low phosphate.

422

423 One experiment using cultures of different phytoplankton groups found that 424 chlorophyll content was lower during both nitrogen and phosphorus depletion 425 (Riemann *et al.*, 1989). Additionally, phytoplankton cells grown under low nutrients 426 have been observed to decrease their photosynthesis rates (Litchman et al., 2003; 427 Spilling *et al.*, 2015). This may be due to the cells allocating resources to synthesizing chloroplasts under nutrient limitation. In our study the control environment was the 428 429 preferred one, and it is possible that cells were unable to synthesise large quantities of 430 chlorophyll in the other (less permissive) environments since their energy was 431 allocated to growth. It is possible that under elevated temperature, the metabolism of 432 O. tauri was increased, leading the cells to synthesise more chlorophyll. Temperature 433 did not affect chlorophyll a content in diatoms (Sigaud and Aidar, 1993). Salinity appears to affect different phytoplankton species differently, with some species 434 435 showing no change in chlorophyll content across a range of salinities, and others 436 having higher chlorophyll contents at the optimum salinity for growth (McLachlan, 437 1961; Sigaud and Aidar, 1993).

438

#### 439 *Concluding remarks*

440

441 Resistance of microbes to virus infection often comes at a cost, with one common 442 observation being a reduction in growth compared to susceptible cells in the 443 population. In this study, our aim was to measure resistance to viruses in O. tauri 444 across different environments and to determine whether the magnitude of a cost of 445 resistance depends on environmental context. We did not observe a cost of resistance 446 as measured by cell divisions, cell size or chlorophyll content in the present study. 447 Growth rates of O. tauri were reduced when grown in low phosphate, however this 448 did not affect the ability of OtV5 to lyse susceptible cells in this environment. 449 Additionally, although growth rates were lower than the controls in high temperature, 450 low light and low salinity, OtV5 still caused cell lysis of susceptible cells. Indeed,

451 some populations that were tolerant to infection (RP populations) had evolved high 452 growth rates, and some RP populations also increased their growth rates after 453 exposure to viruses. Both observations suggest that resistance strategy could have 454 interesting ecological consequences by changing the relative fitness of different 455 populations.

#### 456 EXPERIMENTAL PROCEDURES

457

#### 458 *Susceptible and resistant populations used in this experiment*

459

O. tauri populations were obtained from N. Grimsley, Observatoire Océanologique, 460 461 Banyuls-sur-Mer. Three susceptible populations (NG'2, NG'3 and NG'4), three 462 resistant producer populations (NG'10, NG'16 and NG27) and three resistant 463 populations (NG5, NG'13 and NG26) were used in this study. We used three 464 biological replicates for each population in each environment. All populations were 465 derived from a single clone of O. tauri (RCC 4221) and therefore had the same 466 starting genotype (see Thomas et al., 2011). All populations have since been maintained separately. All RP populations were tested for viral production prior to the 467 468 start of the experiment (See Supporting Information Figure S1).

469

#### 470 Culturing conditions

471

472 Populations were grown in batch culture. Culture medium was prepared using 0.22 473  $\mu$ m filtered Instant Ocean artificial seawater (salinity 30 ppt) aerated with 400 ppm 474 CO<sub>2</sub> and supplemented with Keller and f/2 vitamins. Control cultures were maintained 475 in a 14:10 light:dark cycle at 85  $\mu$ mol photon m<sup>-2</sup> s<sup>-1</sup> and at a constant temperature of 476 18°C (Table 1).

477

For the selection experiments, *O. tauri* populations were grown without exposure to
viruses in the control environment and four selection environments. The selection
regimes used were high temperature, low light, low phosphate and low salinity (Table

1). Cultures were acclimated in each selection environment for one week, followed by

482 one week of growth in each environment.

483

Environment	Control	Treatment
Phosphate (µM)	10	5
Salinity (ppt)	30	25
Light ( $\mu$ mol m <sup>-2</sup> s <sup>-1</sup> )	85	60
Temperature (°C)	18	20

484

For the low phosphate environment, phosphate was reduced by preparing Keller 485 486 media with only half the amount of  $\beta$ -glycerophosphate that would normally be used. 487 Although the phosphate concentration in the low phosphate environment is not low compared to natural seawater (0.01-2.99  $\mu$ mol l<sup>-1</sup> in the Leucate lagoon where *O. tauri* 488 489 and OtV5 inhabit (Clerissi et al., 2014)), it is low compared to the control media in 490 which the populations had been maintained prior to the experiment. For culture 491 medium with a lower salinity than the control, Instant Ocean was added to reach a 492 salinity of 25 ppt. For the low light condition, culture flasks were wrapped in 0.15 493 neutral density foil to give a light intensity of 1000 lux. Cultures in the high 494 temperature condition were maintained on a heat mat (Exo Terra Heat Wave substrate 495 heat mat) set at 20°C.

496

497 *The effect of viral exposure on cell division rates* 

498

Following one week of growth in the selection environment, each sample wasinoculated with a fresh suspension of OtV5 particles to test whether it was susceptible

501 or resistant to the virus. Samples were tested by inoculating 1 ml cell culture at a 502 density of  $10^5$  with  $10 \,\mu$ l OtV5 in 48-well plates with three replicates for each sample. 503 Controls that were not inoculated with viruses were used as a control for cell growth. 504 Cell density was measured using a FACSCanto flow cytometer 3 days after 505 inoculation.

506

507 Population growth of susceptible and resistant populations across different
508 environments

509

Following the acclimation period, average cell densities per day of all cultures were measured over one week of growth in each environment. Cells were counted using a BD FACSCanto II (BD Biosciences) flow cytometer before the first transfer and after seven days of growth. Each population was counted in triplicate. The cell counts were converted to cells per millilitre and the number of divisions per day was calculated using equation (1).

516

517 (1) 
$$\mu(d^{-1}) = \frac{\log_2(\frac{N_t}{N_0})}{t-t_0}$$

518

where  $N_t$  and  $N_0$  are the cell densities (cells ml<sup>-1</sup>) at times  $t_1$  and  $t_0$  (days). This measures the average number of cell divisions per ancestor over a single growth cycle and allows a comparison of offspring production between environments (Brennan and Collins, 2015). This is useful if different environments produce different growth curves since populations with different growth strategies can be compared. This calculation is also not sensitive to small differences in  $N_0$ , which is important if the 525 population size reached during the acclimation period differs between environments526 or resistance types.

527

528 Cell size and chlorophyll content of populations with different resistance types across529 environments

530

531 Cell size and relative chlorophyll content per cell volume were determined using a 532 FACSCanto flow cytometer. Cell size was inferred from FSC (forward scatter), which was calibrated using beads of known sizes (1µm, 3µm and 6.6µm). Chlorophyll 533 534 fluorescence was inferred by measuring PerCP-Cy5.5 emission with excitation at 535 488nm. Relative chlorophyll was analysed by taking the average chlorophyll fluorescence for all susceptible populations in the control environment and setting this 536 537 to a value of 1, with chlorophyll measurements of all populations relative to this 538 value.

539

540 Statistical analysis

541

542 Data were analysed with linear mixed effects models using the statistical package 543 nlme in R (version 3.2.0) to identify differences in growth rates between the different 544 environments after one week of growth and after virus inoculation. Environment and 545 resistance type were fixed effects when analyzing growth under different 546 environments, and environment, resistance type and treatment were fixed effects 547 when analyzing virus inoculation under different environments. Population was a 548 random effect in both models.

549

*Post hoc* mixed effects models were used to examine whether growth rate had an effect on cell size and chlorophyll content in cells. Environment, resistance type and growth rate (cells divisions per day) were set as fixed effects with populations as the only random effect.

554

#### 555 ACKNOWLEDGEMENTS

556

This work was funded by an EASTBIO DTP to SEH. SC is supported by a Royal
Society (UK) University Research Fellowship and an ERC starting grant under the
FP7 framework.

560

#### 561 **REFERENCES**

562

563 Atkinson, D., Ciotti, B.J., and Montagnes, D.J.S. (2003) Protists decrease in size

564 linearly with temperature: ca. 2.5% degrees C(-1). *Proc. Biol. Sci.* 270: 2605–11.

- Avrani, S. and Lindell, D. (2015) Convergent evolution toward an improved growth
- rate and a reduced resistance range in Prochlorococcus strains resistant to phage.
- 567 *Proc. Natl. Acad. Sci. U. S. A.* **112**: E2191–200.
- 568 Bellec, L., Grimsley, N., Derelle, E., Moreau, H., and Desdevises, Y. (2010)

Abundance, spatial distribution and genetic diversity of Ostreococcus tauri

- 570 viruses in two different environments. *Environ. Microbiol. Rep.* **2**: 313–21.
- 571 Bidle, K.D., Haramaty, L., Barcelos E Ramos, J., and Falkowski, P. (2007) Viral
- activation and recruitment of metacaspases in the unicellular coccolithophore,
- 573 Emiliania huxleyi. Proc. Natl. Acad. Sci. U. S. A. 104: 6049–54.
- 574 Blanc-Mathieu, R., Verhelst, B., Derelle, E., Rombauts, S., Bouget, F.-Y., Carré, I., et

- al. (2014) An improved genome of the model marine alga Ostreococcus tauri
- unfolds by assessing Illumina de novo assemblies. *BMC Genomics* **15**: 1103.
- 577 Bohannan, B.J.M., Kerr, B., Jessup, C.M., Hughes, J.B., and Sandvik, G. (2002)
- 578 Trade-offs and coexistence in microbial microcosms. *Antonie van Leeuwenhoek,*579 *Int. J. Gen. Mol. Microbiol.* 81: 107–115.
- 580 Børsheim, K.Y. (1993) Native marine bacteriophages. *FEMS Microbiol. Lett.* 102:
  581 141–159.
- 582 Bratbak, G., Egge, J.K., and Heldal, M. (1993) Viral mortality of the marine alga
- 583 Emiliania huxleyi (Haptophyceae) and termination of algal blooms. *Mar. Ecol.*584 *Prog. Ser.* 93: 39–48.
- 585 Bratbak, G., Jacobsen, A., Heldal, M., Nagasaki, K., and Thingstad, F. (1998) Virus
- production in Phaeocystis pouchetii and its relation to host cell growth and
  nutrition. *Aquat. Microb. Ecol.* 16: 1–9.
- Brennan, G. and Collins, S. (2015) Growth responses of a green alga to multiple
  environmental drivers. *Nat. Clim. Chang.* 5: 892–897.
- 590 Brussaard, C.P.D., Bratbak, G., Baudoux, A.C., and Ruardij, P. (2007) Phaeocystis

and its interaction with viruses. *Biogeochemistry* **83**: 201–215.

- 592 Bruussard, C.P. (2004) Viral control of phytoplankton populations a review. J.
  593 *Eukarvot. Microbiol.* 51: 125–138.
- 594 Chen, S., Gao, K., and Beardall, J. (2014) Viral attack exacerbates the susceptibility
- of a bloom-forming alga to ocean acidification. *Glob. Chang. Biol.* 629–636.
- 596 Clerissi, C., Desdevises, Y., and Grimsley, N. (2012) Prasinoviruses of the marine
- 597 green alga Ostreococcus tauri are mainly species specific. J. Virol. 86: 4611–9.
- 598 Clerissi, C., Grimsley, N., Subirana, L., Maria, E., Oriol, L., Ogata, H., et al. (2014)
- 599 Prasinovirus distribution in the Northwest Mediterranean Sea is affected by the

- 600 environment and particularly by phosphate availability. *Virology* **466-467**: 146–
- 601 157.
- 602 Danovaro, R., Corinaldesi, C., Dell'Anno, A., Fuhrman, J. a., Middelburg, J.J., Noble,
- R.T., and Suttle, C. a. (2011) Marine viruses and global climate change. *FEMS Microbiol. Rev.* 35: 993–1034.
- 605 Derelle, E., Ferraz, C., Escande, M.-L., Eychenié, S., Cooke, R., Piganeau, G., et al.
- 606 (2008) Life-cycle and genome of OtV5, a large DNA virus of the pelagic marine
  607 unicellular green alga Ostreococcus tauri. *PLoS One* **3**: e2250.
- 608 Finkel, Z. V., Beardall, J., Flynn, K.J., Quigg, A., Rees, T.A. V, and Raven, J.A.
- 609 (2010) Phytoplankton in a changing world: Cell size and elemental
- 610 stoichiometry. J. Plankton Res. **32**: 119–137.
- Frada, M., Probert, I., Allen, M.J., Wilson, W.H., and de Vargas, C. (2008) The
- 612 "Cheshire Cat" escape strategy of the coccolithophore Emiliania huxleyi in
- 613 response to viral infection. *Proc. Natl. Acad. Sci. U. S. A.* **105**: 15944–9.
- 614 Frickel, J., Sieber, M., and Becks, L. (2016) Eco-evolutionary dynamics in a
- 615 coevolving host-virus system. *Ecol. Lett.* **19**: 450–459.
- 616 Geider, R., Platt, T., and Raven, J. (1986) Size dependence of growth and
- 617 photosynthesis in diatoms: a synthesis. *Mar. Ecol. Ser.* **30**: 93–104.
- 618 Haaber, J. and Middelboe, M. (2009) Viral lysis of Phaeocystis pouchetii:
- 619 implications for algal population dynamics and heterotrophic C, N and P cycling.
- 620 *Isme J* **3**: 430–441.
- 621 Jacquet, S. and Bratbak, G. (2003) Effects of ultraviolet radiation on marine virus-
- 622 phytoplankton interactions. *FEMS Microbiol. Ecol.* **44**: 279–289.
- 623 Larsen, J.B., Larsen, a., Thyrhaug, R., Bratbak, G., and Sandaa, R. -a. (2007) Marine
- 624 viral populations detected during a nutrient induced phytoplankton bloom at

- 625 elevated pCO<sub>2</sub> levels. *Biogeosciences Discuss.* **4**: 3961–3985.
- Lennon, J.T., Khatana, S.A.M., Marston, M.F., and Martiny, J.B.H. (2007) Is there a
  cost of virus resistance in marine cyanobacteria? *ISME J.* 1: 300–312.
- 628 Lenski, R.E. (1988) Experimental studies of pleiotropy and epistasis in Escherichia
- 629 coli. I. variation in competitive fitness among mutants resistant to virus T4.
- 630 *Evolution (N. Y).* **42**: 425–432.
- 631 León, M. and Bastías, R. (2015) Virulence reduction in bacteriophage resistant
  632 bacteria. *Front. Microbiol.* 6: 1–7.
- 633 Litchman, E., Steiner, D., and Bossard, P. (2003) Photosynthetic and growth
- 634 responses of three freshwater algae to phosphorus limitation and daylength.
- 635 *Freshw. Biol.* **48**: 2141–2148.
- Maat, D., Crawfurd, K., Timmermans, K., and Brussard, C. (2014) Elevated CO2 and
  phosphate limitation favor Micromonas pusilla through stimulated growth and
  reduced viral impact. *Appl. Environ. Microbiol.*
- Marston, M.F., Pierciey, F.J., Shepard, A., Gearin, G., Qi, J., Yandava, C., et al.
- 640 (2012) Rapid diversification of coevolving marine Synechococcus and a virus.
- 641 *Pnas* **109**: 4544–4549.
- 642 McLachlan, J. (1961) The effect of salinity on growth and chlorophyll content in
- representative classes of unicellular marine algae. *Can. J. Microbiol.* 7: 399–406.
- 644 Mojica, K.D.A. and Brussaard, C.P.D. (2014) Factors affecting virus dynamics and
- 645 microbial host-virus interactions in marine environments. *FEMS Microbiol*.
- 646 *Ecol.* **89**: 495–515.
- 647 Morán, X.A.G., López-Urrutia, Á., Calvo-Díaz, A., and Li, W.K.W. (2010)
- 648 Increasing importance of small phytoplankton in a warmer ocean. *Glob. Chang.*
- *Biol.* **16**: 1137–1144.

- 650 Nagasaki, K. and Yamaguchi, M. (1998) Effect of temperature on the algicidal
- activity and the stability of HaV (Heterosigma akashiwo virus). *Aquat. Microb. Ecol.* 15: 211–216.
- 653 Peter, K.H. and Sommer, U. (2015) Interactive effect of warming, nitrogen and
- 654 phosphorus limitation on phytoplankton cell size. *Ecol. Evol.* **5**: 1011–1024.
- Raven, J.A. (1998) The twelfth Tansley Lecture. Small is beautiful: The
- picophytoplankton. *Funct. Ecol.* **12**: 503–513.
- 657 Riemann, B., Simonsen, P., and Stensgaard, L. (1989) The carbon and chlorophyll
- content of phytoplankton from various nutrient regimes. *J. Plankton Res.* 11:
  1037–1045.
- 660 Seed, K.D., Faruque, S.M., Mekalanos, J.J., Calderwood, S.B., Qadri, F., and Camilli,
- A. (2012) Phase Variable O Antigen Biosynthetic Genes Control Expression of
  the Major Protective Antigen and Bacteriophage Receptor in Vibrio cholerae O1.
- 663 *PLoS Pathog.* 8.:
- Sigaud, T.C.S. and Aidar, E. (1993) Salinity and temperature effects on the growth
  and chlorophyll-a content of some planktonic aigae. *Bol. do Inst. Ocean.* 41: 95–
  103.
- 667 Spilling, K., Ylöstalo, P., Simis, S., and Seppälä, J. (2015) Interaction effects of light,
- temperature and nutrient limitations (N, P and Si) on growth, stoichiometry and
- 669 photosynthetic parameters of the cold-water diatom Chaetoceros wighamii. *PLoS*
- 670 *One* **10**: 1–18.
- 671 Stoddard, L.I., Martiny, J.B.H., and Marston, M.F. (2007) Selection and
- 672 characterization of cyanophage resistance in marine Synechococcus strains.
- 673 *Appl. Environ. Microbiol.* **73**: 5516–22.
- 674 Šupraha, L., Gerecht, A.C., Probert, I., and Henderiks, J. (2015) Eco-physiological

- adaptation shapes the response of calcifying algae to nutrient limitation. *Sci. Rep.*5: 16499.
- 677 Tarutani, K., Nagasaki, K., and Yamaguchi, M. (2006) Virus adsorption process

determines virus susceptibility in Heterosigma akashiwo (Raphidophyceae).

- 679 *Aquat. Microb. Ecol.* **42**: 209–213.
- Thomas, R., Grimsley, N., Escande, M.-L., Subirana, L., Derelle, E., and Moreau, H.
- 681 (2011) Acquisition and maintenance of resistance to viruses in eukaryotic

682 phytoplankton populations. *Environ. Microbiol.* **13**: 1412–20.

- 683 Weinbauer, M.G. (2004) Ecology of prokaryotic viruses. 28: 127–181.
- 684 Wells, L.E. and Deming, J.W. (2006) Effects of temperature, salinity and clay
- particles on inactivation and decay of cold-active marine Bacteriophage 9A.
- 686 *Aquat. Microb. Ecol.* **45**: 31–39.
- 687 Weynberg, K.D., Allen, M.J., Ashelford, K., Scanlan, D.J., and Wilson, W.H. (2009)
- From small hosts come big viruses: the complete genome of a second
- 689 Ostreococcus tauri virus, OtV-1. *Environ. Microbiol.* **11**: 2821–39.
- 690 Weynberg, K.D., Allen, M.J., Gilg, I.C., Scanlan, D.J., and Wilson, W.H. (2011)
- 691 Genome sequence of Ostreococcus tauri virus OtV-2 throws light on the role of
- 692 picoeukaryote niche separation in the ocean. J. Virol. 85: 4520–9.
- 693 Wilhelm, S.W. and Suttle, C.A. (1999) Viruses and nutrient cycles in the sea.
- 694 *Bioscience* **49**: 781–788.
- 695
- 696
- 697 TABLE AND FIGURE LEGENDS
- 698

Table 1. A comparison of the control environment and the environment treatmentsthat were used for each environmental condition in this study.

701

Figure 1. Mean cell densities ml<sup>-1</sup> of resistant (R), resistant producer (RP) and 702 703 susceptible (S) O. tauri cells. Inoculated = cells inoculated with OtV5, Not inoculated 704 = control cultures that were grown for the same amount of time, but not inoculated 705 with OtV5. The dashed line represents the starting densities of the cultures at  $10^5$  cells 706 ml<sup>-1</sup>. There were three biological replicates for each populations. Boxes represent the 707 interquartile range with the median indicated as the thick black line inside the box, 708 and whiskers extend to the highest and lowest values within  $1.5 \times$  the inter-quartile 709 range from the edge of the box. Outlier data beyond the end of the whiskers are 710 plotted as points.

711

712 Figure 2. Mean growth rates, measured as average number of cell divisions per day 713 over 7 days, of susceptible (S), resistant (R) and resistant producer (RP) O. tauri cells 714 grown in five environments in the absence of OtV5. There were three populations for 715 each resistance type, with three biological replicates for each population. Boxes 716 represent the interquartile range with the median indicated as the thick black line 717 inside the box, and whiskers extend to the highest and lowest values within  $1.5 \times$  the 718 inter-quartile range from the edge of the box. Outlier data beyond the end of the 719 whiskers are plotted as points.

720

Figure 3. Ranked environment by average cell divisions per day over 7 days (±
SEM) for susceptible (S), resistant (R) and resistant producer (RP and RPfast) cells.
Environments were ranked in order from best to worst for each resistance type based

on growth rate in the absence of OtV5, where 1 is the environment with the highest
growth rate. Fast and normal growing resistant producers have been plotted separately
for visual purposes.

727

Figure 4. Mean cell size for susceptible (S), resistant (R) and resistant producer (RP) cells after seven days of growth in the absence of viruses in five environments. There were three populations for each resistance type, with three biological replicates for each populations. Boxes represent the interquartile range with the median indicated as the thick black line inside the box, and whiskers extend to the highest and lowest values within  $1.5 \times$  the inter-quartile range from the edge of the box. Outlier data beyond the end of the whiskers are plotted as points.

735

Figure 5. Mean relative chlorophyll to cell size for susceptible (S), resistant (R) and resistant producer (RP) cells after seven days of growth in the absence of viruses in five environments. There were three populations for each resistance type, with three biological replicates for each population. Boxes represent the interquartile range with the median indicated as the thick black line inside the box, and whiskers extend to the highest and lowest values within  $1.5 \times$  the inter-quartile range from the edge of the box. Outlier data beyond the end of the whiskers are plotted as points.

743

Table S1. ANOVA results of a linear mixed effects model to analyse interaction

resistance type and treatment (with or without OtV5 effects of environment, resistance type and treatment (with or without OtV5

746 inoculation) on *O. tauri* cell density. Population was a random effect.

747

Table S2. ANOVA results of a linear mixed effects model to analyse interaction

effects of environment, resistance type, treatment (with or without OtV5 inoculation)

and growth rate (fast RP or normal) on *O. tauri* growth rate, as measured by cell

divisions per day. Population was set as a random effect in all models.

752

Table S3. Ranked environments by fitness as measured by cell divisions per day for each resistance type. Environments were ranked in order from best to worst, where 1 is the environment with the highest growth rate. Fast and normal growing resistant producers were ranked separately to compare slopes.

757

Figure S1. Mean cell densities ml<sup>-1</sup> (±SEM) of O. tauri strain RCC4221 three days 758 759 after inoculation with supernatant from Resistant Producing populations (NG'10, 760 NG'16 and NG27). To ensure that the RP populations being used in this experiment 761 were producing infectious viruses and releasing them to their external surroundings, 762 we used the supernatant of these strains to infect susceptible O. tauri cells. Populations NG'10, NG'16 and NG'27 were aliquoted into 2ml Eppendorf tubes and 763 centrifuged at 8000  $\times$  g for 15 min. Next, 400 µl of supernatant was carefully 764 765 removed without drawing up any cells from the pellet at the bottom of the tube, and 766 used to inoculate 1 ml of susceptible O. tauri strain RCC4221. Eight replicates were 767 performed. A positive control was performed using known OtV5, and a negative 768 control was performed by adding Keller media. Controls were performed in 769 quadruplicate. Cells were left to grow for 3 days after which their densities were 770 measured using a FACSCanto flow cytometer. We observed cell lysis resulting from 771 inoculation with supernatant from all three RP populations, showing that there was 772 active virus in the media taken from these cultures.