THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Maximizing Coordination Capsule-Guest Polar Interactions in Apolar Solvents Reveals Significant Binding

Citation for published version:

August, DP, Nichol, GS & Lusby, PJ 2016, 'Maximizing Coordination Capsule-Guest Polar Interactions in Apolar Solvents Reveals Significant Binding', *Angewandte Chemie International Edition*.
<https://doi.org/10.1002/anie.201608229>

Digital Object Identifier (DOI):

[10.1002/anie.201608229](https://doi.org/10.1002/anie.201608229)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Angewandte Chemie International Edition

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Maximizing Coordination Capsule-Guest Polar Interactions in Apolar Solvents Reveals Significant Binding**

David P. August, Gary S. Nichol and Paul J. Lusby*

Abstract: Guest encapsulation underpins the functional properties of self-assembled capsules yet identifying systems capable of strongly binding small organic molecules in solution remains a challenge. Most coordination capsules rely on the hydrophobic effect to ensure effective solution-phase association. In contrast, we show that using non-interacting anions in apolar solvents can maximize favorable interactions between a cationic Pd_2L_4 host and charge-neutral guests resulting in a dramatic increase in binding strength. With quinone-type guests, association constants in excess of 10^8 M^{-1} were observed, comparable to the highest previously recorded for a metallocapsule. Modulation of guests' optoelectronic properties was also observed, with encapsulation either changing or switching-on luminescence not present in the bulk-phase.

Supramolecular capsules appear at the forefront of research efforts because their propensity to partition whole molecules from the bulk-phase produces interesting properties ranging from sensing^[1] through catalysis^[2] to the stabilization of reactive species.^[3] With coordination systems, binding charge neutral guests provides a notable challenge because of the competition with associated counter-anions or cations.^[4] As a result, polar solvents are typically favored as these stabilize the counter-charged species outside of the cavity.^[5] Certain solvents, such as water, can also provide a strong and universal driving-force for guest encapsulation through solvophobic desolvation pathways.^[6] However, metallo-organic capsules often possess a mix of hydrophobic and hydrophilic regions—usually large apolar aromatic surfaces linked by polar coordination vertices—such that binding can be difficult to predict and also require a trade-off with possible favorable polar interactions.^[7] Here we show that it is possible to attain significant binding, comparable with the strongest previously reported by a coordination capsule in water^[5a,8]—almost 10^9 M^{-1} for a charge-neutral guest—by maximizing non-covalent interactions in apolar solvents.^[9]

The system we selected to study was the Pd_2L_4 capsule, 1^{4+} (Figure 1a), first reported by Hooley,^[10] in anticipation that (a) the low charge would aid investigation in apolar solvents; (b) the strong Pd-pyridine interactions would ensure the integrity of the anion-free cavity; (c) it would be possible to better the modest binding ($<20 \text{ M}^{-1}$) previously reported for various aromatic guests in DMSO.^[10a] Molecular modelling also indicated that the o-pyridyl positions (H_a) are polarized by the Pd^{II} ions creating pockets of H-

bond donors that can form complementary interactions with guests such as quinones (Figure 1b).^[11] Promisingly, when excess naphthoquinone, G^1 , was added to $1\cdot 4\text{OTf}$ in CD_3CN , the ^1H NMR spectrum of the mixture showed significant changes when compared to the individual species (Figure 2). While the single set of host-guest signals indicated that the interaction was dynamic relative to the NMR timescale, it was notable that the inside cage resonances (H_a , H_e) and two of the guest (H_y , H_z) were most shifted. Also, whereas H_e , H_y and H_z all moved upfield due to mutual shielding by host and guest aromatic surfaces, H_a was downfield shifted, supporting the initial supposition that binding would be driven by multiple $\text{CH}\cdots\text{O}$ H-bonds.

Figure 1. (a) Chemical structure of the $\text{Pd}_2\text{L}_4^{4+}$ cage, 1^{4+} ; (b) Energy-minimised model of naphthoquinone G^1 within the cavity of 1^{4+} showing attractive electrostatic surface potentials between the electron deficient CH regions of the capsule (shown in blue) and electron-rich areas provided by the guest (shown in red).

Figure 2. Partial ^1H NMR spectra (500 MHz, CD_3CN , 300 K) of a) naphthoquinone, G^1 , only; b) a mixture of $1\cdot 4\text{OTf}$ with excess G^1 ; c) $1\cdot 4\text{OTf}$ only. The lettering refers to those shown in Figures 1 and 3.

We next sought to assess the strength of binding between G^1 and 1^{4+} (Table 1). Starting with $1\cdot 4\text{OTf}$ in CD_3CN , plotting the change in chemical shifts ($\Delta\delta$) of the host when titrated with G^1 produced multiple curves that fitted a 1:1 binding isotherm, which

[*] Mr. D. P. August, Dr. G. S. Nichol, Dr. P. J. Lusby
EaStCHEM School of Chemistry, Joseph Black Building, David
Brewster Road, Edinburgh, UK, EH9 3FJ
E-mail: Paul.Lusby@ed.ac.uk

[**] We thank the University of Edinburgh for a Principal's Career
Development Scholarship (D.P.A.).

Supporting information for this article (including synthesis,
characterization data, X-ray analysis and details of titration
experiments) is given via a link at the end of the document.

COMMUNICATION

gave a global association constant, K_a , of 210 M^{-1} (Table 1, Entry 1; see Supporting Information for details). Encouraged that the affinity for **G**¹ was ten-fold higher than the previous best guest,^[10a] several different solvents were screened (Table 1, Entries 2–5), which indicated that apolar solvents promote better binding (Table 1, Entries 4–5). Surmising that even stronger binding was possibly being masked by tight ion-pairing, other capsule salts, **1**·**4X**, were then prepared either directly from the relevant Pd^{II} source ($X^- = \text{BF}_4^-$) or by adding excess NaX or KX to **1**·**4OTf** ($X^- = \text{PF}_6^-$, SbF_6^- , BArF^- [$\text{BArF} = \text{B}(3,5\text{-(CF}_3)_2\text{C}_6\text{H}_3)_4^-$]).^[12] Non-capsule salts were removed by exploiting the low solubility of **1**·**4X** in either methanol or water, while ¹⁹F NMR spectroscopy confirmed anion metathesis. Interestingly, comparing the ¹H NMR spectra of **1**·**4X** (Figure S3, S27) indicates that the stronger coordinating anions, OTf[−] and BF₄[−] in particular, are likely to reside within the capsule's cavity, with internal signals H_a and H_b being notably deshielded by up to 0.2 ppm in the case of both **1**·**4OTf** and **1**·**4BF**₄.^[13] The affinity of **1**⁴⁺ for the different anions was also qualitatively observed using ESI-MS; **1**·**4OTf** exhibited dominant 2+ and 3+ charge states with two and one associated anions, respectively, while the “naked” **1**⁴⁺ was the major ion with **1**·**4BArF** (Figures S28–32). Measuring the K_a for **G**¹ with the additional ion-pair capsules **1**·**4X** in CD₃NO₂—the optimal solvent to balance solubility whilst maximizing favourable interactions—revealed that, as anticipated, replacing OTf[−] with weaker interacting anions (Table 1, Entries 5–9) increases the binding strength, with a significant 25-fold increase in the case of BArF[−].

Table 1. Association constants, K_a , for naphthoquinone, **G**¹, with various capsule ion-pairs, **1**·**4X**, in different solvents.^a

	X	Solvent	K_a / M^{-1}	$\Delta G / \text{kJ mol}^{-1}$
Entry 1	OTf	CD ₃ CN	210	13.2
Entry 2	OTf	CD ₃ OD	26	8.1
Entry 3	OTf	[D ₈]THF	290	14.1
Entry 4	OTf	CD ₂ Cl ₂	1800	18.7
Entry 5	OTf	CD ₃ NO ₂	2000	18.8
Entry 6	BF ₄	CD ₃ NO ₂	6500	21.7
Entry 7	PF ₆	CD ₃ NO ₂	13000	23.5
Entry 8	SbF ₆	CD ₃ NO ₂	22000	24.8
Entry 9	BArF	CD ₃ NO ₂	50000	26.8
Entry 10	BArF	CD ₃ OD	530	15.5
Entry 11	BArF	CD ₃ CN	1600	18.3
Entry 12	BArF	CD ₂ Cl ₂	350000 ^b	31.1

[a] Determined by ¹H NMR titration, errors are estimated to be <10%. [b] Competitive ¹H NMR titration with **G**³.

The affinity of **G**¹ for **1**·**4BArF** has also been measured in different solvents (Table 1, Entries 9–12). This analysis was more complicated with CD₂Cl₂ as a solvent (Table 1, Entry 12) because of capsule signal broadening during the titration, indicating guest exchange was occurring close to the NMR timescale. In this case,

K_a was determined using a competitive binding experiment with a stronger, slow exchange guest (see below and Supporting Information).^[14] The trend of increased binding with **1**·**4OTf** in solvents of decreasing polarity (Table 1, Entries 1–4) was mirrored by **1**·**4BArF** (Table 1, Entries 9–12), however, the latter produced globally higher affinities, from a factor of ten in more polar solvents through to a greater than 100-fold increase in CD₂Cl₂. Overall, the combination of weakly interacting anions and a non-polar solvent dramatically increases the K_a between **1**⁴⁺ and **G**¹ by 10⁴ (Table 1, Entry 2 vs. Entry 1) thus indicating that a major contribution to the binding free energy are the polar CH⋯O H-bonds.

Using the optimized ion-pair and solvent combination (**1**·**4BArF** in CD₂Cl₂), different potential guests were explored (Figure 3). Notably, **G**^{3–5} all showed slow in-out kinetics, which was most apparent with **G**⁵ due to the reduction in capsule symmetry caused by the different benzo rings of the guest (Figure S35). Addition of sub-stoichiometric **G**^{3–5} to **1**·**4BArF** also revealed that they were very tight binders as no free guest was detectable at concentrations above 50 μM.^[15] Strong association was also evident by preservation of the inclusion complexes under ESI-MS conditions (Figures S57–59). Consequently, association constants were obtained using ¹H NMR competitive titration experiments; K_a for **G**³ was measured using a large excess of the fast exchange guest **G**², while **G**⁴ was competed against **G**³ (see Supporting Information). Attempts to obtain a binding constant for **G**⁵ using competitive binding produced data of insufficient quality, however, the same experiment showed it was better than **G**³.

Figure 3. The log K_a values for selected molecules, with binding strength energies (kJ mol^{−1}) shown in parenthesis. Association constants measured in CD₂Cl₂ using **1**·**4BArF**, except **G**⁶, which was obtained in CD₃CN.

With the quinone series (**G**¹–**G**⁵), increasing the number of fused aromatic rings results in a significant increase in K_a . The difference between **G**¹ vs. **G**² and **G**² vs. **G**³ are fairly similar, with each additional aromatic ring adding about 10 kJ mol^{−1} to the binding strength.^[16] These energetic contributions are likely a result of additional edge-to-face interactions (CH–π H-bonds^[9a], see below), which is consistent with the significant shielding of H_a observed by ¹H NMR spectroscopy following host-guest complexation. With pentacenedione, **G**⁴, the extra two rings

produce a smaller increase, perhaps not unsurprisingly as these protrude further into the void between adjacent ligands. Nonetheless, the log K_a of 8.9 for $\mathbf{G}^4$ is, as far as we are aware, comparable to the highest for a charge neutral guest inside a coordination capsule. The crystal structure of $[\mathbf{G}^4 \subset \mathbf{1}]4\text{OTf}$ has also been obtained, using single crystals grown from CH_3CN and Et_2O (Figure 4).^[17] The solid state structure confirms the solution binding model with the oxygen atoms of $\mathbf{G}^4$ clearly located in the two pockets of four H_a atoms, with C—O distances ranging from 3.3 to 3.8 Å, indicating multiple $\text{CH}\cdots\text{O}$ H-bonds. Edge-to-face interactions between the extended aromatic surface of $\mathbf{G}^4$ and the four H_a atoms are also apparent (see above). In addition to quinones, $\mathbf{1}^{4+}$ also binds other guests with suitably disposed H-bond acceptor groups (e.g. $\mathbf{G}^{6-8}$). The log K_a of 4.0 for $\mathbf{G}^6$ was measured in CD_3CN to alleviate problems of intermediate exchange; a comparison with $\mathbf{G}^1$ under similar conditions (Table 1, entry 11) is consistent with the better H-bond acceptor properties of amides vs. enones, not least considering $\mathbf{G}^6$ lacks the additional benzo ring that adds 10 kJ mol⁻¹ to the binding strength of $\mathbf{G}^1$. A further interesting comparison can also be made to the classic tetraamide macrocycle reported by Hunter and co-workers,^[18] which binds $\mathbf{G}^2$, $\mathbf{G}^6$ and $\mathbf{G}^9$. Whereas the K_a for $[\mathbf{G}^2 \subset \mathbf{1}]^{4+}$ is an order of magnitude higher than the tetraamide macrocycle under similar conditions, and a solvent/anion adjusted value for $[\mathbf{G}^6 \subset \mathbf{1}]^{4+}$ would be at least comparable with the covalent host, in contrast $\mathbf{G}^9$ shows no evidence of encapsulation inside $\mathbf{1}^{4+}$.^[19] A molecular model of $\mathbf{G}^9$ revealed that the preferred chair conformation results in only a marginally smaller distance between H-bond acceptor oxygen atoms in comparison to $\mathbf{G}^2$ ($\Delta(\text{O}—\text{O}) = 0.1$ Å). Instead, the lack of binding could possibly be due to the non-linear orientation of carbonyl groups, coupled to the relative rigidity of the metallosupramolecular framework, thus not allowing an optimal arrangement of H-bonding interactions with both sets of CH donor pockets.

Figure 4. X-ray crystal structure of $[\mathbf{G}^4 \subset \mathbf{1}]4\text{OTf}$ (counteranions, solvent and non-interacting H atoms omitted for clarity). Color code: carbon of $\mathbf{1}^{4+}$, green; carbon of $\mathbf{G}^4$, orange; hydrogen, white; nitrogen, blue; oxygen, red; palladium, magenta.

The optoelectronic properties of guests $\mathbf{G}^{4-5}$ are modulated upon encapsulation within $\mathbf{1}^{4+}$. With $\mathbf{G}^5$ both the λ_{max} of the absorption and emission spectra are redshifted with respect to the free guest, by 70 and 34 nm, respectively (see Figures S66), a

possible consequence of the LUMO being stabilized by H-bonding to the capsule. Similar yet even more dramatic effects are seen with $\mathbf{G}^4$. Whereas both $\mathbf{1}^{4+}\text{BarF}$ and $\mathbf{G}^4$ are virtually colorless to the naked eye under ambient lighting, $[\mathbf{G}^4 \subset \mathbf{1}]\text{4BarF}$ is clearly yellow (Figure 5a, left). When held under a UV lamp, the difference is even more stark, with $[\mathbf{G}^4 \subset \mathbf{1}]\text{4BarF}$ showing strong emission whereas $\mathbf{G}^4$ alone shows little (Figure 5a, right). The switch-on emission of the host-guest complex has also been confirmed spectroscopically, both by titrating $\mathbf{1}^{4+}\text{BarF}$ into $\mathbf{G}^4$ (Figure 5b) and also $\mathbf{G}^4$ into $\mathbf{1}^{4+}\text{BarF}$ (Figure S63-64). In both cases, the emission intensity increases until a 1:1 ratio of $\mathbf{1}^{4+}\text{BarF}$ and $\mathbf{G}^4$ is reached, where after it remains constant, strongly indicating that the luminescence is due to the formation of $[\mathbf{G}^4 \subset \mathbf{1}]\text{4BarF}$. While many coordination cages have been shown to quench the emission of guests, due to heavy-atom effects and/or charge-transfer processes, those that either maintain or even enhance the optoelectronic properties of the encapsulated species are rare.^[20] In the case of $[\mathbf{G}^4 \subset \mathbf{1}]\text{4BarF}$, we likely attribute the increase in fluorescence with respect to the free guest due to preventing the formation of weakly-emissive aggregates.^[20a]

Figure 5. a) Images of 100 μM CD_2Cl_2 solutions of i) $\mathbf{G}^4$; ii) $[\mathbf{G}^4 \subset \mathbf{1}]\text{4BarF}$; iii) $\mathbf{1}^{4+}\text{BarF}$ under ambient lighting (left) and under a 365 nm UV lamp (right); b) Fluorescence titration of $\mathbf{1}^{4+}\text{BarF}$ into 100 μM of $\mathbf{G}^4$ in CH_2Cl_2 with excitation at 412 nm (isosbestic point of $\mathbf{G}^4$ and $[\mathbf{G}^4 \subset \mathbf{1}]\text{4BarF}$). A quantum yield enhancement factor of 15.6 was calculated from the relative peak intensities of $\mathbf{G}^4$ and $[\mathbf{G}^4 \subset \mathbf{1}]\text{4BarF}$. No further increase in emission intensity was observed upon addition of excess $\mathbf{1}^{4+}\text{BarF}$.

In conclusion, we have shown that minimizing the competitive interactions between a charged cationic cage and its associated anions can lead to a dramatic increase in the strength of charge-neutral guest binding in apolar solvents, giving association constants comparable to the highest previously observed for a metallosupramolecular capsule system. We are currently investigating how such electronic manipulation of guest molecules can be exploited for various applications.

Keywords: Host-guest • coordination capsule • self-assembly • non-coordinating anions • quinone

- [1] For representative examples, see a) A. Suzuki, K. Kondo, Y. Sei, M. Akita, M. Yoshizawa, *Chem. Commun.* **2016**, 52, 3151-3154; b) P. D. Frischmann, V. Kunz, F. Würthner *Angew. Chem. Int. Ed.* **2015**, 54, 7285–7289; *Angew. Chem.* **2015**, 127, 7393–7397; c) C. He, J. Wang, P. Wu, L. Jia, Y. Bai, Z. Zhangab, C. Duan, *Chem. Commun.* **2012**, 48, 11880–11882; d) O. Chepelin, J. Ujma, X. Wu, A. M. Z. Slawin, M. B. Pitak, S. J.

- Coles, J. Michel, A. C. Jones, P. E. Barran, P. J. Lusby, *J. Am. Chem. Soc.* **2012**, *134*, 19334–19337.
- [2] For representative examples, see a) P. Howlader, P. Das, E. Zangrando, P. S. Mukherjee, *J. Am. Chem. Soc.* **2016**, *138*, 1668–1676; b) W. Cullen, M. C. Misuraca, C. A. Hunter, N. H. Williams, M. D. Ward, *Nature Chem.* **2016**, *8*, 231–236; c) D. M. Kaphan, M. D. Levin, R. G. Bergman, K. N. Raymond, F. D. Toste, *Science*, **2015**, *350*, 1235–1238; d) A. G. Salles, S. Zarra, R. M. Turner, J. R. Nitschke, *J. Am. Chem. Soc.* **2013**, *135*, 19143–19146; e) T. Murase, Y. Nishijima, M. Fujita, *J. Am. Chem. Soc.* **2012**, *134*, 162–164.
- [3] For representative examples, see a) M. Yamashina, Y. Sei, M. Akita, M. Yoshizawa, *Nat. Commun.* **2014**, *5*, 4662; b) S. Horiuchi, T. Murase, M. Fujita, *J. Am. Chem. Soc.* **2011**, *133*, 12445–12447; c) P. Mal, B. Breiner, K. Rissanen, J. R. Nitschke, *Science* **2009**, *324*, 1697–1699; d) T. Iwasawa, R. J. Hooley, J. Rebek, Jr *Science* **2007**, *317*, 493–496.
- [4] For examples of counteranion / cation binding, see a) M. D. Johnstone, E. K. Schwarze, J. Ahrens, D. Schwarzer, J. J. Holstein, B. Dittich, F. M. Pfeffer, G. H. Clever, *Chem. Eur. J.* **2016**, DOI: 10.1002/chem.201602497; b) S. Freye, J. Hey, A. Torras-Galán, D. Stalke, R. Herbst-Irmer, M. John, G. H. Clever, *Angew. Chem. Int. Ed.* **2012**, *51*, 2191–2194; *Angew. Chem.* **2012**, *124*, 2233–2237; c) I. A. Riddell, M. M. J. Smulders, J. K. Clegg, Y. R. Hristova, B. Breiner, J. D. Thoburn, J. R. Nitschke, *Nature Chem.* **2012**, *4*, 751–756; d) C. R. K. Glasson, J. K. Clegg, J. C. McMurtrie, G. V. Meehan, L. F. Lindoy, C. A. Motti, B. Moubaraki, K. S. Murray, J. D. Cashion, *Chem. Sci.* **2011**, *2*, 540–543; e) Y. R. Hristova, M. M. J. Smulders, J. K. Clegg, B. Breiner, J. R. Nitschke, *Chem. Sci.* **2011**, *2*, 638; f) G. H. Clever, S. Tashiro, M. Shionoya, *Angew. Chem. Int. Ed.* **2009**, *48*, 7010–7012; *Angew. Chem.* **2009**, *121*, 7144–7146; g) J. S. Fleming, K. L. V. Mann, C.-A. Carraz, E. Psillakis, J. C. Jeffery, J. A. McCleverty, M. D. Ward, *Angew. Chem. Int. Ed.* **1998**, *37*, 1279–1281; *Angew. Chem.* **1998**, *110*, 1315–1318; h) D. L. Caulder, R. E. Powers, T. N. Parac, K. N. Raymond, *Angew. Chem. Int. Ed.* **1998**, *37*, 1840–1843; *Angew. Chem.* **1998**, *110*, 1940–1943; i) M. Fujita, D. Oguro, M. Miyazawa, H. Oka, K. Yamaguchi, K. Ogura, *Nature* **1995**, *378*, 469.
- [5] For examples of charge neutral-guest binding, see a) W. Cullen, S. Turega, C. A. Hunter, M. D. Ward, *Chem. Sci.* **2015**, *6*, 2790–2794; b) P. R. Symmers, M. J. Burke, D. P. August, P. I. T. Thomson, G. S. Nichol, M. R. Warren, C. J. Campbell, P. J. Lusby, *Chem. Sci.* **2015**, *6*, 756–760; c) S. Turega, W. Cullen, M. Whitehead, C. A. Hunter, M. D. Ward, *J. Am. Chem. Soc.* **2014**, *136*, 8475–8483; d) J. L. Bolliger, T. K. Ronson, M. Ogawa, J. R. Nitschke, *J. Am. Chem. Soc.* **2014**, *136*, 14545–14553; e) N. Kishi, Z. Li, Y. Sei, M. Akita, K. Yoza, J. S. Siegel, M. Yoshizawa, *Chem. Eur. J.* **2013**, *19*, 6313–6320; f) F. Schmitt, J. Freudenreich, N. P. E. Barry, L. Juillerat-Jeanneret, G. Süß-Fink, B. Therrien, *J. Am. Chem. Soc.* **2012**, *134*, 754–757; g) P. Mal, D. Schultz, K. Beyeh, K. Rissanen, J. R. Nitschke, *Angew. Chem. Int. Ed.* **2008**, *47*, 8297–8301; *Angew. Chem.* **2008**, *120*, 8421–8425; h) S. M. Biro, R. G. Bergman, K. N. Raymond, *J. Am. Chem. Soc.* **2007**, *129*, 12094–12095; i) T. Kusakawa, M. Fujita, *Angew. Chem. Int. Ed.* **1998**, *37*, 3142–3144; *Angew. Chem.* **1998**, *110*, 3327–3329.
- [6] a) F. Biedermann, W. M. Nau, H.-J. Schneider, *Angew. Chem. Int. Ed.* **2014**, *53*, 11158–11171; *Angew. Chem.* **2014**, *126*, 11338–11352; b) L. Yang, C. Adam, S. L. Cockroft, *J. Am. Chem. Soc.* **2015**, *137*, 10084–10087.
- [7] M. Whitehead, S. Turega, A. Stephenson, C. A. Hunter, M. D. Ward, *Chem. Sci.* **2013**, *4*, 2744–2751.
- [8] For strong binding of charge-neutral guests by a cofacial porphyrin dimeric coordination capsule in mixed dichloromethane / methanol, see T. Nakamura, H. Ube, M. Shionoya, *Angew. Chem. Int. Ed.* **2013**, *52*, 12096–12100; *Angew. Chem.* **2013**, *125*, 12318–12322.
- [9] For other examples of maximizing non-covalent interactions in non-polar solvents, see a) S. L. Cockroft, C. A. Hunter, *Chem. Commun.* **2006**, 3806–3808; b) G. M. Prentice, S. I. Pascu, S. V. Filip, K. R. West, G. D. Pantoş, *Chem. Commun.* **2015**, *51*, 8265–8268.
- [10] a) P. Liao, B. W. Langloss, A. M. Johnson, E. R. Knudsen, F. S. Tham, R. R. Julian, R. J. Hooley, *Chem. Commun.* **2010**, *46*, 4932–4934. For cisplatin guest binding in a structurally similar Pd₂L₄ cage system, see b) J. E. M. Lewis, E. L. Gavey, S. A. Cameron, J. D. Crowley, *Chem. Sci.* **2012**, *3*, 778–784. For a recent reviews on Pd₂L₄ capsules, see c) M. Han, D. M. Engelhard, G. H. Clever, *Chem. Soc. Rev.* **2014**, *43*, 1848–1860; d) A. Schmidt, A. Casini, F. E. Kühn, *Coord. Chem. Rev.* **2014**, *275*, 19–36.
- [11] For examples of metal ion polarised CH H-bonding, see a) A. J. Metherell, W. Cullen, A. Stephenson, C. A. Hunter, M. D. Ward, *Dalton Trans.* **2014**, *43*, 71–84. For examples of similar host-guest interactions in coordinations cages, see refs 4a, 5a and 8.
- [12] For recent examples of organocatalysis using cationic H-bond donors that employ weakly coordinating anions, see a) W. Xu, M. Arieno, H. Löw, K. Huang, X. Xie, T. Cruchter, Q. Ma, J. Xi, B. Huang, O. Wiest, L. Gong, E. Meggers, *J. Am. Chem. Soc.* **2016**, *138*, 8774–8780; b) Y. Fan, S. R. Kass, *Org. Lett.* **2016**, *18*, 188–191; c) S. Shirakawa, S. Liu, S. Kaneko, Y. Kumatabara, A. Fukuda, Y. Omagari, K. Maruoka, *Angew. Chem. Int. Ed.* **2015**, *54*, 15767–15770; *Angew. Chem.* **2015**, *127*, 15993–15996.
- [13] For anions such as OTf and BF₄, it is likely that strong coulombic attraction is supplemented by a well defined secondary coordination sphere that involves multiple *o*-pyridyl CH $\cdots$ O or CH $\cdots$ F H-bonds. For the X-ray structure of a similar primary and secondary coordination sphere, see a) C. Bianchini, J. Filippi, A. Lavacchi, W. Oberhauser, *Eur. J. Inorg. Chem.* **2011**, 1797–1805. With 1·4OTf and 1·4BF₄, it is anticipated that, in solution, two anions interact with the external *o*-pyridyl pockets (noticeable shifts of H_b in the ¹H NMR spectra are observed, Figure S27), however, it is suspected that both the coulombic repulsion and size limitation of the capsules cavity means that only one of the internal *o*-pyridyl sites is occupied at any one time. Similar anion binding within capsules has been observed in the solid state, see ref [9a] and also, b) J. E. M. Lewis, J. D. Crowley, *Supramol. Chem.* **2014**, *26*, 173–181.
- [14] While the errors associated with binding constants determined by competitive titration experiments are likely to be greater than 10%, such uncertainty is offset by the significant difference in K_a values, see Figure S38.
- [15] At concentrations less than 50 μ M, 1⁴⁺ starts to disassemble, see Figure S33.
- [16] A slightly larger increase in the binding strength is observed between the first and second additional benzo groups (9 vs. 12 kJ mol⁻¹), possibly indicating that for G² there is free rotation of the guest in the cavity around the C₂ axis that connects both CO groups. With G¹ the additional CH- π interactions would be partially offset by the loss in entropy caused by the lack of rotation for this larger guest, hence the subsequent difference between G¹ and G³ is larger.
- [17] CCDC 1492902 [G⁴ $\subset$ 1]4OTf contains the supplementary crystallographic data for this paper. This data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [18] H. Adams, F. J. Carver, C. A. Hunter, N. J. Osborne, *Chem. Commun.* **1996**, 2529–2530.
- [19] While small changes to the ¹H NMR signals of 1⁴⁺ signals were observed upon titration with G⁹, attempts to fit this data to a 1:1 binding model gave a meaningless K_a value (0.1 $\pm$ 96 M⁻¹). We attribute the small changes to other possible, weak binding modes, such as interactions with the external *o*-pyridyl H-bond donors.
- [20] a) K. Ono, J. K. Klosterman, M. Yoshizawa, K. Sekiguchi, T. Tahara, M. Fujita, *J. Am. Chem. Soc.* **2009**, *131*, 12526–12527; b) P. P. Neelakandan, A. Jiménez, J. R. Nitschke, *Chem. Sci.* **2014**, *5*, 908–915; c) M. Yamashina, M. M. Sartin, Y. Sei, M. Akita, S. Takeuchi, T. Tahara, M. Yoshizawa, *J. Am. Chem. Soc.* **2015**, *137*, 9266–9269.

WILEY-VCH

COMMUNICATION

A combination of apolar solvents and weakly interacting anions have been used to maximize the non-covalent interactions between a simple Pd₂L₄ host and various charge-neutral guest molecules, giving association constants comparable with the highest previously reported for a coordination capsule. Modulation of the guest's optoelectronic properties, notably either changing or switching-on luminescence not present in the bulk-phase, was also observed.

David P. August, Gary S. Nichol and Paul J. Lusby*

Page No. – Page No.

Maximizing Coordination Capsule-Guest Polar Interactions in Apolar Solvents Reveals Significant Binding