



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Characterisation of *Ramularia collo-cygni* laboratory mutants resistant to Succinate Dehydrogenase Inhibitors

Citation for published version:

Piotrowska, MJ, Fountaine, JM, Ennos, R, Kaczmarek, M & Burnett, F 2016, 'Characterisation of *Ramularia collo-cygni* laboratory mutants resistant to Succinate Dehydrogenase Inhibitors', *Pest Management Science*.
<https://doi.org/10.1002/ps.4442>

Digital Object Identifier (DOI):

[10.1002/ps.4442](https://doi.org/10.1002/ps.4442)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Pest Management Science

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Characterisation of *Ramularia collo-cygni* laboratory mutants resistant to Succinate Dehydrogenase Inhibitors

Marta J Piotrowska ^{a,*}, James M Fountaine ^{a,1}, Richard A Ennos ^b, Maciej Kaczmarek ^{a,2}, Fiona J Burnett ^a

^aCrop and Soil Systems Research Group, Scotland's Rural College, West Mains Road, Edinburgh, EH9 3JG, UK

^bInstitute of Evolutionary Biology, University of Edinburgh, Charlotte Auerbach Rd, Edinburgh, EH9 3FL, UK

¹Syngenta, Jealott's Hill International Research Centre, Bracknell, Berkshire RG42 6EY, UK

²Forest Research, Alice Holt Lodge, Farnham, Surrey, GU10 4LH, UK

*Corresponding author: Marta J. Piotrowska, Crop and Soil Systems Research Group, Scotland's Rural College, West Mains Road, Edinburgh, EH9 3JG, UK, Marta.Piotrowska@sruc.ac.uk, +441315354294, +441315354144 (fax)

Running title: SDHIs resistance in *Rcc* lab mutants

Keywords: *Ramularia* leaf spot, fungicide insensitivity, UV mutagenesis, pesticides, plant protection

Accession numbers:

- 1 *Rcc_SdhB*: KU758973
- 2 *Rcc_SdhC*: KU758974
- 3 *Rcc_SdhD*: KU758975

Abstract

BACKGROUND: *Ramularia collo-cygni* (*Rcc*) is responsible for Ramularia leaf spot (RLS), a foliar disease of barley contributing to serious economic losses. Protection against the disease has been almost exclusively based on fungicide applications, including Succinate Dehydrogenase Inhibitors (SDHIs). In 2015 the first field isolates of *Rcc* with reduced sensitivity to SDHIs were recorded in some European countries. This study established baseline sensitivity of *Rcc* to SDHIs in the UK and characterised mutations correlating with resistance to SDHIs in UV-generated mutants.

RESULTS: Five SDHI resistant isolates were generated by UV mutagenesis. In four of these mutants a single amino acid change in a target succinate dehydrogenase (*Sdh*) protein was associated with decrease in sensitivity to SDHIs. Three of these mutations were stably inherited in the absence of SDHI fungicide and resistant isolates did not demonstrate a fitness penalty. There were no detectable declines in sensitivity in field populations in years 2010-2012 in the UK.

CONCLUSIONS: SDHIs remain effective in controlling *Rcc* in the UK, however, given that the first isolates of *Rcc* with reduced sensitivity appeared in 2015 in other European countries in 2015, robust anti-resistance strategies need to be implemented to maintain effective disease control.

1 Introduction

Ramularia collo-cygni (*Rcc*) is the causal agent of Ramularia leaf spot (RLS), a major barley disease in the UK.¹ It can cause yield losses of up to 1 t ha⁻¹, corresponding to around 18% of average yield in the UK.² Although there

1 is an increasing interest in breeding for host resistance, there currently are no
2 lines of barley fully resistant to RLS, although varieties differ in their level of
3 susceptibility to the pathogen.^{1,3-6} Therefore protection against RLS remains
4 based on foliar fungicide applications. *Ramularia collo-cygni* has already
5 developed resistance to Quinone outside Inhibitors (QoIs), a fungicide class
6 which initially provided good control of the disease.⁷⁻⁹ Currently RLS is
7 controlled by a wide range of fungicides comprising Succinate Dehydrogenase
8 Inhibitors (SDHIs), Demethylation Inhibitors (DMIs) and a multisite inhibitor
9 chlorothalonil.¹ Declines in field efficacy to both SDHIs and DMIs have been
10 detected for several plant pathogens.¹⁰⁻¹³ Equally concerning are new
11 directives introduced by the European Commission on pesticide registration
12 (Regulation (EC) No 1107/2009). These directives may restrict future use of
13 some of the DMIs and chlorothalonil¹⁴, leading to increasing concern about the
14 provision of effective plant protection in the near future.

15 SDHIs are rapidly becoming one of the most important fungicide groups
16 for plant protection with resistance to other fungicide classes reported in many
17 crop pathogens.¹⁵ They were initially introduced in 1966 as two active
18 ingredients carboxin and oxycarboxin that showed a good spectrum of activity
19 against a range of basidiomycete pathogenic fungi.¹⁶ Modern SDHIs are
20 broad-spectrum products, with 19 different active ingredients available, used
21 both as foliar applications and seed treatments (FRAC MOA Poster 2016
22 (www.frac.info)). The current generation of SDHIs was introduced in 2005 for
23 use on cereals in the UK (CRD (<https://secure.pesticides.gov.uk/pestreg/>)) and
24 are now a mainstay in disease control programmes. In [the](#) 2014 growing

1 season 77% of winter barley and 40% of spring barley received SDHI
2 treatment (summing up all reported actives), and the use of some ingredients
3 such as bixafen increased by 94%, fluopyram by 382% and fluxapyroxad by
4 173% as compared to [the](#) 2012 growing season (all crops surveyed).¹⁷ The
5 extensive use of SDHIs in plant protection combined with the availability of
6 products containing individual SDHI active ingredients has raised concerns
7 over the evolution of pathogen resistance to SDHIs. Straight SDHI products do
8 not provide anti-resistance strategy 'in the can' and whether cereals growers
9 obey the label guidelines on their proper use, using effective mixing partners
10 at the proper dosage remains uncertain.

11 SDHIs are inhibitors of the mitochondrial respiratory complex II
12 (succinate dehydrogenase, Sdh, EC 1.3.5.1). The target protein of SDHI
13 fungicides, Sdh, consists of four subunits, labelled A-D and it is responsible for
14 oxidising succinate to fumarate and reducing ubiquinone to ubiquinol in the
15 mitochondrial electron transport chain and citric acid cycle.¹⁸⁻²⁴ SDHIs inhibit
16 fungal respiration by blocking the ubiquinone binding site, which is formed by
17 residues of subunits B (SdhB), C (SdhC) and D (SdhD).^{19,22,23,25,26} Sdh subunit
18 A (SdhA) is not involved in forming the ubiquinone binding pocket, and no
19 resistance mutations in this subunit have been described.^{15,19} Single amino
20 acid substitutions in SdhB, SdhC and SdhD have been shown to confer
21 resistance to SDHI fungicides. [Replacement of the highly conserved histidine](#)
22 [residue in the third cysteine rich cluster \[3Fe-4S\] of SdhB has been linked with](#)
23 [reduced sensitivity in lab mutants of *Zymoseptoria tritici* \(B: H267Y/L/F/N/Q\),²⁷⁻](#)
24 [lab and field isolates of *Botrytis cinerea* \(B: H272Y/R/L\),^{13,30-32} and field](#)

isolates of *Alternaria alternata* (B: H277Y/R).³³ Field resistance to SDHIs had not been commonly detected in cereal pathogens, with the exception of loss of sensitivity mutants to carboxin seed treatment reported for *Ustilago nuda*³⁴ but recently examples of mutations conferring reduced sensitivity in isolates of *Z. tritici*¹¹ and *Pyrenophora teres*^{Rehfus et al. 2016} pathogens of wheat and barley, respectively, have been reported. Moreover in 2015 the first isolates of *Rcc* showing strong decrease in sensitivity to SDHIs in bioassays, carrying a point mutation in the *SdhC* gene C: H142R and C: H149R were detected in Germany.— An additional mutation C: N83S, conferring a low resistance factor *in vitro*, was reported in single isolates originating from Germany, Ireland and Slovenia (FRAC 2015 SDHI Working Group (www.frac.info)). This brings a concern about SDHIs field performance in the coming years and the long-term effective protection against RLS.

Given that the first isolates with decreased sensitivity to SDHIs in lab assays have evolved recently in *Rcc* recently (FRAC 2015 SDHI Working Group (www.frac.info)), it is important to obtain the baseline data to which subsequent testing could-can be compared to-and investigate-to investigate possible consequences that resistant population could have on SDHI's field performance. This study reported-reports the current level of sensitivity to SDHI fungicides in *Rcc* in the UK and explored-explores the molecular basis of SDHI resistance in UV-induced mutants. Possible mutations in the target *Sdh* gene related to the resistance phenotype were examined at the nucleotide and protein level and fitness tests were conducted to see whether resistance mutations conferred any fitness penalty.

2 Experimental Methods

2.1 *In vitro* sensitivity testing of SDHI-resistant UV mutants and field isolates

In total 62 isolates sampled from barley in the UK in 2010 (n =7), 2011 (n =18) and 2012 (n =37) were tested in fungicide inhibition assays. Samples collected in 2010 originated from untreated plots in spring barley fungicide performance trial at Bush Estate, Scotland. Samples in 2011 originated from both untreated and treated plots of spring barley fungicide performance trial at Bush Estate, commercial fields in West Sussex, England and random plots in field trials at Lanark, Scotland. In 2012 samples were collected from a spring barley pathology SDHI *Ramularia* trial, including both untreated and treated plots. Single spore cultures of *Rcc* were isolated from leaves using a slight modification of the method described by Frei,³⁵ excluding leaf incubation prior to conidia isolation and using a fine sterile needle instead of a sterile blade. All of the *Rcc* isolates were maintained on potato dextrose agar (PDA, Oxoid, Basingstoke, UK) media amended with streptomycin 5 µg ml⁻¹ and/or kanamycin 50 µg ml⁻¹, in a growth cabinet (Sanyo Incubator, MIR-254, Osaka, Japan), in the dark, at 15°C.

Fungal cultures for inhibition assay were cultivated in alkyl ester (AE) broth²⁷ in 250 ml Erlenmeyer flask containing 150 ml of media. Each flask was inoculated with 150 µl of homogenised mycelium and cultured for 10-12 days in the dark at 16°C with shaking at 120 rpm. Subsequently 5 ml of each culture was homogenised for two minutes at 24000 rpm using an Ultra-Turrax T25 basic homogenizer (IKA®-Werke, GmbH&Co.KG, Staufen, Germany) with reusable plastic blades (T25 S18D, IKA®-Werke). The suspension was

1 vortexed for an additional minute and filtered through sterile nylon filters with
2 a pore size of 100 μm (Millipore, Darmstadt, Germany). Five SDHI fungicides:
3 isopyrazam, bixafen, boscalid, fluopyram and carboxin (Sigma-Aldrich, Saint
4 Louis, USA) were used in the assay. Each test was performed in a 96 well
5 plate, with three replicates per isolate. To each well 100 μl of mycelial
6 suspension and 100 μl of media containing fungicide at a range of
7 concentrations were added. The final concentration of fungal fragments in the
8 assay was 2.5×10^3 pieces of mycelium ml^{-1} . Final concentrations of
9 isopyrazam, bixafen, boscalid and fluopyram for field isolates were 10, 5, 1,
10 0.5, 0.1, 0.05, 0.01, 0 mg litre^{-1} and for carboxin were 50, 10, 5, 1, 0.5, 0.1,
11 0.05, 0 mg litre^{-1} . For SDHI-resistant mutants the same range of concentrations
12 plus one additional higher concentration of each fungicide was used. This
13 additional concentration was 50 mg litre^{-1} for isopyrazam, bixafen, boscalid,
14 fluopyram and 100 mg litre^{-1} for carboxin. All the mycelium and fungicide
15 dilutions were made in AE broth. Fungicide stocks were prepared in DMSO.
16 The final highest concentration of DMSO in wells was equal to 1% v/v when
17 the highest concentration of carboxin was used (50 mg litre^{-1}) and 0.2% v/v for
18 the highest concentration of the remaining four SDHI fungicides (10 mg litre^{-1}).
19 Plates were incubated in the dark for seven days at 16°C shaking at 120 rpm
20 (Gallenkamp, cooled orbital incubator, Weiss Technik Königswinter,
21 Germany). OD₄₀₀ measurements, with 20 flashes per well, were taken at day
22 zero and day seven on a spectrophotometer FLUOstar Omega (BMG Labtech,
23 Offenburg, Germany). Data were analysed using MARS Data Analysis

1 Software (BMG Labtech). EC₅₀ values were calculated from the 4-parameter
2 fit of the standard curve.

3 Resistance factors (RFs) were calculated as: $RF = (EC_{50} \text{ value of}$
4 $\text{mutant}) / (EC_{50} \text{ of parental isolate})$. The classification of resistance levels was
5 based on Leroux *et al.*¹³ However it was calibrated separately for each
6 fungicide using the RFs for the least sensitive isolates from the UK field
7 population. Resistance factors <0.5 were considered as hypersensitive.
8 Normal sensitivity for isopyrazam and bixafen was in the range $\geq 0.5 < 2$,
9 $\geq 0.5 < 5.5$ for boscalid, $\geq 0.5 < 7.5$ for fluopyram and $\geq 0.5 < 3.0$ for carboxin. Weak
10 resistance for isopyrazam and bixafen was classified as $\geq 2 < 10$, for boscalid
11 $\geq 5.5 < 10$, for fluopyram $\geq 7.5 < 10$ and for carboxin $\geq 3 < 10$. Resistance factors
12 $\geq 10 < 100$ were considered as moderate resistance and ≥ 100 as high
13 resistance.

14 2.2 Generation of SDHI-resistant UV mutants of *Rcc*

15 Mutants were developed using fungal mycelium fragments because we
16 failed to reliably generate *Rcc* spores *in vitro*.^{36,37} *Rcc* isolate DK05Rcc001ss2
17 (DK05) was used as the parental isolate for UV mutant generation. It was
18 isolated in Denmark in 2005 prior to the commercial launch of SDHIs (Lise
19 Nistrup Jorgensen pers. com.), from spring barley variety Braemar and was
20 sensitive to Qols. The genome and transcriptome of this particular isolate were
21 have been sequenced^{McGrann et al. 2016} (genome browser:
22 <http://ramularia.org/jbrowse>). Isopyrazam was chosen as the selection agent
23 to isolate resistant mutants. To define the minimum inhibitory concentration
24 (MIC), isopyrazam in concentrations ranging between 0.0001 and 20 mg litre⁻¹

1 ¹ was added to AE plates.²⁷ Each plate was then inoculated with 1.5×10^4
2 mycelial fragments and cultivated in a phytotron in the dark at 15°C for 18
3 days. The MIC of isoprazam was the lowest concentration for which growth
4 of wild type isolate was not observed after 18 days.

5 Selection for SDHI resistance was performed in AE agar amended with
6 0.05 mg litre⁻¹ (MIC) and 0.1 mg litre⁻¹ (2x MIC) of isopyrazam. Isolate DK05
7 was cultivated in AE broth at 16°C in the dark whilst shaking at 120 rpm for
8 seven days. The culture was homogenised, filtered and adjusted to a final
9 concentration of 10^5 pieces of mycelium ml⁻¹. Pieces of mycelium were
10 counted in Improved Neubauer C-Chip Disposable haemocytometers (Digital
11 Bio, Seoul, Korea) under the compound microscope using a 40x objective
12 (Leica, PL Fluotar 40x/0.70). Each isopyrazam amended agar plate was
13 inoculated with 1.5×10^4 of mycelial fragments and exposed to UV energy
14 between 12000 and 23000 $\mu\text{J cm}^{-2}$ in an UV Stratalinker 2400 (Stratagen, San
15 Diego, USA) leading to between approximately 20% and 50% survival.
16 Immediately after UV treatment, Petri dishes were sealed with parafilm and
17 transferred to the dark to avoid the activation of DNA repair systems in the
18 treated mycelial fragments. Samples were incubated for at least 18 days in the
19 dark at 15°C and; any colonies growing on agar after this period were collected
20 (between 22-33 days after UV treatment).

21 2.3 Characterisation of the *Sdh* gene

22 Prior to DNA extraction, fungal material was freeze dried overnight and
23 tissue lysed (Tissue Lyser LT, Qiagen, Hilden, Germany). DNA extraction was
24 performed using an Illustra Nucleon PhytoPure Genomic DNA Extraction Kit

1 according to the manufacturer's guidelines (GE Healthcare Life Sciences, Little
2 Chalfont, UK). Primers for amplifying subunits B, C and D of *Rcc*
3 (SdhC/D_Rcc_Final, ~~Table 4~~Table 4) were designed using data from the *Rcc*
4 genome sequence McGrann et al. 2016 (genome browser:
5 <http://ramularia.org/ibrowse>). Full *SdhB*, *SdhC* and *SdhD* sequences can be
6 found in GenBank database under accession numbers: KU758973,
7 KU758974, KU758975 respectively. PCR reactions were performed in the
8 following thermocycler: GeneAmp® PCR System 9700 (Applied Biosystems,
9 Foster City, USA). The PCRs were carried out using Go Taq® Green Master
10 Mix (Promega, Madison, USA). The PCR mix comprised 1x Master Mix, 0.2
11 µM of each primer, 6.25 ng of DNA and sterile distilled water (SDW) up to final
12 volume of 25 µl. To confirm the position of mutations, additional amplification
13 of the final *Sdh* subunits was performed using the FastStart High Fidelity PCR
14 System (Roche, Mannheim, Germany), containing proofreading polymerase.
15 The PCR mix comprised 1x buffer, 0.4 µM of each primer, 200 µM of each
16 dNTPs, 2.5 mM of MgCl₂, 2.5 U per reaction of an enzyme blend, 25 ng of
17 DNA and SDW up to 50 µl. Thermocycler conditions included an initial
18 denaturation at 95°C for two minutes, followed by 30 cycles of denaturation at
19 95°C for 15 seconds, annealing at 58°C for 30 seconds, extension at 72°C for
20 one minute and a final extension at 72°C for ten minutes. After sequencing, all
21 of the DNA fragments were analysed using Sequence Scanner Software v1.0
22 (Applied Biosystems).

Formatted: Font: (Default) Arial

2.4 Fitness tests on SDHI-resistant UV mutants

To verify the stability of mutations, mutants were sub-cultured six times on AE agar, not amended with SDHI fungicide and antibiotics. Plates were incubated in the dark at 15°C for seven to ten days between subculture steps. In addition the stability of mutants retrieved from long term storage in 0.25% v/v PDB was verified. Growth of mutants was verified *in vitro* on agar plates in the dark at 15°C. Cultures of the parental isolate DK05 and mutants were cultivated on AE agar without antibiotics and fungicides for three to four weeks. From each isolate an 8mm plug was excised and transferred into the center of a fresh AE agar plate. Five replicates for each culture were prepared. The growth of a colony was measured using an electronic digital caliper after two and four weeks, in four directions, excluding the mycelium plug.

For the detached leaf assay, barley plants of cultivar Optic were cultivated in pots in a Micro Clima Plant Growth Chamber MC1000E (Snijders Scientific, Tilburg, Netherlands) for up to four weeks under the following conditions: 16 hours light at 20°C, 80% humidity (day) and 8 hours dark at 16°C, 90% humidity (night). A detached leaf assay was performed using a modified method described by Thirugnanasambandam *et al.*³⁶ after Newton *et al.*³⁸ F-1 and F-2 leaves were used in the experiment. The assay was divided into two parts: untreated control and leaves sprayed with 1 mg litre⁻¹ of isopyrazam. Fungicide solutions were prepared in AE broth and, control material was sprayed with AE broth not containing fungicide. Sections around 4 cm long were cut and placed with the abaxial part downwards on 0.5% (w/v) water agar (Oxoid), amended with 1 mM benzimidazole (Sigma-Aldrich). Up to

1 six leaves were placed into push fit polystyrene boxes of dimensions 79x47x22
2 mm (Steward-Solutions, Croydon, UK).

3 *Rcc* inoculum was prepared from two week old AE agar plates, cultured
4 at 15°C in the dark. SDW (0.5-1 ml) was added to the plate and mycelium was
5 scraped from the colony surface and centrifuged for three minutes at 4000
6 rpm. It was washed three times with SDW and finally diluted in 1.5 ml of SDW.
7 Each leaf was inoculated in two places with 10 µl of mycelial suspension as
8 described by Thirugnanasambandam *et al.*³⁶ Both of the drops were inoculated
9 on the same, adaxial part of the leaf, approximately 1.5-2 cm apart, avoiding
10 the midrib. Boxes were incubated in the phytotron under 12 hours dimmed light
11 and 12 hours dark at 15°C, in high humidity conditions to promote fungal
12 growth. Fungal hyphae were stained dark blue with Aniline Blue:ethanol (50:50
13 v/v) and the infection process was observed under the compound microscope
14 (DM RBE Research Microscope, Leica, Wetzlar, Germany) using 10x, 20x and
15 40x objectives (PL Fluotar 10x/0.30, 20x/0.50, 40x/0.70) up to 29 days post
16 inoculation (dpi). Images were acquired using a CMEX DC.5000 5Mpix camera
17 (Euromex, Arnhem, Netherlands) and edited using ImageJ³⁹ and Adobe
18 Photoshop® CS5 (Adobe Systems, San Jose, USA) softwares. The
19 experiment was repeated twice.

20 **2.5 Statistical analysis**

21 Statistical analysis was performed in Minitab v16 (Minitab Inc., State
22 College, USA). One way ANOVA was used to examine differences between
23 the group means in *in vitro* fungicide sensitivity assay. If significant differences
24 between group means were indicated, Tukey's pairwise comparisons were

1 conducted. Pearson product moment correlation coefficients (r) were used to
2 verify the cross resistance patterns between SDHI fungicides. A correlation of
3 ≤ 0.35 was categorised as weak, 0.36 to 0.67 as moderate, 0.68 to 1.00 as
4 strong, with correlation coefficients ≥ 0.90 described as very strong.⁴⁰
5

3 Results

3.1 Baseline sensitivity of *Rcc* populations to SDHIs

A fungicide inhibition assay was used to screen 62 UK isolates collected in 2010, 2011 and 2012, for sensitivity to five SDHI fungicides: isopyrazam, bixafen, boscalid, fluopyram and carboxin (Table 1). Isopyrazam and bixafen most effectively inhibited *Rcc* growth, with mean EC₅₀ values of 0.019 mg litre⁻¹ and 0.015 mg litre⁻¹ respectively. Boscalid (EC₅₀ = 0.137 mg litre⁻¹) and fluopyram (EC₅₀ = 0.151 mg litre⁻¹) also showed good control of the pathogen *in vitro*. Carboxin (EC₅₀ = 1.120 mg litre⁻¹) was the least effective fungicide. There were no significant differences between years in sensitivity of *Rcc* populations (Table 1) to isopyrazam ($P = 0.216$), bixafen ($P = 0.216$), boscalid ($P = 0.262$), fluopyram ($P = 0.110$) or carboxin ($P = 0.079$).

3.2 Identification of the target mutations conferring resistance to SDHIs

Twenty two *Rcc* colonies with putative resistance to SDHIs were isolated after UV mutagenesis (Table S 1). In total 112.5 mycelial fragments were plated out, resulting in overall mutation frequency (collected colonies/total no of plated mycelial fragments) of 2×10^{-5} . These included five isolates (designated Mut1, Mut2, Mut7, Mut8 and Mut11) showing a notable decrease in sensitivity to SDHI fungicides and 17 false positives. False positives were initially isolated from agar plates after UV mutagenesis, however in further testing did not show a decrease in sensitivity to SDHIs *in vitro* and as a consequence were eliminated from the analysis. UV treatments to create verified SDHI-resistant mutants used energy inputs of 18000 $\mu\text{J cm}^{-2}$ and 22000 $\mu\text{J cm}^{-2}$, corresponding to around 20% of colony survival (Figure S 1,

Formatted: Font: (Default) Arial

1 Figure S 2). Mutants were successfully selected both using the MIC (four
2 isolates) and 2x MIC (one isolate) of isopyrazam.

3 Sequencing of the *Rcc* Sdh subunit genes revealed that in Mut2 there
4 were neither nucleotide nor amino acid changes in genes *SdhB*, *SdhC* or
5 *SdhD*. However the four remaining isolates each possessed a single
6 nucleotide mutation positioned either in locus *SdhB* (Figure 1, Mut1, Mut11) or
7 locus *SdhC* (Figure 1, Mut7, Mut8). The single nucleotide mutation in *SdhB* of
8 isolate Mut1 conferred an amino acid change from serine (tca) to leucine (tta)
9 at position 217 (B: S217L). In isolate Mut11 the single nucleotide mutation in
10 *SdhB* conferred a change from asparagine (aac) to isoleucine (atc) at position
11 224 (B: N224I). Both of these mutations are positioned in a region of subunit
12 B that is conserved across the species (Table 2, Figure 1). Mutation of a two
13 distinct nucleotides in the same codon in *SdhC* was observed in Mut7 and
14 Mut8. In the case of Mut7 the wild type histidine (cat) residue at position 142
15 was substituted with arginine (cgt) (C: H142R), while in the case of isolate Mut8
16 it was substituted with glutamine (caa) (C: H142Q). As was the case above,
17 this particular residue of histidine in *SdhC* at position 142 (C: H142) is highly
18 conserved across the species (Table 2, Figure 1).

19 3.3 Assessment of SDHI resistance associated with mutations

20 Mut7 (C: H142R) was highly resistant to boscalid fungicide (resistance
21 factor (RF) =1114), and moderately resistant to four other SDHI active
22 ingredients, with high RFs for bixafen (RF =55.31), isopyrazam (RF =44.10),
23 carboxin (RF =32.80) and fluopyram (RF =16.77), compared to the parental
24 isolate DK05 (RF =1) and the least sensitive isolate from the UK field

1 population (Table 3). Mut8 (C: H142Q) similarly showed the same high level
2 of resistance to boscalid (RF =1114) and moderate resistance to carboxin (RF
3 =19.19), bixafen (RF =10.69) and fluopyram (RF =15.91). However we
4 observed no differences in sensitivity to isopyrazam (RF =0.688) as compared
5 to the parental isolate and the least sensitive isolate from the UK field
6 population (Table 3).

7 In the case of Mut11 (B: N224I) a moderate level of resistance was
8 observed to most of the tested SDHI fungicides (RF =37.28 for boscalid, RF
9 =12.65 for bixafen, RF =21.82 for fluopyram and RF =13.23 for carboxin), with
10 the exception of isopyrazam to which weak resistance was found (RF =6.758).

11 In contrast, for Mut1 (B: S217L) a moderate resistance was shown only in the
12 case of fluopyram (RF =49.90). For isopyrazam (RF =9.239), boscalid (RF
13 =8.999) and bixafen (RF =2.246) only weak resistance was detected and for
14 carboxin (RF =1.481) there were no changes in sensitivity as compared to the
15 parental isolate and the UK field population (Table 3).

16 Mut2, which had no detectable mutations in SdhB, C or D, showed
17 moderate level of resistance to isopyrazam (RF =31.55) and bixafen (RF
18 =24.51). For the remaining three active ingredients no differences in sensitivity
19 were observed compared to the DK05 and the UK field population (boscalid
20 RF =1.532, fluopyram RF =1.177, carboxin RF =2.752), (Table 3).

21 We observed a very strong cross resistance ($r = 0.901$, $P = 0.037$) only
22 between isopyrazam and bixafen (Figure S 3). For the remainder of the
23 fungicides the correlations in sensitivity to different fungicides were not
24 significant ($P > 0.05$).

3.4 Fitness tests on SDHI-resistant mutants of *Rcc*

1.1.1 Culture characteristics

There were no morphological differences between the mutants and DK05 when grown on agar plates (data not shown). There was no difference in the AE broth liquid culture growth phenotype of the Mut1, 7, 8 and 11 compared to the wild type isolates. However Mut2 liquid cultures had a much darker colouration than wild type (Figure S 4). At this time it is unclear if other uncharacterised mutations are responsible for this aberrant phenotype in Mut2.

1.1.2 Stability of mutations

Most of the mutations correlating with resistance to SDHIs in *Rcc* were stable, with the exception of Mut8 (C: H142Q) in which the target mutation was not detected after the subculturing process. In long term storage, we noted a mixture of resistant and wild type alleles during the sequencing.

1.1.3 *In vitro* plate growth assay

Significant differences in growth on agar plates between DK05, Mut1, Mut7, Mut8 and Mut11 were indicated at both time points, after 14 days ($P < 0.001$) and 28 days ($P < 0.001$), (Figure 2). After 14 days we observed significantly faster growth than for the wild type for Mut7 (C: H142R) and Mut11 (B: N224I), ($P < 0.05$); after 28 days faster growth was observed only for Mut11 (B: N224I), ($P < 0.05$), (Figure 2). The test was performed separately for Mut2 because it failed to grow in the first experiment. In the case of Mut2 we detected no significant differences in growth on agar plates compared to the wild type after 14 days ($P = 0.532$) and 28 days ($P = 0.916$), (Figure 3).

1 1.1.4 *In planta* leaf assay

2 The infection process was examined for two mutants, Mut11 (B: N224I)
3 and Mut7 (C: H142R) and compared to the development of parental isolate
4 DK05. Colonisation of untreated barley leaves by both *Rcc* mutant isolates and
5 isolate DK05 occurred in a very similar manner, typical of the infection process
6 described previously.³⁶ *Rcc* infection of barley began with formation of an
7 extensive hyphal network on the leaf surface (Figure 4a) and entry through
8 stomatal pores. Sporulation occurred from 8 dpi onwards from distinctive swan
9 neck conidiophores on top of which spores developed (Figure 4b). Disease
10 symptoms, initial pepper-like spots expanding to small, brown to blackish
11 necrotic lesions, on the untreated leaf segments were observed
12 macroscopically from 25 dpi for all three isolates used in this study (data not
13 shown). This showed that Mut11 and Mut7 could infect the host plant barley,
14 reproduce successfully by generating spores and complete their life cycle,
15 further suggesting that there was no measurable fitness penalty associated
16 with the target mutations conferring resistance to SDHIs.

17 Infection by isolate Mut11 and Mut7 was not affected by foliar isopyrazam
18 application at a concentration of 1 mg litre⁻¹. Both mutants were able to form
19 an epiphytic hyphal network on the leaf surface and penetrate multiple stomata
20 (Figure 4d). Conidiophores developed, resulting in abundant sporulation, both
21 in stomata and on the leaf surface (Figure 4e). The progressing infection of
22 both mutants led to red discolouration of the guard and surrounding epidermal
23 cells (Figure 4g), followed by rapid development of the typical RLS symptoms
24 from 28 dpi by both Mut7 (Figure 4h) and M11 (Figure 4i). In contrast the

1 growth of the parental isolate DK05, was clearly inhibited after treatment with
2 isopyrazam (Figure 4f). Some hyphae attempted to colonise the leaf surface.
3 However there was a lack of highly branched and controlled epiphytic growth
4 of hyphae, no subsequent infection of stomata, and no disease development
5 was observed (results not shown).
6

4 Discussion and conclusions

This study ~~has shown that presents the baseline sensitivity of *Rcc* populations to SDHIs~~ in the UK ~~presently remain sensitive to SDHIs. This is, which is~~ of high relevance given that ~~in 2015 the~~ first field isolates with decreased sensitivity to SDHIs in lab assays ~~have been were~~ reported in some European countries ~~in 2015~~ (FRAC 2015 SDHI Working Group (www.frac.info)). Furthermore using UV induced mutants we ~~have~~ characterised mutations correlating with resistance to SDHIs in *Rcc* and assessed some of ~~its their~~ fitness parameters, giving an insight into ~~the~~ possible behaviour of ~~a~~ resistant population.

Two mutants Mut7 (C: H142R) and Mut8 (C: H142Q) generated by UV mutagenesis carry a mutation of the highly conserved histidine residue in SdhC and are highly resistant to boscalid. The exact same mutation as in the case of Mut7 (C: H142R) was reported in *Rcc* isolates showing strong decrease in sensitivity to SDHIs in *in vitro* assays in Germany (FRAC 2015 SDHI Working Group (www.frac.info)). Additionally the replacement of this particular amino acid in SdhC has previously been linked with resistance to SDHIs in both *A. alternata* (C: H134R)⁴¹ and *P. teres* (C: H134R)^{Reffhus et al. 2016} field isolates, as well as in a laboratory mutant of *Z. tritici* (C: H145R).²⁸ Histidine residue C: H142 was not predicted to be involved directly in ubiquinone binding and reduction in the *Z. tritici* Sdh protein model.^{28,29} However, this histidine residue has been shown to ligate with heme b and its polar propionate side chains form an integral part of the ubiquinone binding pocket, explaining the loss of sensitivity to SDHIs in mutants carrying variant

1 C: H145R.²⁸ In the light of this evidence we suggest that histidine C: H142 acts
2 as [a](#) ligand for heme b in *Rcc*, and this explains the reduced sensitivity to
3 inhibitors of mitochondrial respiratory complex II in mutant *Rcc* isolates
4 carrying variant C: H142R/Q.

5 Mutation of the serine residue found in Mut1 (B: S217L) has so far only
6 been correlated with resistance to SDHIs in a laboratory mutant of *Z. tritici* (B:
7 S218F)²⁸ and has not to date been found for any field resistant pathogen. The
8 mutation we detected in *Rcc* Mut11 (B: N224I) has been reported to confer
9 SDHI resistance in both artificially induced mutants and in naturally occurring
10 fungal isolates. The *Z. tritici* laboratory mutant carrying the corresponding
11 asparagine mutation (B: N225I) exhibits reduced SDHI sensitivity,²⁸ while *B.*
12 *cinerea* laboratory mutants⁴⁴ and field isolates^{13,30} carrying the equivalent
13 mutation (B: N230I) are resistant to SDHIs. Substitution of the same
14 asparagine residue, this time by threonine, was described recently for SDHI
15 resistant field isolates of *Z. tritici* (B: N225T), (FRAC 2014 SDHI Working
16 Group (www.frac.info)). Although these particular amino acid positions in SdhB
17 in the *Z. tritici* Sdh model were not predicted to be involved in forming the
18 ubiquinone binding pocket, they were positioned in the vicinity of key residues
19 involved in SDHI binding.^{28,29} Both of the mutations in Mut1 and Mut11 could
20 have a long-distance effect on the architecture of the ubiquinone binding
21 pocket, which then explains the sensitivity loss they cause towards inhibitors
22 of mitochondrial respiratory complex II as proposed by Scalliet *et al.*²⁸ None of
23 the substitutions corresponded to a replacement of a conserved histidine
24 residue (B: H266 in *Rcc*) in a third cysteine rich cluster [3Fe-4S] of SdhB, found

1 to be responsible for resistant development to SDHIs in lab mutants of *Z. tritici*
2 (B: H267Y/L/F/N/Q).²⁷⁻²⁹ However, given the limited number of mutants
3 generated in this study, it cannot be excluded that such mutations could
4 develop and contribute to SDHIs efficacy loss in the field.

5 In the case of Mut2 we observed no amino acid changes in the target
6 Sdh enzyme which could be linked with the resistance to some of the SDHIs,
7 isopyrazam and bixafen. Although alterations of the target gene are the most
8 common mechanism responsible for sensitivity loss towards SDHIs among
9 plant pathogenic fungi,^{15,19} they are only one of the possible known
10 mechanisms conferring resistance to such fungicides. SDHI-resistant isolates
11 of *Corynespora cassiicola*,⁴⁸ *A. alternata*⁴⁹ and *B. cinerea*¹³ have been
12 reported with no sequence mutation of the Sdh subunit genes. For the At
13 present moment there is no evidence as to whether which of these mechanisms could
14 be responsible for the resistance patterns in Mut2 and further work is needed
15 to investigate this phenomenon. However given that Mut2 was only resistant
16 to some of the SDHIs tested, overexpression of the target gene or its multiple
17 copies seem to be the two most likely possibilities.

18 We noted positive cross resistance profiles in this study only between
19 isopyrazam and bixafen; for the remaining SDHI active ingredients tested we
20 observed a lack of cross resistance. Although FRAC classifies inhibitors of
21 mitochondrial respiratory complex II as cross resistant¹⁵, recent studies have
22 demonstrated a lack of cross resistance between newer SDHIs.^{19,28,29,44,50,51}
23 In *Rcc* SDHI-resistant mutants the resistance profiles varied notably between
24 the mutated isolates and were strongly associated with the particular position

1 of amino acids. For example Mut7 (C: H142R) was highly resistant to boscalid
2 but moderately resistant to the other four tested SDHIs. Mut11 (B: N224I) was
3 moderately resistant to all SDHIs tested, with the exception of isopyrazam to
4 which weak resistance was observed. This suggests that different mutations
5 could differently influence the affinity of SDHIs to the target site, explaining the
6 limited positive cross resistance among mutated isolates. Additionally it cannot
7 be ruled out that additional mutations, outside the *Sdh* gene, incurred-occurred
8 as a consequence of UV mutagenesis studies, and have an impact on the
9 sensitivity profiles of *Rcc* UV mutants.

10 No measurable fitness penalty associated with resistance to SDHIs was
11 observed in terms of radial colony growth on agar plates in any of the mutated
12 isolates. Additionally *in planta* assays performed for two isolates, Mut7 (C:
13 H142R) and Mut11 (B: N224I), indicated that both of the mutants were able to
14 colonise the leaf and effectively reproduce in untreated barley leaves as well
15 as barley leaves treated with isopyrazam. These results are consistent with
16 previous studies on SDHI-resistant mutants of *Z. tritici*, which were able to
17 colonise the leaf, cause symptoms and produce spores, despite the impaired
18 enzyme activity due to mutation.^{28,29} Although the concentration of isopyrazam
19 used in this study of 1mg litre⁻¹ may not give a good measure of fitness in the
20 presence of commercial rates of fungicide application, it provides a good
21 estimation of fitness in the presence of rates sufficient to eliminate non-
22 resistant genotypes. Thus the extrapolation of this data to field conditions
23 should be taken with caution. More fitness tests on a wider variety of traits,
24 especially on recently emerged reduced sensitivity field isolates of *Rcc*, should

Commented [MP1]: I had to change this section. Assuming that Mut8 might have been picked up as mixed colony our previous discussion in that point was irrelevant and too speculative

1 be performed to fully understand the behaviour of resistant population and their
2 influence on SDHIs field performance. Additionally it will be important to verify
3 if the recorded field mutations reappear in the following seasons, and if ~~yes-so~~
4 in what frequency. In this study most of the mutations were stable in the
5 absence of fungicide, except Mut8 (C: H142Q). Mut8 was either undergoing
6 the process of reversion to the wild type haplotype at the *SdhC* gene or was
7 originally picked up as a mixed colony of a wild type and a mutant which could
8 explain differences in its sensitivity profiles as compared to Mut7 (C: H142R).
9 This suggests that mutations responsible for SDHI resistance in *Rcc* ~~could~~ can
10 be stable, a possibility which needs further verification.

11 This study presents baseline sensitivity of *Rcc* populations to SDHIs in
12 the UK and ~~analysis-analyses~~ the possible behaviour of a resistant population
13 using UV mutants. Further UV mutagenesis studies on *Rcc* population are
14 required to verify the possibility of ~~the~~ other mutations occurring. These should
15 then be incorporated into a molecular monitoring assay, together with any
16 mutations occurring in the field conditions, allowing detection of any amino acid
17 changes associated with resistance at the field level. At the same time it is
18 important to carry on *in vitro* sensitivity testing in order to detect any possible
19 non-target site mechanisms of resistance, which could have been missed ~~due~~
20 to reliance f-relying only on molecular screening tests alone. *Ramularia collo-*
21 *cygni* has been exposed to SDHI fungicides since 2005 in the UK and recent
22 population genetic studies have suggested that it has a high potential for
23 evolutionary adaptation.^{9,55,56} Based on the current evidence and the recent
24 reports of *Rcc* isolates with decreased sensitivity to SDHIs in some European

1 countries (FRAC 2015 SDHI Working Group (www.frac.info)), we conclude
2 that the risk of resistance development in *Rcc* to SDHIs in the UK is high and
3 robust anti-resistance strategies should be implemented in order to prolong
4 SDHIs' effective life span. The long term effective control of RLS in barley
5 requires integrated management systems, which cannot be based exclusively
6 on fungicide applications but should include a combination of chemical control
7 and resistant varieties, a strategy that needs to be revised and implemented
8 in a relatively short time.

5 Acknowledgements

We thank SRUC and Syngenta Ltd. for providing the funding for this research.

6 References

- 1 Havis ND, Brown JK, Clemente G, Frei P, Jedryczka M, Kaczmarek J, Kaczmarek M, Matusinsky P, McGrann GR, Pereyra S, Piotrowska MJ, Sghyer H, Tellier A and Hess M, *Ramularia collo-cygni*- an emerging pathogen of barley crops. *Phytopathology* **105**: 895-904 (2015).
- 2 Oxley SJP, Havis N, Evans A, Waterhouse S and Tonguc L, A guide to the recognition and understanding of *Ramularia* and other leaf spots of barley. BASF plc & SAC (2010).
- 3 Oxley SJP, Havis ND, Managing *Ramularia collo-cygni* through varietal resistance, seed health and forecasting. Project report No. 463. HGCA, Kenilworth, UK (2010).
- 4 Havis ND, Oxley SJP and Burnett FJ, Advances in control of *Ramularia collo-cygni*. In *Proceedings Crop Protection in Northern Britain 2012*, ed(s). Heilbronn TD, pp. 125-130. Page Bros, Norwich, UK (2012).
- 5 Matusinsky P, Hanusova M, Stemberkova L, Marik P, Minarikova V, Tvaruzek L, Langer I and Spitzer T, Response of spring barley cultivars to *Ramularia* leaf spot in conditions of the Czech Republic. *Cereal Res. Commun.* **41**: 126-132 (2013).
- 6 McGrann GRD, Stavrinides A, Russell J, Corbitt MM, Booth A, Chartrain L, Thomas WTB and Brown JKM, A trade off between mlo resistance to powdery mildew and increased susceptibility of barley to a newly important disease, *Ramularia* leaf spot. *J. Exp. Bot.* **65**: 1025-1037 (2014).
- 7 Fountaine JM, Fraaije BA, Development of Qol resistant alleles in populations of *Ramularia collo-cygni*. *Aspects of Applied Biology* **92**: 123-126 (2009).
- 8 Matusinsky P, Svobodova-Leisova L, Marik P, Tvaruzek L, Stemberkova L, Hanusova M, Minarikova V, Vysohlidova M and Spitzer T, Frequency of a mutant allele of cytochrome b conferring resistance to Qol fungicides in the Czech population of *Ramularia collo-cygni*. *J. Plant Dis. Prot.* **117**: 248-252 (2010).

- 1 9 Leisova-Svobodova L, Matusinsky P and Kucera L, Variability of the
2 *Ramularia collo-cygni* population in Central Europe. *J. Phytopathol.*
3 **160**: 701-709 (2012).
- 4 10 Cools HJ, Fraaije BA, Are azole fungicides losing ground against
5 Septoria wheat disease? Resistance mechanisms in *Mycosphaerella*
6 *graminicola*. *Pest Manage. Sci.* **64**: 681-684 (2008).
- 7 11 Dooley H, Shaw MW, Mehenni-Ciz J, Spink J and Kildea S, Detection
8 of *Zymoseptoria tritici* SDHI insensitive field isolates carrying the SdhC-
9 H152R and SdhD-R47W substitutions. *Pest Manage. Sci.* DOI:
10 10.1002/ ps.4269 (2016).
- 11 12 Latorre BA, Torres R, Prevalence of isolates of *Botrytis cinerea* resistant
12 to multiple fungicides in Chilean vineyards. *Crop Prot.* **40**: 49-52 (2012).
- 13 13 Leroux P, Gredt M, Leroch M and Walker AS, Exploring mechanisms of
14 resistance to respiratory inhibitors in field strains of *Botrytis cinerea*, the
15 causal agent of gray mold. *Appl. Environ. Microbiol.* **76**: 6615-6630
16 (2010).
- 17 14 Leadbeater A, The impact of the new European Regulations on the
18 management of crop diseases. In *Modern Fungicides and Antifungal*
19 *Compounds VI*, ed(s). Dehne HW, Deising HB, Gisi U, Kuck KH, Russell
20 PE, Lyr H, pp. 1-10. DPG, Braunschweig, Germany (2014).
- 21 15 Sierotzki H, Scalliet G, A review of current knowledge of resistance
22 aspects for the next-generation succinate dehydrogenase inhibitor
23 fungicides. *Phytopathology* **103**: 880-887 (2013).
- 24 16 von Schmeling B, Kulka M, Systemic fungicidal activity of 1,4-oxathiin
25 derivatives. *Science* **152**: 659-660 (1966).
- 26 17 Monie C, Reay G and Wardlaw J, Pesticide usage in Scotland- arable
27 crops and potato stores 2014. SASA, Edinburgh, UK (2014).
- 28
29 18 Hagerhall C, Succinate: quinone oxidoreductases. Variations on a
30 conserved theme. *Biochim. Biophys. Acta* **1320**: 107-141 (1997).
- 31 19 Avenot HF, Michailides TJ, Progress in understanding molecular
32 mechanisms and evolution of resistance to succinate dehydrogenase
33 inhibiting (SDHI) fungicides in phytopathogenic fungi. *Crop Prot.* **29**:
34 643-651 (2010).
- 35 20 Lancaster CRD, Succinate: quinone oxidoreductases: an overview.
36 *Biochim. Biophys. Acta*, **1553**: 1-6 (2002).
- 37 21 Ackrell BA, Progress in understanding structure-function relationships
38 in respiratory chain complex II. *FEBS Lett.* **469**: 1-5 (2000).

- 1 22 Yankovskaya V, Horsefield R, Tornroth S, Luna-Chavez C, Miyoshi H,
2 Leger C, Byrne B, Cecchini G and Iwata S, Architecture of succinate
3 dehydrogenase and reactive oxygen species generation. *Science* **299**:
4 700-704 (2003).
- 5 23 Horsefield R, Yankovskaya V, Sexton G, Whittingham W, Shiomi K,
6 Omura S, Byrne B, Cecchini G and Iwata S, Structural and
7 computational analysis of the quinone-binding site of complex II
8 (succinate-ubiquinone oxidoreductase). A mechanism of electron
9 transfer and proton conduction during ubiquinone reduction. *J. Biol.*
10 *Chem.* **281**: 7309-7316 (2006).
- 11 24 Ulrich JT, Mathre DE, Mode of action of oxathiin systemic fungicides. V.
12 Effect on electron transport system of *Ustilago maydis* and
13 *Saccharomyces cerevisiae*. *J. Bacteriol.* **110**: 628-632 (1972).
- 14 25 Huang LS, Sun G, Cobessi D, Wang AC, Shen JT, Tung EY, Anderson
15 VE and Berry EA, 3-Nitropropionic acid is a suicide inhibitor of
16 mitochondrial respiration that, upon oxidation by complex II, forms a
17 covalent adduct with a catalytic base arginine in the active site of the
18 enzyme. *J. Biol. Chem.* **281**: 5965-5972 (2006).
- 19 26 Tran QM, Rothery RA, Maklashina E, Cecchini G and Weiner JH, The
20 quinone binding site in *Escherichia coli* succinate dehydrogenase is
21 required for electron transfer to the heme b. *J. Biol. Chem.* **281**: 32310-
22 32317 (2006).
- 23 27 Skinner W, Bailey A, Renwick A, Keon J, Gurr S and Hargreaves J, A
24 single amino-acid substitution in the iron-sulphur protein subunit of
25 succinate dehydrogenase determines resistance to carboxin in
26 *Mycosphaerella graminicola*. *Curr. Genet.* **34**: 393-398 (1998).
- 27 28 Scalliet G, Bowler J, Luksch T, Kirchhofer-Allan L, Steinhauer D, Ward
28 K, Niklaus M, Verras A, Csukai M, Daina A and Fonne-Pfister R,
29 Mutagenesis and functional studies with succinate dehydrogenase
30 inhibitors in the wheat pathogen *Mycosphaerella graminicola*. *Plos One*
31 **7**: 1-20 (2012).
- 32 29 Fraaije BA, Bayon C, Atkins S, Cools HJ, Lucas JA and Fraaije MW,
33 Risk assessment studies on succinate dehydrogenase inhibitors, the
34 new weapons in the battle to control Septoria leaf blotch in wheat. *Mol.*
35 *Plant Pathol.* **13**: 263-275 (2012).
- 36 30 Veloukas T, Leroch M, Hahn M and Karaoglanidis GS, Detection and
37 molecular characterization of boscalid-resistant *Botrytis cinerea*
38 isolates from strawberry. *Plant Dis.* **95**: 1302-1307 (2011).

- 1 31 De Miccolis Angelini RM, Habib W, Rotolo C, Pollastro S and Faretra F,
2 Selection, characterization and genetic analysis of laboratory mutants
3 of *Botryotinia fuckeliana* (*Botrytis cinerea*) resistant to the fungicide
4 boscalid. *Eur. J. Plant Pathol.* **128**: 185-199 (2010).
- 5 32 Yin Y, Kim Y and Xiao C, Molecular characterization of boscalid
6 resistance in field isolates of *Botrytis cinerea* from apple.
7 *Phytopathology* **101**: 986-995 (2011).
- 8 33 Avenot HF, Sellam A, Karaoglanidis G and Michailides TJ,
9 Characterization of mutations in the iron-sulphur subunit of succinate
10 dehydrogenase correlating with boscalid resistance in *Alternaria*
11 *alternata* from California pistachio. *Phytopathology* **98**: 736-742 (2008).
- 12 34 Newcombe G, Thomas PL, Incidence of carboxin resistance in *Ustilago*
13 *nuda*. *Phytopathology* **81**: 247-250 (1991).
- 14 35 Frei P, *Ramularia collo-cygni*: cultivation, storage, and artificial infection
15 of barley and weed grasses under controlled conditions. In *Proceedings*
16 *of the Second International Workshop of Barley Leaf Blights- meeting*
17 *the challenges of barley blights*, ed(s). Yahyaoui AH, Brader L, Tekauz
18 A, Wallwork H, Steffenson B, pp. 351-354. ICARDA, Aleppo, Syria
19 (2004).
- 20 36 Thirugnanasambandam A, Wright K, Havis N, Whisson S and Newton
21 A, *Agrobacterium*-mediated transformation of the barley pathogen
22 *Ramularia collo-cygni* with fluorescent marker tags and live tissue
23 imaging of infection development. *Plant Pathol.* **60**: 929-937 (2011).
- 24 37 Sutton BC, Waller JM, Taxonomy of *Ophioclaadium hordei*, causing leaf
25 lesions on Triticale and other Graminaeae. *Trans. Br. Mycol. Soc.* **90**:
26 55-61 (1988).
- 27 38 Newton AC, Searle J, Guy DC, Hackett CA and Cooke DEL, Variability
28 in pathotype, aggressiveness, RAPD profile, and rDNA ITS1 sequences
29 of UK isolates of *Rhynchosporium secalis*. *J. Plant Dis. Prot.* **108**: 446-
30 458 (2001).
- 31 39 Schneider CA, Rasband WS and Eliceiri KW, NIH Image to ImageJ: 25
32 years of image analysis. *Nat. Methods* **9**: 671-675 (2012).
- 33 40 Taylor R, Interpretation of the correlation coefficient: a basic review.
34 *Journal of Diagnostic Medical Sonography* **6**: 35-39 (1990).
- 35 41 Avenot HF, Sellam A and Michailides TJ, Characterization of mutations
36 in the membrane-anchored subunits AaSDHC and AaSDHD of
37 succinate dehydrogenase from *Alternaria alternata* isolates conferring
38 field resistance to the fungicide boscalid. *Plant Pathol.* **58**: 1134-1143
39 (2009).

- 1 42 Vibat CRT, Cecchini G, Nakamura K, Kita K and Gennis RB,
2 Localization of histidine residues responsible for heme axial ligation in
3 cytochrome b (556) of complex II (succinate:ubiquinone
4 oxidoreductase) in *Escherichia coli*. *Biochemistry* **37**: 4148-4159
5 (1998).
- 6 43 Maklashina E, Rothery RA, Weiner JH and Cecchini G, Retention of
7 heme in axial ligand mutants of succinate-ubiquinone oxidoreductase
8 (complex II) from *Escherichia coli*. *J. Biol. Chem.* **276**: 18968-18976
9 (2001).
- 10 44 Lavee A, Gamet S, Walker AS, Debieu D, Toquin V and Fillinger S,
11 Site-directed mutagenesis of the P225, N230 and H272 residues of
12 succinate dehydrogenase subunit B from *Botrytis cinerea* highlights
13 different roles in enzyme activity and inhibitor binding. *Environ.*
14 *Microbiol.* **16**:2253-2266 (2014).
- 15 45 Brent KJ, Hollomon DW, in *Fungicide resistance in crop pathogens:*
16 *how can it be managed? FRAC Monograph No. 1. 2nd (revised) edition.*
17 CropLife International, Brussels, Belgium (2007).
- 18 46 Cools HJ, Hawkins N and Fraaije B, Constraints on the evolution of
19 azole resistance in plant pathogenic fungi. *Plant Pathol.* **62**: 36-42
20 (2013).
- 21 47 Miguez M, Reeve C, Wood PM and Hollomon DW, Alternative oxidase
22 reduces the sensitivity of *Mycosphaerella graminicola* to QoI fungicides.
23 *Pest Manage. Sci.* **60**: 3-7 (2004).
- 24 48 Miyamoto T, Ishii H, Stammler G, Koch A, Ogawara T, Tomita Y,
25 Fountaine J, Ushio S, Seko T and Kobori S, Distribution and molecular
26 characterization of *Corynespora cassicola* isolates resistant to
27 boscalid. *Plant Pathol.* **59**: 873-881 (2010).
- 28 49 Avenot HF, van den Biggelaar H, Morgan D, Moral J, Joosten M and
29 Michailides TJ, Sensitivities of baseline isolates and boscalid-resistant
30 mutants of *Alternaria alternata* from pistachio to fluopyram,
31 penthiopyrad, and fluxapyroxad. *Plant Dis.* **98**: 197-205 (2014).
- 32 50 Avenot HF, Thomas A, Gitaitis RD, Langston DB and Stevenson KL,
33 Molecular characterization of boscalid- and penthiopyrad-resistant
34 isolates of *Didymella bryoniae* and assessment of their sensitivity to
35 fluopyram. *Pest Manage. Sci.* **68**: 645-651 (2012).
- 36 51 Ishii H, Miyamoto T, Ushio S and Kakishima M, Lack of cross-resistance
37 to a novel succinate dehydrogenase inhibitor, fluopyram, in highly
38 boscalid-resistant isolates of *Corynespora cassicola* and *Podosphaera*
39 *xanthii*. *Pest Manage. Sci.* **67**: 474-482 (2011).

- 1 52 Miyamoto T, Ishii H, Seko T, Kobori S and Tomita Y, Occurrence of
2 *Corynespora cassiicola* isolates resistant to boscalid on cucumber in
3 Ibaraki Prefecture, Japan. *Plant Pathol.* **58**: 1144-1151 (2009).
- 4 53 Kim Y, Xiao C, Stability and fitness of pyraclostrobin- and boscalid-
5 resistant phenotypes in field isolates of *Botrytis cinerea* from apple.
6 *Phytopathology* **101**: 1385-1391 (2011).
- 7 54 Avenot HF, Michailides TJ, Resistance to boscalid fungicide in
8 *Alternaria alternata* isolates from pistachio in California. *Plant Dis.* **91**:
9 1345-1350 (2007).
- 10 55 Hjortshoj R, Ravnsboj A, Nyman M, Orabi J, Backes G, Pinnschmidt H,
11 Havis N, Stougaard J and Stukenbrock E, High levels of genetic and
12 genotypic diversity in field populations of the barley pathogen *Ramularia*
13 *collo-cygni*. *Eur. J. Plant Pathol.* **136**: 51-60 (2013).
- 14 56 Piotrowska MJ, Ennos RA, Fountaine JM, Burnett FJ, Kaczmarek M
15 and Hoebe PN, Development and use of microsatellite markers to study
16 diversity, reproduction and population genetic structure of the cereal
17 pathogen *Ramularia collo-cygni*. *Fungal Genet. Biol.* **87**: 64-71 (2016).
18
- 19 57 Rehfus A, Miessner S, Achenbach J, Strobel D, Bryson R, Stammli G,
20 Emergence of succinate dehydrogenase inhibitor resistance of
21 *Pyrenophora teres* in Europe. *Pest Manage. Sci.* DOI 10.1002/ps.4244
22 (2016).
23
- 24 McGrann GRD, Andongabo A, Sjökvist E, Trivedi U, Dussart F, Kaczmarek M,
25 Mackenzie A, Fountaine JM, Taylor JMG, Paterson LJ, Gorniak K,
26 Burnett F, Kanyuka K, Hammond-Kosack KE, Rudd JJ, Blaxter M,
27 Havis ND, The genome of the emerging barley pathogen *Ramularia*
28 *collo-cygni*. *BMC Genomics*. **In press** (2016).
29
30

1 **7 Tables**

2 **Table 1 Mean EC₅₀ values (mg litre⁻¹) of UK populations of *Ramularia collo-cygni* tested**
 3 **for five SDHI fungicides.**

| Number of <i>Rcc</i> isolates tested from UK | | | | | |
|--|----|--|--|--|--|
| 2010 | 7 | | | | |
| 2011 | 18 | | | | |
| 2012 | 37 | | | | |
| Total | 62 | | | | |

| Mean EC ₅₀ values in each year | | | | | |
|---|------------|---------|----------|-----------|----------|
| | Isopyrazam | Bixafen | Boscalid | Fluopyram | Carboxin |
| 2010 | 0.028 | 0.016 | 0.079 | 0.106 | 0.959 |
| 2011 | 0.018 | 0.013 | 0.127 | 0.128 | 0.891 |
| 2012 | 0.019 | 0.016 | 0.152 | 0.172 | 1.252 |
| Over a period of 3 years | 0.019 | 0.015 | 0.137 | 0.151 | 1.120 |

| Range of EC ₅₀ values for UK population over 3 years | | | | | |
|---|--------|--------|-------|-------|-------|
| Min. | <0.001 | <0.001 | 0.006 | 0.024 | 0.113 |
| Max. | 0.056 | 0.056 | 0.475 | 0.551 | 3.981 |

4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

1 **Table 2 Overview of amino acid and nucleotide substitutions in *Sdh* subunits genes**
2 **responsible for resistance to SDHIs in *Ramularia collo-cygni*.**

| Mutant | AA changes in the subunits B and C | Corresponding codon change | Mutations described at the same position for other plant pathogens |
|--------|------------------------------------|----------------------------|--|
| Mut1 | B: S217L | tca>tta | <i>Z. tritici</i> : B: S218F (lab mutant) ²⁸ |
| Mut11 | B: N224I | aac>atc | <i>Z. tritici</i> : B: N225I (lab mutant) ²⁸ <i>Z. tritici</i> : B: N225T (field isolate) ^a <i>B. cinerea</i> : B: N230I (field isolate) ^{13,30} |
| Mut7 | C: H142R | cat>cgt | <i>Rcc</i> : C: H142R (field isolate) ^b <i>Z. tritici</i> : C: H145R (lab mutant) ²⁸ <i>A. alternata</i> : C: H134R (field isolate) ⁴¹ <i>P. teres</i> : C: H134R (field isolate) ^{Reithus et al. 2016} |
| Mut8 | C: H142Q | cat>caa | As above |
| Mut2 | no AA changes | na | na |

^a FRAC 2014

^b FRAC 2015

1 **Table 3 Resistance factors of five SDHI-resistant *Ramularia collo-cygni* mutants.**

| | EC ₅₀ values of parental isolate and UK population (mg litre ⁻¹) | | | | |
|--|---|--------------|--------------|-------------|-------------|
| | Isopyrazam | Bixafen | Boscalid | Fluopyram | Carboxin |
| DK05 | 0.044 | 0.036 | 0.090 | 0.075 | 1.489 |
| UK pop. range ^a | <0.001-0.056 | <0.001-0.056 | 0.006- 0.475 | 0.024-0.551 | 0.113-3.981 |
| RFs (EC ₅₀ of mutant/ EC ₅₀ of parental isolate) | | | | | |
| DK05 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 |
| Mut1 B: S217L | 9.239 | 2.246 | 8.999 | 49.90 | 1.481 |
| Mut11 B: N224I | 6.758 | 12.65 | 37.28 | 21.82 | 13.23 |
| Mut7 C: H142R | 44.10 | 55.31 | 1114 | 16.77 | 32.80 |
| Mut8 C: H142Q | 0.688 | 10.69 | 1114 | 15.91 | 19.19 |
| Mut2 No AA subst. | 31.55 | 24.51 | 1.532 | 1.177 | 2.752 |
| Highest EC ₅₀ from UK pop. | 1.274 | 1.560 | 5.294 | 7.320 | 2.673 |

2 ^a UK population range = 62 isolates

3

4

5 **Table 4 Primer sets used to amplify *Ramularia collo-cygni* *SdhB*, *SdhC* and *SdhD*.**

| Subunit | Primer name | Primer sequence 5'-3' |
|-------------|------------------|------------------------|
| <i>SdhB</i> | SdhB_Rcc_Final_F | CAAATCACACACCATCCAGT |
| | SdhB_Rcc_Final_R | CCAGCCCTCTTTACATCCTC |
| <i>SdhC</i> | SdhC_Rcc_Final_F | CACTCCAGCAAACCACGACC |
| | SdhC_Rcc_Final_R | TAAAGCAGTTCTGTTGCTCT |
| <i>SdhD</i> | SdhD_Rcc_Final_F | TTCCACCACAACACCACCCACC |
| | SdhD_Rcc_Final_R | TCATCTCATCACCACACCCT |

6