THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Different parts, different stories: climate sensitivity of growth is stronger in root collars versus stems in tundra shrubs

Citation for published version:

Ropars, P, Angers-blondin, S, Gagnon, M, Myers-smith, IH, Lévesque, E & Boudreau, S 2017, 'Different parts, different stories: climate sensitivity of growth is stronger in root collars versus stems in tundra shrubs', *Global Change Biology*. <https://doi.org/10.1111/gcb.13631>

Digital Object Identifier (DOI):

[10.1111/gcb.13631](https://doi.org/10.1111/gcb.13631)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Global Change Biology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Different parts, different stories: climate sensitivity of growth is stronger in root collars versus stems in tundra shrubs

Running head: Climate sensitivity among tundra shrub parts

Pascale Ropars^{1,2,*}, Sandra Angers-Blondin^{2,3}, Marianne Gagnon^{2,4}, Isla H. Myers-Smith³, Esther Lévesque^{2,5} and Stéphane Boudreau^{2,4}

¹ Chaire de recherche du Canada en biodiversité nordique and Département de biologie, chimie et géographie, Université du Québec à Rimouski, 300 allée des Ursulines, Rimouski, QC, G5L 3A1

² Centre d'études nordiques, 2405 av. de la Terrasse, Québec, QC, G1V 0A6

³ School of GeoSciences, University of Edinburgh, Edinburgh, UK, EH9 3FF

⁴ Département de biologie, Université Laval, 1045 av. de la Médecine, Québec, QC, G1V 0A6

⁵ Département des sciences de l'environnement, Université du Québec à Trois-Rivières, 3351 boul. des Forges, Trois-Rivières, QC, G9A 5H7

*Corresponding author: pascale.ropars@uqar.ca; T: +1-418-656-2131 # 12004; F: +1-418-656-2043

Keywords *Betula glandulosa*, Climate sensitivity, Dendrochronology, Dwarf birch, Shrub densification, Subarctic, Tundra

Type of paper Primary Research Article

Abstract

Shrub densification has been widely reported across the circumpolar arctic and subarctic biomes in recent years. Long-term analyses based on dendrochronological techniques applied to shrubs have linked this phenomenon to climate change. However, the multi-stemmed structure of shrubs makes them difficult to sample and therefore leads to non-uniform sampling protocols among shrub ecologists, who will favor either root collars or stems to conduct dendrochronological analyses. Through a comparative study of the use of root collars and stems of *Betula glandulosa*, a common North American shrub species, we evaluated the relative sensitivity of each plant part to climate variables, and assessed if this sensitivity is consistent across three different types of environments in northwestern Québec, Canada (terrace, hilltop and snowbed). We found that root collars had greater sensitivity to climate than stems, and that these differences were maintained across the three types of environment. Growth at the root collar was best explained by spring precipitation and summer temperature, whereas stem growth showed weak and inconsistent responses to climate variables. Moreover, sensitivity to climate was not consistent among plant parts, as individuals having climate sensitive root collars did not tend to have climate sensitive stems. These differences in sensitivity of shrub parts to climate highlight the complexity of resource allocation in multi-stemmed plants. Whereas stem initiation and growth are driven by micro-environmental variables such as light availability and competition, root collars integrate the growth of all plant parts instead, rendering them less affected by mechanisms such as competition and more responsive to signals of global change. Although further investigations are required to determine the degree to which these findings are generalizable across the tundra biome, our results indicate that consistency and caution in the choice of plant parts are a key consideration for the success of future dendroclimatological studies on shrubs.

Introduction

Shrubs are one of the most responsive plant functional groups to recent climate change (Elmendorf *et al.*, 2012) and their increase in abundance in tundra ecosystems have been cited as driver of the satellite-observed greening of the Arctic (Raynolds *et al.*, 2006; Tape *et al.*, 2006; McManus *et al.*, 2012). The increase in shrub cover and abundance is widespread across northern circumpolar regions and has been recorded at both high latitude and high altitude sites in North America (Sturm *et al.*, 2001a; Tape *et al.*, 2006; Myers-Smith *et al.*, 2011a; Ropars & Boudreau, 2012), northern Europe (Bär *et al.*, 2008; Hallinger *et al.*, 2010) and Russia (Forbes *et al.*, 2010). Shrubs have either expanded in patch size (Sturm *et al.*, 2001a; Tape *et al.*, 2006; Ropars & Boudreau, 2012; Tremblay *et al.*, 2012), recruited in areas where they were less abundant in the past (Sturm *et al.*, 2001a; Frost *et al.*, 2013; Frost & Epstein, 2014; Buntgen *et al.*, 2015) or have increased in height (Myers-Smith *et al.*, 2011b; Paradis *et al.*, 2016). Though this increase in tundra shrubs has been attributed to recent warming in tundra ecosystems (Elmendorf *et al.*, 2012; Ropars *et al.*, 2015), the climate sensitivity of shrub growth has been demonstrated to be variable across the tundra biome (Myers-Smith *et al.*, 2015a).

Understanding how shrub growth is controlled by climate is key to predicting vegetation change and its associated impacts on ecosystem function in a warming tundra biome. In treeless ecosystems, erect shrubs provide structure for both plant and animal communities (Tape *et al.*, 2016) and modify important ecological processes and physical characteristics of the environment (Cushman *et al.*, 2010). In tundra regions, shrubs are expected to have complex and sometimes seasonally contrasting or species-specific effects on permafrost dynamics or soil temperature regime (Sturm *et al.*, 2001b; Blok *et al.*, 2011; Myers-Smith & Hik, 2013; Paradis *et al.*, 2016) and absorption of solar radiation (Chapin *et al.*, 2005; Williamson *et al.*, 2016). Shrubs provide important habitat and food sources for moose, caribou, ptarmigan, hare and other wildlife species

(Tape *et al.*, 2010, 2016; Boelman *et al.*, 2015; Christie *et al.*, 2015), but might also reduce lichen availability in caribou winter ranges (Joly *et al.*, 2007). Since erect shrub species have a significant impact on the dynamics and physical properties of arctic and subarctic ecosystems, it is essential to better understand the factors driving their growth and rapid expansion.

Dendrochronological approaches have successfully been applied to shrub species to analyze temporal and spatial processes in fast-changing tundra ecosystems (e.g. Bär *et al.*, 2008; Forbes *et al.*, 2010; Blok *et al.*, 2011; Myers-Smith *et al.*, 2015a; Ropars *et al.*, 2015). Like trees, shrubs form annual growth rings that can be related to climate variables (e.g., Myers-Smith *et al.*, 2015a) or disturbances in ecosystem processes such as landslides (e.g., Gers *et al.*, 2001) and permafrost degradation (e.g., Gaertner-Roer *et al.*, 2013). However, the multi-stemmed growth form of shrubs has several ecological and physiological implications that may influence climate sensitivity among their different parts. For example, shrubs have the potential to allocate resources to different stems and roots depending on environmental conditions (Pajunen, 2009) and competition for light and nutrients in dense shrub patches could lead to differential stem growth within an individual (Hallinger *et al.*, 2010; Myers-Smith *et al.*, 2015b). Moreover, there is still very little known about how stems are initiated and replaced within a shrub individual (but see Charles-Dominique, 2011). All of these complex interactions could result in different radial growth patterns between the stems of a single individual. Despite these considerations, stems and root collars (i.e. the oldest part of a shrub, found at the junction between roots and stems) are often used without distinction both across (Myers-Smith *et al.*, 2011a) and within studies (Dumais *et al.*, 2014).

A review of shrub dendrochronological studies reveals that stems are more frequently used than root collars (44/69 studies compared to 15/69; Fig. 1 and Table S1). The strong preference towards stems likely arises from some of the following reasons: root collars are hard to access in

certain environments and in areas of continuous shrub cover, their sampling is destructive and their identification can be difficult, especially when shrub individuals display complex morphologies such as adventitious roots and buried stems (Myers-Smith *et al.*, 2015b). As a consequence, it is often easier to sample the largest and/or longest stems of an individual, even if it does not track the entire shrub growth record (De Witte *et al.*, 2012). However, comparing the climate sensitivity of shrubs across different sites or regions could be potentially problematic if different shrub parts exhibit different growth trends that are influenced not only by climate but also by within-individual competition and resource allocation.

The growing number of studies dealing with the sensitivity of shrub growth to climate change highlights the need to converge towards standardized methods in dendroecology applied to shrubs, especially in the fast-changing tundra biome. Here, we present a comparative study of the use of root collars and stems in dendroclimatology analyses. Specifically, we aim (1) to evaluate the relative sensitivity of stems and root collars to climate, and (2) to evaluate if this relative sensitivity is consistent across different types of environments. We hypothesize that climate sensitivity will be higher for root collars, given that they integrate growth from the whole individual, and that this difference will be maintained across the landscape.

Materials and methods

Study area and climatic trends

The study area is located around the Boniface River research station (Centre d'études nordiques, <http://www.cen.ulaval.ca/page.aspx?lien=stationboniface>; 57° 45' N, 76° 20' W), about 10 km south of the Arctic treeline in subarctic Québec (Canada). The area lies within the discontinuous permafrost zone (Payette, 2001) and belongs to the shrub subzone of the forest-tundra ecotone (Payette, 1983). At the landscape scale, shrub tundra covers *ca.* 70% of the well-drained sites (Payette *et al.*, 2008). Wetlands, which include palsas and snowbeds, cover 7% of the terrestrial area. *Betula glandulosa* Michx. (hereafter referred to as dwarf birch), a multi-stemmed species widely distributed in North America, is the most abundant shrub species and is responsible for most of the shrub expansion observed in the study area (Ropars & Boudreau, 2012). According to Ju & Masek (2016), this is the Canadian region where the greening was the most extensive in the last three decades. Large herbivores (caribou: *Rangifer tarandus* L.) are present in the area, but browsing is mostly restricted to *Salix* species.

The closest meteorological station (Inukjuak Meteorological Station, Environment Canada; 58° 28' N, 78° 05' W; 130 km northwest of the study site) recorded an annual mean temperature of -7 °C for the 1971-2000 period, with the highest and lowest mean monthly temperatures recorded in July (9.4° C) and February (-25.8 °C), respectively (Canadian climate normal; Environment Canada, 2016). Annual precipitation averaged 460 mm, of which 42% fell as snow (Canadian climate normal; Environment Canada, 2016). Between 2000 and 2009, mean annual temperature has reached - 5.5 °C, with July and February temperatures averaging 11.7 °C and - 24.8 °C. Like many regions in northern Québec, Inukjuak is experiencing a warming trend that began in the early 1990s (Chouinard *et al.*, 2007; Bhiry *et al.*, 2011).

Site selection and field sampling

The root collar and the two main stems of dwarf birch were sampled in three different environments to assess potential site-driven differences in climate sensitivity: sandy terraces (hereafter referred to as terraces), low-altitude hilltops (hereafter referred to as hilltops) and snowbeds. The terraces are well-drained, low-altitude sites located on the banks of the Boniface River. They are characterized by lichens, graminoids and large patches of shrub species, mainly dwarf birch, although willows and alders can be found close to the river. The hilltops are characterized by the presence of arctic-alpine species and exposed mineral soil. Snowbeds are periglacial environments where snow accumulates preferentially during winter and melts later in the growing season, sometimes as late as mid-July. Consequently, they are characterized by plant species well-adapted to winter conditions and to an excess of moisture in the first half of the growing season (Payette & Lajeunesse, 1980; Filion & Payette, 1982; Morin & Payette, 1986). One site in each environment type was randomly selected from a pool of 147 identified sites (50 terraces, 50 hilltops, and 47 snowbeds; see Ropars *et al.*, 2015 for more details on site selection).

In summer 2010, we selected and uprooted 20 dwarf birch individuals at each site. We prioritized large and relatively isolated individuals with a circular form, as they were thought to be older individuals grown from seed. When such individuals were not available, particular attention was taken to discard individuals for which the root collar could not be easily identified. Dwarf birch individuals grown from seeds usually display a short (< 3cm) but well-defined root collar, located between the root and stem systems (Fig. 1c). From the three to six main stems directly emerging just above the root collar, we selected the two largest ones and cut them as close as possible to the root collar and the soil surface. The two stem samples and the root collar

of each dwarf birch individual were collected, carefully cleaned, and left to fully dry at room temperature.

Radial growth measurement

Dwarf birch root collars were sliced (*ca.* 25 μm) using a rotary microtome after being boiled for at least 3 hours. Thin sections were then stained with safranin (1% solution, Safranin O, Fischer Science Education), dried and permanently mounted with a 66% toluene solution (SHUR/mount^{TM} liquid cover glass, Triangle biomedical sciences). Digital photographs of each sample were taken using a binocular-mounted camera (Olympus SZ61 with a SC100 camera). Root collars were discarded if they could not be sliced perpendicularly (stems and roots were too intermingled), or if they were rotten. Of the 60 root collars processed for the three sites, 46 were kept for further analyses (terrace: 15, hilltop: 16, snowbed: 15). The two largest stems of each of these 46 individuals (92 stems) were processed following the above procedures.

Using digital photographs, we aged each root collar and stem sample with the ImageJ freeware (v. 1.40g) while ring widths were measured using the dendrochronological software LignoVision (v. 1.36, Rinntech). If possible, ring widths were measured along two opposite radii and averaged for each sample. Growth measurements were visually examined and statistically verified with COFECHA, a widely used statistical crossdating program (Holmes, 1983). We used the mean detrending method for every root collar and stem ring-width series to allow comparison between individuals. Incomplete rings were frequently encountered, but easily detectable. Unlike other willow and birch species growing at their altitudinal limit (Wilmking *et al.*, 2012), no missing rings were identified in the root collar samples.

185 *Measuring climate sensitivity*

186 The climate sensitivity of root collars and stems was measured using mixed-effects models. We
187 used ring-width data for the period 1947-2009, for which monthly temperature and precipitation
188 data were available from the Inukjuak Meteorological Station. We ran a series of linear mixed-
189 effects models using standardised (mean centered) ring width as a response variable, monthly
190 temperature and/or precipitation variables (singly or in combination; see Table S2 for the full list
191 of models) and the sample type (root collar vs stem) as interacting predictors, and with year as a
192 random effect. We also ran a null model, i.e. a model with the same random effect structure that
193 does not include any climate variables as fixed effects to serve as baseline for the climate
194 sensitivity analysis (Ettinger *et al.*, 2011; Myers-Smith *et al.*, 2015a). We allowed for interactions
195 between fixed effects because we were specifically interested in the potentially different climate
196 responses of shrub root collars and stems. We ran a separate series of models for each site, as the
197 added complexity of including a site level in the model structure while allowing for random
198 slopes caused convergence issues. We ran the analyses with the lme4 package (R version 3.2.2),
199 using the maximum likelihood estimation for model selection and the restricted maximum
200 likelihood estimation to estimate slopes.

201 As our interaction models (sample type * climate) were driven mostly by the stronger
202 response of the root collars, we also ran separate models for stems vs root collars at each site to
203 check whether stems might be more responsive to a different climate variable. We additionally
204 ran the same model structure with the nlme package to compare model results to models with a
205 temporal autocorrelation structure, and using raw (non-standardized) ring-width data and a subset
206 of the data restricted to the common period covered by both stem and root collar chronologies
207 (1960-2009). These complementary analyses yielded the same results, and thus are not presented

in the main text (see Table S3 for results obtained with raw data, the 1960-2009 subset and using the nlme package).

We used model selection analysis to identify the best climate model(s), i.e. the variables that best explain variations in radial growth. We ranked the models according to the Akaike information criterion (AIC) and calculated the difference between each model and the null model, which we denoted by ΔAIC_{null} . ΔAIC_{null} is used as a measure of the sensitivity of radial growth to specific climate variables (Myers-Smith *et al.*, 2015a). As the AIC only assesses the relative quality of competing models, but not the absolute goodness of fit, we also calculated the marginal and conditional R^2 of each model with the r.squaredGLMM function of the MuMin package (R version 3.2.2). The conditional R^2 represents the amount of variance explained by all the fixed and random effects in the model, while the marginal R^2 is the amount of variance explained by the fixed effects only (Nakagawa & Schielzeth, 2012).

Linear models

To investigate whether stem and root collar sensitivity were correlated or decoupled within an individual, we ran a linear model with ring width as a function of the two best climate variables (July temperature and March precipitation) identified through the model selection described above for each individual. We then compared whether the slopes differed between plant parts for each individual. We also used these slope values to test whether there is a significant relationship between climate sensitivity of root collars and their stem counterparts. To do so, we ran a linear model with the slope values obtained for stems as a function of the ones obtained for root collars. We applied a squared-root transformation to the slope values to attain normality (Legendre & Legendre, 1998). A constant was added to the slope values prior to transformation to cope with negative numbers.

Results

Sensitivity of stems and root collars to climate

We found greater sensitivity of *Betula glandulosa* root collars versus stems across our sampling sites (Fig. 2). Climate sensitivity of stems was generally weak, the null model ranking between the 4th and 7th best model (Table S1). Moreover, the proportion of the variance in growth explained by climate variables (i.e. marginal R^2) for root collars was at least twice as high as for stems in the snowbed and hilltop sites, and there was a similar but weaker trend for the terrace site.

Climate variables best explaining growth

We found that March precipitation and July temperature best explained shrub growth, but with a strong interaction with sample type. Root collars showed a consistent positive response to these two climate variables, but not stems (Fig. 2). Moreover, when stems and root collars were analyzed separately, the latter showed greater climate sensitivity than stems to most climate variables (monthly mean temperature and total precipitation; Fig. 3). The difference in AIC score between the null model and the best climate model was systematically higher for root collars than for stems in each type of environment (Table 1). March precipitation and July temperature best explained the growth at the root collar, and this relationship was maintained across the landscape. On the other hand, stems showed weak and inconsistent responses among the three environments; August, May and April temperature best explained stem growth in snowbed, hilltop and terrace, respectively.

Climate sensitivity of individuals among shrub parts

Individual-level climate sensitivity did not show a consistent trend among root collar and stems (Fig. 4). Individuals having a climate sensitive root collar did not tend to have climate sensitive stems, as only five climate sensitive root collars out of 42 had one or two of their stem counterparts sensitive to climate as well. Of the 11 climate sensitive stems, seven were associated with climate sensitive root collars. Moreover, root collar and stems from the same individual showed opposite significant relation with July temperature in one particular case (individual S27, Fig. 4). The climate sensitivity of stems to both July temperature (linear model, $F_{1,90} = 2.927$, $P = 0.091$) and March precipitation (linear model, $F_{1,88} = 1.624$, $P = 0.206$) was not explained by the climate sensitivity of their root collar counterparts.

Root collar and stem samples characteristics

We found that root collar ages were greater than stem ages and that ages varied across the different environments sampled. Root collar samples averaged 41 years of age, but ranged between 10 and 96 (Table 2). The oldest root collar was from an individual growing on a hilltop, whereas the youngest were found in the snowbed environment. There was no difference in root collar samples age between the three types of environment ($F_{1,3} = 1.02$, $P = 0.369$). Stem samples averaged 23 years of age, but some reached over 60 years in the hilltop environment. The stems of dwarf birch individuals growing in the snowbed site were younger (18 years old compared to 27 and 25 for hilltop and terrace; Tukey multiple comparison, $P < 0.01$) and longer (from the root collar to the tip of the stem) than those of individuals growing on the hilltop and terrace sites (Table 2; 133 cm compared to 92 cm and 102 cm for hilltop and terrace, Tukey multiple comparison, $P < 0.01$).

Discussion

In this study, we demonstrated that different parts of an important tundra shrub species in North America, *Betula glandulosa*, have contrasting responses to climate, with root collars exhibiting far greater sensitivity than stems. Growth at the root collar was best explained by spring precipitation and summer temperature, and this relationship was maintained across the landscape. Stems, on the other hand, showed weak and inconsistent responses. These results highlight the complexity of resource allocation in multi-stemmed shrubs and call for caution in sampling when aiming to assess the climate sensitivity of shrub growth.

What explains the difference in sensitivity of root collars vs stems?

The multi-stemmed structure of shrubs results in complex resource allocation patterns among plant parts that could dampen the influence of climate on stems growth. In their review of intra-plant competition, Sadras & Denison (2009) concluded that plant individuals might use competition-like mechanisms to allocate resources such as nutrients, but only if it increases overall individual fitness. Competition within plant individuals is known to lead to trade-offs between size and number of plant units (Sadras, 1995; Chikov, 2008; Sadras & Denison, 2009) as well as between vegetative and reproductive structures (e.g. Liu *et al.*, 2007; Quilot & Genard, 2008). The vertical growth of shrubs can also be limited by the abrasive effect of windblown ice particles (Sonesson & Callaghan, 1991), resulting in differential growth patterns among stems. Moreover, the drivers of stem initiation at the root collar are poorly understood. Light availability promotes stem initiation of *Rhamnus cathartica* L., a deciduous shrub growing in southern Québec (Charles-Dominique *et al.*, 2012) and an increased herbivory pressure is known to alter the vertical structure of *Salix* and *Betula* species in northern Alaska (Tape *et al.* 2010; Christie *et al.*, 2014). The growth of each stem is therefore affected not only by climatic and other

environmental conditions, but also by differential resources allocation among plant parts that enhances the individual's global fitness. On the other hand, root collars integrate the growth of each plant part. Thus, they might be less affected by competition-like mechanisms and more responsive to climate.

Root collars showed consistent responses to climate variables across the landscape, with both July temperature and March precipitation promoting enhanced radial growth. Warmer temperatures during the short growing season have been identified as key factors for promoting shrub growth in different Arctic regions (e.g. Bär *et al.*, 2008; Forbes *et al.*, 2010; Hallinger *et al.*, 2010; Myers-Smith *et al.* 2015a). Higher precipitation in March could reduce the risk of frost damage following early leaf-out, and protect buds and stems from the abrasive effect of drifting ice particles (Sonesson & Callaghan, 1991). Among other possible explanations, an increase in snow precipitation could also promote a better radial growth by increasing water discharge and thus water availability in early summer. Greater snow accumulation may also lead to greater insulation, higher soil temperature and increased microbial activity, which in turn could lead to higher nutrient availability (Chapin *et al.*, 2005).

To our knowledge, the present study is the first to compare climate sensitivity among different parts of shrub individuals. The harsh climatic conditions of arctic and alpine regions can result in intra-plant growth irregularities that could influence climate sensitivity, such as missing or incomplete rings (Hallinger *et al.*, 2010; Buchwal *et al.*, 2013). The distribution of these growth irregularities has been shown to be heterogeneous among shrub parts (Hallinger *et al.*, 2010). For *Salix polaris* Wahlenb for instance, the growth at the root collar and in the belowground system was consistently greater than the growth in stems (Buchwal *et al.*, 2013). While serial sectioning (Kolishchuk, 1990), a technique consisting in repeated tree-ring width measurements along shrub stems, allows to properly date woody plant individuals (Hallinger *et*

al., 2010; Wilmking *et al.*, 2012; Buchwal *et al.*, 2013; Hollesen *et al.*, 2015), it also emphasizes growth heterogeneity among plant parts. For different northern pine and spruce species, differences in growth-climate relationships have been observed among plant parts, breast-height and upper stem samples being more sensitive to previous and current season temperatures and moisture, respectively (Kozłowski *et al.*, 1991; Chhin & Wang, 2008; Chhin *et al.*, 2008, 2010). Our findings suggest that complex controls are also at play in determining the growth of shorter-statured but multi-stemmed plants.

Implication of the study

Changes in shrub cover will play an important ecological role in arctic regions, emphasizing the need to better understand their drivers and to predict their magnitude. Large scale meta-analyses focusing on many taxa are useful to address these questions (see Myers-Smith *et al.*, 2015a), but their conclusions could be challenged by a lack of consistency in sampling methods across studies. Our results show that root collars are consistently more climate sensitive than stems for a widely distributed shrub species, *B. glandulosa*, one of the most commonly reported shrub species to be expanding in tundra ecosystems (Tape *et al.*, 2006; Myers-Smith *et al.*, 2011a; Ropars & Boudreau, 2012; Tremblay *et al.*, 2012). Moreover, a climate sensitive root collar was not necessarily associated with climate sensitive stems (Fig. 4). In their recent meta-analysis, Myers-Smith and collaborators (2015a) found greater climate sensitivity in northwest Russia compared to North America. Based on our literature review (Fig. 1 and Table S1), these differences could be partly explained by the differential use of stems and root collars across studies. Indeed, most Russian studies have used root collars (8 out of 11 studies, 73%), whereas only a third of North American ones did (10 out of 30 studies, 33%; Table S1). The sensitivity of North American shrubs could therefore have been underestimated and could overlook the

potential increase in shrub abundance and canopy height. While the differences among shrub parts have been overlooked in the past, our study highlights the importance of establishing consistent sampling protocols for shrub dendrochronological analysis.

How do we move forward with future dendroecological analyses?

While this study clearly demonstrates that different plant parts show different climate sensitivity, further investigations are required to understand how generalizable our findings are across the tundra biome. In our study, we focused on one shrub species, *Betula glandulosa*, growing in a specific subarctic area of North America. Shrub species having similar growth forms and branching architecture (like alders and willows) could potentially also show similar differences in climate sensitivity among plant parts, but a multispecies comparison would be needed to confirm this hypothesis. Architectural studies of shrubs are arising in the scientific literature and could bring us a step further in our comprehension of energy allocation among plant parts (Charles-Dominique, 2011). We therefore propose that architectural and comparative dendrochronological studies be combined to account for potential differences in growth patterns among shrub parts. If dissimilarities between plant parts are also observed in other sites and species, we urge for the establishment of a standardized sampling methodology for future studies. Based on the fact that root collars were more climate sensitive than stems, we propose that future studies systematically sample shrub individuals as close as possible to the root collar.

Understanding the dynamic of shrub populations and the drivers of their growth is key to assess and predict global ecosystem processes under climate change, especially in Arctic regions where they structure plant communities. Our study clearly demonstrates that root collars provide a better integration of the climate sensitivity of shrub individuals for a widely distributed species

in North America, *Betula glandulosa*. Root collars were more climate sensitive than stems and showed consistent sensitivity to July temperature and March precipitation across three different types of environments in the Boniface River region. Finally, we found no consistency between shrub parts, individuals having a climate sensitive root collar not necessarily having climate sensitive stems. While different shrub parts have been sampled for dendrochronological analyses in the past, this study urges for a standardized sampling methods. To reveal reliable climate sensitivity of tundra shrub species and, more importantly, to conduct multi-sites and multi-species analysis, we propose to systematically sample at root collar.

Acknowledgements

This research was founded by Natural Sciences and Engineering Research Council of Canada (NSERC), by the Fonds de Recherche Québec - Nature et Technologies (FRQNT), by the Northern Scientific Training Program, by the NSERC CREATE training Program in Northern Environmental Sciences (EnviroNorth) and by the Northern Research Chair on Disturbance Ecology. The authors are grateful to Caroline Mercier for help in the field as well as to Annie Girard, Mélody Dos Santos and Mélissa Paradis for lab assistance. The authors would also like to thank Ann Delwaide for valuable help with dendrochronological analysis and the Centre d'études nordiques for logistical support.

References

- Bär A, Pape R, Brauning A, Löffler J (2008) Growth-ring variations of dwarf shrubs reflect regional climate signals in alpine environments rather than topoclimatic differences. *Journal of Biogeography*, **35**, 625-636.
- Bhury N, Delwaide A, Allard M *et al.* (2011) Environmental change in the Great Whale River region, Hudson Bay: five decades of multidisciplinary research by Centre d'études nordiques (CEN). *Ecoscience*, **18**, 182-203.
- Blok D, Sass-Klaassen U, Schaepman-Strub G, Heijmans MMPD, Sauren P, Berendse F (2011) What are the main climate drivers for shrub growth in Northeastern Siberian tundra? *Biogeosciences*, **8**, 1169–1179.
- Boelman NT, Gough L, Wingfield J *et al.* (2015) Greater shrub dominance alters breeding habitat and food resources for migratory songbirds in Alaskan arctic tundra. *Global Change Biology*, **21**, 1508-1520.
- Buchwal A, Rachlewicz G, Fonti P, Cherubini P, Gartner H (2013) Temperature modulates intra-plant growth of *Salix polaris* from a high Arctic site (Svalbard). *Polar Biology* **36**, 1305-1318.
- Büntgen U, Hellmann L, Tegel W, Normand S, Myers-Smith I, Kirdyanov AV, Nievergelt D, Schweingruber FH (2015) Temperature-induced recruitment pulses of Arctic dwarf shrub communities. *Journal of Ecology*, **103**, 489-501.

416

417 Chapin FS, Sturm M, Serreze MC *et al.* (2005) Role of land-surface changes in Arctic summer
 418 warming. *Science*, **310**, 657-660.

419

420 Charles-Dominique T (2011) Analyse des relations entre plasticité architecturale des buissons et
 421 prolifération de leurs populations. Thesis. Université Montpellier II – Sciences et Technologies
 422 du Languedoc, Montpellier, France.

423

424 Charles-Dominique T, Edelin C, Brisson J, Bouchard A (2012) Architectural strategies of
 425 *Rhamnus cathartica* (Rhamnaceae) in relation to canopy openness. *Botany*, **90**, 976-989.

426

427 Chhin S, Hogg EHT, Lieffers VJ, Huang S (2008) Potential effects of climate change on the
 428 growth of lodgepole pine across diameter size classes and ecological regions. *Forest Ecology and*
 429 *Management*, **256**, 1692-1703.

430

431 Chhin S, Wang GG (2008) Climatic response of *Picea glauca* seedlings in a forest-prairie
 432 ecotone of western Canada. *Annals of Forest Science*, **65**, 207.

433

434 Chhin S, Hogg EH, Lieffers VJ, Huang S (2010) Growth-climate relationships vary with height
 435 along the stem in lodgepole pine. *Tree Physiology*, **30**, 335-345.

436

437 Chikov VI (2008) Evolution of notions about relationships between photosynthesis and plant
 438 productivity. *Russian Journal of Plant Physiology*, **55**, 130-143.

439

Chouinard C, Fortier R, Mareschal JC (2007) Recent climate variations in the Subarctic inferred from three borehole temperature profiles in northern Québec, Canada. *Earth and Planetary Science Letters*, **263**, 355-369.

Christie KS, Ruess RW, Lindberg MS, Mulder CP (2014) Herbivores influence the growth, reproduction, and morphology of a widespread arctic willow. *Plos One*, **9**, e101716.

Christie KS, Bryant JP, Gough L, Ravolainen VT, Ruess RW, Tape KD (2015) The role of vertebrate herbivores in regulating shrub expansion in the Arctic: a synthesis. *BioScience*, **65**, 1123-1133.

Cushman JH, Waller JC, Hoak DR (2010) Shrubs as ecosystem engineers in a coastal dune: influences on plant populations, communities and ecosystems. *Journal of Vegetation Science*, **21**, 821-831.

De Witte LC, Armbruster GFJ, Gielly L, Taberlet P, Stocklin J (2012) AFLP markers reveal high clonal diversity and extreme longevity in four key arctic-alpine species. *Molecular Ecology*, **21**, 1081-1097.

Dumais C, Ropars P, Denis MP, Dufour-Tremblay G, Boudreau S (2014) Are the low altitude alpine tundra ecosystems under threat? A case study from the Parc National de la Gaspésie, Québec. *Environmental Research Letters*, **9**, 094001.

463 Elmendorf SC, Henry GHR, Hollister RD *et al.* (2012) Global assessment of experimental
 464 climate warming on tundra vegetation: heterogeneity over space and time. *Ecology Letters*, **15**,
 465 164-175.
 466
 467 Environment Canada (2016) *Canadian climate normal of averages 1971-2000*. Available at
 468 http://climate.weather.gc.ca/climate_normals/index_e.html.
 469
 470 Ettinger AK, Ford KR, HilleRisLambers J (2011) Climate determines upper, but not lower,
 471 altitudinal range limits of Pacific Northwest conifers. *Ecology*, **92**, 1323-1331.
 472
 473 Filion L, Payette S (1982) Régime nival et végétation chionophile à Poste-de-la-Baleine,
 474 Nouveau-Québec. *Naturaliste Canadien*, **109**, 557-571.
 475
 476 Forbes BC, Macias Fauria M, Zetterberg P (2010) Russian Arctic warming and “greening” are
 477 closely tracked by tundra shrub willows. *Global Change Biology*, **16**, 1542-1554.
 478
 479 Frost GV, Epstein HE, Walker DA, Matyshak G, Ermokhina K (2013) Patterned-ground
 480 facilitates shrub expansion in Low Arctic tundra. *Environmental Research Letters*, **8**, 015035.
 481
 482 Frost GV, Epstein HE (2014) Tall shrub and tree expansion in Siberian tundra ecotones since the
 483 1960s. *Global Change Biology*, **20**, 1264-1277.
 484
 485 Gaertner-Roer I, Heinrich I, Gaertner H (2013) Wood anatomical analysis of Swiss willow (*Salix*
 486 *helvetica*) shrubs growing on creeping mountain permafrost. *Dendrochronologia*, **31**, 97-104.

487

488 Gers E, Florin N, Gaertner H, Glade T, Dikau R, Schweingruber FH (2001) Application of
 489 shrubs for dendrogeomorphological analysis to reconstruct spatial and temporal landslide
 490 movement patterns – a preliminary study. *Zeitschrift fur Geomorphologie*, **125**, 163-175.

491

492 Hallinger M, Manthey M, Wilmking M (2010) Establishing a missing link: warm summers and
 493 winter snow cover promote shrub expansion into alpine tundra in Scandinavia. *New Phytologist*,
 494 **186**, 890–899.

495

496 Hollesen J, Buchwal A, Rachlewicz G, Hansen BU, Hansen MO, Stecher O, Elberling B (2015)
 497 Winter warming as an important co-driver for *Betula nana* growth in western Greenland during
 498 the past century. *Global Change Biology*, **21**, 2410-2423.

499

500 Holmes RL (1983) Computer assisted quality control in tree ring dating and measurement. *Tree-*
 501 *Ring Bulletin*, **43**, 69-78.

502

503 Joly K, Jandt RR, Meyers CR, Cole MJ (2007) Changes in vegetation cover on Western Arctic
 504 Herd winter range from 1981 to 2005: potential effects of grazing and climate change. *Rangifer*,
 505 **17**, 199-206.

506

507 Ju J, Masek JG (2016) The vegetation greenness trend in Canada and US Alaska from 1984-2012
 508 Landsat data. *Remote Sensing of Environment*, **176**, 1-16.

509

510 Kolishchuk V (1990) Dendroclimatological study of prostrate woody plant. In: *Methods of*
 511 *dendrochronology applications in the environmental sciences* (eds Cook ER, Kairiukstis LA), pp.
 512 51-54, Kluwer Academia Publishers, Dordrecht, The Netherlands.
 513
 514 Kozłowski TT, Kramer PJ, Pallardy SG (1991) The physiological ecology of woody plants.
 515 Academic Press, San Diego, USA.
 516
 517 Legendre P, Legendre L (1998) *Numerical Ecology*, pp. 39-45, Elsevier, Amsterdam, The
 518 Netherlands.
 519
 520 Liu HF, Genard M, Guichard S, Bertin N (2007) Model-assisted analysis of tomato fruit growth
 521 in relation to carbon and water fluxes. *Journal of Experimental Botany*, **58**, 3567-3580.
 522
 523 McManus K, Morton DC, Masek JG, Wang D, Sexton JO, Nagol JR, Ropars P, Boudreau S
 524 (2012) Satellite-based evidence for shrub and graminoid tundra expansion in northern Quebec
 525 from 1986 to 2010. *Global Change Biology*, **18**, 2313–2323.
 526
 527 Morin H, Payette S (1986) The vegetation of snow patches from the Richmond Gulf, northern
 528 Québec. *Canadian Journal of Botany*, **64**, 1515-1524.
 529
 530 Myers-Smith IH, Forbes BC, Wilmking M *et al.* (2011a) Shrub expansion in tundra ecosystems:
 531 dynamics, impacts and research priorities. *Environmental Research Letters*, **6**, 045509.
 532

533 Myers-Smith IH, Hik DS, Kennedy C, Cooley D, Johnstone JF, Kenney AJ, Krebs CJ (2011b)
534 Expansion of canopy-forming willows over the twentieth century on Hershel Island, Yukon
535 Territory, Canada. *Ambio*, **40**, 610-623.
536

537 Myers-Smith IH, Hik DS (2013) Shrub canopies influence soil temperatures but not nutrient
538 dynamics: an experimental test of tundra snow-shrub interactions. *Ecology and Evolution*, **3**,
539 3683-3700.
540

541 Myers-Smith IH, Elmendorf SC, Beck PSA *et al.* (2015a) Climate sensitivity of shrub growth
542 across the tundra biome. *Nature Climate Change*, **5**, 887-891.
543

544 Myers-Smith IH, Hallinger M, Blok D *et al.* (2015b) Methods for measuring arctic and alpine
545 shrub growth: a review. *Earth-Science Reviews*, **140**, 1-13.
546

547 Nakagawa S, Schielzeth H (2012) A general and simple method for obtaining R^2 from
548 generalized linear mixed-effects models. *Methods in Ecology and Evolution*, **4**, 133-142.
549

550 Pajunen AM (2009) Environmental and biotic determinants of growth and height of Arctic
551 willow shrubs along a latitudinal gradient. *Arctic, Antarctic and Alpine Research*, **41**, 478-485.
552

553 Paradis M, Lévesque E, Boudreau S (2016) Greater effect of increasing shrub height on winter
554 versus summer soil temperature. *Environmental Research Letters*, **11**, 085005.
555

556 Payette S (1983) The forest tundra and present tree-lines of the northern Québec-Labrador
557 peninsula. Tree-line Ecology, Proceedings of the Northern Québec Tree-line Conference
558 (Québec: Nordica).

559

560 Payette S (2001) Les processus et les formes périglaciaires. In: *Écologie des tourbières du*
561 *Québec-Labrador* (eds Payette S, Rochefort L), pp. 199-239, Presses de l'Université Laval,
562 Québec, Canada.

563

564 Payette S, Lajeunesse R (1980) Les combes à neige de la rivière aux Feuilles (Nouveau-
565 Québec) : indicateurs paléoclimatiques holocènes. Géographie physique et Quaternaire, **34**, 209-
566 220.

567

568 Payette S, Filion L, Delwaide A (2008) Spatially explicit fire-climate history of the boreal forest-
569 tundra (eastern Canada) over the last 2000 years. Philosophical Transactions of the Royal Society
570 Biological Sciences, **363**, 2301-2316.

571

572 Quilot B, Genard M (2008) Is competition between mesocarp cells of peach fruits affected by the
573 percentage of wild species genome? Journal of Plant Research, **121**, 55-63.

574

575 Raynolds MK, Walker DA, Maier HA (2006) NDVI patterns and phytomass distribution in the
576 circumpolar Arctic. Remote Sensing of Environment, **102**, 271–281.

577

578 Ropars P, Boudreau S (2012) Shrub expansion at the forest tundra ecotone: spatial heterogeneity
579 linked to local topography. Environmental Research Letters, **7**, 015501.

580

581 Ropars P, Lévesque E, Boudreau S (2015) How do climate and topography influence the

582 greening of the forest tundra ecotone in northern Québec? A dendrochronological analysis of

583 *Betula glandulosa*. Journal of Ecology, **103**, 401-412.

584

585 Sadras VO (1995) Compensatory growth in cotton after loss of reproductive organs: a review.

586 Field Crops Research, **40**, 1-18.

587

588 Sadras VO, Denison RF (2009) Do plant compete for resources? An evolutionary viewpoint.

589 New Phytologist, **183**, 565-574.

590

591 Sonesson M, Callaghan TV (1991) Strategies of survival in plants of the Fennoscandian tundra.

592 Arctic, **44**, 95-105.

593

594 Sturm M, Racine CH, Tape KD (2001a) Increasing shrub abundance in the Arctic. Nature, **411**,

595 546–547.

596

597 Sturm M, McFadden JP, Liston GE, Chapin FS, Racine CH, Holmgren J (2001b) Snow-shrub

598 interactions in Arctic tundra: a hypothesis with climatic implications. Journal of Climate, **14**,

599 336-344.

600

601 Tape KD, Sturm M, Racine CH (2006) The evidence for shrub expansion in Northern Alaska and

602 the Pan-Arctic. Global Change Biology, **12**, 686–702.

603

604 Tape KD, Lord R, Marshall HP, Ruess RW (2010) Snow-mediated ptarmigan browsing and
605 shrub expansion in Arctic Alaska. *Ecoscience*, **17**, 186-193.

606
607 Tape KD, Christie K, Carroll G, O'Donnell JA (2016) Novel wildlife in the Arctic: the influence
608 of changing riparian ecosystems and shrub habitat expansion on snowshoe hares. *Global Change*
609 *Biology*, **22**, 208-219.

610
611 Tremblay B, Lévesque E, Boudreau S (2012) Recent expansion of erect shrubs in the Low Arctic:
612 evidence from Eastern Nunavik. *Environmental Research Letters*, **7**, 035501.

613
614 Williamson SN, Barrio IC, Hik DS, Gamon JA (2016) Phenology and species determine
615 growing-season albedo increase at the altitudinal limit of shrub growth in the sub-Arctic. *Global*
616 *Change Biology*, doi: 10.1111/gcb.13297.

617
618 Wilmking M, Hallinger M, Van Bogaert R *et al.* (2012) Continuously missing outer rings in
619 woody plants at their distributional margins. *Dendrochronologia*, **30**, 213-222.

620

Tables

Table 1 Results from the model selection using Akaike information criterion. The sensitivity (i.e. difference between the best and the null model, in terms of ΔAIC), the marginal R^2 (i.e. the amount of variance explained by fixed effects) and the conditional R^2 (i.e. the amount of variance explained by both fixed and random effects) are shown only for the more plausible model(s) for stem and collar in each type of environment. T and P stands for temperature and precipitation, respectively.

Site	Shrub part	Best climate model	Sensitivity (ΔAIC_{null})	Marginal R^2	Conditional R^2
Snowbed	Root collar	July T + March P	19.0	0.29	0.53
	Stem	August T	7.1	0.15	0.31
Hilltop	Root collar	July T + March P	16.2	0.13	0.24
	Stem	May T	2.2	0.02	0.09
Terrace	Root collar	July and August T	12.2	0.07	0.13
		July T + March P	11.8	0.06	0.12
	Stem	April T	3.8	0.04	0.15

Table 2 Information on individual root collar and stem samples in three different types of environments in the Boniface River region, Nunavik, Québec, Canada. The stem length refers to the measure (in cm) from the root collar to the tip of the stem. The numbers in brackets (second column) represent the sample size.

		Age		Length	
		Mean value (years)	SD	Mean value (cm)	SD
Root collar	Snowbed (15)	35.3	17.5	-	-
	Hilltop (16)	45.2	20.7	-	-
	Terrace (15)	43.5	21.1	-	-
	<i>Mean value</i>	41.4	20.5	-	-
Stem	Snowbed (30)	17.9	9.5	146.8	31.1
	Hilltop (32)	26.9	14.5	92.4	16.4
	Terrace (30)	25.3	12.2	101.8	21.3
	<i>Mean value</i>	23.4	13.0	109.1	28.9

Figure captions

Figure 1 (a) Map of shrub sampling methods for dendroclimatological analyses in northern hemisphere. The black star represents our study area near the Boniface River research station, Nunavik, Québec, Canada; darker dots represent regions for which more than one study took place. (b) General landscape around the Boniface River research station, with extensive *Betula glandulosa* thickets. (c) An uprooted *B. glandulosa* individual with large proportion of its stems and roots removed. The location of the root collar, roots and stems are shown.

Figure 2 Differences in climate sensitivity of growth for March precipitation (a – c) and July temperature (d – f), the two best climate models, between stems (red) and collars (blue). Solid lines and shaded areas indicate the slope estimates and 95% confidence interval for the mixed model analyses.

Figure 3 Sensitivity of collar and stem growth to monthly (a) precipitation and (b) temperature, as expressed by the difference between the AIC of each linear models and the AIC of the null model. The metric "sensitivity" was obtained by multiplying the ΔAIC_{null} by +1 or -1 depending on the sign of the slope of the regression to represent positive and negative growth responses, respectively. "P" preceding a month stands for previous year.

Figure 4 The linear model results in climate sensitivity for each individual chronology, for both stems (red) and root collars (blue). Climate sensitivity to July temperature (a, c, e) and March precipitation (b, d, f) are shown for individuals growing on snowbed (a, b), hilltop (c, d) and

659 terrace (**e**, **f**). Bars are the slope and standard error (linear models). Filled bars indicate regression
660 slopes that are significantly different from 0 ($p < 0.05$).