THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Synaptic integrative mechanisms for spatial cognition

Citation for published version:

Schmidt-Hieber, C & Nolan, M 2017, 'Synaptic integrative mechanisms for spatial cognition', *Nature Neuroscience*. <https://doi.org/10.1038/nn.4652>

Digital Object Identifier (DOI):

[10.1038/nn.4652](https://doi.org/10.1038/nn.4652)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Nature Neuroscience

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Title: Synaptic integrative mechanisms for spatial cognition

Author names: Christoph Schmidt-Hieber¹ & Matthew F Nolan²

Affiliations: 1. Institut Pasteur, Paris, France. Email: christoph.schmidt-hieber@pasteur.fr
2. Centre for Integrative Physiology, University of Edinburgh, Edinburgh. Email: mattnolan@ed.ac.uk

Abstract

Synaptic integrative mechanisms have profound impacts on electrical signaling in the brain that, while largely hidden from recording methods that observe spiking activity of neurons, may be critical for how information is encoded, stored and retrieved. Here, we review roles for synaptic integrative mechanisms in selection, generation and plasticity of place and grid fields, and in related temporal codes for representation of space. We outline outstanding questions and challenges in testing hypothesized models for spatial computation and memory.

Main text

The spatial firing patterns of neurons in the hippocampal formation are central to neurobiological theories of spatial cognition^{1,2}. How the spatial modulation of place, grid, head-direction, border and other spatial cell types emerges from their synaptic input, while largely hidden from view when observing spike firing, is likely to be critical for spatial computation. In simple models the details of this process of synaptic integration are of limited importance - spike output is assumed to be a stable, linear function of synaptic input. In contrast, considerable experimental evidence demonstrates that synaptic integration is often non-linear, can be spatially compartmentalised within a cell and is controlled by diverse mechanisms, suggesting it has key computational roles³⁻⁵. Here, we consider evidence that specific mechanisms for integration of synaptic input are critical for spatial cognition. We will focus on aspects of hippocampal spatial firing fields and temporal codes for which recent experiments give insights into roles of these integrative mechanisms.

What cellular mechanisms does a neuron have available to determine integration of its synaptic input? To influence spatial firing a synaptic event must influence action potential initiation. Several cellular properties determine the impact of a synaptic event³⁻⁵. First, neuronal excitability is established by ion channels that set a neuron's resting membrane potential, voltage threshold for triggering an action potential and membrane conductance. The difference between the resting potential and threshold potential gives the voltage change that must be achieved to trigger a spike. The membrane conductance, in conjunction with the capacitance established by the lipid bilayer of the cell membrane, determines how easily and rapidly a synaptic input can change the membrane potential. Second, various voltage-dependent ionic currents, including those mediated by Na⁺, Ca²⁺, and NMDA receptor (NMDAR) channels, can amplify synaptic responses, while other ion channels and inhibitory synaptic receptors suppress synaptic responses. Third, most synaptic inputs are made onto dendrites, which can extend

hundreds of microns from a neuron's soma. All other things being equal, more distant synapses are less effective because attenuation of synaptic responses increases as they propagate further. Finally, spatially extended dendrites enable compartmentalisation. For example, different dendritic domains may be endowed with distinct combinations of voltage-gated ion channels and the particular signalling mechanisms that modify synaptic responses may be directed to specific locations.

These synaptic integrative mechanisms suggest considerable cellular complexity, but why should we consider them when trying to understand spatial computation? Powerful artificial neural networks can be assembled from simple neurons that linearly sum their synaptic inputs⁶. Why then do networks for spatial cognition employ neurons with diverse and complex integrative properties? One possibility is that specialisation enables neurons to adapt to fundamental limits imposed by their cellular hardware⁷. For example, to integrate many synaptic inputs a neuron requires an extensive dendritic tree, but this comes at a cost in that distal inputs will evoke smaller and less temporally precise somatic responses. This cost can be compensated by integrative mechanisms that boost the strength of distal synapses and that normalise their time course at the soma. A second possibility is that diversity in subthreshold properties reflects the selection of distinct building blocks for specialised computations^{5,7}. According to this view, specific mechanisms for synaptic integration may be necessary to the cognitive function implemented in a circuit. While these viewpoints are not mutually exclusive, we will focus here primarily on computational roles for synaptic integrative mechanisms.

For which aspects of spatial cognition might synaptic integrative mechanisms be important? We will address roles in key elements of spatial computation in the hippocampus and medial entorhinal cortex (MEC). We focus on hippocampal place and entorhinal grid cells, which use their spike firing rate to encode locations with a high signal to noise ratio; firing within fields usually peaks at frequencies > 10 Hz, whereas firing rates outside of fields are less than 1 Hz^{8,9}. We will first consider recent evidence that integrative mechanisms determine selection of active place cells and how this might form a basis for allocation of memory engrams.

In both place and grid cells the approximate Gaussian firing rate distribution of a single firing field is driven by a ramp-like membrane potential depolarization (Fig. 1a-b)¹⁰⁻¹². This is at first glance consistent with models in which straightforward linear integration of excitatory drive is sufficient to explain place firing¹³⁻¹⁵. However, more recent observations that we discuss below argue that active integrative mechanisms are essential for the emergence of ramp-like depolarizations driving place fields, and may influence the spacing and stability of grid fields.

Beyond moment to moment computation, synaptic integrative mechanisms may contribute to spatial memory by influencing the induction of synaptic plasticity. Specifically, encoding and recall of spatial memories are associated with plasticity in the spatial firing of hippocampal neurons^{16,17}. A critical issue is how patterns of spatially modulated synaptic input couple appropriately to plasticity mechanisms. We will consider evidence that synaptic integrative mechanisms establish rules for plasticity of spatial firing, by both promoting and suppressing synaptic plasticity.

Finally, the relative timing of action potentials fired by place and grid cells may be of particular importance for spatial memories. In particular, phase precession of action potentials relative to the network theta rhythm (Fig. 1c) leads to the emergence of population level spike sequences that may be structured to support associative memory storage^{18–21}. We will consider how synaptic integrative mechanisms may contribute to temporal codes that are linked to theta activity and that may be important for episodic memory.

Excitability, place cell selection and memory allocation

In a given environment, only subsets of CA1 pyramidal cells have place fields. Estimates range from as low as 20% rising to 65% for larger environments^{22–24}. While additional cells are recruited to encode larger environments, the number of silent cells and the number of cells with multiple firing fields is greater than expected if all cells have a similar probability of generating a place field²³. Rather, the probability that cells will have place fields is described by a gamma distribution, suggesting a population-level code for environmental context²³. Such a code requires mechanisms to determine which cells within the population become active. Recent studies point to pyramidal cell excitability as critical for selection of active cells and suggest how this population-level code may contribute to temporal components of episodic memories.

How are active place cells selected? Differences in excitable properties are an attractive candidate mechanism, but it has been difficult to directly relate excitable integrative properties of neurons to their firing fields during a behaviour. Technically demanding experiments involving patch-clamp recordings from behaving rodents of the membrane potential of CA1 pyramidal cells, and simultaneous measurement and manipulation of their electrical integrative properties, have met this challenge. These studies reveal two differences in excitability between silent cells and CA1 pyramidal cells that go on to have place fields in a novel environment; future place cells have a lower threshold for action potential firing and a greater likelihood of firing bursts of action potentials^{25,26} (Fig. 2a). By making action potential firing in response to synaptic input more likely, both differences should promote the emergence of firing fields. Therefore, whether a pyramidal cell becomes a place cell may in part be determined a priori by its intrinsic electrical properties.

What consequences might a priori selection of place cells have for memory functions of the hippocampus? Selection of active cells through differences in excitability has been suggested to underpin temporal features of episodic memories²⁷. Elegant investigations of memory allocation in the amygdala provide support for this general idea^{28–30}, but whether hippocampal-dependent memories employ similar mechanisms has only recently been explored. When the activity of populations of CA1 pyramidal cells are imaged in the same environment over multiple days, the ensemble representation slowly evolves; some cells leave the ensemble, whereas others join^{31,32}. If this subset of place cells is predetermined by differences in their excitability, then memories formed on the same day are likely to be allocated to overlapping groups of neurons, whereas memories formed on different days should be allocated to different populations. Consistent with this prediction, cell populations tagged with a genetically encoded activity-

dependent reporter in a first context overlapped with cells labelled by a second activity-dependent marker following exposure to a second context several hours later, but not several days later³². Moreover, memories formed by exposure to contexts several hours apart, but not several days apart, interact with one another³². Thus, the temporal properties of active place cell assemblies, and of contextual memory storage, are consistent with there being an active subset of excitable CA1 pyramidal cells that changes over a time-scale of days.

Selection of active place cells provides a potentially powerful mechanism for encoding temporal components of memory, but what drives the subset of active neurons to change over time? One possibility is that during memory formation neuronal activity leads in itself to transient (hours long) increases in excitability, providing a mechanism for association of a second memory formed within a time window defined by increases in excitability^{32,33}(Fig. 2b). This general scheme is supported by investigations in which virally mediated expression, in subsets of amygdala neurons, of the transcription factor CREB increases their excitability causing them to be selected to participate in the engram of fear memories^{28,29}. Evidence that in hippocampal neurons synaptic activity or spike firing activate CREB^{34,35}, and that CREB activation increases excitability³⁶, is consistent with this idea. In this scheme, later periods of lowered activity that facilitate memory dissociation may be established by self-regulatory mechanisms that come into play after initial activation of CREB^{27,30}. A complementary possibility is that the identity of active subsets of neurons provides a code from which the timeline of events can be read out³¹ (Fig. 2c). This idea is supported by observations that ensemble place field maps of different environments on the same day share representations, and that decoders trained on one environment can infer the day on which ensemble patterns were recorded from a second environment³¹. According to this view, the active subset of place cells could be established independently from neural activity, either through network wide coordination of the excitable set of CA1 pyramidal cells, or perhaps through stochastic switching of CA1 pyramidal cells between more and less excitable states.

Together these observations are consistent with excitability of CA1 pyramidal neurons selecting place cell firing and memory allocation. Nevertheless, important questions remain to be addressed. What is the ionic mechanism that controls which cells become excitable? The difference in spike threshold between place cells and inactive CA1 pyramidal cells points to voltage-gated ion channels that control action potential initiation^{25,26}. In contrast, activation of CREB^{28,36}, and recent learning³⁷, both reduce afterhyperpolarization currents in CA1 neurons. These differences may reflect multiple mechanisms acting across different timescales. Does excitability predict ensemble membership over days? Correlating changes in a cell's excitability with its firing fields will be critical here. Is the probability of a CA1 pyramidal cell forming a place field a cell autonomous property, or does it depend on whether other cells are active? When excitability of subsets of cells in the amygdala is increased, these cells predominate in engrams that are formed, but the overall number of engram cells does not increase, implying that the proportion of cells that form an engram is fixed by reciprocal inhibition³⁰. Similar mechanisms may be present in hippocampal circuits (e.g. ^{38,39}). Does selection of place cells through differences in excitability extend to other hippocampal areas? Unlike CA1, ensemble codes in CA3 appear to be stable over days⁴⁰, whereas ensemble codes in CA2 evolve even more

rapidly than in CA1⁴¹. If stability of intrinsic excitability is used for place cell selection, then we expect this to be reflected in differential control of excitability in each area.

Membrane potential dynamics driving spatial firing

How are synaptic inputs converted into action potential outputs that form a neuron's spatial firing field? In vitro studies demonstrate that dendritic active conductances can either amplify or suppress synaptic responses in hippocampal neurons (e.g. ^{42–45}). Recent experiments probing the membrane potential of spatial cells in awake animals, in the real world and using virtual environments, show that ramp-like depolarizations drive spatial firing and have begun to reveal roles for synaptic integrative mechanisms.

Membrane potential ramps in CA1 pyramidal cells. CA1 pyramidal cells provide a striking example of how computation emerges through interaction between synaptic integrative mechanisms, neuronal morphology and circuit connectivity. Excitatory inputs from layer 3 of entorhinal cortex target distal dendrites of CA1 pyramidal cells, whereas local inputs from CA3 target their proximal dendrites⁴⁶ and diverse interneuron populations provide spatially restricted inhibition⁴⁷. Either excitatory pathway appears to be sufficient to drive place firing^{48,49}, and active integrative mechanisms may control responses to either or both pathways^{3,50–53}. How then is spatial firing in place cells shaped by active synaptic integration?

Direct evidence that non-linear integrative mechanisms contribute to place firing comes from experiments in which the membrane potential of silent CA1 pyramidal cells was continuously depolarized while rats navigated an oval track⁵⁴. This manipulation caused place cells to emerge. Importantly, the location of the induced field could not be predicted from the membrane potential prior to injection of the depolarizing current. This finding argues against simple models for the membrane potential ramp in which synaptic inputs within the firing field are stronger than those outside the field, as these models predict that prior to continuous depolarization there should be subthreshold ramps at the location of the firing field (Fig. 3a). Instead, in these experimental conditions the emergence of place fields appears to be determined by a voltage-dependent gating mechanism.

What might be the nature of this mechanism? One possibility is that excitatory synaptic inputs within the field are indeed stronger than those outside, but that in silent cells the depolarization they generate is insufficient to produce a measureable change in the somatic membrane potential (Fig. 3b). This implies substantial attenuation of EPSPs as they propagate along dendrites towards the soma, as has for example been reported for the basal dendrites of neocortical pyramidal cells⁵⁵. In this scenario, continuous somatic depolarization may activate voltage-dependent dendritic Na⁺, Ca²⁺, or NMDAR channels to amplify the local EPSPs, or cause inactivation of K⁺ channels that would otherwise suppress EPSPs, either way enabling the EPSPs to propagate to the soma. Another possibility is that synapses active within the firing field have similar strength to those outside, but face a lower threshold to engage amplifying dendritic conductances (Fig. 3c). This situation may be favored by clustering of synapses with similar spatial preferences onto CA1 pyramidal cell dendrites⁵⁶. Although the signals encoded

by individual synapses on place cell dendrites are not yet clear, recent *in vivo* spine imaging studies in visual cortex support the hypothesis that functionally similar inputs preferentially target nearby locations on the dendritic tree of a neuron^{57,58}. Future experiments might address this by imaging, during behavior, of synaptic terminals on identified place cells.

The voltage dependence of firing fields that emerges during prolonged membrane potential depolarization provides strong evidence for functional engagement of integrative mechanisms during place cell firing. However, whether active integrative mechanisms are also essential for the generation of place fields under more physiological conditions is unknown. Recent experiments in which the membrane potential of place cells was recorded in novel and familiar environments suggest that new place fields emerge in the absence of a sustained depolarization²⁶. Whether place fields in these conditions require voltage-dependent gating mechanisms is not yet clear.

Regardless of the role of active integration, additional mechanisms are likely to shape the membrane potential ramp driving place firing. For example, the membrane potential ramps underlying receptive fields in other brain regions are substantially shaped by synaptic inhibition⁵⁹. Specifically, orientation tuning curves in visual cortex are transformed linearly, with a threshold, by inhibition from parvalbumin expressing interneurons^{60,61}. Input from local inhibitory interneurons to CA1 place cells affects the shape of sub- and suprathreshold place fields in a strikingly similar manner (Fig. 4), likely by suppressing firing and opposing active mechanisms that amplify synaptic responses outside of the place field⁶². In visual cortex, different interneuron subpopulations are thought to play specific roles in shaping orientation selectivity^{63–65}. Similarly, the effects of inhibition on the rising and falling parts of the place field ramp may be respectively mediated by parvalbumin and somatostatin expressing interneurons⁶⁶.

Membrane potential ramps in medial entorhinal cortex. While grid firing fields of entorhinal neurons are also driven by slow ramp-like depolarizations^{11,12}, the underlying integrative mechanisms may be fundamentally different. For example, in contrast to hippocampal place cells, the relative location of grid cell firing fields is stable across environmental manipulations, suggesting circuit level interactions constrain grid cell firing fields^{67,68}. At the cellular level, the dendritic morphology of hippocampal pyramidal cells differs substantially from stellate and pyramidal cells in layer 2 of the MEC^{69,70}. Proposed mechanisms for generation of ramp depolarizations also differ. Thus, the ramp depolarization recorded from grid cells is consistent with predictions of continuous attractor network models^{11,12}. When these models are implemented so that they reflect evidence that stellate cells interact primarily via local inhibitory neurons^{71,72}, they predict that the depolarizing ramp results from disinhibition⁷².

Although network mechanisms are good candidates for generation of the membrane potential ramp underlying grid firing, there is evidence that synaptic integrative mechanisms contribute to the spacing and stability of grid fields. First, deletion of HCN1 channels, which mediate a major component of the hyperpolarization-activated currents (I_h) in entorhinal stellate cells⁷³, increases the width and spacing of grid cell firing fields⁷⁴. I_h is a mixed Na^+ and K^+ current that is unusual in that it is activated by membrane hyperpolarization⁷⁵. Along with leak K^+ channels, I_h

generates a dorsoventral gradient in synaptic integration by stellate cells⁷⁶. At more dorsal locations, where grid cells have closely spaced firing fields, a high density of both currents reduces the width of synaptic potentials and opposes their temporal summation, whereas at more ventral locations where grid cells typically have widely spaced firing fields, synaptic potentials are broader and temporal summation is greater because the density of each current is lower⁷⁶. Gradients in I_h are also associated with dorsoventral differences in intrinsic oscillatory properties of stellate cells⁷⁷, which we discuss further below. Second, entorhinal stellate and pyramidal cells are endowed with active conductances that produce a supralinear transformation of synaptic inputs into action potential output⁷⁸. Simulations suggest that a slow, NMDAR-mediated supralinear integration mechanism can promote the robustness of the grid cell rate code. While direct recordings of NMDAR-mediated responses have not yet been obtained from grid cells *in vivo*, NMDARs have been shown to be engaged during behaviour in other brain regions⁵, where they contribute to receptive field tuning of somatosensory^{79,80} and visual responses⁸¹.

What are the implications of these biophysical data for computations carried out by place and grid cells? In place cells, non-linear synaptic integrative mechanisms may enable gating of place firing⁵⁴, and maximize memory storage capacity⁸². For grid cells, differences in grid scale may maximise the representational capacity of grid networks⁸³, but whether dorsoventral tuning of synaptic integration plays a necessary or a modulatory role is unclear.

Active synaptic integration and plasticity of spatial representations

Successful learning requires plasticity of behaviorally relevant connections between neurons, which in the case of spatial memory is thought to lead to stabilisation of place fields^{83–85}. Considerable evidence supports a necessary role for NMDAR-dependent synaptic plasticity in this process⁸⁵. For example, pharmacological and genetic manipulations of NMDARs disrupt long-term potentiation (LTP) of synaptic responses⁸⁵, spatial learning^{86,87}, the stability of place cells⁸⁸, and spatial representation by place cells^{87,89}. Plasticity driven by activation of voltage-gated Ca^{2+} channels may also play important roles (e.g. ⁹⁰). By determining the effects of synaptic inputs on the membrane potential, active synaptic integration may interact with several proposed mechanisms for recruitment of NMDARs and voltage-gated Ca^{2+} channels to either facilitate or oppose induction of synaptic plasticity.

Synaptic plasticity during place field formation and stabilization. If NMDARs are indeed instrumental for the stabilization of place cells, the Ca^{2+} influx that is associated with postsynaptic depolarization and NMDAR channel opening should be detectable in the dendritic tree during crossings of future or existing place fields. In support of this hypothesis, regenerative Ca^{2+} events occur in basal dendritic branches during place field crossings and are associated with the precision and stability of place fields⁹¹, suggesting that they represent postsynaptic plasticity signals. A second type of regenerative calcium event is generated in the apical dendrites of CA1 pyramidal cells. Precisely timed, coincident entorhinal cortex and CA3 inputs evoke NMDAR-dependent dendritic plateau potentials *in vitro* and *in vivo*^{92,93} that can trigger synaptic plasticity at least *in vitro*⁹². These complex spikes are associated with stabilization of

membrane potential maps in novel environments²⁶. However, while evoked plateau potentials may be sufficient to induce place fields under some conditions⁹³, they do not appear to be necessary for new place field generation in novel environments²⁶.

Further clues to the forms of plasticity promoting place field formation and stability come from intracellular recordings from CA1 pyramidal cells in novel and familiar virtual environments²⁶. In these experiments place field formation appears not to require firing of action potentials, suggesting that the initial place field ramp is generated by sub-threshold forms of plasticity^{94,95}. For example, isolated dendritic spikes in conjunction with presynaptic activity are sufficient to induce LTP of the CA3 input to CA1 pyramidal cells⁹⁴. This form of spike-independent, localized plasticity could explain why place cells appear rapidly in a novel environment^{96,97}. A similar spike-independent LTP mechanism has also been described for CA3 pyramidal cells: powerful proximal inputs from mossy fiber axons can induce synaptic plasticity even in the absence of postsynaptic somatic spikes^{98,99}. This may enable sparse inputs from dentate gyrus granule cells to efficiently generate active assemblies of CA3 pyramidal cells.

Constraints on synaptic plasticity. Distinct and spatially localised integrative mechanisms may oppose synaptic plasticity. For example, HCN1 channels, which are highly enriched in the distal dendrites of CA1 pyramidal neurons¹⁰⁰, suppress LTP of distal synaptic inputs⁵¹. By depolarizing distal dendrites HCN1 channels prevent synaptically driven calcium transients mediated by T-type Ca²⁺ channels, suggesting a mechanism to account for their actions on LTP⁵². At a behavioural level deletion of HCN1 from forebrain neurons enhances hippocampal-dependent forms of learning⁵¹, and increases the size and stability of CA1 place cell firing fields¹⁰¹. Conversely, cannabinoid mediated enhancement of HCN1 channels reduces LTP and suppresses hippocampal-dependent learning¹⁰². Together, these observations reinforce the idea that compartmentalisation of synaptic integration contributes to spatial computations, and suggest that HCN1 channels in distal dendrites control spatial firing and memory by gating plasticity of direct cortical inputs.

A challenge in establishing roles of synaptic integrative mechanisms in memory is that the ion channels implicated in control of synaptic plasticity may also influence membrane potential ramps that drive spatial firing fields. For example, while HCN1 channels oppose distal synaptic plasticity through their contribution to the resting membrane potential, HCN1 channels also affect the waveform and temporal summation of distally originating post-synaptic potentials as they propagate to the soma^{43,51,103}. Similarly, voltage-dependent gating of NMDARs contributes directly to postsynaptic integration as well as providing a Ca²⁺ source for induction of plasticity.

Theta oscillations and temporal codes

The rate coded representations provided by place and grid fields are multiplexed with codes that represent location through the timing of action potentials relative to the network theta (4-10 Hz) rhythm^{20,104,105}. The theta rhythm is entrained by GABAergic projections from the medial septum to interneurons in the hippocampus and entorhinal cortex^{106–110}. Because the relative delay between theta cycles in the hippocampus and entorhinal cortex is greater than expected from

the synaptic delays between each area, the theta rhythm may establish temporal windows for local circuit interactions¹¹¹. We will focus here on mechanisms by which the hippocampal formation responds to theta modulated inputs and generates population level theta sequences.

Responses to theta modulated signals. How do neurons in the hippocampus and entorhinal cortex respond to theta frequency synaptic inputs? The membrane potential response of hippocampal CA1 pyramidal cells and stellate cells in the MEC to oscillating current inputs are largest for oscillation frequencies in the theta range, whereas fast spiking interneurons may prefer higher input frequencies^{112–115}(Fig. 5a). At resting potentials the theta frequency selectivity, or resonance, of pyramidal and stellate cells requires HCN1 channel mediated I_h currents^{51,73,113,116}, whereas at depolarized potentials around spike threshold M-type K^+ channels appear to be critical^{116,117}. Relatively slow voltage-dependent gating of both types of ion channel leads to the appearance of resonance by opposing responses to input currents with frequencies < 5 Hz. Resonance mechanisms directly affect spike output, by causing neurons to generate greater numbers of spikes in response to inputs active near a cell's resonant frequency¹¹², and may also modify the timing of action potentials driven by synaptic inputs at different phases of the theta cycle¹¹⁸.

Do these single cell resonance phenomena manifest in vivo? Two lines of evidence suggest that HCN1-dependent resonance is engaged during theta states. First, the amplitude of theta frequency field potential oscillations recorded from CA1 is increased following genetic deletion of HCN1^{51,101}. This is consistent with models of the contribution of dendritic HCN channels to the local field potential¹¹⁹. Second, in behaving animals CA1 pyramidal cells respond preferentially to activation of PV interneurons at theta frequencies¹²⁰ (Fig. 5b). This resonance effect is abolished by pharmacological block of HCN channels¹²⁰. Interestingly, pyramidal cells did not show theta frequency resonance upon direct optogenetic activation, suggesting that HCN channel-dependent resonance, which would more effectively be engaged by hyperpolarizing inhibition, may be more prevalent in behaving animals than peri-threshold resonance, which in vitro does not require HCN channels¹¹⁶.

How does membrane potential resonance affect spatial computation? Ion channels contributing to membrane potential resonance participate in sub-threshold theta frequency membrane potential oscillations observed during in vitro recordings^{121,122}. These intrinsic oscillations have been suggested to contribute to rate and temporal codes through oscillatory interference mechanisms^{77,105,123}. However, this intrinsic oscillatory activity is suppressed by background synaptic activity¹²⁴ and has not been observed in recordings from hippocampal or entorhinal neurons in awake animals^{10–12,25}. Alternatively, by filtering signals with frequency outside the theta band, resonance mechanisms may promote the emergence of temporal computations within windows defined by theta oscillations¹¹¹.

Theta phase precession and theta sequences. As an animal moves through a cell's firing field, an advance in the timing of the cell's action potentials relative to the theta rhythm (phase precession) leads to the emergence of population level theta sequences²⁰. How theta rhythms interact with synaptic inputs to cause phase precession and sequences is unresolved¹²⁵.

Several classes of model include components implemented by active integration mechanisms (e.g.^{21,105,126–128}). For example, ion channels that mediate spike frequency adaptation promote symmetry of firing in models in which phase precession involves asymmetric ramp-like synaptic inputs^{10,127,129,130}. In these models place fields are driven by an input current that rises slowly and falls rapidly, with adaptation causing the spike rate to fall before the peak of the ramp. In a detailed pyramidal cell model, sub-threshold membrane potential oscillations and resonance promote phase precession that is generated by shifting the balance between oscillating excitatory and inhibitory synaptic inputs¹²⁶. In simulations of grid cell firing a fast supralinear dendritic integration mechanism sharpens phase precession by restricting time windows for spike firing⁷⁸.

Experiments that focus on model predictions at the level of synaptic integration may help distinguish between these various models and test roles for phase precession in spatial behaviors. Because numerous models generate phase precession, an approach to evaluate model predictions may be to also consider dependence of spatial codes on factors in addition to location¹³¹. In this spirit, a recent model suggests integrative properties may be tuned to account for dorsoventral differences in theta phase, and to maintain phase precession when running speed varies²¹. While experiments with knockout mice show that HCN1 channels are not required for either theta oscillations or for phase precession^{51,101,132}, identifying which ion channels do play roles in phase precession may provide targets for testing contributions of theta sequences to spatial memory.

Concluding remarks

The phenomena of rate and temporally coded spatial firing, when considered at the level of membrane potential dynamics, appear to arise from complex and multi-layered mechanisms, with synaptic integration playing critical roles at multiple key points. For example, in CA1 pyramidal cells, active integrative mechanisms contribute to selection of active cells, membrane potential dynamics driving firing, plasticity of synaptic inputs, and responses to oscillatory network activity. Entorhinal grid cells also appear to engage specific integrative mechanisms, but intriguingly these may be distinct from those used in CA1. In our view, experiments to date may only be scratching the surface of a rich diversity of dynamic integrative mechanisms underlying the well defined spatial firing properties of neurons in the hippocampal formation. We end by outlining areas that may be of importance to future investigation of mechanisms for spatial computation in these circuits.

Spatial cognition involves numerous cell types not considered here. For example, principal cells in CA3 and the dentate gyrus employ distinct integrative mechanisms. Dendritic regenerative events can be readily evoked in CA3 pyramidal neurons^{133,134}. In contrast, while distinct regenerative events have not been observed in direct recordings from granule cell dendrites¹³⁵, their NMDAR-dependent sensitivity to sequences of synaptic inputs¹³⁶ and pronounced dendritic Ca^{2+} transients during backpropagating action potentials¹³⁷ suggest that nonlinear dendritic conductances can be recruited. In agreement with this view, selective deletion of NMDARs in the dentate gyrus causes deficits in rapidly producing a unique memory of a novel context, and

discriminating it from previously encountered contexts¹³⁸. The dendrites of granule cells may support this “pattern separation” function by increasing the sparsity of firing; assuming the same number and weights of synaptic inputs, a granule cell is less likely to fire a spike if it has more dendrites¹³⁹. This sparsification of firing may further be enhanced by short coincidence detection windows for EPSPs in granule cell dendrites¹⁴⁰.

How synaptic integrative properties contribute to the spatial codes of border, head-direction and other spatially modulated neurons is an open target for future investigation. Likewise, network activity patterns such as sharp wave ripples and associated spike sequences, and gamma oscillations that co-occur with theta states, may also be shaped by active synaptic integrative mechanisms. For example, during sharp wave ripples synaptic inhibition may dynamically re-configure synaptic integration by CA1 pyramidal cells^{141,142}.

Synaptic integrative properties are dynamically regulated by neuromodulatory systems according to brain state and behavioural demands¹⁴³. Ion channels that mediate integration of synaptic responses are prime targets for neuromodulators¹⁴⁴, raising questions about how these systems influence spatial computations in the behaving animal. Recent studies indicate highly selective roles for certain neuromodulators. For example, cholinergic inputs increase excitability of dentate gyrus granule cells by stimulating interactions between axonal T-type Ca^{2+} channels and Kv7 channels¹⁴⁵. At a systems level, computational models incorporating neuromodulatory systems provide frameworks for predicting how modulation of ion channels important for synaptic integration contributes to circuit computations and behaviour¹⁴⁶. Future investigation of interactions between neuromodulation and synaptic integration may lead to important insights into control mechanisms for spatial cognition.

New tools will be critical to the twin challenges of selective experimental manipulation of synaptic integrative mechanisms and observation of the subcellular membrane potential dynamics on which they act. Promising strategies for manipulation include optical control of native or engineered light-sensitive ion channels¹⁴⁷. For example, rapid optical block of HCN1 channels in distal dendrites of CA1 pyramidal cells may help resolve the question of whether their influence on spatial firing fields is through control of synaptic plasticity, or by effects on the waveform of synaptic responses that propagate to the soma. Novel imaging approaches, including miniaturization of microscope technologies¹⁴⁸ and development of fluorescent voltage-sensors¹⁴⁹ will facilitate exploration of the impacts of synaptic integrative mechanisms on sub-cellular and network level computations. For example, measurement of resting membrane potential and spike threshold across populations of neurons in behaving animals will facilitate direct testing of the contributions of excitability to selection of active place cells.

Finally, it is intriguing to consider whether synaptic integrative mechanisms contributing to spatial computations are similarly used in other neural systems. On the one hand, the evidence we have considered points towards diversity in strategies for synaptic integrative mechanisms to influence neural computation. On the other hand, nonlinear synaptic integrative mechanisms engaged to drive receptive fields in visual cortex⁸¹ appear similar to those used by CA1 pyramidal cells. As we highlight above, the effect of inhibition on the shape of sub- and

suprathreshold spatially receptive fields in place cells^{62,66} also bears resemblance with observations from orientation-sensitive neurons in visual cortex^{60,63,64}. Establishing how common synaptic integrative mechanisms are adapted to the specific computations carried out by different circuits would be a major achievement for cellular and systems neuroscience.

Acknowledgements

We thank Nathalie Rochefort and Gülşen Sürmeli for comments on the manuscript, and Albert Lee for sharing data prior to publication (ref. 25). This work was funded in part by grants to M.F.N. from the BBSRC (BB/M025454/1 and BB/L010496/1), Human Frontiers Science Program (RGP0062/2014), Wellcome Trust (200855/Z/16/Z), and Simons Initiative for the Developing Brain, and to C.S.-H. from the ERC (StG 678790 NEWRON).

References

1. O'Keefe, J. & Nadel, L. *The hippocampus as a cognitive map*. (Oxford: Clarendon Press, 1978).
2. Redish, A. D. *Beyond the Cognitive Map: From Place Cells to Episodic Memory*. (MIT Press, 1999).
3. Spruston, N. Pyramidal neurons: dendritic structure and synaptic integration. *Nat. Rev. Neurosci.* **9**, 206–221 (2008).
4. Häusser, M., Spruston, N. & Stuart, G. J. Diversity and dynamics of dendritic signaling. *Science* **290**, 739–744 (2000).
5. Chadderton, P., Schaefer, A. T., Williams, S. R. & Margrie, T. W. Sensory-evoked synaptic integration in cerebellar and cerebral cortical neurons. *Nat. Rev. Neurosci.* **15**, 71–83 (2014).
6. LeCun, Y., Bengio, Y. & Hinton, G. Deep learning. *Nature* **521**, 436–444 (2015).
7. Häusser, M. & Mel, B. Dendrites: bug or feature? *Curr. Opin. Neurobiol.* **13**, 372–383 (2003).
8. O'Keefe, J., Burgess, N., Donnett, J. G., Jeffery, K. J. & Maguire, E. A. Place cells, navigational accuracy, and the human hippocampus. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **353**, 1333–1340 (1998).

- 517 9. Hafting, T., Fyhn, M., Molden, S., Moser, M.-B. & Moser, E. I. Microstructure of a spatial
518 map in the entorhinal cortex. *Nature* **436**, 801–806 (2005).
- 519 10. Harvey, C. D., Collman, F., Dombeck, D. A. & Tank, D. W. Intracellular dynamics of
520 hippocampal place cells during virtual navigation. *Nature* **461**, 941–946 (2009).
- 521 11. Schmidt-Hieber, C. & Häusser, M. Cellular mechanisms of spatial navigation in the medial
522 entorhinal cortex. *Nat. Neurosci.* **16**, 325–331 (2013).
- 523 12. Domnisoru, C., Kinkhabwala, A. A. & Tank, D. W. Membrane potential dynamics of grid
524 cells. *Nature* **495**, 199–204 (2013).
- 525 13. Hartley, T., Burgess, N., Lever, C., Cacucci, F. & O'Keefe, J. Modeling place fields in terms
526 of the cortical inputs to the hippocampus. *Hippocampus* **10**, 369–379 (2000).
- 527 14. Burgess, N. & O'Keefe, J. Models of place and grid cell firing and theta rhythmicity. *Curr.*
528 *Opin. Neurobiol.* **21**, 734–744 (2011).
- 529 15. Solstad, T., Moser, E. I. & Einevoll, G. T. From grid cells to place cells: a mathematical
530 model. *Hippocampus* **16**, 1026–1031 (2006).
- 531 16. Eichenbaum, H., Dudchenko, P., Wood, E., Shapiro, M. & Tanila, H. The hippocampus,
532 memory, and place cells: is it spatial memory or a memory space? *Neuron* **23**, 209–226
533 (1999).
- 534 17. Burgess, N., Maguire, E. A. & O'Keefe, J. The human hippocampus and spatial and
535 episodic memory. *Neuron* **35**, 625–641 (2002).
- 536 18. Chrobak, J. J., Lörcincz, A. & Buzsáki, G. Physiological patterns in the hippocampo-
537 entorhinal cortex system. *Hippocampus* **10**, 457–465 (2000).
- 538 19. O'Keefe, J. Place units in the hippocampus of the freely moving rat. *Exp. Neurol.* **51**, 78–
539 109 (1976).
- 540 20. Skaggs, W. E., McNaughton, B. L., Wilson, M. A. & Barnes, C. A. Theta phase precession
541 in hippocampal neuronal populations and the compression of temporal sequences.
542 *Hippocampus* **6**, 149–172 (1996).

21. Chadwick, A., van Rossum, M. C. & Nolan, M. F. Flexible theta sequence compression mediated via phase precessing interneurons. *Elife* **5**, (2016).
22. Ziv, Y. *et al.* Long-term dynamics of CA1 hippocampal place codes. *Nat. Neurosci.* **16**, 264–266 (2013).
23. Rich, P. D., Liaw, H.-P. & Lee, A. K. Place cells. Large environments reveal the statistical structure governing hippocampal representations. *Science* **345**, 814–817 (2014).
24. Thompson, L. T. & Best, P. J. Place cells and silent cells in the hippocampus of freely-behaving rats. *J. Neurosci.* **9**, 2382–2390 (1989).
25. Epsztein, J., Brecht, M. & Lee, A. K. Intracellular determinants of hippocampal CA1 place and silent cell activity in a novel environment. *Neuron* **70**, 109–120 (2011).
26. Cohen, J. D., Bolstad, M. & Lee, A. K. Experience-dependent shaping of hippocampal CA1 intracellular activity in novel and familiar environments. *Elife* **6**, e23040 (2017).
27. Silva, A. J., Zhou, Y., Rogerson, T., Shobe, J. & Balaji, J. Molecular and cellular approaches to memory allocation in neural circuits. *Science* **326**, 391–395 (2009).
28. Zhou, Y. *et al.* CREB regulates excitability and the allocation of memory to subsets of neurons in the amygdala. *Nat. Neurosci.* **12**, 1438–1443 (2009).
29. Yiu, A. P. *et al.* Neurons are recruited to a memory trace based on relative neuronal excitability immediately before training. *Neuron* **83**, 722–735 (2014).
30. Rashid, A. J. *et al.* Competition between engrams influences fear memory formation and recall. *Science* **353**, 383–387 (2016).
31. Rubin, A., Geva, N., Sheintuch, L. & Ziv, Y. Hippocampal ensemble dynamics timestamp events in long-term memory. *Elife* **4**, (2015).
32. Cai, D. J. *et al.* A shared neural ensemble links distinct contextual memories encoded close in time. *Nature* **534**, 115–118 (2016).
33. Kastellakis, G., Silva, A. J. & Poirazi, P. Linking Memories across Time via Neuronal and Dendritic Overlaps in Model Neurons with Active Dendrites. *Cell Rep.* **17**, 1491–1504

- 569 (2016).
- 570 34. Dudek, S. M. & Fields, R. D. Somatic action potentials are sufficient for late-phase LTP-
571 related cell signaling. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 3962–3967 (2002).
- 572 35. Deisseroth, K., Bito, H. & Tsien, R. W. Signaling from synapse to nucleus: postsynaptic
573 CREB phosphorylation during multiple forms of hippocampal synaptic plasticity. *Neuron* **16**,
574 89–101 (1996).
- 575 36. Lopez de Armentia, M. *et al.* cAMP response element-binding protein-mediated gene
576 expression increases the intrinsic excitability of CA1 pyramidal neurons. *J. Neurosci.* **27**,
577 13909–13918 (2007).
- 578 37. Disterhoft, J. F. & Oh, M. M. Learning, aging and intrinsic neuronal plasticity. *Trends*
579 *Neurosci.* **29**, 587–599 (2006).
- 580 38. Lovett-Barron, M. *et al.* Dendritic inhibition in the hippocampus supports fear learning.
581 *Science* **343**, 857–863 (2014).
- 582 39. Stefanelli, T., Bertollini, C., Lüscher, C., Muller, D. & Mendez, P. Hippocampal Somatostatin
583 Interneurons Control the Size of Neuronal Memory Ensembles. *Neuron* **89**, 1074–1085
584 (2016).
- 585 40. Mankin, E. A. *et al.* Neuronal code for extended time in the hippocampus. *Proc. Natl. Acad.*
586 *Sci. U. S. A.* **109**, 19462–19467 (2012).
- 587 41. Mankin, E. A., Diehl, G. W., Sparks, F. T., Leutgeb, S. & Leutgeb, J. K. Hippocampal CA2
588 activity patterns change over time to a larger extent than between spatial contexts. *Neuron*
589 **85**, 190–201 (2015).
- 590 42. Lipowsky, R., Gillessen, T. & Alzheimer, C. Dendritic Na⁺ channels amplify EPSPs in
591 hippocampal CA1 pyramidal cells. *J. Neurophysiol.* **76**, 2181–2191 (1996).
- 592 43. Magee, J. C. Dendritic Ih normalizes temporal summation in hippocampal CA1 neurons.
593 *Nat. Neurosci.* **2**, 508–514 (1999).
- 594 44. Ngo-Anh, T. J. *et al.* SK channels and NMDA receptors form a Ca²⁺-mediated feedback

595 loop in dendritic spines. *Nat. Neurosci.* **8**, 642–649 (2005).

596 45. Losonczy, A., Makara, J. K. & Magee, J. C. Compartmentalized dendritic plasticity and
597 input feature storage in neurons. *Nature* **452**, 436–441 (2008).

598 46. Amaral, D. G. & Witter, M. P. The three-dimensional organization of the hippocampal
599 formation: a review of anatomical data. *Neuroscience* **31**, 571–591 (1989).

600 47. Freund, T. F. & Buzsáki, G. Interneurons of the hippocampus. *Hippocampus* **6**, 347–470
601 (1996).

602 48. Middleton, S. J. & McHugh, T. J. Silencing CA3 disrupts temporal coding in the CA1
603 ensemble. *Nat. Neurosci.* **19**, 945–951 (2016).

604 49. Schlesiger, M. I. *et al.* The medial entorhinal cortex is necessary for temporal organization
605 of hippocampal neuronal activity. *Nat. Neurosci.* **18**, 1123–1132 (2015).

606 50. Magee, J. C. Dendritic integration of excitatory synaptic input. *Nat. Rev. Neurosci.* **1**, 181–
607 190 (2000).

608 51. Nolan, M. F. *et al.* A behavioral role for dendritic integration: HCN1 channels constrain
609 spatial memory and plasticity at inputs to distal dendrites of CA1 pyramidal neurons. *Cell*
610 **119**, 719–732 (2004).

611 52. Tsay, D., Dudman, J. T. & Siegelbaum, S. A. HCN1 channels constrain synaptically evoked
612 Ca²⁺ spikes in distal dendrites of CA1 pyramidal neurons. *Neuron* **56**, 1076–1089 (2007).

613 53. Cai, X. *et al.* Unique roles of SK and Kv4.2 potassium channels in dendritic integration.
614 *Neuron* **44**, 351–364 (2004).

615 54. Lee, D., Lin, B.-J. & Lee, A. K. Hippocampal place fields emerge upon single-cell
616 manipulation of excitability during behavior. *Science* **337**, 849–853 (2012).

617 55. Nevian, T., Larkum, M. E., Polsky, A. & Schiller, J. Properties of basal dendrites of layer 5
618 pyramidal neurons: a direct patch-clamp recording study. *Nat. Neurosci.* **10**, 206–214
619 (2007).

620 56. Druckmann, S. *et al.* Structured synaptic connectivity between hippocampal regions.

621 *Neuron* **81**, 629–640 (2014).

622 57. Wilson, D. E., Whitney, D. E., Scholl, B. & Fitzpatrick, D. Orientation selectivity and the
623 functional clustering of synaptic inputs in primary visual cortex. *Nat. Neurosci.* **19**, 1003–
624 1009 (2016).

625 58. Iacaruso, M. F., Gasler, I. T. & Hofer, S. B. Synaptic organization of visual space in primary
626 visual cortex. *Nature* (2017). doi:10.1038/nature23019

627 59. Haider, B., Häusser, M. & Carandini, M. Inhibition dominates sensory responses in the
628 awake cortex. *Nature* **493**, 97–100 (2013).

629 60. Atallah, B. V., Bruns, W., Carandini, M. & Scanziani, M. Parvalbumin-expressing
630 interneurons linearly transform cortical responses to visual stimuli. *Neuron* **73**, 159–170
631 (2012).

632 61. Atallah, B. V., Scanziani, M. & Carandini, M. Atallah et al. reply. *Nature* **508**, E3 (2014).

633 62. Grienberger, C., Milstein, A. D., Bittner, K. C., Romani, S. & Magee, J. C. Inhibitory
634 suppression of heterogeneously tuned excitation enhances spatial coding in CA1 place
635 cells. *Nat. Neurosci.* (2017). doi:10.1038/nn.4486

636 63. Wilson, N. R., Runyan, C. A., Wang, F. L. & Sur, M. Division and subtraction by distinct
637 cortical inhibitory networks in vivo. *Nature* **488**, 343–348 (2012).

638 64. Lee, S.-H. *et al.* Activation of specific interneurons improves V1 feature selectivity and
639 visual perception. *Nature* **488**, 379–383 (2012).

640 65. Cottam, J. C. H., Smith, S. L. & Häusser, M. Target-specific effects of somatostatin-
641 expressing interneurons on neocortical visual processing. *J. Neurosci.* **33**, 19567–19578
642 (2013).

643 66. Royer, S. *et al.* Control of timing, rate and bursts of hippocampal place cells by dendritic
644 and somatic inhibition. *Nat. Neurosci.* **15**, 769–775 (2012).

645 67. Yoon, K. *et al.* Specific evidence of low-dimensional continuous attractor dynamics in grid
646 cells. *Nat. Neurosci.* **16**, 1077–1084 (2013).

- 647 68. Fyhn, M., Hafting, T., Treves, A., Moser, M.-B. & Moser, E. I. Hippocampal remapping and
648 grid realignment in entorhinal cortex. *Nature* **446**, 190–194 (2007).
- 649 69. Pastoll, H., Ramsden, H. L. & Nolan, M. F. Intrinsic electrophysiological properties of
650 entorhinal cortex stellate cells and their contribution to grid cell firing fields. *Front. Neural*
651 *Circuits* **6**, 17 (2012).
- 652 70. Ray, S. *et al.* Grid-layout and theta-modulation of layer 2 pyramidal neurons in medial
653 entorhinal cortex. *Science* **343**, 891–896 (2014).
- 654 71. Couey, J. J. *et al.* Recurrent inhibitory circuitry as a mechanism for grid formation. *Nat.*
655 *Neurosci.* **16**, 318–324 (2013).
- 656 72. Pastoll, H., Solanka, L., van Rossum, M. C. W. & Nolan, M. F. Feedback inhibition enables
657 θ -nested γ oscillations and grid firing fields. *Neuron* **77**, 141–154 (2013).
- 658 73. Nolan, M. F., Dudman, J. T., Dodson, P. D. & Santoro, B. HCN1 channels control resting
659 and active integrative properties of stellate cells from layer II of the entorhinal cortex. *J.*
660 *Neurosci.* **27**, 12440–12451 (2007).
- 661 74. Giocomo, L. M. *et al.* Grid cells use HCN1 channels for spatial scaling. *Cell* **147**, 1159–
662 1170 (2011).
- 663 75. Robinson, R. B. & Siegelbaum, S. A. Hyperpolarization-activated cation currents: from
664 molecules to physiological function. *Annu. Rev. Physiol.* **65**, 453–480 (2003).
- 665 76. Garden, D. L. F., Dodson, P. D., O'Donnell, C., White, M. D. & Nolan, M. F. Tuning of
666 synaptic integration in the medial entorhinal cortex to the organization of grid cell firing
667 fields. *Neuron* **60**, 875–889 (2008).
- 668 77. Giocomo, L. M., Zilli, E. A., Fransén, E. & Hasselmo, M. E. Temporal frequency of
669 subthreshold oscillations scales with entorhinal grid cell field spacing. *Science* **315**, 1719–
670 1722 (2007).
- 671 78. Schmidt-Hieber, C. *et al.* Active dendritic integration as a mechanism for robust and precise
672 grid cell firing. *Nat. Neurosci.* (2017). doi:10.1038/nn.4582

673 79. Palmer, L. M. *et al.* NMDA spikes enhance action potential generation during sensory input.
674 *Nat. Neurosci.* **17**, 383–390 (2014).

675 80. Lavzin, M., Rapoport, S., Polsky, A., Garion, L. & Schiller, J. Nonlinear dendritic processing
676 determines angular tuning of barrel cortex neurons in vivo. *Nature* **490**, 397–401 (2012).

677 81. Smith, S. L., Smith, I. T., Branco, T. & Häusser, M. Dendritic spikes enhance stimulus
678 selectivity in cortical neurons in vivo. *Nature* **503**, 115–120 (2013).

679 82. Poirazi, P. & Mel, B. W. Impact of active dendrites and structural plasticity on the memory
680 capacity of neural tissue. *Neuron* **29**, 779–796 (2001).

681 83. Buzsáki, G. & Moser, E. I. Memory, navigation and theta rhythm in the hippocampal-
682 entorhinal system. *Nat. Neurosci.* **16**, 130–138 (2013).

683 84. Martin, S. J., Grimwood, P. D. & Morris, R. G. Synaptic plasticity and memory: an
684 evaluation of the hypothesis. *Annu. Rev. Neurosci.* **23**, 649–711 (2000).

685 85. Bliss, T. V. & Collingridge, G. L. A synaptic model of memory: long-term potentiation in the
686 hippocampus. *Nature* **361**, 31–39 (1993).

687 86. Morris, R. G., Anderson, E., Lynch, G. S. & Baudry, M. Selective impairment of learning
688 and blockade of long-term potentiation by an N-methyl-D-aspartate receptor antagonist,
689 AP5. *Nature* **319**, 774–776 (1986).

690 87. Tsien, J. Z., Huerta, P. T. & Tonegawa, S. The essential role of hippocampal CA1 NMDA
691 receptor-dependent synaptic plasticity in spatial memory. *Cell* **87**, 1327–1338 (1996).

692 88. Kentros, C. *et al.* Abolition of long-term stability of new hippocampal place cell maps by
693 NMDA receptor blockade. *Science* **280**, 2121–2126 (1998).

694 89. McHugh, T. J., Blum, K. I., Tsien, J. Z., Tonegawa, S. & Wilson, M. A. Impaired
695 hippocampal representation of space in CA1-specific NMDAR1 knockout mice. *Cell* **87**,
696 1339–1349 (1996).

697 90. Moosmang, S. *et al.* Role of hippocampal Cav1.2 Ca²⁺ channels in NMDA receptor-
698 independent synaptic plasticity and spatial memory. *J. Neurosci.* **25**, 9883–9892 (2005).

91. Sheffield, M. E. J. & Dombeck, D. A. Calcium transient prevalence across the dendritic
arbour predicts place field properties. *Nature* **517**, 200–204 (2015).
92. Takahashi, H. & Magee, J. C. Pathway interactions and synaptic plasticity in the dendritic
tuft regions of CA1 pyramidal neurons. *Neuron* **62**, 102–111 (2009).
93. Bittner, K. C. *et al.* Conjunctive input processing drives feature selectivity in hippocampal
CA1 neurons. *Nat. Neurosci.* **18**, 1133–1142 (2015).
94. Golding, N. L., Staff, N. P. & Spruston, N. Dendritic spikes as a mechanism for cooperative
long-term potentiation. *Nature* **418**, 326–331 (2002).
95. Dudman, J. T., Tsay, D. & Siegelbaum, S. A. A role for synaptic inputs at distal dendrites:
instructive signals for hippocampal long-term plasticity. *Neuron* **56**, 866–879 (2007).
96. Frank, L. M. Hippocampal Plasticity across Multiple Days of Exposure to Novel
Environments. *Journal of Neuroscience* **24**, 7681–7689 (2004).
97. Hill, A. J. First occurrence of hippocampal spatial firing in a new environment. *Exp. Neurol.*
62, 282–297 (1978).
98. Brandalise, F. & Gerber, U. Mossy fiber-evoked subthreshold responses induce timing-
dependent plasticity at hippocampal CA3 recurrent synapses. *Proc. Natl. Acad. Sci. U. S.*
A. **111**, 4303–4308 (2014).
99. Brandalise, F., Carta, S., Helmchen, F., Lisman, J. & Gerber, U. Dendritic NMDA spikes are
necessary for timing-dependent associative LTP in CA3 pyramidal cells. *Nat. Commun.* **7**,
13480 (2016).
100. Lörincz, A., Notomi, T., Tamás, G., Shigemoto, R. & Nusser, Z. Polarized and
compartment-dependent distribution of HCN1 in pyramidal cell dendrites. *Nat. Neurosci.* **5**,
1185–1193 (2002).
101. Hussaini, S. A., Kempadoo, K. A., Thuaux, S. J., Siegelbaum, S. A. & Kandel, E. R.
Increased size and stability of CA1 and CA3 place fields in HCN1 knockout mice. *Neuron*
72, 643–653 (2011).

725 102. Maroso, M. *et al.* Cannabinoid Control of Learning and Memory through HCN Channels.
 726 *Neuron* **89**, 1059–1073 (2016).
 727 103. Magee, J. C. Dendritic Ih normalizes temporal summation in hippocampal CA1 neurons.
 728 *Nat. Neurosci.* **2**, 848 (1999).
 729 104. Buzsáki, G. Theta oscillations in the hippocampus. *Neuron* **33**, 325–340 (2002).
 730 105. O'Keefe, J. & Recce, M. L. Phase relationship between hippocampal place units and the
 731 EEG theta rhythm. *Hippocampus* **3**, 317–330 (1993).
 732 106. Bender, F. *et al.* Theta oscillations regulate the speed of locomotion via a hippocampus to
 733 lateral septum pathway. *Nat. Commun.* **6**, 8521 (2015).
 734 107. Freund, T. F. & Antal, M. GABA-containing neurons in the septum control inhibitory
 735 interneurons in the hippocampus. *Nature* **336**, 170–173 (1988).
 736 108. Boyce, R., Glasgow, S. D., Williams, S. & Adamantidis, A. Causal evidence for the role of
 737 REM sleep theta rhythm in contextual memory consolidation. *Science* **352**, 812–816
 738 (2016).
 739 109. Gonzalez-Sulser, A. *et al.* GABAergic projections from the medial septum selectively inhibit
 740 interneurons in the medial entorhinal cortex. *J. Neurosci.* **34**, 16739–16743 (2014).
 741 110. Tóth, K., Freund, T. F. & Miles, R. Disinhibition of rat hippocampal pyramidal cells by
 742 GABAergic afferents from the septum. *J. Physiol.* **500 (Pt 2)**, 463–474 (1997).
 743 111. Mizuseki, K., Sirota, A., Pastalkova, E. & Buzsáki, G. Theta oscillations provide temporal
 744 windows for local circuit computation in the entorhinal-hippocampal loop. *Neuron* **64**, 267–
 745 280 (2009).
 746 112. Pike, F. G. *et al.* Distinct frequency preferences of different types of rat hippocampal
 747 neurones in response to oscillatory input currents. *J. Physiol.* **529 Pt 1**, 205–213 (2000).
 748 113. Zemankovics, R., Káli, S. & Paulsen, O. Differences in subthreshold resonance of
 749 hippocampal pyramidal cells and interneurons: the role of h⁺ current and passive
 750 membrane characteristics. *The Journal of* (2010).

114. Leung, L. S. & Yu, H. W. Theta-frequency resonance in hippocampal CA1 neurons in vitro demonstrated by sinusoidal current injection. *J. Neurophysiol.* **79**, 1592–1596 (1998).
115. Erchova, I., Kreck, G., Heinemann, U. & Herz, A. V. M. Dynamics of rat entorhinal cortex layer II and III cells: characteristics of membrane potential resonance at rest predict oscillation properties near threshold. *J. Physiol.* **560**, 89–110 (2004).
116. Hu, H., Vervaeke, K. & Storm, J. F. Two forms of electrical resonance at theta frequencies, generated by M-current, h-current and persistent Na⁺ current in rat hippocampal pyramidal cells. *J. Physiol.* **545**, 783–805 (2002).
117. Peters, H. C., Hu, H., Pongs, O., Storm, J. F. & Isbrandt, D. Conditional transgenic suppression of M channels in mouse brain reveals functions in neuronal excitability, resonance and behavior. *Nat. Neurosci.* **8**, 51–60 (2005).
118. Borel, M., Guadagna, S., Jang, H. J., Kwag, J. & Paulsen, O. Frequency dependence of CA3 spike phase response arising from h-current properties. *Front. Cell. Neurosci.* **7**, 263 (2013).
119. Ness, T. V., Remme, M. W. H. & Einevoll, G. T. Active subthreshold dendritic conductances shape the local field potential. *J. Physiol.* **594**, 3809–3825 (2016).
120. Stark, E. *et al.* Inhibition-induced theta resonance in cortical circuits. *Neuron* **80**, 1263–1276 (2013).
121. Alonso, A. & Llinás, R. R. Subthreshold Na⁺-dependent theta-like rhythmicity in stellate cells of entorhinal cortex layer II. *Nature* **342**, 175–177 (1989).
122. Leung, L. W. & Yim, C. Y. Intrinsic membrane potential oscillations in hippocampal neurons in vitro. *Brain Res.* **553**, 261–274 (1991).
123. O'Keefe, J. & Burgess, N. Dual phase and rate coding in hippocampal place cells: theoretical significance and relationship to entorhinal grid cells. *Hippocampus* **15**, 853–866 (2005).
124. Fernandez, F. R. & White, J. A. Artificial synaptic conductances reduce subthreshold

oscillations and periodic firing in stellate cells of the entorhinal cortex. *J. Neurosci.* **28**, 3790–3803 (2008).

125. Jaramillo, J. & Kempter, R. Phase precession: a neural code underlying episodic memory? *Curr. Opin. Neurobiol.* **43**, 130–138 (2017).

126. Leung, L. S. A model of intracellular θ phase precession dependent on intrinsic subthreshold membrane currents. *J. Neurosci.* **31**, 12282–12296 (2011).

127. Harris, K. D. *et al.* Spike train dynamics predicts theta-related phase precession in hippocampal pyramidal cells. *Nature* **417**, 738–741 (2002).

128. Magee, J. C. Dendritic mechanisms of phase precession in hippocampal CA1 pyramidal neurons. *J. Neurophysiol.* **86**, 528–532 (2001).

129. Mehta, M. R., Lee, A. K. & Wilson, M. A. Role of experience and oscillations in transforming a rate code into a temporal code. *Nature* **417**, 741–746 (2002).

130. Mehta, M. R., Quirk, M. C. & Wilson, M. A. Experience-dependent asymmetric shape of hippocampal receptive fields. *Neuron* **25**, 707–715 (2000).

131. Chadwick, A., van Rossum, M. C. W. & Nolan, M. F. Independent theta phase coding accounts for CA1 population sequences and enables flexible remapping. *Elife* **4**, (2015).

132. Eggink, H., Mertens, P., Storm, E. & Giocomo, L. M. Hyperpolarization-activated cyclic nucleotide-gated 1 independent grid cell-phase precession in mice. *Hippocampus* **24**, 249–256 (2014).

133. Kim, S., Guzman, S. J., Hu, H. & Jonas, P. Active dendrites support efficient initiation of dendritic spikes in hippocampal CA3 pyramidal neurons. *Nat. Neurosci.* **15**, 600–606 (2012).

134. Makara, J. K. & Magee, J. C. Variable dendritic integration in hippocampal CA3 pyramidal neurons. *Neuron* **80**, 1438–1450 (2013).

135. Krueppel, R., Remy, S. & Beck, H. Dendritic Integration in Hippocampal Dentate Granule Cells. *Neuron* **71**, 512–528 (2011).

136. Branco, T., Clark, B. A. & Häusser, M. Dendritic discrimination of temporal input sequences in cortical neurons. *Science* **329**, 1671–1675 (2010).
137. Stocca, G., Schmidt-Hieber, C. & Bischofberger, J. Differential dendritic Ca²⁺ signalling in young and mature hippocampal granule cells. *J. Physiol.* **586**, 3795–3811 (2008).
138. McHugh, T. J. *et al.* Dentate gyrus NMDA receptors mediate rapid pattern separation in the hippocampal network. *Science* **317**, 94–99 (2007).
139. Chavlis, S., Petrantonis, P. C. & Poirazi, P. Dendrites of dentate gyrus granule cells contribute to pattern separation by controlling sparsity. *Hippocampus* **27**, 89–110 (2017).
140. Schmidt-Hieber, C., Jonas, P. & Bischofberger, J. Subthreshold dendritic signal processing and coincidence detection in dentate gyrus granule cells. *J. Neurosci.* **27**, 8430–8441 (2007).
141. Valero, M. *et al.* Determinants of different deep and superficial CA1 pyramidal cell dynamics during sharp-wave ripples. *Nat. Neurosci.* **18**, 1281–1290 (2015).
142. Gan, J., Weng, S.-M., Pernía-Andrade, A. J., Csicsvari, J. & Jonas, P. Phase-Locked Inhibition, but Not Excitation, Underlies Hippocampal Ripple Oscillations in Awake Mice In Vivo. *Neuron* **93**, 308–314 (2017).
143. Cobb, S. & Lawrence, J. J. Neuromodulation of Hippocampal Cells and Circuits. in *Hippocampal Microcircuits* 187–246 (2010).
144. Marder, E. Neuromodulation of neuronal circuits: back to the future. *Neuron* **76**, 1–11 (2012).
145. Martinello, K. *et al.* Cholinergic afferent stimulation induces axonal function plasticity in adult hippocampal granule cells. *Neuron* **85**, 346–363 (2015).
146. Hasselmo, M. E., Hay, J., Ilyn, M. & Gorchetnikov, A. Neuromodulation, theta rhythm and rat spatial navigation. *Neural Netw.* **15**, 689–707 (2002).
147. Kramer, R. H., Mouro, A. & Adesnik, H. Optogenetic pharmacology for control of native neuronal signaling proteins. *Nat. Neurosci.* **16**, 816–823 (2013).

148. Hamel, E. J. O., Grewe, B. F., Parker, J. G. & Schnitzer, M. J. Cellular level brain imaging in behaving mammals: an engineering approach. *Neuron* **86**, 140–159 (2015).
149. Gong, Y. *et al.* High-speed recording of neural spikes in awake mice and flies with a fluorescent voltage sensor. *Science* **350**, 1361–1366 (2015).
150. Koenig, J., Linder, A. N., Leutgeb, J. K. & Leutgeb, S. The spatial periodicity of grid cells is not sustained during reduced theta oscillations. *Science* **332**, 592–595 (2011).

Figure 1. Membrane potential ramp and intracellular phase precession during place and grid field crossings. (a) Top panels show example recordings from a place cell (left, adapted from ref. ¹⁵⁰) and from a grid cell (right, adapted from ref. ⁹). Bottom panels show simultaneous membrane potential and LFP recordings during firing field crossings. In both CA1 pyramidal cells (left, adapted from ref. ⁶²) and in MEC stellate cells (right, adapted from ref. ¹¹), firing during field crossings is driven by a sustained membrane potential depolarization^{10–12}. (b) Average firing rate (upper) and membrane potential (lower) of CA1 pyramidal cells (left, adapted from ref. ¹⁰) and MEC stellate cells (right, adapted from ref. ¹¹) during field crossings. (c) Action potential phase relative to the local field potential, membrane potential theta oscillation phase with respect to the local field potential and action potential phase with respect to the membrane potential theta oscillation, are each plotted as a function of position within the rate coded firing field, for a CA1 pyramidal cell (left, adapted from ref. ¹⁰) and MEC stellate cells (right, adapted from ref. ¹¹).

Figure 2. Excitability and place cell selection. (a) Schematised differences between excitable and silent cells recorded in a novel environment²⁵. Excitable cells are more likely to acquire place firing fields as the threshold depolarization required to trigger an action potential is reduced, enabling them to respond with action potentials to a ramp depolarization that is insufficient to trigger output from a silent cell. When activated the excitable cells tend to fire spike bursts, whereas the silent cells do not. (b) Hypothesized model for roles of excitability differences in linking of memories^{27,33}. The excitability of neurons storing information about a recent event is selectively increased. Because spikes required for activity-dependent associative plasticity are more likely to occur in excitable neurons, subsequent events occurring within a period determined by the duration of the increase in excitability are captured to the same neurons. Events occurring after excitability of the previously activated neurons has decayed to baseline are stored by different neurons. (c) Hypothesized model for roles of excitability differences in establishing a temporal context for spatial codes. In this scheme the identity of excitable CA1 pyramidal cells evolves on a time scale of days. The probability of a neuron firing action potentials within its place field is greatest during the periods in which it is most excitable. In this way, the set of a neurons representing a location on a particular day can

be used to generate a timestamp for when an event takes place³¹.

Figure 3. Candidate models of nonlinear integration during firing field crossings in hyperpolarized or depolarized neurons. (a) Schematised synaptic integration in a model that computes the arithmetic (“linear”) sum of its inputs. Synaptic inputs are stronger inside (IN) than outside (OUT) of the field. While this model produces a firing field when the cell is depolarized (right), it predicts a subthreshold membrane potential field in a hyperpolarized neuron (left), contradicting experimental data⁵⁴. (b) Same scheme as in (a) for a model neuron that integrates inputs nonlinearly, and strongly attenuates EPSPs as they propagate along the dendritic tree. The strong attenuation produces subthreshold membrane potential fields that are indistinguishable inside and outside of the field, while the nonlinear mechanism boosts EPSPs sufficiently to produce a firing field when the cell is depolarized, consistent with experimental recordings⁵⁴. (c) Same scheme as in (a) for a model neuron with distinct nonlinear integration functions for inputs arriving inside or outside of the field. Although synaptic weights are the same inside and outside the field, EPSPs inside the field are boosted when the neuron is depolarized because of a lower threshold for engaging nonlinear mechanisms. In hyperpolarized neurons, synaptic inputs inside and outside the field produce similar somatic depolarization, resulting in a lack of a distinct membrane potential field, consistent with experimental data⁵⁴.

Figure 4. Comparison of the effect of local inhibition on visual and spatial receptive fields. (a) Left panel, experimentally recorded IPSCs (blue) and EPSCs (red) during presentation of six different grating orientations. Inhibition (blue curve) is less tuned to orientation than excitation (red curve), leading to an increase in the excitation-inhibition ratio in the center of the receptive field. Middle and right panels, membrane potential (middle) and firing rate (right) tuning to grating orientation in a model of a L2/3 visual cortex pyramidal cell. Model responses are derived from experimental recordings of IPSCs and EPSCs shown in the left panel. Black: control; green: suppression of PV+ interneurons by light-activating archeorhodopsin. The model reproduces an experimentally observed linear-threshold transformation of firing rate tuning curves by inhibition from PV+ interneurons. Adapted from ⁶⁰. (b) Left panel, ratio of excitatory and inhibitory currents (E-I ratio) during a firing field crossing in a place cell model constrained by experimental data. Spatially uniform inhibition leads to an increase in the E-I ratio during the field crossing. Middle and right panels, experimentally determined spatial tuning of the membrane potential (middle) and firing rate (right) to spatial position in place cells. Data are aligned to the place field center. Black: control; orange: suppression of GAD2+ or VGAT+ interneurons by light-activating archeorhodopsin. Adopted from ⁶². Note the similarity of the effect of suppressing local inhibition on visual (a) and spatial (b) tuning curves.

Figure 5. Theta resonant responses of hippocampal neurons. (a) When recorded in brain slices CA1 pyramidal neurons preferentially respond to inputs oscillating at theta frequencies, whereas fast spiking interneurons prefer higher frequency oscillatory inputs (left panels)¹¹². (b) To

912 investigate resonant firing in behaving animals parvalbumin interneurons expressing
913 channelrhodopsin were stimulated at frequencies up to 30 Hz¹²⁰. The coherence between the
914 stimulus and the spiking response is plotted as a function of frequency for CA1 pyramidal cells
915 (upper right) and interneurons (lower right). Consistent with in vitro data CA1 pyramidal cells
916 prefer inputs in the theta frequency band.

917

918

a **b** **c**

a **b** **c**

a Orientation tuning in visual cortex
(Atallah et al., 2012)

b Spatial tuning in hippocampus
(Grienberger et al., 2017)

a

Brain slice

b

Behaving animal